# Bursting firing in ventromedial hypothalamic neurons exerts dual control of anxiety-like behavior and energy expenditure

Jie Shao[1,2,4], Dashuang Gao[1,2,4], Yunhui Liu[1,2,4], Shanping Chen[1], Nian Liu[1], Lu Zhang[1], Xinyi Zhou[1,2], Qian Xiao[1,2], Liping Wang[1], Hailan Hu[3], Fan Yang[1,5,]*

[1]The Brain Cognition and Brain Disease Institute, Shenzhen Institutes of Advanced Technology, Chinese Academy of Sciences, Shenzhen-Hong Kong Institute of Brain Science-Shenzhen Fundamental Research Institutions, Shenzhen 518055, China.

[2]University of Chinese Academy of Sciences, Beijing 100049, China.

[3]Department of Psychiatry of First Affiliated Hospital, Zhejiang University School of Medicine, Hangzhou 310058, China.

[4]These authors contributed equally.

[5]Lead contact

*Correspondence to: Fan Yang, Shenzhen Institutes of Advanced Technology, Chinese Academy of Sciences, 1068 Xueyuan Boulevard, University Town of Shenzhen, Xili Nanshan, Shenzhen 518055, China. Email: fan.yang@siat.ac.cn;

**Abstract**

Exposure to chronic stress induces anxiety-like behavior and metabolic changes in animals, resulting in adaptive or maladaptive responses to the stressful environment. Recent studies have indicated the dorsomedial ventromedial hypothalamus (dmVMH) as an important hub that regulates both anxiety and energy homeostasis. However, up to now, how dmVMH neurons exert dual control of chronic stress-induced anxiety and energy expenditure remains poorly understood. Here, we established a chronic-stress mouse model that induced anxiety-like behavior, reduced food consumption, and decreased energy expenditure. We found that *c-fos* expression increased and theta band power is higher in the dmVMH after chronic stress, and the proportion of burst firing neurons significantly increased, which was mediated by elevated expression of T-type calcium channel Cav 3.1. Optogenetically evoked burst firing of dmVMH neurons induced anxiety-like behavior, shifted the respiratory exchange ratio toward fat oxidation, and decreased food intake, while knockdown of Cav3.1 in the dmVMH had the opposite effects. Collectively, our study first revealed an important role of dmVMH burst firing in the dual regulation of anxiety-like behavior and energy expenditure, and identified Cav 3.1 as a crucial regulator of the activity of the burst firing neurons in dmVMH. These molecular and cellular level findings will advance our understanding of the chronic stress-induced emotional malfunction and energy expenditure disorders.

**Keywords:** Chronic stress; VMH; Burst firing; Anxiety; Energy expenditure

**Introduction**

Exposure to stressors regulate neural activity that integrate the aversive behavior and energy expenditure to facilitate stress coping and survival[1-3], however long-term stress causes adverse effects, impairing both mental and physiological functions [4,5] including anxiety-like behavior [6,7] and metabolic malfunctions in glucose [8,9] and lipid tissues [3,10]. Particular effort has been expended to identify the crucial central nodes that interconnect the regulation of emotion and metabolic processes [3,9,11]. The hypothalamus, especially the ventromedial hypothalamus (VMH), has been found to play crucial roles in controlling both innate survival behaviors [12,13] and energy homeostasis [11,14,15].

The VMH is an evolutionarily conserved deep subcortical nucleus[16]. While the ventrolateral part (vl) of the VMH modulates a series of social behaviors [17-19], the dorsomedial part (dm) is specifically involved in maintaining energy homeostasis [14,15,20,21] and stress response[13,22]. Oscillations in the dmVMH can act on sympathetic excitation and regulate energy expenditure [23,24]. Steroidogenic factor-1 (SF-1) -expressing neurons are enriched in the dmVMH and play important roles in controlling food intake and energy expenditure [14,15,20,21]. Optogenetic activation of VMH SF-1 neurons induces aversive behavior in mice [13,22]. Furthermore, SF-1 neurons can act as a nutrient-sensitive switch between feeding and anxiety states when facing potential stresses [2]. Given the complicated functions of SF-1 neurons [19], it is important to dissect how molecular or electrophysiological distinct neuronal subtypes regulate

emotional state and energy homeostasis. However up to now, how dmVMH neurons

control neuronal firing to regulate chronic stress-induced anxiety and energy metabolic

disorders is elusive.

Recent studies have demonstrated the role of burst firing neurons in regulating

emotional state and brain functions; neuronal burst firing and oscillation are critical for

the transmission of information and regulation of specific physiological processes in

crucial brain areas[25-30]. In the hippocampus, burst firing of CA1 pyramidal neurons play

an important role in regulating N-methyl-d-aspartate (NMDA)-mediated transmission

and the epileptogenic mechanism [25]. In the ventral subiculum, chronic social defeat

stress increases both T-type calcium channel (T-VGCC)-mediated currents and

expression of Cav3.1 protein [29]. In the lateral habenula, NMDA-receptor-and- T-VGCC

-dependent bursting activity are increased after chronic stress and, most strikingly, can

drive depression-like behaviors [26]. Based on these evidences, we decided to investigate

the specific function of bursting firing in the dmVMH, and whether it may be involved

in chronic stress-induced anxiety-like behavior and energy metabolic changes.

In this study, we first identified that dmVMH neurons can be classified into silent,

tonic-firing, and bursting subtypes. We also established a chronic-stress mouse model

that induced anxiety-like behavior, reduced food consumption, and decreased energy

expenditure. We found a significantly increased proportion of bursting neurons under

chronic stress conditions, which was mainly caused by elevated T-type calcium channel

Cav3.1 expression. Importantly, optogenetically-induced dmVMH burst firing drive

anxiety-like behaviors, shifted respiratory exchange ratio and decreased average

energy expenditure in naïve mice. Conversely, knockdown of Cav3.1 in the dmVMH

rescued chronic stress-induced anxiety-like behavior and energy expenditure changes.

Taken together, our study first identified Cav3.1-mediated burst firing pattern in the

dmVMH, and demonstrated its crucial role in regulating anxiety-like behavior and

energy expenditure during the chronic stress.

**Results**

**Chronic stressors induced anxiety-like behavior and energy expenditure change**

Repeated exposure to stressors modifies the activity of neuronal circuits associated

with a variety of abnormalities, including behavior, emotional state, and energy

expenditure [2], which adaptively or maladaptively respond to the stressful situation. To

probe the impact of chronic stress on behavior and energy metabolism, we applied

unpredictable chronic stressors to establish a stress mouse model. During the chronic

stress period, mice were exposed to random stressors for four weeks, as illustrated in

Fig. 1a, with no stressors used in the control group. Both the wild-type and stress

groups were tested in a comprehensive phenotyping paradigm. Consistent with

previous reports [6,7], we found that chronic stress induced a state of anxiety (Fig. 1b-e).

Compared with the control group, mice in the stress group spent less time in the central

area of the open field and open arm of the elevated plus maze without any change in

locomotion distance (Fig. 1c and e). The number of entries into the central area and

open arms also decreased in the stress group compared with the control group (Fig. 1c

and e).

We next explored whether chronic stress affected energy expenditure in stressed mice.

The body weight of each mouse was measured after the chronic stress period, and no

significant difference was observed between the control and stress groups (Fig. 1f).

Both the respiratory exchange ratio (RER) and food intake were monitored for 24 h

using comprehensive laboratory animal monitoring system (CLAMS) cages. Results

showed that the average RER of the stress group was much lower than that of the

control group, which represented a shift toward fat oxidation (Fig. 1g). Energy

expenditure was calculated from oxygen consumption and RER. Average energy

expenditure also showed a significant decrease in stressed mice (Fig. 1h). To determine

the underlying cause of different RER in the treatment groups, we also analyzed food

intake. As expected, caloric intake in stressed mice was significantly lower than that in

the controls, consistent with the known effects of anxiety on food intake (Fig. 1f).

Taken together, these findings suggest that chronic stress can induce anxiety-like

behavior, reduce food consumption, and shift the RER toward fat oxidation.

Given its important role in regulating anxiety, feeding, and energy expenditure [2,21], we

hypothesized that the dmVMH may be involved in the above chronic stress-induced

changes. After chronic stress, mice were perfused, and *c-fos* expression was determined

via immunostaining. Higher *c-fos* expression was found in the dmVMH of chronically

stressed mice than in that of the control group (Fig. 1i), suggesting that more dmVMH

neurons were activated under chronic stress conditions. We then employed *in vivo*

electrophysiology to record neuronal activity in the dmVMH. Mice were implanted

with Ni-Cr electrodes into the dmVMH after chronic stress induction (Fig. 1j), and

dmVMH neuronal activity was monitored after recovery from surgery. The local field

potential (LFP) results demonstrated that stressed mice exhibited higher theta band

power in the dmVMH than that of control mice (Fig. 1k-l). Together with the *c-fos*

staining results, our data suggest that dmVMH neuronal activity is significantly

changed after chronic stress.

**Chronic stressors induced burst firing in dmVMH**

The above experiments revealed enhanced neuronal activity in the dmVMH under

chronic stress, but the underlying mechanism was unclear. To explore which

electrophysiological characteristics of dmVMH neurons changed after chronic stress,

we recorded 84 dmVMH neurons from 38 stressed mice and 85 dmVMH neurons from

32 control mice under whole-cell patch clamp configuration and analyzed the

electrophysiological characteristics. Our data demonstrated that these neurons

displayed shorter onset time and depolarized RMP, on average, compared with the

control group (Fig. 1a). The alterations implied that the overall excitability of dmVMH

neurons was enhanced under chronic stress, consistent with the *c-fos* immunostaining

and *in vivo* electrode recording results.

We carried out a comprehensive analysis of the membrane properties of dmVMH

neurons to quantitatively determine their electrophysiological diversity (Fig. 2b). To

classify these neurons, we measured and analyzed eight electrophysiological

parameters (Table 1) and performed cluster analysis. The resulting dendrogram in Fig.

2b (with rescaled distance shown along the vertical axis) illustrates the similarity

between clusters. Furthermore, cluster analysis showed that all neurons could be

divided into three subtypes: i.e., silent, tonic-firing, bursting. These three dmVMH

neuronal subtypes displayed distinct electrophysiological properties (Fig. 2b, Extended

Data Fig. 1).

We then applied cluster analysis using the same methods as depicted above (Fig. 2d) to

explore overall changes in the electrophysiological properties of dmVMH neurons after

chronic stress. We found that dmVMH neurons in stressed mice could also be divided

into three subtypes. Onset time, RMP and recording site of these three subtypes were

also compared with those of the control group (Extended Data Fig. 2). By analyzing

data from behavior tests and electrophysiological recordings, we found the proportion

of burst firing neurons in stressed group, especially in mice with obvious anxiety-like

behavior (anxiety group), increased significantly compared with that in the control

group, whereas the proportion of the other subtypes decreased (Fig. 2d-e). This

alteration could explain why dmVMH neurons from stressed mice demonstrated a

shorter average onset time of action potentials induced by 50 pA current injection (Fig.

2a), as burst firing neurons displayed a similar onset time as tonic-firing neurons but a

shorter onset time compared with silent neurons (Fig. 2f, Extended Data Fig. 2b). In

addition, the increased proportion of bursting neurons and more depolarized

tonic-firing neurons (Fig. 2f, Extended Data Fig. 2a) may have contributed to the higher

average RMP in dmVMH neurons after chronic stress (Fig. 2a). Furthermore, the

average inter-spike interval (ISI, between first and second spike of a burst) of dmVMH

burst firing neurons in anxiety group was shorter than that of the control group

(unpaired Student's $t$-test, $P = 0.0173$), and a shorter ISI represents higher frequency

firing in a single burst (Fig. 2g). We also found that ISI of dmVMH bursting neurons

was more correlated with the time in central area or open arms in mice of

stress-induced anxiety compared with the mice in control group (Fig. 2h,i). Taken

together, these results demonstrate a possible link between increased bursting activity

in the dmVMH and chronic stress induced anxiety.

**Optogenetic manipulation of burst firing neurons in dmVMH**

Given the enhancement of burst firing in the dmVMH after chronic stress, we

investigated whether induced burst firing of dmVMH neurons alone in unstressed mice

was sufficient to produce a similar response in behavior and energy metabolism. We

first applied optogenetic manipulation to elicit a low-threshold spike and mimic the

generation of bursting in dmVMH neurons (Fig. 3a). *In vitro* patch clamp experiments

on brain slices indicated that yellow light illumination at 0.1 Hz could slightly

hyperpolarize and depolarize bursting neurons periodically, which enabled the

re-activation of calcium channels and generation of burst firing during the light

"ON-OFF" intervals in bursting neurons (Fig. 3b). The evoked burst firing did not

follow the "ON-OFF" intervals when the illumination frequency was over 0.1 Hz (Fig.

3b). In addition, illumination at 0.1 Hz exerted no significant effects on the tonic-firing

or silent neuronal subtypes (Fig. 3c).

We then applied low-frequency yellow light illumination to induce burst firing *in vivo*

and performed behavioral and energy metabolic tests in NpHR-expressing mice (Fig.

3d). This led to decreased residence time in the central area of the open field and open

arm of the elevated plus maze, mimicking the effects of chronic stress-induced

anxiety-like behavior (Fig. 3e,f). We also applied wireless optogenetics to induce burst

firing of neurons and simultaneously monitored the metabolism of mice using CLAMS.

We applied 0.1-Hz yellow light illumination at the start of the test period for 2 h and

repeated the stimulation after 12 h, with consecutive monitoring of mouse energy

metabolism for 24 h. Results indicated that average RER, energy expenditure, and food

intake in the test period decreased significantly compared with the baseline level before

the test period (Fig. 3g-i). Taken together, our results indicated that enhancement of

burst firing induced by local optogenetic manipulation in the dmVMH was sufficient to

affect anxiety-like behavior and energy expenditure changes, which simulated the

phenotypes of chronically stressed mice.

**Burst firing in dmVMH is mediated by T-VGCC**

Bursting is an important firing pattern in neural systems and is essential for specific

information transmission and function regulation[30]. The T-VGCC, including its three

isoforms (Cav3.1, Cav3.2, and Cav3.3), is a pacemaker that can generate LTS [26,31].

Unlike hyperpolarization-activated cyclic nucleotide-gated channel (HCN), the

T-VGCC can induce higher frequency firing and has a threshold near to RMP. It has

been reported that Cav 3.1 and Cav 3.2 alone display strong burst firing with a

low-voltage threshold, whereas Cav3.3 contributes to burst firing in a different way [31].

The expression of T-VGCC is highly correlated with T-type calcium currents, which

directly affect the strength and width of bursting [29,32]. Given the known role of T-VGCC

in burst firing, we examined T-VGCC expression in the dmVMH after chronic stress.

Based on immunohistochemical analysis, Cav3.1, Cav3.2, and Cav3.3 were all found to

be expressed in the dmVMH of naïve mice. The signals of Cav3.1 were much stronger

than those of Cav3.2 and Cav3.3 (Fig. 4a). To investigate whether differential

expression of the T-VGCC occurs after chronic stress, we performed acute dissection of

dmVMH tissue from brain slices and harvested single dmVMH cells after patch

recordings to extract RNA, then performed qRT-PCR to quantify T-VGCC expression.

The expression of Cav3.1 in the dmVMH of stressed mice was significantly higher than

that of the control group, whereas no obvious changes were observed in Cav 3.2 or Cav

3.3 at either the tissue or single-cell level (Fig. 4b-c).

To further confirm the contribution of the T-VGCC in dmVMH burst generation, we

applied mibefradil, an antagonist of T-VGCC, in whole-cell recording experiments.

Results indicated that the burst firing of dmVMH neurons elicited by a 10-pA current

injection was indeed inhibited (Fig. 4d). Furthermore, we found that application of

mibefradil increased the onset time in the dmVMH neurons of the stress group, but not

in the control group, but the change in RMP was similar between the two groups (Fig.

4e). Moreover, as we collected single neurons after whole-cell recordings for T-VGCC

quantification, we combined the single-cell qRT-PCR and electrophysiological

classification data to determine the differential expression of Cav3.1 in the dmVMH

neuronal subtypes. Results showed that Cav3.1 expression was much more enriched in

dmVMH bursting neurons, especially after chronic stress (Fig. 4f). Consistently, the

Cav3.2 antagonist ascorbate [33] did not significantly affect burst firing of dmVMH

neurons (Extended Data Fig. 3). We also investigated the effects of mibefradil on the

frequency-current curves using the same stimulus protocol, and found that the firing

rate of bursting neurons was decreased by bath application of mibefradil($P = 0.007$, Fig.

4g, right), but the changes in the firing rate were much more significant in the stress

group than in the control group ($P = 0.0047$, Fig. 4h, upper). These data support the idea

that the T-VGCC, especially the Cav3.1 isoform, mediates burst firing in the dmVMH.

To investigate the *in vivo* effects of T-VGCC blockade, we bilaterally implanted a

cannula and precisely delivered mibefradil or saline into the dmVMH of chronically

stressed mice to block burst firing (Extended Data Fig. 4a). Results showed that

mibefradil infusion increased the time spent in the central area of the open field and

open arm of the elevated plus maze compared with the control group treated with saline

(Extended Data Fig. 4b). We also investigated energy metabolic changes after oral

administration of mibefradil, and found it to have no effect on food intake. However,

administration of mibefradil significantly rescued the decreased RER found in stressed

mice. Taken together, these data indicate that the T-VGCC in the dmVMH mediates

chronic stress-induced behavioral and energy expenditure changes.

The function of the T-VGCC is tightly related to postsynaptic excitatory/inhibitory state

[34,35]. Several studies have suggested that the generation of burst firing mediated by

T-VGCC activation in certain regions is highly dependent on membrane potential [35,36],

and the NMDA receptor plays a critical role in regulating burst firing [26]. Glutamate is a

common excitatory neurotransmitter and its concentration around neurons may affect

membrane potential [37]. We applied different glutamate receptor antagonists into

artificial cerebrospinal fluid (ACSF) during whole-cell recordings of burst firing neurons. Consistent with previous studies [26], our results indicated that blockade of NMDA receptors, but not AMPA receptors, significantly inhibited burst firing and caused membrane hyperpolarization, which made it more difficult to reach the T-VGCC threshold (Extended Data Fig. 5).

**Knockdown of Cav3.1 in dmVMH decreased burst firing and ameliorated chronic stress response**

Given the important role of Cav 3.1 in mediating burst firing in the dmVMH, we next tested whether down-regulation of Cav 3.1 expression in the dmVMH was sufficient to ameliorate anxiety-like behavior and energy metabolic changes induced by chronic stress. We developed a lentivirus-mediated RNAi method to interfere with expression of Cav3.1 in the dmVMH under chronic stress, with the shRNA-expressing vector injected into the dmVMH bilaterally prior to chronic-stress exposure (Fig. 5a). To test the efficacy of shRNA-mediated Cav3.1 silencing, we used immunofluorescence to confirm the efficient knockdown of Cav3.1 in the dmVMH of stressed mice (Fig. 5b). The effects of Cav3.1 knockdown on burst firing were tested using whole-cell recording on brain slices obtained from Cav 3.1 knockdown mice. Results showed that 7 out of 17 dmVMH neurons from stressed mice displayed evoked burst firing, whereas only 2 out of 13 dmVMH neurons succeeded after Cav3.1 knockdown (Fig. 5b), suggesting burst firing inhibition induced by RNAi of Cav3.1.

To investigate phenotypic changes in Cav3.1 knockdown mice, we carried out

behavioral and energy metabolic tests after viral expression of shRNA of Cav3.1 for

4–5 weeks. Importantly, behavioral tests showed that Cav 3.1 knockdown significantly

increased residence time and entries into the central area of the open field and open arm

of the elevated plus-maze of stressed mice (Fig. 5c,d), suggesting that local knockdown

of Cav 3.1 in the dmVMH was sufficient to rescue stress-induced anxiety-like behavior.

For energy metabolic tests, we fasted all mice overnight and analyzed energy

expenditure using CLAMS. Data showed that Cav 3.1 silencing effectively rescued the

chronic stress-induced decrease in RER (Fig. 5f), as well as the reduction in food intake

(Fig. 5f) and energy expenditure (Fig. 5a and h). No obvious body weight differences

were observed among the control, stress, and RNAi groups (Fig. 5e).

Together with our optogenetic manipulation experiments, our results consistently

demonstrated that Cav 3.1 in the dmVMH was both sufficient and necessary to elicit

burst firing of dmVMH neurons. Furthermore, knockdown of Cav 3.1 in the dmVMH

ameliorated chronic stress-induced phenotypic abnormalities, including anxiety-like

behavior, lower RER, and decreased food intake.

**Discussion and conclusion**

Previous studies have indicated that the dmVMH is an important stress coping center to

balance aversive behavior and energy seeking [2] and may be a possible hub connecting

stress-induced emotion and energy expenditure. Here, we reported that chronic stress

induced burst firing enhancement in the dmVMH, which was critical for inducing anxiety-like behavior and energy expenditure alteration in stressed mice. Bursting is an important firing pattern in the central nervous system and is also important for specific information transmission. Based on our data, we determined that dmVMH burst firing neurons play an important role in connecting the emotional state of anxiety and energy homeostasis, and that T-VGCC Cav 3.1 is essential for dmVMH burst firing and contributes to chronic stress-induced behavior and energy expenditure changes (Figure S6).

**dmVMH is responsive in chronic stress**

Chronic unpredictable stress is detrimental to both mental and physical health. Responses to chronic stress can vary from emotional disorders, like anxiety and depression, in the central nervous system to disturbed metabolism and energy homeostasis in peripheral organs [3-5]. However, how these physiological processes integrate in the brain and the underlying neural mechanisms remain poorly understood. The dmVMH is implicated in integrating information in the limbic system and in maintaining energy balance [38]. Furthermore, SF-1 neurons in the dmVMH constitute a nutritionally sensitive switch, which modulates the competing motivations of feeding and avoidance of environments full of acute stress [2]. Given the important role of the dmVMH in balancing acute stress inputs and food-seeking, the effects of persistent unpredictable stress on the function of dmVMH neurons require further research.

In the current study, the dmVMH was found to be involved in integrating chronic stress inputs to regulate anxiety-like behavior and energy expenditure. We established an unpredictable and persistent stress mouse model and characterized the model with multiple energy metabolic, electrophysiological, and behavioral approaches. Consistent with previous reports [3], we found that unpredictable persistent stress induced aversive behavior, decreased food intake, and shifted the RER toward fat oxidation, without obvious effects on body weight. Furthermore, *in vivo* electrophysiology demonstrated that the power density of the theta band of local field potential was higher in the stress group than in the control group. To dissect the alteration in dmVMH neuronal activity under chronic stress conditions, we studied the electrophysiology of dmVMH neurons. Based on differences in electrical activity, we classified the dmVMH neurons into three subtypes: i.e., silent, tonic-firing, and bursting. By comparing these subtypes in the control and stress groups, we found an enhancement of burst firing in the dmVMH after chronic stress, including increased percentage of burst firing neurons, enhanced LFP, and decreased ISI of bursts. Using patch clamp recordings, our data demonstrated that chronic stressors indeed induced obvious electrophysiological changes in dmVMH neurons at the cellular level, further supporting the important role of the dmVMH in chronic stress.

**Burst firing of dmVMH is critical for chronic stress response**

Action potentials that arrive in bursts provide more precise information than action potentials that arrive singly and further enhance the efficacy of neuromodulator release, thus revealing the special role of bursts in information transmission and processing [30]. Many previous studies have indicated that burst firing of specific neuronal subpopulations is critical for performing specific functions. In the hippocampus, a single burst can produce long-term synaptic modifications [39]. In the hypothalamus, burst firing is found in several regions, including the preoptic area and VMH [40]. In the lateral habenula, bursting activity depends on the NMDA receptor and T-VGCC; and most importantly, can drive behavioral aversion and depression-like symptoms [26]. To date, however, no specific study has reported on the functions of dmVMH burst firing under chronic stress. Here, we performed electrophysiological and gene expression studies and demonstrated that burst firing neurons in the dmVMH are actively involved in chronic stress response, and the ISI of burst firing was significantly correlated with the anxiety-like behavior in anxious mice after chronic stress.

To mimic burst firing *in vivo*, we applied optogenetic inhibition with an NpHR-expressing AAV vector, as optogenetic inhibition on the lateral habenula or hippocampal NpHR-expressing neurons can elicit burst firing in yellow light illumination intervals [26,41]. However, burst firing in the dmVMH could not be elicited under an illumination frequency greater than 0.2 Hz in our electrophysiological studies, indicating regional specificity of T-VGCC activation kinetics. Importantly, we

applied wireless optogenetics to successfully achieve burst manipulation in mice

residing within a closed cage, followed by behavioral and energy metabolic analyses.

Our data indicated that optogenetic-elicited burst firing was sufficient to promote the

chronic stress-induced phenotypes described above. Taken together, these data show

that VMH burst firing is an important hub for integrating and interacting anxiety-like

emotional state and energy expenditure regulation.

**Mechanism of burst firing in dmVMH**

The mechanism underlying neuronal burst firing is complicated and could explain

specific functions in different brain regions. Previous studies have reported that

various modulatory substances can promote depolarization of membranes and evoke

burst firing of neurons. Oxytocin binds to its receptor in CA2 region and can activate

the G protein-coupled pathway, and thus induce activation of several spike channels

and closure of KCNQ potassium channels [39]. Among the various mechanisms

underlying burst initiation, activation of a low-threshold ion channel can induce burst

firing in diverse brain regions [26,31]. Typically, low-threshold T-VGCC is widely

expressed in the central nervous system and can be activated by stimuli near the RMP

to elicit burst firing. Molecular cloning has revealed three isoforms of T-type channel

genes: i.e., Cav3.1, Cav3.2, and Cav3.3, which make distinct contributions to cellular

electrical properties [31,32,42]. Selective expression of Cav3.1 is sufficient to generate a

strong rebound burst in deep cerebellar nuclear neurons, whereas expression of

Cav3.2 or Cav3.3 alone does not generate a rebound discharge under normal

conditions [31]. Several studies have demonstrated that substitution of Cav3.1 or Cav3.2

for the native channel in model thalamic relay neurons causes elimination of

high-frequency bursts, further confirming the role of Cav3.1 in burst firing [32,42].

Here, we identified the existence of Cav3.1, Cav3.2, and Cav3.3 in the dmVMH

region. Cav3.1 and Cav3.2 displayed high expression in the dmVMH, whereas Cav3.3

showed lower expression, consistent with previous study [40]. The T-VGCC antagonist

mibefradil blocked burst firing of dmVMH neurons, thus suggesting the existence of

T-VGCC-mediated burst firing in the dmVMH. Furthermore, combined analysis of

electrophysiological classification and single-cell qRT-PCR, we found that Cav3.1

expression increased under chronic stress, which contributed to the enhancement of

burst firing in the dmVMH. Furthermore, Cav3.1 demonstrated greater contribution to

stress-induced enhancement of bursting in the dmVMH than the other isoforms.

We further investigated the necessity of Cav3.1 in chronic stress-induced anxiety-like

behavior and energy metabolic disorders. We found that microinjection of mibefradil

ameliorated anxiety-like behavior. Using RNAi methods, the expression of Cav3.1 in

the dmVMH was specifically knocked down, resulting in the significant inhibition of

neuronal bursting activity. Behavioral and energy expenditure experiments further

demonstrated that the chronic stress responses described above were partially rescued.

Thus, our findings consistently demonstrated that Cav3.1 may be a potential drug

target for the treatment of anxiety and related energy metabolic disorders.

The current study has several limitations. Firstly, we focused on exploring neuronal

activity changes in the dmVMH after chronic stress. As such, further research should

focus on identifying the mechanism regulating Cav3.1 expression in the dmVMH,

especially the receptors regulating Cav3.1 expression. Several factors might contribute

to the increased percentage of burst firing neurons after stress, including stress-induced

hormones or feeding-related peptides [24]. Previous studies have reported that estrogen

can regulate the expression and function of T-VGCC in vlVMH neurons, but not

dmVMH neurons, through estrogen receptors [40]. However, the mechanism underlying

Cav3.1 expression influenced by hormones or neuropeptides after chronic stress needs

further study.

Secondly, GABAergic neural circuits and astrocytes have been found in the VMH and

are implicated in regulating energy expenditure [43,44], but their specific contributions to

bursting activity need to be further investigated. We demonstrated that glutamate

receptors (especially NMDA receptors) affect the generation of burst firing; however,

whether synaptic glutamate uptake mediated by astrocytes affects burst firing requires

further exploration. Moreover, as the function of the T-VGCC is highly membrane

potential dependent, how upstream inputs integrate with the surrounding

microenvironment to affect membrane potential, and hence regulate burst firing,

needs to be further explored.

In summary, our study first identified bursting firing neurons in the dmVMH as a hub

regulating emotional state and energy metabolic disorders. We also identified Cav 3.1

as the crucial regulator of bursting firing of dmVMH neurons. The results of this

molecular and electrophysiological study should provide a more complete

understanding of the chronic stress-induced emotional malfunction and peripheral

metabolism disorders, and provides potential therapeutic targets for treating such

malfunctions.

## Competing interests

452 The authors declare no competing interests.

## Acknowledgements
This project was partly supported by the National Natural Science Foundation of

China (81471164, 31800881), Key Research Program of Frontier Sciences of Chinese

Academy of Sciences (QYZDB-SSW-SMC056), and Shenzhen Governmental Basic

Research Grant (JCYJ20170413164535041, JCYJ20180507182301299). We also

thank Z.B Xu and B.F Liu for their help in transgenic mice husbandry and

phenotyping. We are grateful to N.N Li and X.L Liu for the help in virus packaging.

## Author contributions
F.Y. and H.-L.H. conceived the idea and designed the experiments. J.S. D.-S.G and

Y.-H.L performed all electrophysiological recordings, immunostaining, qRT-PCR and

behavioral experiments, S.-P.C. and X.-Y.Z. helped with patch clamp recording and

qRT-PCR, L.Z. helped with immunostaining and surgery,  N.L. helped with the

RNAi construction, Q.X. helped with in vivo electrophysiological recording, F.Y. and

J.S. interpreted the results and wrote the manuscript with critical inputs from H.-L.H.

and L.-P.W..

**Methods**

**Animals**

All procedures were carried out in accordance with protocols approved by the Ethics

Committee of the Shenzhen Institutes of Advanced Technology, Chinese Academy of

Sciences. Male C57BL/6 mice (4–8 weeks old) were purchased from the Guangdong

Medical Laboratory Animal Center (Guangzhou, China). The SF-1-Cre mice (stock no:

012462) were obtained from Jackson Laboratories (Bar Harbor, ME, USA). Mice were

housed at 22–25 °C on a circadian cycle of 12-hour light and 12-hour dark with

ad-libitum access to food and water.

**Chronic stress procedures**

All animals used in this study were male, and randomly assigned to either control or

stress groups for chronic stress study. The stress group was daily subjected to one

stressor which randomly chosen from following: (i) 10 mice squeezing in a relatively

small cage (15 cm×10 cm×4 cm) for two hours. (ii) wet bedding in home cage

overnight (200 ml water was added to moisten bedding). (iii) each mouse was tightly

restraint in a tube for two hours. The chronic stress protocol lasted for 28 consecutive

days. Mice in different stress groups received the same number of each stressor.

Control animals were subjected to no stressors.

**Behavioral tests**

*Elevated plus maze (EPM).* Exposure (5-min) to EPM was used to assess locomotor

activity and anxiety-related behaviors after chronic stress. The stress and control

groups (n = 18–20/group) were placed in the center of the plus maze facing an open arm

and behavior was recorded for the entire 5 min using an overhead-mounted camera.

Videos recorded during the EPM test were analyzed with ANY-maze software

(Stoelting Co., Wood Dale, USA) to acquire data on time spent in open and closed arms,

locomotor activity (total distance travelled in maze), and entries into the open arm.

Anxiety-related behavior is associated with less exploration in the open arm relative to

overall exploration of all arms.

*Open field test (OFT).* Open field exposure (10-min) was used to assess locomotor

activity and anxiety-related behaviors after chronic stress. Mice (n = 18–20/group)

were placed in the center of an open field and behavior was recorded for the entire 10

min. Videos were analyzed to acquire data on time spent in the center and corner areas,

total locomotor activity, and number of entries into the center area. Anxiety-related

behavior is associated with less exploration of the center area.

**CLAMS and energy expenditure**

To characterize metabolic changes caused by chronic stress or Cav 3.1 knockdown, RER was measured by indirect calorimetry using a four-chamber open-circuit calorimeter (Oxymax Series; Columbus Instruments, Columbus, OH, USA). Mice were food deprived overnight, with body weight and chow in each cage weighed prior to the experiment. During the experiment, mice were housed individually in specially built Plexiglas cages (40 × 25 × 20 cm). Temperature was maintained at 22 °C with an airflow of 0.5 per min. Food and water were available *ad libitum*. Mice were subsequently monitored in the system for 24 h (whole light-dark cycle). Oxygen consumption (VO2) and carbon dioxide production (VCO2) were measured every 10 min. The RER was calculated as a quotient of VCO2/VO2, with 1 representing 100% carbohydrate oxidation and 0.7 representing 100% fat oxidation. Energy expenditure (kcal heat produced) was calculated as calorific value (CV) × VO2, where CV is 3.815 + 1.232 × RER [3]. Metabolic data collected from the 24-h monitoring period were averaged for energy expenditure and RER. After the experiment, chow in each cage was weighed to calculate food intake.

**Slice preparation**

Mice were deeply anesthetized with isoflurane and decapitated rapidly. Brains were then removed and transferred to chilled cutting solution within 3 min. Cutting solution contained (in mM): choline chloride 110; KCl 2.5; Na-pyruvate 0.6; MgCl$_2$ 7.0; CaCl$_2$

0.5; $NaH_2PO_4$ 1.3; $NaHCO_3$ 25; glucose 20 (pH 7.4). The chilled cutting solution was

bubbled with carbogen (95% $O_2$ and 5% $CO_2$) for at least 30 min before use. Coronal

slices (250–300 mm thick) were prepared on a vibratome (Series 1000, Warner

Instruments, Berlin, Germany), and incubated in artificial cerebrospinal fluid (ACSF)

containing (in mM): NaCl 125; KCl 2.5; Na-pyruvate 0.6; $MgCl_2$ 1.3; $CaCl_2$ 2.0;

$NaH_2PO_4$ 1.3; $NaHCO_3$ 25; glucose 10 (pH 7.4) bubbled with carbogen at 34 °C for 30

min. After incubation, all slices were equilibrated in ACSF at room temperature

(24–26 °C) for at least 40 min. Single slices were placed on the recording chamber

perfused with ACSF bubbled with carbogen at room temperature. Unless stated

otherwise, drugs were applied with perfused ACSF.

**Electrophysiology**

Whole-cell patch clamp recording was performed on VMH neurons. The dmVMH was

identified based on landmarks (third ventricle). Recordings were obtained with

multi-clamp 700B amplifiers (Molecular Devices, San Jose, USA) under visual

guidance using a Nikon FN1 microscope (Tokyo, Japan). Electrophysiological data

were acquired and analyzed using pClamp 10 software (Molecular Devices, San Jose,

USA ). Whole-cell recordings were performed with borosilicate glass electrodes (0.69

mm OD, 5–7 MΩ) with internal solution containing (in mM): K-gluconate 135.0; KCl

4.0; NaCl 2.0; HEPES 10; EGTA 4.0; Mg-ATP 4.0; Na-GTP 5.0. Osmolality was

adjusted to 290–310 mOsm $kg^{-1}$ with sucrose and pH was adjusted to 7.4 with KOH.

After forming a high-resistance seal (GΩ), the cell was held in current-clamp mode for

7–10 min until access resistance stabilized. Resting membrane potential (RMP) was

assessed at the beginning of the recording period after stabilization of access resistance,

and periodically monitored throughout the recording by momentarily relieving the

direct current injection. To elucidate differences among neurons, 800-ms current

injections (-100 to 100 pA in 10 pA increments; 5 s interstimulus interval) were applied,

and the number of action potentials evoked by each current injection, input resistance,

and half-width of action potentials were determined.

**Drugs and reagents**

All chemicals included in ACSF prescription were purchased from Sigma-Aldrich

(Merck KGaA, Darmstadt, Germany). Mibefradil (40 μM) were purchased from Tocris

(Bio-Techne, Minneapolis, USA) and used for electrophysiology and microinjection.

NBQX (1,2,3,4-tetrahydro-6-nitro-2,3-dioxo- benzo[f]quinoxaline-7-sulfonamide, 30

μM) and AP5 (2-amino-5-phosphono-pentanoic acid, 30 μM) were acquired from Med

Chem Express (MCE, Shanghai, China).

**Cell harvesting and single-cell Quantitative real-time polymerase chain reaction**

**(qRT-PCR)**

Single Cell-to-CT™ Kits (Thermo Fisher, USA) were used to perform single-cell

qRT-PCR. After recording, single cells were collected and lysed to acquire total RNA.

A Mastercycler 5333 PCR thermal cycler (Eppendorf, Germany) was used to perform

reverse transcription and pre-amplification. TaqMan gene expression assay was used to

assay each gene. Gene expression levels were normalized to the expression of

housekeeping gene *Rn18s*. All protocols were performed according to the

manufacturer's instructions. qRT-PCR was performed with a Light cycler 480 (Roche,

Switzerland).

**Tissue collection and qRT-PCR**

To acquire dmVMH tissue, mice were first anesthetized with isoflurane. After bathing

in cold 1% diethyl pyrocarbonate (DEPC) phosphate-buffered saline (PBS) solution,

brains were acute cut on a vibratome. VMH tissues were dissected microscopically

from those sections, and quickly transferred to TRIzol reagent. The manufacturer's

standard protocols for RNA extraction (TransGene, China) and synthesis (TOYOBO,

Japan) were followed. Primers used for qRT-PCR included: Cav 3.1 (5'-TGG

CCTTCTTCGTCCTGAAC-3' and 5'-TTCTCCAGCCTCTTTAGTCGC-3'). Cav 3.2

(5'-CGGCCCTACT ACGCAGACTA-3' and 5'-TTAAGGGCCTCGTCCAGAGA-3'),

Cav 3.3 (5'-CTGCTATTCTCCAGCCCAGG-3' and 5'-AGCTGCACCTCTTG

CTTGT-3'). Expression of these gene was normalized to the expression of

housekeeping gene β-Actin (*Act-b*).

*In vivo* **electrophysiology**

Mice were implanted with two nickel-chromium wires (25-μm diameter; AM Systems,

Sequim, USA) targeting the dmVMH, connector was bind to wires and fixed on skull

with dental cement. After total recovery from surgery, mice were returned to the

recording sessions. Neuronal activity was collected using a Plexon Multichannel

Acquisition Processor system (Plexon, Dallas, USA). Local field potentials (digitized

at 1 kHz sampling rate, low-pass filtered up to 250 Hz) were recorded simultaneously

for 90 min with a gain of 5 000×. After the recording sessions, mice were anesthetized

by pentobarbital sodium and perfused intracardially. The electrode recording position

was marked by histological staining.

**Stereotaxic surgery and viral injection**

For all stereotaxic surgery, 12–16-week-old mice were anesthetized by pentobarbital

sodium (0.3% in saline, 1 ml/100 g, intraperitoneally) and placed in a stereotaxic

apparatus for surgery. Stereotaxic surgical procedures were performed using standard

protocols. To target the dmVMH, bilateral brain injection coordinates relative to

bregma were chosen (AP, -1.58 mm; ML, ±0.3 mm; DV, -5.5 mm). Unless stated

otherwise, 0.4 µL of viral vector was injected into the VMH at a rate of 0.1 µL/min

using a 10-µL Hamilton syringe and a syringe infusion pump (World Precision

Instruments, USA).

For RNAi knockdown study, Cav3.1-shRNA (U6-shRNA-EF1a-copGFP-PGK-Puro)

was injected into the VMH with lenti-viral vector. Mice were housed for four weeks

following injection for viral expression. Anxiety-like behavior and metabolic tests were

performed at the end of the paradigm. To assess the knockdown effectiveness of Cav

3.1 shRNA, mice were perfused with 4% paraformaldehyde (PFA) and brain tissues

were removed for immunostaining analysis after the final session.

For microinjection, a stainless-steel guide-cannula (0.6 mm outer diameter and 0.4 mm

inner diameter, RWD, Shenzhen, China) was implanted into the diencephalon to target

the VMH (AP, -1.58 mm; ML, ±0.3 mm; DV, -5.0 mm). The guide-cannula was fixed to

the skull using dental cement and three stainless steel screws. Each guide-cannula was

sealed with a stainless-steel wire for protection from obstruction. Before behavioral

testing, the stainless-steel wire was replaced with an injection cannula (0.5 mm deeper

than guide-cannula to break astrocyte aggregation), through which the drug dissolved

in saline (400 nl) was delivered at 100 nl/min into the VMH. After microinjection, mice

were rested for 30 min before tests.

**Wireless optogenetic manipulation**

To achieve wireless optogenetic manipulation of burst firing neurons, AAV9-DIO

-NpHR-mCherry was injected unilaterally into the VMH in the left hemisphere. Mice

were housed for four weeks following injection for viral expression before initiation of

experiments. A copper coil with a light emitting diode (LED) on the right side of the tip

(Inper, Hangzhou, China) was implanted into the brain of NpHR-expressing mice, with

the LED located 0.2 μm left of the dmVMH. The coil and indicator were fixed on the

surface of the skull bone with Vetbond Tissue Adhesive (3M, USA). Charging the coil

was achieved through antennas surrounding the cage. To initiate burst firing of

dmVMH neurons, 550-nm yellow light stimulation was performed at 0.1 Hz (2 s/pulse)

in the NpHR-mCherry group during the metabolic (one illumination trial/12 h, each

trial lasted for 2 h) and behavioral tests (OFT, 10 min; EPM, 5 min).

**Immunohistochemistry**

Mice were first anesthetized by chloral hydrate (10% in saline, 1 ml/100 g,

intraperitoneally), then perfused with 0.01 M PBS and 50 mL of 4% PFA transcardially.

Brains were dissected and post-fixed in 4% PFA overnight and then transferred to 30%

w/v sucrose solution for cryoprotection until sinking. Sections from the entire

anterior-posterior range of the VMH were stained using an antibody specific to Cav3.1

(1: 100, Alomone labs, Cat# ACC-021, RRID: AB_2039779), Cav3.2 (1: 100,

Alomone labs, Cat#ACC-025, RRID: AB_2039781), or Cav 3.3(1: 100, Alomone labs,

Cat# ACC-009, RRID: AB_2039783). Briefly, sections were washed, permeabilized in

0.1% Triton X-100/PBS for 15 min/three times, washed again, and blocked in 10%

normal goat serum (NGS) (w/v)/0.1% Triton X-100/PBS for 30 min. Primary antibody

was added, and sections were incubated overnight at 4 ℃. The following day, sections

were washed with 0.01 mM PBS, incubated with incubated with secondary antibody

(Alexa Fluor 594 Goat Anti-Rabbit, Cat#115-585-003, RRID: AB_2338059 or Alexa

Fluor 488 Goat Anti-Rabbit, Cat#111-547-003 RRID: AB_2338058) in 1% NGS/0.1%

Triton X-100/PBS for 1 h at room temperature, then washed, mounted, and cover

slipped with mounting medium containing DAPI.

**Statistical analysis**

Slice electrophysiological data were analyzed with pCLAMP (Molecular Device, San

Jose, CA, USA). *In vivo* electrophysiology data were analyzed with NeuroExplorer 5.0

(Plexon, Dallas, USA). All data were imported into Prism 7 (GraphPad Software, La

Jolla, CA, USA) and normality was assessed using D'Agostino-Pearson tests to verify

the appropriateness of the following statistical analyses. Unless stated otherwise, the

data are presented as means ± SEM. Statistical significance was determined using

two-tailed unpaired Student's *t*-tests when comparing two groups, paired Student's

*t*-test when comparing the effects of different treatments in the same group. When

multiple measures were compared between groups (e.g., current-frequency curves),

repeated measures two-way analysis of variance (ANOVA) with Bonferroni's test was

used. Differences were considered significant at $P < 0.05$.

To classify dmVMH neurons, unsupervised cluster analysis was performed with SPSS

v19 (Chicago, IL, USA) using squared Euclidean distances. The parameters for cluster

analysis were chosen based on their lack of linear correlation with each other. The

following electrophysiological parameters were included for analysis: onset time,

evoked firing rate, input resistance, and potential sag by H-current

(Hyperpolarization-activated current). Resting potential were excluded from the

parameters for cluster analysis because of their linear correlation with onset times. All

electrophysiological parameters were converted into standardized z-scores before

clustering.

## References

**References**

1    Koolhaas, J. M. *et al.* Coping styles in animals: current status in behavior and
stress-physiology. *Neuroscience & Biobehavioral Reviews* **23**, 925-935,
doi:https://doi.org/10.1016/S0149-7634(99)00026-3 (1999).

2    Viskaitis, P. *et al.* Modulation of SF1 Neuron Activity Coordinately Regulates
Both Feeding Behavior and Associated Emotional States. *Cell Reports* **21**,
3559-3572, doi:https://doi.org/10.1016/j.celrep.2017.11.089 (2017).

3    Hardaway, J. A. *et al.* Central Amygdala Prepronociceptin-Expressing Neurons
Mediate Palatable Food Consumption and Reward. *Neuron* **102**, 1088,
doi:https://doi.org/10.1016/j.neuron.2019.04.036 (2019).

4    Gross, C. & Hen, R. The developmental origins of anxiety. *Nature Reviews*
*Neuroscience* **5**, 545-552, doi:10.1038/nrn1429 (2004).

5    Lupien, S. J., McEwen, B. S., Gunnar, M. R. & Heim, C. Effects of stress
throughout the lifespan on the brain, behaviour and cognition. *Nature*
*Reviews Neuroscience* **10**, 434-445, doi:10.1038/nrn2639 (2009).

6    Furay, A. R., Bruestle, A. E. & Herman, J. P. The role of the forebrain
glucocorticoid receptor in acute and chronic stress. *Endocrinology* **149**,
5482-5490, doi:10.1210/en.2008-0642 (2008).

7    Isingrini, E. *et al.* Resilience to chronic stress is mediated by noradrenergic
regulation of dopamine neurons. *Nature Neuroscience* **19**, 560-563,
doi:10.1038/nn.4245 (2016).

8    Kuperman, Y. *et al.* Perifornical Urocortin-3 mediates the link between
stress-induced anxiety and energy homeostasis. *Proceedings of the National*
*Academy of Sciences of the United States of America* **107**, 8393-8398,
doi:10.1073/pnas.1003969107 (2010).

9    Cheung, C. C. *et al.* Sex-dependent changes in metabolism and behavior, as
well as reduced anxiety after eliminating ventromedial hypothalamus
excitatory output. *Molecular metabolism* **4**, 857-866,
doi:10.1016/j.molmet.2015.09.001 (2015).

10   Guo, M. *et al.* Role of the adipose PPARγ-adiponectin axis in susceptibility to
stress and depression/anxiety-related behaviors. *Molecular psychiatry* **22**,
1056-1068, doi:10.1038/mp.2016.225 (2017).

11   Grayson, B. E., Seeley, R. J. & Sandoval, D. A. Wired on sugar: the role of the
CNS in the regulation of glucose homeostasis. *Nature reviews. Neuroscience*
**14**, 24-37, doi:10.1038/nrn3409 (2013).

12   Silva, B. A. *et al.* Independent hypothalamic circuits for social and predator
fear. *Nature Neuroscience* **16**, 1731-1733, doi:10.1038/nn.3573 (2013).

13   Wang, L., Chen, I. Z. & Lin, D. Collateral pathways from the ventromedial
hypothalamus mediate defensive behaviors. *Neuron* **85**, 1344-1358,
doi:10.1016/j.neuron.2014.12.025 (2015).

14   Coutinho, E. A. *et al.* Activation of SF1 Neurons in the Ventromedial

Hypothalamus by DREADD Technology Increases Insulin Sensitivity in
Peripheral Tissues. *Diabetes* **66**, 2372-2386, doi:10.2337/db16-1344 (2017).

15   Yadav, V. K. *et al.* A serotonin-dependent mechanism explains the leptin
regulation of bone mass, appetite, and energy expenditure. *Cell* **138**, 976-989,
doi:10.1016/j.cell.2009.06.051 (2009).

16   McClellan, K. M., Parker, K. L. & Tobet, S. Development of the ventromedial
nucleus of the hypothalamus. *Frontiers in Neuroendocrinology* **27**, 193-209,
doi:https://doi.org/10.1016/j.yfrne.2006.02.002 (2006).

17   Anderson, D. J. Circuit modules linking internal states and social behaviour in
flies and mice. *Nature Reviews Neuroscience* **17**, 692,
doi:10.1038/nrn.2016.125 (2016).

18   Hashikawa, K. *et al.* Esr1+ cells in the ventromedial hypothalamus control
female aggression. *Nat Neurosci* **20**, 1580-1590, doi:10.1038/nn.4644 (2017).

19   Kim, D. W. *et al.* Multimodal Analysis of Cell Types in a Hypothalamic Node
Controlling Social Behavior. *Cell* **179**, 713-728.e717,
doi:10.1016/j.cell.2019.09.020 (2019).

20   Wang, C., Bomberg, E., Billington, C. J., Levine, A. S. & Kotz, C. M.
Brain-derived neurotrophic factor (BDNF) in the hypothalamic ventromedial
nucleus increases energy expenditure. *Brain research* **1336**, 66-77,
doi:https://doi.org/10.1016/j.brainres.2010.04.013 (2010).

21   Klöckener, T. *et al.* High-fat feeding promotes obesity via insulin
receptor/PI3K-dependent inhibition of SF-1 VMH neurons. *Nature*
*Neuroscience* **14**, 911-918, doi:10.1038/nn.2847 (2011).

22   Kunwar, P. S. *et al.* Ventromedial hypothalamic neurons control a defensive
emotion state. *eLife* **4**, doi:10.7554/eLife.06633 (2015).

23   Iigaya, K., Okazaki, S., Minoura, Y. & Onimaru, H. Interaction between novel
oscillation within the ventromedial hypothalamus and the sympathetic
nervous system. *Neuroscience* **343**, 213-221,
doi:10.1016/j.neuroscience.2016.11.048 (2017).

24   Iigaya, K., Minoura, Y., Onimaru, H., Kotani, S. & Izumizaki, M. Effects of
Feeding-Related Peptides on Neuronal Oscillation in the Ventromedial
Hypothalamus. *J Clin Med* **8**, 292, doi:10.3390/jcm8030292 (2019).

25   Wang, G. *et al.* CaV3.2 calcium channels control NMDA receptor-mediated
transmission: a new mechanism for absence epilepsy. *Genes & development*
**29**, 1535-1551, doi:10.1101/gad.260869.115 (2015).

26   Cain, S. M. *et al.* CaV3.2 drives sustained burst-firing, which is critical for
absence seizure propagation in reticular thalamic neurons. *Epilepsia* **59**,
778-791, doi:10.1111/epi.14018 (2018).

27   Yuan, Y. *et al.* Reward Inhibits Paraventricular CRH Neurons to Relieve Stress.
*Current Biology* **29**, 1243-1251.e1244, doi:10.1016/j.cub.2019.02.048 (2019).

28   Yang, Y. *et al.* Ketamine blocks bursting in the lateral habenula to rapidly
relieve depression. *Nature* **554**, 317-322, doi:10.1038/nature25509 (2018).

29 Lee, S., Lee, C., Woo, C., Kang, S. J. & Shin, K. S. Chronic social defeat
stress-induced enhancement of T-type calcium channels increases burst-firing
neurons in the ventral subiculum. *Biochemical and biophysical research*
*communications* **508**, 1182-1187, doi:10.1016/j.bbrc.2018.12.073 (2019).

30 Lisman, J. E. Bursts as a unit of neural information: making unreliable
synapses reliable. *Trends Neurosci* **20**, 38-43,
doi:10.1016/s0166-2236(96)10070-9 (1997).

31 Molineux, M. L. *et al.* Specific T-type calcium channel isoforms are associated
with distinct burst phenotypes in deep cerebellar nuclear neurons.
*Proceedings of the National Academy of Sciences of the United States of*
*America* **103**, 5555-5560, doi:10.1073/pnas.0601261103 (2006).

32 McRory, J. E. *et al.* Molecular and functional characterization of a family of rat
brain T-type calcium channels. *The Journal of biological chemistry* **276**,
3999-4011, doi:10.1074/jbc.M008215200 (2001).

33 Nelson, M. T. *et al.* Molecular mechanisms of subtype-specific inhibition of
neuronal T-type calcium channels by ascorbate. *The Journal of neuroscience :*
*the official journal of the Society for Neuroscience* **27**, 12577-12583,
doi:10.1523/jneurosci.2206-07.2007 (2007).

34 Sherman, S. M. Tonic and burst firing: dual modes of thalamocortical relay.
*Trends in Neurosciences* **24**, 0-126 (2001).

35 Cui, Y. *et al.* Astroglial Kir4.1 in the lateral habenula drives neuronal bursts in
depression. *Nature* **554**, 323-327, doi:10.1038/nature25752 (2018).

36 Cain, S. M. *et al.* GABAB receptors suppress burst-firing in reticular thalamic
neurons. *Channels (Austin, Tex.)* **11**, 574-586,
doi:10.1080/19336950.2017.1358836 (2017).

37 Matott, M. P., Kline, D. D. & Hasser, E. M. Glial EAAT2 regulation of
extracellular nTS glutamate critically controls neuronal activity and
cardiorespiratory reflexes. *The Journal of physiology* **595**, 6045-6063,
doi:10.1113/jp274620 (2017).

38 Steffens, A. B., Scheurink, A. J. W., Luiten, P. G. M. & Bohus, B. Hypothalamic
food intake regulating areas are involved in the homeostasis of blood glucose
and plasma FFA levels. *Physiology & Behavior* **44**, 581-589,
doi:https://doi.org/10.1016/0031-9384(88)90322-8 (1988).

39 Tirko, N. N. *et al.* Oxytocin Transforms Firing Mode of CA2 Hippocampal
Neurons. *Neuron* **100**, 593-608.e593, doi:10.1016/j.neuron.2018.09.008
(2018).

40 Qiu, J. *et al.* Estrogen upregulates T-type calcium channels in the
hypothalamus and pituitary. *The Journal of neuroscience : the official journal*
*of the Society for Neuroscience* **26**, 11072-11082,
doi:10.1523/jneurosci.3229-06.2006 (2006).

41 Berglind, F. *et al.* Optogenetic inhibition of chemically induced
hypersynchronized bursting in mice. *Neurobiology of Disease* **65**, 133-141,

doi:https://doi.org/10.1016/j.nbd.2014.01.015 (2014).

42    Cain, S. M. & Snutch, T. P. Contributions of T-type calcium channel isoforms to
neuronal firing. *Channels* **4**, 475-482, doi:10.4161/chan.4.6.14106 (2010).

43    Bouyakdan, K. *et al.* The gliotransmitter ACBP controls feeding and energy
homeostasis via the melanocortin system. *The Journal of clinical investigation*
**129**, 2417-2430, doi:10.1172/jci123454 (2019).

44    Chan, O. *et al.* Increased GABAergic Output in the Ventromedial
Hypothalamus Contributes to Impaired Hypoglycemic Counterregulation in
Diabetic Rats. *Diabetes* **60**, 1582-1589, doi:10.2337/db10-1579 (2011).

**Fig. 1| Chronically stressed mice exhibited anxiety-like behavior, altered**

**metabolism, and *in vivo* electrophysiological dmVMH activity.**

**(a)** Illustration of unpredictable chronic stress protocol and phenotype assessment, with

one stressor treatment randomly chosen per day, lasting for four weeks.    **(b)** Residence

time in each site of open field (blue, less time; red, more time). **(c)** Behavioral analysis

of control (n = 33) and stress group mice (n = 38) in open field test showed significant

decrease in both time spent in central area (unpaired Student's *t*-test, *P* = 0.0035) and

entries into central area (unpaired Student's *t*-test, *P* = 0.0012), but no obvious change

in travelling distance (unpaired Student's *t*-test, *P* = 0.3029). **(d)** Residence time in each

site of elevated plus-maze (blue, less time; red, more time) of control and stress groups.

**(e)** Control (n = 32) and stress groups (n = 38) in elevated plus-maze showed significant

decrease in time spent in open arm (unpaired Student's *t*-test, *P* = 0.0194) and entries

into open arm (unpaired Student's *t*-test, *P* < 0.0001), but no obvious change in

locomotion (unpaired Student's *t*-test, *P* = 0.9302). **(f)** Body weight monitored after

chronic stress period showed no obvious change compared with naïve mice (n = 9 in

each group, unpaired Student's *t*-test, *P* = 0.7125), and 24-h food intake after overnight

fasting decreased in stress group (n = 9 in each group, unpaired Student's *t*-test, *P* =

0.0018) compared with control group (n = 9). **(g)** Average respiration exchange ratio

(RER) decreased in stress group compared with control group (unpaired student's *t*-test,

*P* = 0.0003, n = 9 mice in each group); RER curve shifted after chronic stress (two-way

ANOVA, *P* = 0.0002, *F*(1, 16) = 21.95). **(h)** Significant decreases in 24-h energy

expenditure (EE) curve and average EE were observed in stress group (two-way

ANOVA, *P* = 0.0057, *F*(1, 16) = 20.2; unpaired Student's *t*-test, *P* = 0.0060). **(i)**

Increased *c-fos* expression in dmVMH under chronic stress (unpaired Student's *t*-test, *P*

< 0.001, n = 5 mice in each group). **(j)** Schematic of electrode implantation and *in vivo*

electrophysiology recordings. **(k** and **l)** Power spectral density of local field potential

(LFP) in dmVMH after chronic stress, with significant power improvement observed in

theta band (two-way ANOVA, $P = 0.0298$, $F(1, 14) = 5.845$,; unpaired Student's $t$-test,

$P = 0.0243$, n = 4). Data are means ± SEM. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$.

CLAMS, comprehensive laboratory animal monitoring system.

**Fig. 2| Chronic stress-induced enhancement of burst firing in dmVMH neurons.**

**(a)** Increased average onset time (unpaired Student's *t*-test, *P* = 0.0233) and depolarized

average resting membrane potential (RMP) (unpaired Student's *t*-test, *P* = 0.0434) in 84

dmVMH neurons from stressed mice compared with 85 neurons from wild-type mice.

The box plotted at the median extending from the 25-75th percentile, and the whisker

represents Min to Max distribution. **(b)** Cluster analysis of 85 dmVMH neurons from

35 normal mice. Dendrogram of cluster analysis shows that dmVMH neurons could be

classified into three subtypes: i.e., silent, tonic-firing, and bursting. **(c)**

Electrophysiological properties of silent, tonic-firing, and bursting dmVMH neuronal

subtypes. left: whole-cell recording traces of three neuronal subtypes without current

injection; right: frequency-current curve of three subtypes at current injections of 0 to

100 pA and 10 pA/step. **(d)** Cluster analysis of 84 dmVMH neurons from 39 stressed

mice. Dendrogram of cluster analysis shows these dmVMH neurons can be classified

into three subtypes: i.e., silent (n = 13), tonic-firing (n = 35), and bursting (n = 38). See

also Figure S1 and S2. **(e)** Pie chart of percentages of neuronal dmVMH subtypes in

control group and stressed mice with obvious anxiety-like behavior (anxiety group). **(f)**

Distribution of 85 dmVMH neurons in control mice (left) and 57 dmVMH neurons in

anxiety group (right) using onset time-RMP coordinate system, with coordinate of

centroid (★) indicating average onset time and RMP. Blue, silent; Green, tonic-firing;

Red, bursting. Centroid coordinate was determined by average onset time and RMP.

Shift in centroid after chronic stress represents shorter average onset time and more

depolarized average RMP, caused by changes in proportion of three subtypes. **(g)**

Inter-spike interval (ISI) of bursts in dmVMH neurons of control and stressed mice.

Left, Example of burst firing and ISI; right, ISI of burst firing dmVMH neurons (n = 30)

in anxiety group decreased significantly compared with that in control group (n = 28,

unpaired Student's *t*-test, $P = 0.0099$). **(h)** ISI of dmVMH bursting neurons in control

group is slightly correlated with the residing time in open arms of EPM, but not with

the time spent in central area of open field (n = 28 cells from 21 mice). **(i)** ISI of

dmVMH bursting neurons in stressed mice which displayed obvious anxiety behavior

is significantly correlated with both the residing time in open arms and the time in

central area of open field (n = 30 cells from 20 mice). Data are means ± SEM * $P <$

0.05, ** $P < 0.01$.

**Fig. 3| Optogenetic activation of burst firing neurons in dmVMH induced anxiety-like behavior and energy expenditure changes.**

**(a)** Schematic of dmVMH injection of NpHR AAV viral vector to induce burst firing *in vivo*. **(b)** Whole-cell recordings of yellow light-evoked burst firing in brain slices (yellow light: 590 nm, left: 0.1 Hz, 2 s; middle: 0.2 Hz, 1 s; right: 1 Hz, 200 ms), 0.1 Hz successfully induced activation of burst firing neurons. **(c)** 0.1 Hz and 2 s yellow light illumination exerted no significant influence on silent or tonic-firing dmVMH neurons. **(d)** Illustration of wireless optogenetic manipulation of dmVMH neurons and

behavioral analysis in free-moving mice. **(e)** Open field test before, during, and after

light illumination: residence time in central area decreased during 10-min yellow light

illumination (n = 5, paired Student's $t$-test, $P = 0.0237$), though no obvious changes

were observed in number of entries into central area (paired Student's $t$-test, $P =$

0.5932). **(f)** Elevated plus-maze test before, during, and after 5-min light illumination.

Residence time in open arms decreased during yellow light illumination (n = 5, paired

Student's $t$-test, $P = 0.0144$) and recovered after light-off; no obvious changes were

observed in number of entries into open arms (paired Student's $t$-test, $P = 0.6974$). **(g)**

Food intake and metabolism were monitored during optogenetic manipulation of

dmVMH neurons in free-moving mice. Food intake decreased during light stimulation

(paired Student's $t$-test, $P = 0.0039$, n = 5 mice in each group), also average RER

(paired Student's $t$-test, $P = 0.0169$) and EE (paired Student's $t$-test, $P = 0.0274$). **(h and**

**i)** RER and EE curve shifted (two-way ANOVA, RER: $P = 0.0053$, $F(1, 8) = 14.38$; EE:

$P = 0.0227$, $F(1, 8) = 7.7918$ ) when applying two yellow light stimulation trials (0.1 Hz,

2 s; for 2 h). Data are means ± SEM; * $P < 0.05$, ** $P < 0.01$.

**Fig. 4| T-VGCC mediated enhancement of burst firing in dmVMH neurons under**

**chronic stress.**

**(a)** Schematic of structure of T-VGCC located on cell membrane (left top).

Representative immunofluorescence showing Cav 3.1 (left bottom), Cav 3.2 (right top),

and Cav 3.3 (right bottom) expression in dmVMH, respectively, with stronger Cav3.1

expression observed. Scale bar is 200 μm. **(b)** Quantification of Cav 3.1, Cav 3.2, and

Cav 3.3 expression in dmVMH tissue between control (n = 5 mice) and chronic stress

groups (n = 6 mice). Expression of Cav 3.1 was significantly up-regulated under

chronic stress conditions (unpaired Student's *t*-test, *P* = 0.0232). **(c)** Single-cell

qRT-PCR analysis of Cav 3.1, Cav 3.2, and Cav 3.3 expression in dmVMH neurons

between control (n = 16 cells) and chronic stress groups (n = 14 cells). Expression of

Cav 3.1 was significantly up-regulated under chronic stress conditions (unpaired

Student's $t$-test, $P = 0.0390$). **(d)** Evoked burst firing trace of dmVMH neurons without

and with T-VGCC antagonist (mibefradil, 10 μM), 10 pA current injection was given in

cosine waveform. **(e)** Effects of mibefradil on onset time and RMP of dmVMH neurons

from wild-type (n = 15) and chronic stress groups (n = 13). Significant differences were

observed in both onset time and RMP (paired Student's $t$-test, $P = 0.0388$ and $P =$

$0.0395$) in the stress group, whereas the control group demonstrated obvious changes in

RMP but not onset time (paired Student's $t$-test, $P = 0.0275$ and $P = 0.3125$). **(f)**

Single-cell qRT-PCR analysis of Cav 3.1 expression among three neuronal subtypes in

dmVMH. Upper: burst firing neurons (n = 11) showed higher Cav 3.1 expression than

other two subtypes (unpaired Student's $t$-test, $P = 0.0133$ compared with silent neurons

(n = 7), $P = 0.0139$ compared with tonic-firing neurons (n = 12)); bottom: Cav 3.1

expression in burst firing subtype showed significant differences between control and

chronic-stress groups (unpaired Student's $t$-test, silent: control, n = 3, stress, n = 3, $P =$

$0.5574$; tonic-firing: control, n = 7, stress, n = 5, $P = 0.9339$; bursting: control, n = 5,

stress, n = 6, $P = 0.0165$). **(g)** Effects of mibefradil on suprathreshold and subthreshold

activity in dmVMH burst firing neurons in control (n = 5) and chronic stress groups (n =

7). Mibefradil inhibited T-VGCC and caused a right frequency-current curve shift

(two-way ANOVA, control, $P = 0.2140$, $F(1, 8) = 1.854$; stress, $P = 0.0077$, $F(1, 12) =$

$10.22$); lower, mibefradil application exerted no obvious influence on current-voltage

curve of burst neurons (two-way ANOVA, control, $P = 0.8188$, $F(1, 10) = 0.0573$;

stress, $P = 0.4487$, $F(1, 10) = 0.6218$). **(h)** Obvious differences were observed in

suprathreshold activity (two-way ANOVA, $P = 0.0047$, $F(1, 110) = 20.53$), but not in

subthreshold (two-way ANOVA, $P = 0.9913$, $F(1, 33) = 4.958$) membrane potential of

dmVMH burst firing neurons between control (n = 5) and chronic stress groups (n = 7)

after application of mibefradil. Data are means ± SEM. * $P < 0.05$, ** $P < 0.01$.

**Fig. 5| Knockdown of Cav3.1 in dmVMH decreased burst firing, and rescued anxiety-like behavior and metabolic alteration induced by chronic stress.**

(a) Schematic of Cav3.1 shRNA construct and injection of shRNA-expressing lenti-viral vector into dmVMH to interfere with Cav 3.1 expression. (b) Representative images of dmVMH Cav 3.1 immunostaining in chronic stress and RNAi (under chronic stress) animals (left). Proportion of burst firing neurons decreased in RNAi group (2/13, 15.38%) compared with stress group (7/17, 41.18%) (right). Scale bar is 100 μm. (c)

Time spent in central area and number of entries into central area of open field in

control, chronic stress, and RNAi groups. Residence time: control (n = 8) versus

chronic stress group (n = 8), $P = 0.0068$; chronic stress versus RNAi group (n = 7), $P =$

0.0017. Number of entries: control versus chronic stress group, $P = 0.4830$; chronic

stress versus RNAi group, $P = 0.8437$ (unpaired Student's $t$-test). **(d)** Time spent in

open arm and number of entries into open arm of elevated plus-maze in control, chronic

stress, and RNAi groups. Residence time: control versus chronic stress group, $P =$

0.0009; chronic stress versus RNAi group, $P = 0.0379$. Number of entries: control

versus chronic stress group, $P < 0.0001$; chronic stress versus RNAi group, $P = 0.0607$

(unpaired Student's $t$-test, 8 mice in each group). **(e)** No significant differences in

average body weights of control (n = 8), chronic stress (n = 8) and RNAi (n = 8) groups

were observed after four weeks of chronic stress (unpaired Student's $t$-test, control

versus stress group, $P = 0.5857$; chronic stress versus RNAi group, $P = 0.6937$). **(f)**

Food intake of mice in control, chronic stress, and RNAi groups: control versus chronic

stress group, $P = 0.0022$; chronic stress versus RNAi group, $P < 0.0001$ (unpaired

Student's $t$-test, 8 mice in each group). Average RER of control, chronic stress, and

RNAi groups (unpaired Student's $t$-test: control versus chronic stress group, $P = 0.0095$;

chronic stress versus RNAi group, $P = 0.0032$). Average EE of control, chronic stress,

and RNAi groups (unpaired Student's $t$-test: control versus chronic stress group, $P =$

0.0077; chronic stress versus RNAi group, $P = 0.0005$). **(g)** 24-h RER curve of control,

chronic stress, and RNAi groups (two-way ANOVA: control versus chronic stress

group, $P = 0.0123$, $F(1, 14) = 8.244$; chronic stress versus RNAi group, $P = 0.0198$, $F(1,$

$13) = 7.055$; 8 mice in each group). **(h)** 24-h EE curve of control, chronic stress, and

RNAi groups (two-way ANOVA: control versus chronic stress group, $P = 0.0005$, $F(1,$

$14) = 20.21$; chronic stress versus RNAi group, $P = 0.0073$, $F(1, 14) = 9.851$; 8 mice in

each group).Data are means ± SEM, * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$.

     **Table 1. Electrophysiological properties of three dmVMH neuronal subtypes**

| | Silent (n = 19) | Tonic-firing (n = 38) | Bursting (n = 28) |
|---|---|---|---|
| Input resistance (MΩ) | 535.30 ± 21.83 | 661.72 ± 19.84 | 662.72 ± 24.21 |
| Onset Time (ms, 50 pA) | 37.17 ± 2.53 | 18.20 ± 1.28 | 25.84 ± 1.87 |
| RMP (mV) | -65.27 ± 1.08 | -57.50 ± 0.92 | -63.39 ± 1.07 |
| Overshoot by H-current (mv, -100 pA) | 7.26 ± 1.17 | 10.71 ± 0.94 | 12.01 ± 1.19 |
| Evoke firing rate (Hz, 100 pA) | 17.53 ± 0.60 | 21.76 ± 0.50 | 22.00 ± 0.73 |
| AHP (mV) | -12.38 ± 0.53 | -15.2 ± 0.50 | -14.26 ± 0.65 |
| Half width (ms) | 2.17 ± 0.08 | 2.00 ± 0.04 | 2.04 ± 0.05 |
| AP amplitude (mV) | 71.79 ± 1.88 | 73.29 ± 1.25 | 76.27 ± 1.28 |

972   Data are represented as mean ± SEM. RMP: resting membrane potential; H-current:

973   hyperpolarization-activated current; AHP: after-hyperpolarization potential; AP: action

974   potential.

    **Supplemental Information**

**Extended Data Fig. 1| Electrophysiological properties of dmVMH neurons**

**subtypes.** Related to Fig. 2.

(**a**) Schematic of location of dmVMH in coronal section slice of mouse brain, mCherry

was specifically expressed in SF-1 (specific dmVMH marker) neurons. Scale bar is 300

μm. (**b**) Whole-cell recording trace from a dmVMH neuron, with current injection of

-100 pA to 100 pA and 10 pA/step.   (**c-e**), Representative traces of whole-cell

recordings showing electrophysiological properties of silent (n = 19), tonic-firing (n =

38), and bursting (n = 28) dmVMH neuronal subtypes. Three subtypes exhibit different

electrophysiological activity at 50, 10, and -100 pA current injection.

**Extended Data Fig. 2| Electrophysiological comparison of three dmVMH**

**neuronal subtypes in control and chronic stress groups.** Related to Fig. 2.

**(a)** Resting membrane potential (RMP) of three neuronal subtypes in control and

stressed mice. Top: silent neurons in control (n = 19) and chronically stress mice (n =

13), *P* = 0.5549; Middle: tonic-firing neurons in control (n = 38) and chronically

stress mice (n = 33), *P* = 0.0047; Bottom: bursting neurons in control (n = 29) and

chronically stress mice (n = 38), *P* = 0.9452. **(b)** Onset time of three neuronal

subtypes in control and stressed mice. Top: silent neurons in control (n = 19) and

chronically stress mice (n = 13), *P* = 0.3029; Middle: tonic-firing neurons in control

(n = 39) and chronically stress mice (n = 33), $P = 0.0767$; Bottom: bursting neurons in

control (n = 27) and chronically stress mice (n = 38), $P = 0.7704$ (unpaired Student's

*t*-test). **(c)** Location of each recorded neuron in dmVMH of control and stressed group,

no region specificity was observed among three subtypes. Data are means ± SEM; *$P$

< 0.05, **$P$ < 0.01. The box plotted at the median extending from the 25-75th

percentile, and the whisker represents Min to Max distribution.

1004

1005

**Extended Data Fig. 3| Ascorbate elicited no obvious effects on burst firing in**

**dmVMH neurons.** Related to Fig. 4.

(**a**) Ascorbate (antagonist of Cav3.2) did not block burst firing of dmVMH neurons (n =

4) induced by 10-pA current injection in cosine waveform.

(**b**) Ascorbate imposed no obvious influence on burst firing of dmVMH neurons, with

no significant shifts found in frequency-current curve after ascorbate application

(two-way ANOVA, $P = 0.6590$, $F(1, 6) = 0.2152$ ). *$P < 0.05$.

1013

1014

**Extended Data Fig. 4| Injection of T-VGCC antagonist partially ameliorated**

**anxiety-like behavior**. Related to Fig. 4.

**(a)** Schematic of microinjection of mibefradil in dmVMH through cannula. **(b)**

Behavioral test before and after drug delivery; left, residence time in central area of

open field of saline group decreased compared with mibefradil group (n = 4 in each

group, $P = 0.0169$, unpaired Student's $t$-test); right, residence time in open arm of saline

group decreased compared with mibefradil group (n = 4 in each group, $P = 0.0156$,

unpaired Student's $t$-test). **(c-d)** Oral administration of mibefradil exerted no obvious

change on food intake (n = 4 in each group, $P = 0.9507$, unpaired Student's $t$-test) but

1024    increased average RER in stressed mice ($P = 0.0171$, unpaired Student's $t$-test). Data

1025    are means ± SEM; *$P < 0.05$.

1026

1027

1028

**Extended Data Fig. 5| Glutamate receptor is critical for generation of burst firing**

**in dmVMH.** Related to Fig. 4.

**(a)** Application of APV blocked generation of burst firing induced by cosine waveform

current injection (-10 pA–10 pA, n = 4), even with depolarizing current injection (10pA)

to eliminate the hyperpolarization caused by APV (lower). **(b)** Application of NBQX

did not abolish generation of burst firing (n = 3).

1035

**Naïve**

**Chronic stress**

Anxiety ↑
Food intake ↓
Energy expenditure ↓

dmVMH

dmVMH

Tonic

Burst

Cav3.1

Cav3.1 ↑

1036
**Extended Data Fig. 6| Schematic representation of the enhancement of burst firing in dmVMH neurons after chronic stress, which was mediated by the elevated expression of Cav3.1.**