Pastures and Climate Extremes: Impacts of warming and drought on the productivity

and resilience of key pasture species in a field experiment

Amber C. Churchill*[a,1], Haiyang Zhang[a], Kathryn J. Fuller[a], Burhan Amiji[a], Ian C.

Anderson[a], Craig V. M. Barton[a], Yolima Carrillo[a], Karen L. M. Catunda[a], Manjunatha H.

Chandregowda[a], Chioma Igwenagu[a], Vinod Jacob[a], Gil Won Kim[a,b] Catriona A. Macdonald[a],

Belinda E. Medlyn[a], Benjamin D. Moore[a], Elise Pendall[a], Jonathan M. Plett[a], Alison K.

Post[a,c], Jeff R. Powell[a], David T. Tissue[a], Mark G. Tjoelker[a], Sally A. Power[a]

* Corresponding author: achurchi@umn.edu

[a]Hawkesbury Institute for the Environment

Western Sydney University

Hawkesbury Campus

Ground Floor, Building R2

Locked Bag 1797

Penrith 2751, NSW Australia

Emails: ACC- achurchi@umn.edu, HZ- h.zhang3@westernsydney.edu.au, KJF-

k.fuller@westernsydney.edu.au, BA- b.amiji@westernsydney.edu.au, ICA-

i.anderson@westernsydney.edu.au, CVMB- c.barton@westernsydney.edu.au, YC-

y.carrillo@westernsydney.edu.au, KLMC- k.marinhocatunda@westernsydney.edu.au, MHC-

m.chandregowda@westernsydney.edu.au, CI- c.igwenagu@westernsydney.edu.au, VJ-

v.jacob@westernsydney.edu.au, GWK- gilwonkim74@gmail.com, CAM-

c.macdonald@westernsydney.edu.au, BEM- b.medlyn@westernsydney.edu.au, BJM-

b.moore@westernsydney.edu.au, EP- e.pendall@westernsydney.edu.au, JMP-

j.plett@westernsydney.edu.au, AKP- akpost@colostate.edu, JRP-

jeff.powell@westernsydney.edu.au, DTT- d.tissue@westernsydney.edu.au, MGT-

m.tjoelker@westernsydney.edu.au, SAP- s.power@westernsydney.edu.au

[b]Institute of Agriculture and Life Sciences

Gyeongsang National University,

501 Jinju Daero

Jinju, 52828, South Korea

[c]Department of Biology and Graduate Degree Program in Ecology

Colorado State University

1878 Campus Delivery

Fort Collins, CO 80523-1878 USA

[1]Present address: Department of Ecology, Evolutionary Biology and Behavior

University of Minnesota

140 Gortner Laboratory

1479 Gortner Ave.

St. Paul, MN 55108 USA

Abstract

Shifts in the timing and frequency of climate extremes, such as drought and heatwaves, can

generate sustained shifts in ecosystem function with important ecological and economic

impacts for rangelands and managed pastures. The Pastures and Climate Extremes

experiment (PACE) in southeast Australia used a factorial combination of elevated

temperature (ambient +3 °C) and winter/spring extreme drought (60% rainfall reduction) to

evaluate the impacts of increased frequency of climate extremes on pasture productivity and

subsequent summer/autumn recovery. The experiment included nine species comprising

three plant functional groups (C_3 grasses, C_4 grasses, and legumes) in monoculture and three

two-species mixtures. The winter/spring drought resulted in productivity declines of up to

73% (*Digitaria eriantha*) during the 6-month treatment period, with nine of the twelve

plantings exhibiting significant yield reductions. Functional group identity was not an

important predictor of yield response to drought. Many species recovered rapidly once the

drought ended, although there were carry-over effects on warm season (summer/autumn)

growth for four species/mixtures, spanning all functional groups. Cool season drought

translated into significant reductions in annual biomass production for four species/mixtures,

ranging from 33% (*Medicago sativa*) to 70% (*Festuca arundinacea*). Additionally, warming

had neutral to negative effects on productivity during both winter/spring and summer/autumn

periods, resulting in annual yield declines of up to 58%, driven at least partially by indirect

effects on soil water content. The combination of winter/spring drought and year-round

warming resulted in net yield reductions that were either additive or less-than-additive,

compared to ambient plots. This study demonstrates that predicted extreme climate

conditions will have substantial negative impacts on productivity of common pasture and

rangeland species.

Key words (6-10)

Climate warming, drought, grassland, rangeland, aboveground production, resistance and

resilience, PACE research facility

1 Introduction

Climate change is identified as a dominant driver of ecosystem change across the globe (Sage, 2019; Steffen et al., 2015). Exposure to high temperatures or severe drought has been shown to disrupt physiological function and alter plant species interactions, ultimately driving changes in productivity and ecological processes such as nutrient and water cycling (Backhaus et al., 2014; Frank et al., 2015; Ma et al., 2015; Parmesan et al., 2000). Predicting how altered climate will impact these processes is difficult as climate extremes likely play a significant role in driving change in ecosystem function whereas models tend to focus on differences in mean annual temperature or precipitation (Easterling et al., 2000; Jentsch et al., 2009; Kreyling et al., 2008a). Despite a recent increase in the number of studies focusing on climate extremes (De Boeck et al., 2019, 2015; Hanson and Walker, 2019; Knapp et al., 2017), relatively few have addressed the ecological implications of seasonal shifts in patterns of extreme conditions for contrasting plant functional groups (i.e. C3 vs. C4, legumes; Beier et al., 2012). This is especially true for studies considering the impact of multiple climate variables simultaneously (Beier et al., 2012).

Ecosystems differ considerably in their sensitivity to drought. Managed grasslands and rangelands dominated by temperate grasses are generally considered to be drought sensitive, while tropical grasslands and legumes tend to be more resistant (Adams et al., 2016; Arredondo et al., 2016; Pearcy et al., 1987; Pearcy and Ehleringer, 1984; Wilcox et al., 2016). Ecosystem and plant responses to drought can be considered in terms of resistance - defined as the degree to which processes of interest remain unchanged during drought - or resilience, considered as the time required for recovery to pre-drought conditions (Kreyling et al., 2017; Pimm, 1984). Many studies have examined the potential for resistance versus resilience trade-offs among species in the context of drought (Hoover et al., 2014; Mitchell et

al., 2016) and there are a wide range of plant traits hypothesized to drive these responses.

Examples include photosynthetic pathways and phenology (C_3 vs. C_4 grasses), growth forms

(grass vs. forb), life history (perennial vs. annual) and ecosystem factors such as community

species richness or functional diversity; indeed species- and/or functional group diversity is

typically associated with both higher productivity and increased stress resistance in

herbaceous communities, compared to lower diversity systems (Isbell et al., 2015; Vitasse et

al., 2019; Vogel et al., 2012). Assessing both ecosystem resistance and resilience can be

difficult in field settings due to requirements for long-term monitoring, inter-annual

variability in abiotic conditions and ecological processes such as succession. Consequently,

resilience has been assessed in less than half of all studies evaluating ecological

consequences of an extreme event (Maxwell et al., 2019), although this aspect of plant

response is key to understanding long-term impacts and consequences for ecosystem services.

In addition to changing rainfall regimes, rising temperatures are predicted to have direct and

indirect impacts on many ecosystem processes (Bai et al., 2013; Novem Auyeung et al.,

2013; Song et al., 2019). Direct impacts range from shifts in the timing of peak productivity

relative to the start of the growing season (typically earlier) to extending the length of the

growing season such that production occurs earlier and/or later in the year (Keatinge et al.,

1998; Walker et al., 2006). The majority of field-based temperature manipulations have been

carried out in temperate or cold-climate ecosystems where low temperatures constrain growth

for at least a portion of the year. These studies often find increased total productivity and

shifts in growth phenology associated with warming (Aerts et al., 2006; Kueppers et al.,

2017; Lu et al., 2013; Walker et al., 2006), although there are exceptions (Arnone et al.,

2011; Deutsch et al., 2011; Dukes et al., 2005). Warming experiments in warm-temperate or

subtropical climates with comparatively short mild winters and hot summers are relatively

rare but needed to test whether climate warming exceeds optimal temperatures for growth

(Dukes et al., 2005; Song et al., 2019; Volder et al., 2013).

Plant functional groups are predicted to differ in their response to warming, with warm-

season species such as tropical grasses likely to benefit from increased winter or spring

temperatures, while temperate grasses may be particularly vulnerable to higher temperatures

(Munson and Long, 2017). Legume response to warming is likely to vary between seasons.

For example, as with grasses, warming during the cooler months can bring temperatures

closer to thermal optima (Peng et al., 2020; Sanz-Sáez et al., 2012; Whittington et al., 2013).

However, higher summer temperatures may result in growth reductions as a consequence of

thermal constraints on nitrogen fixation and increased respiratory carbon losses as

temperatures exceed temperature optima for photosynthesis (Aranjuelo et al., 2007;

Whittington et al., 2012). Understanding how warmer temperatures will affect the timing and

amount of different grassland species' biomass production across seasons is an important

knowledge gap in the context of both managed pastures and native grassland ecology.

While studies considering the impact of either precipitation or temperature on plant

physiology and ecosystem function are valuable, it is important to note that warmer

temperatures associated with extreme heatwaves frequently co-occur with drought (Boer et

al., 2020; Dai, 2011; De Boeck et al., 2010; Yuan et al., 2016). Consequently, it is necessary

to evaluate the impacts of co-occurring climate extremes on plant community structure and

ecosystem function (Hoeppner and Dukes, 2012; Sherry et al., 2008). The **PA**stures and

Climate **E**xtremes (PACE) field experiment sought to address this key area of research by

exposing pasture and rangeland plant species to extreme winter and spring drought (60%

rainfall reduction), in combination with elevated temperature (ambient + 3°C). Climate

models predict that temperate/sub-tropical Australia will be subject to large reductions in

winter and spring rainfall (CSIRO, 2020). Increasing frequency and severity of drought is a

consistent prediction for many grassland regions across the globe. Given the importance of

pastures and rangelands for sustaining food security (Godde et al., 2020; Nábrádi, 2016;

O'Mara, 2012), we established a large-scale, field experiment (the PACE facility) to evaluate

plant species' responses to extreme winter/spring rainfall reduction and year-round warming,

including intensification of heatwaves. Specifically, we asked: 1) How does extreme

winter/spring drought affect species' productivity in monoculture and two-species mixtures?

2) Do the effects of winter/spring drought persist through the following summer and autumn

(i.e. species' drought resilience) and what is the effect on annual productivity? 3) How does

warming affect species' productivity in combination with drought? and, 4) Are there

consistent functional group differences in response to drought and warming among pasture

species? Here, we show that tropical and temperate grasses and legumes exhibited large and,

in some cases persistent, reductions in productivity associated with both drought and elevated

temperatures, and highlight the role of species mixtures in mitigating some of the effects of

climate extremes on pasture productivity.

2 Methods

2.1 Site description

The PAstures and Climate Extremes (PACE) facility was constructed in 2017 at the

Hawkesbury Campus of Western Sydney University, in Richmond, NSW, Australia

(S33.60972, E150.73833, elevation 25 m). Mean annual precipitation at this location is 800

mm (Australian Government Bureau of Meteorology, Richmond - UWS Hawkesbury Station

1980-2010); however, there is large inter-annual variability (between 500 mm and over 1400

mm over the past 30 years). Winter/spring precipitation accounts for 40% of annual rainfall.

Mean annual temperature is 17.2 °C, with the monthly maximum and minimum occurring in

January (22.9° C) and July (10.2° C), respectively. The site is fenced to prevent access by

large herbivores. The soil is a loamy sand with a volumetric water holding capacity of 15-

20% (**Table S1**).

The field facility comprised six replicate polytunnel shelters constructed from galvanized

steel frames. These were covered with a single layer of polyethylene 180-micron plastic

(Argosee, Australia) to intercept all ambient precipitation, although the long sides were open

to a height of 1.5 m to allow free flow of air (**Figure 1**). Shelters were 48 m long by 8 m

wide, with a maximum height of 4.6 m, and were oriented along a SW-NE axis with the open

ends facing into the direction of the prevailing wind. Each shelter had eight treatment plots (4

m by 4 m; **Figure 1**) that were further subdivided into four subplots (**Figure 1B; Figure S1**),

with different grassland species or two-species mixtures (hereafter referred to as "plantings")

assigned at the subplot level (total 192 subplots). All surface soils were rotary-tilled to a

depth of 12 cm to homogenize the upper soil profile prior to pasture establishment.

Additionally, all 4 x 4 m plots had a full root barrier installed to a depth of 90 cm to ensure

hydrological isolation between treatments and the 2 x 2 m subplots had an additional root

barrier between them to a depth of 30 cm to minimise root ingress.

Nine plant species were grown in monoculture subplots along with three sets of two-species

mixtures, for a total of twelve different plantings replicated in six independent shelters.

Mixed two-species plots were included to examine drought resistance and resilience of a

range of functional group pairings, compared to their respective monocultures. Species

encompassed a range of functional diversity (C_3/C_4 grasses, legumes, annuals and perennials)

and origins (native grasses, tropical pasture, temperate pasture; **Table 1**) that are all

commonly used in improved grasslands (pastures) or rangelands. For comparisons among

common groupings, we considered three plant functional groups - legumes, C_3 grasses

(temperate) and C_4 grasses (tropical); a fourth category comprised the mixed two-species

plantings. Sward establishment was initiated during early spring 2017 (**Table S2**). Initial

sowing included a fertilizer addition of diammonium phosphate (DAP;110 kg ha^{-1}), and

swards were subsequently managed via hand-weeding, herbicide and, where needed,

insecticide application to maintain target species following industry practice. Plantings with a

legume component received appropriate rhizobium inoculant during sward establishment:

ALOSCA granular inoculant for *Biserrula* subplots (Group BS; ALOSCA Technologies,

Western Australia, Australia); Easy Rhiz soluble legume inoculant and protecting agent to

Medicago subplots (Group AL; New Edge Microbials, New South Wales, Australia); and

NoduleN for the *Trifolium* subplots (Group C; New Edge Microbials). Plantings received top-

up fertilization seasonally to replace nutrients removed from the soil (55 kg ha^{-1}; Cal-Gran

Aftergraze, Incitec Pivot Fertilisers, Australia).

2.2 Experimental treatments

All 12 planting combinations were exposed to a winter/spring drought regime implemented

with reduced irrigation (Drought), and a subset of four plantings also received a warming

treatment in a factorial combination with drought (**Figure S1**). The drought treatment

comprised a control (Control: C) and a drought (Drought: D) watering regime that was

applied during the 6-month austral winter/spring period (1 June to 30 November). For the

following 6-month summer/autumn period (1 December to 31 May) all plots received the

control irrigation amount. A 60% reduction in winter/spring rainfall was chosen for the

drought manipulation as representing the upper end of climate model predictions for end-of-

century seasonal rainfall change for south-eastern Australia, under RCP8.5 (CSIRO, 2020).

Furthermore, a 60% reduction in winter/spring was also in line with historical climate

extremes for key pasture growing regions across south-eastern Australia (BOM, 2019); this

treatment therefore represented historically relevant rainfall extremes, which are predicted to

increase in both frequency and duration. The control was set to represent a typical

precipitation regime for the local area, accounting for long-term patterns in seasonality and in

the statistical distribution of event sizes and timing within seasons (**Figure S2**). The drought

treatment was applied across all plantings of a species/mixture once swards were fully

established, which generated staggered starting times and durations of drought among

different species/mixtures (**Table 1**).

Four plantings consisting of two monocultures (*Festuca, Medicago*) and two two-species

mixtures (*Phalaris/Trifolium and Themeda/Rhytidosperma*) were also exposed to a warming

treatment in a factorial combination with the winter/spring drought. These four plantings

were grouped together as 2 x 2 m subplots in a single 4 x 4 m plot. The warming treatment

comprised a year-round temperature increase of +3 °C (**Table 1**) achieved using infra-red

(IR) heaters. Elevated temperatures (eT) were applied to two 4 x 4 m plots within each

shelter; one plot received the control irrigation (eT-C) the other plot the drought treatment

(eT-D) and these were paired with two ambient temperature (aT) plots, one receiving control

irrigation (aT-C) and the other the drought treatment (aT-D). Each warmed plot had a heating

array comprising eight 1000W ceramic heaters (FTE 1000W, Ceramicx, Ireland) mounted on

an aluminium frame (4 m x 4 m) suspended 1.4 m above ground level (**Figure 1C**). Lamps

were angled to give uniform coverage of IR radiation across the four composite subplots. The

power level to the heating lamps was adjusted each minute, via pulse width modulation using

a solid-state relay controlled by a data logger (CR1000, Campbell Scientific) using a

proportional-integral control algorithm. Target temperatures for these plots were controlled

via feedback from IR-sensors (SI-100, Apogee Instruments, Logan, UT, USA) mounted at a

height of 3.8 m, recording canopy surface temperatures every five minutes. The ambient + 3

°C warming treatment was applied with reference to canopy temperatures for the relevant

control and drought treatments (i.e. aT-C paired to eT-C and aT-D paired to eT-D) to account

for differences in soil moisture which can have an indirect effect on surface temperature.

2.3 Environmental monitoring

Each shelter had a data logger (CR1000, Campbell Scientific) that recorded environmental

conditions and regulated the heating lamps. Soil moisture sensors (16 per shelter; Time

Domain Reflectometers; CS616, Campbell Scientific) recorded soil moisture (0-15 cm) every

15 min in all six replicates of four different species monoculture subplots and treatment

combinations (**Table S3**); in the *Medicago* subplot type soil moisture was also monitored at a

second depth (15-30 cm). Soil temperature probes (T107, Campbell Scientific) were installed

into the top 6-12 cm of the soil of eight subplots per shelter (all four drought and warming

treatment combinations of *Festuca* and *Medicago*) to record soil temperature every 15 min

(**Table S3**). Air temperature and humidity sensors (Series RHP-2O3B, Dwyer Instruments

Inc, USA) mounted in force-ventilated radiation shields were installed inside three of the

rainout shelters at 0.6 m height, with records collected every 5 min to determine shelter

effects on environmental conditions. Additionally, three sets of sensors were installed at the

same height outside the shelters. Photosynthetically active radiation (PAR) was recorded at 5-

min intervals using PAR sensors (Apogee quantum sensor, USA) installed at a height of 6 m

outside two shelters, with two additional sensors located within shelters at a height of 3 m.

2.4 Biomass harvests

All subplots were regularly harvested to determine aboveground productivity during active growing periods. This 'surrogate grazing' involved use of hand shears and a sickle mower. Timing of harvests was based on grazing recommendations for individual species and mixtures (Clements et al., 2003); hence, we conducted 3-5 harvests per year for perennial species (Clark et al., 2016) and 2-3 harvests for annual species that died back at the end of spring. Plants were cut to 5 cm above the soil surface and weighed (fresh and dry mass), with a sub-sample of harvested material sorted to determine the proportion of live and dead biomass as well as the weed fraction from each subplot. All materials were oven-dried at 70 °C for at least 48 hours.

2.5 Calculations and statistical analyses

The response of temperature and soil water content to applied drought and warming treatments were determined based on subplot-level (soil moisture, soil temperature) or plot-level (surface temperature) daily mean values. Aboveground production responses were determined based on subplot totals, summed across all harvests conducted within three defined time periods. Data were analysed separately for the 6-month winter/spring drought period (Drought period), the following 6-month summer/autumn recovery period (Recovery period), and for the combined 12-month period from June 2018-May 2019 (Annual).

Statistical analyses of treatment effects were carried out using linear mixed effects models with climate treatments ('drought': D or C, 'warming': aT or eT) as fixed effects and random effects defined as 'subplot nested within plot' or 'plot' (to account for non-independence among continuous measurements) nested within 'shelter' (to account for the blocked design), and 'date' (to account for temporal variation). Soil moisture data in *Medicago* subplots were

also analysed for differences between upper (0-15 cm) and lower (15-30 cm) depths among

climate treatments by including 'depth' as an additional fixed effect. Climate treatment

effects on aboveground production were analysed using two different linear mixed effects

models based on our hypotheses comparing responses among different functional plant

groups, and mixtures, using a 'functional group' categorical predictor fixed effect. For these

analyses, 'drought' and 'warming' treatments were included as fixed effects; 'planting type'

was also included as random effect for models examining differences among 'functional

groups' as a fixed effect, to account for variability within plantings and 'plot nested within

shelter' was included as a nested random effect for all models. Aboveground production was

transformed using the natural logarithm to meet assumptions of normality where indicated in

statistical tables. All analyses were conducted in R version 4.0.0 (R Core Team, 2020) using

the package lme4 (Bates, 2015). Pairwise comparisons to determine treatment effects on

temperature of soil or canopy and soil water content, or aboveground production among

plantings or functional groups, were conducted using the R package emmeans (Length, 2019)

and using the Tukey method for P-value adjustment.

To compare responses to drought among species/mixtures (i.e. "plantings") and functional

groups, log response ratios between control and droughted subplots paired within a shelter

were calculated on aboveground production data for each period (winter/spring,

summer/autumn, and annual). These ratios were used to evaluate differences in the effect of

drought among 1) plantings and 2) functional groups using linear mixed effects models, with

both specified as fixed effects. Random effects included 'shelter' for both models, while the

model for 2) also included 'planting type' as an independent random effect. We additionally

calculated mean treatment effect size (Equation 1) for each species on log-transformed

production, with mean treatment level effects back-transformed and scaled such that 0 values

indicate no yield reduction. Standard error for the mean effect size was calculated as the

product of the back-transformed mean and the standard error of the effect size.

Equation 1.   *Effect size* $=e^{\left(ln\left(\frac{Drought}{Control}\right)-1\right)}$

Finally, to account for differences in drought duration among plantings (Table 1), we

calculated a drought sensitivity metric to provide a standardized comparison of production

responses as a function of the amount of rainfall withheld (mm), compared with control

treatments during the six-month winter/spring drought and annual periods (Smith et al., 2017;

Equation 2). Drought sensitivity responses were also analysed using linear mixed effects

models following the statistical model structure of the log response ratios above to examine

differences among plantings and functional groups.

Equation 2.   *Sensitivity Index* $= -\dfrac{Control\ production\text{-}\ drought\ production\ (kg\ ha^{-1})}{Control\ irrigation\text{-}drought\ irrigation\ (mm)}$

Interactive effects underlying the effect of warming on drought responses, the effect of

drought on warming responses, and the effect of warming on drought effects, were calculated

as the ratio between single factor treatment effects within a shelter for each period (Dieleman

et al., 2012). These ratios were calculated such that the effect of drought was examined by

comparing the effect size (Equation 1) for each warming level (aT-D / aT-C, eT-D / eT-C),

the effect of warming was examined as the effect size for each drought level (eT-C / aT-C,

eT-D / aT-D) and the effect of warming on the drought response as the ratio between these

two effects ((eT-D / aT-D)/(eT-C / aT-C)).

3 Results

3.1 Drought and warming treatment effects on environmental conditions

Experimental drought and warming treatments significantly altered soil water content and

temperature during the winter/spring period when drought was applied. This effect lasted

into the summer/autumn drought recovery period resulting in overall treatment differences in

soil moisture, soil temperature and surface temperature at the annual scale (**Figure 2**, **Figure**

S3-6, **Tables S4**, **S5**). During the winter/spring period, droughted subplots of *Biserrula* and

Festuca had significantly reduced soil moisture in the upper 15 cm (**Figure 2**; **Table S4**; $P <$

0.001) compared with controls, while *Lolium* and *Medicago* soils were unaffected ($P = 0.12$,

$P = 0.16$, respectively). Additionally, soil moisture contents in all droughted subplots were

less variable throughout the winter/spring drought treatment period compared to their

respective control subplots (**Figure 2A, B**). There were no residual effects of drought on soil

moisture during the subsequent summer/autumn period for the two cool season annuals

(*Biserrula* and *Lolium*, the latter effectively functioning as an annual at the study site due to

high summertime temperatures; **Figure S2B**). However, both *Festuca* and *Medicago* had

higher soil moisture in formerly droughted, compared to control, subplots throughout

summer/autumn (*Festuca*: $P = 0.01$, *Medicago*: $P = 0.02$; **Figure 2B**, **Figure S2B**). Across

the full 12-month period, only *Medicago* plantings exhibited a significant drought treatment

effect, with droughted subplots having higher soil moisture levels than controls ($P = 0.05$;

Figure S2C).

Soil moisture was only moderately affected by the warming treatment applied to *Festuca* and

Medicago, with eT reducing soil moisture during the winter/spring drought period for

Medicago ($P = 0.03$; **Figure S2A**), although there were intermittent short-term positive

effects of warming on soil moisture following irrigation events in *Festuca* (**Figure 2B**). There

was no significant effect of the warming treatment on soil moisture during the

summer/autumn period for either of the monitored species, nor interactions with the drought

treatment. Additionally, differences in soil moisture between upper (0-10 cm) and lower (15-

30 cm) depths for *Medicago* soils varied among the different drought and warming treatments

throughout the drought and recovery periods (**Figure S4**).

Plot-level surface temperatures of the plant canopy and soil surface were consistently

increased by both drought ($P < 0.001$, $+ 0.9$ °C) and warming ($P < 0.001$, $+ 3.0$ °C)

treatment, and interactive effects between drought and warming were not significant (**Figure**

S5; **Table S4**). Additionally, warming altered the temperature range within plots (**Figure S5**;

Table S5). The warmed monoculture plots of *Festuca* and *Medicago* had no days when

minimum temperatures fell below freezing (compared to three such days in ambient plots)

and only 20 days when temperatures fell below 5 °C (compared to 62 days in ambient plots).

At the other end of the scale, the warming treatment resulted in an additional 28 days when

summer temperatures exceeded 40 °C and an extra 23 days where temperatures exceeded 45

°C (**Figure 2C, D**). Subsurface soil temperatures were also affected by treatments, with both

drought (*Festuca*) and warming (*Festuca* and *Medicago*) consistently increasing average

temperatures (**Figure S6**). Overall, soil temperature responses were similar among species,

although drought amplified the warming effect in *Festuca* but not *Medicago* soils.

3.2 Aboveground productivity response to drought

During the 6-month winter/spring drought period, total productivity ranged from 1,800 to

7,400 kg ha^{-1} under control conditions (**Figure 3A**). In the control treatment, the most

productive plantings were *Themeda*, *Themeda*/*Rytidosperma*, *Phalaris* and *Chloris,* while

Lolium and *Biserrula* produced the least biomass. Droughted subplots were significantly less

productive than their respective controls, with an average yield reduction of ~40% across the

twelve plantings (**Figure 3B**; **Table 2**). There was also a significant interaction between

drought and planting type on production, reflecting both differences in drought response and

drought duration, the latter associated with the length of time taken for initial sward

establishment before imposition of the drought treatment (**Table 2**). Production in nine of the

twelve plantings was significantly reduced by drought. The remaining three included the

legume monoculture *Biserrula* ($P = 0.15$), which only had three months of drought, and two

that had only a one-month drought (*Rytidosperma*: $P = 0.96$ and *Phalaris/Trifolium*: $P =$

0.77).

The drought effect size, quantified as the log response ratio between control and droughted

subplots compared during the 6-month winter/spring treatment period, varied among

plantings (**Table 3**) with those that received the full six months of drought generally showing

the greatest loss of productivity (e.g. *Digitaria:* 70% and *Fescue:* 59% yield reduction). An

exception to this pattern was *Chloris*, which also received six months of drought, but

exhibited only a 37% reduction. In contrast, *Phalaris* received only one month of drought

treatment but had yield reductions of 34% (**Figure 3B**). This latter result suggests a high

drought-sensitivity for *Phalaris*, as was evident when standardizing the change in

productivity between control and droughted subplots by the amount of irrigation withheld

during the winter/spring drought treatment period (i.e. the species' "drought sensitivity" -

Figure 3C). Indeed, drought sensitivities ranged from high (*Digitaria, Phalaris* and

Themeda), with 6-month winter/spring yield losses of ~ 20 kg ha^{-1} mm^{-1}, to low (*Biserrula*,

Medicago, *Rytidosperma*, and the *Phalaris*/*Trifolium* mixture; **Figure 3C**), with an overall

average 6-month loss of ~9 kg ha^{-1} mm^{-1} across the twelve plantings. The potential benefit of

growing grasses in two-species mixtures versus in monoculture with respect to drought

effects on winter/spring production was species-dependent. For the C_3 grass-legume mixture,

the drought associated decline in winter/spring productivity was lower than that measured in

the dominant grass monoculture (**Figure 3B**; *Phalaris* in *Phalaris/Trifolium* mixture). Once

this effect was standardized by the amount of water withheld, the *Phalaris*/*Trifolium* mixed

species plots had significantly lower drought sensitivity than the *Phalaris* monoculture

(**Figure 3C**; $P = 0.04$). In contrast, for both the grass-grass mixture (*Themeda*/*Rytidosperma*)

and the C_4 grass-legume mixture (*Digitaria*/*Biserrula*), the drought sensitivity of mixtures

was intermediate between that of the component species (**Figure 3C**).

Plant functional group was not a significant predictor of winter/spring yield reduction under

drought treatment (**Table 3**), nor of drought sensitivity as measured by yield loss per mm

irrigation withheld (**Table 4**), although it is noteworthy that all subplots dominated by C_4

grasses experienced significant cool season biomass reductions under drought. During the

summer/autumn (drought recovery) period, the 6-month productivity of perennial species

ranged from 1,300-10,500 kg ha^{-1} in control subplots and 470-7,800 kg ha^{-1} in formerly

droughted subplots. This variation reflects the differing growth phenologies among the C_3

and C_4 species, as well as their varying capacities for recovery following the end of the

experimental drought (**Figure 4A**). Four plantings experienced significant carry-over effects

of drought, including *Medicago*, *Themeda*, *Themeda*/*Rytidosperma* and *Festuca*, (**Figure**

4A), the latter exhibiting the largest 6-month summer/autumn yield reduction in formerly

droughted plots (i.e. lowest resilience, (**Table 3**; **Figure 4B**). There were no significant

differences among functional groups in terms of carry-over effects of drought on

summer/autumn productivity (**Table 3**).

Annual biomass production among perennial plantings reflected their combined drought

resistance in terms of 6-month yield reduction during the winter/spring drought and capacity

to recover production in the following 6-month summer/autumn, once conditions improved.

Annual productivity varied almost four-fold (from 4,300-16,500 kg ha^{-1}yr^{-1}) in control

subplots among the twelve plantings and by a factor of six (1,700-10,700 kg ha^{-1}yr^{-1}; **Figure**

4C) in the droughted subplots. Four of the perennial species experienced significant drought-

related yield reductions across the full 12 months of the study (**Figure 4C**). Although there

were no significant differences in annual productivity) among functional groups, C$_4$ grass

species were more consistently affected by drought compared to temperate C$_3$ grasses, the

latter group including species that were strongly negatively affected (*Festuca*) and others that

were not (*Rytidosperma*; **Figure 4D**).

Table 1. Plant species traits and treatment details for the PACE experiment

Planting	Species*	Origin	Growth Form	Photosynthetic pathways	Drought Started#	Warming Started
Festuca	*F. arundinacea*	Temperate, introduced	Grass	C_3	01/06/2018	16/04/2018
Lolium	*L. perenne*	Temperate, introduced	Grass	C_3	23/08/2018	NA
Phalaris	*P. aquatica*	Temperate, introduced	Grass	C_3	14/10/2018	NA
Rytidosperma	*R. caespitosum*	Temperate, native	Grass	C_3	14/10/2018	NA
Chloris	*C. gayana*	Tropical, introduced	Grass	C_4	01/06/2018	NA
Digitaria	*D. eriantha*	Tropical, introduced	Grass	C_4	01/06/2018	NA
Themeda	*T. triandra*	Tropical, native	Grass	C_4	01/06/2018	NA
Biserrula	*B. pelecinus*	Temperate, introduced	Legume	C_3	23/08/2018	NA
Medicago	*M. sativa*	Temperate, introduced	Legume	C_3	23/08/2018	16/04/2018
Digitaria/ Biserrula	*D. eriantha, B. pelecinus*	Tropical/temperate, introduced mixture	Grass Legume	C_4 C_3	23/08/2018	NA
Phalaris/ Trifolium	*P. aquatica, T. subterraneum*	Temperate, introduced mixture	Grass Legume	C_3 C_3	14/10/2018	16/04/2018
Themeda/ Rytidosperma	*R. caespitosum, T. triandra*	Tropical/temperate, native mixture	Grass Grass	C_3 C_4	01/06/2018	16/04/2018

Plantings referenced by genus name in the text

 # Date (dd/mm/year) of first watering event with 60% reduced irrigation input for drought plots

Table 2. Linear mixed effects model output for aboveground productivity during the 6-month winter/spring drought period (June-Nov. 2018), the 6-month summer/autumn (drought-recovery) period (Dec. 2018- May 2019 and annually (June 2018-May 2019).

Period	Fixed Effects	F value	*P* value	R^2m[#]	R^2c[§]
Drought*	Drought	100.9	<0.01	0.66	0.71
	Plantings[^]	17.0	< 0.01		
	Drought x Plantings	3.9	< 0.01		
Recovery	Drought	9.4	0.01	0.86	0.86
	Plantings	75.3	<0.01		
	Drought x Plantings	2.4	0.01		
Annual*	Drought	48.7	<0.01	0.81	0.83
	Plantings	51.8	<0.01		
	Drought x Plantings	4.5	<0.01		

Biomass data were log transformed to meet assumptions of constant variance
[^]Refers to different species/mixture types
[#]R^2m indicates marginal error associated with linear model fixed effects
[§]R^2c indicates conditional error or the total variation described by the full model including nested random effects (plots within a shelter).

Table 3. Linear mixed effects model output for comparison of log response ratios (LRR) and sensitivity of aboveground production in response to drought treatment among plantings (i.e. different species/mixtures) and functional groups (FG) for the 6-month winter/spring drought period, the 6-month summer/autumn (drought-recovery) period and the entire 12 months (annual).

Response	Period	Fixed Effects	F value	*P* value	R^2m	R^2c
LRR	Drought	Plantings*	5.1	<0.01	0.43	0.46
		FG*[#]	1.1	0.39	0.12	0.49
	Recovery	Plantings*	14.2	<0.01	0.67	0.69
		FG*[#]	0.2	0.87	0.04	0.64
	Annual	Plantings*	8.1	<0.01	0.55	0.55
		FG*[#]	0.4	0.75	0.08	0.63
Sensitivity	Drought	Plantings*	3.1	<0.01	0.31	0.35
		FG*[#]	1.1	0.40	0.09	0.34
	Annual	Plantings*	2.9	<0.01	0.30	0.34
		FG*[#]	0.4	0.76	0.04	0.35

Includes shelter as a random effect
[#]Includes plantings as a random effect

3.3 Effects of warming and drought-warming interaction on aboveground biomass

production

The effects of warming varied across time periods and among the four different planting

types, with significant warming x planting interactions in each period (**Table 4**). Warming

resulted in a significant decline (28 %) in winter/spring productivity in *Phalaris/Trifolium*

(**Figure 5A, Figure 6B**; **Table 4**) that translated to net annual reductions (29 %) associated

with warming (**Figure 5C**). In contrast, warming reduced productivity for *Festuca* during the

6-month summer/autumn warm season (**Figure 5B**), but not during the winter/spring months,

an effect that also resulted in net annual reductions in aboveground productivity for this

species (**Figure 5C**). Importantly, no plantings experienced a significant productivity benefit

associated with warmer temperatures at any time period (**Figure 6B**).

There was a significant interaction between warming and drought treatments for aboveground

productivity during the winter/spring period and at the annual scale; there were also

significant three-way interactions between drought, warming, and planting type (**Table 4**)

across both 6-month time periods and the entire year, highlighting the species-contingency of

climate treatment impacts on plant productivity. During the summer/autumn, warming

amplified the carry-over effects of winter/spring drought on *Festuca* (**Figure 5B**) with

formerly droughted and warmed subplots producing significantly less biomass than other

treatments. In contrast, for winter/spring and annual *Festuca* productivity, warming partially

offset the negative effects of drought (winter/spring- aT: -63% yield, eT: -22% yield, **Figure**

6A). There was no evidence of greater-than-additive (i.e. synergistic) impacts (**Figures 5A,**

6C) of the combined climate manipulation treatments for any species.

Table 4. Linear mixed effects model output for aboveground production in response to drought (D) and warming (W) treatments among plantings (P)

Period	Fixed Effects	F value	*P* value	R^2m	R^2c
Drought[*#]	Drought	61.4	*<0.01*	0.66	0.72
	Warming	2.0	0.16		
	Planting	36.9	*<0.01*		
	D * W	3.9	0.05		
	D * P	8.3	*<0.01*		
	W * P	4.3	*0.01*		
	D * W * P	3.5	*0.02*		
Recovery[*#]	Drought	37.1	*<0.01*	0.88	0.88
	Warming	44.3	*<0.01*		
	Planting	136.2	*<0.01*		
	D * W	3.0	0.10		
	D * P	27.5	*<0.01*		
	W * P	26.0	*<0.01*		
	D * W* P	4.6	*0.01*		
Annual[*#]	Drought	45.7	*<0.01*	0.86	0.88
	Warming	17.0	*<0.01*		
	Planting	180.9	*<0.01*		
	D * W	6.3	*0.01*		
	D * P	9.3	*<0.01*		
	T * P	4.4	*0.01*		
	D * W * P	3.6	*0.02*		

*Includes plots within shelter as a nested random effect

[#] *Log transformation applied*

4 Discussion

Projections of future climate change, including increases in the frequency and magnitude of extreme climate events, are likely to disrupt ecosystem functioning (Jentsch and Beierkuhnlein, 2008; Knapp et al., 2008; Nijp et al., 2015), with important consequences for pastures and rangelands (Godde et al., 2020; Jiménez et al., 2011). We found that most of the C_3 and C_4 pasture species and mixtures exposed to extreme winter/spring drought experienced significant reductions in cool season productivity, with declines of up to 73%, relative to controls. Grass-legume plantings were more drought-resistant than their respective monocultures, significantly so for one species pair. Despite substantial impacts of winter/spring drought on cool season production, most species recovered quickly, with only four plantings experiencing carry-over effects into the subsequent summer/autumn. We found no evidence of a positive effect of warming on temperate species' productivity during the 6-month winter/spring period and instead observed significant reductions in productivity across the year for two of the four plantings exposed to increased temperatures. Furthermore, the combination of warming and drought generally resulted in additive or less-than-additive effects on biomass production, although always with net negative impacts, relative to controls.

4.1 Productivity responses to winter/spring drought

Decades of research focused on examining grassland responses to various drought regimes have emphasized the importance of regional climate context (Heisler-White et al., 2009; Knapp et al., 2017; Slette et al., 2019) and of standardizing observed drought effects relative to the amount of precipitation applied and the magnitude of drought imposed (Felton et al., 2020; Gao et al., 2019; Knapp et al., 2015). When we accounted for differences in drought duration by expressing yield reductions relative to the magnitude of rainfall reduction (Smith

et al., 2017; Wilcox et al., 2017), *Phalaris* and *Themeda* emerged as the more drought-

sensitive species, while *Rytidosperma* and the *Phalaris*/*Trifolium* mixture were least affected;

all species sensitivities (in terms of yield reductions per rainfall reduction) were, however,

within the range reported from grassland studies elsewhere (Wilcox et al., 2017).

On average, cool season growth of C_4 grasses was more affected by winter/spring drought

than legumes, with C_3 grasses showing the greatest range of responses. Interestingly,

significant effects of cool season drought on the (predominately summer active) C_4 grasses

also translated to declines in annual biomass production, demonstrating strong, negative

effects of drought occurring prior to the main period of active growth in these tropical

grasses. These apparent ecological mis-matches are often difficult to observe in temperate

systems where the majority of growth occurs during a well-defined warm season (Knapp et

al., 2015). There is, however, evidence for winter precipitation offsetting the effects of

spring/summer drought (Fry et al., 2013), as well as cool season rainfall driving soil water

storage and long-term patterns of productivity in semi-arid grasslands and regions with more

continuous patterns of plant growth (Derner et al., 2020). These findings have important

implications for the seasonality of plant productivity and subsequent short-term stocking

decisions made by land managers when faced with extreme climate conditions (Murray-

Tortarolo and Jaramillo, 2019). Locally in south eastern Australia, projections for increased

severity of winter/spring drought (CSIRO, 2020) are likely to have large impacts on the

seasonality of pasture and rangeland productivity, and similar shifts in precipitation may have

far reaching implications for livestock production around the world (Derner et al., 2020).

Other plant functional groups at PACE had less consistent drought responses, including

contrasting responses between legume species and high variability among the C_3 grasses.

Deep tap roots for the legume *Medicago* may have allowed access to deep soil water

supporting sustained growth during dry periods or, alternatively, facilitated shifts in moisture

depth profiles due to hydraulic lift (Li et al., 2012; Nie et al., 2015; Raza et al., 2013) thereby

reducing drought effects at the sward level (Liste and White, 2008; Pang et al., 2013). Even

so, this species experienced significant declines in productivity in all time periods, although

total production was still comparable to C_3 grasses under control conditions. On the other

hand, *Biserrula*, also a deep-rooted legume (Haling et al., 2016; Loi et al., 2005), experienced

a strong effect of the drought treatment on soil moisture without a significant reduction in

biomass. Differences in growth phenology relative to the timing of drought may explain this

disparity as *Biserrula*, an annual species, reached peak growth in early spring when the

drought treatment was first implemented for this species, suggesting that residual soil water

from winter may have been sufficient to support strong early growth despite the spring

drought (Fay et al., 2000; Hofer et al., 2017). These differences in legume response, along

with the large effects of winter/spring drought on cool season productivity of our tropical

grasses- effects that persisted during the more productive, warm season for one (*Themeda*) of

the three C_4 species - provide support for the importance of drought timing in relation to

species growth phenology (Wilcox et al., 2017; Yao et al., 2019) rather than functional group

classifications.

Grass-legume mixtures were generally more drought resistant in terms of maintaining

production during drought than their respective dominant monocultures, a phenomenon that

has been observed in other systems contrasting different levels of species- (Bloor and

Bardgett, 2012; Isbell et al., 2015) or functional diversity (Gherardi and Sala, 2015; Jentsch et

al., 2009; Kreyling et al., 2008b). The lack of difference in drought response for *Themeda*

grown in (grass-grass) mixture *versus* monoculture suggests that increasing the number of

species was less important than increasing functional diversity. Increased functional diversity,

typically involving legume-grass mixtures, is a management strategy used widely in pasture

systems to increase soil nitrogen (N) availability and improve the nutritional quality of

companion grasses (Nyfeler et al., 2011; Suter et al., 2015). While we have not examined the

mechanism(s) for reduced sensitivity of production to drought here, legume-associated

increases in N availability and its uptake by grasses could potentially improve productivity

under drier conditions by increasing photosynthesis and hence water-use efficiency (Abid et

al., 2016; Adams et al., 2016; Hofer et al., 2017). This management strategy therefore has the

potential to reduce climate change-driven losses in productivity and promote sustainable

pastures under future, more variable climates.

4.2 Species recovery following drought and effects on annual productivity

Despite global interest in the resistance of ecosystems to extreme drought, far fewer studies

have examined the factors driving post-drought recovery or resilience (Maxwell et al., 2019),

especially among seasons in a year. Recovery of the original community following drought

can be rapid (Hoover et al., 2014; Nicholas et al., 2004; Power et al., 2016), prolonged (Niu

et al., 2014) or may never occur (Smith, 2011), with such differences often reflecting the

severity and duration of the drought, land-use history and management regimes (Ruppert et

al., 2015; Vogel et al., 2012). In our experiment, six of the ten perennial pastures fully

recovered from winter/spring drought during the following summer and autumn.

Monoculture plantings that showed full recovery included *Chloris* and *Digitaria*, both C_4

summer-active grasses, as well as the two temperate C_3 grasses that experienced a short (one-

month) duration drought (*Phalaris* and *Rytidosperma*). Previous studies imposing short-

duration or mild/moderate drought conditions also observed full recovery for temperate

grassland species (Nicholas et al., 2004). The predominantly spring-focussed growth

phenology of these temperate grasses - partly associated with very warm summer

temperatures at the experimental site - likely contributed to the minimal legacy effects of

winter/spring drought observed in this study. On the other hand, the complete recovery of the

two C_4 grasses suggests that these species may be particularly well-adapted to maintain

summer production and thus cover a summer feed-gap for livestock (Johnston, 1996), even in

the context of predicted increases in the severity and duration of winter/spring drought.

Extreme drought experiments conducted in North American grasslands have also shown

rapid recovery in aboveground productivity for C_4-dominated systems (Hoover et al., 2014),

providing further support for a role for these grasses in maintaining year-round forage

production under future, drier climates (Godde et al., 2020).

Annual productivity is one of the most commonly used metrics for assessing ecosystem

responses to environmental drivers (Gao et al., 2019; Huxman et al., 2004). Despite generally

high levels of drought resilience, four plantings experienced significant drought-associated

declines in annual yield. These included species with wide geographic distributions (*Festuca*

and *Medicago*), as well as an Australia native grass (*Themeda*), such that our findings are

likely to be broadly applicable to a wide range of grasslands. Climate-associated declines in

seasonal- and annual productivity have been widely reported (Godde et al., 2020; McKeon et

al., 2009; Murray-Tortarolo and Jaramillo, 2020); increased understanding of these patterns

will help guide climate risk management strategies such as stocking rates of livestock, as well

as sustainability initiatives such as carbon farming (Kragt et al., 2012).

4.3 Plant responses to warming and drought x warming interactions

Model predictions of beneficial impacts of warming on grassland productivity globally (Gao

et al., 2016) and locally (Cullen et al., 2009), especially during cool season months in higher

latitude ecosystems (Mori et al., 2014; Naudts et al., 2011), were not borne out in our study.

This was despite an increase in winter growing degree days and a reduction in frost exposure,

changes that are expected to increase winter and spring growth (Chang et al., 2017; Piao et

al., 2019). While our cool season findings were unexpected, they do, however, align with

model simulations of the effects of moderate (+3 °C) warming on Australian rangelands

(McKeon et al., 2009).

Treatment-related increases in maximum temperatures and the intensity and duration of

summer heatwave events were expected to reduce summer productivity as species exceeded

physiological thresholds, especially for temperate species (Reyer et al., 2013). The observed

neutral (*Medicago* and *Themeda/Rytidosperma*) to negative (*Festuca* and *Phalaris/Trifolium*)

impacts of warming during cool and warm seasons is likely to result from both direct

(physiological) and indirect (soil moisture/VPD) mechanisms. Indeed, we found that warmed

plots had consistently lower soil water contents, associated with temperature-driven increases

in evapotranspiration (Brookshire and Weaver, 2015; Gao et al., 2016; Kuster et al., 2013).

This connection between temperature and evapotranspiration is likely to amplify the intensity

of ecological drought under future climates (Adams et al., 2009; Dai, 2011; De Boeck et al.,

2016; Dreesen et al., 2012). Interestingly, the combination of drought and warming in our

study resulted in additive or less-than-additive (although still net negative) effects on

productivity for all four plantings, that varied by season. Similar less-than-additive

temperature and drought interactions have been observed in temperate grasslands and more

generally across major biomes in meta-analyses (De Boeck et al., 2011; Song et al., 2019;

Wu et al., 2011). Further work investigating the mechanisms and nature of these interactions

will help refine predictions relating to multiple, co-occurring changes in future climate. This

study provides important, new experimental, field-based data on the realized effects of

extreme drought and continuous warming on the productivity of a variety of commonly used

pasture/rangeland species. These data add to information from modelling studies (Cullen et

al., 2009; Kaine and Tozer, 2005) and comparisons across precipitation gradients (Clark et

al., 2016) to quantify impacts of future and more extreme rainfall regimes on the productivity

of the pasture and rangeland feed base that underpins livestock grazing in many parts of the

world (Godde et al., 2020). Future work with these species will investigate relationships

between productivity responses and plant traits and strategies relating to the acquisition and

use of water (root traits, plant-microbial interactions, plant hydraulics) and carbon (above-

versus below-ground allocation, vegetative versus reproductive growth, root and crown

carbohydrate storage), to elucidate mechanisms responsible for species differences in

sensitivity. This information will be key to extrapolating findings from this study to a wider

range of locations and grassland species, including many of international relevance in

production systems across the globe.

5 Conclusions

This paper introduces a new experimental facility used to simulate future, more extreme

climates, under field conditions. Our findings of consistent negative effects of winter/spring

drought and year-round warming on many of the pasture species, including two native

rangeland grasses, that play a crucial role in livestock and dairy industries highlights the

challenges associated with future climate risk management for both pasture-related industries

and native grasslands. Observed productivity declines associated with warming, even in the

cooler winter/spring months, indicate the important role of temperature in altering ecosystem

water balance, and associated increases in the severity of ecological drought in a future,

warmer world. This is likely to be an issue for many grassland systems across the globe, as

higher temperatures exacerbate the effects of predicted increases in the magnitude, frequency

and duration of meteorological drought. Species with physiological, morphological and/or

phenological traits that support sustained productivity under more challenging climate

conditions will become increasingly important as climate variability and extremes undermine

the future performance of traditional pasture and rangeland species.

6 Author Contributions/Funding/Acknowledgements

The authors would like to acknowledge individuals whose efforts supported the data

collection and site management for this work, including Angelica Vårhammar, Jackson

Bridges-Parlett, Alexandra Boyd, Shania Therese Didier Serre, Samantha Weller, Jinyan

Yang and Arjunan Krishnananthaselvan. We also gratefully acknowledge expert input to

early project discussions from Cath Lescun, Tom Davison, Tom Dickson, Allen Newman,

Warwick Badgery, Suzanne Boschma, Richard Eckard, Brendan Cullen and Alan Humphries,

and to Heritage Seeds for providing seeds. This project was supported by funding from Meat

and Livestock Australia's Donor Company (P.PSH. 0793), Dairy Australia (C100002357)

and Western Sydney University. Authors ACC and SAP wrote the paper with input from all

co-authors on draft iterations; ACC conducted statistical analyses with input from BEM, HZ,

JRP and KJF; SAP, EP, MGT, DTT, CVMB, BA, JRP, JMP, CAM, BDM, YC, ICA

designed the field experiment with input from ACC, HZ and KJF; ACC, HZ, KJF, KLMC,

MC, CI, VJ, GWK and AKP collected field samples.

7 References

Abid, M., Tian, Z., Ata-Ul-Karim, S.T., Cui, Y., Liu, Y., Zahoor, R., Jiang, D., Dai, T., 2016. Nitrogen nutrition improves the potential of wheat (*Triticum aestivum* L.) to alleviate the effects of drought stress during vegetative growth periods. Front. Plant Sci. 7, 1–14. https://doi.org/10.3389/fpls.2016.00981

Adams, H.D., Guardiola-Claramonte, M., Barron-Gafford, G.A., Villegas, J.C., Breshears, D.D., Zou, C.B., Troch, P.A., Huxman, T.E., 2009. Temperature sensitivity of drought-induced tree mortality portends increased regional die-off under global-change-type drought. Proc. Natl. Acad. Sci. U. S. A. 106, 7063–6. https://doi.org/10.1073/pnas.0901438106

Adams, M.A., Turnbull, T.L., Sprent, J.I., Buchmann, N., 2016. Legumes are different: Leaf nitrogen, photosynthesis, and water use efficiency. Proc. Natl. Acad. Sci. 113, 4098–4103. https://doi.org/10.1073/pnas.1523936113

Aerts, R., Cornelissen, J.H.C., Dorrepaal, E., 2006. Plant performance in a warmer world: General responses of plants from cold, northern biomes and the importance of winter and spring events. Plant Ecol. 182, 65–77. https://doi.org/10.1007/s11258-005-9031-1

Aranjuelo, I., Irigoyen, J.J., Sánchez-Díaz, M., 2007. Effect of elevated temperature and water availability on CO_2 exchange and nitrogen fixation of nodulated alfalfa plants. Environ. Exp. Bot. 59, 99–108. https://doi.org/10.1016/j.envexpbot.2005.10.008

Arnone, J.A., Jasoni, R.L., Lucchesi, A.J., Larsen, J.D., Leger, E.A., Sherry, R.A., Luo, Y., Schimel, D.S., Verburg, P.S.J., 2011. A climatically extreme year has large impacts on C4 species in tallgrass prairie ecosystems but only minor effects on species richness and other plant functional groups. J. Ecol. 99, 678–688. https://doi.org/10.1111/j.1365-2745.2011.01813.x

Arredondo, T., Garcìa-Moya, E., Huber-Sannwald, E., Loescher, H.W., Delgado-Balbuena, J., Luna-Luna, M., 2016. Drought manipulation and its direct and legacy effects on productivity of a monodominant and mixed-species semi-arid grassland. Agric. For. Meteorol. 223, 132–140. https://doi.org/10.1016/j.agrformet.2016.03.011

Backhaus, S., Kreyling, J., Grant, K., Beierkuhnlein, C., Walter, J., Jentsch, A., 2014. Recurrent mild drought events increase resistance toward extreme drought stress. Ecosystems 17, 1068–1081. https://doi.org/10.1007/s10021-014-9781-5

Bai, E., Li, S., Xu, W., Li, W., Dai, W., Jiang, P., 2013. A meta-analysis of experimental warming effects on terrestrial nitrogen pools and dynamics. New Phytol. 199, 431–40. https://doi.org/10.1111/nph.12252

Bates, D., Maechler, M., Bolker, B.M., Walker, S., 2015. lme4: Linear mixed-effects models
using Eigen and S4.

Beier, C., Beierkuhnlein, C., Wohlgemuth, T., Penuelas, J., Emmett, B.A., Körner, C., De
Boeck, H.J., Christensen, J.H., Leuzinger, S., Janssens, I. a., Hansen, K., 2012.
Precipitation manipulation experiments - challenges and recommendations for the
future. Ecol. Lett. 15, 899–911. https://doi.org/10.1111/j.1461-0248.2012.01793.x

Bloor, J.M.G., Bardgett, R.D., 2012. Stability of above-ground and below-ground processes
to extreme drought in model grassland ecosystems: Interactions with plant species
diversity and soil nitrogen availability. Perspect. Plant Ecol. Evol. Syst. 14, 193–204.
https://doi.org/10.1016/j.ppees.2011.12.001

Boer, M.M., Resco De Dios, V., Bradstock, R.A., 2020. Unprecedented burn area of
Australian mega forest fires. Nat. Clim. Chang. 10, 170. https://doi.org/10.1038/s41558-
020-0710-7

BOM, 2019. Daily Rainfall Richmond-UWS Hawkesbury [WWW Document]. Clim. Data
Online. URL http://www.bom.gov.au/climate/data/index.shtml?bookmark=136

Brookshire, E.N.J., Weaver, T., 2015. Long-term decline in grassland productivity driven by
increasing dryness. Nat. Commun. 6. https://doi.org/10.1038/ncomms8148

Chang, J., Ciais, P., Viovy, N., Soussana, J.F., Klumpp, K., Sultan, B., 2017. Future
productivity and phenology changes in European grasslands for different warming
levels: Implications for grassland management and carbon balance. Carbon Balance
Manag. 12. https://doi.org/10.1186/s13021-017-0079-8

Clark, S.G., Nie, Z.N., Culvenor, R.A., Harris, C.A., Hayes, R.C., Li, G.D., Norton, M.R.,
Partington, D.L., 2016. Field evaluation of cocksfoot, tall fescue and *Phalaris* for dry
marginal environments of South-Eastern Australia. 1. Establishment and herbage
production. J. Agron. Crop Sci. 202, 96–114. https://doi.org/10.1111/jac.12152

Clements, B., Ayres, L., Langford, C., McGarva, L., Simpson, P., Hennessy, G., Keys, M.,
Upjohn, B., Leech, F., 2003. The grazier's guide to pastures: sowing and managing
profitable pastures in the central and southern Tablelands, Monaro and Upper South
West Slopes of New South Wales. NSW.

CSIRO, 2020. State of the Climate, Technical Report. https://doi.org/ISBN:978-1-4863-
1509-3

Cullen, B.R., Johnson, I.R., Eckard, R.J., Lodge, G.M., Walker, R.G., Rawnsley, R.P.,
McCaskill, M.R., 2009. Climate change effects on pasture systems in south-eastern
Australia. Crop Pasture Sci. 60, 933–942. https://doi.org/10.1071/CP09019

Dai, A., 2011. Drought under global warming: A review. Clim. Chang. 2, 45–65.

https://doi.org/10.1002/wcc.81

De Boeck, H.J., Bassin, S., Verlinden, M., Zeiter, M., Hiltbrunner, E., 2016. Simulated heat

waves affected alpine grassland only in combination with drought. New Phytol. 209,

531–541. https://doi.org/10.1111/nph.13601

De Boeck, H.J., Bloor, J.M.G., Aerts, R., Bahn, M., Beier, C., Emmett, B.A., Estiarte, M.,

Grünzweig, J.M., Halbritter, A.H., Holub, P., Jentsch, A., Klem, K., Kreyling, J., Kröel-

Dulay, G., Larsen, K.S., Milcu, A., Roy, J., Sigurdsson, B.D., Smith, M.D., Sternberg,

M., Vandvik, V., Wohlgemuth, T., Nijs, I., Knapp, A.K., 2019. Understanding

ecosystems of the future will require more than realistic climate change experiments – A

response to Korell et al. Glob. Chang. Biol. 19–20. https://doi.org/10.1111/gcb.14854

De Boeck, H.J., Dreesen, F.E., Janssens, I.A., Nijs, I., 2011. Whole-system responses of

experimental plant communities to climate extremes imposed in different seasons. New

Phytol. 189, 806–817. https://doi.org/10.1111/j.1469-8137.2010.03515.x

De Boeck, H.J., Dreesen, F.E., Janssens, I.A., Nijs, I., 2010. Climatic characteristics of heat

waves and their simulation in plant experiments. Glob. Chang. Biol. 16, 1992–2000.

https://doi.org/10.1111/j.1365-2486.2009.02049.x

De Boeck, H.J., Vicca, S., Roy, J., Nijs, I., Milcu, A., Kreyling, J., Jentsch, A., Chabbi, A.,

Campioli, M., Callaghan, T. V., Beierkuhnlein, C., Beier, C., 2015. Global Change

Experiments: Challenges and Opportunities. Bioscience 65, 922–931.

https://doi.org/10.1093/biosci/biv099

Derner, J.D., Raynor, E.J., Reeves, J.L., Augustine, D.J., Milchunas, D.G., 2020. Climatic

and management determinants of large herbivore production in semiarid grassland.

Agric. Ecosyst. Environ. 290, 106761. https://doi.org/10.1016/j.agee.2019.106761

Deutsch, E.S., Bork, E.W., Cahill, J.F., Chang, S.X., 2011. Short-term plant community

responses to warming and defoliation in a northern temperate grassland. ISRN Ecol.

2011, 1–8. https://doi.org/10.5402/2011/926061

Dieleman, W.I.J., Vicca, S., Dijkstra, F.A., Hagedorn, F., Hovenden, M.J., Larsen, K.S.,

Morgan, J.A., Volder, A., Beier, C., Dukes, J.S., King, J., Leuzinger, S., Linder, S., Luo,

Y., Oren, R., De Angelis, P., Tingey, D.T., Hoosbeek, M.R., Janssens, I.A., 2012.

Simple additive effects are rare: a quantitative review of plant biomass and soil process

responses to combined manipulations of CO_2 and temperature. Glob. Chang. Biol. 18,

2681–2693. https://doi.org/10.1111/j.1365-2486.2012.02745.x

Dreesen, F.E., De Boeck, H.J., Janssens, I.A., Nijs, I., 2012. Summer heat and drought

extremes trigger unexpected changes in productivity of a temperate annual/biannual

plant community. Environ. Exp. Bot. 79, 21–30.

https://doi.org/10.1016/j.envexpbot.2012.01.005

Dukes, J.S., Chiariello, N.R., Cleland, E.E., Moore, L.A., Shaw, M.R., Thayer, S., Tobeck,

T., Mooney, H.A., Field, C.B., 2005. Responses of grassland production to single and

multiple global environmental changes. PLoS Biol. 3, e319.

https://doi.org/10.1371/journal.pbio.0030319

Easterling, D.R., Evans, J.L., Groisman, P.Y., Karl, T.R., Kunkel, K.E., Ambenje, P., 2000.

Observed variability and trends in extreme climate events: a brief review. Bull. Am.

Meteorol. Soc. 81, 417–425. https://doi.org/10.1175/1520-

0477(2000)081<0417:OVATIE>2.3.CO;2

Fay, P.A., Carlisle, J.D., Knapp, A.K., Blair, J.M., Collins, S.L., 2000. Altering rainfall

timing and quantity in a mesic grassland ecosystem: design and performance of rainfall

manipulation shelters. Ecosystem 3, 208319. https://doi.org/10.1007/s

Felton, A.J., Slette, I.J., Smith, M.D., Knapp, A.K., 2020. Precipitation amount and event size

interact to reduce ecosystem functioning during dry years in a mesic grassland. Glob.

Chang. Biol. 26, 658–668. https://doi.org/10.1111/gcb.14789

Frank, D., Reichstein, M., Bahn, M., Frank, D.C., Mahecha, M.D., Smith, P., Thonicke, K.,

van der Velde, M., Vicca, S., Babst, F., Beer, C., Buchmann, N., Canadell, J.G., Ciais,

P., Cramer, W., Ibrom, A., Miglietta, F., Poulter, B., Rammig, A., Seneviratne, S.I.,

Walz, A., Wattenbach, M., Zavala, M.A., Zscheischler, J., 2015. Effects of climate

extremes on the terrestrial carbon cycle: concepts, processes and potential future

impacts. Glob. Chang. Biol. n/a-n/a. https://doi.org/10.1111/gcb.12916

Fry, E.L., Manning, P., Allen, D.G.P., Hurst, A., Everwand, G., Rimmler, M., Power, S.A.,

2013. Plant functional group composition modifies the effects of precipitation change on

grassland ecosystem function. PLoS One 8.

https://doi.org/10.1371/journal.pone.0057027

Gao, J., Zhang, L., Tang, Z., Wu, S., 2019. A synthesis of ecosystem aboveground

productivity and its process variables under simulated drought stress. J. Ecol. 107, 2519–

2531. https://doi.org/10.1111/1365-2745.13218

Gao, Q., Schwartz, M.W., Zhu, W., Wan, Y., Qin, X., Ma, X., Liu, S., Williamson, M.A.,

Peters, C.B., Li, Y., 2016. Changes in global grassland productivity during 1982 to 2011

attributable to climatic factors. Remote Sens. 8, 1–12. https://doi.org/10.3390/rs8050384

Gherardi, L.A., Sala, O.E., 2015. Enhanced interannual precipitation variability increases

plant functional diversity that in turn ameliorates negative impact on productivity. Ecol.

Lett. 18, 1293–1300. https://doi.org/10.1111/ele.12523

Godde, C.M., Boone, R., Ash, A.J., Waha, K., Sloat, L., Thornton, P.K., Herrero, M., 2020.

Global rangeland production systems and livelihoods at threat under climate change and

variability. Environ. Res. Lett. 15, 44021. https://doi.org/10.1088/1748-9326/ab7395

Haling, R.E., Yang, Z., Shadwell, N., Culvenor, R.A., Stefanski, A., Ryan, M.H., Sandral,

G.A., Kidd, D.R., Lambers, H., Simpson, R.J., 2016. Growth and root dry matter

allocation by pasture legumes and a grass with contrasting external critical phosphorus

requirements. Plant Soil 407, 67–79. https://doi.org/10.1007/s11104-016-2808-2

Hanson, P.J., Walker, A.P., 2019. Advancing global change biology through experimental

manipulations: Where have we been and where might we go? Glob. Chang. Biol. 26,

287–299. https://doi.org/10.1111/gcb.14894

Heisler-White, J.L., Blair, J.M., Kelly, E.F., Harmoney, K., Knapp, A.K., 2009. Contingent

productivity responses to more extreme rainfall regimes across a grassland biome. Glob.

Chang. Biol. 15, 2894–2904. https://doi.org/10.1111/j.1365-2486.2009.01961.x

Hoeppner, S.S., Dukes, J.S., 2012. Interactive responses of old-field plant growth and

composition to warming and precipitation. Glob. Chang. Biol. 18, 1754–1768.

https://doi.org/10.1111/j.1365-2486.2011.02626.x

Hofer, D., Suter, M., Buchmann, N., Lüscher, A., 2017. Nitrogen status of functionally

different forage species explains resistance to severe drought and post-drought

overcompensation. Agric. Ecosyst. Environ. 236, 312–322.

https://doi.org/10.1016/j.agee.2016.11.022

Hoover, D.L., Knapp, A.K., Smith, M.D., 2014. Resistance and resilience of a grassland

ecosystem to climate extremes. Ecology 95, 2646–2656. https://doi.org/10.1890/13-

2186.1

Huxman, T.E., Smith, M.D., Fay, P.A., Knapp, A.K., Shaw, M.R., Lolk, M.E., Smith, S.D.,

Tissue, D.T., Zak, J.C., Weltzin, J.F., Pockman, W.T., Sala, O.E., Haddad, B.M., Harte,

J., Koch, G.W., Schwinning, S., Small, E.E., Williams, D.G., 2004. Convergence across

biomes to a common rain-use efficiency. Nature 429, 651–654.

https://doi.org/10.1038/nature02561

Isbell, F.I., Craven, D., Connolly, J., Loreau, M., Schmid, B., Beierkuhnlein, C., Bezemer,

T.M., Bonin, C., Bruelheide, H., de Luca, E., Ebeling, A., Griffin, J.N., Guo, Q.,

Hautier, Y., Hector, A., Jentsch, A., Kreyling, J., Lanta, V., Manning, P., Meyer, S.T.,

Mori, A.S., Naeem, S., Niklaus, P.A., Polley, H.W., Reich, P.B., Roscher, C., Seabloom,

E.W., Smith, M.D., Thakur, M.P., Tilman, D.G., Tracy, B.F., van der Putten, W.H., van

Ruijven, J., Weigelt, A., Weisser, W.W., Wilsey, B.J., Eisenhauer, N., 2015.

Biodiversity increases the resistance of ecosystem productivity to climate extremes.

Nature 526, 574–577. https://doi.org/10.1038/nature15374

Jentsch, A., Beierkuhnlein, C., 2008. Research frontiers in climate change: Effects of extreme

meteorological events on ecosystems. Comptes Rendus - Geosci. 340, 621–628.

https://doi.org/10.1016/j.crte.2008.07.002

Jentsch, A., Kreyling, J., Boettcher-Treschkow, J., Beierkuhnlein, C., 2009. Beyond gradual

warming: Extreme weather events alter flower phenology of European grassland and

heath species. Glob. Chang. Biol. 15, 837–849. https://doi.org/10.1111/j.1365-

2486.2008.01690.x

Jiménez, M.A., Jaksic, F.M., Armesto, J.J., Gaxiola, A., Meserve, P.L., Kelt, D.A., Gutiérrez,

J.R., 2011. Extreme climatic events change the dynamics and invasibility of semi-arid

annual plant communities. Ecol. Lett. 14, 1227–1235. https://doi.org/10.1111/j.1461-

0248.2011.01693.x

Johnston, W.H., 1996. The place of C4 grasses in temperate pastures in Australia. New Zeal.

J. Agric. Res. 39, 527–540. https://doi.org/10.1080/00288233.1996.9513213

Kaine, G.W., Tozer, P.R., 2005. Stability, resilience and sustainability in pasture-based

grazing systems. Agric. Syst. 83, 27–48. https://doi.org/10.1016/j.agsy.2004.03.001

Keatinge, J.D.H., Qi, A., Wheeler, T.R., Ellis, R.H., Summerfield, R.J., 1998. Effects of

temperature and photoperiod on phenology as a guide to the selection of annual legume

cover and green manure crops for hillside farming systems. F. Crop. Res. 57, 139–152.

https://doi.org/10.1016/S0378-4290(97)00122-6

Knapp, A.K., Avolio, M.L., Beier, C., Carroll, C.J.W., Collins, S.L., Dukes, J.S., Fraser,

L.H., Griffin-Nolan, R.J., Hoover, D.L., Jentsch, A., Loik, M.E., Phillips, R.P., Post,

A.K., Sala, O.E., Slette, I.J., Yahdjian, L., Smith, M.D., 2017. Pushing precipitation to

the extremes in distributed experiments: recommendations for simulating wet and dry

years. Glob. Chang. Biol. 23, 1774–1782. https://doi.org/10.1111/gcb.13504

Knapp, A.K., Beier, C., Briske, D.D., Classen, A.T., Luo, Y., Reichstein, M., Smith, M.D.,

Smith, S.D., Bell, J.E., Fay, P.A., Heisler, J.L., Leavitt, S.W., Sherry, R.A., Smith, B.,

Weng, E., 2008. Consequences of more extreme precipitation regimes for terrestrial

ecosystems. Bioscience 58, 811. https://doi.org/10.1641/B580908

Knapp, A.K., Carroll, C.J.W., Denton, E.M., La Pierre, K.J., Collins, S.L., Smith, M.D.,

2015. Differential sensitivity to regional-scale drought in six central US grasslands.

Oecologia 177, 949–957. https://doi.org/10.1007/s00442-015-3233-6

Kragt, M.E., Pannell, D.J., Robertson, M.J., Thamo, T., 2012. Assessing costs of soil carbon

sequestration by crop-livestock farmers in Western Australia. Agric. Syst. 112, 27–37.

https://doi.org/10.1016/j.agsy.2012.06.005

Kreyling, J., Beierkuhnlein, C., Ellis, L., Jentsch, A., 2008a. Invasibility of grassland and

heath communities exposed to extreme weather events- additive effects of diversity

resistance and fluctuating physical environment. Oikos 117, 1542–1554.

https://doi.org/10.1111/j.2008.0030-1299.16653.x

Kreyling, J., Dengler, J., Walter, J., Velev, N., Ugurlu, E., Sopotlieva, D., Ransijn, J., Picon-

Cochard, C., Nijs, I., Hernandez, P., Güler, B., von Gillhaussen, P., De Boeck, H.J.,

Bloor, J.M.G., Berwaers, S., Beierkuhnlein, C., Arfin Khan, M.A.S., Apostolova, I.,

Altan, Y., Zeiter, M., Wellstein, C., Sternberg, M., Stampfli, A., Campetella, G., Bartha,

S., Bahn, M., Jentsch, A., 2017. Species richness effects on grassland recovery from

drought depend on community productivity in a multisite experiment. Ecol. Lett. 20,

1405–1413. https://doi.org/10.1111/ele.12848

Kreyling, J., Wenigmann, M., Beierkuhnlein, C., Jentsch, A., 2008b. Effects of extreme

weather events on plant productivity and tissue die-back are modified by community

composition. Ecosystems 11, 752–763. https://doi.org/10.1007/s10021-008-9157-9

Kueppers, L.M., Conlisk, E., Castanha, C., Moyes, A.B., Germino, M.J., de Valpine, P.,

Torn, M.S., Mitton, J.B., 2017. Warming and provenance limit tree recruitment across

and beyond the elevation range of subalpine forest. Glob. Chang. Biol. 23, 2383–2395.

https://doi.org/10.1111/gcb.13561

Kuster, T.M., Arend, M., Bleuler, P., Günthardt-Goerg, M.S., Schulin, R., 2013. Water

regime and growth of young oak stands subjected to air-warming and drought on two

different forest soils in a model ecosystem experiment. Plant Biol. 15, 138–147.

https://doi.org/10.1111/j.1438-8677.2011.00552.x

Length, R., 2019. emmeans: Estimated Marginal Means, aka Least-Squares Means.

Li, F., Kautz, T., Pude, R., Köpke, U., 2012. Nodulation of lucerne (*Medicago sativa* L.)

roots: Depth distribution and temporal variation. Plant, Soil Environ. 58, 424–428.

https://doi.org/10.17221/314/2012-pse

Liste, H.-H., White, J.C., 2008. Plant hydraulic lift of soil water – implications for crop

production and land restoration. Plant Soil 313, 1–17. https://doi.org/10.1007/s11104-

008-9696-z

Loi, A., Howieson, J.G., Nutt, B.J., Carr, S.J., 2005. A second generation of annual pasture

legumes and their potential for inclusion in Mediterranean-type farming systems. Aust.

J. Exp. Agric. 45, 289–299. https://doi.org/10.1071/ea03134

Lu, M., Zhou, X., Yang, Q., Li, H., Fang, C., Chen, J., Yang, X., Li, B., 2013. Responses of

ecosystem carbon cycle to experimental warming: a meta-analysis. Ecology 94, 726–

738.

Ma, X., Huete, A., Moran, S., Ponce-Campos, G., Eamus, D., 2015. Abrupt shifts in

phenology and vegetation productivity under climate extremes. J. Geophys. Res.

Biogeosciences 120, 2036–2052. https://doi.org/10.1002/2015JG003144

Maxwell, S.L., Butt, N., Maron, M., McAlpine, C.A., Chapman, S., Ullmann, A., Segan,

D.B., Watson, J.E.M., 2019. Conservation implications of ecological responses to

extreme weather and climate events. Divers. Distrib. 25, 613–625.

https://doi.org/10.1111/ddi.12878

McKeon, G.M., Stone, G.S., Syktus, J.I., Carter, J.O., Flood, N.R., Ahrens, D.G., Bruget,

D.N., Chilcott, C.R., Cobon, D.H., Cowley, R.A., Crimp, S.J., Fraser, G.W., Howden,

S.M., Johnston, P.W., Ryan, J.G., Stokes, C.J., Day, K.A., 2009. Climate change

impacts on northern Australian rangeland livestock carrying capacity: A review of

issues. Rangel. J. 31, 1–29. https://doi.org/10.1071/RJ08068

Mitchell, P.J., O'Grady, A.P., Pinkard, E.A., Brodribb, T.J., Arndt, S.K., Blackman, C.J.,

Duursma, R.A., Fensham, R.J., Hilbert, D.W., Nitschke, C.R., Norris, J., Roxburgh,

S.H., Ruthrof, K.X., Tissue, D.T., 2016. An ecoclimatic framework for evaluating the

resilience of vegetation to water deficit. Glob. Chang. Biol. 22, 1677–1689.

https://doi.org/10.1111/gcb.13177

Mori, A.S., Fujii, S., Kurokawa, H., 2014. Ecological consequences through responses of

plant and soil communities to changing winter climate. Ecol. Res. 29, 547–559.

https://doi.org/10.1007/s11284-013-1091-4

Munson, S.M., Long, A.L., 2017. Climate drives shifts in grass reproductive phenology

across the western USA. New Phytol. 213, 1945–1955.

https://doi.org/10.1111/nph.14327

Murray-Tortarolo, G.N., Jaramillo, V.J., 2020. Precipitation extremes in recent decades

impact cattle populations at the global and national scales. Sci. Total Environ. 736,

139557. https://doi.org/10.1016/j.scitotenv.2020.139557

Murray-Tortarolo, G.N., Jaramillo, V.J., 2019. The impact of extreme weather events on

livestock populations: the case of the 2011 drought in Mexico. Clim. Change 153, 79–

89. https://doi.org/10.1007/s10584-019-02373-1

Nábrádi, A., 2016. The economic value of grassland products. Appl. Stud. Agribus. Commer.
1, 19–28. https://doi.org/10.19041/apstract/2007/1/2

Naudts, K., Van den Berge, J., Janssens, I.A., Nijs, I., Ceulemans, R., 2011. Does an extreme
drought event alter the response of grassland communities to a changing climate?
Environ. Exp. Bot. 70, 151–157. https://doi.org/10.1016/j.envexpbot.2010.08.013

Nicholas, P.K., Kemp, P.D., Barker, D.J., 2004. Stress and recovery of hill pastures in the
North Island of New Zealand. Grass Forage Sci. 59, 250–263.

Nie, Z., Mitchell, M., Clark, S., Smith, K., 2015. Distribution of lucerne roots in summer-dry
environments of southern Australia, in: Proceedings of the 17th ASA Conference.
Hobart, Australia, pp. 1–4.

Nijp, J.J., Limpens, J., Metselaar, K., Peichl, M., Nilsson, M.B., van der Zee, S.E.A.T.M.,
Berendse, F., 2015. Rain events decrease boreal peatland net CO_2 uptake through
reduced light availability. Glob. Chang. Biol. 21, 2309–2320.
https://doi.org/10.1111/gcb.12864

Niu, S., Luo, Y., Li, D., Cao, S., Xia, J., Li, J., Smith, M.D., 2014. Plant growth and mortality
under climatic extremes: An overview. Environ. Exp. Bot. 98, 13–19.
https://doi.org/10.1016/j.envexpbot.2013.10.004

Novem Auyeung, D.S., Suseela, V., Dukes, J.S., 2013. Warming and drought reduce
temperature sensitivity of nitrogen transformations. Glob. Chang. Biol. 19, 662–676.
https://doi.org/10.1111/gcb.12063

Nyfeler, D., Huguenin-Elie, O., Suter, M., Frossard, E., Lüscher, A., 2011. Grass-legume
mixtures can yield more nitrogen than legume pure stands due to mutual stimulation of
nitrogen uptake from symbiotic and non-symbiotic sources. Agric. Ecosyst. Environ.
140, 155–163. https://doi.org/10.1016/j.agee.2010.11.022

O'Mara, F.P., 2012. The role of grasslands in food security and climate change. Ann. Bot.
110, 1263–1270. https://doi.org/10.1093/aob/mcs209

Pang, J., Wang, Y., Lambers, H., Tibbett, M., Siddique, K.H.M., Ryan, M.H., 2013.
Commensalism in an agroecosystem: Hydraulic redistribution by deep-rooted legumes
improves survival of a droughted shallow-rooted legume companion. Physiol. Plant.
149, 79–90. https://doi.org/10.1111/ppl.12020

Parmesan, C., Root, T.L., Willig, M.R., 2000. Impacts of Etreme Weather and Climate on
Terrestrial Biota. Bull. Am. Meteorol. Soc. 81, 443–450.

Pearcy, R.W., Caldwell, M.M., Keeley, J.E., Monson, R.K., Strain, B.R., 1987. Carbon gain
by plants in natural environments. Bioscience 37, 21–29.

https://doi.org/10.2307/1310174

Pearcy, R.W., Ehleringer, J., 1984. Comparative ecophysiology of C3 and C4 plants. Plant.

Cell Environ. 7, 1–13. https://doi.org/10.1111/j.1365-3040.1984.tb01194.x

Peng, A., Klanderud, K., Wang, G., Zhang, L., Xiao, Y., Yang, Y., 2020. Plant community

responses to warming modified by soil moisture in the Tibetan Plateau. Arctic, Antarct.

Alp. Res. 52, 60–69. https://doi.org/10.1080/15230430.2020.1712875

Piao, S., Liu, Q., Chen, A., Janssens, I.A., Fu, Y., Dai, J., Liu, L., Lian, X., Shen, M., Zhu,

X., 2019. Plant phenology and global climate change: Current progresses and

challenges. Glob. Chang. Biol. 25, 1922–1940. https://doi.org/10.1111/gcb.14619

Pimm, L.S., 1984. The complexity and stability of ecosystems. Nature 307, 321–326.

Power, S.A., Barnett, K.L., Ochoa-hueso, R., Facey, S.L., Gibson-Forty, E.V.J., Hartley,

S.E., Nielsen, U.N., Tissue, D.T., Johnson, S.N., 2016. DRI-Grass: A new experimental

platform for addressing grassland ecosystem responses to future precipitation scenarios

in South-East Australia. Front. Plant Sci. 7, 1–14.

https://doi.org/10.3389/fpls.2016.01373

R Core Team, 2020. R: A language and environment for statistical computing.

Raza, A., Friedel, J.K., Moghaddam, A., Ardakani, M.R., Loiskandl, W., Himmelbauer, M.,

Bodner, G., 2013. Modeling growth of different lucerne cultivars and their effect on soil

water dynamics. Agric. Water Manag. 119, 100–110.

https://doi.org/10.1016/j.agwat.2012.12.006

Reyer, C.P.O., Leuzinger, S., Rammig, A., Wolf, A., Bartholomeus, R.P., Bonfante, A., de

Lorenzi, F., Dury, M., Gloning, P., Abou Jaoudé, R., Klein, T., Kuster, T.M., Martins,

M., Niedrist, G., Riccardi, M., Wohlfahrt, G., de Angelis, P., de Dato, G., François, L.,

Menzel, A., Pereira, M., 2013. A plant's perspective of extremes: Terrestrial plant

responses to changing climatic variability. Glob. Chang. Biol. 19, 75–89.

https://doi.org/10.1111/gcb.12023

Ruppert, J.C., Harmoney, K., Henkin, Z., Snyman, H.A., Sternberg, M., Willms, W.,

Linstädter, A., 2015. Quantifying drylands' drought resistance and recovery: The

importance of drought intensity, dominant life history and grazing regime. Glob. Chang.

Biol. 21, 1258–1270. https://doi.org/10.1111/gcb.12777

Sage, R.F., 2019. Global Change Biology: A Primer. Glob. Chang. Biol. gcb.14893.

https://doi.org/10.1111/gcb.14893

Sanz-Sáez, Á., Erice, G., Aguirreolea, J., Irigoyen, J.J., Sánchez-Díaz, M., 2012. Alfalfa yield

under elevated CO_2 and temperature depends on the Sinorhizobium strain and growth

season. Environ. Exp. Bot. 77, 267–273.

https://doi.org/10.1016/j.envexpbot.2011.11.017

Sherry, R.A., Weng, E., Arnone, J.A., Johnson, D.W., Schimel, D.S., Verburg, P.S., Wallace,

L.L., Luo, Y., 2008. Lagged effects of experimental warming and doubled precipitation

on annual and seasonal aboveground biomass production in a tallgrass prairie. Glob.

Chang. Biol. 14, 2923–2936. https://doi.org/10.1111/j.1365-2486.2008.01703.x

Slette, I.J., Post, A.K., Awad, M., Even, T., Punzalan, A., Williams, S., Smith, M.D., Knapp,

A.K., 2019. How ecologists define drought, and why we should do better. Glob. Chang.

Biol. 25, 3193–3200. https://doi.org/10.1111/gcb.14747

Smith, M.D., 2011. An ecological perspective on extreme climatic events: A synthetic

definition and framework to guide future research. J. Ecol. 99, 656–663.

https://doi.org/10.1111/j.1365-2745.2011.01798.x

Smith, M.D., Wilcox, K.R., Power, S.A., Tissue, D.T., Knapp, A.K., 2017. Assessing

community and ecosystem sensitivity to climate change – toward a more comparative

approach. J. Veg. Sci. 28, 235–237. https://doi.org/10.1111/jvs.12524

Song, J., Wan, S., Piao, S., Knapp, A.K., Classen, A.T., Vicca, S., Ciais, P., Hovenden, M.J.,

Leuzinger, S., Beier, C., Kardol, P., Xia, J., Liu, Q., Ru, J., Zhou, Z., Luo, Y., Guo, D.,

Adam Langley, J., Zscheischler, J., Dukes, J.S., Tang, J., Chen, J., Hofmockel, K.S.,

Kueppers, L.M., Rustad, L., Liu, L., Smith, M.D., Templer, P.H., Quinn Thomas, R.,

Norby, R.J., Phillips, R.P., Niu, S., Fatichi, S., Wang, Y., Shao, P., Han, H., Wang, D.,

Lei, L., Wang, J., Li, Xiaona, Zhang, Q., Li, Xiaoming, Su, F., Liu, B., Yang, F., Ma, G.,

Li, G., Liu, Yanchun, Liu, Yinzhan, Yang, Z., Zhang, K., Miao, Y., Hu, M., Yan, C.,

Zhang, A., Zhong, M., Hui, Y., Li, Y., Zheng, M., 2019. A meta-analysis of 1,119

manipulative experiments on terrestrial carbon-cycling responses to global change. Nat.

Ecol. Evol. 3, 1309–1320. https://doi.org/10.1038/s41559-019-0958-3

Steffen, W., Richardson, K., Rockström, J., Cornell, S., Fetzer, I., Bennett, E., Biggs, R.,

Carpenter, S.R., de Wit, C. a., Folke, C., Mace, G.M., Persson, L.M., Veerabhadran, R.,

Reyers, B., Sörlin, S., 2015. Planetary Boundaries: Guiding human development on a

changing planet. Science (80-.). 347, 1259855. https://doi.org/10.1126/science.1259855

Suter, M., Connolly, J., Finn, J.A., Loges, R., Kirwan, L., Sebastià, M.T., Lüscher, A., 2015.

Nitrogen yield advantage from grass-legume mixtures is robust over a wide range of

legume proportions and environmental conditions. Glob. Chang. Biol. 21, 2424–2438.

https://doi.org/10.1111/gcb.12880

Vitasse, Y., Bottero, A., Cailleret, M., Bigler, C., Fonti, P., Gessler, A., Lévesque, M.,

1053  Rohner, B., Weber, P., Rigling, A., Wohlgemuth, T., 2019. Contrasting resistance and

1054    resilience to extreme drought and late spring frost in five major European tree species.

1055    Glob. Chang. Biol. 25, 3781–3792. https://doi.org/10.1111/gcb.14803

1056  Vogel, A., Scherer-Lorenzen, M., Weigelt, A., 2012. Grassland resistance and resilience after

1057    drought depends on management intensity and species richness. PLoS One 7.

1058    https://doi.org/10.1371/journal.pone.0036992

1059  Volder, A., Briske, D.D., Tjoelker, M.G., 2013. Climate warming and precipitation

1060    redistribution modify tree-grass interactions and tree species establishment in a warm-

1061    temperate savanna. Glob. Chang. Biol. 19, 843–857. https://doi.org/10.1111/gcb.12068

1062  Walker, M.D., Wahren, C.-H., Hollister, R.D., Henry, G.H.R., Ahlquist, L.E., Alatalo, J.M.,

1063    Bret-Harte, M.S., Calef, M.P., Callaghan, T. V., Carroll, A.B., Epstein, H.E., Jónsdóttir,

1064    I.S., Klein, J.A., Magnússon, B., Molau, U., Oberbauer, S.F., Rewa, S.P., Robinson,

1065    C.H., Shaver, G.R., Suding, K.N., Thompson, C.C., Tolvanen, A., Totland, Ø., Turner,

1066    P.L., Tweedie, C.E., Webber, P.J., Wookey, P.A., 2006. Plant community responses to

1067    experimental warming across the tundra biome. Proc. Natl. Acad. Sci. U. S. A. 103,

1068    1342–6. https://doi.org/10.1073/pnas.0503198103

1069  Whittington, H.R., Deede, L., Powers, J.S., 2012. Growth responses, biomass partitioning,

1070    nitrogen isotopes of prairie legumes in response to elevated temperature and varying

1071    nitrogen source in a growth chamber experiment. Am. J. Bot. 99, 838–846.

1072    https://doi.org/10.3732/ajb.1100283

1073  Whittington, H.R., Tilman, D., Powers, J.S., 2013. Consequences of elevated temperatures on

1074    legume biomass and nitrogen cycling in a field warming and biodiversity experiment in

1075    a North American prairie. Funct. Plant Biol. 40, 1147–1158.

1076    https://doi.org/10.1071/FP12345

1077  Wilcox, K.R., Blair, J.M., Smith, M.D., Knapp, A.K., 2016. Does ecosystem sensitivity to

1078    precipitation at the site- level conform to regional- scale predictions? Ecology 97, 561–

1079    568. https://doi.org/10.1007/s13398-014-0173-7.2

1080  Wilcox, K.R., Shi, Z., Gherardi, L.A., Lemoine, N.P., Koerner, S.E., Hoover, D.L., Bork, E.,

1081    Byrne, K.M., Cahill, J., Collins, S.L., Evans, S., Gilgen, A.K., Holub, P., Jiang, L.,

1082    Knapp, A.K., LeCain, D., Liang, J., Garcia-Palacios, P., Peñuelas, J., Pockman, W.T.,

1083    Smith, M.D., Sun, S., White, S.R., Yahdjian, L., Zhu, K., Luo, Y., 2017. Asymmetric

1084    responses of primary productivity to precipitation extremes: A synthesis of grassland

1085    precipitation manipulation experiments. Glob. Chang. Biol. 23, 4376–4385.

1086    https://doi.org/10.1111/gcb.13706

Wu, Z., Dijkstra, P., Koch, G.W., Peñuelas, J., Hungate, B.A., 2011. Responses of terrestrial

ecosystems to temperature and precipitation change: A meta-analysis of experimental

manipulation. Glob. Chang. Biol. 17, 927–942. https://doi.org/10.1111/j.1365-

2486.2010.02302.x

Yao, X., Wu, J., Gong, X., 2019. Precipitation and seasonality affect grazing impacts on

herbage nutritive values in alpine meadows on the Qinghai-Tibet Plateau. J. Plant Ecol.

12, 993–1008. https://doi.org/10.1093/jpe/rtz027

Yuan, W., Cai, W., Chen, Y., Liu, Shuguang, Dong, W., Zhang, H., Yu, G., Chen, Z., He, H.,

Guo, W., Liu, D., Liu, Shaoming, Xiang, W., Xie, Z., Zhao, Z., Zhou, G., 2016. Severe

summer heatwave and drought strongly reduced carbon uptake in Southern China. Sci.

Rep. 6, 1–12. https://doi.org/10.1038/srep18813

1098

1099

Figure Captions

Figure 1. The Pastures and Climate Extremes (PACE) field facility located at Western Sydney University in Richmond, NSW Australia. A) There were six open-sided polytunnels, each with eight experimental plots; B) Experimental plots (4 x 4 m) received a drought, warming, or drought + warming treatment and were each divided into four discrete 2 x 2 m subplots comprising nine different pasture species in monoculture or three select two-species mixtures; C) Heater (infra-red: IR) arrays were mounted above the vegetation canopy and warm the plot surface an average of 3 °C above paired ambient temperature plots.

Figure 2. Experimental treatment effects of winter/spring drought (control: C, drought: D) applied during the period between 1 June to 30 November 2018 and year-round warming (ambient temperature: aT, elevated temperature: eT) treatments on soil moisture (panels A and B) and plot surface temperature (panels C and D) averaged across the six shelters, from 1 June 2018 to 31 May 2019. A) Average soil moisture content in *Festuca* subplots (additional species are shown in Figure S2) with 95% confidence intervals as well as individual irrigation events as daily rainfall totals for control (black) and droughted (red) plots over the entire year; B) Treatment differences in soil moisture from aT-C plots over time in the *Festuca* subplots, noting the winter/spring drought treatment period between 1 June and 30 November 2018 and subsequent summer/autumn without drought treatment; C) Daily maximum plot surface temperature, relative to maximum air temperature (black line) and 40 °C (representing extreme temperatures, dashed lines); D) Daily minimum temperature compared with minimum air temperature (black line) and 0 °C (dashed lines).

Figure 3. A) Aboveground production summed for all harvests during the 6-month winter/spring drought period. Values shown are means ± 1 standard error (control = solid symbol, droughted = outline); B) drought effect size (log response ratio of drought vs control production during the 6-month drought treatment period), and C) drought sensitivity (yield reduction in the drought treatment relative to controls per mm irrigation withheld, to account for varying lengths of drought treatments among planting types, see Table 1). Values for panels B and C are mean values with 95% confidence intervals and same letter designations indicate non-significant differences among plantings. Abbreviations are as follows: *Biserrula* (Bis), *Chloris* (Chl), *Digitaria* (Dig), *Digitaria & Biserrula* mixture (DigBis), *Festuca* (Fes), *Lolium* (Lol), *Medicago* (Med), *Phalaris* (Pha), *Phalaris & Trifolium* mixture (PhaTri), *Rytidosperma* (Ryt), *Themeda* (The), *Themeda & Rytidosperma* mixture (TheRyt). Significant pairwise comparisons for the effect of drought treatment are indicated as follows: NS= not significant, # = $P < 0.1$, * $P < 0.05$, ** $P = <0.01$, *** $P = < 0.001$.

Figure 4. A) Aboveground production summed for all harvests during the 6-month summer/autumn recovery period following the winter/spring drought treatment (control = solid symbol, formerly droughted = outline), B) effect size of summer/autumn legacy effects of winter/spring drought on productivity (shown as log response ratios), C) annual aboveground production summed for all harvests June 2018-May 2019 and D) effect size (log response ratios) of winter/spring drought on total annual productivity (note that for panels C and D annual species are not included, as annual totals are the same as winter/spring values). Panels B and D are back-transformed, following analysis on log-transformed data. All symbols and abbreviations follow Figure 3.

Figure 5. Aboveground production for individual plantings exposed to drought (control: C and droughted: D) and warming (ambient: aT and elevated: eT) treatments during A) the 6-month winter/spring drought period (June 1st- Nov 30th 2018), B) the subsequent

1150   summer/autumn recovery period (Dec 1^{st} 2018- May 31^{st} 2019), and C) annual production.
1151   Drt and W indicate significant drought and warming effects, respectively. Values are means $\pm$
1152   1 SE and all spp. abbreviations and significance levels follow Fig. 3.
1153
Figure 6. Treatment effect sizes during the winter/spring drought period for A) drought under
ambient (aT) and elevated (eT) temperatures, B) warming, under control (C) and droughted
(D) conditions, and C) the effect of warming on biomass responses to drought (for panel C,
positive values indicate a reduction in drought impact under warming) among plantings
exposed to both drought and warming. Values shown are means with 95% confidence
intervals and all spp. abbreviations follow Fig 3.

Figure 1. The Pastures and Climate Extremes (PACE) field facility located at Western Sydney University in Richmond, NSW Australia. A) There were six open-sided polytunnels, each with eight experimental plots; B) Experimental plots (4 x 4 m) received a drought, warming, or drought + warming treatment and were each divided into four discrete 2 x 2 m subplots comprising nine different pasture species in monoculture or three select two-species mixtures; C) Heater (infra-red: IR) arrays were mounted above the vegetation canopy and warm the plot surface an average of 3 °C above paired ambient temperature plots

1160

Figure 2. Experimental treatment effects of winter/spring drought (control: C, drought: D) applied during the period between 1 June to 30 November 2018 and year-round warming (ambient temperature: aT, elevated temperature: eT) treatments on soil moisture (panels A and B) and plot surface temperature (panels C and D) averaged across the six shelters, from 1 June 2018 to 31 May 2019. A) Average soil moisture content in *Festuca* subplots (additional species are shown in Figure S2) with 95% confidence intervals as well as individual irrigation events as daily rainfall totals for control (black) and droughted (red) plots over the entire year; B) Treatment differences in soil moisture from aT-C plots over time in the *Festuca* subplots, noting the winter/spring drought treatment period between 1 June and 30 November 2018 and subsequent summer/autumn without drought treatment; C) Daily maximum plot surface temperature, relative to maximum air temperature (black line) and 40 °C (representing extreme temperatures, dashed lines); D) Daily minimum temperature compared with minimum air temperature (black line) and 0 °C (dashed lines).

Figure 3. A) Aboveground production summed for all harvests during the 6-month winter/spring drought period. Values shown are means ± 1 standard error (control = solid symbol, droughted = outline); B) drought effect size (log response ratio of drought vs control production during the 6-month drought treatment period), and C) drought sensitivity (yield reduction in the drought treatment relative to controls per mm irrigation withheld, to account for varying lengths of drought treatments among planting types, see Table 1). Values for panels B and C are mean values with 95% confidence intervals and same letter designations indicate non-significant differences among plantings. Abbreviations are as follows: *Biserrula* (Bis), *Chloris* (Chl), *Digitaria* (Dig), *Digitaria & Biserrula* mixture (DigBis), *Festuca* (Fes), *Lolium* (Lol), *Medicago* (Med), *Phalaris* (Pha), *Phalaris & Trifolium* mixture (PhaTri), *Rytidosperma* (Ryt), *Themeda* (The), *Themeda & Rytidosperma* mixture (TheRyt). Significant pairwise comparisons for the effect of drought treatment are indicated as follows: NS= not significant, # = $P < 0.1$, * $P < 0.05$, ** $P = <0.01$, *** $P = < 0.001$.

Figure 4. A) Aboveground production summed for all harvests during the 6-month summer/autumn recovery period following the winter/spring drought treatment (control = solid symbol, formerly droughted = outline), B) effect size of summer/autumn legacy effects of winter/spring drought on productivity (shown as log response ratios), C) annual aboveground production summed for all harvests June 2018-May 2019 and D) effect size (log response ratios) of winter/spring drought on total annual productivity (note that for panels C and D annual species are not included, as annual totals are the same as winter/spring values). Panels B and D are back-transformed, following analysis on log-transformed data. All symbols and abbreviations follow Figure 3.

Figure 5. Aboveground production for individual plantings exposed to drought (control: C and droughted: D) and warming (ambient: aT and elevated: eT) treatments during A) the 6-month winter/spring drought period (June 1st-Nov 30th 2018), B) the subsequent summer/autumn recovery period (Dec 1st 2018-May 31st 2019), and C) annual production. Drt and W indicate significant drought and warming effects, respectively. Values are means ± 1 SE and all spp. abbreviations and significance levels follow Fig. 3.

Figure 6. Treatment effect sizes among plantings exposed to both drought and warming during the winter/spring drought period for A) drought under ambient (aT) and elevated (eT) temperatures, B) warming under control (C) and drought (D) conditions and C) the effect of warming on biomass responses to rainfall (positive values indicate a reduction in drought impacts under warming). Values shown are means with 95% confidence intervals and all spp. abbreviations follow Fig 3.