Pastures and Climate Extremes: Impacts of cool season warming and drought on the

productivity of key pasture species in a field experiment

Amber C. Churchill*[a,1], Haiyang Zhang[a], Kathryn J. Fuller[a], Burhan Amiji[a], Ian C.

Anderson[a], Craig V. M. Barton[a], Yolima Carrillo[a], Karen L. M. Catunda[a], Manjunatha H.

Chandregowda[a], Chioma Igwenagu[a], Vinod Jacob[a], Gil Won Kim[a,b] Catriona A. Macdonald[a],

Belinda E. Medlyn[a], Ben D. Moore[a], Elise Pendall[a], Jonathan M. Plett[a], Alison K. Post[a,c], Jeff

R. Powell[a], David T. Tissue[a,d], Mark G. Tjoelker[a], Sally A. Power[a]

[a]Hawkesbury Institute for the Environment, Western Sydney University, Richmond, NSW

Australia

[b]Institute of Agriculture and Life Sciences, Gyeongsang National University, Jinju, South

Korea

[c]The Center for Ecosystem Science and Society, Northern Arizona University, Flagstaff, AZ,

USA

[d]Global Centre for Land-based Innovation, Western Sydney University, Hawkesbury

Campus, Richmond, NSW, Australia

[1]Present address: Department of Ecology, Evolutionary Biology and Behavior

University of Minnesota, St. Paul, MN, USA

*Correspondence: Corresponding author achurchi@umn.edu

Abstract

Shifts in the timing, intensity and/or frequency of climate extremes, such as severe drought

and heatwaves, can generate sustained shifts in ecosystem function with important ecological

and economic impacts for rangelands and managed pastures. The Pastures and Climate

Extremes experiment (PACE) in Southeast Australia was designed to investigate the impacts

of a severe winter/spring drought (60% rainfall reduction) and, for a subset of species, a

factorial combination of drought and elevated temperature (ambient +3 °C) on pasture

productivity. The experiment included nine common pasture and Australian rangeland

species from three plant functional groups (C_3 grasses, C_4 grasses and legumes) planted in

monoculture. Winter/spring drought resulted in productivity declines of 45% on average and

up to 74% for the most affected species (*Digitaria eriantha*) during the 6-month treatment

period, with eight of the nine species exhibiting significant yield reductions. Despite

considerable variation in species' sensitivity to drought, C_4 grasses were more strongly

affected by this treatment than C_3 grasses or legumes. Warming also had negative effects on

cool-season productivity, associated at least partially with exceedance of optimum growth

temperatures in spring and indirect effects on soil water content. The combination of

winter/spring drought and year-round warming resulted in the greatest yield reductions. We

identified responses that were either additive such that there was only as significant warming

effect under drought (*Festuca*), or less-than-additive, where there was no drought effect under

warming (*Medicago*), compared to ambient plots. Results from this study highlight the

sensitivity of diverse pasture species to increases in winter and spring drought severity

similar to those predicted for this region, and that anticipated benefits of cool-season

warming are unlikely to be realised. Overall, the substantial negative impacts on productivity

suggest that future, warmer, drier climates will result in shortfalls in cool-season forage

availability, with profound implications for the livestock industry and natural grazer

communities.

1 Introduction

Climate change is a dominant driver of ecosystem change across the globe (Steffen et al., 2015; Sage, 2019). Exposure to high temperatures and changes in rainfall regimes have been shown to disrupt physiological function and alter plant species' interactions, ultimately driving changes in productivity and ecological processes such as nutrient and water cycling (Parmesan et al., 2000; Backhaus et al., 2014; Frank et al., 2015; Ma et al., 2015). Predicting the impact of climate change on these processes is challenging, as temperature profiles (including minimum and maximum values) and the frequency, timing and size of rainfall events play major roles in driving change in ecosystem function, whereas climate models and projections often focus on only changes in mean annual temperature or precipitation (Easterling et al., 2000; Kreyling et al., 2008; Jentsch et al., 2009). Despite a recent increase in the number of studies focusing on climate extremes (De Boeck et al., 2015, 2019; Knapp et al., 2017; Hanson and Walker, 2019), relatively few have addressed the ecological implications of seasonal shifts in climate, which is important for understanding the underlying trade-offs between plant phenology and associated plant functional group responses (i.e. C_3 vs. C_4, legumes; Beier et al., 2012). This is especially true for studies considering the impact of multiple climate variables simultaneously as the combination may generate contrasting predictions for different plant functional groups (Beier et al., 2012).

Shifts in the seasonality of precipitation associated with climate change are predicted for many terrestrial ecosystems (IPCC, 2021). These changes are likely to have serious economic ramifications in biomes where productivity is tightly coupled with seasonal rainfall patterns (O'Mara, 2012; Godde et al., 2020), especially in warmer regions where the absence of low-temperature constraints allows year-round plant growth that depends on seasonal rainfall (Arredondo et al., 2016; Zeiter et al., 2016). Grassland phenology is often predicted based on

classifications of plant traits or functional groups, with different optimum temperatures for

photosynthesis (C_3 vs. C_4 grasses; Munson and Long, 2017; Ode and Tieszen, 1980; Winslow

et al., 2003), timing of flowering in relation to peak summer temperatures (Sherry et al.,

2007), life history strategies (annual vs. perennial; Cleland et al., 2006; Enloe et al., 2004;

Veenendaal et al., 1996) and life forms (grasses/legumes/forbs etc.; König et al., 2018; Lesica

and Kittelson, 2010; Rathcke and Lacey, 1985). In temperate and subtropical climates,

grasslands dominated by C_3 grasses mainly accrue biomass during cooler periods of the year

(i.e. winter or spring), while C_4-dominated grasslands typically accumulate biomass later in

the growing season, when temperatures are higher (Pearcy and Ehleringer, 1984; Pearcy et

al., 1987; Adams et al., 2016; Wilcox et al., 2016). These broad patterns are based on key

differences in optimum temperatures for photosynthesis (**Figure 1**), as well as differences in

water-use-efficiency and strategies for nutrient acquisition. Field-based manipulations

examining impacts of drought on grassland production have primarily been conducted at the

community scale, where both direct and indirect (inter-specific competition) responses

combine to determine community-level productivity. For pasture systems, however, it is

important to evaluate climate impacts for a wide range of forage species grown in

monoculture in order to predict productivity responses and capacity to expand into new

regions or vulnerability within current ranges as local climate shifts (Johnston, 1996; Bindi

and Olesen, 2011).

In addition to changing rainfall regimes, rising temperatures are expected to have direct and

indirect impacts on many ecosystem processes (Bai et al., 2013; Novem Auyeung et al.,

2013; Song et al., 2019). In temperate climates, direct effects range from shifts in the timing

of peak productivity relative to the start of the growing season (typically earlier) to extending

the length of the growing season (Keatinge et al., 1998; Walker et al., 2006). The majority of

field-based temperature manipulations have been conducted in temperate or cold-climate

ecosystems, where low temperatures constrain growth for part of the year. These studies

often find increased productivity and shifts in growth phenology, notably growth starting

earlier and/or continuing later into the year, associated with warming (Aerts et al., 2006;

Kueppers et al., 2017; Lu et al., 2013; Walker et al., 2006; **Figure 1**), although there are

exceptions (Dukes et al., 2005; Arnone et al., 2011; Deutsch et al., 2011). Warming

experiments in warm-temperate or subtropical climates with comparatively short mild winters

and hot summers are relatively rare. These are, however, needed to evaluate the effects of

rising temperatures in circumstances where warming results in exceedance of thermal optima

for growth (Dukes et al., 2005; Volder et al., 2013; Song et al., 2019) and for regions where

forage species grow throughout the year.

Plant functional groups are predicted to differ in their responses to warming, with warm-

season species such as tropical grasses likely to benefit from increased winter or spring

temperatures, while temperate grasses may be particularly vulnerable to higher spring or

summer temperatures, especially when soil moisture is limiting (Munson and Long, 2017;

Figure 1). Legume responses to warming are also likely to vary between seasons. For

example, as with grasses, warming during the cooler months can bring temperatures closer to

thermal optima (Sanz-Sáez et al., 2012; Whittington et al., 2013; Peng et al., 2020) ultimately

promoting greater production (**Figure 1**). However, higher late spring and summer

temperatures may result in growth reductions as a consequence of thermal constraints on

nitrogen fixation and increased respiratory carbon losses or reduced photosynthesis if

temperatures exceed temperature optima (Aranjuelo et al., 2007; Whittington et al., 2012).

The impact of warmer temperatures on the timing and amount of biomass production in

different species across seasons is an important knowledge gap, both in pastures and native

grasslands and rangelands.

While studies that address the impact of either precipitation or temperature on plant

physiology and ecosystem function are valuable, warmer temperatures frequently co-occur

with drought (De Boeck et al., 2010; Dai, 2011; Yuan et al., 2016; Boer et al., 2020).

Consequently, it is necessary to evaluate the impacts of co-occurring climate stressors on

plant community structure and ecosystem function (Sherry et al., 2008; Hoeppner and Dukes,

2012). Climate models predict that temperate and sub-tropical Australia will be subject to

large reductions in winter and spring rainfall (CSIRO, 2020) and grasslands across the globe

are expected to experience more frequent and severe drought (Wang et al., 2021). Given the

importance of pastures and rangelands for maintaining food security (O'Mara, 2012; Nábrádi,

2016; Godde et al., 2020), we established a large-scale, field experiment (at the **PA**stures and

Climate **E**xtremes- PACE- field facility) to evaluate plant species' responses to winter/spring

rainfall reduction and year-round warming. Specifically, we asked: 1) How does

winter/spring drought affect productivity in a range of pasture species? 2) Do drought

responses in pasture grasses depend on their functional group? 3) Does cool-season warming

enhance productivity and/or exacerbate the impacts of drought in the considered species?

2 Methods

2.1 Site description

The PACE facility was constructed in 2017 at the Hawkesbury Campus of Western Sydney

University, in Richmond, New South Wales, Australia (S33.60972, E150.73833, elevation 25

m). Mean annual precipitation at this location is 800 mm (Australian Government Bureau of

Meteorology, Richmond - UWS Hawkesbury Station 1980-2010); however, there is large

inter-annual variability (annual precipitation 500-1400 mm over the past 30 years). Winter

and spring precipitation accounts for 40% of annual rainfall. Mean annual temperature is 17.2

°C, with the monthly maximum and minimum occurring in January (22.9° C) and July (10.2°

C), respectively (BOM, 2020). The site is fenced to prevent access by mammalian herbivores.

The soil is a loamy sand with a volumetric water holding capacity of 15-20% (**Table S1**).

The field facility comprises six replicate polytunnel shelters constructed from galvanized

steel frames. These are covered with a single layer of 180 µm polyethylene (Argosee,

Australia) to intercept all ambient precipitation, although the long sides are open to a height

of 1.5 m to allow free flow of air (**Figure 1**). Shelters are 48 m long by 8 m wide, with a

maximum height of 4.6 m, and are oriented along a SW-NE axis with the open ends facing

into the direction of the prevailing wind. Each shelter has eight treatment plots (4 m by 4 m;

Figure 1) that are further subdivided into four subplots (**Figure 1B; Figure S1**), with

different plant species assigned at the subplot level (total 192 subplots). All surface soils were

rotary-tilled to a depth of 12 cm to homogenize the upper soil profile prior to pasture

establishment. All 4 x 4 m plots have a full root barrier installed to a depth of 90 cm to ensure

hydrological isolation between treatments; the nested 2 x 2 m subplots have an additional root

barrier between them to a depth of 30 cm to minimise root ingress.

Nine plant species were grown in monoculture subplots, along with three sets of two-species

mixtures, for a total of twelve different planting combinations replicated in six independent

shelters. However, for the purposes of this study, we focused on the nine monoculture species

only. Species encompassed a range of functional diversity (C_3 and C_4 grasses, legumes;

annuals and perennials) and origins (native grasses, and tropical or temperate introduced

pasture grasses and legumes; **Table 1**) that are commonly found in improved grasslands

(pastures) or rangelands (Clements et al., 2003). Sward establishment was initiated during

early spring 2017 and yield data for the pilot year (2018-2019) are included in the supplement

(**Table S2**). Initial sowing included a fertilizer addition in the form of diammonium

phosphate (110 kg ha^{-1}), and swards were subsequently managed via hand-weeding, herbicide

and, where needed, insecticide application to maintain target species, in line with industry

practice. Subplots with legumes received appropriate rhizobium inoculant during sward

establishment: ALOSCA granular inoculant for *Biserrula* subplots (Group BS; ALOSCA

Technologies, Western Australia, Australia); Easy Rhiz soluble legume inoculant and

protecting agent for *Medicago* subplots (Group AL; New Edge Microbials, New South

Wales, Australia). Subplots received top-up fertilization seasonally to replace nutrients

removed from the soil (55 kg ha^{-1}; Cal-Gran Aftergraze, Incitec Pivot Fertilisers, Australia).

2.2 Experimental treatments

All nine species were exposed to a winter/spring drought treatment, and a subset of two

species (*Festuca*, *Medicago*) received a warming treatment in a factorial combination with

drought (**Figure S1**). The drought treatment comprised a control (Control: C) and a drought

(Drought: D) watering regime that was applied during the 6-month austral winter/spring

period (1 June to 30 November 2019). A 60% reduction in winter/spring rainfall was chosen

for the drought manipulation as representing the upper end of climate model predictions for

end-of-century seasonal rainfall change for south-eastern Australia, under RCP8.5 (CSIRO,

2020). Furthermore, a 60% reduction in winter/spring rainfall aligns with historical climate

extremes for key pasture growing regions across south-eastern Australia (BOM, 2019); this

treatment therefore represented historically relevant rainfall extremes, which are predicted to

increase in both frequency and duration. The control was set to represent a typical

precipitation regime for the local area, accounting for long-term patterns in seasonality and in

the statistical distribution of event sizes and timing within seasons (**Figure S2**).

The warming treatment comprised a year-round temperature increase of +3 °C (**Table 1**)

achieved using infra-red (IR) heaters, approximating predicted changes in temperature for

Australia by 2080 under RCP7.0 (IPCC, 2017) and SSP3-7.0 (IPCC, 2021) scenarios.

Elevated temperatures (eT) were applied to two 4 x 4 m plots within each shelter; one plot

received the control irrigation (eT-C), while the other plot received the drought treatment

(eT-D) and these were paired with two ambient temperature (aT) plots, one receiving control

irrigation (aT-C) and the other the drought treatment (aT-D). Each warmed plot had a heating

array comprising eight 1000W ceramic heaters (FTE 1000W, Ceramicx, Ireland) mounted on

an aluminium frame (4 m x 4 m) suspended 1.4 m above ground level (**Figure 2C**). Lamps

were positioned to give uniform coverage of IR radiation across the four composite subplots.

The power level to the heating lamps was adjusted each minute, via pulse width modulation

using a solid-state relay controlled by a data logger (CR1000, Campbell Scientific), based on

a proportional-integral control algorithm. Target temperatures for these plots were controlled

via feedback from IR-sensors (SI-100, Apogee Instruments, Logan, UT, USA) mounted at a

height of 3.8 m, recording plot surface temperatures every five minutes; temperatures thus

represent plot-level means for the plant canopy and, where visible, soil, and are henceforth

referred to as canopy temperatures. The + 3 °C warming treatment was applied with reference

to canopy temperatures for the relevant control and drought treatments (i.e., aT-C paired to

eT-C and aT-D paired to eT-D) to account for differences in soil moisture and vegetation

cover between these plots.

2.3 Environmental monitoring

Each shelter had a data logger (CR1000, Campbell Scientific) that recorded environmental

conditions and regulated the heating lamps. Soil moisture sensors (16 per shelter; Time

Domain Reflectometers; CS616, Campbell Scientific) recorded volumetric soil water content

(0-15 cm) every 15 min in all six replicates of four different species subplots and treatment

combinations (**Table S3**); in the *Medicago* subplots soil water content was also monitored at

a second depth (15-30 cm). Soil temperature probes (T107, Campbell Scientific) were

installed in the top 6-12 cm of the soil of eight subplots per shelter (all four drought and

warming treatment combinations of *Festuca* and *Medicago*) to record soil temperature every

15 min (**Table S3**). Air temperature and humidity sensors (Series RHP-2O3B, Dwyer

Instruments Inc, USA) mounted in force-ventilated radiation shields were installed inside

three of the rainout shelters at 0.6 m height, with records collected every 5 min to determine

any shelter effects on environmental conditions. Additionally, three sets of sensors were

installed at the same height outside the shelters. Photosynthetically active radiation (PAR)

was recorded at 5-min intervals using PAR sensors (Apogee quantum sensor, USA) installed

at a 6 m height outside two shelters, with two additional sensors located within shelters at 3

m.

2.4 Biomass harvests

All subplots were regularly harvested by clipping to determine aboveground productivity

during active growing periods. This 'surrogate grazing' involved use of hand shears and a

sickle mower. Timing of harvests was based on grazing recommendations for individual

species (Clements et al., 2003); hence, there were 2 (*Chloris* and *Digitaria*) or 3 (all other

species) harvests per species during the six-month winter and spring period in 2019 (Clark et

al., 2016). During each harvest plants were cut to 5 cm above the soil surface and weighed

(fresh and dry mass), with a sub-sample of harvested material sorted to determine the

proportion of live and dead biomass. The weed (i.e. non-target species) fraction from each

subplot was also assessed and was excluded from aboveground biomass measurements (<

5%). All materials were oven-dried at 70 °C for at least 48 hours prior to determining dry

mass.

2.5 Calculations and statistical analyses

The responses of temperature and soil water content to drought and warming treatments were

determined based on subplot-level (soil moisture, soil temperature) or plot-level (canopy

temperature) daily mean values. Aboveground production responses were determined based

on subplot totals, summed across all harvests conducted during the 6-month winter/spring

drought period.

Statistical analyses of treatment effects were conducted using linear mixed effects models

with climate treatments ('drought' or 'control': D or C, 'warming' or 'ambient': eT or aT) as

fixed effects and random effects defined as 'subplot nested within plot' or 'plot' (to account

for non-independence among continuous measurements) nested within 'shelter' (to account

for the blocked design), and 'date' (to account for temporal variation). Volumetric water

content data in *Medicago* subplots were also analysed for differences between upper (0-15

cm) and lower (15-30 cm) depths among climate treatments by including 'depth' as a fixed

effect. Climate treatment effects on aboveground production were analysed using two

different linear mixed effects models based on our hypotheses comparing responses among

different functional plant groups using a 'functional group' categorical predictor fixed effect.

For the first set of these analyses, 'drought' and 'warming' treatments were included as fixed

effects; 'Species' was also included as a random effect for models examining differences

among 'functional groups' (as a fixed effect), and 'plot nested within shelter' was included as

a nested random effect for all models. Aboveground production was natural log-transformed

to meet assumptions of constant variance, where indicated in statistical tables. All analyses

were conducted in R version 4.0.0 (R Core Team, 2020) using the package lme4 (Bates,

2015) and Kenward-Roger degrees of freedom were calculated using the Anova function in

the 'car' package (Fox and Weisberg, 2019). Pairwise comparisons to determine treatment

effects on soil or canopy temperature and soil water content, or aboveground production

among species or functional groups, were conducted using the package 'emmeans' (Length,

2020) and he Tukey method for *P*-value adjustment.

To compare responses to drought among species and functional groups, log response ratios

between paired control and droughted subplots within a shelter were calculated for

aboveground biomass summed across all harvests to derive a treatment effect size:

$$Effect\ size = \exp\left(\overline{\ln\left(\frac{Drought}{Control}\right)}\right) - 1 \qquad\qquad (1)$$

Linear mixed effects models were used for this second set of analyses evaluating differences

in the effect size among 1) species and 2) functional groups, with both specified as fixed

effects. Random effects included 'shelter' for both models, while the model for 2) also

included 'species' as an independent random effect. Modelled data were back transformed

prior to visualization of species- and functional group-level differences. The standard error of

the mean effect size was calculated as the product of the back-transformed mean and the

standard error of the effect size.

Interactions between warming and drought, the effect of drought on warming responses, and

the effect of warming on drought effects, were calculated as the log response ratio between

single factor treatment effects within a shelter for each period (Dieleman et al., 2012). These

ratios were calculated such that the effect of drought was examined by comparing the effect

size (Equation 1) for each warming level (aT-D / aT-C, eT-D / eT-C). The effect of warming

was examined as the effect size for each drought level (eT-C / aT-C, eT-D / aT-D) and the

effect of warming on the drought response as the ratio between these two effects ((eT-D / aT-

D)/(eT-C / aT-C)).

3 Results

3.1 Drought and warming treatment effects on environmental conditions

Drought and warming significantly altered soil water content and temperature across the six-

month winter/spring study period (**Figure 3**, **Figure S3-6**, **Tables S4** and **S5**). During this

time, droughted subplots of all continuously monitored species (*Biserrula*, *Festuca*,

Medicago, and *Lolium*) had significantly reduced soil moisture in the upper 15 cm (**Figure 3**;

Figure S3; $P < 0.05$) compared with controls. Additionally, soil water contents in all

droughted subplots were less variable compared to their respective control subplots (**Figure**

3A, B).

Soil water content was reduced by warming in *Festuca* subplots, particularly under drought

conditions ($P = 0.03$; **Figure S2**), although there were intermittent short-term positive effects

of warming on soil moisture following irrigation events for this species (**Figure 3B**). Soil

water content (0-15 cm) in *Medicago* subplots was unaffected by warming. There was,

however, evidence of greater soil water content at depth (15-30 cm) in droughted *Medicago*

soils, compared with surface levels (**Figure S4**).

Plot-level canopy temperatures were consistently increased by both drought ($P < 0.001$, + 1.0

°C) and warming ($P < 0.001$, + 3.0 °C). There were, however, no interactions between

drought and warming due to the experimental design which referenced warming treatments to

ambient temperatures for the respective droughted or control plot (**Figure S5**; **Table S4**).

Additionally, warming altered the maximum and minimum temperatures and thus the

temperature range within plots (**Figure S5**; **Table S5**). Warmed plots of *Festuca* and

Medicago had no days when minimum canopy temperatures fell below freezing (compared to

one day in ambient plots) and only 15 days when canopy temperatures fell below 5 °C

(compared to 45 days in ambient plots). At the other end of the scale, the warming treatment

resulted in an additional 16 days when canopy temperatures exceeded 40 °C and an extra 8

days where temperatures exceeded 45 °C (**Figure 3C, D**). Subsurface soil temperatures were

also significantly increased by both drought (*Festuca*) and warming (*Festuca* and *Medicago*)

(**Figure S6**). Overall, soil temperature responses were similar between species, although

drought amplified the warming effect in *Festuca* but not *Medicago* soils.

Shelter effects on air temperature, relative air humidity, and PAR levels were even across

shelters; on average there was an 11% decrease in temperature, a 6% increase in relative

humidity and a 22% reduction in PAR under shelters, as compared to outside (**Figure S7**).

Shelter effect on air temperature changed from winter to spring, with a slight cooling effect

during the winter, likely due to lower levels of radiation, and gradually transitioned to a

neutral and then positive effect during late spring (**Figure S7b**).

3.2 Aboveground production response to drought

During the 6-month winter/spring drought period, total aboveground production ranged from

2,800 to 8,300 kg ha^{-1} under control conditions (**Figure 4A**). The most productive species

were *Themeda*, *Medicago* and *Rytidosperma*, while *Festuca*, *Chloris* and *Biserrula* were the

least productive. Droughted subplots were significantly less productive than their respective

controls, with an average yield reduction of ~45% across the nine species (**Figure 4B**; **Table**

2). There was also a significant interaction between drought and species, with eight of the

nine species having significantly lower productivity under drought (**Table 2**). The remaining

species (*Lolium*, a perennial C_3 grass) experienced late spring die back and had a non-

significant 11% reduction in total productivity summed across the six months ($P = 0.51$).

The drought effect size, quantified as the log response ratio between control and droughted

subplots, varied among species (**Figure 4B**) with the greatest reductions being shown by two

of the C_4 grasses (*Digitaria:* 74% and *Themeda:* 59% yield reduction). The two legume

species showed intermediate reductions (*Biserrula*: 56% and *Medicago*: 37%), while C_3

grasses were the most variable group (*Lolium*: 11% to *Phalaris*: 48% reduction). All plant

functional groups were negatively impacted by drought, with C_4 grasses experiencing

significantly greater loss of productivity than C_3 grasses or legumes (**Figure 5; Table 2**).

Table 1. The origin, growth form and photosynthetic pathway of pasture and rangeland species selected for study in a drought and warming field experiment in the PACE facility

Species*	Origin	Growth Form	Photosynthetic pathway	Warming Treatment
Biserrula pelecinus	Temperate, introduced	Legume	C_3	
Medicago sativa	Temperate, introduced	Legume	C_3	Yes
Festuca arundinacea	Temperate, introduced	Grass	C_3	Yes
Lolium perenne	Temperate, introduced	Grass	C_3	
Phalaris aquatica	Temperate, introduced	Grass	C_3	
Rytidosperma caespitosum	Temperate, native	Grass	C_3	
Chloris gayana	Tropical, introduced	Grass	C_4	
Digitaria eriantha	Tropical, introduced	Grass	C_4	
Themeda triandra	Tropical, native	Grass	C_4	

*Species referenced by genus names in the text

Table 2. Linear mixed effects model output for the effects of drought on aboveground productivity of nine pasture and rangeland species during the 6-month winter/spring period.

Response	Fixed Effects	F value	P value	R²m[#]	R²c[§]
Biomass	Drought	$131.8_{1,23}$	<0.01	0.75	0.78
	Species	$24.0_{8,69}$	< 0.01		
	Drought x Species	$4.2_{8,69}$	< 0.01		
Biomass	Drought	$114.8_{1,23}$	<0.01	0.27	0.78
	FG[*]	$0.0_{2,6}$	0.97		
	Drought x FG	$6.4_{2,84}$	<0.01		

All biomass data were ln transformed to meet assumptions of constant variance. F value subscripts indicate degrees of freedom, [*]FG refers to functional group, [#]R²m indicates marginal error associated with linear model fixed effects, [§]R²c indicates conditional error or the total variation described by the full model including nested random effects (plots within a shelter).

3.3 Effects of warming and drought-warming interactions on aboveground biomass production

Warming during the 6-month winter/spring period generally reduced productivity (**Table 3**). While there was a broadly consistent response to the combination of drought and warming (**Table 3**), the absolute magnitude of productivity decline differed between the two species (**Figure 6A**). Warming resulted in a significant decline in winter/spring productivity in *Festuca* that was greater under drought (control: 18 %, drought: 31% reduction; **Figure 6B, C**). *Medicago* swards were not significantly affected by warming under either precipitation treatment (**Figure 6C**). Additionally, while *Medicago* swards that experienced ambient temperatures were negatively impacted by drought, productivity of swards that were exposed to both warming and drought did not differ significantly from those receiving the control precipitation regime (**Figure 6B**). This indicates that warming slightly reduces the negative effects of drought for *Medicago* (**Figure 6A**, **C**), although interactions between warming and drought were not significant for either species (**Figure 6D**).

Table 3. Linear mixed effects model output for the combined effects of drought and warming on aboveground productivity of both *Medicago sativa* and *Festuca arundinacea* during the 6-month winter/spring period.

Response	Fixed Effects	F value	*P* value	$R^2m^{\#}$	$R^2c^{\S}$
	Drought	$23.2_{1,15}$	<0.01	0.79	0.86
	Warming	$6.9_{1,15}$	0.02		
	Species	$208.3_{1,20}$	<0.01		
Biomass	D x W	$0.0_{1,15}$	0.86		
	D x S	$0.7_{1,20}$	0.42		
	W x S	$1.6_{1,20}$	0.22		
	D x W x S	$2.5_{1,20}$	0.13		

Biomass data were ln transformed to meet assumptions of constant variance, F value subscripts indicate degrees of freedom, $^{\#}R^2m$ indicates marginal error associated with linear model fixed effects, $^{\S}R^2c$ indicates conditional error or the total variation described by the full model including nested random effects (plots within a shelter).

4 Discussion

Projections of future climate change, including increases in the frequency and magnitude of

extreme climate events, are likely to disrupt ecosystem functioning (Jentsch and

Beierkuhnlein, 2008; Knapp et al., 2008; Nijp et al., 2015), with important consequences for

the productivity of pastures and rangelands (Jiménez et al., 2011; Godde et al., 2020). We

found that eight of the nine C_3 and C_4 species exposed to extreme winter/spring drought

experienced significant reductions in cool season productivity, with 45% declines on average

among species and losses of up to 74%, relative to controls. Despite large species differences

in drought sensitivity, we did find evidence for functional group-specific responses between

C_4 grasses, C_3 grasses, and legumes, such that C_4 grasses had the greatest response to

drought. Furthermore, we found no evidence of a positive effect of warming on the

productivity of two temperate species' (*Festuca* and *Medicago*) and the combination of

warming and drought resulted in the greatest biomass declines, which were either additive

(*Festuca*) or less-than-additive (*Medicago*). Taken together, these results demonstrate the

utility of evaluating responses of a range of species to single and compound climate extremes

for improved forecasts of future grassland vulnerability to climate change.

4.1 Productivity responses to winter/spring drought

Decades of research focused on examining grassland responses to drought have emphasized

the importance of regional climate context (Heisler-White et al., 2009; Knapp et al., 2017;

Slette et al., 2019) including rainfall seasonality (Padrón et al., 2020). The timing of drought

in relation to plant growth is a key factor influencing species' responses to changing rainfall

regimes, with evidence that shifts in rainfall seasonality can be more important than changes

in the amount of rainfall (Belovsky and Slade, 2020). Cool season rainfall has been found to

drive soil water storage and long-term patterns of productivity in many grasslands and

regions with year-round patterns of plant growth (Derner et al., 2020). Furthermore, there is

evidence that winter precipitation can offset the effects of spring/summer drought (Fry et al.,

2013). Locally in South-eastern Australia, projections for increased severity of winter/spring

drought (CSIRO, 2020) are likely to have large impacts on the seasonality of pasture and

rangeland productivity. Similar shifts in precipitation may have far reaching implications for

decisions on stocking densities and associated livestock production around the world (Derner

et al., 2020).

Changes in the seasonality of precipitation and the timing of when drought occurs with a

growing season, is particularly important in ecosystems that comprise a mixture of C_3 and C_4

plant species with distinct growth phenologies. In our warm temperate/subtropical region, we

predicted that cool-season drought would more negatively impact C_3 species due, in part, to

the timing of high soil water deficits during their period of active growth as well as

differences in water use efficiency. We found that while some individual C_3 species (e.g.

Festuca) experienced large growth reductions in response to cool-season drought, as a group,

C_4 species were more negatively affected than their temperate C_3 counterparts. This finding

has numerous implications for the sustainability of forage production, particularly in systems

where growth occurs year-round due to a lack of cold temperature constraints. Firstly,

sustained rainfall and soil water deficits during this period will have direct impacts on

productivity of temperate species that are active in winter and early spring, contributing to

winter feed-gaps. Secondly, the accumulated soil water deficit during periods of winter and

early spring drought can have negative impacts on the spring growth of warm-season (C_4)

grasses, as seen in our study and those of others in the USA (Prevéy and Seastedt, 2014;

Arredondo et al., 2016). If this response is consistent across C_4 grass species, management

recommendations that promote switching to more heat-tolerant C_4 species to accommodate

increased temperatures in traditionally temperate/sub-tropical cropping regions (Johnston,

1996) may not result in yield gains during the spring and early summer.

Legumes had the most consistent drought response, although broad generalizations are

limited with results from only two species. Compared with other species in this study,

Medicago experienced only modest impacts of drought on aboveground production. This may

have been a consequence of deep tap roots (Li et al., 2012; Nie et al., 2015), allowing access

to deep soil water to sustain growth during dry periods. Deep rooting strategies can also

facilitate shifts in moisture depth profiles due to hydraulic lift (Raza et al., 2013), thereby

reducing drought effects at the sward level (Liste and White, 2008; Pang et al., 2013).

Biserrula, also a deep-rooted legume (Loi et al., 2005; Haling et al., 2016), experienced

greater productivity losses in response to drought, compared to Medicago. However, unlike

Medicago, a perennial whose swards develop over multiple years (Li et al., 2012), *Biserrula*

is an annual species, regenerating from seed early in spring each year. These life history

differences that influence growth seasonality along with the significant effect of winter/spring

drought on productivity of the tropical (C_4) grasses, highlight the importance of drought

timing in relation to species' growth phenology (Wilcox et al., 2017; Yao et al., 2019), which

is at least partially reflected in their functional group classifications.

4.2 Plant responses to warming and drought × warming interactions

Gradual warming associated with climate change is predicted to affect ecosystems through a

variety of mechanisms, including direct impacts of increased air and soil temperature on

species' physiology and indirect impacts on soil water content via increases in evaporation or

transpiration (Deutsch et al., 2011). Many temperate and cold-climate systems report gains in

productivity associated with warming (Bloor et al., 2010; Wang et al., 2012), due to reduced

exposure to growth-limiting cold temperatures (Mori et al., 2014; Naudts et al., 2011; Reyer

et al., 2013) or greater nutrient availability resulting from increased microbial activity (Bloor

et al., 2010; Dellar et al., 2018). The beneficial impacts of warming on grassland productivity

have been reported both globally (Gao et al., 2016) and locally (Cullen et al., 2009). In other

systems, however, warming can shift temperatures beyond critical physiological thresholds,

resulting in reduced growth or even tissue die-back (Bastos et al., 2014; Cremonese et al.,

2017). In our sub-tropical system, we found no increase in productivity associated with cool-

season warming for either pasture species. Rather, we found a significant overall decline in

productivity in response to elevated temperature. This is despite an increase in winter

growing degree days and a reduction in frost exposure, changes that are expected to increase

winter and early spring growth (Chang et al., 2017; Piao et al., 2019). It is likely, therefore,

that lower productivity was associated with supra-optimal temperatures for these species

during spring and/or warming-associated reductions in water availability. In addition to these

productivity changes, changes to the nutritional quality of pastures can be anticipated and will

impact livestock production under future, more extreme climate conditions (Catunda et al.,

2021).

Festuca and *Medicago* are widely planted across Europe, North and South America,

Australasia and Africa (Gibson and Newman, 2001; Ghaleb et al., 2021), contributing to the

pasture feed-base that underpins global livestock production. These temperate species have

optimum temperatures for photosynthesis in the region of 20-29^{o}C (*Festuca;* (Sasaki et al.,

2002; Sinclair et al., 2007; Jacob et al., 2020) and 15-30^{o}C (*Medicago;* Al-hamdani and

Todd, 1990; Jacob et al., 2020). Whilst winter warming is, therefore, likely to stimulate gross

photosynthetic rates, spring temperatures were regularly above these thresholds, especially in

the later part of the season where daily maxima of over 45^{o}C were recorded in warmed plots.

The exceedance of thermal optima, along with increased respiratory carbon losses at warmer

temperatures (Heskel et al., 2016; Chandregowda, 2021) may explain observed productivity

declines in response to warming. There was, however, also evidence of small reductions in

soil water content associated with the warming treatment, particularly for *Festuca*. Given the

generally low levels of soil water availability in these well-drained sandy soils, this increased

soil moisture stress may have contributed to the large decline in aboveground productivity

observed in warmed plots for this species. Although higher temperatures are generally

associated with increased productivity in northern hemisphere grasslands (Craine et al.,

2012), warming can have both positive (winter) and negative (summer) impacts, depending

on ambient temperature (Kreyling et al., 2019). Contrary to expectations, our findings

indicate that perceived benefits of winter and spring warming may not be realized under field

conditions, especially where ambient levels of soil water availability are sub-optimal for

growth. Similar negative relationships between grassland productivity and cool season

temperatures have also been reported elsewhere from long term survey data (Wu et al., 2021).

High temperatures and drought are strongly coupled (Seneviratne et al., 2010) and their co-

occurrence can exacerbate soil water deficits as a result of evaporation from surface soils and

increased requirements for transpirational cooling (Ciais et al., 2005; Kirschbaum and

McMillan, 2018). However, the ecological impacts of these co-occurring stressors depend on

the physiological thermal optima and drought adaptation strategies of individual species, with

additive, greater-than-additive or less-than-additive responses all reported (Zavalloni et al.,

2008; Wu et al., 2011; Dreesen et al., 2012; Yu et al., 2012; De Boeck et al., 2016). In our

study both species exposed to warming alongside drought experienced the greatest

productivity declines in this combined treatment, with effects being either additive (*Festuca*)

or less-than-additive (*Medicago*) such that the effects of winter-spring drought were lower

reduced under continuous warming. An exacerbation of drought effects at higher

temperatures has been reported across biomes and plant functional groups, often associated

with reductions in soil moisture (Adams et al., 2009; Orsenigo et al., 2014; De Boeck et al.,

2016), such as we found for *Festuca*. This connection between temperature and water

availability is likely to amplify the intensity of ecological drought under future climates (Dai,

2011; IPCC, 2021).

Alternatively, temperature-driven reduction in available water has the potential to provide

drought-priming effects, through 'stress-memory', that reduce impacts of subsequent water

stress (Zavalloni et al., 2008). The less-than-additive productivity response in *Medicago*

exposed to the combined warming and drought treatment suggests the possibility of plant

acclimation to water stress, following prolonged exposure to low-severity droughts associated

with the warming treatment (Schwinning et al., 2004; Walter et al., 2013; Backhaus et al.,

2014). This acclimation may be important for the persistence and profitability of this pasture

species in the future. Similar less-than-additive temperature and drought interactions have

been observed in temperate grasslands and more generally across major biomes in meta-

analyses (De Boeck et al., 2011; Wu et al., 2011; Song et al., 2019). Importantly, our results

align with model predictions of productivity declines in Australian rangelands in response to

moderate (+3 °C) warming combined with reduced rainfall (McKeon et al., 2009).

Many studies have highlighted the importance of a species' persistence in drought and high

temperatures (Culvenor et al., 2016), and this is especially true for perennial grasses and

legumes in managed grasslands (Norton et al., 2009). Further work investigating the

mechanisms underpinning species' responses will help refine predictions about the impacts of

multiple, co-occurring changes under future climates. This study provides important, new

experimental, field-based data on the effects of extreme drought on a wide variety of pasture

species and two native grasses, and, for a sub-set of species, in combination with continuous

warming. These data complement information from modelling studies (Kaine and Tozer,

2005; Cullen et al., 2009) and comparisons across precipitation gradients (Clark et al., 2016)

to quantify impacts of future, more extreme rainfall regimes on the productivity of the pasture

and rangeland feed base that underpins livestock grazing in many parts of the world (Godde

et al., 2020). Work is underway to characterise relationships between productivity losses and

plant traits and strategies relating to the acquisition and use of water (root traits, plant-

microbial interactions, plant hydraulics) and the allocation of carbon (above- versus

belowground, root and crown carbohydrate storage), to determine the mechanisms

responsible for the observed species' differences to drought and warming. This information

will be key to extrapolating findings from this study to a wider range of locations and

grassland species, including many of international relevance in production systems across the

globe.

5 Conclusions

This paper introduces a new experimental facility used to simulate future, more extreme

climates, under field conditions. We found consistent negative effects of severe winter/spring

drought on seven pasture species and two native Australian rangeland grasses, highlighting

the challenges associated with future climate risk management for livestock production

systems. Strong reductions in cool season productivity for all functional groups highlight

potential climate constraints on winter forage availability, but also species' persistence

throughout the warmer summer months. Furthermore, the expanded use of tropical C_4 grasses

to mitigate declines in temperate C_3 species with rising global temperatures may lead to lower

cool-season productivity due to the relatively high seasonal drought sensitivity of the C_4

grasses examined in this study. Trade-offs are therefore implicit in planting decisions aimed

at enhancing pasture drought tolerance, given the increases in mean and maximum

temperatures that are already being observed globally. Additionally, substantial productivity

declines associated with warming, even in the cooler months, highlight the important role of

temperature in altering ecosystem water balance and, potentially, carbon dynamics,

suggesting limited benefits from future warming in warm temperate and sub-tropical systems.

Selection of species and cultivars with the physiological and/or phenological traits that

support sustained productivity under more extreme climate conditions will become

increasingly important as climate change undermines the performance of traditional pasture

and rangeland species.

6 Conflict of Interest

The authors declare that the research was conducted in the absence of any commercial or

financial relationships that could be construed as a potential conflict of interest.

7 Author Contributions/Funding/Acknowledgements

The authors would like to acknowledge individuals whose efforts supported the data

collection and site management for this work, including Angelica Vårhammar, Jackson

Bridges-Parlett, Alexandra Boyd, Shania Therese Didier Serre, Samantha Weller, Jinyan

Yang and Arjunan Krishnananthaselvan. We also gratefully acknowledge expert input to

early project discussions from Cath Lescun, Tom Davison, Tom Dickson, Allen Newman,

Warwick Badgery, Suzanne Boschma, Richard Eckard, Brendan Cullen and Alan Humphries,

and to Heritage Seeds for providing seeds. This project was supported by funding from Meat

and Livestock Australia's Donor Company (P.PSH. 0793), Dairy Australia (C100002357)

and Western Sydney University. Authors ACC and SAP wrote the paper with input from all

co-authors on draft iterations; ACC conducted statistical analyses with input from BEM, HZ,

JRP and KJF; SAP, EP, MGT, DTT, CVMB, BA, JRP, JMP, CAM, BDM, YC, ICA

designed the field experiment with input from ACC, HZ and KJF; ACC, HZ, KJF, KLMC,

MC, CI, VJ, GWK and AKP collected field samples.

8 Data Availability Statement

The datasets analysed for this study can be found in the Dryad Repository upon publication

of this manuscript.

8 References

Adams, H. D., Guardiola-Claramonte, M., Barron-Gafford, G. A., Villegas, J. C., Breshears, D. D., Zou, C. B., et al. (2009). Temperature sensitivity of drought-induced tree mortality portends increased regional die-off under global-change-type drought. *Proc. Natl. Acad. Sci. U. S. A.* 106, 7063–6. doi:10.1073/pnas.0901438106.

Adams, M. A., Turnbull, T. L., Sprent, J. I., and Buchmann, N. (2016). Legumes are different: leaf nitrogen, photosynthesis, and water use efficiency. *Proc. Natl. Acad. Sci.* 113, 4098–4103. doi:10.1073/pnas.1523936113.

Aerts, R., Cornelissen, J. H. C., and Dorrepaal, E. (2006). Plant performance in a warmer world: general responses of plants from cold, northern biomes and the importance of winter and spring events. *Plant Ecol.* 182, 65–77. doi:10.1007/s11258-005-9031-1.

Al-hamdani, S., and Todd, G. W. (1990). Effects of temperature on photosynthesis on photosynthate partitioning in alfalfa. *Can. J. Plant Sci.* 70, 203–208.

Aranjuelo, I., Irigoyen, J. J., and Sánchez-Díaz, M. (2007). Effect of elevated temperature and water availability on CO_2 exchange and nitrogen fixation of nodulated alfalfa plants. *Environ. Exp. Bot.* 59, 99–108. doi:10.1016/j.envexpbot.2005.10.008.

Arnone, J. A., Jasoni, R. L., Lucchesi, A. J., Larsen, J. D., Leger, E. A., Sherry, R. A., et al. (2011). A climatically extreme year has large impacts on C_4 species in tallgrass prairie ecosystems but only minor effects on species richness and other plant functional groups. *J. Ecol.* 99, 678–688. doi:10.1111/j.1365-2745.2011.01813.x.

Arredondo, T., Garcìa-Moya, E., Huber-Sannwald, E., Loescher, H. W., Delgado-Balbuena, J., and Luna-Luna, M. (2016). Drought manipulation and its direct and legacy effects on productivity of a monodominant and mixed-species semi-arid grassland. *Agric. For. Meteorol.* 223, 132–140. doi:10.1016/j.agrformet.2016.03.011.

Backhaus, S., Kreyling, J., Grant, K., Beierkuhnlein, C., Walter, J., and Jentsch, A. (2014). Recurrent mild drought events increase resistance toward extreme drought stress. *Ecosystems* 17, 1068–1081. doi:10.1007/s10021-014-9781-5.

Bai, E., Li, S., Xu, W., Li, W., Dai, W., and Jiang, P. (2013). A meta-analysis of experimental warming effects on terrestrial nitrogen pools and dynamics. *New Phytol.* 199, 431–40. doi:10.1111/nph.12252.

Bastos, A., Gouveia, C. M., Trigo, R. M., and Running, S. W. (2014). Analysing the spatio-temporal impacts of the 2003 and 2010 extreme heatwaves on plant productivity in Europe. *Biogeosciences* 11, 3421–3435. doi:10.5194/bg-11-3421-2014.

Bates, D., Maechler, M., Bolker, B. M., and Walker, S. (2015). Fitting linear mixed-effects

models using lme4. *J. Stat. Softw.* 67, 1–48. Available at: http://arxiv.org/abs/1406.5823.

Beier, C., Beierkuhnlein, C., Wohlgemuth, T., Penuelas, J., Emmett, B. A., Körner, C., et al.
(2012). Precipitation manipulation experiments - challenges and recommendations for
the future. *Ecol. Lett.* 15, 899–911. doi:10.1111/j.1461-0248.2012.01793.x.

Belovsky, G. E., and Slade, J. B. (2020). Climate change and primary production: Forty years
in a bunchgrass prairie. *PLoS One* 15, e0243496. doi:10.1371/journal.pone.0243496.

Bindi, M., and Olesen, J. E. (2011). The responses of agriculture in Europe to climate change.
Reg. Environ. Chang. 11, 151–158. doi:10.1007/s10113-010-0173-x.

Bloor, J. M. G., Pichon, P., Falcimagne, R., Leadley, P., and Soussana, J.-F. (2010). Effects
of warming, summer drought, and CO_2 enrichment on aboveground biomass production,
flowering phenology, and community structure in an upland grassland ecosystem.
Ecosystems 13, 888–900. doi:10.1007/s10021-010-9363-0.

Boer, M. M., Resco De Dios, V., and Bradstock, R. A. (2020). Unprecedented burn area of
Australian mega forest fires. *Nat. Clim. Chang.* 10, 170. doi:10.1038/s41558-020-0710-
7.

BOM (2019). Daily rainfall Richmond-UWS Hawkesbury. *Clim. Data Online*. Available at:
http://www.bom.gov.au/climate/data/index.shtml?bookmark=136.

Chandregowda, M. (2021). Aboveground-belowground linkages and carbon allocation in
pasture grasses during climate extremes.Western Sydney University Ph.D. Thesis.

Chang, J., Ciais, P., Viovy, N., Soussana, J. F., Klumpp, K., and Sultan, B. (2017). Future
productivity and phenology changes in European grasslands for different warming
levels: implications for grassland management and carbon balance. *Carbon Balance*
Manag. 12. doi:10.1186/s13021-017-0079-8.

Ciais, P., Reichstein, M., Viovy, N., Granier, A., Ogée, J., Allard, V., et al. (2005). Europe-
wide reduction in primary productivity caused by the heat and drought in 2003. *Nature*
437, 529–533. doi:10.1038/nature03972.

Clark, S. G., Nie, Z. N., Culvenor, R. A., Harris, C. A., Hayes, R. C., Li, G. D., et al. (2016).
Field evaluation of cocksfoot, Tall Fescue and Phalaris for dry marginal environments of
South-Eastern Australia. 1. Establishment and herbage production. *J. Agron. Crop Sci.*
202, 96–114. doi:10.1111/jac.12152.

Cleland, E. E., Chiariello, N. R., Loarie, S. R., Mooney, H. A., and Field, C. B. (2006).
Diverse responses of phenology to global changes in a grassland ecosystem. *PNAS* 103,
13740–13744. doi:10.1073/pnas.0600815103.

Clements, B., Ayres, L., Langford, C., McGarva, L., Simpson, P., Hennessy, G., et al. (2003).

662   The grazier's guide to pastures: sowing and managing profitable pastures in the central

663    and southern Tablelands, Monaro and Upper South West Slopes of New South Wales.

664    NSW.

665 Craine, J. M., Nippert, J. B., Elmore, A. J., Skibbe, A. M., Hutchinson, S. L., and Brunsell, N.

666    A. (2012). Timing of climate variability and grassland productivity. *Proc. Natl. Acad.*

667    *Sci.* 109, 3401–3405. doi:10.1073/pnas.1118438109.

668 Cremonese, E., Filippa, G., Galvagno, M., Siniscalco, C., Oddi, L., Morra di Cella, U., et al.

669    (2017). Heat wave hinders green wave: The impact of climate extreme on the phenology

670    of a mountain grassland. *Agric. For. Meteorol.* 247, 320–330.

671    doi:10.1016/j.agrformet.2017.08.016.

672 CSIRO (2020). State of the Climate. doi:ISBN:978-1-4863-1509-3.

673 Cullen, B. R., Johnson, I. R., Eckard, R. J., Lodge, G. M., Walker, R. G., Rawnsley, R. P., et

674    al. (2009). Climate change effects on pasture systems in south-eastern Australia. *Crop*

675    *Pasture Sci.* 60, 933–942. doi:10.1071/CP09019.

676 Culvenor, R. A., Clark, S. G., Harris, C. A., Hayes, R. C., Li, G. D., Nie, Z. N., et al. (2016).

677    Field evaluation of cocksfoot, tall fescue and Phalaris for dry marginal environments of

678    South-Eastern Australia. 2. Persistence. *J. Agron. Crop Sci.* 202, 355–371.

679    doi:10.1111/jac.12141.

680 Dai, A. (2011). Drought under global warming: A review. *Clim. Chang.* 2, 45–65.

681    doi:10.1002/wcc.81.

682 De Boeck, H. J., Bassin, S., Verlinden, M., Zeiter, M., and Hiltbrunner, E. (2016). Simulated

683    heat waves affected alpine grassland only in combination with drought. *New Phytol.*

684    209, 531–541. doi:10.1111/nph.13601.

685 De Boeck, H. J., Bloor, J. M. G., Aerts, R., Bahn, M., Beier, C., Emmett, B. A., et al. (2019).

686    Understanding ecosystems of the future will require more than realistic climate change

687    experiments – A response to Korell et al. *Glob. Chang. Biol.*, 19–20.

688    doi:10.1111/gcb.14854.

689 De Boeck, H. J., Dreesen, F. E., Janssens, I. A., and Nijs, I. (2010). Climatic characteristics

690    of heat waves and their simulation in plant experiments. *Glob. Chang. Biol.* 16, 1992–

691    2000. doi:10.1111/j.1365-2486.2009.02049.x.

692 De Boeck, H. J., Dreesen, F. E., Janssens, I. A., and Nijs, I. (2011). Whole-system responses

693    of experimental plant communities to climate extremes imposed in different seasons.

694    *New Phytol.* 189, 806–817. doi:10.1111/j.1469-8137.2010.03515.x.

695 De Boeck, H. J., Vicca, S., Roy, J., Nijs, I., Milcu, A., Kreyling, J., et al. (2015). Global

change experiments: challenges and opportunities. *Bioscience* 65, 922–931.

doi:10.1093/biosci/biv099.

Dellar, M., Topp, C. F. E., Banos, G., and Wall, E. (2018). A meta-analysis on the effects of

climate change on the yield and quality of European pastures. *Agric. Ecosyst. Environ.*

265, 413–420. doi:10.1016/j.agee.2018.06.029.

Derner, J. D., Raynor, E. J., Reeves, J. L., Augustine, D. J., and Milchunas, D. G. (2020).

Climatic and management determinants of large herbivore production in semiarid

grassland. *Agric. Ecosyst. Environ.* 290, 106761. doi:10.1016/j.agee.2019.106761.

Deutsch, E. S., Bork, E. W., Cahill, J. F., and Chang, S. X. (2011). Short-term plant

community responses to warming and defoliation in a northern temperate grassland.

ISRN Ecol. 2011, 1–8. doi:10.5402/2011/926061.

Dieleman, W. I. J., Vicca, S., Dijkstra, F. A., Hagedorn, F., Hovenden, M. J., Larsen, K. S., et

al. (2012). Simple additive effects are rare: a quantitative review of plant biomass and

soil process responses to combined manipulations of CO_2 and temperature. *Glob.*

Chang. Biol. 18, 2681–2693. doi:10.1111/j.1365-2486.2012.02745.x.

Dreesen, F. E., De Boeck, H. J., Janssens, I. A., and Nijs, I. (2012). Summer heat and drought

extremes trigger unexpected changes in productivity of a temperate annual/biannual

plant community. *Environ. Exp. Bot.* 79, 21–30. doi:10.1016/j.envexpbot.2012.01.005.

Dukes, J. S., Chiariello, N. R., Cleland, E. E., Moore, L. A., Shaw, M. R., Thayer, S., et al.

(2005). Responses of grassland production to single and multiple global environmental

changes. *PLoS Biol.* 3, e319. doi:10.1371/journal.pbio.0030319.

Easterling, D. R., Evans, J. L., Groisman, P. Y., Karl, T. R., Kunkel, K. E., and Ambenje, P.

(2000). Observed variability and trends in extreme climate events: a brief review. *Bull.*

Am. Meteorol. Soc. 81, 417–425. doi:10.1175/1520-

0477(2000)081<0417:OVATIE>2.3.CO;2.

Enloe, S. F., DiTomaso, J. M., Orloff, S. B., and Drake, D. J. (2004). Soil water dynamics

differ among rangeland plant communities dominated by yellow starthistle (*Centaurea*

solstitialis), annual grasses, or perennial grasses. *Weed Sci.* 52, 929–935.

doi:10.1614/ws-03-156r.

Fox, J., and Weisberg, S. (2019). *An R companion to applied regression*. Third. Thousand

Oaks, CA: Sage.

Frank, D., Reichstein, M., Bahn, M., Frank, D. C., Mahecha, M. D., Smith, P., et al. (2015).

Effects of climate extremes on the terrestrial carbon cycle: concepts, processes and

potential future impacts. *Glob. Chang. Biol.*, n/a-n/a. doi:10.1111/gcb.12916.

Fry, E. L., Manning, P., Allen, D. G. P., Hurst, A., Everwand, G., Rimmler, M., et al. (2013).
Plant functional group composition modifies the effects of precipitation change on
grassland ecosystem function. *PLoS One* 8. doi:10.1371/journal.pone.0057027.

Gao, Q., Schwartz, M. W., Zhu, W., Wan, Y., Qin, X., Ma, X., et al. (2016). Changes in
global grassland productivity during 1982 to 2011 attributable to climatic factors.
Remote Sens. 8, 1–12. doi:10.3390/rs8050384.

Ghaleb, W., Ahmed, L. Q., Escobar-Gutiérrez, A. J., and Julier, B. (2021). The history of
domestication and selection of Lucerne: a new perspective from the genetic diversity for
seed germination in response to temperature and scarification. *Front. Plant Sci.* 11, 1–
11. doi:10.3389/fpls.2020.578121.

Gibson, D. J., and Newman, J. A. (2001). *Festuca arudinacea* Schreber (*F. elatior* L. ssp.
arundinacea (Schreber) Hackel). *J. Ecol.* 89, 304–324. Available at:
https://www.jstor.org/stable/3599410.

Godde, C. M., Boone, R., Ash, A. J., Waha, K., Sloat, L., Thornton, P. K., et al. (2020).
Global rangeland production systems and livelihoods at threat under climate change and
variability. *Environ. Res. Lett.* 15, 44021. doi:10.1088/1748-9326/ab7395.

Haling, R. E., Yang, Z., Shadwell, N., Culvenor, R. A., Stefanski, A., Ryan, M. H., et al.
(2016). Growth and root dry matter allocation by pasture legumes and a grass with
contrasting external critical phosphorus requirements. *Plant Soil* 407, 67–79.
doi:10.1007/s11104-016-2808-2.

Hanson, P. J., and Walker, A. P. (2019). Advancing global change biology through
experimental manipulations: Where have we been and where might we go? *Glob.*
Chang. Biol. 26, 287–299. doi:10.1111/gcb.14894.

Heisler-White, J. L., Blair, J. M., Kelly, E. F., Harmoney, K., and Knapp, A. K. (2009).
Contingent productivity responses to more extreme rainfall regimes across a grassland
biome. *Glob. Chang. Biol.* 15, 2894–2904. doi:10.1111/j.1365-2486.2009.01961.x.

Heskel, M. A., O'Sullivan, O. S., Reich, P. B., Tjoelker, M. G., Weerasinghe, L. K.,
Penillard, A., et al. (2016). Convergence in the temperature response of leaf respiration
across biomes and plant functional types. *Proc. Natl. Acad. Sci. U. S. A.* 113, 3832–
3837. doi:10.1073/pnas.1520282113.

Hoeppner, S. S., and Dukes, J. S. (2012). Interactive responses of old-field plant growth and
composition to warming and precipitation. *Glob. Chang. Biol.* 18, 1754–1768.
doi:10.1111/j.1365-2486.2011.02626.x.

IPCC (2017). IPCC fifth assessment report (AR5) observed climate change impacts database,

Version 2.01. Palisades, NY doi:https://doi.org/10.7927/H4FT8J0X.

IPCC (2021). *Climate Change 2021: The Physical Science Basis. Contribution of Working*

Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate

Change. 6th ed. , eds. V. Masson-Delmotte, P. Zhai, A. Pirani, S. . Connors, C. Pean, S.

Berger, et al. Cambridge University Press.

Jacob, V., Zhang, H., Churchill, A. C., Yang, J., Choat, B., Medlyn, B. E., et al. (2020).

Warming reduces net carbon gain and productivity in *Medicago sativa* L. and *Festuca*

arundinacea. *Agronomy* 10. doi:10.3390/agronomy10101601.

Jentsch, A., and Beierkuhnlein, C. (2008). Research frontiers in climate change: effects of

extreme meteorological events on ecosystems. *Comptes Rendus - Geosci.* 340, 621–628.

doi:10.1016/j.crte.2008.07.002.

Jentsch, A., Kreyling, J., Boettcher-Treschkow, J., and Beierkuhnlein, C. (2009). Beyond

gradual warming: extreme weather events alter flower phenology of European grassland

and heath species. *Glob. Chang. Biol.* 15, 837–849. doi:10.1111/j.1365-

2486.2008.01690.x.

Jiménez, M. A., Jaksic, F. M., Armesto, J. J., Gaxiola, A., Meserve, P. L., Kelt, D. A., et al.

(2011). Extreme climatic events change the dynamics and invasibility of semi-arid

annual plant communities. *Ecol. Lett.* 14, 1227–1235. doi:10.1111/j.1461-

0248.2011.01693.x.

Johnston, W. H. (1996). The place of C_4 grasses in temperate pastures in Australia. *New Zeal.*

J. Agric. Res. 39, 527–540. doi:10.1080/00288233.1996.9513213.

Kaine, G. W., and Tozer, P. R. (2005). Stability, resilience and sustainability in pasture-based

grazing systems. *Agric. Syst.* 83, 27–48. doi:10.1016/j.agsy.2004.03.001.

Keatinge, J. D. H., Qi, A., Wheeler, T. R., Ellis, R. H., and Summerfield, R. J. (1998). Effects

of temperature and photoperiod on phenology as a guide to the selection of annual

legume cover and green manure crops for hillside farming systems. *F. Crop. Res.* 57,

139–152. doi:10.1016/S0378-4290(97)00122-6.

Kirschbaum, M. U. F., and McMillan, A. M. S. (2018). Warming and elevated CO_2 have

opposing influences on transpiration. Which is more important? *Curr. For. Reports* 4,

51–71. doi:10.1007/s40725-018-0073-8.

Knapp, A. K., Avolio, M. L., Beier, C., Carroll, C. J. W., Collins, S. L., Dukes, J. S., et al.

(2017). Pushing precipitation to the extremes in distributed experiments:

recommendations for simulating wet and dry years. *Glob. Chang. Biol.* 23, 1774–1782.

doi:10.1111/gcb.13504.

Knapp, A. K., Beier, C., Briske, D. D., Classen, A. T., Luo, Y., Reichstein, M., et al. (2008).

Consequences of more extreme precipitation regimes for terrestrial ecosystems.

Bioscience 58, 811. doi:10.1641/B580908.

König, P., Tautenhahn, S., Cornelissen, J. H. C., Kattge, J., Bönisch, G., and Römermann, C.

(2018). Advances in flowering phenology across the Northern Hemisphere are explained

by functional traits. *Glob. Ecol. Biogeogr.* 27, 310–321. doi:10.1111/geb.12696.

Kreyling, J., Beierkuhnlein, C., Ellis, L., and Jentsch, A. (2008). Invasibility of grassland and

heath communities exposed to extreme weather events- additive effects of diversity

resistance and fluctuating physical environment. *Oikos* 117, 1542–1554.

doi:10.1111/j.2008.0030-1299.16653.x.

Kreyling, J., Grant, K., Hammerl, V., Arfin-Khan, M. A. S., Malyshev, A. V., Peñuelas, J., et

al. (2019). Winter warming is ecologically more relevant than summer warming in a

cool-temperate grassland. *Sci. Rep.* 9, 1–9. doi:10.1038/s41598-019-51221-w.

Kueppers, L. M., Conlisk, E., Castanha, C., Moyes, A. B., Germino, M. J., de Valpine, P., et

al. (2017). Warming and provenance limit tree recruitment across and beyond the

elevation range of subalpine forest. *Glob. Chang. Biol.* 23, 2383–2395.

doi:10.1111/gcb.13561.

Length, R. (2020). emmeans: estimated marginal means, aka least-squares means.

https://cran.r-project.org/package=emmeans. Available at: https://cran.r-

project.org/package=emmeans.

Lesica, P., and Kittelson, P. M. (2010). Precipitation and temperature are associated with

advanced flowering phenology in a semi-arid grassland. *J. Arid Environ.* 74, 1013–1017.

doi:10.1016/j.jaridenv.2010.02.002.

Li, F., Kautz, T., Pude, R., and Köpke, U. (2012). Nodulation of lucerne (Medicago sativa L.)

roots: Depth distribution and temporal variation. *Plant, Soil Environ.* 58, 424–428.

doi:10.17221/314/2012-pse.

Liste, H.-H., and White, J. C. (2008). Plant hydraulic lift of soil water – implications for crop

production and land restoration. *Plant Soil* 313, 1–17. doi:10.1007/s11104-008-9696-z.

Loi, A., Howieson, J. G., Nutt, B. J., and Carr, S. J. (2005). A second generation of annual

pasture legumes and their potential for inclusion in Mediterranean-type farming systems.

Aust. J. Exp. Agric. 45, 289–299. doi:10.1071/ea03134.

Lu, M., Zhou, X., Yang, Q., Li, H., Fang, C., Chen, J., et al. (2013). Responses of ecosystem

carbon cycle to experimental warming: a meta-analysis. *Ecology* 94, 726–738.

Ma, X., Huete, A., Moran, S., Ponce-Campos, G., and Eamus, D. (2015). Abrupt shifts in

phenology and vegetation productivity under climate extremes. *J. Geophys. Res.*

Biogeosciences 120, 2036–2052. doi:10.1002/2015JG003144.

McKeon, G. M., Stone, G. S., Syktus, J. I., Carter, J. O., Flood, N. R., Ahrens, D. G., et al.

(2009). Climate change impacts on northern Australian rangeland livestock carrying

capacity: A review of issues. *Rangel. J.* 31, 1–29. doi:10.1071/RJ08068.

Meterology, B. of (2020). Bureau of Meterology. *Gov. Aust.* Available at:

http://www.bom.gov.au/ [Accessed August 15, 2018].

Mori, A. S., Fujii, S., and Kurokawa, H. (2014). Ecological consequences through responses

of plant and soil communities to changing winter climate. *Ecol. Res.* 29, 547–559.

doi:10.1007/s11284-013-1091-4.

Munson, S. M., and Long, A. L. (2017). Climate drives shifts in grass reproductive

phenology across the western USA. *New Phytol.* 213, 1945–1955.

doi:10.1111/nph.14327.

Nábrádi, A. (2016). The economic value of grassland products. *Appl. Stud. Agribus. Commer.*

1, 19–28. doi:10.19041/apstract/2007/1/2.

Naudts, K., Van den Berge, J., Janssens, I. A., Nijs, I., and Ceulemans, R. (2011). Does an

extreme drought event alter the response of grassland communities to a changing

climate? *Environ. Exp. Bot.* 70, 151–157. doi:10.1016/j.envexpbot.2010.08.013.

Nie, Z., Mitchell, M., Clark, S., and Smith, K. (2015). Distribution of lucerne roots in

summer-dry environments of southern Australia. in *Proceedings of the 17th ASA*

Conference (Hobart, Australia), 1–4.

Nijp, J. J., Limpens, J., Metselaar, K., Peichl, M., Nilsson, M. B., van der Zee, S. E. A. T. M.,

et al. (2015). Rain events decrease boreal peatland net CO_2 uptake through reduced light

availability. *Glob. Chang. Biol.* 21, 2309–2320. doi:10.1111/gcb.12864.

Norton, M. R., Volaire, F., Lelièvre, F., and Fukai, S. (2009). Identification and measurement

of summer dormancy in temperate perennial grasses. *Crop Sci.* 49, 2347–2352.

doi:10.2135/cropsci2009.06.0319.

Novem Auyeung, D. S., Suseela, V., and Dukes, J. S. (2013). Warming and drought reduce

temperature sensitivity of nitrogen transformations. *Glob. Chang. Biol.* 19, 662–676.

doi:10.1111/gcb.12063.

O'Mara, F. P. (2012). The role of grasslands in food security and climate change. *Ann. Bot.*

110, 1263–1270. doi:10.1093/aob/mcs209.

Ode, D. J., and Tieszen, L. L. (1980). The seasonal contribution of C_3 and C_4 plant species to

primary production in a mixed prairie. *Ecology* 61, 1304–1311.

Orsenigo, S., Mondoni, A., Rossi, G., and Abeli, T. (2014). Some like it hot and some like it
cold, but not too much: Plant responses to climate extremes. *Plant Ecol.* 215, 677–688.
doi:10.1007/s11258-014-0363-6.

Padrón, R. S., Gudmundsson, L., Decharme, B., Ducharne, A., Lawrence, D. M., Mao, J., et
al. (2020). Observed changes in dry-season water availability attributed to human-
induced climate change. *Nat. Geosci.* 13, 477–481. doi:10.1038/s41561-020-0594-1.

Pang, J., Wang, Y., Lambers, H., Tibbett, M., Siddique, K. H. M., and Ryan, M. H. (2013).
Commensalism in an agroecosystem: hydraulic redistribution by deep-rooted legumes
improves survival of a droughted shallow-rooted legume companion. *Physiol. Plant.*
149, 79–90. doi:10.1111/ppl.12020.

Parmesan, C., Root, T. L., and Willig, M. R. (2000). Impacts of extreme weather and climate
on terrestrial biota. *Bull. Am. Meteorol. Soc.* 81, 443–450.

Pearcy, R. W., Caldwell, M. M., Keeley, J. E., Monson, R. K., and Strain, B. R. (1987).
Carbon gain by plants in natural environments. *Bioscience* 37, 21–29.
doi:10.2307/1310174.

Pearcy, R. W., and Ehleringer, J. (1984). Comparative ecophysiology of C_3 and C_4 plants.
Plant. Cell Environ. 7, 1–13. doi:10.1111/j.1365-3040.1984.tb01194.x.

Peng, A., Klanderud, K., Wang, G., Zhang, L., Xiao, Y., and Yang, Y. (2020). Plant
community responses to warming modified by soil moisture in the Tibetan Plateau.
Arctic, Antarct. Alp. Res. 52, 60–69. doi:10.1080/15230430.2020.1712875.

Piao, S., Liu, Q., Chen, A., Janssens, I. A., Fu, Y., Dai, J., et al. (2019). Plant phenology and
global climate change: current progresses and challenges. *Glob. Chang. Biol.* 25, 1922–
1940. doi:10.1111/gcb.14619.

Power, S. A., Barnett, K. L., Ochoa-hueso, R., Facey, S. L., Gibson-Forty, E. V. J., Hartley,
S. E., et al. (2016). DRI-Grass: A new experimental platform for addressing grassland
ecosystem responses to future precipitation scenarios in South-East Australia. *Front.*
Plant Sci. 7, 1–14. doi:10.3389/fpls.2016.01373.

Prevéy, J. S., and Seastedt, T. R. (2014). Seasonality of precipitation interacts with exotic
species to alter composition and phenology of a semi-arid grassland. *J. Ecol.* 102, 1549–
1561. doi:10.1111/1365-2745.12320.

R Core Team (2020). R: A language and environment for statistical computing. http://www.r-
project.org/. Available at: http://www.r-project.org/.

Rathcke, B., and Lacey, E. P. (1985). Phenological patterns of terrestrial plants. *Annu. Rev.*
Ecol. Syst. 16, 179–214.

Raza, A., Friedel, J. K., Moghaddam, A., Ardakani, M. R., Loiskandl, W., Himmelbauer, M.,
et al. (2013). Modeling growth of different lucerne cultivars and their effect on soil
water dynamics. *Agric. Water Manag.* 119, 100–110. doi:10.1016/j.agwat.2012.12.006.

Reyer, C. P. O., Leuzinger, S., Rammig, A., Wolf, A., Bartholomeus, R. P., Bonfante, A., et
al. (2013). A plant's perspective of extremes: terrestrial plant responses to changing
climatic variability. *Glob. Chang. Biol.* 19, 75–89. doi:10.1111/gcb.12023.

Sage, R. F. (2019). Global change biology: a primer. *Glob. Chang. Biol.*, gcb.14893.
doi:10.1111/gcb.14893.

Sanz-Sáez, Á., Erice, G., Aguirreolea, J., Irigoyen, J. J., and Sánchez-Díaz, M. (2012).
Alfalfa yield under elevated CO_2 and temperature depends on the *Sinorhizobium* strain
and growth season. *Environ. Exp. Bot.* 77, 267–273.
doi:10.1016/j.envexpbot.2011.11.017.

Sasaki, H., Fukuyama, M., Onoue, T., Suyama, T., and Ahoji, A. (2002). Effects of
increasing CO_2 concentration and leaf temprature on the photosynthesis of Tall Fescue
(*Festuca arundinacea* Schreb.). *Grassl. Sci.* 48, 12–16.

Schwinning, S., Sala, O. E., Loik, M. E., and Ehleringer, J. R. (2004). Thresholds, memory,
and seasonality: understanding pulse dynamics in arid/semi-arid ecosystems. *Oecologia*
141, 191–193. doi:10.1007/s00442-004-1683-3.

Seneviratne, S. I., Corti, T., Davin, E. L., Hirschi, M., Jaeger, E. B., Lehner, I., et al. (2010).
Investigating soil moisture-climate interactions in a changing climate: a review. *Earth-*
Science Rev. 99, 125–161. doi:10.1016/j.earscirev.2010.02.004.

Sherry, R. A., Weng, E., Arnone, J. A., Johnson, D. W., Schimel, D. S., Verburg, P. S., et al.
(2008). Lagged effects of experimental warming and doubled precipitation on annual
and seasonal aboveground biomass production in a tallgrass prairie. *Glob. Chang. Biol.*
14, 2923–2936. doi:10.1111/j.1365-2486.2008.01703.x.

Sherry, R. A., Zhou, X., Gu, S., Arnone, J. A., Schimel, D. S., Verburg, P. S., et al. (2007).
Divergence of reproductive phenology under climate warming. *PNAS* 104, 198–202.

Sinclair, T., Fiscus, E., Wherley, B., Durham, M., and Rufty, T. (2007). Atmospheric vapor
pressure deficit is critical in predicting growth response of "cool-season" grass *Festuca*
arundinacea to temperature change. *Planta* 227, 273–276. doi:10.1007/s00425-007-
0645-5.

Slette, I. J., Post, A. K., Awad, M., Even, T., Punzalan, A., Williams, S., et al. (2019). How
ecologists define drought, and why we should do better. *Glob. Chang. Biol.* 25, 3193–
3200. doi:10.1111/gcb.14747.

Song, J., Wan, S., Piao, S., Knapp, A. K., Classen, A. T., Vicca, S., et al. (2019). A meta-
analysis of 1,119 manipulative experiments on terrestrial carbon-cycling responses to
global change. *Nat. Ecol. Evol.* 3, 1309–1320. doi:10.1038/s41559-019-0958-3.

Steffen, W., Richardson, K., Rockström, J., Cornell, S., Fetzer, I., Bennett, E., et al. (2015).
Planetary boundaries: guiding human development on a changing planet. *Science (80-.).*
347, 1259855. doi:10.1126/science.1259855.

Veenendaal, E. M., Ernst, W. H. O., and Modise, G. S. (1996). Reproductive effort and
phenology of seed production of savanna grasses with different growth form and life
history. *Vegetatio* 123, 91–100. doi:10.1007/BF00044891.

Volder, A., Briske, D. D., and Tjoelker, M. G. (2013). Climate warming and precipitation
redistribution modify tree-grass interactions and tree species establishment in a warm-
temperate savanna. *Glob. Chang. Biol.* 19, 843–857. doi:10.1111/gcb.12068.

Walker, M. D., Wahren, C.-H., Hollister, R. D., Henry, G. H. R., Ahlquist, L. E., Alatalo, J.
M., et al. (2006). Plant community responses to experimental warming across the tundra
biome. *Proc. Natl. Acad. Sci. U. S. A.* 103, 1342–6. doi:10.1073/pnas.0503198103.

Walter, J., Jentsch, A., Beierkuhnlein, C., and Kreyling, J. (2013). Ecological stress memory
and cross stress tolerance in plants in the face of climate extremes. *Environ. Exp. Bot.*
94, 3–8. doi:10.1016/j.envexpbot.2012.02.009.

Wang, C., Sun, Y., Chen, H. Y. H., Yang, J., and Ruan, H. (2021). Meta-analysis shows non-
uniform responses of above- and belowground productivity to drought. *Sci. Total*
Environ. 782, 146901. doi:10.1016/j.scitotenv.2021.146901.

Wang, S., Duan, J., Xu, G., Wang, Y., Zhang, Z., Rui, Y., et al. (2012). Effects of warming
and grazing on soil N availability , species composition , and ANPP in an alpine
meadow. *Ecology* 93, 2365–2376. doi:10.1890/11-1408.1.

Whittington, H. R., Deede, L., and Powers, J. S. (2012). Growth responses, biomass
partitioning, nitrogen isotopes of prairie legumes in response to elevated temperature
and varying nitrogen source in a growth chamber experiment. *Am. J. Bot.* 99, 838–846.
doi:10.3732/ajb.1100283.

Whittington, H. R., Tilman, D., and Powers, J. S. (2013). Consequences of elevated
temperatures on legume biomass and nitrogen cycling in a field warming and
biodiversity experiment in a North American prairie. *Funct. Plant Biol.* 40, 1147–1158.
doi:10.1071/FP12345.

Wilcox, K. R., Blair, J. M., Smith, M. D., and Knapp, A. K. (2016). Does ecosystem
sensitivity to precipitation at the site- level conform to regional- scale predictions?

Ecology 97, 561–568. doi:10.1007/s13398-014-0173-7.2.

Wilcox, K. R., Shi, Z., Gherardi, L. A., Lemoine, N. P., Koerner, S. E., Hoover, D. L., et al.

(2017). Asymmetric responses of primary productivity to precipitation extremes: A

synthesis of grassland precipitation manipulation experiments. *Glob. Chang. Biol.* 23,

4376–4385. doi:10.1111/gcb.13706.

Winslow, J. C., Hunt, E. R., and Piper, S. C. (2003). The influence of seasonal water

availability on global C_3 versus C_4 grassland biomass and its implications for climate

change research. *Ecol. Modell.* 163, 153–173. doi:10.1016/S0304-3800(02)00415-5.

Wu, G. L., Cheng, Z., Alatalo, J. M., Zhao, J., and Liu, Y. (2021). Climate warming

consistently reduces grassland ecosystem productivity. *Earth's Futur.* 9, 1–14.

doi:10.1029/2020EF001837.

Wu, Z., Dijkstra, P., Koch, G. W., Peñuelas, J., and Hungate, B. A. (2011). Responses of

terrestrial ecosystems to temperature and precipitation change: A meta-analysis of

experimental manipulation. *Glob. Chang. Biol.* 17, 927–942. doi:10.1111/j.1365-

2486.2010.02302.x.

Yao, X., Wu, J., and Gong, X. (2019). Precipitation and seasonality affect grazing impacts on

herbage nutritive values in alpine meadows on the Qinghai-Tibet Plateau. *J. Plant Ecol.*

12, 993–1008. doi:10.1093/jpe/rtz027.

Yu, J., Chen, L., Xu, M., and Huang, B. (2012). Effects of elevated CO_2 on physiological

responses of Tall Fescue to elevated temperature, drought stress, and the combined

stresses. *Crop Sci.* 52, 1848–1858. doi:10.2135/cropsci2012.01.0030.

Yuan, W., Cai, W., Chen, Y., Liu, S., Dong, W., Zhang, H., et al. (2016). Severe summer

heatwave and drought strongly reduced carbon uptake in Southern China. *Sci. Rep.* 6, 1–

12. doi:10.1038/srep18813.

Zavalloni, C., Gielen, B., Lemmens, C. M. H. M., De Boeck, H. J., Blasi, S., Van Den Bergh,

S., et al. (2008). Does a warmer climate with frequent mild water shortages protect

grassland communities against a prolonged drought? *Plant Soil* 308, 119–130.

doi:10.1007/s11104-008-9612-6.

Zeiter, M., Schärrer, S., Zweifel, R., Newbery, D. M., and Stampfli, A. (2016). Timing of

extreme drought modifies reproductive output in semi-natural grassland. *J. Veg. Sci.* 27,

238–248. doi:10.1111/jvs.12362.

1000

Figure Captions

Figure 1. Schematic representation of expected changes in aboveground plant biomass during winter and spring associated with concurrent drought for A) plant functional groups including legumes and grasses relying on either the C_3 or C_4 photosynthetic pathways. Legumes are predicted to be less affected by drought (shallower slope) due to their higher nutrient and water use efficiency compared with C_3 grasses, while C_3 grasses are predicted to be most affected by drought due to a comparatively lower water use efficiency. C_4 grasses are predicted to have lower biomass due to low temperature constraints on growth but lower drought sensitivity than C_3 species due to greater water use efficiency. B) For legumes (green) and C_3 grasses (blue) winter and spring warming (eT) is predicted to increase productivity under the control precipitation treatment relative to ambient temperature (aT; solid line); potential interactions between drought and warming can include an amplifying effect resulting in increased loss of biomass (C_3 grass) or a stabilizing effect where warming offsets the impact of drought on biomass (legume).

Figure 2. The Pastures and Climate Extremes (PACE) field facility located at Western Sydney University in Richmond, New South Wales, Australia. A) There are six open-sided polytunnels, each with eight experimental plots; B) Experimental plots (4 x 4 m) received a drought, warming, or drought + warming treatment and are each divided into four discrete 2 x 2 m subplots comprising nine different pasture species in monoculture; C) Heater (infra-red: IR) arrays were mounted above the vegetation canopy and warmed the plot surface an average of 3 °C above paired (control, drought) ambient temperature plots.

Figure 3. Experimental treatment effects of winter/spring drought (control: C, drought: D) applied during the period between 1 June to 30 November 2019 and year-round warming (ambient temperature: aT, elevated temperature: eT) treatments on soil moisture (panels A and B) and canopy temperature (panels C and D) averaged across the six shelters, from 1 May 2019 to 30 November 2019. A) Average soil volumetric water content in *Festuca* sub-plots (additional species are shown in Figure S2) with 95% confidence intervals as well as individual irrigation events as daily rainfall totals for control (black bars) and drought (red bars) plots over the 6-month period of study; B) Treatment differences in soil water content from aT-C plots over time in the *Festuca* subplots, noting the winter/spring drought treatment period between 1 June and 30 November 2019; C) Daily maximum canopy temperature, relative to maximum ambient air temperature (black line) and 40 °C (representing extreme temperatures, dashed lines); D) Daily minimum temperature compared with minimum air temperature (black line) and 0 °C (dashed lines).

Figure 4. A) Aboveground production summed for all harvests during the 6-month winter/spring drought period. Large points shown are means ± 1 standard error (control = solid symbol, droughted = open symbol) and opaque points show species level variability in biomass. Significant pairwise comparisons for the effect of drought treatment are indicated as follows: NS= not significant, * $P < 0.05$, ** $P = <0.01$, *** $P = < 0.001$. B) drought effect size (log response ratio of drought vs control production during the 6-month drought treatment period). Values for panel B are mean values with 95% confidence intervals and same letter designations indicate non-significant differences among plantings. Abbreviations are as follows: *Biserrula* (Bis), *Chloris* (Chl), *Digitaria* (Dig), *Festuca* (Fes), *Lolium* (Lol), *Medicago* (Med), *Phalaris* (Pha), *Rytidosperma* (Ryt), *Themeda* (The).

Figure 5. Aboveground production summed for all harvests during the 6-month winter/spring drought period by functional group. Large points shown are means ± 1 standard error (control =

solid symbol, droughted = open symbol) and opaque points show subplot level variability in biomass for each species within the functional group. Significant pairwise comparisons for the effect of drought within a functional group are indicated matching Fig 4.

Figure 6. Aboveground production for *Festuca* and *Medicago* exposed to the combined effects of drought (control: C and droughted: D) and warming (ambient: aT and elevated: eT) treatments during A) the 6-month winter/spring drought period. Treatment effect sizes during the winter/spring drought period for B) drought under ambient (also shown in Fig 4) and elevated temperatures, C) warming, under control and droughted conditions, and D) the effect of warming on biomass responses to drought (for panel D, positive values indicate a reduction in drought impact under warming). Values in panel A are means ± 1 SE; while panels B-D are means and 95% confidence intervals. Spp. abbreviations and significance levels follow Fig. 4.

Figure 1. Schematic representation of expected changes in aboveground plant biomass during winter and spring associated with concurrent drought for A) plant functional groups including legumes and grasses relying on either the C_3 or C_4 photosynthetic pathways. Legumes are predicted to be less affected by drought (shallower slope) due to their higher nutrient and water use efficiency compared with C_3 grasses, while C_3 grasses are predicted to be most affected by drought due to a comparatively lower water efficiency. C_4 grasses are predicted to have lower biomass due to low temperature constraints on growth but lower drought sensitivity than C_3 species due to greater water use efficiency. B) For legumes (green) and C_3 grasses (blue) winter and spring warming (eT) is predicted to increase productivity under the control precipitation treatment relative to ambient temperature (aT; solid line); potential interactions between drought and warming can include an amplifying effect resulting in increased loss of biomass (C_3 grass) or a stabilizing effect where warming offsets the impact of drought on biomass (legume).

1064

Figure 2. The Pastures and Climate Extremes (PACE) field facility located at Western Sydney University in Richmond, New South Wales, Australia. A) There are six open-sided polytunnels, each with eight experimental plots; B) Experimental plots (4 x 4 m) received a drought, warming, or drought + warming treatment and are each divided into four discrete 2 x 2 m subplots comprising nine different pasture species in monoculture; C) Heater (infra-red: IR) arrays were mounted above the vegetation canopy and warmed the plot surface an average of 3 °C above paired (control, drought) ambient temperature plots.

1065

1066

Figure 3. Experimental treatment effects of winter/spring drought (control: C, drought: D) applied during the period between 1 June to 30 November 2019 and year-round warming (ambient temperature: aT, elevated temperature: eT) treatments on soil moisture (panels A and B) and canopy temperature (panels C and D) averaged across the six shelters, from 1 May 2019 to 30 November 2019. A) Average soil volumetric water content in *Festuca* sub-plots (additional species are shown in Figure S2) with 95% confidence intervals as well as individual irrigation events as daily rainfall totals for control (black bars) and drought (red bars) plots over the 6-month period of study; B) Treatment differences in soil water content from aT-C plots over time in the *Festuca* subplots, noting the winter/spring drought treatment period between 1 June and 30 November 2019; C) Daily maximum canopy temperature, relative to maximum ambient air temperature (black line) and 40 °C (representing extreme temperatures, dashed lines); D) Daily minimum temperature compared with minimum air temperature (black line) and 0 °C (dashed lines).

1067
1068

Figure 4. A) Aboveground production summed for all harvests during the 6-month winter/spring drought period. Large points shown are means ± 1 standard error (control = solid symbol, droughted = open symbol) and opaque points show species level variability in biomass; B) drought effect size (log response ratio of drought vs control production during the 6-month drought treatment period). Values for panel B are mean values with 95% confidence intervals and same letter designations indicate non-significant differences among plantings. Abbreviations are as follows: *Biserrula* (Bis), *Chloris* (Chl), *Digitaria* (Dig), *Festuca* (Fes), *Lolium* (Lol), *Medicago* (Med), *Phalaris* (Pha), *Rytidosperma* (Ryt), *Themeda* (The). Significant pairwise comparisons for the effect of drought treatment are indicated as follows: NS= not significant, * $P < 0.05$, ** $P = <0.01$, *** $P = < 0.001$.

Figure 5. Aboveground production summed for all harvests during the 6-month winter/spring drought period by functional group. Large points shown are means ± 1 standard error (control = solid symbol, droughted = open symbol) and opaque points show subplot level variability in biomass for each species within the functional group. Significant pairwise comparisons for the effect of drought within a functional group are indicated matching Fig 4.

1071

Figure 6. Aboveground production for *Festuca* and *Medicago* exposed to the combined effects of drought (control: C and droughted: D) and warming (ambient: aT and elevated: eT) treatments during A) the 6-month winter/spring drought period. Treatment effect sizes during the winter/spring drought period for B) drought under ambient (also shown in Fig 4) and elevated temperatures, C) warming, under control and droughted conditions, and D) the effect of warming on biomass responses to drought (for panel D, positive values indicate a reduction in drought impact under warming). Values in panel A are means ± 1 SE; while panels B-D are means and 95% confidence intervals. Spp. abbreviations and significance levels follow Fig. 4.

1072

1073