GNS561, a new autophagy inhibitor active against cancer stem cells in hepatocellular carcinoma and hepatic metastasis from colorectal cancer

S. Brun[a, *], J.M. Pascussi[b, 1], E.P. Gifu[c, 1], E. Bestion[a, 1], Z. Macek-Jilkova[d, e, f], G. Wang[c], F. Bassissi[a], S. Mezouar[a], J. Courcambeck[a], P. Merle[c, g], T. Decaens[d, e, f], J. Pannequin[b], P. Halfon[a, 1, *] and C. Caron de Fromentel[c, 1]

[a]Genoscience Pharma, Marseille, France

[b]IGF, University of Montpellier, CNRS, INSERM, Montpellier, France

[c]CRCL, INSERM U1052, CNRS 5286, Université Lyon 1 - Centre Léon Bérard, Lyon, France

[d]Institute for Advanced Biosciences, Research Center UGA, Inserm U 1209, CNRS 5309, La Tronche, France

[e]University of Grenoble Alpes, Faculté de Médecine, France

[f]Clinique Universitaire d'Hépato-gastroentérologie, Pôle Digidune, CHU Grenoble, France

[g]Hepatology and Gastroenterology Unit, Croix-Rousse Hospital, Hospices Civils de Lyon, France

[1]These authors contributed equally

[*]Correspondence to: S. Brun and P. Halfon, Genoscience Pharma, 10 Rue d'Iéna, 13006 Marseille, France; brun.sonia@hotmail.fr and p.halfon@genosciencepharma.com

ABSTRACT

Patients with advanced hepatocellular carcinoma (HCC) or metastatic colorectal cancer (mCRC) have a very poor prognosis due to the lack of efficient treatments. As observed in several other tumors, the effectiveness of treatments is mainly hampered by the presence of a highly tumorigenic subpopulation of cancer cells called cancer stem cells (CSCs). Indeed, CSCs are resistant to chemotherapy and radiotherapy and have the ability to regenerate the tumor bulk. Hence, innovative drugs that are efficient against both bulk tumor cells and CSCs would likely improve cancer treatment. In this study, we demonstrated that GNS561, a new autophagy inhibitor that induces lysosomal cell death, showed significant activity against not only the whole tumor population but also a subpopulation displaying CSC features (high ALDH activity and tumorsphere formation ability) in HCC and in liver mCRC cell lines. These results were confirmed *in vivo* in an HCC-induced cirrhotic rat model in which GNS561 decreased tumor growth and reduced the frequency of CSCs ($CD90^+CD45^-$). Accordingly, GNS561, which was in a global phase 1b clinical trial in liver cancers that was recently successful, offers great promise for cancer therapy by exterminating both the tumor bulk and the CSC subpopulation.

Keywords: GNS561, cancer stem cell, liver cancer, colorectal cancer, lysosome, therapy

1. Introduction

Patients with advanced hepatocellular carcinoma (HCC) or metastatic colorectal cancer (mCRC) have a very poor prognosis (Bray et al. 2018). They suffer a lack of efficient therapy for HCC or treatment failures for CRC, both resulting in low survival rates. Since the late 1990s, a subpopulation of poorly differentiated cancer cells, known as cancer stem cells (CSCs), have raised substantial interest, as they are believed to play a crucial role in cancer initiation, propagation, relapse and metastasis (Bonnet 2005, Vitale et al. 2015, Sun et al. 2016, Nazio et al. 2019). Due to their pluripotency and self-renewal characteristics, CSCs have the ability to regenerate tumor bulk ad infinitum (Clarke et al. 2006). It has also been shown that CSCs are resistant to various types of stresses, including those generated by treatments such as radiotherapy or chemotherapy (Wang and Sun 2018, Zhou et al. 2018). Recently, it was shown that apoptosis was induced in the majority of HCC tumor cells following treatment with sorafenib, a standard of care, whereas a subpopulation of CSCs remained and led to tumor metastases (Chen et al. 2015, Cheng et al. 2019). In the same extent, colorectal CSCs are not sensitive to radiotherapy or chemotherapy with oxaliplatin, irinotecan or 5-FU, leading to tumor metastasis and recurrence after chemotherapy (Das et al. 2020). Thus, innovative drugs that simultaneously exterminate both bulk tumor cells and CSC subpopulation can improve cancer treatment and result in long-term tumor remission (Sun et al. 2016).

We previously reported that GNS561, a new lysosomotropic molecule with high hepatic tropism, was efficient against intrahepatic cholangiocarcinoma and HCC (Brun et al. 2019, Brun et al. 2020). Based on robust preclinical data, the potential use of GNS561 against primary and secondary liver cancers was evaluated in a global phase 1b clinical trial (NCT03316222), which was recently successful (ClinicalTrials.gov). Regarding its mode of action, GNS561 was shown to suppress cancer cell growth *via* the induction of apoptosis resulting from inhibition of palmitoyl-protein thioesterase 1 (PPT1) activity, lysosomal unbound Zn^{2+} accumulation, impairment of cathepsin activity, blockage of autophagic flux,

altered localization of mTOR, lysosomal membrane permeabilization and caspase activation

(Brun et al. 2020).

The inhibition of autophagy induced by GNS561 suggests that this molecule could be

effective killing CSCs. Indeed, autophagy, as a catabolic and prosurvival pathway, appears

to be critical for the survival of cancer cells and especially for the maintenance, plasticity, and

chemoresistance of CSCs and their adaptation to tumor microenvironment changes (TME)

(Ojha et al. 2015, Vitale et al. 2015, Sun et al. 2016, Sharif et al. 2017). An increase in

autophagy levels as a source of nutrient replenishment was reported in cancer cells and

CSCs in response to increased metabolic demands triggered by environmental stress

signals, such as hypoxia, starvation, metabolic and oxidative stresses and therapy (El Hout

et al. 2020, Mandhair et al. 2020). Therefore, autophagy inhibition might be a useful strategy

for targeting and killing both cancer cells and CSCs (Li and Zhu 2019, Nazio et al. 2019,

Smith and Macleod 2019, Wu et al. 2020).

In this study, we investigated the effect of GNS561 on CSCs. We first showed that *in vitro,*

GNS561 induced a dramatic increase of CSC death in both HCC and liver mCRC cell lines.

We then confirmed its efficacy on CSCs *in vivo* in an HCC-induced cirrhotic rat model.

2. Materials and methods

Details of the materials and methods are provided in the Supplementary Methods.

2.1. Cell lines culture

Hep3B cells were maintained in EMEM (ATCC) supplemented with 10% heat-inactivated fetal bovine serum (FBS) (Invitrogen) and 1% penicillin/streptomycin (Gibco). Huh7 cells were cultured in DMEM-F12 medium (Gibco) supplemented with 10% heat-inactivated FBS (Invitrogen), 1% nonessential amino acids (Gibco), 1% penicillin/streptomycin (Gibco), 1% GlutaMAX (Invitrogen) and 1 mM sodium pyruvate (Gibco). Liver mCRC patient-derived cell lines (CPP19, 30, 36 and CPP45) (Grillet et al. 2017) were maintained in DMEM (Gibco) with 10% FBS. All cells were cultured in a humidified atmosphere at 37 °C and 5% CO_2.

2.2. Diethylnitrosamine-induced cirrhotic immunocompetent rat model of HCC

Rats with HCC were treated with vehicle, GNS561 or sorafenib, by oral gavage for a period of six weeks. The animals were checked daily for clinical signs, effects of tumor growth and any other abnormal effects. All rats received humane care in accordance with the Guidelines on the Humane Treatment of Laboratory Animals (Directive 2010/63/EU), and experiments were approved by the animal Ethics Committee: GIN Ethics Committee No. 004.

2.3. Statistical analysis

Statistical analyses were performed using Prism 8.4.3 software (GraphPad Software, Inc.). For datasets with a normal distribution, multiple comparisons were performed using one-way ANOVA with Dunnett's post hoc analysis. For datasets without normal distribution, multiple comparisons were performed using the non-parametric Kruskal-Wallis test. Data are presented as the mean values ± standard error mean (SEM) of three independent

109   experiments unless stated otherwise. Statistical significance was defined as a P-value < 0.05

110   and is indicated by an asterisk in all figures.

3. Results

3.1. GNS561 is efficient against the whole tumor

We first investigated the activity of GNS561 in the whole population of two HCC cell lines and four liver mCRC cell lines. GNS561 decreased viability in a dose-dependent manner, with very similar IC_{50} values for all the tested cell lines and < 3 µM (Table 1).

	Name	IC_{50} (µM)	95% confidence interval
HCC cell lines	Hep3B	2.33	2.358 to 2.535
	Huh7	1.48	1.238 to 1.699
Liver mCRC cell lines	CPP19	1.88	1.424 to 2.487
	CPP30	1.22	0.9210 to 1.624
	CPP36	1.45	0.6990 to 3.025
	CPP45	1.12	0.9555 to 1.318

Table 1: *In vitro* activity (IC_{50}) of GNS561 on two HCC (Hep3B and Huh7) and four liver mCRC cell lines (CPP19, CPP30, CPP36 and CPP45) after 72 h of treatment with GNS561.

3.2. GNS561 is also active against a subpopulation displaying CSC features

Aldehyde dehydrogenase (ALDH) is known to be a marker of CSCs in numerous solid cancers, including CRC (Huang et al. 2009, Vassalli 2019). To evaluate the sensitivity to GNS561 of the CSC/cancer progenitor cell-enriched subpopulations of liver mCRC cell lines, the percentage of ALDH^{bright} cells was quantified by flow cytometry after 72 h of incubation with GNS561. As shown in Fig. 1A, GNS561 significantly decreased the percentage of ALDH^{bright} cells in a dose-dependent manner in all three mCRC cell lines.

Unlike CRC cells, ALDH-overexpressing HCC cells seem to exhibit a differentiated rather

than a CSC phenotype (Tanaka et al. 2015). Moreover, there is a lack of consensus

biomarkers for CSCs in HCC (Flores-Tellez et al. 2017, Wu et al. 2020). Consequently, the

CSC subpopulation is commonly identified in HCC based on functional criteria such as their

capacity to efflux Hoechst 33342 dye (side population) (Chiba et al. 2006) or to form

tumorspheres (Uchida et al. 2010, Cao et al. 2011), as only these cells have the ability to

proliferate in nonadherent conditions (Johnson et al. 2013, Ma et al. 2019). Tumorsphere

formation was used to isolate the CSC subpopulation of HCC cell lines, the immature

phenotype of which was confirmed by the overexpression of the stemness-associated factors

Sox2, Oct4, Klf4 or Nanog compared with the whole cell population (data not shown). To

avoid chemoresistance because cells within the interior of tumorspheres are protected from

drug penetration by neighboring cells on the periphery of the sphere, we compared the

impact of GNS561 between the whole and CSC-enriched populations of Hep3B and HuH7

cell lines in monolayer cultures. GNS561 had the same effect on the viability of the two

populations from Hep3B (Fig.1B) or Huh7 (Fig.1C) cells treated in monolayer cultures for 48

or 72 h. Since cells with an immature phenotype have high plasticity when cultured *in vitro*,

we hypothesized that they could have derived after 48 or 72 h of monolayer culture and

spontaneously restored the initial heterogeneous population (Clarke et al. 2006). Thus, we

repeated the experiment in Hep3B cells and assessed the antitumor activity of GNS561 at

shorter times (6, 8 and 24 h) and at higher doses (0.5 to 32 µM). Similarly, GNS561 was as

efficient against the bulk population as the CSC-enriched population (Supplementary Fig. 1).

Fig. 1: GNS561 is efficient against subpopulation displaying CSC features. (A) Box and whisker representation (min to max) of the percentage of ALDH[bright] cells after 72 h of treatment with GNS561 in three liver mCRC cell lines with the indicated concentrations. For comparison with the control, one-way ANOVA with Dunnett's post hoc analysis was performed. *, $P < 0.05$. Antitumor activity of GNS561 on whole tumor (circle) or CSC-enriched populations (square) in Hep3B (B) and Huh7 (C) HCC cell lines after 48 or 72 h of treatment with GNS561. The curves represent the means of two independent experiments in triplicate.

3.3. GNS561 blocks tumorsphere formation

Furthermore, the ability of GNS561 to impair tumorsphere formation was investigated in both

liver mCRC and HCC cell lines. GNS561 (0.3 or 3 µM) was added after seeding a low

number of liver mCRC cells in CSC medium. Ten days later, tumorspheres > 50 µm were

counted. As shown in Fig. 2A and 2B, GNS561 induced significant dose-dependent

decreases in tumorspheres in all three cell lines, with complete suppression of tumorspheres

at the highest dose (3 µM). Similar results were obtained with the Hep3B HCC cell line at 96

h (Fig. 2A and B) and 11 days (data not shown) of GNS561 treatment. As illustrated in Fig. 2,

GNS561 completely inhibited tumorsphere formation, even at low concentration (0.8 µM

corresponding to 1/3 IC_{50} in the adherent whole population). These results showed that

GNS561 has an antitumor effect not only on the bulk, but also on the CSC-enriched

subpopulation. In order to further enrich the CSC population, secondary spheres were

generated from primary Hep3B spheres and treated with the same doses of GNS561. A

slight difference in sensitivity was observed between primary and secondary spheres, with a

complete inhibition of sphere formation reached at 0.8 and 1.6 µM, respectively (Fig 2B and

Supplementary Fig.2). These results confirm the significant activity of GNS561 on CSCs. In

contrast, sorafenib had only a mild effect on the CSC-enriched subpopulation since it failed

to completely abolish tumorsphere formation whether from adherent cells or primary spheres

(Supplementary Fig. 2 and 3).

A

Fig. 2: GNS561 alters the self-renewal of liver mCRC and HCC cell lines. (A) Tumorsphere forming efficiency of three liver mCRC (CPP19, CPP30 and CPP36) and one HCC (Hep3B) cell lines treated with the indicated GNS561 concentrations. Scale bars represent 50 µm. (B) Number of spheres > 50 µm. For comparison with the control, statistical significance was determined using one-way ANOVA and Dunnett's post hoc test. *, P < 0.05.

3.4. GNS561 decreases tumor growth and CSCs in a diethylnitrosamine-induced cirrhotic rat model of HCC

As demonstrating the antitumor activity of a drug on CSCs by tumorsphere formation in vitro

could reflect it *in vivo* anti-tumor efficiency (Lee et al. 2016), we investigated its efficacy on

CSCs in vivo.

GNS561 has been previously shown to have potent antitumor activity in two HCC *in vivo*

models including a diethylnitrosamine-induced cirrhotic rat model of HCC (Brun et al. 2020).

Here, we used the same model to study the impact of GNS561 on the frequency of CSCs *in*

vivo. Rats with already developed HCC were either treated with vehicle, sorafenib (a

standard of care of HCC) at 10 mg/kg or GNS561 at 15 mg/kg. The $CD90^+CD45^-$ phenotype

was used to identify and quantify CSCs, since it has been reported to be associated with

CSCs in liver tumor tissues and with circulating CSCs in blood samples (Yang et al. 2008,

Yang et al. 2008, Motawi et al. 2016). In our rat model, no significant decrease was observed

in the frequency of CSCs from the liver tumor tissue between the GNS561-treated group and

the control group (Fig. 3A). However, the decrease in circulating CSCs became significant in

the GNS561-treated group compared with the control group in the circulating CSCs (Fig. 3B).

Notably, sorafenib, which is widely used to treat advanced HCC in humans, had no

significant effect on the percentage of CSCs.

Fig. 3: GNS561 decreases CSC frequency *in vivo*. Frequency of CSCs in liver tumor (A)

and blood (B) samples in a diethylnitrosamine-induced cirrhotic rat model of HCC. Rats

received vehicle (control), sorafenib at 10 mg/kg or GNS561 at 15 mg/kg. Values are means

± SEM. For comparison with the control nontreated group, the non-parametric Kruskal-Wallis

test was performed. *, P < 0.05.

4. Discussion

HCC and mCRC are refractory to most conventional chemotherapies. One of the reasons for

the high mortality rate is the existence of a small population of CSCs with high tumorigenicity

and resistance to drugs. Accordingly, many cancer therapies, while killing the bulk of tumor

cells, may ultimately fail because they do not eliminate CSCs that reform heterogeneous

tumors.

GNS561 is a small molecule that showed potent antitumor activity *in vitro* in a panel of

human cancer cells, including HCC patient-derived cells, and *in vivo* in two HCC different

models (Brun et al. 2020). The *in vitro* studies performed in this study clearly showed that

GNS561 displays potent dose response activity when assayed against a panel of human

colon tumor cells isolated from liver metastases. In addition, GNS561 was efficient against

subpopulation displaying ALDH^{bright} activity and produced a dramatic decrease in the ability of

these cells to form tumorspheres. Similarly, in two HCC cell lines, GNS561 was equally

effective in both populations (bulk and CSCs) and dramatically decreased the cell capacity to

form spheres, even at low doses, unlike sorafenib, a standard of care. This latter observation

was in agreement with the literature. Indeed, CSC resistance to sorafenib has been widely

reported and has been cited as responsible for treatment failure and relapse (Cao et al.

2019, Feng et al. 2020).

Taken together, our results strongly suggest that GNS561 was efficient not only against the

whole tumor population but also against cell subpopulations displaying CSC features

(ALDH^{bright} cells and tumorsphere initiating cells). These original results were confirmed *in*

vivo, in an HCC-induced cirrhotic rat model, in which GNS561 showed antitumor effects and

reduced the frequency of CSCs in liver tumor tissue and circulating CSCs. To go further, it

would be informative to xenograft tumorsphere-derived CSCs or serial transplant tumor to

determine GNS561 activity on the tumor growth.

The antitumor effect of GNS561 on tumor bulk cells and on CSCs can be explained by its

capacity to inhibit autophagy and lysosomal functions and to disrupt lysosomes (Brun et al.

2020). In fact, once established, tumors require an uninterrupted nutritional supply for

maintaining their proliferative needs in stressful conditions such as hypoxia, nutrient and

growth factor starvation, and oxidative and metabolic stress. (Ojha et al. 2015). In this

context, autophagy has a protumoral activity by providing recycled bioenergetic substrates

and consequently overcoming nutritional deficiency and favoring the survival of cancer cells.

Moreover, in the CSC population, autophagy was shown to be a major factor for the

preservation of cell homeostasis and the maintenance of stemness properties (Vitale et al.

2015). Consistently, it was reported that metastatic cells and CSCs are vulnerable to

lysosomal inhibition and disruption (Salerno et al. 2014, Morgan et al. 2018, Cash et al.

2020, Versini et al. 2020). For example, inhibition of autophagic flux and lysosomal functions

by salinomycin interferes with the maintenance and expansion of breast cancer stem-

like/progenitor cells (Yue et al. 2013). Mefloquine showed activity on acute myeloid leukemia

cells and progenitors by disrupting lysosomes (lysosomal membrane permeabilization and

cytosolic release of cathepsins) (Sukhai et al. 2013).

Further research will be necessary to better understand the underlying mechanisms

implicated in GNS561 activity against CSCs. It would be interesting to evaluate whether

GNS561 treatment results from GNS561-induced inhibition of PPT1 and the mTOR pathway

in CSCs, as demonstrated in whole tumors (Brun et al., 2020). In fact, knockout of PPT1 in

tumor cells inhibited tumor growth, tumorsphere formation and tumorigenesis *in vivo*

(Rebecca et al. 2019), suggesting that PPT1 may be implicated in CSC maintenance. In turn,

the mTOR pathway has been shown to be one of the most important pathways involved in

the development and progression of cancer and in the maintenance and hallmarks of CSCs

(Xia and Xu 2015), implying that inhibition of the mTOR pathway could be a good therapeutic

strategy to eradicate CSCs. Moreover, as autophagy is an adaptive mechanism modulating

the TME surrounding CSCs to support their stemness and cancer propagation (Mandhair et

al. 2020), studying the effect of GNS561 on the TME would be of broad interest.

In conclusion, GNS561 showed activity against not only the whole tumor population but also

against subpopulations displaying CSC features *in vitro* and *in vivo*. Thus, a strategy that

simultaneously exterminates the CSC subpopulation and tumor bulk by using autophagy

inhibitors and lysosomal disruptors, such as GN561, is promising as a new direction to

achieve cure and to prevent relapse of liver primary cancer and metastasis. Moreover, as

many tumor types have been reported to contain CSCs, GNS561 could be useful to target

these CSCs in a wide variety of human tumors.

Author contributions

S. Brun: conceptualization, methodology, visualization, project administration, writing –
original draft preparation, writing – review and editing; J.M. Pascussi: investigation, formal
analysis, writing – review and editing; E.P. Gifu: investigation, formal analysis; E. Bestion:
investigation, formal analysis; Z. Macek-Jilkova: investigation, formal analysis; G. Wang:
investigation; F. Bassissi: conceptualization, methodology, supervision; S. Mezouar:
investigation; J. Courcambeck: resources; P. Merle: supervision; T. Decaens: supervision; J.
Pannequin: supervision, writing – review and editing; P. Halfon: funding acquisition,
supervision, writing – review and editing; C. Caron de Fromentel: conceptualization,
methodology, supervision, writing – review and editing

Disclosure of potential conflicts of interest

SB, EB, FB, SM, JC and PH are employees of Genoscience Pharma. SB, FB and PH are
shareholders of Genoscience Pharma. SB, FB, JC and PH are co-inventors of a pending
patent. The other authors declare that they have no conflicts of interest to report.

Funding

This research did not receive any specific grant from funding agencies in the public,
commercial, or not-for-profit sectors.

Acknowledgement

The authors are very grateful to Dr. Michel Prudhomme and Dr. Jean-François Bourgaux
from CHU Carémeau, Nîmes, France, for the use of liver mCRC cells.

Appendix A. Supplementary data

Supplementary data for this article can be found online.

References

Bonnet, D. (2005). Cancer stem cells: lessons from leukaemia. Cell Prolif 38(6): 357-361.

Bray, F., Ferlay, J., Soerjomataram, I., Siegel, R.L., Torre, L.A. and Jemal, A. (2018). Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. CA: a cancer journal for clinicians 68(6): 394-424.

Brun, S., Bassissi, F., Serdjebi, C., Novello, M., Tracz, J., Autelitano, F., Guillemot, M., Fabre, P., Courcambeck, J., Ansaldi, C., et al. (2019). GNS561, a new lysosomotropic small molecule, for the treatment of intrahepatic cholangiocarcinoma. Invest New Drugs 37(6): 1135-1145.

Brun, S., Raymond, E., Bassissi, F., Jilkova, Z.M., Mezouar, S., Rachid, M., Novello, M., Tracz, J., Hamaï, A., Lalmanach, G., et al. (2020). GNS561, a clinical-stage PPT1 inhibitor, has powerful antitumor activity against hepatocellular carcinoma via modulation of lysosomal functions. bioRxiv.

Cao, J., Zhao, M., Liu, J., Zhang, X., Pei, Y., Wang, J., Yang, X., Shen, B. and Zhang, J. (2019). RACK1 Promotes Self-Renewal and Chemoresistance of Cancer Stem Cells in Human Hepatocellular Carcinoma through Stabilizing Nanog. Theranostics 9(3): 811-828.

Cao, L., Zhou, Y., Zhai, B., Liao, J., Xu, W., Zhang, R., Li, J., Zhang, Y., Chen, L., Qian, H., et al. (2011). Sphere-forming cell subpopulations with cancer stem cell properties in human hepatoma cell lines. BMC Gastroenterol 11: 71.

Cash, T.P., Alcala, S., Rico-Ferreira, M.D.R., Hernandez-Encinas, E., Garcia, J., Albarran, M.I., Valle, S., Munoz, J., Martinez-Gonzalez, S., Blanco-Aparicio, C., et al. (2020). Induction of Lysosome Membrane Permeabilization as a Therapeutic Strategy to Target Pancreatic Cancer Stem Cells. Cancers (Basel) 12(7).

Chen, J., Jin, R., Zhao, J., Liu, J., Ying, H., Yan, H., Zhou, S., Liang, Y., Huang, D., Liang, X., et al. (2015). Potential molecular, cellular and microenvironmental mechanism of sorafenib resistance in hepatocellular carcinoma. Cancer Lett 367(1): 1-11.

Cheng, C.C., Chao, W.T., Liao, C.C., Shih, J.H., Lai, Y.S., Hsu, Y.H. and Liu, Y.H. (2019). The Roles Of Angiogenesis And Cancer Stem Cells In Sorafenib Drug Resistance In Hepatocellular Carcinoma. Onco Targets Ther 12: 8217-8227.

Chiba, T., Kita, K., Zheng, Y.W., Yokosuka, O., Saisho, H., Iwama, A., Nakauchi, H. and Taniguchi, H. (2006). Side population purified from hepatocellular carcinoma cells harbors cancer stem cell-like properties. Hepatology 44(1): 240-251.

Clarke, M.F., Dick, J.E., Dirks, P.B., Eaves, C.J., Jamieson, C.H., Jones, D.L., Visvader, J., Weissman, I.L. and Wahl, G.M. (2006). Cancer stem cells--perspectives on current status and future directions: AACR Workshop on cancer stem cells. Cancer Res 66(19): 9339-9344.

ClinicalTrials.gov. " Study of GNS561 in Patients With Liver Cancer." ClinicalTrials.gov.

Das, P.K., Islam, F. and Lam, A.K. (2020). The Roles of Cancer Stem Cells and Therapy Resistance in Colorectal Carcinoma. Cells 9(6).

El Hout, M., Cosialls, E., Mehrpour, M. and Hamai, A. (2020). Crosstalk between autophagy and metabolic regulation of cancer stem cells. Mol Cancer 19(1): 27.

Feng, Y., Jiang, W., Zhao, W., Lu, Z., Gu, Y. and Dong, Y. (2020). miR-124 regulates liver
cancer stem cells expansion and sorafenib resistance. Exp Cell Res 394(2): 112162.

Flores-Tellez, T.N., Villa-Trevino, S. and Pina-Vazquez, C. (2017). Road to stemness in
hepatocellular carcinoma. World J Gastroenterol 23(37): 6750-6776.

Grillet, F., Bayet, E., Villeronce, O., Zappia, L., Lagerqvist, E.L., Lunke, S., Charafe-Jauffret,
E., Pham, K., Molck, C., Rolland, N., et al. (2017). Circulating tumour cells from patients with
colorectal cancer have cancer stem cell hallmarks in ex vivo culture. Gut 66(10): 1802-1810.

Huang, E.H., Hynes, M.J., Zhang, T., Ginestier, C., Dontu, G., Appelman, H., Fields, J.Z.,
Wicha, M.S. and Boman, B.M. (2009). Aldehyde dehydrogenase 1 is a marker for normal
and malignant human colonic stem cells (SC) and tracks SC overpopulation during colon
tumorigenesis. Cancer Res 69(8): 3382-3389.

Johnson, S., Chen, H. and Lo, P.K. (2013). In vitro Tumorsphere Formation Assays. Bio
Protoc 3(3).

Lee, C.-H., Yu, C.-C., Wang, B.-Y. and Chang, W.-W. (2016). Tumorsphere as an effective in
vitro platform for screening anti-cancer stem cell drugs. Oncotarget 7(2): 1215-1226.

Li, N. and Zhu, Y. (2019). Targeting liver cancer stem cells for the treatment of hepatocellular
carcinoma. Therap Adv Gastroenterol 12: 1756284818821560.

Ma, X.L., Sun, Y.F., Wang, B.L., Shen, M.N., Zhou, Y., Chen, J.W., Hu, B., Gong, Z.J.,
Zhang, X., Cao, Y., et al. (2019). Sphere-forming culture enriches liver cancer stem cells and
reveals Stearoyl-CoA desaturase 1 as a potential therapeutic target. BMC Cancer 19(1): 760.

Mandhair, H.K., Arambasic, M., Novak, U. and Radpour, R. (2020). Molecular modulation of
autophagy: New venture to target resistant cancer stem cells. World J Stem Cells 12(5): 303-
322.

Morgan, M.J., Fitzwalter, B.E., Owens, C.R., Powers, R.K., Sottnik, J.L., Gamez, G.,
Costello, J.C., Theodorescu, D. and Thorburn, A. (2018). Metastatic cells are preferentially
vulnerable to lysosomal inhibition. Proc Natl Acad Sci U S A 115(36): E8479-E8488.

Motawi, T.K., El-Boghdady, N.A., El-Sayed, A.M. and Helmy, H.S. (2016). Comparative study
of the effects of PEGylated interferon-alpha2a versus 5-fluorouracil on cancer stem cells in a
rat model of hepatocellular carcinoma. Tumour Biol 37(2): 1617-1625.

Nazio, F., Bordi, M., Cianfanelli, V., Locatelli, F. and Cecconi, F. (2019). Autophagy and
cancer stem cells: molecular mechanisms and therapeutic applications. Cell Death Differ
26(4): 690-702.

Ojha, R., Bhattacharyya, S. and Singh, S.K. (2015). Autophagy in Cancer Stem Cells: A
Potential Link Between Chemoresistance, Recurrence, and Metastasis. Biores Open Access
4(1): 97-108.

Rebecca, V.W., Nicastri, M.C., Fennelly, C., Chude, C.I., Barber-Rotenberg, J.S., Ronghe,
A., McAfee, Q., McLaughlin, N.P., Zhang, G., Goldman, A.R., et al. (2019). PPT1 Promotes
Tumor Growth and Is the Molecular Target of Chloroquine Derivatives in Cancer. Cancer
Discov 9(2): 220-229.

Salerno, M., Avnet, S., Bonuccelli, G., Hosogi, S., Granchi, D. and Baldini, N. (2014). Impairment of lysosomal activity as a therapeutic modality targeting cancer stem cells of embryonal rhabdomyosarcoma cell line RD. PLoS One 9(10): e110340.

Sharif, T., Martell, E., Dai, C., Kennedy, B.E., Murphy, P., Clements, D.R., Kim, Y., Lee, P.W. and Gujar, S.A. (2017). Autophagic homeostasis is required for the pluripotency of cancer stem cells. Autophagy 13(2): 264-284.

Smith, A.G. and Macleod, K.F. (2019). Autophagy, cancer stem cells and drug resistance. J Pathol 247(5): 708-718.

Sukhai, M.A., Prabha, S., Hurren, R., Rutledge, A.C., Lee, A.Y., Sriskanthadevan, S., Sun, H., Wang, X., Skrtic, M., Seneviratne, A., et al. (2013). Lysosomal disruption preferentially targets acute myeloid leukemia cells and progenitors. J Clin Invest 123(1): 315-328.

Sun, J.H., Luo, Q., Liu, L.L. and Song, G.B. (2016). Liver cancer stem cell markers: Progression and therapeutic implications. World J Gastroenterol 22(13): 3547-3557.

Tanaka, K., Tomita, H., Hisamatsu, K., Nakashima, T., Hatano, Y., Sasaki, Y., Osada, S., Tanaka, T., Miyazaki, T., Yoshida, K., et al. (2015). ALDH1A1-overexpressing cells are differentiated cells but not cancer stem or progenitor cells in human hepatocellular carcinoma. Oncotarget 6(28): 24722-24732.

Uchida, Y., Tanaka, S., Aihara, A., Adikrisna, R., Yoshitake, K., Matsumura, S., Mitsunori, Y., Murakata, A., Noguchi, N., Irie, T., et al. (2010). Analogy between sphere forming ability and stemness of human hepatoma cells. Oncol Rep 24(5): 1147-1151.

Vassalli, G. (2019). Aldehyde Dehydrogenases: Not Just Markers, but Functional Regulators of Stem Cells. Stem Cells Int 2019: 3904645.

Versini, A., Colombeau, L., Hienzsch, A., Gaillet, C., Retailleau, P., Debieu, S., Muller, S., Caneque, T. and Rodriguez, R. (2020). Salinomycin Derivatives Kill Breast Cancer Stem Cells by Lysosomal Iron Targeting. Chemistry 26(33): 7416-7424.

Vitale, I., Manic, G., Dandrea, V. and De Maria, R. (2015). Role of autophagy in the maintenance and function of cancer stem cells. Int J Dev Biol 59(1-3): 95-108.

Wang, K. and Sun, D. (2018). Cancer stem cells of hepatocellular carcinoma. Oncotarget 9(33): 23306-23314.

Wu, Y., Zhang, J., Zhang, X., Zhou, H., Liu, G. and Li, Q. (2020). Cancer Stem Cells: A Potential Breakthrough in HCC-Targeted Therapy. Front Pharmacol 11: 198.

Xia, P. and Xu, X.Y. (2015). PI3K/Akt/mTOR signaling pathway in cancer stem cells: from basic research to clinical application. Am J Cancer Res 5(5): 1602-1609.

Yang, Z.F., Ho, D.W., Ng, M.N., Lau, C.K., Yu, W.C., Ngai, P., Chu, P.W., Lam, C.T., Poon, R.T. and Fan, S.T. (2008). Significance of CD90+ cancer stem cells in human liver cancer. Cancer Cell 13(2): 153-166.

Yang, Z.F., Ngai, P., Ho, D.W., Yu, W.C., Ng, M.N., Lau, C.K., Li, M.L., Tam, K.H., Lam, C.T., Poon, R.T., et al. (2008). Identification of local and circulating cancer stem cells in human liver cancer. Hepatology 47(3): 919-928.

461 Yue, W., Hamai, A., Tonelli, G., Bauvy, C., Nicolas, V., Tharinger, H., Codogno, P. and
462 Mehrpour, M. (2013). Inhibition of the autophagic flux by salinomycin in breast cancer stem-
463 like/progenitor cells interferes with their maintenance. Autophagy 9(5): 714-729.
464
465 Zhou, Y., Xia, L., Wang, H., Oyang, L., Su, M., Liu, Q., Lin, J., Tan, S., Tian, Y., Liao, Q., et
466 al. (2018). Cancer stem cells in progression of colorectal cancer. Oncotarget 9(70): 33403-
467 33415.
468