GNS561, a New Autophagy Inhibitor Active against Cancer Stem Cells in Hepatocellular Carcinoma and Hepatic Metastasis from Colorectal Cancer

Sonia Brun[1, *], Jean-Marc Pascussi[2, #], Elena Patricia Gifu[3, #], Eloïne Bestion[1, #], Zuzana Macek-Jilkova[4, 5, 6], Guanxiong Wang[3], Firas Bassissi[1], Soraya Mezouar[1], Jérôme Courcambeck[1], Philippe Merle[3, 7], Thomas Decaens[4, 5, 6], Julie Pannequin[2], Philippe Halfon[1, #, *] and Claude Caron de Fromentel[3, #]

[1]Genoscience Pharma, Marseille, France

[2]IGF, University of Montpellier, CNRS, INSERM, Montpellier, France

[3]CRCL, INSERM U1052, CNRS 5286, Université Lyon 1 - Centre Léon Bérard, Lyon, France

[4]Institute for Advanced Biosciences, Research Center UGA, Inserm U 1209, CNRS 5309, La Tronche, France

[5]University of Grenoble Alpes, Faculté de Médecine, France

[6]Clinique Universitaire d'Hépato-gastroentérologie, Pôle Digidune, CHU Grenoble, France

[7]Hepatology and Gastroenterology Unit, Croix-Rousse Hospital, Hospices Civils de Lyon, France

[#]These authors contributed equally

[*]Correspondence to: S. Brun and P. Halfon, Genoscience Pharma, 10 Rue d'Iéna, 13006 Marseille, France; Telephone: +33 4 91 26 99 50, E-mail: brun.sonia@hotmail.fr and p.halfon@genosciencepharma.com

ABSTRACT

Patients with advanced hepatocellular carcinoma (HCC) or metastatic colorectal cancer (mCRC) have a very poor prognosis due to the lack of efficient treatments. As observed in several other tumors, the effectiveness of treatments is mainly hampered by the presence of a highly tumorigenic sub-population of cancer cells called cancer stem cells (CSCs). Indeed, CSCs are resistant to chemotherapy and radiotherapy and have the ability to regenerate the tumor bulk. Hence, innovative drugs that are efficient against both bulk tumor cells and CSCs would likely improve cancer treatment. In this study, we demonstrated that GNS561, a new autophagy inhibitor that induces lysosomal cell death, showed significant activity against not only the whole tumor population but also a sub-population displaying CSC features (high ALDH activity and tumorsphere formation ability) in HCC and in liver mCRC cell lines. These results were confirmed *in vivo* in HCC from a DEN-induced cirrhotic rat model in which GNS561 decreased tumor growth and reduced the frequency of CSCs (CD90$^+$CD45$^-$). Accordingly, GNS561, which was in a global phase 1b clinical trial in liver cancers that was recently successful, offers great promise for cancer therapy by exterminating both the tumor bulk and the CSC sub-population.

Keywords: GNS561, cancer stem cell, liver cancer, colorectal cancer, lysosome, therapy

Introduction

Patients with advanced hepatocellular carcinoma (HCC) or metastatic colorectal cancer (mCRC) have a very poor prognosis [1]. They suffer from a lack of efficient therapy for HCC or treatment failures for CRC, both resulting in low survival rates. Since the late 1990s, a sub-population of poorly differentiated cancer cells, known as cancer stem cells (CSCs), have raised substantial interest, as they are believed to play a crucial role in cancer initiation, propagation, relapse and metastasis [2–5]. According to the stem cell paradigm, CSCs are able to both self-renew and give rise to progenitors thus to generate the bulk of a tumor ad infinitum [6]. This model has been validated for several tumor types, including HCC and CRC, whereas other ones, such as melanoma [7], do not exactly follow this pattern [8]. It has also been shown that CSCs are resistant to various types of stress, including those generated by treatments such as radiotherapy or chemotherapy [9,10]. Recently, it was shown that apoptosis was induced in the majority of HCC tumor cells following treatment with sorafenib, a standard of care, whereas a subpopulation of CSCs remained and led to tumor metastases [11,12]. In the same line, colorectal CSCs are not sensitive to radiotherapy or chemotherapy with oxaliplatin, irinotecan or 5-FU, leading to tumor metastasis and recurrence after chemotherapy [13]. Thus, innovative drugs that simultaneously exterminate both bulk tumor cells and CSC subpopulation can improve cancer treatment and result in long-term tumor remission [4,8].

We previously reported that GNS561, a new lysosomotropic molecule with high hepatic tropism, was efficient against intrahepatic cholangiocarcinoma and HCC [14,15]. Based on robust preclinical data, the potential use of GNS561 against primary and secondary liver cancers was evaluated in a global phase 1b clinical trial (NCT03316222), which was recently successful [16]. Regarding its mode of action, GNS561 was shown to suppress cancer cell growth *via* the induction of apoptosis resulting from inhibition of palmitoyl-protein thioesterase 1 (PPT1) activity, lysosomal unbound Zn^{2+} accumulation, impairment of cathepsin activity,

blockage of autophagic flux, altered localization of mTOR, lysosomal membrane

permeabilization and caspase activation [14].

The inhibition of autophagy induced by GNS561 suggests that this molecule could be

effective in killing CSCs. Indeed, autophagy, as a catabolic and prosurvival pathway, appears

to be critical for the survival of cancer cells and especially for the maintenance, plasticity, and

chemoresistance of CSCs and their adaptation to tumor microenvironment changes (TME)

[3,5,17,18]. An increase in autophagy levels as a source of nutrient replenishment was

reported in cancer cells and CSCs in response to increased metabolic demands triggered by

environmental stress signals, such as hypoxia, starvation, metabolic and oxidative stresses

and therapy [19,20]. Therefore, autophagy inhibition might be a useful strategy for targeting

and killing both cancer cells and CSCs [3,21–23].

In this study, we investigated the effect of GNS561 on CSCs. We first showed that *in vitro,*

GNS561 induced a dramatic increase of CSC death in both HCC and liver mCRC cell lines.

We then confirmed its efficacy on CSCs *in vivo* in an HCC-induced cirrhotic rat model.

Materials and Methods

Details of the materials and methods are provided in the Supplementary Methods.

Cell line culture

Hep3B cells were maintained in EMEM (ATCC) supplemented with 10% heat-inactivated fetal bovine serum (FBS) (Invitrogen) and 1% penicillin/streptomycin (Gibco). Huh7 cells were cultured in DMEM-F12 medium (Gibco) supplemented with 10% heat-inactivated FBS (Invitrogen), 1% nonessential amino acids (Gibco), 1% penicillin/streptomycin (Gibco), 1% GlutaMAX (Invitrogen) and 1 mM sodium pyruvate (Gibco). Liver mCRC patient-derived cell lines (CPP19, 30, 36 and CPP45) [24] were maintained in DMEM (Gibco) with 10% FBS. All cells were cultured in a humidified atmosphere at 37 °C and 5% CO_2.

Diethylnitrosamine-induced cirrhotic immunocompetent rat model of HCC

Rats with HCC were treated with vehicle, GNS561 or sorafenib, by oral gavage for a period of six weeks. The animals were checked daily for clinical signs, effects of tumor growth and any other abnormal effects. All rats received humane care in accordance with the Guidelines on the Humane Treatment of Laboratory Animals (Directive 2010/63/EU), and experiments were approved by the animal Ethics Committee: GIN Ethics Committee No. 004.

Statistical analysis

Statistical analyses were performed using Prism 8.4.3 software (GraphPad Software, Inc.). For datasets with a normal distribution, multiple comparisons were performed using one-way ANOVA with Dunnett's post hoc analysis. For datasets without normal distribution, multiple comparisons were performed using the non-parametric Kruskal-Wallis test. Data are presented as the mean values ± standard error mean (SEM) of three independent

experiments unless stated otherwise. Statistical significance is indicated by one (P-value <

0.05) or two (P-value < 0.01) asterisks on the figures.

Results

GNS561 is efficient against the whole tumor

We first investigated the activity of GNS561 in the whole population of two HCC cell lines and

four liver mCRC cell lines. GNS561 decreased viability in a dose-dependent manner, with

very similar maximal inhibitory concentration (IC_{50}) values for all the tested cell lines and < 3

µM (Table 1).

	Name	IC_{50} (µM)	95% confidence interval
HCC cell lines	Hep3B	2.33	2.358 to 2.535
	Huh7	1.48	1.238 to 1.699
Liver mCRC cell lines	CPP19	1.88	1.424 to 2.487
	CPP30	1.22	0.9210 to 1.624
	CPP36	1.45	0.6990 to 3.025
	CPP45	1.12	0.9555 to 1.318

Table 1. *In vitro* activity (IC_{50}) of GNS561 on two HCC (Hep3B and Huh7) and four liver

mCRC cell lines (CPP19, CPP30, CPP36 and CPP45) after 72 h of treatment with GNS561.

GNS561 is also active against a subpopulation displaying CSC features

Aldehyde dehydrogenase (ALDH) is known to be a marker of CSCs in numerous solid

cancers, including CRC [25,26]. To evaluate the sensitivity to GNS561 of the CSC/cancer

progenitor cell-enriched subpopulations of liver mCRC cell lines, the percentage of ALDH[bright]

cells was quantified by flow cytometry after 72 h of incubation with GNS561. As shown in Fig.

1A, GNS561 significantly decreased the percentage of ALDH[bright] cells in a dose-dependent

manner in all three mCRC cell lines.

Unlike CRC cells, ALDH-overexpressing HCC cells seem to exhibit a differentiated rather

than a CSC phenotype [27]. Moreover, there is a lack of consensus biomarkers for CSCs in

HCC [23,28]. Consequently, the CSC subpopulation is commonly identified in HCC based on

functional criteria such as their capacity to efflux Hoechst 33342 dye (side population) [29] or

to form tumorspheres [30,31], as only these cells have the ability to proliferate in non-

adherent conditions [32,33]. Tumorsphere formation assay was used to isolate the CSC

subpopulation of HCC cell lines. The significant overexpression of the stemness-associated

factors *SOX2* and *NANOG* in Hep3B tumorspheres compared to the whole cell population

confirmed their immature phenotype (Fig. S1). To avoid chemoresistance because cells

within the interior of tumorspheres are protected from drug penetration by neighboring cells

on the periphery of the sphere, we compared the impact of GNS561 between the whole and

CSC-enriched populations of Hep3B and HuH7 cell lines in monolayer cultures. GNS561 had

the same effect on the viability of the two populations from Hep3B (Fig.1B) or Huh7 (Fig.1C)

cells treated in monolayer cultures for 48 or 72 h. Since cells with an immature phenotype

have high plasticity when cultured *in vitro*, we hypothesized that they could have derived

after 48 or 72 h of monolayer culture and spontaneously restored the initial heterogeneous

population [6]. Thus, we repeated the experiment in Hep3B cells and assessed the antitumor

activity of GNS561 at shorter times (6, 8 and 24 h) and at higher doses (0.5 to 32 µM).

Similarly, GNS561 was as efficient against the bulk population as the CSC-enriched

population (Fig. S2).

Figure 1. GNS561 is efficient against subpopulation displaying CSC features. (A) Box

and whisker representation (min to max) of the percentage of ALDH[bright] cells after 72 h of

treatment with GNS561 in three liver mCRC cell lines with the indicated concentrations. For

comparison with the control, one-way ANOVA with Dunnett's post hoc analysis was

performed. *, $P < 0.05$. Antitumor activity of GNS561 on whole tumor (circle) or CSC-

enriched populations (square) in Hep3B (B) and Huh7 (C) HCC cell lines after 48 or 72 h of

treatment with GNS561. The curves represent the mean of two independent experiments in

triplicate.

GNS561 blocks tumorsphere formation

Furthermore, the ability of GNS561 to impair tumorsphere formation was investigated in both

liver mCRC and HCC cell lines. GNS561 (0.3 or 3 µM) was added after seeding a low

number of liver mCRC cells in CSC medium. Ten days later, tumorspheres > 50 µm were

counted. As shown in Fig. 2A and 2B, GNS561 induced significant dose-dependent

decreases in tumorspheres in all three cell lines, with complete suppression of tumorspheres

at the highest dose (3 µM). Similar results were obtained with the Hep3B HCC cell line at 96

h (Fig. 2A and B) and 11 days (data not shown) of GNS561 treatment. As illustrated in Fig. 2,

GNS561 completely inhibited tumorsphere formation, even at low concentration (0.8 µM

corresponding to 1/3 IC_{50} in the adherent whole population). These results showed that

GNS561 has an antitumor effect not only on the bulk, but also on the CSC-enriched

subpopulation. In order to further enrich the CSC population, secondary spheres were

generated from primary Hep3B spheres and treated with the same doses of GNS561. A

slight difference in sensitivity was observed between primary and secondary spheres, with a

complete inhibition of sphere formation reached at 0.8 and 1.6 µM, respectively (Fig 2B and

Fig.S3). These results confirm the significant activity of GNS561 on CSCs. In contrast,

sorafenib had only a mild effect on the CSC-enriched subpopulation since it failed to

completely abolish tumorsphere formation whether from adherent cells or primary spheres

(Fig. S3 and S4).

178
Figure 2. GNS561 alters the self-renewal of liver mCRC and HCC cell lines. (A)

Tumorsphere forming efficiency of three liver mCRC (CPP19, CPP30 and CPP36) and one

HCC (Hep3B) cell lines treated with the indicated GNS561 concentrations. Scale bars

represent 50 μm. (B) Number of spheres > 50 μm. For comparison with the control, statistical

significance was determined using one-way ANOVA and Dunnett's post hoc test. *, P < 0.05.

GNS561 decreases tumor growth and CSCs in a diethylnitrosamine-induced cirrhotic rat

model of HCC

As demonstrating the antitumor activity of a drug on CSCs by tumorsphere formation *in vitro*

could reflect its *in vivo* anti-tumor efficiency [34], we investigated its efficacy on CSCs in vivo.

GNS561 has been previously shown to have potent antitumor activity in two HCC *in vivo*

models including a diethylnitrosamine-induced cirrhotic rat model of HCC [14]. Here, we used

the same model to study the impact of GNS561 on the frequency of CSCs *in vivo*. Rats with

already developed HCC were either treated with vehicle, sorafenib (a standard of care of

HCC) at 10 mg/kg or GNS561 at 15 mg/kg. The CD90$^+$CD45$^-$ phenotype was used to identify

and quantify CSCs, since it has been reported to be associated with CSCs in liver tumor

tissues and with circulating CSCs in blood samples [35–37]. In our rat model, no significant

decrease was observed in the frequency of CSCs from the liver tumor tissue between the

GNS561-treated group and the control group (Fig. 3A). However, the decrease in circulating

CSCs became significant in the GNS561-treated group compared with the control group in

the circulating CSCs (p = 0.0067, Fig. 3B). Notably, sorafenib, which is widely used to treat

advanced HCC in humans, had no significant effect on the percentage of CSCs.

Figure 3. GNS561 decreases CSC frequency *in vivo*. Box and whisker representation (min

to max) of the percentage of CSCs in liver tumor (A) and blood (B) samples in a

diethylnitrosamine-induced cirrhotic rat model of HCC. Rats received vehicle (control),

sorafenib at 10 mg/kg or GNS561 at 15 mg/kg. For comparison with the control non-treated

group, the non-parametric Kruskal-Wallis test was performed. **, P < 0.01.

Discussion

HCC and mCRC are refractory to most conventional chemotherapies. One of the reasons for the high mortality rate is the existence of a small population of CSCs with high tumorigenicity and resistance to drugs. Accordingly, many cancer therapies, while killing the bulk of tumor cells, may ultimately fail because they do not eliminate CSCs that reform heterogeneous tumors.

GNS561 is a small molecule that showed potent antitumor activity *in vitro* in a panel of human cancer cells, including HCC patient-derived cells, and *in vivo* in two HCC different models [14]. The *in vitro* studies performed in this study clearly showed that GNS561 displays potent dose response activity when assayed against a panel of human colon tumor cells isolated from liver metastases. In addition, GNS561 was efficient against subpopulation displaying ALDH[bright] activity and produced a dramatic decrease in the ability of these cells to form tumorspheres. Similarly, in two HCC cell lines, GNS561 was equally effective in both populations (bulk and CSCs) and dramatically decreased the cell capacity to form spheres, even at low doses, unlike sorafenib, a standard of care. This latter observation was in agreement with the literature. Indeed, CSC resistance to sorafenib has been widely reported and has been cited as responsible for treatment failure and relapse [38,39].

Taken together, our results strongly suggest that GNS561 was efficient against both the whole tumor cell population and cell subpopulations displaying CSC features (ALDH[bright] cells and tumorsphere initiating cells). These original results were confirmed *in vivo*, in a HCC-induced cirrhotic rat model, in which GNS561 showed antitumor effects and reduced the frequency of CSCs in liver tumor tissue and circulating CSCs. To go further, it would be informative to xenograft tumorsphere-derived CSCs or serial transplant tumors to determine GNS561 activity on the tumor growth.

The effectiveness of GNS561 to kill both tumor bulk cells and CSCs can be explained by its capacity to inhibit autophagy and lysosomal functions and to disrupt lysosomes [14]. In fact,

once established, tumors require an uninterrupted nutritional supply for maintaining their

proliferative needs in stressful conditions such as hypoxia, nutrient and growth factor

starvation, and oxidative and metabolic stress [17]. In this context, autophagy has a

protumoral activity by providing recycled bioenergetic substrates and consequently

overcoming nutritional deficiency and favoring the survival of cancer cells [40]. Concerning

the CSC population, autophagy was shown to be a major factor for the preservation of cell

homeostasis and the maintenance of stemness properties [5]. Consistently, it was reported

that metastatic cells and CSCs are vulnerable to lysosomal inhibition and disruption [41–44].

For example, inhibition of autophagic flux and lysosomal functions by salinomycin interferes

with the maintenance and expansion of breast cancer stem-like/progenitor cells [45].

Moreover, a salinomycin derivative has been showed to be able to sequester iron in

lysosomes ultimately resulting in cell death of CSCs [44,46]. Mefloquine showed activity on

acute myeloid leukemia cells and progenitors by disrupting lysosomes (lysosomal membrane

permeabilization and cytosolic release of cathepsins) [47].

Further research will be necessary to better understand the underlying mechanisms

implicated in GNS561 activity against CSCs. It would be interesting to evaluate whether

GNS561 treatment results from GNS561-induced inhibition of PPT1 and the mTOR pathway

and lysosomal zinc sequestration in CSCs, as demonstrated in whole tumors [14]. In fact,

knockout of PPT1 in tumor cells inhibited tumor growth, tumorsphere formation and

tumorigenesis *in vivo* [48], suggesting that PPT1 may be implicated in CSC maintenance. In

turn, the mTOR pathway has been shown to be one of the most important pathways involved

in the development and progression of cancer and in the maintenance and hallmarks of

CSCs [49], implying that inhibition of the mTOR pathway could be a good therapeutic

strategy to eradicate CSCs. Moreover, as autophagy is an adaptive mechanism modulating

the TME surrounding CSCs to support their stemness and cancer propagation [19], studying

the effect of GNS561 on the TME would be of broad interest. Finally, it was recently reported

that interfering with iron homeostasis using small molecules alters epigenetic plasticity of

cells [50]. To evaluate if zinc could also be a regulator of epigenetic plasticity and if GNS561,

by sequestrating lysosomal zinc, impairs this process would also be of particular interest.

In conclusion, GNS561 showed activity against not only the whole tumor population but also

against subpopulations displaying CSC features *in vitro* and *in vivo*. Thus, a strategy that

simultaneously exterminates the CSC subpopulation and tumor bulk by using autophagy

inhibitors and lysosomal disruptors, such as GN561, is promising as a new direction to

achieve cure and to prevent relapse of liver primary cancer and metastasis. GNS561 could

also be used in combination with other drugs, in order to enhance the overall anticancer

effect, as already described for hydroxychloroquine in several tumor types (reviewed in [3]).

In particular, as classical chemo- or radiotherapy mainly fails to eliminate the CSCSs present

in many tumor types, GNS561 could be useful to target these CSCs and then to reduce the

risk of relapse in patients.

Abbreviations

ALDH: Aldehyde dehydrogenase; CSCs: cancer stem cells; FBS: fetal bovine serum; HCC:

hepatocellular carcinoma; IC_{50}: half-maximal inhibitory concentration; mCRC: metastatic

colorectal cancer; PPT1: palmitoyl-protein thioesterase 1; SEM: standard error mean; TME:

tumor microenvironment.

Acknowledgement

The authors are very grateful to Dr. Michel Prudhomme and Dr. Jean-François Bourgaux

from CHU Carémeau, Nîmes, France, for the use of liver mCRC cells.

Conflicts of Interest

SB, EB, FB, SM, JC and PH are employees of Genoscience Pharma. SB, FB and PH are

shareholders of Genoscience Pharma. SB, FB, JC and PH are co-inventors of a pending

patent. The other authors declare that they have no conflicts of interest to report.

Funding

This work did not received external funding

References

1.   Bray F, Ferlay J, Soerjomataram I, Siegel RL, Torre LA, Jemal A. Global cancer statistics 2018:
GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. CA Cancer J
Clin. 2018; 68: 394–424.

2.   Bonnet D. Cancer stem cells: lessons from leukaemia. Cell Prolif. 2005; 38: 357–61.

3.   Nazio F, Bordi M, Cianfanelli V, Locatelli F, Cecconi F. Autophagy and cancer stem cells: molecular
mechanisms and therapeutic applications. Cell Death Differ. 2019; 26: 690–702.

4.   Sun JH, Luo Q, Liu LL, Song GB. Liver cancer stem cell markers: Progression and therapeutic implications.
World J Gastroenterol. 2016; 22: 3547–57.

5.   Vitale I, Manic G, Dandrea V, De Maria R. Role of autophagy in the maintenance and function of cancer
stem cells. Int J Dev Biol. 2015; 59: 95–108.

6.   Clarke MF, Dick JE, Dirks PB, Eaves CJ, Jamieson CH Jones DL, et al. Cancer stem cells--perspectives on
current status and future directions: AACR Workshop on cancer stem cells. Cancer Res. 2006; 66: 9339–
44.

7.   Quintana E, Shackleton M, Sabel MS, Fullen DR, Johnson TM, Morrison SJ. Efficient tumour formation by
single human melanoma cells. Nature. 2008; 456: 593–8.

8.   Pattabiraman DR, Weinberg RA. Tackling the cancer stem cells - what challenges do they pose? Nat Rev
Drug Discov. 2014; 13: 497–512.

9.   Zhou Y, Xia L, Wang H, Oyang L, Su M, Liu Q, et al. Cancer stem cells in progression of colorectal cancer.
Oncotarget. 2018; 9: 33403–15.

10.   Schulte L-A, López-Gil JC, Sainz B, Hermann PC. The Cancer Stem Cell in Hepatocellular Carcinoma.
Cancers. 2020; 12: 684.

11.   Chen J, Jin R, Zhao J, Liu J, Ying H, Yan H, et al. Potential molecular, cellular and microenvironmental
mechanism of sorafenib resistance in hepatocellular carcinoma. Cancer Lett. 2015; 367: 1–11.

12.   Cheng CC, Chao WT, Liao CC, Shih JH, Lai YS, Hsu YH, et al. The Roles Of Angiogenesis And Cancer
Stem Cells In Sorafenib Drug Resistance In Hepatocellular Carcinoma. Onco Targets Ther. 2019; 12:
8217–27.

13.   Das PK, Islam F, Lam AK. The Roles of Cancer Stem Cells and Therapy Resistance in Colorectal
Carcinoma. Cells. 2020; 9.

14.   Brun S, Raymond E, Bassissi F, Jikova ZM, Mezouar S, Rachid M, et al. GNS561, a clinical-stage PPT1
inhibitor, has powerful antitumor activity against hepatocellular carcinoma via modulation of lysosomal
functions. bioRxiv. 2020;

15.   Brun S, Bassissi F, Serdjebi C, Novello M, Tracz J,Autelitano F, et al. GNS561, a new lysosomotropic small
molecule, for the treatment of intrahepatic cholangiocarcinoma. Invest New Drugs. 2019; 37: 1135–45.

16.   ClinicalTrials. gov. Study of GNS561 in Patients With Liver Cancer. ClinicalTrials.gov.

17.   Ojha R, Bhattacharyya S, Singh SK. Autophagy in Cancer Stem Cells: A Potential Link Between
Chemoresistance, Recurrence, and Metastasis. Biores Open Access. 2015; 4: 97–108.

18.   Sharif T, Martell E, Dai C, Kennedy BE, Murphy P, Clements DR, et al. Autophagic homeostasis is required
for the pluripotency of cancer stem cells. Autophagy. 2017; 13: 264–84.

19.   Mandhair HK, Arambasic M, Novak U, Radpour R. Molecular modulation of autophagy: New venture to
target resistant cancer stem cells. World J Stem Cells. 2020; 12: 303–22.

20.   El Hout M, Cosialls E, Mehrpour M, Hamai A. Crosstalk between autophagy and metabolic regulation of
cancer stem cells. Mol Cancer. 2020; 19: 27.

21. Li N, Zhu Y. Targeting liver cancer stem cells for the treatment of hepatocellular carcinoma. Ther Adv
Gastroenterol. 2019; 12: 175628481882156.

22. Smith AG, Macleod KF. Autophagy, cancer stem cells and drug resistance. J Pathol. 2019; 247: 708–18.

23. Wu Y, Zhang J, Zhang X, Zhou H, Liu G, Li Q. Cancer Stem Cells: A Potential Breakthrough in HCC-
Targeted Therapy. Front Pharmacol. 2020; 11: 198.

24. Grillet F, Bayet E, Villeronce O, Zappia L, Lagerqvist EL, Lunke S, et al. Circulating tumour cells from
patients with colorectal cancer have cancer stem cell hallmarks in ex vivo culture. Gut. 2017; 66: 1802–10.

25. Huang EH, Hynes MJ, Zhang T, Ginestier C, Dontu G, Appelman H, et al. Aldehyde dehydrogenase 1 is a
marker for normal and malignant human colonic stem cells (SC) and tracks SC overpopulation during colon
tumorigenesis. Cancer Res. 2009; 69: 3382–9.

26. Vassalli G. Aldehyde Dehydrogenases: Not Just Markers, but Functional Regulators of Stem Cells. Stem
Cells Int. 2019; 2019: 3904645.

27. Tanaka K, Tomita H, Hisamatsu K, Nakashima T, Hatano Y, Sasaki Y, et al. ALDH1A1-overexpressing
cells are differentiated cells but not cancer stem or progenitor cells in human hepatocellular carcinoma.
Oncotarget. 2015; 6: 24722–32.

28. Flores-Tellez TN, Villa-Trevino S, Pina-Vazquez C. Road to stemness in hepatocellular carcinoma. World J
Gastroenterol. 2017; 23: 6750–76.

29. Chiba T, Kita K, Zheng Y-W, Yokosuka O, Saisho H, Iwama A, et al. Side population purified from
hepatocellular carcinoma cells harbors cancer stem cell-like properties. Hepatol Baltim Md. 2006; 44: 240–
51.

30. Cao L, Zhou Y, Zhai B, Liao J, Xu W, Zhang R, et al. Sphere-forming cell subpopulations with cancer stem
cell properties in human hepatoma cell lines. BMC Gastroenterol. 2011; 11: 71.

31. Uchida Y, Tanaka S, Aihara A, Adikrisna R, Yoshitake K, Matsumura S, et al. Analogy between sphere
forming ability and stemness of human hepatoma cells. Oncol Rep. 2010; 24: 1147–51.

32. Johnson S, Chen H, Lo PK. In vitro Tumorsphere Formation Assays. Bio Protoc. 2013; 3.

33. Ma XL, Sun YF, Wang BL, Shen MN, Zhou Y, Chen JW, et al. Sphere-forming culture enriches liver cancer
stem cells and reveals Stearoyl-CoA desaturase 1 as a potential therapeutic target. BMC Cancer. 2019; 19:
760.

34. Lee C-H, Yu C-C, Wang B-Y, Chang W-W. Tumorsphere as an effective in vitro platform for screening anti-
cancer stem cell drugs. Oncotarget. 2016; 7: 1215–26.

35. Motawi TK, El-Boghdady NA, El-Sayed AM, Helmy HS. Comparative study of the effects of PEGylated
interferon-alpha2a versus 5-fluorouracil on cancer stem cells in a rat model of hepatocellular carcinoma.
Tumour Biol. 2016; 37: 1617–25.

36. Yang ZF, Ngai P, Ho DW, Yu WC, Ng MN, Lau CK, et al. Identification of local and circulating cancer stem
cells in human liver cancer. Hepatology. 2008; 47: 919–28.

37. Yang ZF, Ho DW, Ng MN, Lau CK, Yu WC, Ngai P, et al. Significance of CD90+ cancer stem cells in
human liver cancer. Cancer Cell. 2008; 13: 153–66.

38. Cao J, Zhao M, Liu J, Zhang X, Pei Y, Wang J, et al. RACK1 Promotes Self-Renewal and Chemoresistance
of Cancer Stem Cells in Human Hepatocellular Carcinoma through Stabilizing Nanog. Theranostics. 2019;
9: 811–28.

39. Feng Y, Jiang W, Zhao W, Lu Z, Gu Y, Dong Y. miR-124 regulates liver cancer stem cells expansion and
sorafenib resistance. Exp Cell Res. 2020; 394: 112162.

40. Lim J, Murthy A. Targeting Autophagy to Treat Cancer: Challenges and Opportunities. Front Pharmacol.
2020; 11: 590344.

41.   Cash TP, Alcala S, Rico-Ferreira MDR, Hernandez-Encinas E, Garcia J, Albarran MI, et al. Induction of
Lysosome Membrane Permeabilization as a Therapeutic Strategy to Target Pancreatic Cancer Stem Cells.
Cancers Basel. 2020; 12.

42.   Morgan MJ, Fitzwalter BE, Owens CR, Powers RK, Sottnik JL, Gamez G, et al. Metastatic cells are
preferentially vulnerable to lysosomal inhibition. Proc Natl Acad Sci U A. 2018; 115: E8479–88.

43.   Salerno M, Avnet S, Bonuccelli G, Hosogi S, Granchi D, Baldini N. Impairment of lysosomal activity as a
therapeutic modality targeting cancer stem cells of embryonal rhabdomyosarcoma cell line RD. PLoS One.
2014; 9: e110340.

44.   Versini A, Colombeau L, Hienzsch A, Gaillet C, Retailleau P, Debieu S, et al. Salinomycin Derivatives Kill
Breast Cancer Stem Cells by Lysosomal Iron Targeting. Chemistry. 2020; 26: 7416–24.

45.   Mai TT, Hamaï A, Hienzsch A, Cañeque T, Müller S, Wicinski J, et al. Salinomycin kills cancer stem cells by
sequestering iron in lysosomes. Nat Chem. 2017; 9: 1025–33.

46.   Yue W, Hamai A, Tonelli G, Bauvy C, Nicolas V, Tharinger H, et al. Inhibition of the autophagic flux by
salinomycin in breast cancer stem-like/progenitor cells interferes with their maintenance. Autophagy. 2013;
9: 714–29.

47.   Sukhai MA, Prabha S, Hurren R, Rutledge AC, Lee AY, Sriskanthadevan S, et al. Lysosomal disruption
preferentially targets acute myeloid leukemia cells and progenitors. J Clin Invest. 2013; 123: 315–28.

48.   Rebecca VW, Nicastri MC, Fennelly C, Chude CI, Barber-Rotenberg JS, Ronghe A, et al. PPT1 Promotes
Tumor Growth and Is the Molecular Target of Chloroquine Derivatives in Cancer. Cancer Discov. 2019; 9:
220–9.

49.   Xia P, Xu XY. PI3K/Akt/mTOR signaling pathway in cancer stem cells: from basic research to clinical
application. Am J Cancer Res. 2015; 5: 1602–9.

50.   Müller S, Sindikubwabo F, Cañeque T, Lafon A, Versini A, Lombard B, et al. CD44 regulates epigenetic
plasticity by mediating iron endocytosis. Nat Chem. 2020; 12: 929–38.