Coordinated Changes in Gene Expression Kinetics Underlie both Mouse and Human Erythroid Maturation

Melania Barile[1, 2], Ivan Imaz-Rosshandler[1, 2], Isabella Inzani[3], Shila Ghazanfar[4], Jennifer Nichols[2, 5], John C. Marioni[4, 6, 7], Carolina Guibentif[1, 2, 8,*], Berthold Göttgens[1, 2,*]

1. Department of Haematology, University of Cambridge, CB2 0AW Cambridge, UK

2. Wellcome-Medical Research Council Cambridge Stem Cell Institute, University of Cambridge, CB2 0AW Cambridge, UK

3. University of Cambridge Metabolic Research Laboratories and MRC Metabolic Diseases Unit, CB2 0QQ Cambridge, UK

4. Cancer Research UK Cambridge Institute, University of Cambridge, CB2 0RE Cambridge, UK

5. Department of Physiology, Development and Neuroscience, University of Cambridge, CB2 3DY Cambridge, UK

6. Wellcome Sanger Institute, Wellcome Genome Campus, CB10 1SA Cambridge, UK

7. European Molecular Biology Laboratory, European Bioinformatics Institute (EMBL-EBI), Wellcome Genome Campus, CB10 1SD Cambridge, UK

8. Sahlgrenska Center for Cancer Research, Department of Microbiology and Immunology, University of Gothenburg, 413 90 Gothenburg, Sweden

* Co-corresponding authors

BG: bg200@cam.ac.uk

CG: carolina.guibentif@gu.se

Abstract

Background: Single cell technologies are transforming biomedical research, including the recent

demonstration that unspliced pre-mRNA present in single cell RNA-Seq permits prediction of

future expression states. Here we applied this 'RNA velocity concept' to an extended timecourse

dataset covering mouse gastrulation and early organogenesis. **Results:** Intriguingly, RNA velocity

correctly identified epiblast cells as the starting point, but several trajectory predictions at later

stages were inconsistent with both real time ordering and existing knowledge. The most striking

discrepancy concerned red blood cell maturation, with velocity-inferred trajectories opposing the

true differentiation path. Investigating the underlying causes revealed a group of genes with a

coordinated step-change in transcription, thus violating the assumptions behind current velocity

analysis suites, which do not accommodate time-dependent changes in expression dynamics.

Using scRNA-Seq analysis of chimeric mouse embryos lacking the major erythroid regulator

Gata1, we show that genes with the step-changes in expression dynamics during erythroid

differentiation fail to be up-regulated in the mutant cells, thus underscoring the coordination of

modulating transcription rate along a differentiation trajectory. In addition to the expected block

in erythroid maturation, the *Gata1*⁻ chimera dataset revealed induction of PU.1 and expansion of

megakaryocyte progenitors. Finally, we show that erythropoiesis in human fetal liver is similarly

characterized by a coordinated step-change in gene expression. **Conclusions:** By identifying a

limitation of the current velocity framework coupled with *in vivo* analysis of mutant cells, we

reveal a coordinated step-change in gene expression kinetics during erythropoiesis, with likely

implications for many other differentiation processes.

(247 words)

Keywords

Background

Cellular differentiation into diverse cell types underpins all metazoan development. Moreover, cellular differentiation processes are also crucial for stem cell-mediated tissue maintenance, and their perturbation has been implicated in ageing-associated regenerative failure as well as malignant transformation (1, 2). Since cellular differentiation decisions are made at the level of individual cells, elucidation of the underlying molecular mechanisms requires the use of single cell approaches. It is no surprise therefore that recent innovations in single cell molecular profiling technologies have been embraced rapidly by developmental and stem cell biologists, with complete single cell gene expression maps now available for developing embryos of several model organisms (3-5, reviewed in 6), as well as large-scale datasets covering adult tissue homeostasis (7-9).

Comprehensive molecular profiling necessarily entails the generation of snapshot data, because cells need to be fixed to examine their molecular content. This in turn represents a major drawback for the study of differentiation processes, which commonly occur over extended timeframes via complex trajectories underpinned by intricate decision-making processes. Much excitement was therefore generated by a recent seminal study (10), which demonstrated that

unspliced pre-mRNA present in scRNA-Seq datasets can be exploited to predict likely future

expression states. This so-called RNA velocity concept is based on the notion that the ratio

between unspliced and spliced RNA differs depending on whether a gene is in the process of

being up- or downregulated. During upregulation, there is a relative increase in newly transcribed

unspliced RNA, with the converse occurring during downregulation. The RNA velocity framework

has rapidly gained traction across the wider single cell community, being applied across multiple

experimental systems (11-13), and also extended as part of the scVelo analysis suite (14), which

allows inclusion of genes whose transcript levels are not in steady state.

One system where the RNA velocity concept has particular potential is erythropoiesis, the

process whereby oxygen-transporting red blood cells are generated from multipotent

haematopoietic progenitors. Research into the transcriptional control processes of

erythropoiesis led to several paradigmatic discoveries, including the dissection of distal

transcriptional control elements (15-17), as well as antagonistic transcription factor pairings as

executors of lineage choice in multipotent progenitors (18). During embryogenesis, a first so-

called primitive wave of erythropoiesis occurs in the yolk sac, followed by a second definitive

wave, initiated also in the yolk sac, then predominantly in the fetal liver and later in the adult

bone marrow (19). The zinc finger protein Gata1 represents the archetypal erythroid

transcription factor, and is required for the maturation of both primitive and definitive erythroid

cells (20-23), as well as megakaryocyte maturation (24). However, the precise molecular

processes affected by Gata1 deletion in early embryonic erythropoiesis have remained obscure,

principally because conventional biochemical methods are unsuitable for the very small number

of cells present at these early developmental stages.

Here, we have applied RNA velocity to a recently published scRNA-Seq dataset of nine sequential

timepoints, spaced 6 hours apart, which encompass mouse gastrulation and early organogenesis

(25). We observed that some of the inferred trajectories are incompatible with the existing

biological knowledge, as well as with the real time ordering derived from the sequential sampling

timepoints. For erythroid differentiation in particular, we show that failure of the velocity

framework is due to a concerted increase in transcription rate of a subset of erythroid genes,

midway through the red blood cell maturation trajectory. Analysis of *Gata1*- chimeric embryos

underscores the concerted nature of this expression boost, consistent with the notion that such

concerted upregulation events may be a feature of stabilizing a given differentiated cellular state.

Results

Limitations of RNA velocity trajectory inference at organismal scale

To evaluate RNA velocity-based trajectory inference with a complex dataset, we applied the

scVelo analysis pipeline (14) to a recently reported timecourse scRNA-Seq dataset covering

mouse gastrulation and early organogenesis. This mouse gastrulation atlas contains

approximately 120,000 single cell transcriptomes across nine sequential timepoints covering 37

major cell types (25). Prior to scVelo analysis, we removed extraembryonic ectoderm and

extraembryonic endoderm cells, as they derive from early lineage branching events that are not

covered in this dataset. We first applied scVelo to the normalised and batch corrected count

matrix across all embryonic stages (Figure 1A). We observed that scVelo correctly identifies the

epiblast population as the origin of the global differentiation processes that occur during

gastrulation and early organogenesis. In relation to the more differentiated cell types however,

there were several instances where scVelo had difficulty in capturing some of the highly complex

differentiation events that occur across the entire embryo. For instance, scVelo predicted that

E8.0 allantois and mesenchyme cell-types give rise to mesodermal cells from earlier timepoints

rather than the E8.25/E8.5 allantoic and mesenchymal cells. Another inconsistency occurred with

E8.0-E8.25 endoderm cells, which were predicted to give rise to E6.5-E7 visceral endoderm,

rather than the other way round. Most noteworthy, scVelo failed to recapitulate the

erythropoiesis branch, where it predicts a backwards differentiation from later to earlier

populations. We next repeated this analysis using data from each individual time-point (Figure

1B; shown are E7.5 and E8.5). We saw that the pipeline accurately recapitulates known biological

trajectories up to E7.5, but observed the same inconsistency from E7.75 to E8.5, with scVelo

arrows pointing backwards.

Taken together therefore, we have identified that for erythroid development, the output of

scVelo is inconsistent with the timecourse information gathered from the experimental design of

the gastrulation atlas.

Unspliced sequence reads help to discriminate between cell types

We next asked whether this issue is due to a general lack of biologically meaningful information

captured in the unspliced reads.

To this end, we exploited two variance-based dimensionality reduction methods, Principal

Component Analysis (PCA) and Multi-Omics Factor Analysis (MOFA; 26), to interrogate how much

inter-population variability is explained by the spliced and unspliced information layers, whether

considered separately or together. Upon comparing PC1 and PC2 (or MOFA Factors 1 and 2), in

addition to the expected lineage separation obtained using the spliced reads (Figure 2A, left

panel), we could also observe a degree of lineage separation when using the unspliced reads

alone (Figure 2A, middle panel). In addition, we saw a qualitatively improved separation of the

different lineages when spliced and unspliced information is used in combination (Figure 2A, right

panel; see Supplementary Figure 1 for further components/factors). Moreover, the MOFA factors

account for 16% of variation in the spliced data and 4% of the of variation in unspliced data

(Figure 2Bi). Interestingly, a closer look at the MOFA pre-processing and final outcome showed a

minor overlap of genes that are highly variable with respect to spliced or unspliced counts (Figure

2Bii) and a different weight contributed by the two layers to the final factors (Figure 2Biii).

Multiomics factor analysis therefore not only demonstrates that the unspliced reads in the

gastrulation atlas dataset contain biologically relevant information, but also suggests that

integrated analysis of spliced and unspliced reads may more broadly facilitate the interpretation

of complex scRNA-Seq datasets.

Analysis of unspliced reads reveals complex expression kinetics

Having confirmed the utility of unspliced reads, we next explored whether the inability to recover

real-time progression in whole embryo trajectory inference using scVelo might be related to the

assumptions made by the current RNA velocity analysis tools. The derivation of gene-specific

expression kinetics underpins the scVelo analysis pipeline, as illustrated by so-called phase plots

that depict the amounts of spliced versus unspliced reads within a population of cells (14). If a

gene is upregulated during a differentiation timecourse, cells will be placed above the diagonal

between no expression and maximum expression due to the relatively larger amount of newly

produced pre-mRNA during the gene induction process, while the converse is true for

downregulated genes (Figure 3A). Both of these scenarios are readily captured by scVelo, with

the predicted vectors of differentiation agreeing with the actual temporal progression. If a given

gene however experiences an increase in transcription rate midway through a differentiation

timecourse, the sudden increase in unspliced pre-mRNA will result in a phase plot that may be

wrongly classified by scVelo, with predicted vectors of differentiation diametrically opposed to

the true direction of differentiation (Figure 3A). This is indeed what we observed when inspecting

the phase plots of the scVelo driver genes (top-likelyhood genes, Supplementary Table 1), which

display a steep increase of unspliced counts in the Erythroid 3 population, leading to a reverse

velocity prediction, progressing from Erythroid 3 to earlier populations (Supplementary Figure

2A).

We next set out to identify all genes exhibiting this rapid increase in expression levels in the

Erythroid 3 population (Figure 3B). After fitting a linear regression through each population and

each gene and testing whether the inferred slopes reflected the expected order based on

biological knowledge, we found 89 such genes, which we termed Multiple Rate Kinetics or MURK

genes. These genes included *Smim1*, coding for the Vel Blood Group Antigen (27), and *Hba-x*,

where we could confirm an increase in expression kinetics using phase plots (Figure 3C).

Having identified a set of genes with a coordinated increase in expression rate midway through

erythropoiesis, we next asked what function these genes might play in the broader

transcriptional program of red blood cell maturation. Visual inspection of the gene list revealed

it to contain archetypal red blood cell genes including the globin genes *Hba-x*, *Hbb-a1*, *Hba-a2*,

Hbb-bt, *Hbb-bh1*, *Hbb-y* (Supplementary table 2). Unsupervised gene ontology analysis

confirmed that biological functions essential for red blood cells were highly enriched, including

"gas transport" and "heme biosynthetic process" (Figure 3D).

We next removed this set of MURK genes and recalculated the RNA velocity inferred trajectories.

As can be seen in Figure 3E, inferred vectors of differentiation are now in good agreement with

the real time progression of erythropoiesis

The scVelo suite also calculates a so-called latent time, which represents the pseudotime

ordering hidden in the spliced and unspliced dynamics, and is more powerful than previously

described pseudotime inferring approaches since it incorporates both the gene dynamics and the

spliced and unspliced information (14). Using the full gene set, the latent time calculation for the

erythroid lineage is contrary to the know progression of erythroid differentiation (Figure 3E left

panels, Supplementary Figure 2B, left panels). By contrast, removing the MURK genes results in

a latent time prediction that is not only consistent with the major axis of erythropoiesis, but also

identifies the two sequential inputs described previously (25), namely an early wave directly from

posterior mesoderm as well as a second wave coming from yolk sac hemogenic endothelium (see

Figure 3E, Supplementary Figure 2B, right panels).

Taken together therefore, this analysis shows that inconsistent RNA velocity-inferred trajectories

can be remedied by the removal of genes with complex expression kinetics.

Erythroid Multiple Rate Kinetics genes are essential for red blood cell function

To corroborate upregulation of our identified MURK genes during erythropoiesis, we

interrogated a previously published dataset with transcriptomic analysis of a loss of function

model for the erythropoiesis master-regulator *Gata1* (28). *In vitro* differentiation of Gata1 knock-

out embryonic stem cells over-expressing human *BCL2* can produce permanently self-renewing

immature erythroid progenitor cell lines. One such model, G1ER, contains a tamoxifen-inducible

Gata1 transgene, the activation of which triggers erythroid maturation (29, 30; Figure 4A).

Microarray-based differential gene expression was performed, comparing the uninduced and

induced conditions (28). 76 of our 89 MURK genes overlapped with the genes identified by this

microarray-based comparison. Of those, 64 were upregulated, of which 55 showed strong

upregulation, 4 were downregulated, and 8 showed no change in expression following induction

of Gata1 in the G1ER system, demonstrating a highly significant overlap of our identified MURK

genes with the G1ER-induced genes ($p < 10^{-24}$; see Figure 4B).

Our newly identified erythropoietic MURK genes therefore perform key roles in red blood cell

function, and their upregulation was validated in an independent model of red blood cell

maturation.

scRNA-Seq of mouse chimeras reveals the early cellular defects in Gata1 loss of function

The G1ER cell line represents an *in vitro* model, and the published differential gene expression data were from bulk microarray profiling, thus precluding any analysis of single-cell gene expression kinetics. We therefore turned to our recently reported Chimaera-Seq approach, whereby scRNA-Seq is coupled with mouse chimeric embryo technology, to define both cellular and molecular consequences of gene knock-outs *in vivo* (25, 31). We used our standard embryonic stem cells (ESCs) expressing a constitutive tdTomato (tdTom) fluorescent marker gene to generate a Gata1 knock-out line (see Methods). *Gata1*⁻ tdTom⁺ cells were injected into tdTom⁻ wild-type blastocyst and transferred into pseudo-pregnant females, resulting in chimeric embryos that we harvested at E8.5. Six chimeric embryos were pooled, dissociated into a single-cell suspension, and tdTom⁺ and tdTom⁻ cell fractions were sorted for scRNA sequencing. We obtained 8420 tdTom⁻ and 7944 tdTom⁺ cells passing quality control and assigned to a cell type, with an average of 4354 genes being detected per cell.

We then concatenated the chimera data with the Pijuan-Sala et al. (2019) reference dataset and mapped nearest neighbors (see Methods). We observed an overall homogeneous distribution of both mutant and wild-type fractions throughout the later time-points of the landscape, except for the erythroid branch. Indeed, we observed a block in the erythroid lineage of the mutant cells, which were over-represented in the start of the erythroid differentiation branch, while their wild-type counterparts were present throughout erythroid differentiation (Supplementary Figure 3). Identification of the nearest neighbours of chimeric cells within the reference dataset allowed their quick cell-type annotation, which we used to quantify the differences in the hemato-endothelial cell-type representation within the chimera fractions. This analysis confirmed a

severe erythroid differentiation defect of the mutant cells (Figure 4C-E). When examining the

reference dataset sampled-time point of the chimera nearest neighbours we also observed a

temporal shift within the erythroid lineage, with tdTom[+] mutant cells mapping to earlier time-

points than their wildtype tdTom[-] counterparts, further confirming a developmental block of the

mutant cells (Figure 4D, E). In addition, we observed that this erythroid defect was coupled with

an over-representation of cells with a megakaryocyte signature (Figure 4C).

The newly generated *Gata1*[-] Chimaera-Seq data therefore not only recapitulated the expected

block in erythroid maturation, but also revealed an expansion of the megakaryocytic lineage in

the E8.5 yolk sac.

The molecular program affected by Gata1 loss in early embryos

Although the role of Gata1 is well documented in developmental erythropoiesis (21, 23), the early

molecular defects of Gata1 loss of function *in vivo* had not been reported. The Gata1 Chimaera-

Seq dataset therefore presented an opportunity to dissect the early molecular program

controlled by Gata1 *in vivo*. Having registered a defect in erythroid differentiation and an increase

in the megakaryocytic lineage population, we performed differential gene expression testing

between the chimera mutant and wild-type cells in these clusters (Supplementary Table 3).

Regarding the megakaryocytic subset, we observed upregulation of progenitor markers *Kit*,

Gata2 and *Myb* in the *Gata1*[-] cells as well as lower expression of maturation genes for the

megakaryocyte lineage *Gp5*, *Pf4*, *Mpl* and *Plek* (Figure 5A). Hyper-proliferative megakaryocyte

progenitors, detected previously in *Gata1*[-] E12.5 fetal livers, led to compromised platelet

function, and were suggested to originate in the yolk sac (32). Our results showing over-

production of megakaryocytic cells with impaired maturation characteristics in E8.5 *Gata1*⁻

chimera yolk sacs support this notion, and importantly place the megakaryocytic defect within

the very early phase of megakaryocyte formation.

Interestingly, all hemato-endothelial cell subsets displayed up-regulation of *Spi1* (coding for the

PU.1 transcription factor) in the *Gata1*⁻ cell fraction compared to wild-type counterpart (FDR <

0.01; Figure 5A). Given the previously reported Gata1-PU.1 cross-repression in adult bone

marrow (18) and in zebrafish embryonic hematopoiesis (33), we systematically assessed the

effect of *Gata1* knockout in the mouse chimera lineages and observed that in *Gata1*⁻ cells, *Spi1*

was specifically up-regulated in all hematopoietic sub-clusters, with a stronger effect on Mk and

Ery1 subsets. (Supplementary Figure 3).

In the early erythroid subset, Ery1, we again noted that the mutant cells displayed increased

expression of genes characteristic of a progenitor signature. Conversely, erythroid maturation

hallmark genes such as *Hbb-bs* and *Gypa* were downregulated, along with the erythroid Gata1

target *Mllt3* (34; Figure 5A). GO-term enrichment analysis of genes downregulated in *Gata1*⁻ Ery1

cells revealed biological processes essential to red blood cell function (Figure 5B). Furthermore,

we also observed that 48% of the MURK genes identified in Figure 3 overlapped with these genes

that fail to up-regulate in *Gata1*⁻ erythroid cells (Figure 5C; $p < 10^{-24}$).

In addition to the failure of inducing genes associated with erythroid maturation, single cell

resolution molecular analysis also revealed a striking failure to downregulate genes associated

with alternative lineage programs such as Pu.1, consistent with the notion that the earliest wave

of primitive hematopoiesis produces erythroid cells, megakaryocytes and macrophages, with

evidence for at least bipotential progenitor cells (35).

The late erythroid increase in expression rate is downstream of Gata1 function

Having generated the Chimaera-Seq single cell data for both wildtype and Gata1 knock-out cells,

we next used the ratio of spliced/unspliced reads to explore differences in expression kinetics

between the wildtype and mutant cells. As can be seen in Figure 5D, the previously defined MURK

genes failed to display the increased rate of expression characteristic for the later stages of

erythropoiesis in the mutant cells. The examples shown include the embryonic globin gene *Hbb-*

y, as well as the *Fam210b* gene, coding for a putative mitochondrial protein recently implicated

in erythroid differentiation (36; Figure 5D). This result confirms that the erythroid boost in

expression forms part of the transcriptional program downstream of Gata1 function, although it

does not demonstrate a direct regulatory role for Gata1.

However, preliminary modelling analysis suggests that the change observed in MURK gene

dynamics is due to altered transcription rates (see Supplementary Note), indicating a close

association of the coordinated late erythroid increase in transcription rate with the molecular

program downstream of Gata1.

A coordinated increase of expression rate during human fetal liver erythropoiesis

Having identified a coordinated increase in transcription rate during mouse yolk sac

erythropoiesis, we next wanted to ascertain whether the same phenomenon could also be seen

in human cells. Moreover, we were keen to explore an scRNA-Seq dataset generated by a

different laboratory, to exclude any potential technical bias caused by our own experimental

protocols. We therefore turned to a recently published comprehensive dataset of human fetal

liver erythropoiesis (37), and extracted the 49,388 cells annotated to the four clusters

encompassing human fetal liver erythropoiesis. When calculating scVelo-based differentiation

vectors as well as latent time using the full gene set (see methods), both were reversed (Figure

6A, left plots), consistent with the mouse yolk sac results. We therefore again ran our pipeline to

discover genes with a potential increase in expression rate along the differentiation pathway.

The resulting 97 genes again contained archetypal erythroid genes such as the haemoglobin

genes (Figure 6B), with overall gene ontologies demonstrating a functional role in erythropoiesis

(Figure 6C, see also Supplementary Table 4). We then recalculated both the scVelo differentiation

vectors as well as latent time after removing the fetal liver MURK genes. This revealed scVelo

vectors that were consistent with the expected developmental progression (see Figure 6A, right

plots). This analysis therefore demonstrates that complex expression kinetics apply broadly to

erythropoiesis, and their identification can be used to amend the RNA velocity framework to

prevent erroneous predictions.

Discussion

There is no doubt that single cell molecular profiling constitutes a transformative technology. It

suffers however from the major drawback that cells need to be fixed in order to profile them,

with the consequence that measurements are by necessity static snapshots. To decipher complex

biological processes, however, temporal information is commonly required. The single cell RNA

velocity concept raised the prospect of overcoming some of the limitations associated with static

measurements, by providing a strategy that can infer future cellular states. The RNA velocity

framework is based on an explicit model of transcriptional processes (transcription, splicing,

degradation). The notion that physical parameters of gene expression can be deduced from single

cell gene expression data had been explored before the single cell RNA velocity concept was

introduced (38, 39). However, the scVelo implementation provided an attractive framework for

estimating gene-specific expression parameters by taking advantage of the spliced versus

unspliced read counts across large cell populations (14). Using erythropoiesis as an example, we

show here that this current framework needs to be adapted to accommodate more complex

expression kinetics. Importantly, our analysis revealed that sets of genes can show a coordinated

increase in transcription rate along a differentiation pathway. Moreover, deletion of the key

erythroid regulator Gata1 abrogated this coordinated change in expression dynamics, thus

revealing this increase in transcription rate as an important feature of erythropoiesis. Of note,

current RNA velocity frameworks consider only a single reason for the presence of introns,

namely that a pre mRNA has not been fully processed. However, it is known that other processes

such as intron retention can result in the presence of intronic sequences in otherwise fully

processed cytoplasmic mRNA molecules (40, 41), thus suggesting that a more granular approach

towards both the modelling and experimental analysis of spliced versus unspliced reads

represents a promising avenue for future research.

Application of the single cell RNA velocity concept has commonly been "confirmatory", whereby

a differentiation path proposed by other means was shown to be consistent with RNA velocity

inference. When we applied the RNA velocity framework to the entire mouse gastrulation atlas,

some inferred vectors of differentiation agreed with our current understanding of developmental

biology, but others disagreed. Deeper interrogation of predictions that conflicted with our

current understanding of erythropoiesis showed that the RNA velocity predictions could not be

correct, not only because they ran counter to the known expression changes that accompany red

blood cell differentiation, but also because they contradicted the real-time sampling of the data.

Our results thus highlight certain limitations of the current implementation of this framework for

identification of novel trajectories. Importantly however, it is through our observation of the

inconsistent predictions that we were led to identify the previously unrecognized dynamic nature

of the transcriptional control of erythropoiesis. Moreover, it is plausible that coordinated

increases in transcription rate midway through a differentiation process may operate more

widely, as a powerful mechanism for stabilising a cell state. Our extension to the scVelo

implementation reveals the presence of such time-dependent changes of gene expression

parameters and retrieves the concerned MURK genes in developmental trajectories of interest.

As to the precise mechanisms, at this stage we can only confidently assert that this process occurs

downstream of Gata1 during erythropoiesis. Of note, comprehensive analysis of the G1ER

erythroid differentiation model has shown that Gata1-induced maturation triggers increased

enhancer/promoter interactions for upregulated genes, and that the most highly enriched motif

in the promoters of these genes are GATA sites (42). These observations are therefore consistent

with the lineage-determining function of Gata1 involving a coordinated increase in expression

kinetics of a set of genes important for red blood cell function.

Our observations regarding the Gata1 knock-out phenotype also warrant some discussion. With

embryonically lethal phenotypes such as Gata1 knock-out, conventional analysis tends to be

somewhat limited, since the embryos are dead because they have no red blood cells. By contrast,

the Chimaera-Seq assay enables both quantification of cell numbers as well as characterisation

of their molecular profiles. Moreover, there are no secondary effects caused by the dying

embryo, because the wildtype host cells rescue overall fetal development, thus allowing a

focussed analysis of cell-intrinsic molecular defects. One noteworthy observation from our data

is that erythroid differentiation proceeds substantially beyond the stage where *Gata1* expression

itself is first initiated, but fails to proceed to the late erythroid phase where expression of

canonical red blood cell genes is greatly upregulated. However, gene expression prior to the

differentiation block is not normal. In particular, we observed increased Spi1/Pu.1 in the Gata1

knock-out cells, consistent with the previously reported (18) but also disputed (43) antagonistic

relationship between Gata1 and Pu.1.

Within haematopoiesis, Pu.1 is recognised as a key regulator of myeloid and T-cell lineages, but

not erythroid cells, even though a role in the proliferation of immature erythroid progenitors has

been reported (44, reviewed in 45). Upregulation of Pu.1 in our immature *Gata1* knock-out cells

therefore suggests that these cells of the primitive haematopoietic lineage represent progenitors

with multilineage potential, rather than being restricted to just the red cell lineage. Further

evidence for this notion is provided by our observation that the reduction in erythroid cells in the

Gata1 knock-out is accompanied by an increase in megakaryocyte progenitors, consistent with a

model whereby Gata1 levels influence the lineage choice decisions of a multipotent progenitor

cell. Live cell tracking studies have suggested that the primary role of Gata1 and Pu.1 may be fate

stabilization rather than fate choice (43). The increase in transcription rate of erythroid genes

downstream of Gata1 would cohere with stabilizing the erythroid fate, thus suggesting that our

results are consistent with roles in both fate choice and fate stabilization.

Our observation of an expanded pool of megakaryocyte progenitors may also be of direct

relevance to our understanding of the pre-leukaemic transient myeloproliferative disease (TMD)

that is prevalent in newborns with trisomy 21 (46). TMD is thought to arise when a fetal specific

haematopoietic progenitor cell with trisomy 21 acquires a partial loss of function mutation in

GATA1, resulting in a short form of GATA1 (GATA1s). TMD is characterized by expansion of

immature megakaryocyte progenitors, and in 10 to 20% of cases transforms into malignant acute

megakaryoblastic leukaemia (reviewed in 47). Over-expression of GATA1s in mouse models

resulted in the identification of mid-gestation fetal liver megakaryocyte progenitors as uniquely

sensitive to this mutant GATA1s form compared to their adult bone marrow counterparts (48).

The over-represented population of immature megakaryocytic progenitors in our E8.5 *Gata1*⁻

chimeras may correspond to the developmental emergence of this transient precursor, TMD-

initiating cell, in the yolk sac.

Conclusions

Taken together, this study reports how the RNA velocity framework can be extended to delve

into the transcriptional mechanisms of tissue differentiation, complemented with single cell

resolution and *in vivo* analysis of Gata1 function, which revealed a number of previously unknown

facets of this canonical regulator of red blood cell development.

Methods

scVelo implementation

Mouse atlas dataset. To obtain separated count matrices for spliced and unspliced mRNAs, we

ran velocyto 0.17.17 (10) on the .bam files from the mouse atlas in Pijuan-Sala et al. 2019 (25;

GEO accession number: GSE87038). We kept all cells that passed the QC as described in the

original publication, but filtered out from downstream analysis the extraembryonic tissues: ExE

endoderm, ExE ectoderm and Parietal endoderm as well as samples with no timepoint allocation

(labelled as 'mixed gastrulation'). To select highly variable genes (HVGs) we applied both the

scanpy v1.5.1 and the scVelo v0.2.1 (14) pipelines. That is, we removed genes with less than 20

shared counts between spliced and unspliced counts, before normalising and log transforming

the remaining genes. Then, we selected the top 2500 HVGs from each approach (resulting in a

total of 4000, with 1000 overlapping genes) for further calculation of moments; while performing

imputation using the top 30 nearest neighbours from the graph connectivities generated with

the original UMAP coordinates from Pijuan-Sala et al. 2019. The velocity vectors were computed

in dynamical mode rather than steady state.

Human dataset. We first downloaded raw reads from Popescu et al., 2019 (37; GEO accession

number: GSE127980), and aligned them against the human genome hg19-3.0.0 with CellRanger

v3.0.2 to generate the .bam files and obtain separated count matrices for spliced and unspliced

mRNAs as described above. We filtered out cells with less than 3,550 counts, less than 900 genes

and more than 6% mitochondrial counts. Again, we combined scapy and scVelo's pipelines to

select 1,500 HVGs to compute PCA coordinates and applied batch correction using the function

reducedMNN from the batchelor package v1.4.0 (49), followed by the estimation of velocity

vectors in the same way it was done for the mouse dataset.

MOFA+ implementation

We ran MOFA+ v1.4.0 (26) using as input the two single cell experiment objects obtained from

the spliced and unspliced counts independently. Each object was created in R using the scran

v1.16.0 (50) library as follows: we started from the raw counts, normalized them with factor sizes

obtained after pre-clustering, log transformed and reduced to 5000 HVG. We then switched to

Python v3.7.4, where we regressed out the sample effect and scaled the object to generate a

MOFA+ model with standard parameters. Finally, we used reducedMNN to correct the MOFA

Factors for batch effects. The same objects used as MOFA input were used for PCA calculation in

Figure 2A.

MURK genes identification

To identify MURK genes, we considered the imputed counts resulting from the scVelo standard

pipeline. Then, for each gene and each population among the Erythroid lineage, we calculated

the unspliced versus spliced slope with a linear regression, as well as the standard error on the

slope. In the mouse dataset we selected all genes for which the slope in Erythroid3 is significantly

higher than the slope in Erythoid2 (according to a one-sided t-test p-value < 0.05), the average

spliced counts in Erythroid3 is higher than the average spliced counts in every other population,

and the slope in Erythroid3 positive. We found 89 genes that respect all these criteria.

In the human dataset, in order to obtain erythroid populations more comparable to our mouse

data, we re-clustered the erythroid clusters (Figure 6A). We retained the population annotations

from the original paper except for the Late Erythroid population, which we defined after

performing Leiden clustering on the Umap coordinates. Specifically, we re-allocated a subset of

the previously annotated Mid Erythroid population to Late Erythroid, in such a way that they

have a similar numbers of cells. We then calculated the unspliced versus spliced slope with linear

regression and identified MURK genes where the slope in Late Erythroid is significantly higher

than the slope in Mid Erythroid. We found 97 genes respecting these criteria.

Gene ontology enrichment analysis

We performed gene ontology enrichment analysis using the http://geneontology.org website

comparing the MURK genes against all biological processes, with the default all Mus musculus

genes in database as background set (51, 52). We ranked the processes by FDR.

Overlap testing

Overlap was tested with Fisher exact test. We calculated the probability of having m = 55 genes

of our n = 89 MURK genes mapping to the A = 1022 high response genes (out of N = 4195 genes)

in the Wu et al., 2011 publication (GEO accession number: GSE30142) as the probability of

randomly picking m elements of a specific type when randomly choosing n elements out of N,

where the frequency of the special type is A/N.

Gata1⁻ chimera dataset generation and analysis

Embryo collection. All procedures were performed in strict accordance to the UK Home Office

regulations for animal research under the project license number PPL 70/8406. **Chimaera**

generation. TdTomato-expressing mouse embryonic stem cells (ESC) were derived as previously

described (25). Briefly, ESC lines were derived from E3.5 blastocysts obtained by crossing a male

ROSA26tdTomato (Jax Labs – 007905) with a wildtype C57BL/6 female, expanded under the

2i+LIF conditions (53) and transiently transfected with a Cre-IRES-GFP plasmid (54) using

Lipofectamine 3000 Transfection Reagent (ThermoFisher Scientific, #L3000008) according to

manufacturer's instructions. A tdTomato-positive, male, karyotypically normal line, competent

for chimaera generation as assessed using morula aggregation assay, was selected for targeting

Gata1. Two guides were designed using the http://crispr.mit.edu tool (guide 1:

CGGCTACTCCACTGTGGCGG; guide 2: CGCTTCTTGGGCCGGATGAG) and were cloned into the

pX458 plasmid (Addgene, #48138) as previously described (55). The obtained plasmids were then

used to transfect the cells and single transfected clones were expanded and assessed for Cas9-

induced mutations. Genomic DNA was isolated by incubating cell pellets in 0.1 mg/ml of

Proteinase K (Sigma, #03115828001) in TE buffer at 50°C for 2 hours, followed by 5 min at 99°C.

The sequence flanking the guide-targeted sites was amplified from the genomic DNA by

polymerase chain reaction (PCR) in a Biometra T3000 Thermocycler (30 sec at 98°C ; 30 cycles of

10 sec at 98°C, 20 sec at 58°C, 20 sec at 72°C; and elongation for 7 min at 72°C) using the Phusion

High-Fidelity DNA Polymerase (NEB, #M0530S) according to the manufacturer's instructions.

Primers   including   Nextera   overhangs   were   used   (F-

TCGTCGGCAGCGTCAGATGTGTATAAGAGACAGTCTACCCTGCCTCAACTGTG;   R-

GTCTCGTGGGCTCGGAGATGTGTATAAGAGACAGTCTTGTCTTGGGCAGGAACA), allowing library

preparation with the Nextera XT Kit (Illumina, #15052163), and sequencing was performed using

the Illumina MiSeq system according to manufacturer's instructions. An ESC clone showing a 38

base-pair frameshift mutation in exon 4 resulting in the functional inactivation of *Gata1* were

selected for injection into C57BL/6 E3.5 blastocysts. A total of 6 chimaeric embryos were

harvested at E8.5, dissected, and single-cell suspensions were generated by TrypLE Express

dissociation reagent (Thermo Fisher Scientific) incubation for 7-10 minutes at 37°C under

agitation. Single-cell suspensions were sorted into tdTom+ and tdTom- samples using a BD Influx

sorter with DAPI at 1µg/ml (Sigma) as a viability stain for subsequent 10X scRNA-seq library

preparation (version 3 chemistry), and sequencing using an S1 flow cell in the Illumina Novaseq

platform, which resulted in 8420 tdTom$^-$ and 7944 tdTom$^+$ cells that passed quality control (see

"Single-cell RNA sequencing analysis" below).

Single-cell RNA sequencing analysis. Raw files were processed with Cell Ranger 3.0.2 using

default mapping arguments. Reads were mapped to the mm10 genome and counted with

GRCm38.92 annotation, including tdTomato sequence for chimera cells. Cell barcodes with

expression profiles significantly different to the ambient mRNA expression profile were identified

using emptyDrops (56), and cell barcodes with low complexity, i.e. low total mRNA counts and/or

high mitochondrial proportion, were identified by fitting four-component bivariate mixture

models to the $\log_{10}$-transformed total mRNA counts and percentage of mitochondrial counts, and

selecting the components with high total mRNA and low mitochondrial percentage. Gene

expression normalization and doublet cell barcodes were identified using the approach taken by

Pijuan-Sala et al. (2019). Both spliced and unspliced count matrices were extracted using velocyto

0.17.17 (10).

Mapping to the reference dataset. We mapped the chimaera cells to the mouse atlas following

almost exactly the procedure used in the original publication article to map the *Tal1* chimaera.

First, we concatenated the mouse atlas and chimaera counts (both previously controlled for

quality of the cells), normalized the resulting counts matrix with scran, computed HVGs and then

applied multiBatchPCA, and reducedMNN with cosine normalization from batchelor (49) for

batch effect correction within samples (where sample refers to a single lane of a 10x Chromium

chip) as well as between datasets in order to extract a number of nearest neighbours between

the mouse atlas and the chimaera using queryKNN from BiocNeighbors package v1.6.0.

Differential Gene Expression Analysis. For differential gene expression analysis, we took samples

that included at least 7 cells per tdTom status per cell population (eg. Erythroid3). We ran the

analysis in scanpy v1.5.1 (57) with Wilcoxon test and choosing 2 as fold change and 0.1 as false

discovery rate thresholds.

Funding

Research in the authors' laboratories is supported by the Wellcome Trust, MRC, CRUK, Blood

Cancer UK, NIH-NIDDK, the Sanger-EBI Single Cell Centre; by core support grants by the Wellcome

Trust to the Cambridge Institute for Medical Research and Wellcome Trust-MRC Cambridge Stem

Cell Institute; and by core funding from Cancer Research UK and the European Molecular Biology

Laboratory. C.G. was funded by the Swedish Research Council (2017-06278), I.I. was funded by a

British Heart Foundation studentship (FS/18/56/35177), S.G. was supported by a Royal Society

Newton International Fellowship (NIF\R1\181950). This work was funded as part of a Wellcome

Strategic Award (105031/D/14/Z) awarded to Wolf Reik, Berthold Göttgens, John Marioni,

Jennifer Nichols, Ludovic Vallier, Shankar Srinivas, Benjamin Simons, Sarah Teichmann, and

Thierry Voet.

Authors' contributions

M.B. performed scVelo implementations in mouse and human datasets, mathematical modelling,

and analysis of Gata1 embryonic chimera dataset; I.I-R. assisted on the scVelo implementation in

mouse datasets; I.I. performed Gata1 CRISPR/Cas9 targeting and expansion of the resulting

mutant lines; S.G. performed quality controls of the Gata1 embryonic chimera dataset; S.G. and

C.G. performed initial analysis of the Gata1 embryonic chimera dataset; C.G. designed and

optimized the mutant chimera single-cell profiling experiments; B.G. wrote the initial draft of the

manuscript; M.B., C.G., J.C.M. edited the manuscript; J.N., J.C.M., C.G. and B.G. supervised the

study. All authors read and approved the final manuscript.

Acknowledgements

We would like to thank Prof. Fabian Theis and Volker Bergen for discussions and valuable input

on the scVelo implementation. We thank Prof. Ross Hardison for providing the list of Gata1-

regulated genes from Wu et al. 2011 (Figure 4B). We thank William Mansfield and the Gurdon

Institute animal facility for blastocyst injections, the Flow Cytometry Core Facility at the

Cambridge Institute for Medical Research for cell sorting, Katarzyna Kania and the CRUK-CI

genomics core for preparing the 10X libraries and for sequencing.

References

1.   Akunuru S, Geiger H. Aging, Clonality, and Rejuvenation of Hematopoietic Stem Cells.

Trends Mol Med. 2016;22(8):701-12.

2.   Schultz MB, Sinclair DA. When stem cells grow old: phenotypes and mechanisms of stem

cell aging. Development. 2016;143(1):3-14.

3.   Mahadevaiah SK, Sangrithi MN, Hirota T, Turner JMA. A single-cell transcriptome atlas of

marsupial embryogenesis and X inactivation. Nature. 2020;586(7830):612-7.

4.   Gerber T, Murawala P, Knapp D, Masselink W, Schuez M, Hermann S, et al. Single-cell

analysis uncovers convergence of cell identities during axolotl limb regeneration. Science.

2018;362(6413).

5.   Wagner DE, Weinreb C, Collins ZM, Briggs JA, Megason SG, Klein AM. Single-cell

mapping of gene expression landscapes and lineage in the zebrafish embryo. Science.

2018;360(6392):981-7.

6.   Ton MN, Guibentif C, Gottgens B. Single cell genomics and developmental biology:

moving beyond the generation of cell type catalogues. Curr Opin Genet Dev. 2020;64:66-71.

7.   Borrett MJ, Innes BT, Jeong D, Tahmasian N, Storer MA, Bader GD, et al. Single-Cell

Profiling Shows Murine Forebrain Neural Stem Cells Reacquire a Developmental State when

Activated for Adult Neurogenesis. Cell Rep. 2020;32(6):108022.

8.   Weinreb C, Rodriguez-Fraticelli A, Camargo FD, Klein AM. Lineage tracing on

transcriptional landscapes links state to fate during differentiation. Science. 2020;367(6479).

9.   Dahlin JS, Hamey FK, Pijuan-Sala B, Shepherd M, Lau WWY, Nestorowa S, et al. A single-

cell hematopoietic landscape resolves 8 lineage trajectories and defects in Kit mutant mice.

Blood. 2018;131(21):e1-e11.

10.   La Manno G, Soldatov R, Zeisel A, Braun E, Hochgerner H, Petukhov V, et al. RNA velocity

of single cells. Nature. 2018;560(7719):494-8.

11.   Zhang Q, He Y, Luo N, Patel SJ, Han Y, Gao R, et al. Landscape and Dynamics of Single

Immune Cells in Hepatocellular Carcinoma. Cell. 2019;179(4):829-45 e20.

12.   Zhou W, Yui MA, Williams BA, Yun J, Wold BJ, Cai L, et al. Single-Cell Analysis Reveals

Regulatory Gene Expression Dynamics Leading to Lineage Commitment in Early T Cell

Development. Cell Syst. 2019;9(4):321-37 e9.

13.   Kanton S, Boyle MJ, He Z, Santel M, Weigert A, Sanchis-Calleja F, et al. Organoid single-

cell genomic atlas uncovers human-specific features of brain development. Nature.

2019;574(7778):418-22.

14.   Bergen V, Lange M, Peidli S, Wolf FA, Theis FJ. Generalizing RNA velocity to transient cell

states through dynamical modeling. Nat Biotechnol. 2020.

15.   Grosveld F, van Assendelft GB, Greaves DR, Kollias G. Position-independent, high-level

expression of the human beta-globin gene in transgenic mice. Cell. 1987;51(6):975-85.

16.   Higgs DR, Wood WG, Jarman AP, Sharpe J, Lida J, Pretorius IM, et al. A major positive

regulatory region located far upstream of the human alpha-globin gene locus. Genes Dev.

1990;4(9):1588-601.

17.   Mettananda S, Gibbons RJ, Higgs DR. Understanding alpha-globin gene regulation and

implications for the treatment of beta-thalassemia. Ann N Y Acad Sci. 2016;1368(1):16-24.

18.   Zhang P, Behre G, Pan J, Iwama A, Wara-Aswapati N, Radomska HS, et al. Negative

cross-talk between hematopoietic regulators: GATA proteins repress PU.1. Proc Natl Acad Sci U

S A. 1999;96(15):8705-10.

19.   McGrath K, Palis J. Ontogeny of erythropoiesis in the mammalian embryo. Curr Top Dev

Biol. 2008;82:1-22.

20.   Pevny L, Simon MC, Robertson E, Klein WH, Tsai SF, D'Agati V, et al. Erythroid

differentiation in chimaeric mice blocked by a targeted mutation in the gene for transcription

factor GATA-1. Nature. 1991;349(6306):257-60.

21.   Pevny L, Lin CS, D'Agati V, Simon MC, Orkin SH, Costantini F. Development of

hematopoietic cells lacking transcription factor GATA-1. Development. 1995;121(1):163-72.

22.   Gutierrez L, Tsukamoto S, Suzuki M, Yamamoto-Mukai H, Yamamoto M, Philipsen S, et

al. Ablation of Gata1 in adult mice results in aplastic crisis, revealing its essential role in steady-

state and stress erythropoiesis. Blood. 2008;111(8):4375-85.

23.   Fujiwara Y, Browne CP, Cunniff K, Goff SC, Orkin SH. Arrested development of embryonic

red cell precursors in mouse embryos lacking transcription factor GATA-1. Proc Natl Acad Sci U

S A. 1996;93(22):12355-8.

24.   Shivdasani RA, Fujiwara Y, McDevitt MA, Orkin SH. A lineage-selective knockout

establishes the critical role of transcription factor GATA-1 in megakaryocyte growth and platelet

development. EMBO J. 1997;16(13):3965-73.

25.   Pijuan-Sala B, Griffiths JA, Guibentif C, Hiscock TW, Jawaid W, Calero-Nieto FJ, et al. A

single-cell molecular map of mouse gastrulation and early organogenesis. Nature.

2019;566(7745):490-5.

26.   Argelaguet R, Arnol D, Bredikhin D, Deloro Y, Velten B, Marioni JC, et al. MOFA+: a

statistical framework for comprehensive integration of multi-modal single-cell data. Genome

Biol. 2020;21(1):111.

27.   Storry JR, Joud M, Christophersen MK, Thuresson B, Akerstrom B, Sojka BN, et al.

Homozygosity for a null allele of SMIM1 defines the Vel-negative blood group phenotype. Nat

Genet. 2013;45(5):537-41.

28.   Wu W, Cheng Y, Keller CA, Ernst J, Kumar SA, Mishra T, et al. Dynamics of the epigenetic

landscape during erythroid differentiation after GATA1 restoration. Genome Res.

2011;21(10):1659-71.

29.   Tsang AP, Visvader JE, Turner CA, Fujiwara Y, Yu C, Weiss MJ, et al. FOG, a multitype zinc

finger protein, acts as a cofactor for transcription factor GATA-1 in erythroid and

megakaryocytic differentiation. Cell. 1997;90(1):109-19.

30.   Weiss MJ, Yu C, Orkin SH. Erythroid-cell-specific properties of transcription factor GATA-

1 revealed by phenotypic rescue of a gene-targeted cell line. Mol Cell Biol. 1997;17(3):1642-51.

31.   Guibentif C, Griffiths JA, Imaz-Rosshandler I, Ghazanfar S, Nichols J, Wilson V, et al.

Diverse Routes toward Early Somites in the Mouse Embryo. Dev Cell. 2020.

32.   Vyas P, Ault K, Jackson CW, Orkin SH, Shivdasani RA. Consequences of GATA-1 deficiency

in megakaryocytes and platelets. Blood. 1999;93(9):2867-75.

33.   Monteiro R, Pouget C, Patient R. The gata1/pu.1 lineage fate paradigm varies between

blood populations and is modulated by tif1gamma. EMBO J. 2011;30(6):1093-103.

34.   Pina C, May G, Soneji S, Hong D, Enver T. MLLT3 regulates early human erythroid and

megakaryocytic cell fate. Cell Stem Cell. 2008;2(3):264-73.

35.   Palis J. Hematopoietic stem cell-independent hematopoiesis: emergence of erythroid,

megakaryocyte, and myeloid potential in the mammalian embryo. FEBS Lett.

2016;590(22):3965-74.

36.   Kondo A, Fujiwara T, Okitsu Y, Fukuhara N, Onishi Y, Nakamura Y, et al. Identification of

a novel putative mitochondrial protein FAM210B associated with erythroid differentiation. Int J

Hematol. 2016;103(4):387-95.

37.   Popescu DM, Botting RA, Stephenson E, Green K, Webb S, Jardine L, et al. Decoding

human fetal liver haematopoiesis. Nature. 2019;574(7778):365-71.

38.   Ezer D, Moignard V, Gottgens B, Adryan B. Determining Physical Mechanisms of Gene

Expression Regulation from Single Cell Gene Expression Data. PLoS Comput Biol.

2016;12(8):e1005072.

39.   Kim JK, Marioni JC. Inferring the kinetics of stochastic gene expression from single-cell

RNA-sequencing data. Genome Biol. 2013;14(1):R7.

40.   Edwards CR, Ritchie W, Wong JJ, Schmitz U, Middleton R, An X, et al. A dynamic intron

retention program in the mammalian megakaryocyte and erythrocyte lineages. Blood.

2016;127(17):e24-e34.

41.   Pimentel H, Parra M, Gee SL, Mohandas N, Pachter L, Conboy JG. A dynamic intron

retention program enriched in RNA processing genes regulates gene expression during terminal

erythropoiesis. Nucleic Acids Res. 2016;44(2):838-51.

42.   Liu X, Chen Y, Zhang Y, Liu Y, Liu N, Botten GA, et al. Multiplexed capture of spatial

configuration and temporal dynamics of locus-specific 3D chromatin by biotinylated dCas9.

Genome Biol. 2020;21(1):59.

43.   Hoppe PS, Schwarzfischer M, Loeffler D, Kokkaliaris KD, Hilsenbeck O, Moritz N, et al.

Early myeloid lineage choice is not initiated by random PU.1 to GATA1 protein ratios. Nature.

2016;535(7611):299-302.

44.   Choe KS, Ujhelly O, Wontakal SN, Skoultchi AI. PU.1 directly regulates cdk6 gene

expression, linking the cell proliferation and differentiation programs in erythroid cells. J Biol

Chem. 2010;285(5):3044-52.

45.   Carotta S, Wu L, Nutt SL. Surprising new roles for PU.1 in the adaptive immune

response. Immunol Rev. 2010;238(1):63-75.

46.   Roberts I, Alford K, Hall G, Juban G, Richmond H, Norton A, et al. GATA1-mutant clones

are frequent and often unsuspected in babies with Down syndrome: identification of a

population at risk of leukemia. Blood. 2013;122(24):3908-17.

47.   Bhatnagar N, Nizery L, Tunstall O, Vyas P, Roberts I. Transient Abnormal Myelopoiesis

and AML in Down Syndrome: an Update. Curr Hematol Malig Rep. 2016;11(5):333-41.

48.   Li Z, Godinho FJ, Klusmann JH, Garriga-Canut M, Yu C, Orkin SH. Developmental stage-

selective effect of somatically mutated leukemogenic transcription factor GATA1. Nat Genet.

2005;37(6):613-9.

49.   Haghverdi L, Lun ATL, Morgan MD, Marioni JC. Batch effects in single-cell RNA-

sequencing data are corrected by matching mutual nearest neighbors. Nat Biotechnol.

2018;36(5):421-7.

50.   Lun AT, McCarthy DJ, Marioni JC. A step-by-step workflow for low-level analysis of

single-cell RNA-seq data with Bioconductor. F1000Res. 2016;5:2122.

51.   Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, et al. Gene ontology:

tool for the unification of biology. The Gene Ontology Consortium. Nat Genet. 2000;25(1):25-9.

52.   The Gene Ontology C. The Gene Ontology Resource: 20 years and still GOing strong.

Nucleic Acids Res. 2019;47(D1):D330-D8.

53.   Ying QL, Wray J, Nichols J, Batlle-Morera L, Doble B, Woodgett J, et al. The ground state

of embryonic stem cell self-renewal. Nature. 2008;453(7194):519-23.

54.   Wray J, Kalkan T, Gomez-Lopez S, Eckardt D, Cook A, Kemler R, et al. Inhibition of

glycogen synthase kinase-3 alleviates Tcf3 repression of the pluripotency network and increases

embryonic stem cell resistance to differentiation. Nat Cell Biol. 2011;13(7):838-45.

55.   Ran FA, Hsu PD, Wright J, Agarwala V, Scott DA, Zhang F. Genome engineering using the

CRISPR-Cas9 system. Nat Protoc. 2013;8(11):2281-308.

56.   Lun ATL, Riesenfeld S, Andrews T, Dao TP, Gomes T, participants in the 1st Human Cell

Atlas J, et al. EmptyDrops: distinguishing cells from empty droplets in droplet-based single-cell

RNA sequencing data. Genome Biol. 2019;20(1):63.

57.   Wolf FA, Angerer P, Theis FJ. SCANPY: large-scale single-cell gene expression data

analysis. Genome Biol. 2018;19(1):15.

Figure Legends

Figure 1. Inferring Differentiation Trajectories at organismal scale

A. Pijuan-Sala et al. (2019) layout containing single-cell transcriptomes belonging from E6.5

to E8.5, colored by sampled time-point (left) and by cell-type (right). The overlaying

arrows result from applying the scVelo pipeline to the whole embryonic dataset and

represent inferred developmental trajectories. Arrowheads highlight the erythroid

branch, displaying scVelo trajectory predictions that are inconsistent with real-time

sampling.

B. Pijuan-Sala et al. (2019) layout highlighting single-cell transcriptomes belonging to E7.5

(left) and E8.5 (right) and colored by cell-type (see legend in A). The overlaying arrows

result from applying the scVelo pipeline to these individual time-points and represent

inferred developmental trajectories. Arrowheads highlight the erythroid branch.

Figure 2. Unspliced counts contribute to explaining the variability among cell types

A. Dimensionality reduction with the first two principal components/MOFA factors using

spliced reads alone (left), unspliced reads alone (middle) and both spliced and unspliced

(right). Single-cell transcriptomes are colored by cell-type annotation; see Figure 1 for full

legend.

B. MOFA characterization of spliced and unspliced reads assessing proportion of variance

explained (i), overlap in highly variable genes calculating using either spliced or unspliced

reads (ii), and factor weight distributions (iii).

Figure 3. A set of genes with complex expression kinetics confounds velocity estimation in

erythropoiesis

A. Illustration of phase plot representation in datasets of differentiating cell populations,

and associated scVelo predictions

B.   Illustration of strategy for MURK gene identification

C. Phase plots of representative MURK genes. X-axis: normalized imputed counts of spliced

transcript; y-axis: normalized imputed counts of unspliced transcript.

D. GO-term enrichment of MURK genes identified in mouse yolk sac erythropoiesis

E. Zoomed-in UMAP of the erythroid branch (see Figure 1 for full UMAP) with scVelo

734   calculations, before and after removing MURK genes identified in B. Distinct waves of

735   embryonic erythropoiesis are visible upon MURK gene removal, highlighted with

736   arrowheads.

737   **Figure 4. *In vivo* analysis of Gata1 function using a chimaera assay coupled with scRNA-Seq**

738   A. Schematic of the G1ER system (29, 30)

739   B. Behaviour of the 89 MURK genes identified in Figure 3 upon Gata1 induction in the G1ER

740   system (28). Wu et al. report that upon Gata1 induction they obtained a total of 2769

741   upregulated genes, 6079 mildly upregulated, 3566 downregulated, and 3445 with no

742   response.

743   C. UMAPS of *Gata1*[-] chimera cells allocated a hemato-endothelial identity colored by cell-

744   type (sub-clusters defined in Pijuan-Sala et al. (2019) - BP: Blood Progenitors, EC:

745   Endothelial Cells, Haem: Hemato-endothelial Progenitors, Mk: Megakaryocytes, My:

746   Myeloid cells, Ery: Erythroid cells) and split by genotype. Orange arrowheads highlight

747   increased population with megakaryocytic signature in Gata1[-] fraction.

748   D. UMAPS of *Gata1*[-] chimera cells allocated a hemato-endothelial identity colored by

749   sampling timepoint and split by genotype.

750   E. Barplots with the quantification of chimera cells mapping to each hemato-endothelial

751   lineage of the reference dataset (left) and to sampled time-points of the reference dataset

752   (right).

Figure 5. Gata1 chimaera assay reveals disruption of MURK genes and perturbed yolk sac

hematopoiesis

A. Violin plots of representative genes differentially regulated in *Gata1*[-] hematopoietic

lineages.

B. GO-term enrichment of genes downregulated in *Gata1*[-] Ery1 cells compared to their WT

counterparts in chimeras.

C. Venn diagram showing overlap between MURK genes and genes downregulated in *Gata1*[-]

Ery1 cells

D. Phase plots of MURK genes identified along erythroid differentiation, in E8.5 *Gata1*[-]

chimera datasets, colored by tdTom status.

Figure 6. Concept of dual kinetics of gene expression is also revealed in human foetal liver

hematopoiesis

A. UMAP representation of human fetal liver erythroid cell populations. The overlaying

arrows result from applying the scVelo pipeline using all genes (left) or after MURK gene

exclusion (right). Bottom UMAPs are colored by corresponding scVelo-inferred latent

time. In order to facilitate comparison with the mouse data, a new clustering was

performed on the erythroid cells, see Methods. MEMP: megakaryocyte-erythroid-mast

cell progenitor.

B. Phase plots of representative MURK genes identified in human fetal liver erythropoiesis

single-cell RNAseq dataset.

C. GO-term enrichment of MURK genes identified in human fetal liver erythropoiesis.

Supplementary Figures

1. Dimensionality reduction with the first three principal components/MOFA factors using spliced reads alone (left), unspliced reads alone (middle) and both spliced and unspliced (right). Single-cell transcriptomes are colored by cell-type annotation; see Figure 1 for full legend.

2. Identification of MURK genes along yolk sac erythropoiesis. A. Phase plots of representative scVelo driver genes, with scVelo model prediction overlayed (see also Supplementary Table 1). B. Distribution of annotated cell type (top) and sampling time-point (bottom) along scVelo calculated latent time, using all genes (left panels) and after removing the MURK genes identified in Figure 3B-C.

3. Pijuan-Sala et al. (2019) layout highlighting nearest neighbours of *Gata1*⁻ chimeras. In red are nearest neighbours of tdTom+ mutant cells, in black those of tdTom- wildtype cells. To compare with Figure 1A.

4. Impact of Gata1 knockout on *Spi1*/PU.1 expression on the hematoendothelial cell types. X-axis: *Spi1* $\log_2$(fold-change) in *Gata1*⁻ vs WT chimera cells and Atlas nearest neighbours. Y-axis: $\log_{10}$(FDR).

Supplementary Tables

1. Driver genes of the scVelo perdictions along erythroid differentiation, ranked by

likelihood in the dynamic model.

2. List of mouse MURK genes identified in Figure 3B-C, ranked by calculated increase in slope

value.

3. Differential Expression Analysis of Gata1$^-$ tdTom$^+$ vs WT tdTom$^-$ chimera cells. For the Mk

subset, given the low numbers of WT chimera cells present, the nearest neighbors from

the reference Atlas dataset were included in the comparison. LFC: log fold change.

4. List of human MURK genes identified in Figure 6, ranked by calculated increase in slope

value.

Figure 1

A

- E6.5
- E6.75
- E7.0
- E7.25
- E7.5
- E7.75
- E8.0
- E8.25
- E8.5

- Allantois
- Anterior Primitive Streak
- Blood progenitors 1
- Blood progenitors 2
- Cardiomyocytes
- Caudal Mesoderm
- Caudal epiblast
- Caudal neurectoderm
- Def. endoderm
- Endothelium
- Epiblast
- Erythroid1

- Erythroid2
- Erythroid3
- ExE mesoderm
- Forebrain/Midbrain/Hindbrain
- Gut
- Haematoendothelial progenitors
- Intermediate mesoderm
- Mesenchyme
- Mixed mesoderm
- NMP
- Nascent mesoderm

- Neural crest
- Notochord
- PGC
- Paraxial mesoderm
- Pharyngeal mesoderm
- Primitive Streak
- Rostral neurectoderm
- Somitic mesoderm
- Spinal cord
- Surface ectoderm
- Visceral endoderm

B

E7.5

E8.5

Figure 2

A

Spliced | Unspliced | Spliced + Unpliced (MOFA)

PC (or MOFA Factor) 2

PC (or MOFA Factor) 1

B

i) Variance explained

ii) Highly variable genes

iii) Factor weights

Figure 3

A

Upregulation: constant rates

Downregulation: constant rates

Upregulation: non-constant rates

Differentiation order

Differentiation path

scVelo prediction

Linear regression

B

m = 0.3
m = 0.4
m = 0.2
m = 0.5

m = 1.5
m = 0.3
m = 0.4
m = 0.5

C

Smim1

Hba-x

Blood Progenitors 1
Blood Progenitors 2
Erythroid 1
Erythroid 2
Erythroid 3

D

heme biosynthetic process
hydrogen peroxide catabolic process
protoporphyrinogen IX metabolic process
cellular detoxification
iron ion homeostasis
gas transport
erythrocyte maturation

-log10(FDR)

E

All genes

Removing MURK genes (**MU**ltiple **R**ate **K**inetics)

Velocity graph

Latent time

2nd wave

1st wave

Figure 4

Figure 5

Figure 6

A

- MEMP
- Early Erythroid
- Mid Erythroid
- Late Erythroid

B

HBG2   HBA2   HBM

ERMAP   HEMGN   ALAS2

C

oxygen transport
hydrogen peroxide catabolic process
cellular oxidant detoxification
protoporphyrinogen IX biosynthetic process
erythrocyte development
hemoglobin metabolic process
one-carbon compound transport
renal absorption

-log10(FDR)