To See, Not to See, or to See Poorly: Perceptual Quality and Guess Rate as a Function of

Electroencephalography (EEG) Brain Activity in an Orientation Perception Task

Sarah S. Sheldon[1] and Kyle E. Mathewson[1-2]

[1]University of Alberta, Department of Psychology, Faculty of Science

[2]University of Alberta, Neuroscience and Mental Health Institute, Faculty of Medicine and

Dentistry

Author Note

Sarah S. Sheldon https://orcid.org/0000-0002-4026-8939

Kyle E. Mathewson https://orcid.org/0000-0003-4856-3980

This research was supported by start-up funds from the Faculty of Science and an

NSERC Discovery grant (#RES0024267) awarded to Kyle Elliot Mathewson. The authors would

like you thank Erron Jacob Meneses, Jenny Le, and David Lam for assistance in data collection,

and Daniel Robles and Jonathan Kuziek for analysis feedback.

Correspondence concerning this article should be addressed to Sarah S. Sheldon,

Department of Psychology, University of Alberta, Edmonton, Alberta, T6G 2R3, Canada. Email:

ssheldon@ualberta.ca

Abstract

Brain oscillations are known to modulate detection of visual stimuli, but it is unclear if this is due to increased guess rate or decreased precision of the mental representation. Here we estimated quality and guess rate as a function of electroencephalography (EEG) brain activity using an orientation perception task. Errors on each trial were quantified as the difference between the target orientation and the orientation reported by participants with a response stimulus. Response errors were fitted to standard mixed model by Zhang and Luck (2008) to quantify how participants' guess rate and standard deviation parameters varied as a function of brain activity. Twenty-four participants were included in the analysis.

Within subjects, the power and phase of delta and theta post-target oscillatory activity were found to vary along with performance on the orientation perception task in that greater power and phase coherence in the 2-5 Hz band range was measured in trials with more accurate responses. In addition, the phase of delta and theta correlated with the degree of response error while oscillatory power did not have a relationship with trial-by-trial response errors. Analysis of task-related alpha activity yielded no significant results implying that alpha oscillations do not play an important role in orientation perception at single trial level. Across participants, only the standard deviation parameter correlated with oscillatory power in the high alpha and low beta frequency ranges. These results indicate that post-target power is associated with the precision of mental representations rather than the guess rate, both across trials within subjects and across subjects.

Keywords: EEG, time-frequency analysis, orientation perception, standard mixed model, visual perception

To See, Not to See, or to See Poorly: Perceptual Quality and Guess Rate as a Function of

Electroencephalography (EEG) Brain Activity in an Orientation Perception Task

Variations in neural activity give rise to observed variations in our visual perception.

(Chaumon & Busch, 2014; Mathewson et al., 2011; Samaha et al., 2020). While there has been a

plethora of research into the brain activity that drives this process, there remains noticeable gaps

in our understanding of how these processes work. One of the reasons for this might be because

investigators have left basic questions about the underlying neural mechanisms unanswered.

Specifically, does perception have an all-or-none principle where a certain threshold must be met

before a perceptual representation can form; or, does a perceptual representation always form but

the quality depends on the state of the neural activity? It seems unlikely researchers can form a

deep understanding of these perceptual processes without first determining what kind of process

we are looking at.

Traditionally, visual perception has been studied with two-alternative forced-choice (2-

AFC) tasks or similar discrete response paradigms. In these types of paradigms, participants are

required to select one out of two or more possible responses. Sometimes they are asked to choose

the correct stimulus out of an array of different stimuli or to simply report whether they detected

a visual stimulus. Although informative, these paradigms might miss certain aspects of visual

perception that cannot be measured with only two or three discrete categories. Continuous

behavior allows for more interpretable statistical measures since many of the common

descriptive and inferential statistics rely on the assumption of continuous measures. Furthermore,

while there are numerous inferential models based on the discrete response paradigms, it is

unclear whether these models are the best approach for investigating the neural mechanisms

underlying the perceptual responses they were designed for. In particular, the question about

what type of process underlies the formation of perceptual representations cannot be addressed

without a way to independently measure perceptual detection and perceptual quality.

To answer the question about the nature of perceptual processes, we need a task that can

measure visual perception on a continuous scale and a model that can quantify perceptual

responses in a way that will inform our question. With this goal in mind, we developed the

orientation perception task. The orientation perception task is an adaptation of Bae and Luck's

(2018) orientation memory task. The researchers originally used their task to investigate how

well information held in working memory can be decoded from brain activity. What makes the

task useful for the current study is that it allows participants to give a continuous response when

asked to report the orientation of the target. Performance can then be quantified as the angular

difference between the orientation of the target and the orientation reported by the participants,

referred to as response errors. This continuous measure of response errors can utilize a standard

mixed model introduced by Zhang and Luck (2008) to quantify parameters of interest such as

guess rate and precision of the perceptual representation. We here extend this method in order to

look for evidence of either all-or-none or graded perceptual mechanisms, and determine whether

these processes are individually related to electrical brain activity during the task.

To address the question about the type of process underlying visual perception, we

combined our adapted version of the visual orientation task with the standard mixed model and

electroencephalography (EEG). Because there are several novel parts in the current study, the

purpose was two-fold. First, we asked whether the standard mixed model could be extended to a

visual perception task? Second, we tested the hypothesis that there was a relationship between

EEG oscillatory activity, specifically in the alpha frequency band, and the standard mixed model

parameters (guess rate and standard deviation). To address these questions, we modified an

orientation memory task from Bae and Luck's (2018) so that it probed participants' perception of

the target's orientation rather than their ability to remember it. We then recorded EEG activity as

participants performed the adapted task so we can see how brain activity varies with their

perceptual performance.

<p align="center">**Materials and Methods**</p>

Participants

Twenty-eight participants from the University of Alberta community participated in the

study (age range = 17-35 years). Two participants were not included in the analysis due to

excessive movement artifacts (more than 25% of trials rejected due to artifacts). Another two

participants were excluded from the analysis due to having extreme outlying performance on the

task (see Behavioral Analysis in the Results section for more details). Participants were all right-

handed and had normal or corrected normal vision and no history of neurological problems. All

participants gave informed written consent, were either compensated at a rate of $10/hr or given

research credit for their time. The study adhered to the tenets of the Declaration of Helsinki and

was approved by the Internal Ethics Board at the University of Alberta.

Orientation Perception Task

Participants were seated 57 cm away from a 1920 x 1080 pixel^2 ViewPixx/EEG LCD

monitor (VPixx Technologies, Quebec, Canada) with a refresh rate of 120 Hz, simulating a CRT

display with LED backlight rastering. The rastering, along with 8-bit digital TTL output triggers

yoked to the onset and value of the top left pixel, allowed for submillisecond accuracy in pixel

illumination times, which were confirmed with a photocell prior to the experiment. Stimuli were

presented using a Windows 7 PC running MATLAB R2012b with the Psychophysics toolbox

(Version 3; Brainard, 1997; Pelli, 1997). The code running the task was a modified version of the

ColorWorkingMemoryExperiment.m code from MemToolbox (Suchow et al., 2013;

memtoolbox.org). The modified version of color working memory experiment can be found

here: https://github.com/APPLabUofA/OrientTask_paper/tree/master/OrientationTask. Video

output was sent to the ViewPixx/EEG with an Asus Striker GTX760 (Fremont, CA) graphics

processing unit.

Each trial began with a white fixation dot presented at the center of the monitor for 742,

783, 825, or 867 ms (target stimulus onset asynchrony; tSOA) after which the target appeared for

8.33 ms (one monitor refresh). The target was in the shape of a needle and was pointing toward

one of 24 predefined evenly spaced directions so that all the orientations covered 360 degrees.

The direction of the target was randomly selected on each trial. Of all the trials, 20% were

randomly chosen not to have a target. All aspects of the target-present and target-absent trials

were identical except that for the target-absent condition a blank interval replaced target

presentation. A backward mask lasting for 8.33 ms with a constant 41.7 ms target-mask SOA

(mSOA) appeared centrally. The mask was created by overlaying the target orientated in all 24

directions which created a star shape seen in Figure 1A. Following the mask offset, a 500 ms

blank interval period occurred.

After the blank interval, a response screen appeared with the needle in the center of the

screen. Using the computer mouse, participants were asked to rotate the needle so that it was

pointed in the same direction as the previous target. If participants detected a target but could not

remember its orientation, they were asked to guess the orientation of the target. Participants

could provide their response at their own pace. No feedback was given to participants. The next

trial began immediately after a needle's orientation was selected. See Figure 1A for a summary

of the task sequence and the stimulus dimensions.

Participants completed seven blocks consisting of 48 trials each, along with 20 practice

trials at the beginning of the experiment. Participants could rest at their own pace every 48 trials.

Extensive written and verbal instructions were presented to participants prior to the practice trials.

Instructions thoroughly explained and demonstrated each component that would compose a

single trial.

Before the orientation perception task, participants performed a staircased target detection

task that had the same parameters as the orientation perception task except that participants only

reported whether they saw the target or not using the keyboard. The target color was a gray

determined by a scalar value passed to the functions in Psychtoolbox. In the staircased target

detection task, the target color value could range from the background color (making it not

visible; corresponding value of 256) to black (making it the most visible; corresponding value of

0). This target gray value was adjusted throughout the task based on a 1-up/2-down staircasing

procedure targeting a 0.6 target detection rate for each individual (García-Pérez, 1998; Kingdom

& Prins, 2016). The staircased task consisted of three blocks of 32 trials. The target gray value

was determined for each participant by taking the average target gray value across the last two

blocks of trials. These final average values ranged from 70 to 112 and were used as the target

gray value in the orientation perception task.

The MATLAB code for the a staircased target detection task and the orientation

perception task are available at https://osf.io/cw7ux/ and

https://github.com/APPLabUofA/OrientTask_paper.

EEG Recording

During the experiment, EEG data was recorded from each participant with a Brain-Amp

32-channel amplifier (BrainVision) using gelled low-impedance electrodes (actiCAP passive

electrodes kept below 5 kΩ). Inter-electrode impedances were measured at the start of each

experiment. All electrodes were arranged in the same 10-20 positions (Fp2, Fp1, F4, F3, F8, F7,

FC2, FC1, FC6, FC5, C4, C3, CP2, CP1, CP6 CP5, P4, P3 P6, P5, P8, P7, PO4, PO3, O2, O1, Fz,

FCz, Cz, Pz, and Oz). In addition to the 31 EEG sensors, a ground electrode was used, positioned

at AFz. Two reference electrodes and the vertical and horizontal bipolar EOG were recorded

from passive Ag/AgCl easycap disk electrodes affixed on the mastoids, above and below the left

eye, and 1 cm lateral from the outer canthus of each eye. The bipolar channels were recorded

using the AUX ports of the Brain-Amp amplifier. SuperVisc electrolyte gel and mild abrasion

with a blunted syringe tip were used to lower impedances. Gel was applied and inter-electrode

impedances were lowered to less than 5 kΩ for all electrode sites. EEG data was recorded online

referenced to an electrode attached to the left mastoid. Offline, the data were re-referenced to the

arithmetically derived average of the left and right mastoid electrodes.

166   Data were digitized at 1000 Hz with a resolution of 24 bits. Data were filtered with an

online bandpass with cutoffs of 0.1 Hz and 250 Hz. The experiment was run in a dimly lit, sound

and radio frequency-attenuated chamber from Electromedical Instruments, with copper mesh

covering the window. The only electrical devices in the chamber were an amplifier, speakers,

keyboard, mouse, and monitor. The monitor ran on DC power from outside the chamber, the

keyboard and mouse were plugged into USB outside the chamber, and the speakers and amplifier

were both powered from outside the chamber, and nothing was plugged into the internal power

outlets. Any devices transmitting or receiving radio waves (e.g., cell phones) were removed from

the chamber for the duration of the experiment.

EEG Preprocessing

All analyses were completed using Matlab R2018b with the EEGLAB 13.6.5b (Delorme

& Makeig, 2004) and CircStat (Berens, 2009) toolboxes, as well as custom scripts. After the data

had been re-referenced offline, the bandpass FIR filter from EEGLAB was applied with lower

and upper cut-offs of 0.1 Hz and 50 Hz. Data was segmented into 3000 ms epochs aligned to

target onset (-1500 ms pre-target onset to 1500 ms post-target onset). The average voltage in the

200 ms baseline prior to the target was subtracted on each trial for every electrode, and trials

with absolute voltage fluctuations on any channel greater than 1000 μV were discarded. Eye

movements were then corrected with a regression-based procedure developed by Gratton, Coles,

and Donchin (1983). After a second baseline subtraction with 200 ms pre-target, trials with

remaining absolute voltage fluctuations on any channel greater than 500 μV were removed from

further analysis. Data was then subjected to visual inspection and manual rejection of trials

contaminated by artifacts. On average, 3% of trials were rejected during visual inspection. Other

than the two participants mentioned earlier, none of the remaining participants had more than

20% of trials rejected in this manner.

Data Analyses

Data analysis was performed using MATLAB R2018b (The MathWorks Inc, Natick, MA,

USA) and EEGLAB 13.6.5b (Delorme & Makeig, 2004). All statistical analyses were conducted

using MATLAB R2018b. Red-white-blue colormaps were created using the redblue.m function

by Auton (2009) found here: https://www.mathworks.com/matlabcentral/fileexchange/25536-

red-blue-colormap. The MATLAB code for data analysis is available at the GitHub repository

https://github.com/APPLabUofA/OrientTask_paper and the raw data files are available at

https://osf.io/cw7ux/.

Behavioral Data

Response errors on each trial were calculated by subtracting the orientation of the

response stimulus, as reported by the participant, from the orientation of the target stimulus (see

Figure 1B). The standard mixed model proposed by Zhang and Luck (2008) was fit to each

participant's response errors using the maximum likelihood estimation routine in the

MemToolbox (Suchow et al., 2013). According to the standard mixture model, response

deviations from the actual target orientation reflect a mixture of trials where the target's

orientation was detected and trials where participants did not detect the target so guessed

randomly. Therefore, the distribution of response errors consists of a mixture of a von Mises

distribution (representing the trials where the target's orientation was detected) and a uniform

distribution (random guesses (g)). Parameter sigma (σ) is the standard deviation of the von Mises

distribution, which represents the width of the response error distribution of trials that the target's

orientation was detected. Parameter g is the height of the uniform distribution representing the

guessing probability. A third parameter, mu (μ), which is the mean of the von Mises distribution

and represents systematic bias of the response error distribution was included in the standard

mixture model of two participants because the Bayesian information criterion (BIC), calculated

with the model comparison functions provided by MemToolbox (Suchow et al., 2013), indicated

that the three-parameter standard mixture model provided better fits for those two participants

(see Table 1). Although the three-parameter model was used for all analysis of those two

participants, the systematic bias was much smaller than the spacing between adjacent target

orientations (spacing was 15° whereas the two participants' μ was -4.2° and 2.7°) indicating that

those two participants had a slight clockwise and counterclockwise bias, respectively. The two-

parameter standard mixture model was used for all analysis of the remaining 22 participants

because it provided a better fit according to the BIC.

Time-Frequency Analyses

To calculate the phase angle and power for each trial, we used Morlet wavelet transform

of single trials using the newtimef() function of EEGLAB. A Morlet wavelet is a tapered sinusoid,

constructed by modulating a sine wave with a Gaussian envelope. Wavelet transformation was

created with 1.027 Hz linear steps and cycles starting from 2 cycles at 2 Hz to 12 cycles at

40 Hz. The output of this function was a matrix of complex values. The abs() function from

MATLAB was used to get the instantaneous amplitude and the instantaneous phase angle of each

trial was calculated using the angle() function from MATLAB.

Oscillatory power data was converted to Z scores by applying a single-trial normalization

procedure to the data from each participant at each electrode and frequency separately. This was

done because it helps disentangle background from task-related dynamics, allows for comparison

across different frequency bands and electrodes, and facilitates group-level analysis (Cohen,

2014). Each trial's entire epoch was used for the baseline during normalization because it has

been shown to be robust to the effects of noisy trials and is still conducive for analysis of pre-

and post-target activity (Grandchamp & Delorme, 2011). The downside to using the entire epoch

for the baseline normalization is that sustained changes throughout the trial period become

difficult to detect (Cohen, 2014). Ultimately, we considered this an acceptable trade-off for being

able to analyze oscillatory power and behavioral measures across participants without having to

make assumptions about frequency bands and time-windows such as would be required for

"binning" analysis used in previous studies (Barne et al., 2020; Iemi et al., 2017; Limbach &

Corballis, 2016; Samaha et al., 2017).

Accurate vs Guess Trials

To test for significant differences in brain activity on trials where participants had small

response errors compared to large response errors, we separated each participant's trials based on

the sigma value from the fits of their individual response errors to the standard mixed model.

This was done by defining each participant's trials with response errors between -0.75σ and

$+0.75\sigma$ as "accurate," and trials with response errors less than -1.5σ and greater than $+1.5\sigma$ as

"guesses" (Figure 2A). The logic is that trials). Trials where the participant clearly perceives the

target are likely trials with a response error less than the participant's overall response standard

deviation, and trials the participant has little to no perception of the target are likely trials with a

response error greater than just the participant's response standard deviation. It should be noted

that there are various reasons for participants to be accurate when they are guessing or have a

large response error when they accurately perceived the target. However, such events are thought

to be rare enough, or at least not systematic, that they will not unduly affect the overall

distribution.

The main reason we separated trials into "accurate" or "guess" was to see if the standard

mixed model parameters could be used to categorize trials in a meaningful way. Though this

method resulted in excluding some trials, namely ones that fell between the cutoffs for accurate

and guess, the outcome provided insight into how brain activity differs between different levels

of objective perceptual performance.

ERP Analyses. To remove the activity elicited by the mask without removing activity

resulting from the interaction of mask and target, the catch trials (mask-only) average was

subtracted from the orientation detection trials (target-plus-mask) average. ERP data was

submitted to a repeated measures, two-tailed permutation test based on the *tmax* statistic (Blair &

Karniski, 1993) using the mxt_perm1() function from the Mass Univariate ERP Toolbox (Groppe

et al., 2011). The time windows of interest were the P1 (80-140 ms), N1 (140-200 ms), P2 (200-

255 ms), N2 (255-360 ms), and P3 (360-500 ms) components. The ERP component time

windows were selected based on previous literature (Koivisto & Revonsuo, 2003, 2010). All 31

brain electrodes were included in the test. 100,000 random within-participant permutations were

used to estimate the distribution of the null hypothesis and the family-wise alpha (α) was set to

0.05. Based on this estimate, critical *t*-scores of +/-3.68 (df = 23) were derived. Any t-scores that

exceeded the critical *t*-score were considered statistically significant.

Oscillatory Power Analysis. To analyze differences in EEG band power between guess

and accurate trials, nonparametric permutation testing with a pixel-based multiple-comparison

correction procedure (Cohen, 2014) was used to analyze differences in EEG band power between

guess and accurate trials. The pixel-based multiple-comparison correction method involves

creating one distribution of the largest positive pixel value and another distribution of the largest

negative pixel value from each iteration of the permutation testing. After all iterations, the

statistical threshold is defined as the value corresponding to the 2.5[th] percentile of the smallest

values and the value corresponding to the 97.5[th] percentile of the largest values which are the

thresholds corresponding to an α of 0.05. Any pixel that has a value exceeding the upper or lower

value is considered significant. The pixel-based method corrects for multiple comparisons by

creating two distributions based on map-level information instead of pixel-level information. In

other words, this method results in two distributions of the most extreme null-hypothesis test

statistical values across all pixels rather than calculating null-hypothesis distributions for each

pixel (see Cohen (2014) for further details about pixel-based multiple-comparison correction

method). All analysis using nonparametric permutation testing with pixel-based multiple-

comparison correction performed 10,000 iterations per test. To obtain more stable estimates from

permutation testing, we ran a "meta-permutation test" by repeating the pixel-level permutation

procedure 10 times and then averaging the results (Cohen, 2014). It needs to be pointed out that a

"significant effect" determined by pixel-based permutation testing should not be considered a

precise estimate in the temporal and frequency domains. Although pixel-based permutation

testing is more stringent than cluster-based permutation tests (Cohen, 2014), caution should still

be used when interpreting "significant" differences, especially if the temporal and frequency

range of each pixel is relatively small.

297   **Oscillatory Phase Analysis.** To determine whether the mean phase values significantly

differ between accurate and guess trials, we used the circular Watson–Williams (W-W) test

which was calculated using the PhaseOpposition.m function by VanRullen (2016). We chose the

parametric circular W-W test because it has shown to be equivalent to the non-parametric phase

opposition sum (POS) measures under most conditions, and performed better in situations where

either the relative trial number or the ERP amplitude differed between the two trial groups

(VanRullen, 2016). The statistical significance of the W-W test across participants was

determined by combining the individual-level p-value time-frequency map at each electrode

across participants using Stouffer's method (Stouffer et al., 1949; VanRullen, 2016), which

transforms individual p-values into z-scores, combines them across participants, and converts the

resulting z-score to a combined probability. P-values were then corrected for multiple

comparisons across time points and frequencies at each electrode using the false discovery rate

(FDR) procedure described in Benjamini and Yekutieli (2001). Effects that satisfied a 5% FDR

criterion were considered significant.

Single-Trial EEG Activity and Response Errors

312   To test the correlation between time-frequency power and degree of response error,

313   Spearman's rho (r_s) correlation coefficients were calculated using a nonparametric permutation

314   testing approach with the pixel-based multiple-comparison correction procedure described above.

315   The null-hypothesis distribution was created by shuffling power values and response errors on

316   each trial with respect to each other. This provided a data-driven test of the null hypothesis that

317   there is no consistent relationship between degree of response error and EEG power.

318   To look at whether task performance is related to oscillatory phase, and if yes, at what

frequency, we used the weighted inter-trial phase clustering (wITPC) (Cohen, 2014; Cohen &

Voytek, 2013). The logic behind the inter-trial phase coherence (ITPC) is that a systematic

relation between EEG phase and behavioral outcome should result in a higher-than-chance ITPC

in each of the trial subgroups. However, if the phase of the EEG signal is randomized and

unpredictable, the distribution of phases at a given time period should follow a uniform

distribution over all trials. The problem with ITPC is that it assumes EEG phase is relevant to

experimental measures only when phase values are similar across trials (van Diepen & Mazaheri,

2018). Unlike ITPC, wITPC is sensitive to modulations of phase values even if those phases are

randomly distributed across trials as would be expected if response errors (which differs from

trial to trial) were modulated by oscillatory phase (Cohen, 2014; Cohen & Voytek, 2013).

329   The wITPC was computed for each participant as the resultant vector length, or ITPC, of

phase angles across trials once the length of each vector has been weighted by a variable of

interest (in this case, each trial's phase vector is weighted by the degree of response error on that

trial; see Figure 4A for example of computation) (Cohen, 2014; Cohen & Voytek, 2013). For

statistical testing, a null-hypothesis distribution was created by shuffling the phase values

relative to trial response error 10,000 times (see Figure 4A middle and bottom left). The wITPCz

was calculated as the wITPC standardized relative to the null-hypothesis distribution, providing a

z-value corresponding to the probability of finding the observed response error–phase

modulation by chance, given the measured data. To evaluate statistical significance of the

wITPCz across participants, we combined the individual-level p-value, calculated from the z-

values, time-frequency map at each electrode across participants using Stouffer's method

(Stouffer et al., 1949; VanRullen, 2016). P-values were then corrected for multiple comparisons

across time points and frequencies at each electrode using the false discovery rate (FDR)

procedure described in Benjamini and Yekutieli (2001). Effects that satisfied a 5% FDR criterion

were considered significant.

We chose to use the phase opposition measure and the wITPCz even though they are both

quantifying oscillatory phase because they provide different but complementary information

about the effects of phase. The phase opposition measure provides insight into whether there is

an overall consistent difference in the response-related (*i.e.*, "accurate" and "guess" response)

mean oscillation phase values. On the other hand, the weighted single-trial phase modulation

metric (*i.e.*, wITPCz) provides information about process-specific modulations of phase values

even when the phase distribution across trials is not consistent (Cohen, 2014; Cohen & Cavanagh,

2011). In other words, phase values do not need to be consistent over trials, they only need to be

consistently related to response errors (Cohen & Cavanagh, 2011). This is an important

distinction when trying to determine how much of the phase modulation is an artifact of the

stimulus-evoked activity and how much is related to the difference in task performance.

Relationship Between Oscillatory Power and Standard Mixed Model Parameters

For testing the relationship between time-frequency power and each participant's

standard deviation and guess rate parameter, Spearman's rho (r_s) correlation coefficients were

calculated using a nonparametric permutation testing approach with the pixel-based multiple-

comparison correction procedure described above. The null hypothesis correlation coefficients

were produced by randomly reassigning participant's estimated parameter values from the

standard mixed model (guess rate or standard deviation) to other participants then recomputing r_s.

362   **Results**

Accurate vs Guess Trials

Two participants were excluded before further analysis because one had a guess rate more

than three *IQR*s from the median and the other had a guess rate of 3.9e-15 indicating that the

staircasing procedure did not work properly for this individual. These participants were not

included in further analysis. Figure 2B shows the fit of the standard mixed model (or standard

mixed model with a bias parameter for the participants mentioned in the Methods section) to

each participant's response errors as well as the average fit of response errors across participants.

The remaining 24 participants had a mean guess rate of 0.19 ($SD = 0.18$) and mean standard

deviation (σ) parameter of 11.1 ($SD = 2.5$). The boxplots in Figure 2C summarizes the

distributions.

ERP Analysis

The ERPs from accurate and guess trials (Figure 3A) showed no statistical difference for

the first 200 ms following target onset across all electrodes. A divergence in the waveforms can

be seen in the P2 (200-300 ms) component for the frontal, central, and centroparietal electrodes

(Figure 3B left) in that the voltage of the guess trial ERPs was much attenuated compared to the

accurate trial ERPs. On the other hand, the voltage of the N2 (255-360 ms) component was more

negative in the guess trials than accurate trials (Figure 3A) and this difference was only

significant in the right frontocentral, central, centroparietal and parietal electrodes (Figure 3B

middle). Finally, the P3 (360-500 ms) component from the guess trials had a similar attenuation

as was seen in the P2 component (Figure 3A), but the distribution was more posterior with

significant effects seen in the right central, centroparietal and parietal electrodes as well as

bilateral parietooccipital and occipital electrodes (Figure 3B right).

Oscillatory Power Analysis

Pixel-based permutation test indicated significant differences between accurate and guess

trials within the lower part of the theta band and the delta band (2-4 Hz) which was observed to

start around 300 ms post-target onset in P8 and 350 ms post-target onset in P7 and had a duration

of around 75 ms (Figure 3C). Fp2 (not shown) showed a significant difference between accurate

and guess trials at around 325 ms to 385 ms post-target in the delta band (2-3 Hz) while P5 (not

shown) had a significant difference in the delta band (2-3 Hz) starting at around 450 ms post-

target with a duration of about 130 ms (extending to shortly after response screen onset). Finally,

the P3 electrode showed a brief significant difference in power at 6 Hz at about 330 ms post-

target (Figure 3C).

Overall, there was a trend for increased theta power in accurate trials compared to guess

trials, especially in left frontal and right parietal areas, though this difference was not significant

except for what is mentioned above. This lack of significance might be due to too much

variability in when theta power changed during the trial. It is also possible that theta reflects a

perceptual process that could occur in both accurate and guess trials, though more often or to a

greater degree in one type of trial compared to the other.

Oscillatory Phase Analysis

Similar to the oscillatory power results, significant differences in mean phase between

accurate and guess trials were found in the delta and theta frequency ranges following target

onset (Figure 3D). Central and bilateral frontal, frontocentral, and central electrodes show

significant phase differences in theta starting between 150-200 ms and terminating before 350 ms

post-target (not shown). The duration of this effect varies so that the more central electrodes

tended to have longer durations than those placed more laterally on the head. This effect was

more prevalent in the right hemisphere in the early period.

The most lateral parietal and centroparietal electrodes on the left side of the head show

significant phase differences in delta after the response screen onset. On the other hand, Pz, P3

(Figure 3D), P4, and PO3 electrodes had no significant effects in phase while P6 and P8 had

brief periods of significant phase differences starting a little after 300 ms until 500 ms post-target

in the delta frequency range. P7 showed a similar significant difference in mean phase at 2 Hz

starting around 225 ms and continuing until more than 200 ms after the response screen had been

presented (about 770 ms post-target; Figure 3D). The occipital electrodes had significant phase

effects primarily in the low theta and delta range (3-5 Hz) starting a little after 100 ms on the left

(not shown) and 150 ms on the right but were brief time periods until 200 ms post-target which

then had significant differences lasting for about 200 ms (Figure 3D). PO4 electrode showed a

similar difference, but the effect was less continuous and had larger *p*-values (*i.e.*, smaller phase

difference between accurate and guess trials) than the occipital electrodes; however, PO4 also

had significant differences in delta between 400 and 500 ms (Figure 3D).

It should be noted that the time window of significant phase opposition overlaps with the

significant differences in ERP amplitudes. ERP amplitudes have been shown to have a "masking"

effect on phase opposition measures resulting in a decrease in their statistical power (VanRullen,

2016) while at the same time the stimulus-evoked activity results in temporal distortion of

oscillatory activity towards earlier latencies (Brüers & VanRullen, 2017). While the circular W-

W test is relatively robust against the detrimental effect of ERPs, it does not negate their

influence entirely (VanRullen, 2016). Therefore, it is important to be aware that the significant

effects of phase are affected by the stimulus evoked activity and the different ERP amplitudes

between accurate and guess trials.

Single-Trial EEG Activity and Response Errors

To see whether response errors were related to trial-by-trial changes in pre-target alpha

power, the Spearman's rho correlations were calculated between oscillatory power and degree of

response error on each trial. Our results did not allow us to reject the null hypothesis. Follow-up

analysis on the entire time-frequency space also yielded no significant results. This is in line with

the lack of significant correlations between oscillatory power and guess rate (not shown).

To examine the relationship between response errors and the distribution of phase values,

we used the wITPCz. As can be seen in Figure 4B (bottom row), phase was primarily modulated

by degree of response errors in the delta frequency range in the posterior electrodes starting at

around 200 ms post-target and lasting until response screen onset. PO3 was similar except the

effect started at around 250 ms (not shown), but PO4 showed an effect starting at 350 ms and

lasting until almost 100 ms after the response screen onset (Figure 4B, bottom row). Parietal

electrodes show a significant effect in the low frequency bands starting at around 200 ms until

about 800 ms post-target (Figure 4B, top and bottom rows). Centroparietal and central electrodes

had a significant relationship between delta phase and response errors at around 300-400 ms

post-target and lasted until after response screen onset (not shown). All frontocentral electrodes

had response errors significantly related to delta phase starting between 200 ms and 300 ms post-

target and lasting until shortly after response screen onset (not shown). Out of all the frontal

electrodes, only Fz and F3 had a significant delta and theta phase relationship to response errors

(not shown).

Relationship Between Oscillatory Power and Standard Mixed Model Parameters

452   Using pixel-based multiple comparison correction procedure and Spearman's rho (r_s),

power at each pixel across time-frequency space was correlated with participants' parameter

values calculated from their response errors being fit to the standard mixed model. Results

indicate the oscillatory power and standard deviation (σ) parameter were significantly positively

correlated in right parietal, parietooccipital, occipital and central occipital electrodes (Figure 5).

The significant correlations were mostly in the alpha and low beta frequency ranges and occurred

between a time window starting at around 150 ms and ending at 440 ms post-target onset. The

exceptions were F3 and F4 (not shown) that showed significant positive correlations between

with the standard deviation parameter and power in the 3 Hz range during the onset of the

response screen for around 50 ms. The same analysis was done on oscillatory power and guess

rate which yielded no significant correlations though there was a trend towards a negative

correlation between guess rate and high alpha/low beta oscillatory power (not shown), especially

in more frontal electrodes.

465   **Discussion**

466   There were two goals for the present study. First, we asked what might the relationship be

between the parameters of the standard mixed model and EEG activity? We were successful in

this first goal: application of the standard mixed model to a visual perception task and relating

that performance measure and model parameters to EEG brain activity recorded during the task.

The second goal was to try to answer the question of how brain activity modulates perceptual

representation. Specifically, we wanted to see whether perception has an all-or-none principle

where a certain threshold must be met before a perceptual representation can form; or, if a

perceptual representation always forms but the quality depends on the state of the neural activity.

The results from the current study strongly point towards visual perception operating by the latter

principle rather than the former.

Accurate vs Guess Trials: ERP Activity

Significant differences in ERP waveforms of the accurate and guess trials started at

around 200 ms post-target onset in the anterior locations on the head (Figure 3B) which

corresponds to the P2 component (Di Russo et al., 2019; Key et al., 2005; Potts & Tucker, 2001).

The greater amplitude of the P2 for accurate guess trails fits with the P2 being related to the

salience of stimuli (Potts & Tucker, 2001) or stimulus recognition (Harel et al., 2016).

We also observed a more negative N2 for guess trials, that seems to go against current literature

which suggests that the onset of visual consciousness can be marked by a greater negativity in

the N2 time range due to an overlapping posterior negative component called the visual

awareness negativity (VAN) (Förster et al., 2020; Koivisto & Revonsuo, 2003, 2010). However,

the separation of trials based on response performance rather than awareness (according to

Koivisto and Revonsuo (2010), the VAN can only be reliably detected by comparing aware and

unaware conditions), would explain why we would not expect greater negativity in the N2 time

range for accurate ERPs compared to guess ERPs. Furthermore, the distribution of the observed

N2 component (mostly right central electrodes) does not match the typical posterior VAN

distribution (Förster et al., 2020). In fact, the N2 distribution points towards the presence of the

much larger P3 component. The most likely explanation is that the attenuated guess trial ERPs

(compared to accurate trials) results in a smaller or later positive voltage from the much larger P3

overlapping the negative deflection at the N2 time period (Förster et al., 2020; Koivisto &

Revonsuo, 2010) so that the N2 appears more negative in guess trials when it is actually due to

the P3 being smaller. The voltage increase of accurate compared to guess ERPs during the P3

time window follow previous findings on attention and visual perception (Key et al., 2005; Salti

et al., 2012). In the current study the P3 is likely related to processes such as stimulus

classification and saliency evaluation (Doradzińska et al., 2020; Förster et al., 2020; Helfrich &

Knight, 2019; Salti et al., 2012) and that these processes are overwhelmingly more relevant to

accurate orientation perception than processes for conscious perception as reflected by the

relatively smaller N2 ERP. Furthermore, the difference in ERPs between accurate trials and guess

trials during the N2 time period (see middle topographic plot of Figure 3B) indicates that the

processes reflected by the N2 likely occur in both types of trials though to differing degrees.

Instead, it appears that the main difference between accurate and guess trials is the extent that the

target can be classified or evaluated by the brain regardless of consciously perceiving the target.

Accurate vs Guess Trials: EEG Oscillatory Activity

We had originally hypothesized alpha activity, especially prior to target onset, mediating

task performance. However, the results did not support this hypothesis. As a visual perception

task, this might be surprising. There has been a recent accumulation of evidence suggesting pre-

target alpha power modulates visual detection performance by changing baseline neural

excitability (Iemi et al., 2019; Samaha et al., 2017, 2020). This might be attributed to using both

pre- and post-target activity in the pixel-based multiple correction procedure. Relative to post-

target activity, pre-stimulus power would be very small since the only thing for participants to do

was fixate on a white circle. As a result, differences in pre-target activity might not exceed

significance thresholds when those thresholds were based on the background spectra of the

combined pre- and post-target activity. However, we tested this possibility by analyzing power in

the alpha frequency during just the pre-target time period and found no significant differences

between accurate and guess trials (data not shown). Therefore, it is unlikely that a lack of

significant difference in pre-target alpha is due to an exceedingly high statistical threshold set by

the post-target activity.

522   The most likely reason for not finding a pre-target effect of alpha activity is that there

was no information for the participant to use before a target was presented, so there would be no

attention-related changes or task-based preparation prior to the target. The absence of significant

effects in post-target alpha oscillations suggests that alpha activity is not related to the processes

responsible for differences between accurate and guess trials. The fact that post-target alpha

activity was also unrelated to response errors on a trial-by-trial basis suggests that alpha activity

is not related to perception of a target's orientation. This is consistent with Bae and Luck (2018)

who found that alpha activity did not encode a target's orientation when the target's spatial

location was controlled for. An absence of a relationship between alpha power and performance

measures has been noted in other studies though they were not able to rule out changes in

detection bias which can be ruled out in the current study since guess rate was not related to

alpha activity (Benwell et al., 2017; Keitel et al., 2018). However, awareness bias might be a

possibility assuming orientation perception can proceed in the absence of conscious awareness

which seems likely based on previous research (Benwell et al., 2017; Doradzińska et al., 2020;

Koenig & Ro, 2019). In sum, it is likely that alpha oscillations are more related to a global

process that is not dependent on conscious awareness and would not change across trials such as

feature-independent stimulus processing.

The most significant differences between accurate and guess trials were in the delta and

low theta (2-4 Hz) frequency ranges, particularly in the parietal and parietooccipital electrodes

during the latter half of the post-stimulus epoch (Karakaş, 2020). A significant increase of delta

and low theta power in accurate compared to guess trials could be seen at around 300 ms though

their differences in mean phase, especially for theta, started earlier (200 ms post-target).

Although there was a trend in our data for increased theta power in accurate trials compared to

guess trials, this difference was not reliable.

The significant differences in theta and delta seem to correspond to changes in the N2 and

P3 ERP components observed in the time domain. Many studies have shown that delta and theta

are the primary contributors to the formation of the N2 and P3 (Harmony, 2013; Harper et al.,

2014; Karakaş et al., 2000). It has been proposed that delta contributes continuous positivity

throughout the ERP response while theta activity corresponds to the polarity change as the

negative deflection during the N2 shifts to the positive deflection during the P3 (Harper et al.,

2014). This fits with the observed oscillatory activity of accurate and guess trials (Figure 3C and

3D). Furthermore, the interplay between theta and delta oscillations would explain the indistinct

N2 but large P3 waveforms (Karakaş et al., 2000).

The increase in theta power during accurate trials relative to guess trials has the earliest

onset in left frontal and right parietal electrodes. Prolonged theta in right posterior electrodes is

accompanied by significant differences in mean (low) theta phase between accurate and guess

trials. Theta has been interpreted as being correlated with selective attentional processing (Başar

et al., 2001; Karakaş, 2020; Karakaş et al., 2000) and an increase in theta synchronization and

power has been associated with successful memory encoding (Klimesch, 1999) and right

hemispheric theta is greater than left when encoding visuospatial information (Sauseng et al.,

2004). Although the task in this study was not a memory task, encoding information is still a

viable way of looking at how the brain transforms the visual sensory information into perceptual

representations. In fact, theta has been shown to be sensitive to target and non-target stimuli

regardless of memory load (Palomäki et al., 2012). Therefore, it is probable that theta activity

might be related to the "encoding" of the target's orientation or just the detectability of target

itself. Both functions could occur in accurate and guess trials, though more often or to a greater

degree in accurate trials compared to guess trials. This would explain why theta oscillations

showed a for trend increased power in accurate trials compared to guess trials though this

difference was usually not significant.

In contrast to theta, delta showed a more spatially diffuse increase in power in accurate vs

guess trials with maxima over the lateral parietal areas. Delta phase differed significantly

between accurate and guess trials over the occipital and lateral parietal areas with the most robust

phase differences at around 350 ms post-target. Several experiments that showed increases in

delta activity during the performance of different mental tasks and conflict-monitoring paradigms

have led investigators to associate delta with modulation of networks that should be inactive to

accomplish the task (Harmony, 2013; Rawls et al., 2020). Investigators have also noted a close

association between delta oscillations and the P3 ERP waveform (Harper et al., 2014; Rawls et

al., 2020; Schürmann et al., 1995, 2001). If the P3 reflects post-perceptual processes, then it is

likely delta oscillations are also involved. Whether that role is as an inhibitory mechanism or not

remains unknown. Overall, when considering the time and frequency at which reliable

differences were observed between accurate and guess trials, it seems likely that this post-target

activity reflects differences in the level of perceptual encoding of the target orientation and the

subsequent precision of responses on the task.

Single-Trial EEG Activity and Response Errors

We were surprised to find no significant correlation between response error and

oscillatory power in any of the measured frequencies, but this could also be attributed to the lack

of sensitivity of the analysis method. Since failed to find support for our hypothesis regarding

alpha oscillations, we chose to take a conservative approach when analyzing the rest of the time-

frequency space. Using a less conservative method might help answer the question but that

would increase the risk of finding false positives which could be more harmful than accepting

false negatives.

Interestingly, we did find a relationship between response error and phase values in the

delta and theta frequencies similar to the significant differences in mean phase of accurate trials

compared to guess trials. In fact, phase modulation by response errors is even more pronounced

than the differences in phase between accurate and guess trials. This strongly suggests delta and

theta phase plays an important role in the amount of participants' response error on a trial-by-trial

bases. Delta and theta frequencies are usually associated with working memory and cognitive

control, but these results imply that they have an important role in visual perceptual processes as

well.

Relationship Between Oscillatory Power and Standard Mixed Model Parameters

There were no significant correlations between oscillatory power and the guess rate

parameter though there was a negative trend between guess rate and oscillatory power at the

theta and alpha frequencies. However, in the fitting of the standard mixed model, there is always

a slight negative correlation between the guess rate parameter and the standard deviation

parameter (Suchow et al., 2013). Consequently, the trend in correlations between guess rate and

oscillatory power might be an artifact of the relationship between the guess rate parameter and

the standard deviation parameter rather than a true association between oscillatory power and

guess rate. This suggests that there is truly no significant correlation between guess rate and

oscillatory power and points towards perception not operating by an all-or-none principle.

However, correlations between guess rate and oscillatory power were more transient and in

slightly lower frequencies than the correlations between power and the standard deviation

parameter. In other words, the correlation between oscillatory power and guess rate are not what

one would expect if they were mediated by the parameters' relationship. A relationship between

guess rate and oscillatory might be worth investigating further with a task that purposefully

changes the probability of guessing by alternating the detectability of the target.

Interestingly, frontal electrodes showed significant positive correlations between with the

standard deviation parameter and power in the 3 Hz range during the onset of the response screen

for around 50 ms. Accumulation of evidence has led investigators to associate frontal delta

oscillations with top-down control and response inhibition (De Vries et al., 2018; Harmony,

2013; Helfrich et al., 2017; Rawls et al., 2020). Considering the significant relationship between

delta activity and the standard deviation parameter is at the onset of the response screen, delta is

likely being engaged for response inhibition instead of cognitive control. The positive correlation

would further support this interpretation since an increase in response inhibition on a task that

does not benefit from it would, at best, have little impact and, at worst, impair responding and

increase response variability.

The significant positive correlation between alpha/low beta power and the standard

deviation parameter provides further support for the idea that neural activity modulates the

quality of the perceptual representation rather than just its detection. The standard deviation

parameter comes from the part of the standard mixed model representing trials where targets

were seen and is meant to reflect the precision with which a participant perceived the target's

orientation. It can be argued that the parameter also reflects the precision with which a

participant gave responses (*i.e.*, how carefully the participant oriented towards the response

stimulus). However, the location of these significant correlations over non-motor areas (right

medial parietal, parietooccipital and occipital electrodes) points toward the quality of perception

being more of an influence on the standard deviation parameter than the motor-related precision

with which participants responded.

If not motor-related, the question then is how decreases in high alpha/low beta power

improved the quality of the perceptual representation? It has been demonstrated that decreases in

alpha/low beta power correspond to an increase in cortical excitability (de Munck et al., 2007;

Gonçalves et al., 2006; Iemi & Busch, 2018) and an increase in stimulus-specific information

(Griffiths et al., 2019). Thus, our results could be interpreted as less alpha/low beta power mark

enhanced processing of the target resulting in more precise perceptual representations of the

target's orientation. At first these results might seem to go against the findings of the study by

Bae and Luck (2018). However, another way to interpret their results could be that post-target

alpha activity is more related to the encoding of the target's presence rather than specific features

such as how the target was oriented. In that case, our results would be corroborating their

findings and would explain why alpha/low beta tracks spatial location so well in attentional and

working memory studies, but not the actual features of stimuli (Bae & Luck, 2019, 2018; Foster

& Awh, 2019; Griffiths et al., 2019; Palomäki et al., 2012; Pesonen et al., 2007).

Conclusion

Overall, we have shown that the standard mixed model can be extended to a visual

perception task and that doing so provides important insight into how brain activity shapes visual

perception. Within-group results suggest that the difference between accurate responses and

guesses are in the amount of post-target oscillatory activity in the delta and theta frequencies

where greater activity was associated with smaller errors. This is evident by the significant

differences between accurate and guess trials in power and phase at the delta and theta

frequencies as well as the associated ERPs.

At the between-group level, those with lower post-target power in high alpha and low

beta frequencies had more precision in their perceptual performance. This suggests that

participants with greater cortical activity in the posterior regions have enhanced perceptual

processing leading to a more precise perceptual representation of the targets' orientations. The

posterior location of this activity precludes this being a motor-related effect.

Taken together, our results point towards a perceptual representation always forming but

the quality, and thus, detectability, depends on the level of neural activity after stimulus onset.

This is consistent with previous findings indicating that the brain holds stimulus information

even when the stimulus is not consciously accessible to the participant and the difference

between conscious and unconscious processing has to do with the difference in strength of the

stimulus representation (Doradzińska et al., 2020; Mai et al., 2019).

It is important to note that our results are not suggesting pre-target EEG activity has no

effect on visual perception. There is a lot of evidence to contrary. It is possible the analysis

methods were not sensitive enough to pick up the trial-by-trial changes in pre-target activity, but

it more likely there was no task-related modulation of pre-target activity due to participants

having no information about an upcoming trial. A follow-up study that directly manipulates

attention or detectability of the target might be able to better test how differences in brain activity

prior to target onset affect visual perception and subsequent task performance.

677   **References**

Auton, A. (2009). *Red Blue Colormap*. MATLAB Central File Exchange.

https://www.mathworks.com/matlabcentral/fileexchange/25536-red-blue-colormap

Bae, G.-Y., & Luck, S. (2019). Appropriate Correction for Multiple Comparisons in Decoding of

ERP Data: A Re-Analysis of Bae & Luck (2018). *BioRxiv*, Article 672741.

https://doi.org/10.1101/672741

Bae, G.-Y., & Luck, S. J. (2018). Dissociable decoding of spatial attention and working memory

from EEG oscillations and sustained potentials. *Journal of Neuroscience*, *38*(2), 409–422.

https://doi.org/10.1523/JNEUROSCI.2860-17.2017

Barne, L. C., de Lange, F. P., & Cravo, A. M. (2020). Prestimulus alpha power is related to the

strength of stimulus representation. *Cortex*, *132*, 250–257.

https://doi.org/10.1016/j.cortex.2020.08.017

Başar, E., Başar-Eroglu, C., Karakaş, S., & Schürmann, M. (2001). Gamma, alpha, delta, and

theta oscillations govern cognitive processes. *International Journal of Psychophysiology*,

39(2–3), 241–248. https://doi.org/10.1016/S0167-8760(00)00145-8

Benjamini, Y., & Yekutieli, D. (2001). The control of the false discovery rate in multiple testing

under dependency. *Annals of Statistics*, *29*(4), 1165–1188.

https://doi.org/10.1214/aos/1013699998

Benwell, C. S. Y., Tagliabue, C. F., Veniero, D., Cecere, R., Savazzi, S., & Thut, G. (2017).

Prestimulus EEG Power Predicts Conscious Awareness But Not Objective Visual

Performance. *ENeuro*, *4*(6). https://doi.org/10.1523/ENEURO.0182-17.2017

Berens, P. (2009). CircStat: a MATLAB toolbox for circular statistics. *Journal of Statistical*

Software, *31*(10), 679–685. https://doi.org/10.1016/j.amp.2012.05.023

Blair, R. C., & Karniski, W. (1993). An alternative method for significance testing of waveform

difference potentials. *Psychophysiology*, *30*(5), 518–524. https://doi.org/10.1111/j.1469-

8986.1993.tb02075.x

Brainard, D. H. (1997). The Psychophysics Toolbox. *Spatial Vision*, *10*(4), 433–436.

http://www.ncbi.nlm.nih.gov/pubmed/9176952

Brüers, S., & VanRullen, R. (2017). At what latency does the phase of brain oscillations

influence perception? *ENeuro*, *4*(3). https://doi.org/10.1523/ENEURO.0078-17.2017

Chaumon, M., & Busch, N. A. (2014). Prestimulus neural oscillations inhibit visual perception

via modulation of response gain. *Journal of Cognitive Neuroscience*, *26*(11), 2514–2529.

https://doi.org/10.1162/jocn_a_00653

Cohen, M. X. (2014). *Analyzing neural time series data: theory and practice*. MIT Press.

Cohen, M. X., & Cavanagh, J. F. (2011). Single-trial regression elucidates the role of prefrontal

theta oscillations in response conflict. *Frontiers in Psychology*, *2*.

https://doi.org/10.3389/fpsyg.2011.00030

Cohen, M. X., & Voytek, B. (2013). Linking nonlinear neural dynamics to single-trial human

behavior. In M. (Meyer) Z. Pesenson (Ed.), *Multiscale Analysis and Nonlinear Dynamics*

(pp. 217–232). Wiley-VCH Verlag GmbH & Co. KGaA.

https://doi.org/10.1002/9783527671632

de Munck, J. C., Gonçalves, S. I., Huijboom, L., Kuijer, J. P. A., Pouwels, P. J. W., Heethaar, R.

M., & Lopes da Silva, F. H. (2007). The hemodynamic response of the alpha rhythm: An

EEG/fMRI study. *NeuroImage*, *35*(3), 1142–1151.

https://doi.org/10.1016/j.neuroimage.2007.01.022

De Vries, I. E. J., Van Driel, J., Karacaoglu, M., & Olivers, C. N. L. (2018). Priority Switches in

Visual Working Memory are Supported by Frontal Delta and Posterior Alpha Interactions.

Cerebral Cortex, *28*(11), 4090–4104. https://doi.org/10.1093/cercor/bhy223

Delorme, A., & Makeig, S. (2004). EEGLAB: An open source toolbox for analysis of single-trial

EEG dynamics including independent component analysis. *Journal of Neuroscience*

Methods, *134*(1), 9–21. https://doi.org/10.1016/J.JNEUMETH.2003.10.009

Di Russo, F., Berchicci, M., Bianco, V., Perri, R. L., Pitzalis, S., Quinzi, F., & Spinelli, D. (2019).

Normative event-related potentials from sensory and cognitive tasks reveal occipital and

frontal activities prior and following visual events. *NeuroImage*, *196*, 173–187.

https://doi.org/10.1016/j.neuroimage.2019.04.033

Doradzińska, Ł., Wójcik, M. J., Paź, M., Nowicka, M. M., Nowicka, A., & Bola, M. (2020).

Unconscious perception of one's own name modulates amplitude of the P3B ERP

component. *Neuropsychologia*, *147*, 107564.

https://doi.org/10.1016/j.neuropsychologia.2020.107564

Förster, J., Koivisto, M., & Revonsuo, A. (2020). ERP and MEG correlates of visual

consciousness: The second decade. *Consciousness and Cognition*, *80*, Article 102917.

https://doi.org/10.1016/j.concog.2020.102917

Foster, J. J., & Awh, E. (2019). The role of alpha oscillations in spatial attention: limited

evidence for a suppression account. *Current Opinion in Psychology*, *29*, 34–40.

https://doi.org/10.1016/J.COPSYC.2018.11.001

García-Pérez, M. A. (1998). Forced-choice staircases with fixed step sizes: asymptotic and small-

sample properties. *Vision Research*, *38*(12), 1861–1881. https://doi.org/10.1016/S0042-

6989(97)00340-4

Gonçalves, S. I., de Munck, J. C., Pouwels, P. J. W., Schoonhoven, R., Kuijer, J. P. A., Maurits, N.

M., Hoogduin, J. M., Van Someren, E. J. W., Heethaar, R. M., & Lopes da Silva, F. H.

(2006). Correlating the alpha rhythm to BOLD using simultaneous EEG/fMRI: Inter-subject

variability. *NeuroImage*, *30*(1), 203–213. https://doi.org/10.1016/j.neuroimage.2005.09.062

Grandchamp, R., & Delorme, A. (2011). Single-trial normalization for event-related spectral

decomposition reduces sensitivity to noisy trials. *Frontiers in Psychology*, *2*, 236.

https://doi.org/10.3389/fpsyg.2011.00236

Gratton, G., Coles, M. G. ., & Donchin, E. (1983). A new method for off-line removal of ocular

artifact. *Electroencephalography and Clinical Neurophysiology*, *55*(4), 468–484.

https://doi.org/10.1016/0013-4694(83)90135-9

Griffiths, B. J., Mayhew, S. D., Mullinger, K. J., Jorge, J., Charest, I., Wimber, M., & Hanslmayr,

S. (2019). Alpha/beta power decreases track the fidelity of stimulus specific information.

ELife, *8*, Article e49562. https://doi.org/10.7554/eLife.49562

Groppe, D. M., Urbach, T. P., & Kutas, M. (2011). Mass univariate analysis of event-related

brain potentials/fields I: A critical tutorial review. *Psychophysiology*, *48*(12), 1711–1725.

https://doi.org/10.1111/j.1469-8986.2011.01273.x

Harel, A., Groen, I. I. A., Kravitz, D. J., Deouell, L. Y., & Baker, C. I. (2016). The temporal

dynamics of scene processing: A multifaceted EEG investigation. *ENeuro*, *3*(5), Article

e0139-16.2016. https://doi.org/10.1523/ENEURO.0139-16.2016

Harmony, T. (2013). The functional significance of delta oscillations in cognitive processing.

Frontiers in Integrative Neuroscience, *7*, Article 83.

https://doi.org/10.3389/fnint.2013.00083

Harper, J., Malone, S. M., & Bernat, E. M. (2014). Theta and delta band activity explain N2 and

P3 ERP component activity in a go/no-go task. *Clinical Neurophysiology*, *125*(1), 124–132.

https://doi.org/10.1016/j.clinph.2013.06.025

Helfrich, R. F., Huang, M., Wilson, G., & Knight, R. T. (2017). Prefrontal cortex modulates

posterior alpha oscillations during top-down guided visual perception. *Proceedings of the*

National Academy of Sciences of the United States of America, *114*(35), 9457–9462.

https://doi.org/10.1073/pnas.1705965114

Helfrich, R. F., & Knight, R. T. (2019). Cognitive neurophysiology: Event-related potentials. In

Handbook of Clinical Neurology (Vol. 160, pp. 543–558). Elsevier B.V.

https://doi.org/10.1016/B978-0-444-64032-1.00036-9

Iemi, L., & Busch, N. A. (2018). Moment-to-moment fluctuations in neuronal excitability bias

subjective perception rather than strategic decision-making. *ENeuro*, *5*(3), ENEURO.0430-

17.2018. https://doi.org/10.1523/ENEURO.0430-17.2018

Iemi, L., Busch, N. A., Laudini, A., Haegens, S., Samaha, J., Villringer, A., & Nikulin, V. V.

(2019). Multiple mechanisms link prestimulus neural oscillations to sensory responses.

ELife, *8*, Article e43620. https://doi.org/10.7554/eLife.43620.001

Iemi, L., Chaumon, M., Crouzet, S. M., & Busch, N. A. (2017). Spontaneous neural oscillations

bias perception by modulating baseline excitability. *Journal of Neuroscience*, *37*(4), 807–

819. https://doi.org/10.1523/JNEUROSCI.1432-16.2016

Karakaş, S. (2020). A review of theta oscillation and its functional correlates. *International*

Journal of Psychophysiology. https://doi.org/10.1016/j.ijpsycho.2020.04.008

Karakaş, S., Erzengin, Ö. U., & Başar, E. (2000). A new strategy involving multiple cognitive

paradigms demonstrates that ERP components are determined by the superposition of

oscillatory responses. *Clinical Neurophysiology*, *111*(10), 1719–1732.

https://doi.org/10.1016/S1388-2457(00)00418-1

Keitel, C., Benwell, C. S. Y., Thut, G., & Gross, J. (2018). No changes in parieto-occipital alpha

during neural phase locking to visual quasi-periodic theta-, alpha-, and beta-band

stimulation. *European Journal of Neuroscience*, 1–15. https://doi.org/10.1111/ejn.13935

Key, A. P. F., Dove, G. O., & Maguire, M. J. (2005). Linking brainwaves to the brain: An ERP

primer. *Developmental Neuropsychology*, *27*(2), 183–215.

https://doi.org/10.1207/s15326942dn2702_1

Kingdom, F. A. A., & Prins, N. (2016). Chapter 5 – Adaptive Methods. In *Psychophysics*

(Second Edi, pp. 119–148). Academic Press. https://doi.org/10.1016/B978-0-12-407156-

8.00005-0

Klimesch, W. (1999). EEG alpha and theta oscillations reflect cognitive and memory

performance: a review and analysis. *Brain Research Reviews*, *29*(2–3), 169–195.

https://doi.org/10.1016/S0165-0173(98)00056-3

Koenig, L., & Ro, T. (2019). Dissociations of conscious and unconscious perception in TMS-

induced blindsight. *Neuropsychologia*, *128*, 215–222.

https://doi.org/10.1016/j.neuropsychologia.2018.03.028

Koivisto, M., & Revonsuo, A. (2003). An ERP study of change detection, change blindness, and

visual awareness. *Psychophysiology*, *40*(3), 423–429. https://doi.org/10.1111/1469-

8986.00044

Koivisto, M., & Revonsuo, A. (2010). Event-related brain potential correlates of visual

awareness. *Neuroscience & Biobehavioral Reviews*, *34*(6), 922–934.

https://doi.org/10.1016/j.neubiorev.2009.12.002

Limbach, K., & Corballis, P. M. (2016). Prestimulus alpha power influences response criterion in

a detection task. *Psychophysiology*, *53*(8), 1154–1164. https://doi.org/10.1111/psyp.12666

Mai, A. T., Grootswagers, T., & Carlson, T. A. (2019). In search of consciousness: Examining the

temporal dynamics of conscious visual perception using MEG time-series data.

Neuropsychologia, *129*, 310–317. https://doi.org/10.1016/j.neuropsychologia.2019.04.015

Mathewson, K. E., Lleras, A., Beck, D. M., Fabiani, M., Ro, T., & Gratton, G. (2011). Pulsed out

of awareness: EEG alpha oscillations represent a pulsed-inhibition of ongoing cortical

processing. *Frontiers in Psychology*, *2*, Article 99. https://doi.org/10.3389/fpsyg.2011.00099

Palomäki, J., Kivikangas, M., Alafuzoff, A., Hakala, T., & Krause, C. M. (2012). Brain

oscillatory 4-35 Hz EEG responses during an n-back task with complex visual stimuli.

Neuroscience Letters, *516*(1), 141–145. https://doi.org/10.1016/j.neulet.2012.03.076

Pelli, D. G. (1997). The VideoToolbox software for visual psychophysics: transforming numbers

into movies. *Spatial Vision*, *10*(4), 437–442. http://www.ncbi.nlm.nih.gov/pubmed/9176953

Pesonen, M., Hämäläinen, H., & Krause, C. M. (2007). Brain oscillatory 4–30 Hz responses

during a visual n-back memory task with varying memory load. *Brain Research*, *1138*, 171–

177. https://doi.org/10.1016/J.BRAINRES.2006.12.076

Potts, G. F., & Tucker, D. M. (2001). Frontal evaluation and posterior representation in target

detection. *Cognitive Brain Research*, *11*(1), 147–156. https://doi.org/10.1016/S0926-

6410(00)00075-6

Rawls, E., Miskovic, V., & Lamm, C. (2020). Delta phase reset predicts conflict-related changes

in P3 amplitude and behavior. *Brain Research*, *1730*, 146662.

https://doi.org/10.1016/j.brainres.2020.146662

Salti, M., Bar-Haim, Y., & Lamy, D. (2012). The P3 component of the ERP reflects conscious

perception, not confidence. *Consciousness and Cognition*, *21*(2), 961–968.

https://doi.org/10.1016/j.concog.2012.01.012

Samaha, J., Iemi, L., Haegens, S., & Busch, N. A. (2020). Spontaneous Brain Oscillations and

Perceptual Decision-Making. *Trends in Cognitive Sciences*, *24*(8), 639–653.

https://doi.org/10.1016/j.tics.2020.05.004

Samaha, J., Iemi, L., & Postle, B. R. (2017). Prestimulus alpha-band power biases visual

discrimination confidence, but not accuracy. *Consciousness and Cognition*, *54*, 47–55.

https://doi.org/10.1016/J.CONCOG.2017.02.005

Sauseng, P., Klimesch, W., Doppelmayr, M., Hanslmayr, S., Schabus, M., & Gruber, W. R.

(2004). Theta coupling in the human electroencephalogram during a working memory task.

Neuroscience Letters, *354*(2), 123–126. https://doi.org/10.1016/j.neulet.2003.10.002

Schürmann, M., Başar-Eroglu, C., Kolev, V., & Başar, E. (1995). A new metric for analyzing

single-trial event-related potentials (ERPs): application to human visual P300 delta response.

Neuroscience Letters, *197*(3), 167–170. https://doi.org/10.1016/0304-3940(95)11912-G

Schürmann, M., Başar-Eroglu, C., Kolev, V., & Başar, E. (2001). Delta responses and cognitive

processing: Single-trial evaluations of human visual P300. *International Journal of*

Psychophysiology, *39*(2–3), 229–239. https://doi.org/10.1016/S0167-8760(00)00144-6

Stouffer, S. A., Suchman, E. A., Devinney, L. C., Star, S. A., & Williams Jr., R. M. (1949).

Studies in social psychology in World War II: the American soldier. In *Adjustment During*

Army Life (Vol. 1). Princeton University Press. https://psycnet.apa.org/record/1950-00790-

000

Suchow, J. W., Brady, T. F., Fougnie, D., & Alvarez, G. A. (2013). Modeling visual working

memory with the MemToolbox. *Journal of Vision*, *13*(10), 1–8.

https://doi.org/10.1167/13.10.9

van Diepen, R. M., & Mazaheri, A. (2018). The Caveats of observing Inter-Trial Phase-

Coherence in Cognitive Neuroscience. *Scientific Reports*, *8*(1), 2990.

https://doi.org/10.1038/s41598-018-20423-z

VanRullen, R. (2016). How to Evaluate Phase Differences between Trial Groups in Ongoing

Electrophysiological Signals. *Frontiers in Neuroscience*, *10*, 426.

https://doi.org/10.3389/fnins.2016.00426

Zhang, W., & Luck, S. J. (2008). Discrete fixed-resolution representations in visual working

memory. *Nature*, *453*(7192), 233–235. https://doi.org/10.1038/nature06860

Tables

Table 1

List of goodness-of-fit measures comparing the standard mixed model to the standard mixed

model with a bias parameter (µ) for each participant.

	Log Likelihood			Bayesian Information Criterion (BIC)		
	Standard Model	Model + Bias	Difference	Standard Model	Model + Bias	Difference
Participant 1	-1255.82	-1255.33	-0.49	2526.45	2532.88	-6.44
Participant 2	-1213.93	-1212.78	-1.15	2442.77	2447.92	-5.15
Participant 3	-1095.28	-1094.58	-0.70	2205.45	2211.50	-6.05
Participant 4	-1093.29	-1093.24	-0.05	2201.25	2208.50	-7.24
Participant 5	-1191.23	-1190.92	-0.31	2397.23	2404.01	-6.77
Participant 6	-1113.21	-1111.14	-2.07	2241.28	2244.57	-3.29
Participant 7	-965.54	-964.92	-0.62	1945.90	1952.06	-6.16
Participant 8	-1022.30	-1021.07	-1.23	2059.48	2064.47	-4.99
Participant 9	-1109.18	-1109.12	-0.06	2233.12	2240.39	-7.27
Participant 10	-1422.57	-1422.42	-0.15	2859.94	2867.05	-7.11
Participant 11	-1384.47	-1384.46	-0.00	2783.78	2791.20	-7.42
Participant 12	-1017.58	-1015.52	-2.06	2049.58	2052.67	-3.09
Participant 13	-1498.89	-1498.55	-0.34	3012.59	3019.32	-6.73
Participant 14	-1071.57	-1071.55	-0.02	2157.98	2165.36	-7.38
Participant 15	-1124.71	-1123.84	-0.87	2264.26	2269.94	-5.68
Participant 16	-985.46	-985.17	-0.29	1985.36	1991.99	-6.63
Participant 17	-1149.36	-1149.26	-0.11	2313.61	2320.84	-7.23
Participant 18	-1136.58	-1136.50	-0.09	2287.86	2295.02	-7.17
Participant 19	-1133.03	-1132.91	-0.12	2280.88	2288.04	-7.16
Participant 20	-1276.89	-1275.81	-1.08	2568.44	2573.61	-5.17
Participant 21	-1138.27	-1133.04	-5.23	2291.40	2288.37	3.03[a]
Participant 22	-1003.69	-988.80	-14.89	2022.10	1999.69	22.41[a]
Participant 23	-1011.18	-1010.68	-0.50	2037.17	2043.57	-6.40
Participant 24	-1166.38	-1162.88	-3.49	2347.60	2348.04	-0.44

873 *Note*. Difference is calculated as the goodness-of-fit measure of the standard mixed model with a

874 bias model subtracted from the goodness-of-fit measure of the standard mixed model.

875 [a] Positive difference indicates the standard mixed model with a bias parameter (µ) fits better to

876 the participants' response errors (*i.e.*, has a smaller BIC) than the standard mixed model.

877   **Figure Captions**

Figure 1.

A) Sequence of task events with duration of each screen presentation and sizes of fixation, target,

mask, and response stimuli. Sizes are in degrees of visual angle. **B)** Example of response error

calculation. Response errors are reported in degrees.

Figure 2.

A) Representation of how trials were split into the two categories, Accurate and Guess. This was

done by defining each participant's trials with response errors between -0.75σ and $+0.75\sigma$ as

"accurate" (shaded yellow region), and trials with response errors less than -1.5σ and greater

than $+1.5\sigma$ as "guesses" (shaded pink regions). Trials that did not fall in either category were

discarded from this analyses between the two trial categories. **B)** Left, model fit of each

participant. Right, solid purple line is the average of participants' model fit. Light purple area

represents SEM. Zoomed in window shows upward shift of the averaged model fit showing a

non-zero guess rate. **C)** Boxplots of parameter values (left) guess rate and (right) standard

deviation estimated from model fits of participants' response errors.

Figure 3.

A) ERPs of accurate trials and guess trials aligned to target onset at selected electrodes and

averaged across all electrodes (Grand Average). Light yellow and light pink shaded areas around

waveforms represent ±SEM of accurate trials and guess trials, respectively. Mask trial EPRs have

been subtracted out of the ERPs to remove mask-related brain activity except for the Grand

Average + Mask ERPs plot. Time period shaded in green indicates significant difference between

accurate and guess trials 255-360 ms (N2) post-target. Blue shaded time period indicates

significant difference between accurate and guess trials 360-500 ms (P3) post-target. **B)**

Topographies of the voltage distribution difference between accurate and guess trials. The time

periods in each topography are as follows: 200-255 ms (P2), 255-360 ms (N2), and 360-500 ms

(P3) post-target. Stars indicate electrodes with significant differences between accurate and guess

ERPs. Mask trial activity has been subtracted out of the ERPs to remove mask-related brain

activity. **C)** Oscillatory power analysis of accurate vs guess trials. Time-frequency plots showing

the results of the statistical analysis of the difference in power between accurate vs guess trials at

each time-frequency point at selected electrodes and the average of all electrodes (Grand

Average). Black contour denotes statistically significant differences (after applying the pixel-

based multiple-comparison correction procedure) at $p < 0.05$. **D)** Oscillatory phase analysis of

accurate vs guess trials. Time-frequency plots showing the p-values (after applying FDR

correction) from the statistical analysis testing differences in the mean phase of accurate trials vs

guess trials. Plots are from selected electrodes and the average of all electrodes (Grand Average).

Time-frequency points with p-values above the threshold of FDR correction for multiple

comparisons were set to 1.

Figure 4.

A) Example computation of weighted inter-trial phase clustering (wITPC) to relate single-trial

phase to degree of response error. Top left, single-trial prestimulus phase vectors are shown as

black lines and are not clustered across trials due to the randomization of the fixation length,

leading to a low resultant vector length (i.e., low ITPC). Right, histogram of null-hypothesis

vector lengths created by shuffling the mapping of the trial response error values to the trial

phase values. The observed vector length (large arrow) is calculated from the distribution shown

in bottom left panel. Bottom left shows how the length each trial's phase vector is scaled by that

trial's degree response error and a weighted ITPC is computed, reflecting the relationship

between the distribution of phase angles and degree of response error even though the

distribution of the phase angles themselves is uniform (as seen in the top left plot). The wITPCz

value is the wITPC standardized relative to the null-hypothesis distribution seen in the histogram

plot. Example based on figure by Cohen (2014) **B)** Time-frequency plots of analysis relating

single-trial phase activity and response errors. Significant *p*-values indicating that the normalized

distance of the observed wITPC (i.e., wITPCz) is significantly different from the distribution of

null hypothesis wITPC values. This measure represents the relationship between the distribution

of phase angles and the degree of response error on each trial. Plots are only of selected

electrodes and the average of all electrodes (Grand Average). Time-frequency points with *p*-

values at or above 0.05 were set to 1.

Figure 5.

Time-frequency plots showing the relationship between oscillatory power and the standard

mixed model parameter standard deviation (σ) parameter. Standard deviation parameters were

estimated by fitting the standard mixed model to each participants' set of response errors (Figure

2B and 2C). Dark gray contour denotes significant pixel-based corrected effects at $p < 0.05$.

Scatterplots are bootstrap Spearman's rho (r_s) correlation coefficients of mean power across each

statistically significant region outlined in black time. Arrows indicate which region on the time-

frequency maps being plotted. Gray lines are bootstrapped least-squares correlation lines fit to

1000 bootstrap samples.

A

Fixation
0.11°
H
742/783/
825/867 ms

Target
1°
8.3 ms

Blank
41.7 ms

Mask
1.65°
8.3 ms

Blank
500 ms

Response
1°

B

Target   -   Response   =   Error