To See, Not to See, or to See Poorly: Perceptual Quality and Guess Rate as a Function of

Electroencephalography (EEG) Brain Activity in an Orientation Perception Task

Sarah S. Sheldon[1] and Kyle E. Mathewson[1-2]

[1]University of Alberta, Department of Psychology, Faculty of Science

[2]University of Alberta, Neuroscience and Mental Health Institute, Faculty of Medicine and

Dentistry

Author Note

Sarah S. Sheldon https://orcid.org/0000-0002-4026-8939

Kyle E. Mathewson https://orcid.org/0000-0003-4856-3980

Correspondence concerning this article should be addressed to Sarah S. Sheldon,

Department of Psychology, University of Alberta, Edmonton, Alberta, T6G 2R3, Canada. Email:

ssheldon@ualberta.ca

Abstract

Rhythmic brain activity is known to modulate detection of visual stimuli, but it is unclear if this is due to a change in guess rate, change in precision of the mental representation, or both. Here we determined whether the quality of perception randomly varies across trials or is fixed so that the variability is the same across trials. Then we estimated how perceptual quality and guess rate on an orientation perception task relates to EEG activity. Errors on each trial were quantified as the difference between the target orientation and the orientation reported by participants. Response errors were fitted to two variable precision models and the standard mixed model to determine whether perceptual quality is variable or fixed. Overall, the fixed variability parameter of the standard mixed model was the best fit to the response errors. Then we quantified how participants' guess rate and standard deviation parameters from the model varied as a function of EEG activity.

The power and phase of 2-7 Hz post-target activity were found to vary along with task performance in that greater power and phase coherence were measured in trials with more accurate responses. Analysis of task-related alpha activity yielded no significant results. Guess rate and standard deviation were significantly lower on trials with high 2-3 Hz log power than low and the difference started around 250 ms post-target. These results suggest that the precision of perceptual representation is fixed across trials and strongly modulated by the post-target power in the lower frequency bands.

Keywords: EEG, time-frequency analysis, orientation perception, standard mixed model, variable precision model, visual perception

To See, Not to See, or to See Poorly: Perceptual Quality and Guess Rate as a Function of

Electroencephalography (EEG) Brain Activity in an Orientation Perception Task

Variations in neural activity give rise to observed variations in our visual perception.

(Chaumon & Busch, 2014; Mathewson et al., 2011; Samaha et al., 2020). While there has been a

plethora of research into the brain activity that drives this process, there remains noticeable gaps

in our understanding of how these processes work. One of the reasons for this might be because

investigators have left basic questions about the underlying mechanisms unanswered.

Specifically, and the question the current research will address, does a perceptual representation

always form with the same precision or does the quality depend on the state of the neural

activity?

Traditionally, visual perception has been studied with two-alternative forced-choice (2-

AFC) tasks or similar discrete response paradigms. In these types of paradigms, participants are

required to select one out of two or more possible responses. Sometimes they are asked to choose

the correct stimulus out of an array of different stimuli or to simply report whether they detected

a visual stimulus. Although informative, these paradigms might miss certain aspects of visual

perception that cannot be measured with only two or three discrete categories. For example, the

question of whether the quality of visual perception varies from trial to trial or has a precision

that remains constant for a given level of visibility would be difficult to answer without a way to

measure the variability of a response, something that cannot be done with categorical data (in

regards to the traditional concept of variability; see Kader and Perry (2007) for a discussion on

variability in categorical data). To answer this question about the nature of perceptual processes,

we need a task that can measure visual perception on a continuous scale and a model that can

quantify perceptual responses in a way that will inform our question. With this goal in mind, we

adapted the orientation memory task by Bae and Luck's (2018) into an orientation perception

task. The researchers originally used their task to investigate how well information held in

working memory can be decoded from brain activity. What makes the task useful for the current

study is that it allows participants to give a continuous response when asked to report the

orientation of the target. Performance can then be quantified as the angular difference between

the orientation of the target and the orientation reported by the participants, referred to as

response errors. This continuous measure of response errors can utilize models such as the

standard mixture model introduced by Zhang and Luck (2008) or the variable precision model by

Fougnie and colleagues (2012) to quantify parameters of interest such as guess rate and precision.

By extending this method to orientation perception, we can look at how target detection and

perceptual variability are individually related to electrical brain activity during the task.

To address the question about the type of process underlying visual perception, we

combined our adapted version of the visual orientation task with the standard mixed model and

electroencephalography (EEG). Orientation estimation tasks are common in the visual perception

literature (Fischer & Whitney, 2014) and the application of the standard mixture model to

perception, and orientation perception, in particular, has been previously studied (Bays, 2016;

Samaha et al., 2019). However, to our knowledge, this is a novel approach to quantify the effects

of EEG brain activity on visual orientation perception using the standard mixed model to

quantify performance. As a result, the purpose of the study was two-fold. First, we asked whether

the standard mixed model is a good choice for quantifying orientation perception task

performance by comparing the fits of other appropriate working memory models to the data. All

the working memory models we tested made the same assumption that the distribution of errors

could be separated in a uniform distribution representing guesses and a normal distribution of

seen or remembered targets. The main difference between models was how the variability of the

normal distribution got defined. This means that regardless of the model, there would be a guess

rate and standard deviation parameter (the standard deviation parameter could be defined by one

value or two depending on the model). Therefore, our second purpose was to test the relationship

between EEG activity and the model parameters guess rate and standard deviation. We

hypothesized that alpha activity prior to the target onset would be related to whether the target

was later perceived or not which would be reflected as a modulation of the guess rate parameter.

We also hypothesized that the precision of perceptual representations was a fixed ranged (*i.e.*,

based on the same distribution across trials) and that it would be related to post-target activity in

the lower frequency ranges (4-7 Hz) which would be reflected as modulation of the standard

deviation parameter. To address these questions and test our hypotheses, we modified an

orientation memory task from Bae and Luck's (2018) so that it probed participants' perception of

the target's orientation rather than their ability to remember it. We then recorded EEG activity as

participants performed the adapted task so we can see how brain activity varies with their

perceptual performance.

Materials and Methods

Participants

Twenty-eight participants from the University of Alberta community participated in the

study (age range = 17-35 years). Two participants were not included in the analysis due to

excessive movement artifacts (more than 25% of trials rejected due to artifacts). Another two

participants were excluded from the analysis due to having extreme outlying performance on the

task (see Behavioral Analysis in the Results section for more details). Participants were all right-

handed and had normal or corrected normal vision and no history of neurological problems. All

participants gave informed written consent, were either compensated at a rate of $10/hr or given

research credit for their time. The study adhered to the tenets of the Declaration of Helsinki and

was approved by the Internal Ethics Board at the University of Alberta.

Orientation Perception Task

Participants were seated 57 cm away from a 1920 x 1080 pixel[2] ViewPixx/EEG LCD

monitor (VPixx Technologies, Quebec, Canada) with a refresh rate of 120 Hz, simulating a CRT

display with LED backlight rastering. The rastering, along with 8-bit digital TTL output triggers

yoked to the onset and value of the top left pixel, allowed for submillisecond accuracy in pixel

illumination times, which were confirmed with a photocell prior to the experiment. Stimuli were

presented using a Windows 7 PC running MATLAB R2012b with the Psychophysics toolbox

(Version 3; Brainard, 1997; Pelli, 1997). The code running the task was a modified version of the

ColorWorkingMemoryExperiment.m code from MemToolbox (Suchow et al., 2013;

memtoolbox.org). The modified version of color working memory experiment can be found

here: https://github.com/APPLabUofA/OrientTask_paper/tree/master/OrientationTask. Video

output was sent to the ViewPixx/EEG with an Asus Striker GTX760 (Fremont, CA) graphics

processing unit.

Each trial began with a white fixation dot presented at the center of the monitor for 742,

783, 825, or 867 ms (target stimulus onset asynchrony; tSOA) after which the target appeared for

8.33 ms (one monitor refresh). The target was in the shape of a needle and was pointing toward

one of 24 predefined evenly spaced directions so that all the orientations covered 360 degrees.

The direction of the target was randomly selected on each trial. Of all the trials, 20% were

randomly chosen not to have a target. All aspects of the target-present and target-absent trials

were identical except that for the target-absent condition a blank interval replaced target

presentation. A backward mask lasting for 8.33 ms with a constant 41.7 ms target-mask SOA

(mSOA) appeared centrally. The mask was created by overlaying the target orientated in all 24

directions which created a star shape seen in Figure 1A. Following the mask offset, a 500 ms

blank interval period occurred.

After the blank interval, a response screen appeared with the needle in the center of the

screen. Using the computer mouse, participants were asked to rotate the needle so that it was

pointed in the same direction as the previous target. If participants detected a target but could not

remember its orientation, they were asked to guess the orientation of the target. Participants

could provide their response at their own pace. No feedback was given to participants. The next

trial began immediately after a needle's orientation was selected. See Figure 1A for a summary

of the task sequence and the stimulus dimensions.

Participants completed seven blocks consisting of 48 trials each, along with 20 practice

trials at the beginning of the experiment. Participants could rest at their own pace every 48 trials.

Extensive written and verbal instructions were presented to participants prior to the practice trials.

Instructions thoroughly explained and demonstrated each component that would compose a

single trial.

Before the orientation perception task, participants performed a staircased target detection

task that had the same parameters as the orientation perception task except that participants only

reported whether they saw the target or not using the keyboard. The target color was a gray

determined by a scalar value passed to the functions in Psychtoolbox. In the staircased target

detection task, the target color value could range from the background color (making it not

visible; corresponding value of 256) to black (making it the most visible; corresponding value of

0). This target gray value was adjusted throughout the task based on a 1-up/2-down staircasing

procedure targeting a 0.6 target detection rate for each individual (García-Pérez, 1998; Kingdom

& Prins, 2016). The staircased task consisted of three blocks of 32 trials. The target gray value

was determined for each participant by taking the average target gray value across the last two

blocks of trials. These final average values ranged from 70 to 112 and were used as the target

gray value in the orientation perception task.

The MATLAB code for the a staircased target detection task and the orientation

perception task are available at https://osf.io/cw7ux/ and

https://github.com/APPLabUofA/OrientTask_paper.

EEG Recording

During the experiment, EEG data was recorded from each participant with a Brain-Amp

32-channel amplifier (BrainVision) using gelled low-impedance electrodes (actiCAP passive

electrodes kept below 5 kΩ). Inter-electrode impedances were measured at the start of each

experiment. All electrodes were arranged in the same 10-20 positions (Fp2, Fp1, F4, F3, F8, F7,

FC2, FC1, FC6, FC5, C4, C3, CP2, CP1, CP6 CP5, P4, P3 P6, P5, P8, P7, PO4, PO3, O2, O1, Fz,

FCz, Cz, Pz, and Oz). In addition to the 31 EEG sensors, a ground electrode was used, positioned

at AFz. Two reference electrodes and the vertical and horizontal bipolar EOG were recorded

from passive Ag/AgCl easycap disk electrodes affixed on the mastoids, above and below the left

eye, and 1 cm lateral from the outer canthus of each eye. The bipolar channels were recorded

using the AUX ports of the Brain-Amp amplifier. SuperVisc electrolyte gel and mild abrasion

with a blunted syringe tip were used to lower impedances. Gel was applied and inter-electrode

impedances were lowered to less than 5 kΩ for all electrode sites. EEG data was recorded online

referenced to an electrode attached to the left mastoid. Offline, the data were re-referenced to the

arithmetically derived average of the left and right mastoid electrodes.

Data were digitized at 1000 Hz with a resolution of 24 bits. Data were filtered with an

online bandpass with cutoffs of 0.1 Hz and 250 Hz. The experiment was run in a dimly lit, sound

and radio frequency-attenuated chamber from Electromedical Instruments, with copper mesh

covering the window. The only electrical devices in the chamber were an amplifier, speakers,

keyboard, mouse, and monitor. The monitor ran on DC power from outside the chamber, the

keyboard and mouse were plugged into USB outside the chamber, and the speakers and amplifier

were both powered from outside the chamber, and nothing was plugged into the internal power

outlets. Any devices transmitting or receiving radio waves (e.g., cell phones) were removed from

the chamber for the duration of the experiment.

EEG Preprocessing

All analyses were completed using Matlab R2018b with the EEGLAB 13.6.5b (Delorme

& Makeig, 2004) and CircStat (Berens, 2009) toolboxes, as well as custom scripts. After the data

had been re-referenced offline, the bandpass FIR filter from EEGLAB was applied with lower

and upper cut-offs of 0.1 Hz and 50 Hz. Data was segmented into 3000 ms epochs aligned to

target onset (-1500 ms pre-target onset to 1500 ms post-target onset). The average voltage in the

200 ms baseline prior to the target was subtracted on each trial for every electrode, and trials

with absolute voltage fluctuations on any channel greater than 1000 μV were discarded. Eye

movements were then corrected with a regression-based procedure developed by Gratton, Coles,

and Donchin (1983). After a second baseline subtraction with 200 ms pre-target, trials with

remaining absolute voltage fluctuations on any channel greater than 500 μV were removed from

further analysis. Data was then subjected to visual inspection and manual rejection of trials

contaminated by artifacts. On average, 3% of trials were rejected during visual inspection. Other

than the two participants mentioned earlier, none of the remaining participants had more than

20% of trials rejected in this manner.

Data Analyses

Data analysis was performed using MATLAB R2018b (The MathWorks Inc, Natick, MA,

USA) and EEGLAB 13.6.5b (Delorme & Makeig, 2004). All statistical analyses were conducted

using MATLAB R2018b. Red-white-blue colormaps were created using the redblue.m function

by Auton (2009) found here: https://www.mathworks.com/matlabcentral/fileexchange/25536-

red-blue-colormap. The MATLAB code for data analysis is available at the GitHub repository

https://github.com/APPLabUofA/OrientTask_paper and the raw data files are available at

https://osf.io/cw7ux/.

Behavioral Data

Response errors on each trial were calculated by subtracting the orientation of the

response stimulus, as reported by the participant, from the orientation of the target stimulus (see

Figure 1B).

Comparing Model Fits. In addition to the standard mixture model proposed by Zhang

and Luck (2008), the working memory literature has several other models similar to the standard

mixture model but makes different assumptions about some of the parameters. Some of the

working memory models are not appropriate for the current visual orientation perception task

such as those that have an additional Von Mises distributions to account for "swapping" errors or

errors where the participant report a distractor item rather than the target (Bays et al., 2009). On

the other hand, the variable precision models were ones that could be appropriate for the current

study. The original idea behind to model was that the precision of memory varies from trial-to-

trial rather than being fixed as it is in the standard mixture model. This is done by Fougnie and

colleagues (2012) by having the standard deviation parameter be distributed according to a

higher-order distribution, we chose a Gaussian distribution in this case. The variable precision

model proposed by van den Berg and colleagues (2012) has a precision (*i.e.*, the inverse of

variance) parameter drawn from a gamma distribution. Neither paper presented clear justification

for choosing one distribution over another, especially when the set size is always one, so we

tested both distributions. We determined which model better fit the response errors using the

model comparison routine in the MemToolbox (Suchow et al., 2013). We included the standard

mixture model with the bias parameter in addition to the two variable precision models since it

has been shown to fit the participants' data well. The goodness-of-fit measures used were the log

likelihood and the Bayesian information criterion (BIC).

Standard Mixture Model. After determining the standard mixture model proposed by

Zhang and Luck (2008) was the best fit to the current data set, the model was fit to each

participant's response errors using the maximum likelihood estimation routine in the

MemToolbox (Suchow et al., 2013). According to the standard mixture model, response

deviations from the actual target orientation reflect a mixture of trials where the target's

orientation was detected and trials where participants did not detect the target so guessed

randomly. Therefore, the distribution of response errors consists of a mixture of a von Mises

distribution (representing the trials where the target's orientation was detected) and a uniform

distribution (random guesses (g)). Parameter sigma (σ) is the standard deviation of the von Mises

distribution, which represents the width of the response error distribution of trials that the target's

orientation was detected. Parameter g is the height of the uniform distribution representing the

guessing probability. A third parameter, mu (μ), which is the mean of the von Mises distribution

and represents systematic bias of the response error distribution was included in the standard

mixture model of two participants because the Bayesian information criterion (BIC), calculated

with the model comparison functions provided by MemToolbox (Suchow et al., 2013), indicated

that the three-parameter standard mixture model provided better fits for those two participants

(see Table 1). Although the three-parameter model was used for all analysis of those two

participants, the systematic bias was much smaller than the spacing between adjacent target

orientations (spacing was 15° whereas the two participants' μ was -4.2° and 2.7°) indicating that

those two participants had a slight clockwise and counterclockwise bias, respectively. The two-

parameter standard mixture model was used for all analysis of the remaining 22 participants

because it provided a better fit according to the BIC.

Time-Frequency Analyses

To calculate the phase angle and power for each trial, we used Morlet wavelet transform

of single trials using the newtimef() function of EEGLAB. A Morlet wavelet is a tapered sinusoid,

constructed by modulating a sine wave with a Gaussian envelope. Wavelet transformation was

created with 1.027 Hz linear steps and cycles starting from 2 cycles at 2 Hz to 12 cycles at

40 Hz. The output of this function was a matrix of complex values. The abs() function from

MATLAB was used to get the instantaneous amplitude and the instantaneous phase angle of each

trial was calculated using the angle() function from MATLAB.

EEG power data was converted to Z scores by applying a single-trial normalization

procedure to the data from each participant at each electrode and frequency separately. This was

done because it helps disentangle background from task-related dynamics, allows for comparison

across different frequency bands and electrodes, and facilitates group-level analysis (Cohen,

2014). Each trial's entire epoch (-700 ms to 800 ms relative to target onset) was used for the

baseline during normalization because it has been shown to be robust to the effects of noisy trials

(Grandchamp & Delorme, 2011). The downside to using the entire epoch for the baseline

normalization is that sustained changes throughout the trial period become difficult to detect

(Cohen, 2014). Ultimately, we considered this an acceptable trade-off for being able to compare

effects in power and behavioral measures across participants without having to make as many

assumptions about frequency bands and time-windows.

Another possible limitation of using the entire epoch for the baseline during

normalization is that changes in pre-target power may be obscured by post-target activity.

However, comparing effects across participants using raw EEG power is difficult due to

individual differences caused by factors independent of the experimental manipulations (*e.g.*,

skull thickness) (Cohen, 2014). Therefore, when possible, the logarithmically transformed power

was used. EEG power data was logarithmically transformed by applying a single-trial log10

transformation procedure to the data from each participant at each electrode and frequency

separately. When the raw log transformed power is used, it is referred to as log power. EEG

power that was converted to z-scores is called baseline normalized power.

Accurate vs Guess Trials

To test for significant differences in brain activity on trials where participants had small

response errors compared to large response errors, we separated each participant's trials based on

the sigma value from the fits of their individual response errors to the standard mixed model.

This was done by defining each participant's trials with response errors between -0.75σ and

+0.75σ as "accurate," and trials with response errors less than -1.5σ and greater than +1.5σ as

"guesses" (Figure 2A). The logic is that trials). Trials where the participant clearly perceives the

target are likely trials with a response error less than the participant's overall response standard

deviation, and trials the participant has little to no perception of the target are likely trials with a

response error greater than just the participant's response standard deviation. It should be noted

that there are various reasons for participants to be accurate when they are guessing or have a

large response error when they accurately perceived the target. However, such events are thought

to be rare enough, or at least not systematic, that they will not unduly affect the overall

distribution.

The main reason we separated trials into "accurate" or "guess" was to see if the standard

mixed model parameters could be used to categorize trials in a meaningful way. Though this

method resulted in excluding some trials, namely ones that fell between the cutoffs for accurate

and guess, the outcome provided insight into how brain activity differs between different levels

of objective perceptual performance.

ERP Analyses. To remove the activity elicited by the mask without removing activity

resulting from the interaction of mask and target, the catch trials (mask-only) average was

subtracted from the orientation detection trials (target-plus-mask) average. ERP data was

submitted to a repeated measures, two-tailed permutation test based on the *tmax* statistic (Blair &

Karniski, 1993) using the mxt_perm1() function from the Mass Univariate ERP Toolbox (Groppe

et al., 2011). The time windows of interest were the P1 (80-140 ms), N1 (140-200 ms), P2 (200-

255 ms), N2 (255-360 ms), and P3 (360-500 ms) components. The ERP component time

windows were selected based on previous literature (Koivisto & Revonsuo, 2003, 2010). All 31

brain electrodes were included in the test. 100,000 random within-participant permutations were

used to estimate the distribution of the null hypothesis and the familywise alpha (α) was set to

0.05. Based on this estimate, critical *t*-scores of +/-3.68 (df = 23) were derived. Any t-scores that

exceeded the critical *t*-score were considered statistically significant.

Baseline Normalized EEG Power Analysis. To analyze differences in EEG baseline

normalized power between guess and accurate trials, nonparametric permutation testing with a

pixel-based multiple-comparison correction procedure (Cohen, 2014) was used to analyze

differences in EEG band power between guess and accurate trials. The pixel-based multiple-

comparison correction method involves creating one distribution of the largest positive pixel

value and another distribution of the largest negative pixel value from each iteration of the

permutation testing. After all iterations, the statistical threshold is defined as the value

corresponding to the 2.5^{th} percentile of the smallest values and the value corresponding to the

97.5^{th} percentile of the largest values which are the thresholds corresponding to an α of 0.05. Any

pixel that has a value exceeding the upper or lower value is considered significant. The pixel-

based method corrects for multiple comparisons by creating two distributions based on map-level

information instead of pixel-level information. In other words, this method results in two

distributions of the most extreme null-hypothesis test statistical values across all pixels rather

than calculating null-hypothesis distributions for each pixel (see Cohen (2014) for further details

about pixel-based multiple-comparison correction method). All analysis using nonparametric

permutation testing with pixel-based multiple-comparison correction performed 10,000 iterations

per test. To obtain more stable estimates from permutation testing, we ran a "meta-permutation

test" by repeating the pixel-level permutation procedure 10 times and then averaging the results

(Cohen, 2014). It needs to be pointed out that a "significant effect" determined by pixel-based

permutation testing should not be considered a precise estimate in the temporal and frequency

domains. Although pixel-based permutation testing is more stringent than cluster-based

permutation tests (Cohen, 2014), caution should still be used when interpreting "significant"

differences, especially if the temporal and frequency range of each pixel is relatively small.

EEG Phase Analysis. To determine whether the mean phase values significantly differ

between accurate and guess trials, we used the circular Watson–Williams (W-W) test which was

calculated using the PhaseOpposition.m function by VanRullen (2016). We chose the parametric

circular W-W test because it has shown to be equivalent to the non-parametric phase opposition

sum (POS) measures under most conditions, and performed better in situations where either the

relative trial number or the ERP amplitude differed between the two trial groups (VanRullen,

2016). The statistical significance of the W-W test across participants was determined by

combining the individual-level p-value time-frequency map at each electrode across participants

using Stouffer's method (Stouffer et al., 1949; VanRullen, 2016), which transforms individual p-

values into z-scores, combines them across participants, and converts the resulting z-score to a

combined probability. P-values were then corrected for multiple comparisons across time points

and frequencies at each electrode using the false discovery rate (FDR) procedure described in

Benjamini and Yekutieli (2001). Effects that satisfied a 5% FDR criterion were considered

significant.

Single-Trial EEG Activity and Response Errors

To test the correlation between time-frequency log power and degree of response error,

Spearman's rho (r_s) correlation coefficients were calculated using a nonparametric permutation

testing approach with the pixel-based multiple-comparison correction procedure described above.

The null-hypothesis distribution was created by shuffling power values and response errors on

each trial with respect to each other. This provided a data-driven test of the null hypothesis that

there is no consistent relationship between degree of response error and EEG power.

To look at whether task performance is related to oscillatory phase, and if yes, at what

frequency, we used the weighted inter-trial phase clustering (wITPC) (Cohen, 2014; Cohen &

Voytek, 2013). The logic behind the inter-trial phase coherence (ITPC) is that a systematic

relation between EEG phase and behavioral outcome should result in a higher-than-chance ITPC

in each of the trial subgroups. However, if the phase of the EEG signal is randomized and

unpredictable, the distribution of phases at a given time period should follow a uniform

distribution over all trials. The problem with ITPC is that it assumes EEG phase is relevant to

experimental measures only when phase values are similar across trials (van Diepen & Mazaheri,

2018). Unlike ITPC, wITPC is sensitive to modulations of phase values even if those phases are

randomly distributed across trials as would be expected if response errors (which differs from

trial to trial) were modulated by oscillatory phase (Cohen, 2014; Cohen & Voytek, 2013).

363   The wITPC was computed for each participant as the resultant vector length, or ITPC, of

phase angles across trials once the length of each vector has been weighted by a variable of

interest (in this case, each trial's phase vector is weighted by the degree of response error on that

trial; see Figure 4A for example of computation) (Cohen, 2014; Cohen & Voytek, 2013). For

statistical testing, a null-hypothesis distribution was created by shuffling the phase values

relative to trial response error 10,000 times (see Figure 4A middle and bottom left). The wITPCz

was calculated as the wITPC standardized relative to the null-hypothesis distribution, providing a

z-value corresponding to the probability of finding the observed response error–phase

modulation by chance, given the measured data. As was done for the parametric circular W-W

test, statistical significance of the wITPCz across participants was evaluated by combining the

individual-level p-value, calculated from the z-values, time-frequency map at each electrode

across participants using Stouffer's method (Stouffer et al., 1949; VanRullen, 2016). P-values

were then corrected for multiple comparisons across time points and frequencies at each

electrode using the false discovery rate (FDR) procedure described in Benjamini and Yekutieli

(2001). Effects that satisfied a 5% FDR criterion were considered significant.

We chose to use the phase opposition measure and the wITPCz even though they are both

quantifying phase coherence because they provide slightly different but complementary

information about the effects of phase. The phase opposition measure provides insight into

whether there is an overall consistent difference in mean phase values between trials separated

by model parameter defined categories (*i.e*., "accurate" and "guess"). On the other hand, the

weighted single-trial phase modulation metric (*i.e*., wITPCz) provides information about

response error-specific modulations of phase values irrespective of the model. In other words, the

circular W-W test reflects differences between the mean phase of the binned trials while wITPCz

reflects differences in phase as it relates to the continuous measure of response errors. Also, the

wITPCz does not rely on phase values being consistent over trials, they only need to be

consistently related to response errors (Cohen & Cavanagh, 2011). This is an important

distinction when trying to determine how much of the phase modulation is an artifact of the

stimulus-evoked activity and how much is related to the difference in task performance.

Relationship Between Log Power and Standard Mixed Model Parameters

To determine how mixed model parameters standard deviation and guess rate varied as

power varied, a median split of trials according to raw power at each time and frequency point

was done for each participant at each electrode separately. EEG power from the Morlet Wavelet

transformation was logarithmically transformed by applying a single-trial log10 transformation

then trials were split by whether they were above or below the median power at each time point

and frequency. This was done separately for each participant at each electrode. The standard

mixture model was then fit to each set of response errors on the high power and low power trials

to get model parameter values. This meant that every time-frequency point had a standard

deviation and guess rate from trials with high power and low power. The "high power" and "low

power" parameters were then averaged across a frequency band (2-3 Hz, 4-7 Hz, 8-14 Hz, 15-29

Hz or 30-40 Hz) and then tested statistically with the same procedure as the ERP analysis except

each time point was tested rather than averaging across an ERP component time windows and

the alpha level was set to 0.01 to control for the familywise error rate (Bonferroni corrected

alpha level $\alpha_{corr} = 0.05/5$ to account for the five frequency bands). Figure 3 gives a visual

overview of each step in this analysis.

<div align="center">Results</div>

Comparing Model Fits

We tested the best fitting model using the goodness-of-fit measures log likelihood and the

Bayesian information criterion (BIC). The results are presented in Figure 4. Overall, the BIC

indicates the standard mixture model fits the data better than the variable precision models and

the standard mixture model with a bias parameter. The log likelihood indicates the standard

mixture model with a bias parameter is better than the standard mixture model and the variable

precision models. The differences between the BIC and log likelihood metrics can be attributed

to the BIC having a penalty for model complexity whereas the log likelihood does not control for

those factors.

Accurate vs Guess Trials

Two participants were excluded before further analysis because one had a guess rate more

than three *IQR*s from the median and the other had a guess rate of 3.9e-15 indicating that the

staircasing procedure did not work properly for this individual. These participants were not

included in further analysis. Figure 2B shows the fit of the standard mixed model (or standard

mixed model with a bias parameter for the participants mentioned in the Methods section) to

each participant's response errors as well as the average fit of response errors across participants.

The remaining 24 participants had a mean guess rate of 0.19 ($SD = 0.18$) and mean standard

deviation (σ) parameter of 11.1 ($SD = 2.5$). The boxplots in Figure 2C summarizes the

distributions.

ERP Analysis

The ERPs from accurate and guess trials (Figure 3A) showed no statistical difference for

the first 200 ms following target onset across all electrodes. A divergence in the waveforms can

be seen in the P2 (200-300 ms) component for the frontal, central, and centroparietal electrodes

(Figure 3B left) in that the voltage of the guess trial ERPs was much attenuated compared to the

accurate trial ERPs. On the other hand, the voltage of the N2 (255-360 ms) component was more

negative in the guess trials than accurate trials (Figure 3A) and this difference was only

significant in the right frontocentral, central, centroparietal and parietal electrodes (Figure 3B

middle). Finally, the P3 (360-500 ms) component from the guess trials had a similar attenuation

as was seen in the P2 component (Figure 3A), but the distribution was more posterior with

significant effects seen in the right central, centroparietal and parietal electrodes as well as

bilateral parietooccipital and occipital electrodes (Figure 3B right).

EEG Power Analysis

Pixel-based permutation test indicated significant differences between accurate and guess

trials within the 2-4 Hz frequency range which was observed to start around 310 ms post-target

onset in P8 with a duration of about 70 ms and 350 ms post-target onset in P7 with a duration of

around 120 ms (Figure 3C).

Overall, there was a trend for increased 4-7 Hz power in accurate trials compared to

guess trials, especially in left frontal and right parietal areas, though this difference was not

significant. This lack of significance might be due to too much variability in when the 4-7 Hz

power changed during the trial. It is also possible that 4-7 Hz activity reflects a perceptual

process that could occur in both accurate and guess trials, though more often or to a greater

degree in one type of trial compared to the other.

Finally, there were a brief period (20-30 ms) where guess trials had significantly more

baseline normalized power than accurate trials at around 2 Hz right before target onset in FC2

(around -50 ms; Figure 3C) and Cz (around -70 ms; not shown). However, because of the

wavelet parameters used and the timing being around a large evoked response, the timing of the

difference is likely smeared backward so that the effect probably occurred after the target had

been presented (Brüers & VanRullen, 2017; Herrmann et al., 2014; VanRullen, 2011). It should

be noted that no other analysis yielded a significant effect immediately before or after target

onset suggesting that these results are false positives, or the other analyses lacks the power to

detect the effect. The very short duration of an effect at such a low frequency suggests the former

is more likely. If it is the latter, the timing suggests it might have something to do with the mask

onset (*e.g*., anticipation of the mask stimuli) rather than the target. However, the current study

was not designed to investigate the masking stimulus making it difficult to determine the truth

behind the observed effect.

EEG Phase Analysis

Similar to the baseline normalized power results, significant differences in mean phase

between accurate and guess trials were found in the 2-7 Hz frequency ranges following target

onset (Figure 3D). Central and bilateral frontal, frontocentral, and central electrodes show

significant phase differences in 4-7 Hz starting between 150-200 ms and terminating before 350

ms post-target (not shown). The duration of this effect varies so that the more central electrodes

tended to have longer durations than those placed more laterally on the head. This effect was

more prevalent in the right hemisphere in the early period.

The most lateral parietal and centroparietal electrodes on the left side of the head show

significant phase differences within 2-3 Hz after the response screen onset. On the other hand, Pz,

P3 (Figure 3D), P4, and PO3 electrodes had no significant effects in phase while P6 and P8 had

brief periods of significant phase differences starting a little after 300 ms until 500 ms post-target

in the 2-3 Hz frequency range. P7 showed a similar significant difference in mean phase at 2 Hz

starting around 225 ms and continuing until more than 200 ms after the response screen had been

presented (about 770 ms post-target; Figure 3D). The occipital electrodes had significant phase

effects primarily in the 3-5 Hz range starting a little after 100 ms on the left (not shown) and 150

ms on the right but were brief time periods until 200 ms post-target which then had significant

differences lasting for about 200 ms (Figure 3D). PO4 electrode showed a similar difference, but

the effect was less continuous and had larger p-values (*i.e.*, smaller phase difference between

accurate and guess trials) than the occipital electrodes; however, PO4 also had significant

differences within 2-3 Hz frequency between 400 and 500 ms (Figure 3D).

It should be noted that the time window of significant phase opposition overlaps with the

significant differences in ERP amplitudes. ERP amplitudes have been shown to have a "masking"

effect on phase opposition measures resulting in a decrease in their statistical power (VanRullen,

2016) while at the same time the stimulus-evoked activity results in temporal distortion of

oscillatory activity towards earlier latencies (Brüers & VanRullen, 2017). While the circular W-

W test is relatively robust against the detrimental effect of ERPs, it does not negate their

influence entirely (VanRullen, 2016). Therefore, it is important to be aware that the significant

effects of phase are affected by the stimulus evoked activity and the different ERP amplitudes

between accurate and guess trials.

Single-Trial EEG Activity and Response Errors

To see whether response errors were related to trial-by-trial changes in pre-target alpha

power, the Spearman's rho correlations were calculated between log power and degree of

response error on each trial. Our results did not allow us to reject the null hypothesis. Follow-up

analysis on the entire time-frequency space also yielded no significant results (not shown).

To examine the relationship between response errors and the distribution of phase values,

we used the wITPCz. As can be seen in Figure 4B, phase was primarily modulated by degree of

response errors in the 2-3 Hz frequency range in the posterior electrodes starting at around 200

ms post-target and lasting until response screen onset. PO3 was similar except the effect started

at around 250 ms (not shown), but PO4 showed an effect starting at 350 ms and lasting until

almost 100 ms after the response screen onset (not shown). Parietal electrodes show a significant

effect in the low frequency bands starting at around 200 ms until about 800 ms post-target

(Figure 4B, top and bottom rows). Centroparietal and central electrodes had a significant

relationship between phase at 2-3 Hz and response errors at around 300-400 ms post-target and

lasted until after response screen onset. All frontocentral electrodes had response errors

significantly related to 2-3 Hz phase starting between 200 ms and 300 ms post-target and lasting

until shortly after response screen onset. Out of all the frontal electrodes, only Fz and F3 had a

significant 2-7 Hz phase relationship to response errors (not shown).

Relationship Between Log Power and Standard Mixed Model Parameters

A repeated measure, two-tailed permutation test indicated that significant differences in

parameter values between high and low power trials were only within the 2-3 Hz frequency band

following target onset. Figure 6A shows the electrodes that had significant differences in guess

rate and Figure 6B shows the electrodes that significant differences in the standard deviation (σ)

parameter. No other electrodes had significant parameter value differences. Most of the effects

are seen in the occipital and parietal electrodes except for the left frontal electrode which had a

significant difference in guess rate at the later time points than observed in the other electrodes.

The left parietal and occipital electrodes showing significant effects in standard deviation were

not the same electrodes showing significant effects in guess rate. Interestingly, both the guess

rate and standard deviation were higher in trials with low 2-3 Hz log power than high. No

significant differences were seen prior to 250 ms post-target onset and did not occur later than

100 ms before the response screen appeared. It should be noted that, unlike the accurate vs. guess

analysis, these results are based on the log power so the lack of pre-target effects cannot be an

artifact of the normalization process.

526   **Discussion**

There were two goals for the present study. First, we asked what working memory model

best fit to continuous response measures on an orientation perception task? The second goal of

the study was to ask what might the relationship be between the parameters of the best-fitting

model and EEG activity?

Comparing Model Fits

Since perceiving the visual stimuli usually precedes remembering those same stimuli,

some of the assumptions built into the visual working memory models are applicable to visual

perception. Namely, the assumption that there are a set of targets that are remembered and a set

of targets that are not remembered and they can be represented by two distributions which would

translate to the current task as a set of targets that are seen and another set of targets not seen. For

this reason, the first question addressed was what working memory model and their associated

assumption could be extended to the current study's orientation perception task? Based on

goodness-of-fit metrics, we found that the standard mixture model fit the current data set better

than the models that assumed a varying distribution for the standard deviation parameter. It is

important to point out that this does not mean there is no variability in performance over the

course of an experiment. It has been noted by many authors that performance changes as

participants get better at the task or when they get tired towards the end of the experiment.

Rather, these results suggest that a fixed standard deviation fits better to orientation perception

task performance than one that varies from trial to trial according to some distribution. This is

partially in line with the work by Bays (2016) who also noted the success of a standard mixture

model (called it a threshold model in the paper) though it is also shown that the population

encoding model is a better fit though it has limited application to the current data since a model

based on spiking activity does not translate well to a model based on oscillatory activity since

they represent different synaptic mechanisms (Logothetis et al., 2001). It would be interesting to

see if a model based on oscillatory activity within and between different visual regions better

accounts for response errors.

EEG Activity, Model Parameters and Perceptual Behavior

The second goal was to try to answer the question of how brain activity modulates

perceptual representation as it is quantified by the guess rate and standard deviation parameters.

Specifically, we wanted to see if brain activity prior to the target onset in the 8-14 Hz frequency

range (alpha band) modulated whether a target is perceived (*i.e.*, guess rate); and, we tested

whether the quality of target's perceptual representation (*i.e.*, standard deviation parameter) was

related to post-target brain activity in the 4-7 Hz frequency range (theta band). To this end, trials

were categorized as accurate or guesses based on model parameter values. The EEG activity on

accurate and guess trials were then compared across participants. Then, a complementary

approach was used that did a median split of trials based on EEG power. The model was fit to the

high power and low power trials and the resulting parameter values were compared across time

within different frequency bands. Despite our original hypothesis that alpha activity (8-14 Hz)

would be related to guess rate or the standard deviation parameter, the evidence did not support

this idea. Instead, the model-based and power-based approaches both found that 2-7 Hz activity

after target onset modulated perceptual representation as it was quantified by the guess rate and

standard deviation parameters.

Accurate vs Guess Trials: ERP Activity

Significant differences in ERP waveforms of the accurate and guess trials started at

around 200 ms post-target onset in the anterior locations on the head (Figure 3B) which

corresponds to the P2 component (Di Russo et al., 2019; Key et al., 2005; Potts & Tucker, 2001).

The greater amplitude of the P2 for accurate guess trails fits with the P2 being related to the

salience of stimuli (Potts & Tucker, 2001) or stimulus recognition (Harel et al., 2016).

We also observed a more negative N2 for guess trials, that seems to go against current literature

which suggests that the onset of visual consciousness can be marked by a greater negativity in

the N2 time range due to an overlapping posterior negative component called the visual

awareness negativity (VAN) (Förster et al., 2020; Koivisto & Revonsuo, 2003, 2010). However,

the separation of trials based on response performance rather than awareness (according to

Koivisto and Revonsuo (2010), the VAN can only be reliably detected by comparing aware and

unaware conditions), would explain why we would not expect greater negativity in the N2 time

range for accurate ERPs compared to guess ERPs. Furthermore, the distribution of the observed

N2 component (mostly right central electrodes) does not match the typical posterior VAN

distribution (Förster et al., 2020). In fact, the N2 distribution points towards the presence of the

much larger P3 component. The most likely explanation is that the attenuated guess trial ERPs

(compared to accurate trials) results in a smaller or later positive voltage from the much larger P3

overlapping the negative deflection at the N2 time period (Förster et al., 2020; Koivisto &

Revonsuo, 2010) so that the N2 appears more negative in guess trials when it is actually due to

the P3 being smaller. The voltage increase of accurate compared to guess ERPs during the P3

time window follow previous findings on attention and visual perception (Key et al., 2005; Salti

et al., 2012). In the current study the P3 is likely related to processes such as stimulus

classification and saliency evaluation (Doradzińska et al., 2020; Förster et al., 2020; Helfrich &

Knight, 2019; Salti et al., 2012) and that these processes are overwhelmingly more relevant to

accurate orientation perception than processes for conscious perception as reflected by the

relatively smaller N2 ERP. Furthermore, the difference in ERPs between accurate trials and guess

trials during the N2 time period (see middle topographic plot of Figure 3B) indicates that the

processes reflected by the N2 likely occur in both types of trials though to differing degrees.

Instead, it appears that the main difference between accurate and guess trials is the extent that the

target can be classified or evaluated by the brain regardless of consciously perceiving the target.

Accurate vs Guess Trials: EEG Activity

601   We had originally hypothesized activity within the alpha frequency band (8-14 Hz range),

especially prior to target onset, mediating task performance. However, the results did not support

this hypothesis. It is possible a null effect could be attributed to using both pre- and post-target

activity in the pixel-based multiple correction procedure. Relative to post-target activity, pre-

stimulus power would be very small since the only thing for participants to do was fixate on a

white circle. As a result, differences in pre-target activity might not exceed significance

thresholds when those thresholds were based on the background spectra of the combined pre-

and post-target activity. However, we tested this possibility by analyzing power in the 8-14 Hz

frequency range during just the pre-target time period and found no significant differences

between accurate and guess trials (data not shown). Therefore, it is unlikely that a lack of

significant difference in pre-target 8-14 Hz range is due to an exceedingly high statistical

threshold set by the post-target activity.

Another possibility is that the short intertrial interval (ITI) between participants' response

and fixation onset starting the next trial meant alpha desynchronization carried over to the next

trial so that it masked any true effects of spontaneous alpha. While this is a possibility, alpha

activity following the end of a trial is also affected by evaluating their response, task load, and

motivational state (Compton et al., 2011, 2014, 2017). While the short ITI is problematic for

alpha analysis in some regards, it is beneficial in other ways such as discouraging participants

from strategizing or reflecting back on the previous trial.

Similarly, the interstimulus interval (ISI) between fixation onset and target onset is

possibly too short to let event-related desynchronization (ERD) of alpha to recover, thus masking

the effects of spontaneous alpha during the pre-target period. Although potentially problematic,

this should not adversely affect results because the ERD should be equal across all conditions

since fixation is the same on all trials. Therefore, the activity related to the target perception is

equally affected so relative differences can still be detected if they are present.

The most likely reason for not finding a pre-target effect of 8-14 Hz activity is that there

was no information for the participant to use before a target was presented, so there would be no

attention-related changes or task-based preparation prior to the target. This means that the only

activity present during fixation are the spontaneous fluctuations of normal neural activity. Recent

work by Samaha et al. (2020) propose that spontaneous alpha activity modulates neural activity

in a non-specific manner, thus neither facilitating nor inhibiting perceptual activity. The net effect

would be no change to performance which is line with what the model proposed by Samaha and

colleagues (2020) would predict.

The absence of significant effects in post-target 8-14 Hz (alpha band) activity suggests

that alpha activity is not related to the processes responsible for differences between accurate and

guess trials. The fact that post-target 8-14 Hz activity was also unrelated to response errors on a

trial-by-trial basis suggests that 8-14 Hz activity is not related to perception of a target's

orientation. This is consistent with Bae and Luck (2018) who found that alpha activity did not

encode a target's orientation when the target's spatial location was controlled for. An absence of

a relationship between 8-14 Hz power and performance measures has been noted in other studies

though they were not able to rule out changes in detection bias which can be ruled out in the

current study since guess rate was not related to alpha activity (Benwell et al., 2017; Keitel et al.,

2018). However, awareness bias might be a possibility assuming orientation perception can

proceed in the absence of conscious awareness which seems likely based on previous research

(Benwell et al., 2017; Doradzińska et al., 2020; Koenig & Ro, 2019). In sum, it is likely that 8-14

Hz (alpha band) activity is more related to a global process that is not dependent on conscious

awareness and would not change across trials such as feature-independent stimulus processing.

The most significant differences between accurate and guess trials were in the 2-4 Hz

frequency ranges, particularly in the parietal and parietooccipital electrodes during the latter half

of the post-stimulus epoch (Karakaş, 2020). A significant increase of delta and low theta power

in accurate compared to guess trials could be seen at around 300 ms though their differences in

mean phase, especially for theta, started earlier (200 ms post-target). Although there was a trend

in our data for increased theta power in accurate trials compared to guess trials, this difference

was not reliable.

The significant differences in theta and delta seem to correspond to changes in the N2 and

P3 ERP components observed in the time domain. Many studies have shown that delta and theta

are the primary contributors to the formation of the N2 and P3 (Harmony, 2013; Harper et al.,

2014; Karakaş et al., 2000). It has been proposed that activity in the 2-3 Hz range contributes

continuous positivity throughout the ERP response while theta activity corresponds to the

polarity change as the negative deflection during the N2 shifts to the positive deflection during

the P3 (Harper et al., 2014). This fits with the observed EEG activity of accurate and guess trials

(Figure 3C and 3D). Furthermore, the interplay between theta and delta activity would explain

the indistinct N2 but large P3 waveforms (Karakaş et al., 2000).

The increase in baseline normalized 4-7 Hz power during accurate trials relative to guess

trials has the earliest onset in left frontal and right parietal electrodes. Prolonged theta in right

posterior electrodes is accompanied by significant differences in mean (low) theta phase between

accurate and guess trials. Theta has been interpreted as being correlated with selective attentional

processing (Başar et al., 2001; Karakaş, 2020; Karakaş et al., 2000) and an increase in theta

synchronization and power has been associated with successful memory encoding (Klimesch,

1999) and right hemispheric theta is greater than left when encoding visuospatial information

(Sauseng et al., 2004). Although the task in this study was not a memory task, encoding

information is still a viable way of looking at how the brain transforms the visual sensory

information into perceptual representations. In fact, theta has been shown to be sensitive to target

and non-target stimuli regardless of memory load (Palomäki et al., 2012). Therefore, it is

probable that theta activity might be related to the "encoding" of the target's orientation or just

the detectability of target itself. Both functions could occur in accurate and guess trials, though

more often or to a greater degree in accurate trials compared to guess trials. This would explain

why 4-7 Hz activity showed a for trend increased power in accurate trials compared to guess

trials though this difference was usually not significant.

680   In contrast to the 4-7 Hz activity, activity within 2-3 Hz showed a more spatially diffuse

increase in power in accurate vs guess trials with maxima over the lateral parietal areas. Delta

phase differed significantly between accurate and guess trials over the occipital and lateral

parietal areas with the most robust phase differences at around 350 ms post-target. Several

experiments that showed increases in delta activity during the performance of different mental

tasks and conflict-monitoring paradigms have led investigators to associate delta with

modulation of networks that should be inactive to accomplish the task (Harmony, 2013; Rawls et

al., 2020). Investigators have also noted a close association between delta activity and the P3

ERP waveform (Harper et al., 2014; Rawls et al., 2020; Schürmann et al., 1995, 2001). If the P3

reflects post-perceptual processes, then it is likely 2-3 Hz activity is also involved. Whether that

role is as an inhibitory mechanism or not remains unknown. Overall, when considering the time

and frequency at which reliable differences were observed between accurate and guess trials, it

seems likely that this post-target activity reflects differences in the level of perceptual encoding

of the target orientation and the subsequent precision of responses on the task.

Single-Trial EEG Activity and Response Errors

We were surprised to find no significant correlation between response error and

oscillatory power in any of the measured frequencies, but this could also be attributed to the lack

of sensitivity of the analysis method. Since failed to find support for our hypothesis regarding

alpha oscillations, we chose to take a conservative approach when analyzing the rest of the time-

frequency space. Using a less conservative method might help answer the question but that

would increase the risk of finding false positives which could be more harmful than accepting

false negatives.

Interestingly, we did find a relationship between response error and phase values in the 2-

7 Hz frequencies similar to the significant differences in mean phase of accurate trials compared

to guess trials. In fact, phase modulation by response errors is even more pronounced than the

differences in phase between accurate and guess trials. This strongly suggests 2-7 Hz phase

activity plays an important role in the amount of participants' response error on a trial-by-trial

bases. Delta (2-3 Hz) and theta (4-7 Hz) frequencies are usually associated with working

memory and cognitive control, but these results imply that they have an important role in visual

perceptual processes as well.

Relationship Between Log Power and Standard Mixed Model Parameters

There were significant differences in the guess rate and standard deviation parameter

values from trials with high compared to low log power at 2-3 Hz during the post-target time

period. The trend was for trials with low power to have higher guess rate and standard deviation

parameter values than trials with high power. These results are in line with the accurate vs guess

analysis performed on baseline normalized power data. Specifically, trials categorized as

"accurate" had more power in the 2-3 Hz frequency range than trials considered "guesses."

Although a lack of pre-target effects were not hypothesized, it does fit with the baseline sensory

excitability model propose by Samaha and colleagues (2020). Interestingly, Fp1 electrode

showed significant difference in guess rate between high and low power trials prior to the onset

of the response screen. Accumulation of evidence has led investigators to associate frontal delta

(2-3 Hz) activity with top-down control and response inhibition (De Vries et al., 2018; Harmony,

2013; Helfrich et al., 2017; Rawls et al., 2020). Considering that the significant effects are almost

entirely after those in the parietal and occipital electrodes and are only for guess rate, it is likely

top-down control has to do with maintaining the perceptual representation or inhibiting

distracting information during the delay period. In this way, an increase in 2-3 Hz brain activity

would help maintain a target's "perceived" state rather than the target becoming part of the

"unseen" distribution that the guess rate represents.

Conclusion and Future Directions

Overall, we have shown that the standard mixed model can be extended to a visual

perception task and that doing so provides important insight into how brain activity shapes visual

perception. Our results point towards a perceptual representation that has a fixed precision

meaning that while response errors vary with the level of neural activity after stimulus onset,

their variability is according to the same distribution across trials. Whether this is true across

different levels of target visibility is not known but would be an interesting question for future

research.

It is important to note that our results are not suggesting pre-target EEG activity has no

effect on visual perception. There is a lot of evidence to contrary. It has been proposed that pre-

target alpha activity alters participants' confidence rather than performance on visual perception

tasks (Samaha et al., 2017, 2020), a distinction the current study was not designed to test. It is

possible the analysis methods were not sensitive enough to pick up changes in pre-target activity

though there were no significant effects in a follow-up analysis analyzing just the pre-target time

period making a lack of statistical power unlikely. The most likely explanation for an absence of

pre-target effects is that participants did not have information about an upcoming trial so there

was no reason to prepare prior to the target. A follow-up study that directly manipulates attention

or detectability of the target and measures participants' confidence along with the objective

measures of response error, might be able to better test how differences in brain activity prior to

target onset affect visual perception and subsequent task performance.

Acknowledgements

This research was supported by start-up funds from the Faculty of Science and an

NSERC Discovery grant (#RES0024267) awarded to Kyle Elliot Mathewson. The authors would

like you thank Erron Jacob Meneses, Jenny Le, and David Lam for assistance in data collection,

and Daniel Robles and Jonathan Kuziek for analysis feedback.

Conflict of Interest

The author declares that there is no conflict of interest that could be perceived as

prejudicing the impartiality of the research reported.

Author Contributions

KEM and SSS conceived the experiment. SSS collected and analyzed the data. KEM

facilitated the results interpretation. SSS wrote the first draft of the manuscript. SSS and KEM

edited the manuscript.

Data Accessibility

The data that support the findings of this study are openly available through an Open

Science Framework repository at https://doi.org/10.17605/OSF.IO/CW7UX. Scripts to reproduce

all reported analyses and figures are available through a Github repository at

https://github.com/APPLabUofA/OrientTask_paper.

References

Auton, A. (2009). *Red Blue Colormap*. MATLAB Central File Exchange.

https://www.mathworks.com/matlabcentral/fileexchange/25536-red-blue-colormap

Bae, G.-Y., & Luck, S. J. (2018). Dissociable decoding of spatial attention and working memory

from EEG oscillations and sustained potentials. *Journal of Neuroscience*, *38*(2), 409–422.

https://doi.org/10.1523/JNEUROSCI.2860-17.2017

Başar, E., Başar-Eroglu, C., Karakaş, S., & Schürmann, M. (2001). Gamma, alpha, delta, and

theta oscillations govern cognitive processes. *International Journal of Psychophysiology*,

39(2–3), 241–248. https://doi.org/10.1016/S0167-8760(00)00145-8

Bays, P. M. (2016). A signature of neural coding at human perceptual limits. *Journal of Vision*,

16(11), Article 4. https://doi.org/10.1167/16.11.4

Bays, P. M., Catalao, R. F. G., & Husain, M. (2009). The precision of visual working memory is

set by allocation of a shared resource. *Journal of Vision*, *9*(10), 7–7.

https://doi.org/10.1167/9.10.7

Benjamini, Y., & Yekutieli, D. (2001). The control of the false discovery rate in multiple testing

under dependency. *Annals of Statistics*, *29*(4), 1165–1188.

https://doi.org/10.1214/aos/1013699998

Benwell, C. S. Y., Tagliabue, C. F., Veniero, D., Cecere, R., Savazzi, S., & Thut, G. (2017).

Prestimulus EEG Power Predicts Conscious Awareness But Not Objective Visual

Performance. *ENeuro*, *4*(6). https://doi.org/10.1523/ENEURO.0182-17.2017

Berens, P. (2009). CircStat: a MATLAB toolbox for circular statistics. *Journal of Statistical*

Software, *31*(10), 679–685. https://doi.org/10.1016/j.amp.2012.05.023

Blair, R. C., & Karniski, W. (1993). An alternative method for significance testing of waveform

difference potentials. *Psychophysiology*, *30*(5), 518–524. https://doi.org/10.1111/j.1469-

8986.1993.tb02075.x

Brainard, D. H. (1997). The Psychophysics Toolbox. *Spatial Vision*, *10*(4), 433–436.

http://www.ncbi.nlm.nih.gov/pubmed/9176952

Brüers, S., & VanRullen, R. (2017). At what latency does the phase of brain oscillations

influence perception? *ENeuro*, *4*(3). https://doi.org/10.1523/ENEURO.0078-17.2017

Chaumon, M., & Busch, N. A. (2014). Prestimulus neural oscillations inhibit visual perception

via modulation of response gain. *Journal of Cognitive Neuroscience*, *26*(11), 2514–2529.

https://doi.org/10.1162/jocn_a_00653

Cohen, M. X. (2014). *Analyzing neural time series data: theory and practice*. MIT Press.

Cohen, M. X., & Cavanagh, J. F. (2011). Single-trial regression elucidates the role of prefrontal

theta oscillations in response conflict. *Frontiers in Psychology*, *2*.

https://doi.org/10.3389/fpsyg.2011.00030

Cohen, M. X., & Voytek, B. (2013). Linking nonlinear neural dynamics to single-trial human

behavior. In M. (Meyer) Z. Pesenson (Ed.), *Multiscale Analysis and Nonlinear Dynamics*

(pp. 217–232). Wiley-VCH Verlag GmbH & Co. KGaA.

https://doi.org/10.1002/9783527671632

Compton, R. J., Arnstein, D., Freedman, G., Dainer-Best, J., & Liss, A. (2011). Cognitive control

in the intertrial interval: Evidence from EEG alpha power. *Psychophysiology*, *48*(5), 583–

590. https://doi.org/10.1111/j.1469-8986.2010.01124.x

Compton, R. J., Bissey, B., & Worby-Selim, S. (2014). Task motivation influences alpha

suppression following errors. *Psychophysiology*, *51*(7), 585–595.

https://doi.org/10.1111/psyp.12212

Compton, R. J., Heaton, E., & Ozer, E. (2017). Intertrial interval duration affects error

monitoring. *Psychophysiology*, *54*(8), 1151–1162. https://doi.org/10.1111/psyp.12877

De Vries, I. E. J., Van Driel, J., Karacaoglu, M., & Olivers, C. N. L. (2018). Priority switches in

visual working memory are supported by frontal delta and posterior alpha interactions.

Cerebral Cortex, *28*(11), 4090–4104. https://doi.org/10.1093/cercor/bhy223

Delorme, A., & Makeig, S. (2004). EEGLAB: An open source toolbox for analysis of single-trial

EEG dynamics including independent component analysis. *Journal of Neuroscience*

Methods, *134*(1), 9–21. https://doi.org/10.1016/J.JNEUMETH.2003.10.009

Di Russo, F., Berchicci, M., Bianco, V., Perri, R. L., Pitzalis, S., Quinzi, F., & Spinelli, D. (2019).

Normative event-related potentials from sensory and cognitive tasks reveal occipital and

frontal activities prior and following visual events. *NeuroImage*, *196*, 173–187.

https://doi.org/10.1016/j.neuroimage.2019.04.033

Doradzińska, Ł., Wójcik, M. J., Paź, M., Nowicka, M. M., Nowicka, A., & Bola, M. (2020).

Unconscious perception of one's own name modulates amplitude of the P3B ERP

component. *Neuropsychologia*, *147*, 107564.

https://doi.org/10.1016/j.neuropsychologia.2020.107564

Fischer, J., & Whitney, D. (2014). Serial dependence in visual perception. *Nature Neuroscience*,

17(5), 738–743. https://doi.org/10.1038/nn.3689

Förster, J., Koivisto, M., & Revonsuo, A. (2020). ERP and MEG correlates of visual

consciousness: The second decade. *Consciousness and Cognition*, *80*, Article 102917.

https://doi.org/10.1016/j.concog.2020.102917

Fougnie, D., Suchow, J. W., & Alvarez, G. A. (2012). Variability in the quality of visual working

memory. *Nature Communications*, *3*, 1229. https://doi.org/10.1038/ncomms2237

García-Pérez, M. A. (1998). Forced-choice staircases with fixed step sizes: asymptotic and small-

sample properties. *Vision Research*, *38*(12), 1861–1881. https://doi.org/10.1016/S0042-

6989(97)00340-4

Grandchamp, R., & Delorme, A. (2011). Single-trial normalization for event-related spectral

decomposition reduces sensitivity to noisy trials. *Frontiers in Psychology*, *2*, 236.

https://doi.org/10.3389/fpsyg.2011.00236

Gratton, G., Coles, M. G. ., & Donchin, E. (1983). A new method for off-line removal of ocular

artifact. *Electroencephalography and Clinical Neurophysiology*, *55*(4), 468–484.

https://doi.org/10.1016/0013-4694(83)90135-9

Groppe, D. M., Urbach, T. P., & Kutas, M. (2011). Mass univariate analysis of event-related

brain potentials/fields I: A critical tutorial review. *Psychophysiology*, *48*(12), 1711–1725.

https://doi.org/10.1111/j.1469-8986.2011.01273.x

Harel, A., Groen, I. I. A., Kravitz, D. J., Deouell, L. Y., & Baker, C. I. (2016). The temporal

dynamics of scene processing: A multifaceted EEG investigation. *ENeuro*, *3*(5), Article

e0139-16.2016. https://doi.org/10.1523/ENEURO.0139-16.2016

Harmony, T. (2013). The functional significance of delta oscillations in cognitive processing.

Frontiers in Integrative Neuroscience, *7*, Article 83.

https://doi.org/10.3389/fnint.2013.00083

Harper, J., Malone, S. M., & Bernat, E. M. (2014). Theta and delta band activity explain N2 and

P3 ERP component activity in a go/no-go task. *Clinical Neurophysiology*, *125*(1), 124–132.

https://doi.org/10.1016/j.clinph.2013.06.025

Helfrich, R. F., Huang, M., Wilson, G., & Knight, R. T. (2017). Prefrontal cortex modulates

posterior alpha oscillations during top-down guided visual perception. *Proceedings of the*

National Academy of Sciences of the United States of America, *114*(35), 9457–9462.

https://doi.org/10.1073/pnas.1705965114

Helfrich, R. F., & Knight, R. T. (2019). Cognitive neurophysiology: Event-related potentials. In

Handbook of Clinical Neurology (Vol. 160, pp. 543–558). Elsevier B.V.

https://doi.org/10.1016/B978-0-444-64032-1.00036-9

Herrmann, C. S., Rach, S., Vosskuhl, J., & Strüber, D. (2014). Time–frequency analysis of event-

related potentials: A brief tutorial. *Brain Topography*, *27*(4), 438–450.

https://doi.org/10.1007/s10548-013-0327-5

Kader, G. D., & Perry, M. (2007). Variability for Categorical Variables. *Journal of Statistics*

Education, *15*(2). https://eric.ed.gov/?id=EJ842705

Karakaş, S. (2020). A review of theta oscillation and its functional correlates. *International*

Journal of Psychophysiology. https://doi.org/10.1016/j.ijpsycho.2020.04.008

Karakaş, S., Erzengin, Ö. U., & Başar, E. (2000). A new strategy involving multiple cognitive

paradigms demonstrates that ERP components are determined by the superposition of

oscillatory responses. *Clinical Neurophysiology*, *111*(10), 1719–1732.

https://doi.org/10.1016/S1388-2457(00)00418-1

Keitel, C., Benwell, C. S. Y., Thut, G., & Gross, J. (2018). No changes in parieto-occipital alpha

during neural phase locking to visual quasi-periodic theta-, alpha-, and beta-band

stimulation. *European Journal of Neuroscience*, 1–15. https://doi.org/10.1111/ejn.13935

Key, A. P. F., Dove, G. O., & Maguire, M. J. (2005). Linking brainwaves to the brain: An ERP

primer. *Developmental Neuropsychology*, *27*(2), 183–215.

https://doi.org/10.1207/s15326942dn2702_1

Kingdom, F. A. A., & Prins, N. (2016). Chapter 5 – Adaptive Methods. In *Psychophysics*

(Second Edi, pp. 119–148). Academic Press. https://doi.org/10.1016/B978-0-12-407156-

8.00005-0

Klimesch, W. (1999). EEG alpha and theta oscillations reflect cognitive and memory

performance: a review and analysis. *Brain Research Reviews*, *29*(2–3), 169–195.

https://doi.org/10.1016/S0165-0173(98)00056-3

Koenig, L., & Ro, T. (2019). Dissociations of conscious and unconscious perception in TMS-

induced blindsight. *Neuropsychologia*, *128*, 215–222.

https://doi.org/10.1016/j.neuropsychologia.2018.03.028

Koivisto, M., & Revonsuo, A. (2003). An ERP study of change detection, change blindness, and

visual awareness. *Psychophysiology*, *40*(3), 423–429. https://doi.org/10.1111/1469-

8986.00044

Koivisto, M., & Revonsuo, A. (2010). Event-related brain potential correlates of visual

awareness. *Neuroscience & Biobehavioral Reviews*, *34*(6), 922–934.

https://doi.org/10.1016/j.neubiorev.2009.12.002

Logothetis, N. K., Pauls, J., Augath, M., Trinath, T., & Oeltermann, A. (2001).

Neurophysiological investigation of the basis of the fMRI signal. *Nature*, *412*(6843), 150–

157. https://doi.org/10.1038/35084005

Mathewson, K. E., Lleras, A., Beck, D. M., Fabiani, M., Ro, T., & Gratton, G. (2011). Pulsed out

of awareness: EEG alpha oscillations represent a pulsed-inhibition of ongoing cortical

processing. *Frontiers in Psychology*, *2*, Article 99. https://doi.org/10.3389/fpsyg.2011.00099

Palomäki, J., Kivikangas, M., Alafuzoff, A., Hakala, T., & Krause, C. M. (2012). Brain

oscillatory 4-35 Hz EEG responses during an n-back task with complex visual stimuli.

Neuroscience Letters, *516*(1), 141–145. https://doi.org/10.1016/j.neulet.2012.03.076

Pelli, D. G. (1997). The VideoToolbox software for visual psychophysics: transforming numbers

into movies. *Spatial Vision*, *10*(4), 437–442. http://www.ncbi.nlm.nih.gov/pubmed/9176953

Potts, G. F., & Tucker, D. M. (2001). Frontal evaluation and posterior representation in target

detection. *Cognitive Brain Research*, *11*(1), 147–156. https://doi.org/10.1016/S0926-

6410(00)00075-6

Rawls, E., Miskovic, V., & Lamm, C. (2020). Delta phase reset predicts conflict-related changes

in P3 amplitude and behavior. *Brain Research*, *1730*, 146662.

https://doi.org/10.1016/j.brainres.2020.146662

Salti, M., Bar-Haim, Y., & Lamy, D. (2012). The P3 component of the ERP reflects conscious

perception, not confidence. *Consciousness and Cognition*, *21*(2), 961–968.

https://doi.org/10.1016/j.concog.2012.01.012

Samaha, J., Iemi, L., Haegens, S., & Busch, N. A. (2020). Spontaneous Brain Oscillations and

Perceptual Decision-Making. *Trends in Cognitive Sciences*, *24*(8), 639–653.

https://doi.org/10.1016/j.tics.2020.05.004

Samaha, J., Iemi, L., & Postle, B. R. (2017). Prestimulus alpha-band power biases visual

discrimination confidence, but not accuracy. *Consciousness and Cognition*, *54*, 47–55.

https://doi.org/10.1016/J.CONCOG.2017.02.005

Samaha, J., Switzky, M., & Postle, B. R. (2019). Confidence boosts serial dependence in

orientation estimation. *Journal of Vision*, *19*(4). https://doi.org/10.1167/19.4.25

Sauseng, P., Klimesch, W., Doppelmayr, M., Hanslmayr, S., Schabus, M., & Gruber, W. R.

(2004). Theta coupling in the human electroencephalogram during a working memory task.

Neuroscience Letters, *354*(2), 123–126. https://doi.org/10.1016/j.neulet.2003.10.002

Schürmann, M., Başar-Eroglu, C., Kolev, V., & Başar, E. (1995). A new metric for analyzing

single-trial event-related potentials (ERPs): application to human visual P300 delta response.

Neuroscience Letters, *197*(3), 167–170. https://doi.org/10.1016/0304-3940(95)11912-G

Schürmann, M., Başar-Eroglu, C., Kolev, V., & Başar, E. (2001). Delta responses and cognitive

processing: Single-trial evaluations of human visual P300. *International Journal of*

Psychophysiology, *39*(2–3), 229–239. https://doi.org/10.1016/S0167-8760(00)00144-6

Stouffer, S. A., Suchman, E. A., Devinney, L. C., Star, S. A., & Williams Jr., R. M. (1949).

Studies in social psychology in World War II: the American soldier. In *Adjustment During*

Army Life (Vol. 1). Princeton University Press. https://psycnet.apa.org/record/1950-00790-

000

Suchow, J. W., Brady, T. F., Fougnie, D., & Alvarez, G. A. (2013). Modeling visual working

memory with the MemToolbox. *Journal of Vision*, *13*(10), 1–8.

https://doi.org/10.1167/13.10.9

Van Den Berg, R., Shin, H., Chou, W. C., George, R., & Ma, W. J. (2012). Variability in

encoding precision accounts for visual short-term memory limitations. *Proceedings of the*

National Academy of Sciences of the United States of America, *109*(22), 8780–8785.

https://doi.org/10.1073/pnas.1117465109

van Diepen, R. M., & Mazaheri, A. (2018). The Caveats of observing Inter-Trial Phase-

Coherence in Cognitive Neuroscience. *Scientific Reports*, *8*(1), 2990.

https://doi.org/10.1038/s41598-018-20423-z

VanRullen, R. (2011). Four common conceptual fallacies in mapping the time course of

recognition. *Frontiers in Psychology*, *2*, Article 365.

https://doi.org/10.3389/fpsyg.2011.00365

VanRullen, R. (2016). How to Evaluate Phase Differences between Trial Groups in Ongoing

Electrophysiological Signals. *Frontiers in Neuroscience*, *10*, 426.

https://doi.org/10.3389/fnins.2016.00426

Zhang, W., & Luck, S. J. (2008). Discrete fixed-resolution representations in visual working

memory. *Nature*, *453*(7192), 233–235. https://doi.org/10.1038/nature06860

954   **Tables**

955   **Table 1**

956   *List of goodness-of-fit measures comparing the standard mixed model to the standard mixed*

957   *model with a bias parameter (μ) for each participant.*

	Log Likelihood			Bayesian Information Criterion (BIC)		
	Standard Model	Model + Bias	Difference	Standard Model	Model + Bias	Difference
Participant 1	-1255.82	-1255.33	-0.49	2526.45	2532.88	-6.44
Participant 2	-1213.93	-1212.78	-1.15	2442.77	2447.92	-5.15
Participant 3	-1095.28	-1094.58	-0.70	2205.45	2211.50	-6.05
Participant 4	-1093.29	-1093.24	-0.05	2201.25	2208.50	-7.24
Participant 5	-1191.23	-1190.92	-0.31	2397.23	2404.01	-6.77
Participant 6	-1113.21	-1111.14	-2.07	2241.28	2244.57	-3.29
Participant 7	-965.54	-964.92	-0.62	1945.90	1952.06	-6.16
Participant 8	-1022.30	-1021.07	-1.23	2059.48	2064.47	-4.99
Participant 9	-1109.18	-1109.12	-0.06	2233.12	2240.39	-7.27
Participant 10	-1422.57	-1422.42	-0.15	2859.94	2867.05	-7.11
Participant 11	-1384.47	-1384.46	-0.00	2783.78	2791.20	-7.42
Participant 12	-1017.58	-1015.52	-2.06	2049.58	2052.67	-3.09
Participant 13	-1498.89	-1498.55	-0.34	3012.59	3019.32	-6.73
Participant 14	-1071.57	-1071.55	-0.02	2157.98	2165.36	-7.38
Participant 15	-1124.71	-1123.84	-0.87	2264.26	2269.94	-5.68
Participant 16	-985.46	-985.17	-0.29	1985.36	1991.99	-6.63
Participant 17	-1149.36	-1149.26	-0.11	2313.61	2320.84	-7.23
Participant 18	-1136.58	-1136.50	-0.09	2287.86	2295.02	-7.17
Participant 19	-1133.03	-1132.91	-0.12	2280.88	2288.04	-7.16
Participant 20	-1276.89	-1275.81	-1.08	2568.44	2573.61	-5.17
Participant 21	-1138.27	-1133.04	-5.23	2291.40	2288.37	3.03[a]
Participant 22	-1003.69	-988.80	-14.89	2022.10	1999.69	22.41[a]
Participant 23	-1011.18	-1010.68	-0.50	2037.17	2043.57	-6.40
Participant 24	-1166.38	-1162.88	-3.49	2347.60	2348.04	-0.44

958   *Note*. Difference is calculated as the goodness-of-fit measure of the standard mixed model with a

959   bias model subtracted from the goodness-of-fit measure of the standard mixed model.

960   [a] Positive difference indicates the standard mixed model with a bias parameter (μ) fits better to

961   the participants' response errors (*i.e.*, has a smaller BIC) than the standard mixed model.

962   **Figure Captions**

Figure 1.

A) Sequence of task events with duration of each screen presentation and sizes of fixation, target,

mask, and response stimuli. Sizes are in degrees of visual angle. **B)** Example of response error

calculation. Response errors are reported in degrees.

Figure 2.

A) Representation of how trials were split into the two categories, Accurate and Guess. This was

done by defining each participant's trials with response errors between -0.75σ and $+0.75\sigma$ as

"accurate" (shaded brown region), and trials with response errors less than -1.5σ and greater than

$+1.5\sigma$ as "guesses" (shaded green regions). Trials that did not fall in either category were

discarded from this analysis between the two trial categories. **B)** Left, model fit of each

participant. Right, solid purple line is the average of participants' model fit. Light purple area

represents SEM. Zoomed in window shows upward shift of the averaged model fit showing a

non-zero guess rate. **C)** Boxplots of parameter values (left) guess rate and (right) standard

deviation estimated from model fits of participants' response errors.

Figure 3.

Illustration of the analysis looking at how the estimated parameter values from the standard

mixed model varied across an epoch when trials were separated by the power within a frequency

band at each time point. First, time-frequency transformation (e.g., Morlet wavelet

transformation) of single-trial data was used to calculate raw power of every time and frequency

point for each trial. Trials were split by median power at each time and frequency point then the

standard mixed model was fit to the trials' response errors. This was done for each participant

and electrode separately. An average of the parameter values (*i.e.,* standard deviation and guess

rate) from the "high power" and "low power" trials were calculated across the five frequency

bands (2-3 Hz, 4-7 Hz, 8-14 Hz, 15-29 Hz or 30-40 Hz) at each time point and then were

compared statistically using a repeated measures, two-tailed permutation test based on the *tmax*

statistic (Blair & Karniski, 1993).

Figure 4.

Goodness-of-fit measures relative to the standard mixture model. Top, the difference in log

likelihood values compared to the standard mixture model with positive values favoring the

standard mixture model. Bottom, the difference in Bayesian Information Criterion (BIC) values

compared to the standard mixture model with negative values favoring the standard mixture

model. Log likelihood favors the standard mixture model with a bias parameter (μ) over the

standard mixture model and both variable precision models. The BIC favors the standard mixture

model over the other three models. Plots are the means and error bars are ±SEM.

1000

Figure 5.

A) ERPs of accurate trials and guess trials aligned to target onset at selected electrodes and

averaged across all electrodes (Grand Average). Light brown and light green shaded areas around

waveforms represent ±SEM of accurate trials and guess trials, respectively. Mask trial EPRs have

been subtracted out of the ERPs to remove mask-related brain activity except for the Grand

Average + Mask ERPs plot. Time period shaded in purple indicates significant difference

between accurate and guess trials 200-255 ms (P2) post-target. Blue shaded time period indicates

significant difference between accurate and guess trials 360-500 ms (P3) post-target. **B)**

Topographies of the voltage distribution difference between accurate and guess trials. The time

periods in each topography are as follows: 200-255 ms (P2), 255-360 ms (N2), and 360-500 ms

(P3) post-target. Stars indicate electrodes with significant differences between accurate and guess

ERPs. Mask trial activity has been subtracted out of the ERPs to remove mask-related brain

activity. **C)** Analysis results from comparing the baseline normalized power of accurate vs guess

trials. Time-frequency plots showing the results of the statistical analysis of the difference in

power between accurate vs guess trials at each time-frequency point at selected electrodes and

the average of all electrodes (Grand Average). Black contour denotes statistically significant

differences (after applying the pixel-based multiple-comparison correction procedure) at $p < .05$.

D) Phase analysis of accurate vs guess trials. Time-frequency plots showing the p-values (after

applying FDR correction) from the statistical analysis testing differences in the mean phase of

accurate trials vs guess trials. Plots are from selected electrodes and the average of all electrodes

(Grand Average). Time-frequency points with p-values above the threshold of FDR correction for

multiple comparisons were set to 1. **E)** Topographies of the p-value distribution averaged across

the 2-3 Hz frequency (top) and 4-7 Hz frequency (bottom) indicating significant differences in

the mean phase of accurate trials vs guess trials. Time-frequency points with p-values above the

threshold of FDR correction for multiple comparisons were set to 1. The time periods in each

topography are the same as used for the ERP analysis: 200-255 ms (P2), 255-360 ms (N2), and

360-500 ms (P3) post-target. Only the time periods and frequency bands that had significant

effects after averaging over the time window are shown.

1029

Figure 6.

A) Example computation of weighted inter-trial phase clustering (wITPC) to relate single-trial

phase to degree of response error. Top left, single-trial prestimulus phase vectors are shown as

black lines and are not clustered across trials due to the randomization of the fixation length,

leading to a low resultant vector length (i.e., low ITPC). Right, histogram of null-hypothesis

vector lengths created by shuffling the mapping of the trial response error values to the trial

phase values. The observed vector length (large arrow) is calculated from the distribution shown

in bottom left panel. Bottom left shows how the length each trial's phase vector is scaled by that

trial's degree response error and a weighted ITPC is computed, reflecting the relationship

between the distribution of phase angles and degree of response error even though the

distribution of the phase angles themselves is uniform (as seen in the top left plot). The wITPCz

value is the wITPC standardized relative to the null-hypothesis distribution seen in the histogram

plot. Example based on figure by Cohen (2014). **B)** Time-frequency plots of analysis relating

single-trial phase activity and response errors. Significant *p*-values indicating that the normalized

distance of the observed wITPC (i.e., wITPCz) is significantly different from the distribution of

null hypothesis wITPC values. This measure represents the relationship between the distribution

of phase angles and the degree of response error on each trial. Plots are only of selected

electrodes and the average of all electrodes (Grand Average). Time-frequency points with *p*-

values at or above .05 were set to 1. **C)** Topographies of the *p*-value distribution (after applying

FDR correction) averaged across the 2-3 Hz frequency indicating that the normalized distance of

the observed wITPC (i.e., wITPCz) is significantly different from the distribution of null

hypothesis wITPC values. This measure represents the relationship between the distribution of

phase angles and the degree of response error on each trial. Time-frequency points with *p*-values

above the threshold of FDR correction for multiple comparisons were set to 1. The time periods

in each topography are the same as used for the ERP analysis: 120-200 ms (N1), 200-255 ms

(P2), 255-360 ms (N2), and 360-500 ms (P3) post-target. Only showing time periods and

frequency bands that had significant effects after averaging over the time window.

1057

Figure 7.

A) Plots of the fluctuations in the guess rate parameter across time in trials with high and low log

power in the 2-3 Hz frequency band. No other frequency band had significant differences in

guess rate between high and low power trials. Only electrodes with significant effects are shown.

Green shaded regions indicate time points where guess rate in high and low power trials

significantly differed. Shaded regions around waveforms are $\pm SEM$. **B)** Plots of the fluctuations

in the standard deviation (σ) parameter across time in trials with high and low log power in the 2-

3 Hz frequency band. No other frequency band had significant differences in standard deviation

between high and low power trials. Only electrodes with significant effects are shown. Green

shaded regions indicate time points where the standard deviation parameter in high and low

power trials significantly differed. Shaded regions around waveforms are $\pm SEM$.

A

Fixation

0.11°

H

742/783/
825/867 ms

Target

1°

8.3 ms

Blank

41.7 ms

Mask

1.65°

8.3 ms

Blank

500 ms

Response

1°

B

Target   -   Response   =   Error

A

ITPC = .1601

wITPCz = 3.6334

Histogram of H_0 wITPC

Count

wITPC

Observed
wITPC = 2.176

B

Fp1

FC2

P7

P5

P8

O1

O2

Grand Average

Freqency (Hz)

Target
Onset

Response
Screen
Onset

Time (ms)

p-value

C

2-3 Hz Frequency

140-200 ms
(N1)

200-255 ms
(P2)

255-360 ms
(N2)

360-500 ms
(P3)

p-value