HIV reverse transcriptase pre-steady-state kinetic analysis of chain terminators and translocation inhibitors reveals interactions between magnesium and nucleotide 3'-OH

By

Christopher R. Dilmore[a] and Jeffrey J. DeStefano[a,b,#]

[a]Cell Biology and Molecular Genetics, 3130 Bioscience Research Building, University of Maryland, College Park, Maryland, USA

[b]Maryland Pathogen Research Institute, College Park, Maryland, USA

Running Head: 3' hydroxyl is pivotal in physiological Mg^{2+}

[#]Address correspondence to Jeffrey J. DeStefano, jdestefa@umd.edu

Abstract

Deoxythymidine triphosphate analogs with various 3' sugar groups (-OH (dTTP), -H, $-N_3$, $-NH_2$, -F, $-O-CH_3$, no group (2′,3′-didehydro-2′,3′-dideoxythymidine triphosphate (d4TTP)), and those retaining the 3'-OH but with 4' additions (4'-C-methyl, 4'-C-ethyl) or sugar ring modifications (D-carba dTTP) were evaluated using pre-steady-state kinetics in low (0.5 mM) and high (6 mM) Mg^{2+} with HIV reverse transcriptase (RT). Analogs showed diminished incorporation rates (k) compared to dTTP ranging from about 2-fold (3'-H, $-N_3$, and d4TTP with high Mg^{2+}) to >10-fold (3'-NH_2 and 3'-F with low Mg^{2+}), while 3'-$O-CH_3$ dTTP incorporated much slower than other analogs. Illustrating the importance of interactions between Mg^{2+} and the 3'-OH, k using 5 μM dTTP and 0.5 mM Mg^{2+} was only modestly slower (1.6-fold) than with 6 mM Mg^{2+}, while analogs with 3' alterations incorporated 2.8-5.1-fold slower in 0.5 mM Mg^{2+}. In contrast, 4'-C-methyl and D-carb dTTP, which retain the 3'-OH, were not significantly affected by Mg^{2+}. Consistent with these results, analogs with 3' modifications were better inhibitors in 6 mM *vs*. 0.5 mM Mg^{2+}. Equilibrium dissociation constant (K_d) and k_{pol} determinations for dTTP and analogs lacking a 3'-OH indicated that low Mg^{2+} caused a several-fold greater reduction in k_{pol} with the analogs but had little effect on K_d, results consistent with a role for 3'-OH/Mg^{2+} interactions in catalysis rather than nucleotide binding. Overall, results emphasize the importance previously unreported interactions between Mg^{2+} and the 3'-OH of the incoming nucleotide and suggest inhibitors with 3'-OH groups may have advantages in the low free Mg^{2+} in physiological settings.

Introduction

Nucleoside Reverse Transcriptase Inhibitors (NRTIs) are a hallmark of Antiretroviral Therapy (ART). All NRTIs currently approved for ART are "chain terminators" and either lack a hydroxyl group at the 3' ribose position or have a substituted sugar or other group replacing ribose. In all cases there is no group present for addition of the next base by reverse transcriptase (RT) (reviewed in: (1-3)). A second class of emerging NRTIs referred to as "translocation inhibitors" retain the 3'-OH but contain modifications at other positions that dramatically slow the addition of the next base (4-7). The first approved HIV drug, azidothymidine (also referred to as Zidovudine, AZT, or ZDV), is essentially dTTP with the 3'-OH replaced by an azido group. Dideoxy drugs replace the 3'-OH with H (including dideoxyinosine (also referred to as Didanosine or ddI) and dideoxycytidine (a formerly approved HIV drug also referred to as Zalcitabine or ddC). Other drugs remove both the 3'-OH and a hydrogen atom from the 3' position by addition of a carbon double bond to the ribose ring (2',3'-Didehydro-3'-deoxythimidine (also referred to as Stavudine or d4T) and the guanosine analog Abacavir (ABC)). Finally, some approved drugs substantially modify or remove the ribose sugar including Lamivudine (3TC), Emtricitabine (FTC), and Tenofovir (TDF). Despite the success of drugs lacking a 3'-OH, for some translocation inhibitors this group is pivotal in promoting the phosphorylation of the prodrug in cells (8).

Others have examined the effect of different groups at the 3' position of thymidine and adenosine analogs for inhibition of HIV RT (9, 10). For thymidine, inhibition assays using steady-state kinetics and measuring incorporation on poly(rA)-oligo(dT$_{10}$) indicated that substitutions of fluorine (F), amino (NH$_2$), azido (N$_3$), and hydrogen (H) at the 3' position were

all effective inhibitors of dTTP incorporation with modest overall differences, with 3'-F and 3'-

N_3 being the most and least effective, respectively (9). Despite this, only the 3'-N_3 derivative

(marketed as AZT) has been an effective HIV drug. Drug effectiveness is complicated by

several factors including stability, uptake, cellular phosphorylation, target and off-target

utilization, and resistance profiles among others. Even small modification could potentially

affect any of these steps. The F and H 3' substitution (as well as several other changes at other

positions) on dATP were also tested for anti-HIV activity and cell toxicity (10). Both analogs

had lower cytotoxicity than AZT, however, they also had much lower potency. Still, the

selectivity index for 3'-H (ddATP) was only about 4.5-fold lower than AZT suggesting it could

be an effective drug.

Previous results indicated that the level of Mg^{2+} in *in vitro* reactions can affect the fidelity

of HIV RT, efficiency of reverse transcription, and the potency of both NRTIs and

Nonnucleoside Reverse Transcriptase Inhibitors (NNRTIs). Fidelity increased significantly (11,

12) in reactions performed in low Mg^{2+} (0.25-0.5 mM) that more closely mimics cellular free

Mg^{2+} conditions (13-17), in comparison to the high Mg^{2+} (~6 mM) typically used in *in vitro*

reactions with RT. The efficiency of reverse transcription, as judged by the length of products

produced in primer extension reactions, was greater with physiological Mg^{2+} concentrations (18).

The effect on NRTIs and NNRTIs was opposite, with the former being less effective in low Mg^{2+}

and the latter being more effective (18, 19).

In this report we used pre-steady-state kinetic analysis at a fixed nucleotide concentration

(5 µM) that approximated cellular dNTP concentrations (20-22), and high (6 mM) or low (0.5

mM) Mg^{2+} concentrations to examine the incorporation rate of dTTP and several dTTP analogs

with substitutions at the 3' position. This included 3'-OH (dTTP), -N_3 (AZT), -NH_2, -F, -O-CH_3,

-H (ddTTP), and d4T (no group at this position). Three translocation type dTTP analog

inhibitors (4'-C-methyl, 4'-C-ethyl, and D-carba dTTP) that retained the 3'-OH were also tested.

Determinations of k_{cat} and K_d were also performed for some analogs under pre-steady-state

conditions. Overall, the results illustrate a potential benefit for NRTI drugs with 3'-OH groups,

especially under physiological Mg^{2+} concentrations, while also uncovering a previously

unreported direct or indirect interaction between the 3'-OH of incoming nucleotides and Mg^{2+}.

Results

Substitution of the 3'-OH differentially affects the incorporation rate of dTTP analogs. All

the analogs tested, including chain terminators and translocation inhibitors, were incorporated

more slowly than dTTP in both 0.5 and 6 mM Mg^{2+}, in pre-steady-state assays with 5 μM

nucleotide (Table 1, see Fig. 1 for an example of an analysis). This amount was chosen as it

approximately mimics the level of nucleotides in dividing T cells (20-22). The incorporation

rate ordered from highest to lowest in 6 mM Mg^{2+} was: 3'-OH > (-N$_3$, no group (d4TTP), -H) >

(4'-C-methyl, 4'-C-ethyl, -NH$_2$) > (-F, D-carba) >> -O-CH$_3$. In 0.5 mM Mg^{2+} the order was: 3'-

OH > (4'-C-methyl) > (-N$_3$, no group, -H, 4'-C-ethyl, D-carba, -NH$_2$) > -F >> -O-CH$_3$. The 3'-O-

CH$_3$ analog was by far the slowest and no data could be obtained using pre-steady-state

conditions. However, steady-state analysis showed that the analog could be incorporated by RT

(Fig. 2). Among the chain terminating analogs, at 6 mM Mg^{2+}, an -N$_3$, -H, or no group at the 3'

position were most tolerated. The 3'-F was modestly slower than 3'-NH$_2$ and the incorporation

rate of 3'-O-CH$_3$ was negligible in comparison (see above). Note that these results agree with a

previous steady-state analysis of some of these analogs indicating that they were efficiently

incorporated by HIV RT ((9) and Introduction). However, the relative potency of the various

inhibitors did not agree with our pre-steady-state analysis.

Incorporation rates for translocation inhibitors were less dependent on the Mg^{2+} conditions.

Rates for the 4'-C-methyl and 4'-C-ethyl derivatives were similar in 6 mM Mg^{2+} and both were

about 3-fold faster than D-carba dTTP. In 0.5 mM Mg^{2+}, only the incorporation rate for 4'-C-

ethyl dTTP declined significantly.

The effect of lower Mg^{2+} on incorporation is largely independent of the type of 3'

substitution. All analogs that did not have a 3'-OH were incorporated more slowly in 0.5 than 6

mM Mg^{2+}. The magnitudes of the decreases were in a range between 2.8-5.1-fold (Table 1),

with 3'-F and 3'-H (ddTTP) being the lowest and highest, respectively. However, the difference

between the various 3' analogs was small and, in most cases, not significant. This indicates that

the low Mg^{2+} effect results more from the loss of the 3'-OH than a property of the substitution.

The 4'-C-methyl and D-carba translocation inhibitor analogs were not strongly affected by lower

Mg^{2+} while dTTP showed just a 1.6-fold decrease in incorporation rate with low Mg^{2+} (Table 1).

Curiously, 4'-C-ethyl dTTP behaved more like the chain terminators and was incorporated about

3 times more slowly in 0.5 mM Mg^{2+}. It is notable, however, that despite possessing a 3'-OH, 4'-

C-ethyl dTTP actually behaves like a chain terminator as unlike 4'-C-methyl and D-carb dTTP,

nucleotide cannot be added to this moiety ((23) and see Discussion).

Analogs without a 3'-OH demonstrate a more profound loss of catalytic efficiency in lower

Mg^{2+}. The observed incorporation rate of a nucleotide or analog at a particular concentration is

dependent on its affinity for enzyme as well as the overall rate of the catalytic steps. Several

reports have shown that 3'-OH substitutions may alter both the binding affinity (i.e. equilibrium

dissociation constant (K_d)) for HIV RT as well as the maximum incorporation rate (k_{pol}) (29-33).

We choose three dNTPs/analogs, dTTP, AZTTP, and d4TTP, to evaluate more thoroughly and

determine both pre-steady-state K_d and k_{pol} values at 0.5 and 6 mM Mg^{2+}. Although it would

have been more complete to test all 10 analogs, a combination of limiting amounts of material

and time made it more feasible to test a smaller representative set. The decision to focus on

AZTTP and d4TTP was made because there is extensive literature on these compounds, and they

are approved HIV drugs. The K_d values for all 3 compounds were similar with d4TTP showing a

modestly higher K_d than the others at both Mg^{2+} concentrations (Fig. 3 and Table 2). The level

of Mg^{2+} did not strongly affect the K_d value for any of the nucleotides. Values for K_d at 6 mM

obtained hear were comparable to previous results with high Mg^{2+} (33). As expected from the

results in Table 1, changes in k_{pol} were more pronounced, especially for the nucleotides lacking a

3'-OH (AZTTP and d4TTP). The decline in catalytic efficiency (k_{pol}/K_d), with 0.5 vs. 6 mM

Mg^{2+} was also more pronounced for these analogs. For example, AZTTP was incorporated only

1.5-fold less efficiently than dTTP in 6 mM Mg^{2+}, but 7.2-fold less efficiently in 0.5 mM Mg^{2+}

(Table 2). The data again illustrates that the lack of a 3'-OH has a more pronounced negative

affect on incorporation in low, more physiological Mg^{2+}, and a decrease in incorporation rate (k)

as opposed to changes in affinity (K_d) for the analogs is the predominant factor.

Inhibition assays on a heteropolymeric DNA template are generally consistent with the

kinetic analysis. To evaluate the effectiveness of the various analogs at terminating DNA

synthesis, a primer extension assay was performed on a 100-nt DNA template primed with a 20-

nt 5'-[32]P labeled DNA primer (Fig. 4). Assays contained 5 µM dCTP, dGTP, dATP, and dTTP.

Analogs were included at 3 µM when present. Consistent with 6/0.5 mM Mg^{2+} ratios shown in

Table 1, all chain terminators and 4'-C-ethyl dTTP were less effective (as judged by a lower level

of paused (terminated) products and an increase in fully extended products) in 0.5 mM Mg^{2+}

compared to 6 mM Mg^{2+}. Among translocation inhibitors, 4'-C-methyl dTTP was essentially

equivalent at both 0.5 and 6 mM Mg^{2+} while D-carba was a weak inhibitor under both

conditions. The 3'-F analog was the weakest inhibitor among the chain terminators and was also

incorporated the slowest in pre-steady-state reactions (Table 1, note that 3'-O-CH_3 produced no

inhibition in this assay (data not shown) and its incorporation was not measurable in pre-steady-

state reactions (see above)). However, it is important to note that the inhibition observed in these

reactions is a function of both incorporation kinetics and binding affinity of the analog for RT vs.

binding affinity for dTTP. Therefore, a direct correlation between the results in Table 1 and

those in the inhibition assay would not necessarily be expected. Finally, results in Table 1 were

performed using a single sequence position on a primer-template while the experiments in Fig. 4

use a long template with positions for thymidine incorporation presented in several sequence

contexts.

Discussion

In this report, several chain-terminating 3'-OH substitutions of dTTP were tested for

incorporation by HIV RT. All the analogs with 3' alterations were incorporated and several (e.g.

AZT, ddTTP, and d4TTP) with only modestly slower rates (~2-fold) than dTTP in 6 mM Mg^{2+}

(Table 1). In contrast, the rate of incorporation compared to dTTP dropped to ~4.5-16-fold

slower (excluding 3'-O-CH_3 which was much slower at both high and low Mg^{2+}) in 0.5 mM

Mg^{2+}. This is consistent with previous experiments showing that NRTIs are better inhibitors

with higher Mg^{2+} concentrations *in vitro* (19), an observation that was also shown for analogs

tested here (Fig. 4). Overall, the results support a previously unreported interaction, either direct

or indirect, between Mg^{2+} and the 3'-OH of incoming nucleotides. Further, the results suggest

that this interaction plays a role in catalysis rather than nucleotide binding affinity (see below).

The importance of the 3'-OH was supported by experiments with translocation type inhibitors

(4'-C-methyl and D-carba dTTP) which retain the 3'-OH and behaved more like dTTP in that

their rate of incorporation was less affected by low Mg^{2+} (Table 1). However, 4'-C-ethyl dTTP

was an exception, showing a 3.1-fold lower rate of incorporation in low Mg^{2+} compared to high.

As was noted, this analog terminates further extension by RT, in contrast to 4'-C-methyl and D-

carba dTTP which delay extension (23). This suggests that the 3'-OH may be in the wrong

position for catalysis and this position may also weaken the proposed interactions of the 3'-OH

with Mg^{2+}. The translocation inhibitor 4'-ethynyl-2-fluoro-2'-deoxyadenosine triphosphate

(EFdATP) is currently in clinical trials. It also binds strongly to HIV RT as the ethynyl group

can access a hydrophobic pocket in RT which may stabilize binding (7, 34). This, and favorable

kinetics lead to EFdATP showing a competitive advantage over the natural dATP nucleotide

(35). EFdATP inhibition is also insensitive to Mg^{2+} concentration (19), and incorporated EFdA

can accept additional nucleotides at a highly reduced rate (35). This suggests that translocation

inhibitors with the 3'-OH positioned for nucleotide addition behave more like natural nucleotides

with respect to Mg^{2+} interactions.

It was interesting the 4'-C-ethyl dTTP was clearly a more effective inhibitor than 4' C-methyl or

D-carba dTTP at 6 mM Mg^{2+} during extension on the long template (Fig. 4). Even at 0.5 mM

Mg^{2+}, where 4'-C-ethyl dTTP was incorporated much more slowly than 4'-C-methyl dTTP and at

a rate similar to D-carba dTTP (Table 1), it still was modestly more effective than 4'-C-methyl

dTTP and much more effective than D-carba dTTP in the template inhibition assay. This may be

due to two factors. First, 4'-C-ethyl dTTP, as stated above, behaves like a chain terminator, and

does not allow the addition of other nucleotides after incorporation (23). Additional nucleotides

can be added with delayed kinetics with 4'-C-methyl and D-carba dTTP (23, 24) and, for these

analogs, extended products in the reactions in Fig. 4 may include DNA chains containing more

than one inhibitor nucleoside. Second, the K_d for 4'-C-ethyl, 4'-C methyl and D-carba dTTP

binding to HIV RT was measured at 6 mM Mg^{2+} in a separate analysis on a different template

(manuscript in preparation). The K_d value for D-carba dTTP was ~4-8-fold higher than the

others. This helps explain why D-carba dTTP is by far the weakest inhibitor among both

translocation and chain terminating inhibitors in the template extension inhibition assay (Fig. 4).

It binds more weakly to RT than the other translocation inhibitors, and this, coupled with a

relatively low incorporation rate (Table 1), weakens inhibition.

Although positioning of the 3'-OH of an incoming nucleotide is coordinated through many

interactions with HIV RT, glutamine 151 (Q151) presumably interacts directly with this group

through a proposed hydrogen bond between the amide oxygen from Q151 and the 3'-OH

hydrogen atom. Mutations in Q151 like Q151N (glutamine to asparagine) that disrupt this

interaction weaken dNTP binding but do not significantly affect k_{pol} (29). This indicates that

Q151 likely plays a key role in stabilizing dNTP binding. Dideoxy dTTP which lacks a 3'-OH

demonstrated weaker binding to HIV RT than dTTP. However, binding relative to dTTP

improves with Q151N which has weakened association with the 3'-OH. This again predicts a

role for Q151 in stabilizing binding of dNTPs through interactions with the 3'-OH, as the

advantage of a 3'-OH for binding is mitigated when Q151 is altered to a non-interacting amino

acid.

High divalent cation concentrations can also improve incorporation of analogs without 3'-OH

groups with other polymerases, similar to what was found here. The Klenow fragment of *E. coli*

polymerase I (KF) discriminates strongly against ddNTPs relative to dNTPs. However, ddTTP

incorporation improves in high Mg^{2+} showing an optimum of 25 mM which is several-fold

greater than the optimal concentration for incorporation of dTTP (~2 mM) (36). Like Q151N for

HIV RT, an E710A (glutamic acid to alanine) KF mutation reduced ddNTP/dNTP

discrimination, consistent with a role for this amino acid in interactions with the 3'-OH. The

authors hypothesize that Mg^{2+} may bridge an interaction between KF E710 and the 3'-OH group

of incoming nucleotides, although more complex explanations where E710 interacts with Mg^{2+}

via repositioning of other active site residues could not be ruled out by the data. It was also not

clear if the Mg^{2+} ion in question was metal ion A and/or B, the putative metal ions involved in

nucleotide catalysis at polymerase active sites, or another, as yet undescribed metal ion. In this

regard, it is interesting that a third metal ion has been proposed to be involved in polymerase

nucleotide catalysis and may play a role in these interactions (37-41). Interestingly, unlike

Q151N, E710A had a larger effect on k_{cat} rather than on binding, leading the authors to postulate

that its interaction with the 3′-OH does not contribute to ground state binding. Instead, the data

was more consistent for E710-Mg^{2+} interactions having a role in the transition state occurring

after nucleotide binding but before catalysis (36). Our work indicates that lowering the

concentration of Mg^{2+} disproportionately depresses the rate of catalysis for all thymidine analogs

lacking a 3'-OH without strongly affecting affinity (based on K_d values) for a limited set (dTTP,

AZTTP, and d4TTP) of tested nucleotides (Fig. 3, and Tables 1 and 2). This is consistent with

the observations with the KF and suggests that Mg^{2+} may interact with the 3'-OH, either directly

or indirectly, in a step after nucleotide binding during the transition state or the chemistry step

(the latter was ruled for the KF by demonstrating a lack of a sulfur elemental effect (36), but we

did not test this possibility).

It was noteworthy that HIV RT was able to incorporate all 9 analogs that were tested. Only 3'-O-

CH$_3$ dTTP showed very poor incorporation. For chain terminating analogs, there was not a clear

consistent chemical property that explained the different rates of incorporation of the analogs.

However, electronegative groups that, unlike a 3'-OH, could not donate a H atom for hydrogen

bonding (e.g. 3'-F and 3'-O-CH$_3$) were incorporated the slowest. Groups with potential for

positive charge or hydrogen bond formation (e.g. 3'-NH$_2$, and 3'-N$_3$), and small groups (e.g. 3'-H

(ddTTP) or d4TTP (no group)) were better substrates, although all were less efficiently

incorporated than dTTP, especially in low Mg^{2+} (Tables 1 and 2). It is possible that 3'-NH$_2$ and

3'-N$_3$ groups may interact with Q151 through charge interactions or H-bonding and this may help

compensate for loss of the 3'-OH. Both ddTTP and d4TTP could not establish these interactions

but the small size of the 3' substitution and non-charge nature could minimize repulsive

interactions. Also, d4TTP is different than the other tested chain terminators as the 2',3'-

didehydro configuration of the sugar moiety alters the structure of the sugar. This makes it more

difficult to directly compare it to the other 3' modified analogs.

In conclusion, our results support a role for interactions between Mg^{2+} and the 3'-OH of the

incoming nucleotide that enhances the rate of catalysis. Interactions between Mg^{2+} and the 3'-

OH terminus of the growing DNA chain and the incoming nucleotide phosphate backbone have

been well established and are part of polymerase dogma. In these instances, divalent cation is

directly interacting with the reacting moieties of nucleotide catalysis. Presumably, the 3'-OH of

the incoming nucleotide is spatially removed from the catalytic center. This suggests that an

interaction with it that enhances catalysis would likely have a stabilizing (e.g. stabilization of a

transition state) rather than chemical role. An example could be that Mg^{2+} effects established

interactions between Q151 and the 3′-OH. This could occur directly or indirectly through

positioning of the nucleotide in a manner that promotes the Q151/3′-OH interaction. Further

experiments, including those with Q151 mutants and mutations at other positions that effect

Mg^{2+} binding, will be required to better understand the mechanism behind these observations.

Materials and Methods

Materials and Methods

Materials. 3'-fluor (F) and 3'-O-methyl (O-CH_3) dTTP were synthesized by AX Molecules Inc.,

San Marcos, CA. 3'-amino (NH_2) dTTP and dideoxy TTP (ddTTP) were from TriLink Bio

Technologies. AZTTP and d4TTP were from Sierra Bioresearch, Tucson, AZ. 4'-C-methyl, 4'-

C-ethyl, and D-carba dTTP (23, 24) were a generous gift from Dr. Stephen Hughes (National

Institutes of Health). All other nucleotides were from Roche Diagnostics Corporation. All

nucleotides had purity levels >95% (as determined by mass spectroscopy and TLC analysis). T4

polynucleotide kinase (PNK) was from New England Biolabs. Radiolabeled compounds were

from PerkinElmer. DNA oligonucleotides were from Integrated DNA Technologies (IDT). G-

25 spin columns were from Harvard Apparatus. All other chemicals were obtained from Fisher

Scientific, VWR, or Sigma.

Preparation of HIV reverse transcriptase (HIV RT). The expression plasmids (pRT66 and

pRT51) for HIV-1 RT (HXB2 sequence) were used to produce HIV RT (25). The enzyme,

which is a non-tagged heterodimer consisting of equal proportions of p66 and p51, was prepared

as described (26).

Preparation of primer-templates for pre-steady-state kinetics and inhibitor analysis on the

100-nt template. For pre-steady-state kinetics, 5'-^{32}P-labeled 20-nt primer (5'-

TTCCCGTCAGCCTAGCTGAG-3′) and 36-nt template (5'-

ATCGTCGTACTCAGCTAGGCTGACGGGAAATTGTAT-3′) were mixed at a ratio of 1:1.25

(primer:template) in a buffer containing 50 mM Tris-HCl (pH 8), 80 mM KCl, and 1 mM DTT.

For inhibitor analysis during primer extension on a long template, 5'-^{32}P-labeled 20-nt primer (5′-

TTGTTGTCTCTTCCCCAAAC-3′) and 100-nt template (5′-

TGGCCTTCCCACAAGGGAAGGCCAGGGAATTTTCTTCAGAGCAGACCAGAGCCAAC

AGCCCCACCAGAAGAGAGCTTCAGGTTTGGGGAAGAGACAACAA-3′) were mixed at a

at a ratio of 1:1.5 (primer:template). Hybrids were formed by heating to 80□C for 5 min then

slow cooling in a PCR machine to 4□C.

Pre-steady-state kinetic parameters for dTTP or dTTP analog incorporation. Experiments

were performed using a QFM-4000 from Bio-Logic Scientific Instruments. All experiments

were carried out at 37□C. For each reaction, 20 µL of solution containing 50 nM (in primer)

primer-template prepared as described above (final concentration 25 nM in reactions) and 50 nM

HIV-1 RT (HXB2) (final concentration 25 nM in reactions) in buffer 1 (50 mM Tris-HCl (pH 8),

80 mM KCl, 1 mM DTT, and 0.5 or 6 mM MgCl$_2$) were loaded into one loop of the apparatus.

A second loop contained 20 µL of a solution with various concentrations of dTTP or specific

dTTP analogs in the same buffer. Concentrations used ranged from 1 to 100 µM (final

concentrations in reaction) for experiments to determine K$_d$ and k_{cat} (depending on the

nucleotide) and were 5 µM for pre-steady-state rate (k) determinations with various analogs. For

reactions with higher nucleotide concentrations, the starting Mg^{2+} concentration in the 0.5 mM

Mg^{2+} reactions was adjusted to keep the free Mg^{2+} concentration at 0.5 mM (19). A third loop

contained 20 μL of a solution containing 0.5 M EDTA (pH 8). Reactions were initiated by

rapidly mixing the solutions in loops 1 and 2, then terminated by mixing with loop 3 at time

points ranging from 5 milliseconds (ms) to 2 seconds (s). Specific time points depended on the

incorporation rate of the individual analogs. The products were mixed with and equal volume of

2X loading buffer (90% formamide, 0.025% each bromophenol blue and xylene cyanol) and

resolved on 20% polyacrylamide-7M urea denaturing PAGE gels as described (27). Gels were

dried, then exposed to phosphorimager screens. Quantification was performed using a Fujifilm

FLA7000 phosphorimager and Fuji ImageQuant software. First order rate constants for

incorporation (k) were determined using SigmaPlot by plotting the concentration of extended

starting material vs. time, and fitting the data to a single exponential equation (28):

$$y = a(1 - e^{-kt}) + C$$

315   where "a" is the amplitude, k is the rate, and "C" is the end point. In some cases, especially for

316   analogs with slower incorporation rates, the data fit better to a simpler equation:

$$y = a(1 - e^{-kt})$$

317   For some analogs, rate constants at different analog concentrations were used to determine the

318   equilibrium dissociation constant and maximum rate of nucleotide addition (K_d and k_{pol},

319   respectively) by plotting k vs. the concentration of nucleotide and fitting to a hyperbolic equation

320   for ligand binding with single site saturation:

$$y = \frac{B_{max}(x)}{K_d + x}$$

where "B_{max}" corresponds to k_{pol}. Standard error values for K_d and k_{pol} were generated by the

program.

Inhibitor analysis during primer extension on a long template. Primer extension reactions

were performed to study the inhibition of extension by thymidine analogs as described

previously (19). Briefly, 15 nM 5′-^{32}P-labeled primer was hybridized with 22.5 nM template at a

ratio of 1:1.5 as described above. Hybrids were preincubated for 3 min at 37°C in 8.5 μL of

buffer 1, all 4 dNTPs and 1 of the analogs (5 μM each and 3 μM, respectively (final

concentrations in reactions)). Extension was initiated by adding 4 μL of HIV RT (final

concentration 100 nM) in buffer 1. After extension for 30 min, the reactions were terminated by

adding 12.5 μL of 2X gel loading buffer and samples were resolved on an 8% denaturing urea

gel and processed as described above.

Acknowledgements

334 Funding: This work was supported by the National Institutes of Health [grant numbers

335 R01AI150480]. The sponsor was not involved in study design; in the collection, analysis and

336 interpretation of data; in the writing of the report; and in the decision to submit the article for

337 publication. We thank Dr. Stephen Hughes (National Institutes of Health) for 4'-C-methyl, 4'-C-

338 ethyl, and D-carba dTTP.

References

1. Sarafianos SG, Marchand B, Das K, Himmel DM, Parniak MA, Hughes SH, Arnold E. 2009. Structure and function of HIV-1 reverse transcriptase: molecular mechanisms of polymerization and inhibition. J. Mol. Biol. 385:693-713.

2. Singh K, Marchand B, Kirby KA, Michailidis E, Sarafianos SG. 2010. Structural Aspects of Drug Resistance and Inhibition of HIV-1 Reverse Transcriptase. Viruses 2:606-638.

3. Holec AD, Mandal S, Prathipati PK, Destache CJ. 2017. Nucleotide Reverse Transcriptase Inhibitors: A Thorough Review, Present Status and Future Perspective as HIV Therapeutics. Curr. HIV Res. 15:411-421.

4. Kodama EI, Kohgo S, Kitano K, Machida H, Gatanaga H, Shigeta S, Matsuoka M, Ohrui H, Mitsuya H. 2001. 4'-Ethynyl nucleoside analogs: potent inhibitors of multidrug-resistant human immunodeficiency virus variants in vitro. Antimicrob. Agents Chemother. 45:1539-46.

5. Ohrui H, Kohgo S, Hayakawa H, Kodama E, Matsuoka M, Nakata T, Mitsuya H. 2006. 2'-Deoxy-4'-C-ethynyl-2-fluoroadenosine: a nucleoside reverse transcriptase inhibitor with highly potent activity against all HIV-1 strains, favorable toxic profiles and stability in plasma. Nucleic acids symposium series (2004):1-2.

6. Kawamoto A, Kodama E, Sarafianos SG, Sakagami Y, Kohgo S, Kitano K, Ashida N, Iwai Y, Hayakawa H, Nakata H, Mitsuya H, Arnold E, Matsuoka M. 2008. 2'-deoxy-4'-C-ethynyl-2-halo-adenosines active against drug-resistant human immunodeficiency virus type 1 variants. Int. J. Biochem. Cell. Biol. 40:2410-20.

7. Michailidis E, Marchand B, Kodama EN, Singh K, Matsuoka M, Kirby KA, Ryan EM, Sawani AM, Nagy E, Ashida N, Mitsuya H, Parniak MA, Sarafianos SG. 2009.

Mechanism of inhibition of HIV-1 reverse transcriptase by 4'-Ethynyl-2-fluoro-2'-

deoxyadenosine triphosphate, a translocation-defective reverse transcriptase inhibitor. J.

Biol. Chem. 284:35681-91.

8.   Siddiqui MA, Hughes SH, Boyer PL, Mitsuya H, Van QN, George C, Sarafinanos SG,

Marquez VE. 2004. A 4'-C-ethynyl-2',3'-dideoxynucleoside analogue highlights the role

of the 3'-OH in anti-HIV active 4'-C-ethynyl-2'-deoxy nucleosides. J. Med. Chem.

47:5041-8.

9.   Cheng YC, Dutschman GE, Bastow KF, Sarngadharan MG, Ting RY. 1987. Human

immunodeficiency virus reverse transcriptase. General properties and its interactions with

nucleoside triphosphate analogs. J. Biol. Chem. 262:2187-9.

10.   Herdewijn P, Pauwels R, Baba M, Balzarini J, De Clercq E. 1987. Synthesis and anti-

HIV activity of various 2'- and 3'-substituted 2',3'-dideoxyadenosines: a structure-activity

analysis. J. Med. Chem. 30:2131-7.

11.   Okano H, Baba M, Hidese R, Iida K, Li T, Kojima K, Takita T, Yanagihara I, Fujiwara S,

Yasukawa K. 2018. Accurate fidelity analysis of the reverse transcriptase by a modified

next-generation sequencing. Enzyme Microb. Technol. 115:81-85.

12.   Achuthan V, Keith BJ, Connolly BA, DeStefano JJ. 2014. Human immunodeficiency

virus reverse transcriptase displays dramatically higher fidelity under physiological

magnesium conditions in vitro. J. Virol. 88:8514-27.

13.   Gout E, Rebeille F, Douce R, Bligny R. 2014. Interplay of Mg^{2+}, ADP, and ATP in the

cytosol and mitochondria: unravelling the role of Mg^{2+} in cell respiration. Proc. Natl.

Acad. Sci. U. S. A. 111:E4560-7.

14.   Murphy E, Freudenrich CC, Levy LA, London RE, Lieberman M. 1989. Monitoring

cytosolic free magnesium in cultured chicken heart cells by use of the fluorescent

indicator Furaptra. Proc. Natl. Acad. Sci. U. S. A. 86:2981-4.

15.   Delva P, Pastori C, Degan M, Montesi G, Lechi A. 1998. Intralymphocyte free

magnesium and plasma triglycerides. Life Sci. 62:2231-40.

16.   Delva P, Pastori C, Degan M, Montesi G, Lechi A. 2004. Catecholamine-induced

regulation in vitro and ex vivo of intralymphocyte ionized magnesium. J. Membr. Biol.

199:163-71.

17.   Gupta RK, Benovic JL, Rose ZB. 1978. The determination of the free magnesium level in

the human red blood cell by 31P NMR. J. Biol. Chem. 253:6172-6.

18.   Goldschmidt V, Didierjean J, Ehresmann B, Ehresmann C, Isel C, Marquet R. 2006.

Mg2+ dependency of HIV-1 reverse transcription, inhibition by nucleoside analogues and

resistance. Nucleic Acids Res. 34:42-52.

19.   Achuthan V, Singh K, DeStefano JJ. 2017. Physiological Mg(2+) conditions significantly

alter the inhibition of HIV-1 and HIV-2 reverse transcriptases by nucleoside and non-

nucleoside inhibitors in vitro. Biochemistry 56:33-46.

20.   Diamond TL, Roshal M, Jamburuthugoda VK, Reynolds HM, Merriam AR, Lee KY,

Balakrishnan M, Bambara RA, Planelles V, Dewhurst S, Kim B. 2004. Macrophage

tropism of HIV-1 depends on efficient cellular dNTP utilization by reverse transcriptase.

J. Biol. Chem. 279:51545-53.

21.   Kennedy EM, Gavegnano C, Nguyen L, Slater R, Lucas A, Fromentin E, Schinazi RF,

Kim B. 2010. Ribonucleoside triphosphates as substrate of human immunodeficiency

virus type 1 reverse transcriptase in human macrophages. J. Biol. Chem. 285:39380-91.

22.   Amie SM, Noble E, Kim B. 2013. Intracellular nucleotide levels and the control of

retroviral infections. Virology 436:247-54.

23.   Boyer PL, Julias JG, Ambrose Z, Siddiqui MA, Marquez VE, Hughes SH. 2007. The

nucleoside analogs 4'C-methyl thymidine and 4'C-ethyl thymidine block DNA synthesis

by wild-type HIV-1 RT and excision proficient NRTI resistant RT variants. J. Mol. Biol.

371:873-82.

24.   Boyer PL, Vu BC, Ambrose Z, Julias JG, Warnecke S, Liao C, Meier C, Marquez VE,

Hughes SH. 2009. The nucleoside analogue D-carba T blocks HIV-1 reverse

transcription. J. Med. Chem. 52:5356-64.

25.   Fletcher RS, Holleschak G, Nagy E, Arion D, Borkow G, Gu Z, Wainberg MA, Parniak

MA. 1996. Single-step purification of recombinant wild-type and mutant HIV-1 reverse

transcriptase. Protein Expr. Purif. 7:27-32.

26.   Hou EW, Prasad R, Beard WA, Wilson SH. 2004. High-level expression and purification

of untagged and histidine-tagged HIV-1 reverse transcriptase. Protein Expr. Purif. 34:75-

86.

27.   Sambrook J, Russell DW. 2001. Molecular Cloning: A Laboratory Manual, 3rd ed. Cold

Spring Harbor Laboratory Press, Cold Spring Harbor, NY.

28.   Johnson KA. 1995. Rapid quench kinetic analysis of polymerases,

adenosinetriphosphatases, and enzyme intermediates. Methods Enzymol. 249:38-61.

29.   Jamburuthugoda VK, Guo D, Wedekind JE, Kim B. 2005. Kinetic evidence for

interaction of human immunodeficiency virus type 1 reverse transcriptase with the 3'-OH

of the incoming dTTP substrate. Biochemistry 44:10635-43.

30.   Kerr SG, Anderson KS. 1997. Pre-steady-state kinetic characterization of wild type and

434   3'-azido-3'-deoxythymidine (AZT) resistant human immunodeficiency virus type 1

435   reverse transcriptase: implication of RNA directed DNA polymerization in the

436   mechanism of AZT resistance. Biochemistry 36:14064-70.

437   31.   Kellinger MW, Johnson KA. 2011. Role of induced fit in limiting discrimination against

438   AZT by HIV reverse transcriptase. Biochemistry 50:5008-15.

439   32.   Yang G, Wang J, Cheng Y, Dutschman GE, Tanaka H, Baba M, Cheng Y-C. 2008.

440   Mechanism of inhibition of human immunodeficiency virus type 1 reverse transcriptase

441   by a stavudine analogue, 4'-ethynyl stavudine triphosphate. Antimicrob. Agents

442   Chemother. 52:2035-42.

443   33.   Krebs R, Immendorfer U, Thrall SH, Wohrl BM, Goody RS. 1997. Single-step kinetics

444   of HIV-1 reverse transcriptase mutants responsible for virus resistance to nucleoside

445   inhibitors zidovudine and 3-TC. Biochemistry 36:10292-300.

446   34.   Salie ZL, Kirby KA, Michailidis E, Marchand B, Singh K, Rohan LC, Kodama EN,

447   Mitsuya H, Parniak MA, Sarafianos SG. 2016. Structural basis of HIV inhibition by

448   translocation-defective RT inhibitor 4'-ethynyl-2-fluoro-2'-deoxyadenosine (EFdA). Proc.

449   Natl. Acad. Sci. U. S. A. 113:9274-9.

450   35.   Muftuoglu Y, Sohl CD, Mislak AC, Mitsuya H, Sarafianos SG, Anderson KS. 2014.

451   Probing the molecular mechanism of action of the HIV-1 reverse transcriptase inhibitor

452   4'-ethynyl-2-fluoro-2'-deoxyadenosine (EFdA) using pre-steady-state kinetics. Antiviral.

453   Res. 106:1-4.

36.   Astatke M, Grindley ND, Joyce CM. 1998. How E. coli DNA polymerase I (Klenow

fragment) distinguishes between deoxy- and dideoxynucleotides. J. Mol. Biol. 278:147-

65.

37.   Gao Y, Yang W. 2016. Capture of a third Mg□ is essential for catalyzing DNA synthesis.

Science (New York, N Y) 352:1334-7.

38.   Freudenthal BD, Beard WA, Shock DD, Wilson SH. 2013. Observing a DNA polymerase

choose right from wrong. Cell 154:157-68.

39.   Nakamura T, Zhao Y, Yamagata Y, Hua Y-j, Yang W. 2012. Watching DNA polymerase

eta make a phosphodiester bond. Nature 487:196-201.

40.   Vyas R, Reed AJ, Tokarsky EJ, Suo Z. 2015. Viewing Human DNA Polymerase beta

Faithfully and Unfaithfully Bypass an Oxidative Lesion by Time-Dependent

Crystallography. J. Am. Chem. Soc. 137:5225-30.

41.   Freudenthal BD, Beard WA, Perera L, Shock DD, Kim T, Schlick T, Wilson SH. 2015.

Uncovering the polymerase-induced cytotoxicity of an oxidized nucleotide. Nature

517:635-9.

[a]Nucleotide ([Mg^{2+}])	[b]k (s^{-1}) ± standard deviation	Relative rate vs. dTTP same [Mg^{2+}]	6/0.5 mM Mg^{2+} with same nucleotide
dTTP (0.5 mM)	8.6 ± 1.3	1	
dTTP (6 mM)	14.1 ± 1.2	1	1.6
Analogs with substitutions at the 3' position (chain terminators)			
ddTTP (0.5 mM)	1.4 ± 0.7	0.16	
ddTTP (6 mM)	7.2 ± 1.0	0.51	5.1
AZT (0.5 mM)	2.0 ± 0.5	0.23	
AZT (6 mM)	8.9 ± 1.9	0.63	4.5
D4T (0.5 mM)	1.8 ± 0.7	0.21	
D4T (6 mM)	8.3 ± 0.8	0.59	4.6
3'-amino dTTP (0.5 mM)	0.81 ± 0.13	0.094	
3'-amino dTTP (6 mM)	3.1 ± 0.1	0.22	3.8
3'-fluoro dTTP (0.5 mM)	0.54 ± 0.13	0.063	
3'-fluoro dTTP (6 mM)	1.5 ± 0.1	0.11	2.8
[c]3'-O-methyl dTTP (0.5 mM)	[c]3.3 x 10^{-5} ± 0.6 x 10^{-5}	NA	
[c]3'-O-methyl dTTP (6 mM)	[c]1.8 x 10^{-4} ± 0.2 x 10^{-4}	NA	5.4
Analogs with a 3' hydroxyl (translocation inhibitors)			
4'-C-methyl dTTP (0.5 mM)	4.5 ± 1.4	0.52	
4'-C-methyl dTTP (6 mM)	3.9 ± 2.8	0.28	0.88
4'-C-ethyl dTTP (0.5 mM)	1.3 ± 0.1	0.15	
4'-C-ethyl dTTP (6 mM)	3.9 ± 0.4	0.28	3.1
D-carba dTTP (0.5 mM)	1.1 ± 0.5	0.13	
D-carba dTTP (6 mM)	1.3 ± 0.1	0.092	1.2

[a]All nucleotides were tested at 5 µM final concentration. The free Mg^{2+} concentration in reactions is listed (see Methods). The nucleic acid primer-template used was: 20-nt primer, 5'-TTCCCGTCAGCCTAGCTGAG-3◻, and 36-nucleotide template, 5'-ATCGTCGTACTCAGCTAGGCTGACGGGAAATTGTAT-3◻.

[b]Results are averages from 3 or more independent experiments ± standard deviations. See Fig. 1 for an example of an experiment.

[c]Values for 3' O-methyl dTTP are from steady-state experiments as the rate was too low to reliably measure by pre-steady-state methods. Units are nM/min. They cannot be directly compared to other values for k. Conditions were 20 nM primer-template, 20 nM RT, 100 µM 3'-O-methyl dTTP in the same buffer as pre-steady-state reactions and reactions were measured over 30 min.

Table 2. Pre-steady state kinetic data for HIV RT incorporation of TTP, AZTTP, and D4TTP

Nucleotide	$[Mg^{2+}]$ (mM)	[a]k_{pol} (s^{-1})	[a]K_d (μM)	k_{pol}/K_d	[b]Efficiency decline
dTTP	0.5	14.3 ± 0.8	3.3 ± 0.6	4.3	1
AZTTP	0.5	3.3 ± 0.3	5.5 ± 1.7	0.6	7.2
d4TTP	0.5	4.9 ± 0.5	7.6 ± 2.2	0.64	6.7
dTTP	6	25.0 ± 1.3	4.2 ± 0.7	6.0	1
AZTTP	6	17.0 ± 0.6	3.9 ± 0.6	3.9	1.5
d4TTP	6	22.3 ± 1.7	11.6 ± 2.3	1.9	3.2

a- Values were derived from the graphs shown in Fig. 2 and are predicted values base on curve fitting ± standard error (as determined by the fitting program). See Table 1 and Methods for a description of the primer-template used.

b- Numbers represent the fold decline in enzyme efficiency relative to dTTP at the same Mg^{2+} concentration. Enzyme catalytic efficiency was taken as k_{pol}/K_d.

Figure 1. Example of pre-steady state determination of the rate (k) of nucleotide addition with 5 µM dTTP or AZTTP and 0.5 or 6 mM Mg^{2+}. (A) Assays were conducted as described in Materials and Methods with a 20-nt 5'-^{32}P end-labeled DNA primer bound to a 36-nt DNA template using rapid quench analysis. Time points in these assays were 10, 20, 40, 80, 100, 200, and 500 ms and were conducted in duplicate. For nucleotide analogs with slower incorporation rates longer time points were also used. Positions of the primer (P) and primer + 1 nucleotide (P+1) are shown for assays with dTTP and AZTTP at 0.5 or 6 mM Mg^{2+}. FE, full extension; -E, no enzyme added. (B) The data was fitted to an exponential equation as described in Materials and Methods to yield the incorporation rate (k (s^{-1})) used to produce the data in Table 1.

Figure 2. Addition of 3'-O-CH$_3$ dTTP in steady-state assays with HIV RT with 0.5 or 6 mM Mg^{2+}. Reactions were carried out as described in Materials and Methods using a 20-nt 5'-^{32}P end-labeled DNA primer bound to a 36-nt DNA template. All reactions contained 100 µM 3'-O-CH$_3$ dTTP with either 0.5 or 6 mM Mg^{2+}. Reactions were carried out for the indicated times and resolved on a 20% denaturing gel. The proportion of extended primer was quantified on a phosphorimager and used to determine the rate (k) of nucleotide addition. -E, no enzyme added; P, 5'-^{32}P end-labeled DNA primer; P+1, primer extended with 3'-O-CH$_3$ dTTP.

Figure 3. Pre-steady state determination of equilibrium dissociation constant (K_d) and k_{pol} for dTTP, AZTTP, and d4TTP. Assays were conducted as described in Materials and Methods with a 20-nt 5'-[32]P end-labeled DNA primer bound to a 36-nt DNA template using rapid quench analysis. Data points are from independent experiments at various concentrations of dTTP (A), AZTTP (B), or d4TTP (C) (see Fig. 1 for an example). Points from several experiments (typically 2-3, while only 1 point was used for the highest d4TTP concentration), were plotted and fit to an equation to determine values as described in Materials and Methods. The indicated values for K_d and k_{pol} are derived from the curve fit program. Values are ± standard errors as determined by the program.

Figure 4. Primer extension on a 100-nt DNA template with HIV RT. Reactions were carried out as described in Materials and Methods using a 20-nt 5'-^{32}P end-labeled DNA primer bound to a 100-nt DNA template. All reactions contained 5 µM dNTPs and 3 µM of the indicated dTTP analog with either 0.5 or 6 mM Mg^{2+}. Reactions were carried out for 30 min and resolved on an 8% denaturing gel. -E, no enzyme added. *d4TTP lane came from another gel and a separate experiment.