Dynamic Methylome Modification associated with mutational signatures in ageing and etiology of disease

Gayathri Kumar[1&2], Kashyap Krishnasamy[1&2], Naseer Pasha[1&2], Naveenkumar Nagarajan[1&2], Bhavika Mam[1&2], Mahima Kishinani[1], Vedanth Vohra[1], Villoo Morawala Patell[1,2,3]*

¹Avesthagen Limited, Bangalore, India

²The Avestagenome Project® International Pvt Ltd, Bangalore, Karnataka, India-560005

³AGENOME LLC, USA

*Corresponding Author:

Dr.Villoo Morawala Patell,

Avesthagen Limited

Yolee Grande, 2nd Floor,

14, Pottery Road,

Bangalore, 560005, Karnataka, India

Email: villoo@avesthagen.com

Abstract

Epigenetic markers and reversible change in the loci of genes regulating critical cell processes, have recently emerged as important biomarkers in the study of disease pathology. The epigenetic changes that accompany ageing in the context of population health risk needs to be explored. Additionally, the interplay between dynamic methylation changes that accompany ageing in relation to mutations that accrue in an individual's genome over time needs further investigation. Our current study captures the role for variants acting in concurrence with dynamic methylation in an individual analysed over time, in essence reflecting the genome-epigenome interplay, affecting biochemical pathways controlling physiological processes. In our current study, we completed the whole genome methylation and variant analysis in one Zoroastrian-Parsi non-smoking individual, collected at an interval of 12 years apart (at 53 and 65 years respectively) (ZPMetG-Hv2a-1A (old, t_0), ZPMetG-Hv2a-1B (recent, t_0+12)) using Grid-ion Nanopore sequencer at 13X genome coverage overall. We further identified the Single Nucleotide Variants (SNVs) and indels in known CpG islands by employing GATK and MuTect2 variant caller pipeline with GRCh37 (patch 13) human genome as the reference.

We found 5258 disease relevant genes differentially methylated across this individual over 12 years. Employing the GATK pipeline, we found 24,948 genes corresponding to 4,58,148 variants specific to ZPMetG-Hv2a-1B, indicative of variants that accrued over time. 242/24948 gene variants occurred within the CpG regions that were differentially methylated with 67/247 exactly occurring on the CpG site. Our analysis yielded a critical cluster of 10 genes which are significantly methylated and have variants at the CpG site or the ±4bp CpG region window. KEGG enrichment network analysis, Reactome and STRING analysis of mutational signatures of gene specific variants indicated an impact in biological process regulating immune system, disease networks implicated in cancer and neurodegenerative diseases and transcriptional control of processes regulating cellular senescence and longevity.

Our current study provides an understanding of the ageing methylome over time through the interplay between differentially methylated genes and variants in the etiology of disease.

Introduction

Aging is a complex and time-dependent deterioration of physiological processes. Increased human life expectancy have also resulted in higher morbidity rates since advanced age is a predominant risk factor for several diseases including cancer, dementia, diabetes, and cardiovascular disease (CVD)[1]. In addition to molecular and cellular factors like cellular senescence, telomere attrition, epigenetic changes that govern such processes comprise a significant component of the aging process[2]. Methylation signatures more specifically influence gene expression without altering the DNA sequence bridging the intrinsic and extrinsic signals. The most common methylation modifications involves the transfer of a methyl (CH_3) group from S-adenosyl methionine (SAM) to the fifth position of cytosine nucleotides, forming 5-methylcytosine (5mC)[3]. Methylation is a dynamic event wherein instances of reversal in methylation occur at certain sites, while progression with age causes methylation at many CpG sites in inter-genic regions such as TSS[4] (Transcription start sites), etc. The interplay of epigenetic events manifests in the regulation of major biological processes[5], such as development, differentiation, genomic imprinting and X chromosome inactivation (XCI).

DNA methylation both heritable and dynamic exhibits strong correlation with age and age-related outcomes. Additionally, the epigenome responds to a broad range of environmental factors and is sensitive to environmental influences[6] including exercise, stress[7], diet[8] and shift work. Epigenetic profiling of identical twins[9] indicate methylation events to be independent of genome homology with global DNA methylation patterns differing in the older compared to younger twins, consistent with an age-dependent progression[10] of epigenetic change. Global methylation changes over an 11-year span in participants of an Icelandic cohort, and age- and tissue-related alterations in some CpG islands from an array of 1413 arbitrarily chosen CpG sites near gene promoters, further corroborate the evidence for dynamic methylation patterns over time[11]. Promoter hyper-methylation has been shown to increase mutation rates suggesting influence of epigenetics over genetics. These studies argue in support for the role of epigenome changes in aging associated changes. It has been noted that genes related to various tumors can be silenced by heritable epigenetic events involving chromatin remodeling or DNAm (DNA

methylation). In addition to dynamic methylation changes over time, recent studies have also

suggested that epigenetic markers, or their maintenance, are controlled by genes and are and

tightly linked with DNA variants[12] that accrue in an individual genome over time. In fact,

methylation of cytosine to 5mC in the coding regions of genes increases the proneness to

mutations, consequently spontaneous hydrolytic deamination resulting in C to T transitions.

The epigenetic milieu therefore provides a homeostatic mechanism at the molecular level which

allows phenotypic malleability in response to the changing internal and external environments.

The flexibility and dynamic changes in the methylome highlight the importance and influence of

lifestyle changes in the control of gene regulation through epigenetic profiling. This also makes

methylome studies an attractive source to identify hotspots to evaluate the impact of lifestyle

changes on aging as well as provide potential targets for therapeutic intervention[13,14]. Besides

methylation changes, mutational signatures that include somatic variations can change the

preponderance of methylation of CpG islands. CpG-SNPs in the promoter regions have been

found to be associated with various disorders, including type 2 diabetes[15], breast cancer[16],

coronary heart disease[17], and psychosis[13].

Therefore, comparison of comprehensive methylation patterns in healthy individuals at multiple

time points can further our understanding of whether such changes can be early indicators of a

late onset chronic disease. Identifying such indicators earlier, can improve disease prognosis with

potential for reversibility. Our current epigenetic study is unique in analysing the genome-

epigenome interactions from an individual of the endogamous dwindling Zoroastrian Parsi

community, have higher median life span accompanied by a higher incidence of aging-associated

conditions, neurodegenerative conditions like Parkinsons disease, cancers of the breast, colon,

gastrointestinal, prostrate, auto-immune disorders, and rare diseases.

To identify the age and disease associated epigenetic changes in Parsi population, we performed

an intra-individual methylome profile of a Zoroastrian Parsi individual with Nanopore-based

sequencing technology separated by a time span of 12 years. The two samples from the same

non-smoking Zoroastrian Parsi individual, ZPMetG-HV2a-1A (old) and ZPMetG-HV2a-1B (recent)

(12 years apart) were sequenced through Oxford Nanopore Technology (ONT). Between the

samples, we found that aging increased global methylation frequencies for the equivalent CpG

sites, with an overall increase of hyper-methylated regions across genic regions as compared with

non-genic regions. 108 genes were significantly hypermethylated and 304 genes were

significantly hypomethylated. We also identified 10 unique CpG-SNP interactions in the form of

variants occurring at the CpG site and across the CpG region that may affect the biological process

of these occurrences in the genic regions. Enrichment of pathways associated with the genes

showed the implication of pathways involved in immune responses, pathways implicated in

cancer, neurogenerative diseases and physiological processes regulating cellular proliferation,

senescence regulating ageing and longevity.

Materials and Methods

DNA Extraction:

123   The high molecular weight genomic DNA was isolated from the buffy coat of EDTA-whole blood using

124   the Qiagen whole Blood Genomic DNA extraction kit (#69504). Extracted DNA Qubit™ dsDNA BR assay

125   kit (#Q32850) for Qubit 2.0® Fluorometer (Life Technologies™).

DNA QC

128   High molecular weight (HMW) DNA of optimal quality is a prerequisite for Nanopore library

129   preparation. Quality and quantity of the gDNA was estimated using Nanodrop

130   Spectrophotometer and Qubit fluorometer using Qubit™ dsDNA BR assay kit (#Q32850) from Life

131   Technologies respectively.

DNA purification

134   The DNA was subjected for column purification using Zymoclean Large Fragment DNA Recovery

135   kit (Zymo Research, USA) followed by fragmentation using g-TUBE (Covaris, Inc.). The purified

136   DNA samples were used for library preparation.

Library preparation

A total of 2 µg from each sample was used for nanopore library preparation using ligation sequencing kit

(SQK-LSK109). Briefly, 2 µg of gDNA from each sample was end-repaired (NEBnext ultra II end repair kit,

New England Biolabs, MA, USA) and purified using 1x AmPure beads (Beckmann Coulter, USA). Adapter

ligation (AMX) was performed at RT (20 ^{o}C) for 20 minutes using NEB Quick T4 DNA Ligase (New England

Biolabs, MA, USA). The reaction mixture was purified using 0.6X AmPure beads (Beckmann Coulter, USA)

and sequencing library was eluted in 15 µl of elution buffer provided in the ligation sequencing kit (SQK-

LSK109) from Oxford Nanopore Technology (ONT).

Sequencing and sequence processing

Sequencing was performed on GridION X5 (Oxford Nanopore Technologies, Oxford, UK) using

SpotON flow cell R9.4 (FLO-MIN106) as per manufacturers recommendation. Nanopore raw

reads (*'fast5'* format) were base called (*'fastq5'* format) using Guppy v2.3.4 software.

Genomic DNA samples were quantified using the Qubit fluorometer. For each sample, 100 ng of

DNA was fragmented to an average size of 350 bp by ultrasonication (Covaris ME220

ultrasonicator). DNA sequencing libraries were prepared using dual-index adapters with the

TruSeq Nano DNA Library Prep kit (Illumina) as per the manufacturer's protocol. The amplified

libraries were checked on a Tape Station (Agilent Technologies) and quantified by real-time PCR

using the KAPA Library Quantification kit (Roche) with the QuantStudio-7flex Real-Time PCR

system (Thermo). Equimolar pools of sequencing libraries were sequenced using S4 flow cells in

a Novaseq 6000 sequencer (Illumina) to generate 2 x 150-bp sequencing reads for 30x genome

coverage per sample.

Extracting methylation basecalls reads and QC

The sequence information is encoded in signal-level data measured by the Nanopore sequencer. The raw

signal data in Fast5 format is then demultiplexed using the Guppy basecaller and the adapters are

removed using Porechop to obtain the basecalled reads. The quality of the reads (in Fastq format)

post adapter removal is evaluated using Fastqc (Supplementary Figure.1). The adapter trimmed

reads in Fastq format are then mapped to the Human reference genome (GRCh37 (patch 13)

using minimap2 (Version 2.17). The aligned reads obtained are then used to identify the

methylated sites using Nanopolish software (Version 0.13.2). This tool provides a list of reads

with the log-likelihood ratios (wherein a positive value is evidence that the cytosine is methylated

to 5-mC and a negative value indicates a Cytosine is present). From this file, we obtain the

consolidated methylation frequency file for both the sequenced samples. Average read length of

5.77 kb and 12.15 kb was obtained from ZPMetG-HV2a-1A (t0, old) and ZPMetG-HV2a-1B (t0+12,

recent) respectively.

Differential methylation analysis

The reads with methylated CpGs from the methylation frequency files of the two samples

(ZPMetG-HV2a-1A and ZPMetG-HV2a-1B) were compared using bedtools (Version.v2.29.2).

Reads with corresponding entries available in the two samples were taken further for differential

methylation analysis.

To the above, methylated reads common to the samples, ZPMetG-HV2a-1A and ZPMetG-HV2a-

1B annotations were provided from the GRCh37 GTF file.

Annotations were provided in the following manner –

-   Promoter regions were identified as 2Kb upstream of genic regions.

-   TSS (Transcription start sites) for genes on '+' strand, column 2 and column 3 for genes

on '- 'strand

-   Genic and Non-genic regions were extracted based on the tags provided in the third

column of the GTF file.

Hypomethylated and Hypermethylated regions

The porechopped Fastq reads for the two samples were mapped back to the reference genome

GRCh37 (patch 13) using minimap2 and the aligned outputs in BAM (Binary Alignment Map) then

represented in bed format. This was carried out to identify the regions sequenced in both the

samples. The unique regions in each sample that were methylated would correspond to

hypomethylated or hypermethylated regions. A region would be tagged as only Hypermethylated

if there were reads available for the given region in the sample ZPMetG-HV2a-1A but not

reported in the Nanopolish generated methylation frequency file. It would be tagged as only

Hypomethylated if it is reported in the Nanopolish generated methylation frequency file of

ZPMetG-HV2a-1A but no such corresponding record in ZPMetG-HV2a-1B for a region sequenced

with reads available. The gene annotations for these regions were obtained in the same manner

as discussed in the previous section.

Read mapping and Variant calling for illumina (ZPMetG-HV2a-1A and ZPMetG-HV2a-1B)

Single Nucleotide Variants (SNVs) and indels were called using four different pipelines through a

combination of two read mappers and two variant callers. The GRCh37 (patch 13) human genome

was used as the reference genome to map the paired end reads. The two read mappers used

were BWA-MEM (Version 0.7.17) and Bowtie2 (version 2.4.1). The variant calling pipeline was

implemented using GATK (Version 7.3.1).

The GATK pipeline included additional read and variant processing steps such as duplicate

removal using Picard tools, base quality score recalibration, indel realignment, and genotyping

and variant quality score recalibration using GATK, all used according to GATK best practice

recommendations[18]. The MuTect2 tool[19] was employed for identifying the putative somatic

variants, following which Snpeff (build 2017-11-24) was used to annotate the variants along with

functional prediction. As described later in the Results section, variants identified using the BWA+

GATK pipeline were used for all downstream analysis. Variants in the intersection of all four

pipelines (two read mappers and two variant callers) were confidently identified, where the

intersection is defined as variant calls for which the chromosome, position, reference, and

alternate fields in the VCF files were identical.

Pathway mapping of genes to assess physiological implications

A compendium of genes from GSEA were extracted corresponding to tumor suppressors, oncogenes, etc.

Housekeeping genes were collated from (source). Genes associated with aging regulating pathways were

obtained from KEGG and other published work(s). KEGG pathway genes associated with different

neurological and physiological disorders were extracted. For this set of genes, the epigenetic changes

between the two samples were examined. The epigenetic changes were measured in terms of differences

in methylation frequencies for ZPMetG-HV2a-1A and ZPMetG-HV2a-1B**.** The frequency of methylation is

given by the ratio of the number of methylated reads in a gene region (m_r) and the total number of reads

detected in the gene region (T_r).

227                                 $$f_{meth}= m_r/T_r$$

229   The difference in methylation frequency (Δf_{meth}) between the subject at ZPMetG-HV2a-1B (t_0+12)

230   and ZPMetG-HV2a-1B (t_0 years) was obtained by subtracting the methylation frequency for t_0

231   from that of t_0+12.

232                      $$\Delta f_{meth} = (\Delta f_{meth})_{t0+12} - (\Delta f_{meth})_{t0}$$

The degree of methylation was primarily classified as 'hypomethylated', 'hemi-hypomethylated',

'hemi-hypermethylated' and 'hypermethylated' based on the int Δf_{meth} intervals. Here, a term

'background' was also defined corresponding to low-confidence scores of CpG methylation i.e.

Δf_{meth}, greater than or equal to -0.2 but lesser than 0.2. Hypomethylation has been defined as a

gradient of hypomethylation (Δf_{meth} < than 0.6) and hemi-hypomethylation (-0.6 <= Δf_{meth} <-0.2).

Hypermethylation has been defined as a gradient of hypermethylation (Δf_{meth} > than 0.6) and

hemi-hypermethylation (0.2 <= Δf_{meth} <0.6). The weighted difference in methylation frequency

($\Delta f_{meth\_weighted}$) for each read within a gene region was obtained by factoring in the number of

CpG islands (cpg_{num}) detected for that gene region. Thus, the weighted methylated frequency

was obtained as a product of the difference in methylation frequency and the number of CpG

reads in the gene region.

245                        $$\Delta f_{meth\_weighted} = \Delta f_{meth} * cpg_{num}$$

247   The normalized weighted methylation frequency score (F_{meth_norm}) per degree of methylation was

248   obtained by normalizing the summated $\Delta f_{meth\_weighted}$ score with the total number of reads (T_r).

250                  $$F_{meth\_norm} = \Sigma (\Delta f_{meth\_weighted})_{degree} / Tr$$

251   The normalized weighted difference in methylation frequency score was assigned to a specific

252   gene and taken further for downstream analysis.

Results

Analysis of individual methylome modifications over time

To account for difference in coverage depth, we extracted the nucleotide ranges for which base called entries are present in the methylation frequency files for both samples. This enables identifying any distinctive increase or decrease in methylation between the two temporally separated samples. These comparable nucleotide ranges extracted from the methylation base called regions accounted for 18,758,642 regions which covered 98% and 84% of reported regions in fastq files of ZPMetG-HV2a-1A (old) and ZPMetG-HV2a-1B (recent) respectively. The overall analysis pipeline is elaborated in Figure.1A. We examined the overall distribution in the methylation frequency between ZPMetG-HV2a-1A (old) and ZPMetG-HV2a-1B (recent) We applied a non-parametric statistical test (Mann Whitney U) and observed a statistically significant difference between the two sample distributions (Figure.1B). The chromosome wise distribution of the methylated frequencies in CpG dinucleotides was also examined to discern changes in methylation between the samples showed similar profiles for genic and intergenic regions. Notable exceptions were observed for Chr.21, 22 and X for genic regions (Supplementary Figure 2). The median of the distribution is given in Fig.1B for each chromosome while the entire frequency distributions (Figure.1B).

Functional motif-based characterization of CpG's common to ZPMetG-Hv2a-1 and ZPMetG-Hv2a-1

The methylation base called regions compared between ZPMetG-HV2a-1A (old) and ZPMetG-HV2a-1B (recent) were mapped to GRCH37 (The human reference genome from NCBI). Based on the tag 'gene', we extracted putative genic regions, while other regions (excluding even CDS (coding sequence), mRNA, tRNA etc) were extracted to examine the non-genic regions (Figure. 2A). Majority of the CpGs were found in the genic region and the gene promoter regions (Figure. 2B). The distribution across chromosomes for genic and non-genic regions in the two samples indicated majority of the CpGs on genic regions and the lowest for TSS regions. The distribution

of CpG's was similar across all chromosomes for the genic regions compared to the distribution

for TSS, non-genic and genic promoters that displayed a staggered distribution across

chromosomes (Figure. 2C)

Prioritization of genes families:

Characterization of the methylated regions indicated 26,019 genes that are found in the regions

methylated in ZPMetG-HV2a-1B (recent) compared to ZPMetG-HV2a-1A (old) and Selective

filtering and prioritization of the 26,019 genes (based normalized score) based on a curated

dataset of 3050 gene families from the Gene Set Enrichment Analysis GSEA database[20], 168

Epigenetic modifier gene database[21], 2833 Housekeeping genes from HRT Atlas v1.0[22], 3527

pathway and disease specific genes from KEGG database[23] yielded 5214 unique genes. It is

cultural practice that followers of the Zoroastrian faith consider smoking a taboo, resulting in

generations of the community that refrained from smoking. To address the epigenetic changes

associated with smoking related genes and its relevance in the non-smoking Parsi community,

we also curated a list of 44 documented genes that were differentially methylated in smokers.

Taken together, out total gene list comprised of 5258 genes (Supplementary Table.2).

Hierarchical clustering of Differentially Methylated Regions in ZPMetG-Hv2a-1B compared to ZPMetG-
Hv2a-1

We performed hierarchical clustering of the differentially methylated genes by first obtaining the

frequency of CpG methylation across the entire gene length. Based on the weighted score for

CpG methylation (materials and methods), the CpG frequencies across genes were classified as

Hypo, Hemi-Hypo, Hemi-Hyper and Hyper methylated regions. The weighted score common

across all genes was classified as background that served as an internal control.

The hierarchical clustering of row-wise Z score of common Differentially Methylated Gene

regions (Supplementary Table.3) in ZPMetG-Hv2a-1B identified a cluster of 103 genes that were

significantly hyper-methylated and 307 genes that were hypo-methylated (Figure. 3A,

Supplementary Table.7). GSEA based enrichment indicated an enrichment of genes coding for

transcription factors and protein kinases in the hyper and hypomethylated clusters. Among

Hypermethylated genes, Homeodomain proteins were highly represented, while among the

hypomethylated genes, cell differentiation markers, tumor suppressors and translocated cancer

genes dominated in comparison to each other (Figure.3B, Supplementary Figure.3A)

PANTHER based gene annotation classified these genes based on their molecular and protein

function (Supplementary Figure 3B). Hypermethylated and Hypomethylated clusters varied in

their functional classification with a presence of genes that function to increase transcription

regulator activity (Supplementary Figure 3B)

Characterization of variants unique to ZPMetG-Hv2a-1B compared to ZPMetG-Hv2a-1

While dynamic methylation events constitute a major mechanism of epigenetic modification,

variants in the form of SNPs in addition to dynamic methylation events are known to be critical

to epigenome function through the modification of genomic information. To this end, we

employed the GATK variant characterization pipeline to identify variants specific to ZPMetG-

Hv2a-1B viz., variants that have accumulated in the individual because of ageing. We identified

4,58,148 variants corresponding to 24,948 unique genes to ZPMetG-HV2a-1B (recent).

We next sought to identify genes common to both dynamic methylated regions (CpGs) and

variants specific to ZPMetG-Hv2a-1B. We found a direct correlation between gene length, CpG

count and variant count. This association is specifically significant to the correlation between

variants and CpG (Pearson correlation coefficient, r:0.86) (Figure.4A, Supplementary Figure 4A,

B). Analysis of the 5258 ZPMetG-Hv2a-1B specific differentially methylated genes post

disease/GSEA prioritisation filter for somatic variants yielded 4409 differentially methylated

genes (Supplementary Table.5) that are specific to ZPMetG-HV2a-1B (recent) that harbour

somatic variants (Supplementary Figure 4C).

The location of variants with respect to their occurrence on the CpG region is crucial to the

resultant effect on the epigenomic imprinting in the form of methylation changes. Variants that

occur exactly on the CpG site may modulate the resultant methylation, thereby effecting

methylation changes. In this context, we classified ZPMetG-HV2a-1B variants based on their loci

of incidence on the CpG region. We classified variant occurrence as those that occur exactly on

the CpG region and variants that occur at a distance window of 1-4bp either side of the CpG

region (±4bp). Our analysis indicated 242 variants classified as "Modifier variants" occurring

within the CpG region of 177 Differentially Methylated Genes specific to ZPMetG-Hv2a-1B

(Figure. 4B, Supplementary Table.6). 27.68% (67/242 variants) occur exactly at the CpG site and

72% (175/242 variants) occur in the CpG region at a loci window of ±4bp (Fig. 4B). Majority of

classified variants occur in Chr.1 with most variants occurring at ±4bp window size across all

chromosomes except for Chr.4 where more variants are observed exactly at the CpG site

compared to the window size (Supplementary Figure 5A). The number of variants within genes

varied between 1-7 variants (Supplementary Figure 5B). Characterization of the pathway

association and interaction networks differed between the genes harbouring variants at the CpG

site compared to genes with variants in the CpG region (Supplementary Figure.6,7)

Interplay of genome and epigenome in the context of ageing and disease etiology

The presence of variants in the genome can affect the epigenetic modification. We therefore

sought to understand the biological processes of the genes in our study that carried both

methylation changes and harboured variants in the CpG region. To this end, we compared 412

genes with significant hyper (108 genes), hypo methylation (304 genes) changes (Supplementary

Table.7) for gene specific variants within the CpG region. Our analysis yielded a critical cluster of

10 genes which are significantly methylated (hyper/hypo) and have variants at the CpG site or

the ±4bp CpG region window (Figure.5A, Supplementary Table 8). Only 2 genes (*CASP8, PCGF3*)

significantly hypomethylated carried variants at the CpG site, whereas both significantly

hyper/hypo methylated genes carried variants at the CpG window region (3 Hypermethylated

genes and 6 hypomethylated genes). A vast majority of genes were critical housekeeping genes,

followed by transcription factors, protein kinases and genes implicated in neurodegenerative

diseases (Figure. 5B).

To identify the relevance of these 10 genes in pathways related to disease and homeostasis, we

clustered the 10 genes with 1000 genes that harboured variants and CpG's (Supplementary

Table.9, based on normalized cumulative CpG.Variant score). Using Network analyst[24] based

clustering using KEGG based disease enrichment module showed that the majority of the

clustered genes are implicated in cell proliferative pathways regulating cancer, cancer subtypes,

neurodegenerative disease like Parkinsons disease, Alzheimers disease, Huntington disease and

pathways implicated in cellular senescence and longevity regulating gene clusters (Figure 5C,

Supplementary Table.10). Reactome analysis of the biological diversity of the gene function

showed their association to pathways involved in immune system, DNA repair, DNA

transcriptional activity, cell cycle and disease related pathways (Figure.5D).

Mutational signatures associated with ageing (ZPMetG-Hv2a-1B) across all genes and CpG

specific region of functionally relevant genes

We next proceeded to categorise mutational signatures that are characteristic combinations of

mutation types arising from specific mutagenesis process that involve alterations related to DNA

replication, repair and environmental exposure. Our analysis of ZPMetG-Hv2a-1B showed the

presence of 110,616 with a majority associated with C>T transitions followed by T>C transitions.

Analysis of the same in 4409 prioritized genes that show differential methylation and variants

specific to ZPMetG-Hv2a-1B indicated a similar signature for C>T transitions, followed by an

increase in C>A and T>A mutations, while T>C mutations were reduced compared to the

cumulative variants in ZPMetG-Hv2a-1B (Figure.6A). Further categorisation of the approach using

gene sets that consist of epigenetic modifiers, disease specific lists and smoking related

(environmental exposure) showed a high prevalence of C>T transitions especially at the CpG site

for gene sets regulating Alzheimers and Parkinsons pathways, while smoking associated genes

had a decreased incidence of C>T transitions at CpG site but an increase in overall C>T transitions

(Figure.6B)

Discussion:

Aging is one of the important contributors to chronic diseases. Epigenetic modifications in the

form of methylation of CpG di-nucleotides, play a crucial role in regulating physiological

processes which vary as age progresses. Differential methylation at CpG sites between younger

and older subjects have been shown associated with genes - MTOR, ULK1, ADCY6, IGF1R, CREB5

and RELA linked to metabolic traits[25], and CREB5, RELA, and ULK1 were statistically associated

with age. We found 5258 genes significantly altered in terms of methylation differences over

time in an individual assessed at a time of 12 years. DNAm levels of several CpG sites located at

genes involved in longevity-regulating pathways can be examined for DNAm in metabolic

alterations associated with age progression.

Along with epigenetic modifications, variants in the CpG sites have been shown to adversely

affect gene function and activity, especially in the context of disease associated traits[26]. Many

studies have reported that CpG-SNPs are associated with different diseases, such as type 2

diabetes, breast cancer, coronary heart disease, and psychosis which show a clear interaction

between genetic (SNPs) and epigenetic (DNA methylation) regulation. The introduction or

removal of CpG dinucleotides (possible sites of DNA methylation associated with the

environment) has been suggested as a potential mechanism through which SNPs can influence

gene transcription and expression via epigenetics. Our analysis showed the presence of 242

variants classified as "Modifier variants" occurring within the CpG region of 177 Differentially

Methylated Genes specific to ZPMetG-Hv2a-1B. These are genes involved in longevity regulating

pathways, cellular proliferation, and transcriptional regulation.

Recent reports using tissue-specific methylation data describe a strong association between C>T

mutations and methylation at CpG dinucleotides in many cancer types, driving patterns of

mutation formation throughout the genome. In our analysis, the variants occurring at the CpG

sites were C>T transitions, that constitute two genes *CASP8, PCGF3*. Incidentally *PCGF3* in our

analysis displays transitions not only at the CpG site but also transitions within the CpG region.

Pcgf3/5 mainly function as transcriptional activators driving expression of many genes involved

in mesoderm differentiation and *Pcgf3/5* are essential for regulating global levels of the histone

modifier H2AK119ub1 in Embryonic Stem cells[27]. *CASP8* methylation has been shown to be

relevant in cancers and may function as a tumor suppressor gene in neuroendocrine lung

tumors[28]. Our study points to a crucial interplay between CpG methylation levels and variant

incidence. It is therefore critical to extend the study to other individuals to understand the

diversity of mutational signatures that accrue in CpG regions of functionally important genes to

further dissect the epigenome-genome interactions in the context of homeostasis, disease and

ageing.

Mutational signatures associated to ageing and cancer subtypes have been studied and indicate

a strong correlation towards cytosine deamination at CpG sites, that result in an increase of C>T

transitions with age of diagnosis, as age allows more time for deamination events leading to

accumulation of their effects[29]. In line with these observations, we find that C>T transitions

constitute most of the mutational changes in ZPMetG-Hv2a-1B, indicative of their accumulation

over age. In specific, we find an increase of the same transitions at CpG sites in genes that

participate in Alzheimers and Parkinsons disease pathways but their relative contribution is

decreased in smoking associated genes. Further analysis of the same signatures in different

cohorts corresponding to diseases and smoking status would be critical in identifying the

combinatorial effect of mutational signatures associated with epigenome modifications in the

context of ageing.

Our study therefore adds to an important understanding in personal methylome analysis in the

evidence for interactions between epigenome and genome in the form of CpG:variant

interactions. The approach would be significant in identifying significant genome loci that reflect

the effects of methylation modifications in combination with genetic variation that accrue in an

individual, therefore aid in identifying associations with longevity and associated traits like

disease onset.

Declarations:

Ethics approval and consent to participate

The samples of peripheral blood collected in this study involve human healthy donors and were obtained with their informed consent. were in accordance with the ethical standards of the institution (Avesthagen Limited, Bangalore, India) and in line with the 1964 Helsinki declaration and its later amendments. The Zoroastrian Parsi adult participant (>18 years) underwent height and weight measurements and answered an extensive questionnaire designed to capture their medical, dietary, and life history. The subject provided written informed consent for the collection of samples and subsequent analysis. This study was approved by the Avesthagen Ethics Committee constituted under the Department of Biotechnology, Government of India (BLAG-CSP-033).

Availability of data and materials

The authors declare that all data and materials are available upon request.

Competing interests

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Funding

The project was funded by the grant awarded to Dr.Villoo Morawala-Patell "Cancer risk in smoking subjects assessed by next-generation sequencing profile of circulating free DNA and RNA" (GG-0005) by the Foundation for a Smoke-Free World, New York, USA.

Authors contributions

VMP conceptualized, designed, guided the experiments and analysis; VMP founded The Avestagenome Project[TM] and provided access to the dataset for this study; NP, GK, NN, BM and KK analysed the sequences, performed bioinformatics analysis, interpreted the results and

plotted the graphs and figures; MK, VV provided support in the form of literature mining and

database curation. VMP, NP, GK, NN and KK drafted the manuscript. All authors reviewed the

manuscript.

Acknowledgements

We thank the Foundation for Smoke-Free World, who is advancing global progress in smoking

cessation and harm reduction, for funding this project and Dr. Derek Yach for his support of this

project. We thank Genotypic technologies and Dr.Raja Mugasimangalam, Bangalore for their

excellent sequencing services and technical assistance. We thank the Zoroastrian-Parsi

community of India for their enthusiastic cooperation. We thank Dr.Sami Gazder, Kouser

Sonnekhan, and the The Avestagenome Project™ project team.

References

1. Morgan, A. E., Davies, T. J. & McAuley, M. T. The role of DNA methylation in ageing and cancer. in *Proceedings of the Nutrition Society* vol. 77 412–422 (Cambridge University Press, 2018).

2. Aunan, J. R., Watson, M. M., Hagland, H. R. & Søreide, K. Molecular and biological hallmarks of ageing. *British Journal of Surgery* (2016) doi:10.1002/bjs.10053.

3. Moore, L. D., Le, T. & Fan, G. DNA methylation and its basic function. *Neuropsychopharmacology* (2013) doi:10.1038/npp.2012.112.

4. Marttila, S. *et al.* Ageing-associated changes in the human DNA methylome: Genomic locations and effects on gene expression. *BMC Genomics* (2015) doi:10.1186/s12864-015-1381-z.

5. Chaligné, R. & Heard, E. X-chromosome inactivation in development and cancer. *FEBS Letters* (2014) doi:10.1016/j.febslet.2014.06.023.

6. Vaiserman, A. Epidemiologic evidence for association between adverse environmental exposures in early life and epigenetic variation: a potential link to disease susceptibility? *Clinical Epigenetics* (2015) doi:10.1186/s13148-015-0130-0.

7. Burns, S. B., Szyszkowicz, J. K., Luheshi, G. N., Lutz, P. E. & Turecki, G. Plasticity of the epigenome during early-life stress. *Seminars in Cell and Developmental Biology* (2018) doi:10.1016/j.semcdb.2017.09.033.

8. Zhang, Y. & Kutateladze, T. G. Diet and the epigenome. *Nature Communications* (2018) doi:10.1038/s41467-018-05778-1.

9. Fraga, M. F. *et al.* Epigenetic differences arise during the lifetime of monozygotic twins. *Proc. Natl. Acad. Sci. U. S. A.* (2005) doi:10.1073/pnas.0500398102.

10. Talens, R. P. *et al.* Epigenetic variation during the adult lifespan: Cross-sectional and longitudinal data on monozygotic twin pairs. *Aging Cell* (2012) doi:10.1111/j.1474-9726.2012.00835.x.

11. Bjornsson, H. T. *et al.* Intra-individual change over time in DNA methylation with familial clustering. *JAMA - J. Am. Med. Assoc.* (2008) doi:10.1001/jama.299.24.2877.

12. Bonder, M. J. *et al.* Disease variants alter transcription factor levels and methylation of their binding sites. *Nat. Genet.* (2017) doi:10.1038/ng.3721.

13. Van Den Oord, E. J. C. G. *et al.* A whole methylome CpG-SNP association study of psychosis in blood and brain tissue. *Schizophr. Bull.* (2016) doi:10.1093/schbul/sbv182.

14. Zhao, S. G. *et al.* The DNA methylation landscape of advanced prostate cancer. *Nat. Genet.* (2020) doi:10.1038/s41588-020-0648-8.

15. Dayeh, T. A. *et al.* Identification of CpG-SNPs associated with type 2 diabetes and

differential DNA methylation in human pancreatic islets. *Diabetologia* (2013)
doi:10.1007/s00125-012-2815-7.

16. Harlid, S. *et al.* A candidate CpG SNP approach identifies a breast cancer associated ESR1-
SNP. *Int. J. Cancer* (2011) doi:10.1002/ijc.25786.

17. Chen, X. *et al.* Association of six CpG-SNPs in the inflammation-related genes with
coronary heart disease. *Hum. Genomics* (2016) doi:10.1186/s40246-016-0067-1.

18. Van der Auwera, G. A. *et al.* GATK Best Practices. *Curr. Protoc. Bioinformatics* (2002).

19. Tools, V. D. MuTect2. *GATK Man.* (2017).

20. Subramanian, A. *et al.* Gene set enrichment analysis: A knowledge-based approach for
interpreting genome-wide expression profiles. *Proc. Natl. Acad. Sci. U. S. A.* (2005)
doi:10.1073/pnas.0506580102.

21. Nanda, J. S., Kumar, R. & Raghava, G. P. S. dbEM: A database of epigenetic modifiers
curated from cancerous and normal genomes. *Sci. Rep.* (2016) doi:10.1038/srep19340.

22. Hounkpe, B. W., Chenou, F., de Lima, F. & De Paula, E. V. HRT Atlas v1.0 database:
redefining human and mouse housekeeping genes and candidate reference transcripts
by mining massive RNA-seq datasets. *Nucleic Acids Res.* (2020) doi:10.1093/nar/gkaa609.

23. Tanabe, M. & Kanehisa, M. Using the KEGG database resource. *Curr. Protoc. Bioinforma.*
(2012) doi:10.1002/0471250953.bi0112s38.

24. Xia, J., Gill, E. E. & Hancock, R. E. W. NetworkAnalyst for statistical, visual and network-
based meta-analysis of gene expression data. *Nat. Protoc.* (2015)
doi:10.1038/nprot.2015.052.

25. Salas-Pérez, F. *et al.* DNA methylation in genes of longevity-regulating pathways:
Association with obesity and metabolic complications. *Aging (Albany. NY).* (2019)
doi:10.18632/aging.101882.

26. Zhou, D. *et al.* Polymorphisms involving gain or loss of CpG sites are significantly enriched
in trait-associated SNPs. *Oncotarget* (2015) doi:10.18632/oncotarget.5650.

27. Zhao, W. *et al.* Polycomb group RING finger proteins 3/5 activate transcription via an
interaction with the pluripotency factor Tex10 in embryonic stem cells. *J. Biol. Chem.*
292, 21527–21537 (2017).

28. Shivapurkar, N. *et al.* Differential inactivation of caspase-8 in lung cancers. *Cancer Biol.*
Ther. (2002) doi:10.4161/cbt.1.1.45.

29. Alexandrov, L. B. *et al.* Clock-like mutational processes in human somatic cells. *Nat.*
Genet. (2015) doi:10.1038/ng.3441.

Figures

Figure 1: Process workflow and Methylation frequency distribution in an
individual separated by two time points

Figure 1: (A) Flowchart representing the workflow for analysis of individual methylome and mutational
signature (SNP) variations across an individual separated by a duration of 12 years. **(B)** Methylation
frequency distribution for compared base called regions shown for ZPMetG-Hv2a-1B (recent, in green)
and ZPMetG-Hv2a-1A (old, in red). On the right is shown the histogram, identifying the nature of
distribution to be non-Gaussian. Median values of the methylation frequency distribution chromosome
wise for ZPMetG-HV2a-1A (old, in red) and ZPMetG-HV2a-1B (recent, in green)

Figure 2: Distribution and classification of common methylation imprints (CpG)
based on genome architecture

A)

B) C)

Figure 2: (A) Histogram representing the distribution and classification of methylation imprints (CpG) based on genome architecture and classifiers**. (B)** Histogram depicting classification of common methylation imprints (CpG) based on genomic characterization (inlay, left); Chromosome wise distribution of CpG numbers common to ZPMetG-Hv2a-1 and ZPMetG-Hv2a-2 classified based on distribution across genomic motifs (inlay, right)

Figure 3: Heatmap of Differentially Methylated Genic Regions observed in
ZPMetG-HV2a-1B (recent) compared to ZPMetG-HV2a-1A (old)

A)

B)

Figure 3: Heatmap displaying hierarchical clustering of row-wise Z score of common Differentially
Methylated Gene regions in ZPMetG-Hv2a-1B. Clusters of Hypermethylated genes (inlay figure; right, top)
and Hypomethylated (inlay figure; right, bottom). (B) GSEA based classification of hyper and
hypomethylated clusters

Figure 4: Distribution of SNPs across CpG islands (Methylated regions) in ZPMetG-HV2a-1B (recent)

A)

B)

Figure 4: A) Correlation between CpG and variant counts occurring in ZPMetG-Hv2a-1B. Pearson correlation coefficient,r is displayed on the side B) Distribution of ZPMetG-HV2a-1B(recent) specific variants across Differentially Methylated Regions specific to ZPMetG-HV2a-1B (recent)

Figure 5: Biological effect and pathway association of ZPMetG-HV2a-1B (recent) variants occurring exactly in the CpG sites

A)

B)

C)

D)

Figure 5: A) Distribution of CpG variants (exact, non-exact) across significant hyper and hypomethylated genes in ZPMetG-Hv2a-1B B) Categorization of 10 critical genes based on biological acitivity C) Network analyst using KEGG based disease enrichment module for 10 critical and 1000 high priority variants D) Reactome analysis of biologically relevant pathways of 10 critical and 1000 high priority variants

Figure 6: Assessment of mutational signatures and its distribution among CpG sites of ZPMetG-HV2a-1B (recent) in context of disease etiology

A)

B)

Figure 6: A) Distribution of mutational signatures across the genome and specific to CpG regions in 4409 prioritized genes and cumulative variants across ZPMetG-Hv2a-1B B) Mutational signatures indicative of transitions and transversions specific to ZPMetG-Hv2a-1B across different disease categories.