Running Head Anatomy of a phenological mismatch

Title The anatomy of a phenological mismatch: interacting consumer demand and resource

characteristics determine the consequences of mismatching

Authors Luke R. Wilde[1], Josiah E. Simmons[2], Rose J. Swift[3], Nathan R. Senner[1]

Affiliations

[1] Department of Biological Sciences, University of South Carolina, 715 Sumter St., Columbia,

SC 29208

[2] Division of Biological Sciences, University of Montana, Missoula, Montana

[3] U.S. Geological Survey, Northern Prairie Wildlife Research Center, 8711 37th Street SE

Jamestown, ND 58401

Abstract

Climate change has caused shifts in seasonally recurring biological events and the temporal

decoupling of consumer-resource pairs – i.e., phenological mismatching. Despite the

hypothetical risk mismatching poses to consumers, they do not invariably lead to individual- or

population-level effects. This may stem from how mismatches are typically defined, e.g., an

individual or population is 'matched' or 'mismatched' based on the degree of asynchrony with a

resource pulse. However, because both resource availability and consumer demands change over

time, this categorical definition can obscure within- or among-individual fitness effects. We

therefore developed models to identify the effects of resource characteristics on individual- and

population-level processes and determine how the strength of these effects change throughout a

consumer's life. We then measured the effects of resource characteristics on the growth, daily

survival, and fledging rates of Hudsonian godwit (*Limosa haemastica*) chicks hatched near

Beluga River, Alaska. At the individual-level, chick growth and survival improved following

periods of higher invertebrate abundance but were increasingly dependent on the availability of

larger prey as chicks aged. At the population level, seasonal fledging rates were best explained

by a model including age-structured consumer demand. Our study suggests that modelling the

effects of mismatching as a disrupted interaction between consumers and their resources

provides a biological mechanism for how mismatching occurs and clarifies when it matters to

individuals and populations. Given the variable responses to mismatching across consumer

populations, such tools for predicting how populations may respond under future climatic

conditions will be invaluable.

Keywords

mismatch; climate change; Bayesian hierarchical model; ontogeny; resource availability

Introduction

Shifts in the timing of recurring biological events (i.e., phenology) are among the best documented effects of climate change (Parmesan & Yohe, 2003). Higher spring temperatures have led to earlier peaks in seasonal resources (Thackeray et al., 2016), but slower rates of phenological advance at upper trophic levels mean that future climate conditions will likely lead to a greater decoupling of consumer-resource pairs – i.e., 'mismatching' – and heightened extinction risk for consumer populations (Both & Visser, 2001; Both et al., 2009). However, despite the theoretical risks imposed by climate-induced mismatching, mismatches do not invariably lead to reduced individual fitness (Dunn et al., 2011; Corkery et al., 2019) or negative demographic effects for populations (Visser et al., 2012; Reed et al., 2013; Keogan et al., 2020). Recent studies have proposed improved methodologies for studying mismatches (Visser & Gienapp, 2019; Kharouba & Wolkovich, 2020), but overcoming the empirical-theoretical disconnect in phenological studies may first require an improved mechanistic framework to help elucidate how mismatching occurs (Takimoto & Sato, 2020).

The match-mismatch hypothesis presents mismatching as the disrupted interaction between consumer demands and resource availability (Cushing, 1990). Most empirical studies categorize individuals or populations as 'matched' or 'mismatched' depending on the synchrony between the timing of a single life-history event and resource availability (Cushing, 1974; Visser et al., 1998). Contrary to this categorization, however, both resource availability and consumer demands vary over time, and being 'matched' does not guarantee that consumers have sufficient food (Saalfeld et al., 2019; Keogan et al., 2020). Rather, changes to continuous resource characteristics like quantity (i.e., biomass) and quality (i.e., per-capita size) directly affect consumer fitness, but the effects of these factors are rarely measured in studies of mismatching.

Moreover, energetic demand changes throughout an individual's life (Yang & Rudolf, 2010),

meaning that an individual's sensitivity to resource availability is not constant (Dunn et al.,

2011). Viewing mismatching simply as asynchrony in time, instead of as the disrupted

interaction between consumer demand and resource availability, can obscure the cumulative

effects of mismatching and mask population-level consequences (Yang & Rudolf, 2010; Kerby

et al., 2012). Although many conceptual models have been proposed to address this potential

issue, a more robust methodology to model mismatching in relation to the interaction of

consumer demands and resources is still lacking (Chmura et al., 2019; Visser & Gienapp, 2019).

Incorporating both age-structured consumer demand and multiple facets of resource

availability into mismatch models likely requires a re-examination of our statistical concept of

mismatching (Visser & Both, 2005; Kellermann & van Riper, 2015). Phenologies are generally

modelled as frequency curves on a temporal axis (Fig. 1; Cushing, 1974; Visser et al., 1998),

whereby match is estimated as the difference in peak dates (i.e., date models) or proportion of

overlapping area (i.e., overlap models). Both date and overlap models have been criticized in the

literature, however (Lindén, 2018; Ramakers et al., 2020). Furthermore, while date and overlap

models agree if consumer and resource curves are symmetrical (Fig. 1a,b), date models can be

biased when phenologies are skewed or multimodal, or in cases of low resource availability (Fig.

1c,d,e). Because overlap models account for the full interaction of consumer demand and

resource availability posed in the match-mismatch hypothesis (Kerby et al., 2012), they may be

better able to capture the mechanism of mismatching. Even so, overlap models have received

mixed support in empirical tests (Ramakers et al., 2020).

Figure 1. Peak dates (vertical lines) and frequency curves (phenologies) of consumers (solid) and

resources (dashed). Difference in peak dates and peak overlap (shaded area; percent area under

the curve) models are approximately equivalent when both the consumer (solid) and resource

(dashed) curves are symmetrical (a, b). In this case, mismatching is a function of temporal

displacement. However, date and overlap model estimates differ when either curve is skewed (c),

the consumer phenology is multimodal (d), or the curves are aligned but have low overlapping

area due to reduced resource abundance (e).

The inconsistent performance of overlap models may result from an inaccurate

representation of consumer demand. Existing peak overlap models estimate consumer demand

from a single life-history event or timepoint in development, such as when individual growth

rates are maximized (Fig. 2a; Leung et al., 2018). This approach, however, ignores demand prior

to or following this peak, and results in a less realistic demand curve (Fig. 2b; Kerby et al., 2012;

Lindén, 2018). Because animals require increasing energy as they develop, their sensitivity to the

low resource availability associated with mismatching is likely to change over time. As a result,

measuring the consequences of a mismatch from one timepoint could shroud cumulative effects

(Yang & Rudolf, 2010) and mask differences among individuals of differing ages (Reed et al.,

2013). The growing availability of metabolic data and advances in Bayesian survival analyses

now allow for the direct simulation of the age- or stage-specific effects of mismatching. By

modelling cumulative consumer demand as a function of the population age-structure, a 'whole

demand' model incorporates the increasing metabolic demands of individuals as they age (Fig.

2c). As a result, the whole demand curve quantifies overlap at the demand curve's upper tail

when per capita consumer demands are likely greatest (Fig. 2d ; Kerby et al., 2012). Accurately

modelling consumer demand and competing factors of resource availability may be key to

defining *how* mismatching is occurring and *when* it should matter to populations.

Figure 2. The peak demand model estimates consumer phenologies from the daily frequency of

individuals at a single point in development (e.g., peak growth rate; a). Fitting a curve to pseudo-

discrete data of this kind results in a simplified curve (b). However, since resource demand

increases throughout development (c), including the cumulative demand of all individuals for

each day of the season produces a curve with well-defined tails (d). Filled circles are time points

in an individual's development considered by the model. Circle size corresponds with

hypothetical energy requirements at each timepoint. Curves are from predictions from a

generalized additive model (GAM) performed on data collected in our study (see Methods and

Results) 2011.

Migratory birds provide a powerful avenue for re-examining the effects of mismatches

under this new framework. Long-distance migrants represent some of the canonical examples of

mismatches because of their use of endogenous cues to time their migrations and reproduction

(Both & Visser, 2001), and their reliance on seasonal resource pulses to achieve rapid offspring

growth (Schekkerman & Visser, 2001). Yet, while many studies have identified individual-level

fitness effects resulting from mismatches, few have found corresponding population-level

consequences (Visser & Both, 2005; Dunn et al., 2011). Hudsonian Godwits (*Limosa*

haemastica; hereafter, 'godwits') are a case-in-point. Godwits breed in three disjunct populations

spread across the Nearctic (Walker et al., 2011). Like other shorebird species (Kwon et al.,

2019), the godwits breeding in Alaska have kept pace with recent phenological changes in peak

resource availability on their breeding grounds while those breeding in Hudson Bay have not

(Senner, 2012). Despite the mismatch affecting the survival of godwit chicks in Hudson Bay,

there have been few apparent population-level consequences there (Senner et al., 2017).

Furthermore, much of the interannual variation in the fledging rates of Alaskan godwits is not

explained by predation or density-dependent processes (Senner et al., 2017; Swift et al., 2017a,

2018; Wilde et al., in revision). This interannual variation may instead result from a potential

correlation between early snowmelt and low seasonal godwit fledging rates, suggesting that

mismatching may be occurring and having demographic consequences (Saalfeld et al., 2019).

Updating our conceptualization of mismatches may document the effects of mismatching

our previous attempts based on the categorical view of mismatching have missed. Therefore, we

investigated how dynamic consumer demand and resource characteristics interact to drive

mismatching in the Alaskan population of godwits. We developed mechanistic models that

integrate metabolic and resource availability information at the individual- and population-levels.

We first explored how the timing, abundance, and quality of resources have changed over time

for godwits. Then, we investigated the effects of invertebrate abundance and size on the growth

and survival of godwit chicks. We hypothesized that mismatching affects individual fitness

differently throughout development and predicted that more abundant and larger prey would

improve chick growth and survival, with their effect increasing with age. Lastly, we investigated

the influence mismatching has on godwit population dynamics. We hypothesized that

mismatching in godwits is simultaneously a function of both consumer demand and resources.

We therefore predicted that more accurate quantification of both the consumer and resource

curves would explain population-level effects better than alternatives. Identifying how resources

interact with consumer demands will provide evidence for the mechanism underlying

mismatches and help better connect mismatching to demographic process.

Methods

Study area and godwit chick monitoring

During 2009 – 2011, 2014 – 2016, and 2019, we monitored godwits on two plots – North (550

Ha) and South (120 Ha) – near Beluga River, Alaska (61.21°N, 151.03°W; hereafter, 'Beluga';

Supporting information, Fig. S1). Both plots consist of freshwater ponds and black spruce

outcroppings (*Picea mariana*) dominated by dwarf shrub and graminoids surrounded by boreal

forest (Swift et al., 2017a, 2017b).

Each season (early-May to mid-July: $\mu = 78$ days), we censused both plots for godwit

nests (~ 5 nests per km^2, Swift et al., 2017b). We located an average of 23 nests per year (range:

11 – 33). For each found nest, we estimated hatch date and monitored its survival every 2 – 3

days. We moved to daily checks once eggs showed starring or pipping. We captured newly

hatched chicks and uniquely marked each with a leg-flag and USGS metal band. Despite missing

some nests prior to hatch, we are confident that we found all broods and estimated their hatch

dates each year because of the size of the study area and the conspicuousness of godwit broods.

We monitored the survival of 1 – 2 chicks chosen randomly from each brood (range = 7 –

23 per season). We attached a 0.62g VHF-radio transmitter (Holohil Systems Ltd.) above the

uropygial gland. We relocated each radioed chick every 2 – 3 days and attempted to recapture

them weekly to reapply glue and measure their body mass to the nearest gram. Godwits are fully

flight capable, or 'fledged', after ~28 days (Walker et al., 2011). However, the 21-day (range: 17

– 30 d) lifespan of our radios meant we considered those surviving to 21 days to have fledged.

We confirmed mortalities when possible and assumed that chicks had died after three

consecutive failed location attempts.

Resource monitoring

We monitored the abundance and body size of invertebrates for an average of 67 days (range =

61 – 78) in all years with godwit monitoring and for 38 and 5 days during the shortened seasons

of 2012 and 2017, respectively. We collected invertebrates each day along two, 100-m transects

of five traps within godwit breeding habitat (Senner et al., 2017). We used two trap styles: pitfall

traps (10 × 15 cm) filled with 10 cm of 75% ethanol from 2009 – 2012, and modified malaise

traps (see Leung et al., 2018) filled with 3 cm of 75% ethanol from 2014 – 2019. We cleared and

replenished traps every 24 hours.

We identified invertebrates to Order and measured body-lengths to the nearest 0.5 mm.

We converted lengths to dry mass using published, taxon specific length-weight relationships

(Ganihar, 1997; Rogers et al., 1977). Passive traps have been shown to be a good proxy of

resource availability to foraging shorebird chicks (McKinnon et al., 2012).

Statistical Analyses

Interannual resource variation

To examine resource availability over the course of our study, we tested for interannual

differences in the (1) date of the seasonal peak, (2) daily biomass (transect^{-1} day^{-1}; mg), and (3)

daily median body size (i.e., per-capita mass; mg). We excluded 2012 and 2017 from our

analyses of seasonal peaks but included them in tests of daily biomass and body size. We

estimated seasonal peaks using the first derivative of predicted curves of daily invertebrate

biomass within each season. Because godwit chicks are gape limited and rarely consume larval

invertebrates, we subset our data to include only adult invertebrates with lengths of 1.5 – 9 mm

(Schekkerman & Boele, 2009). We used dry mass, not length, for our response variables to

model changes in the cumulative and per-capita energy content available to chicks. We then built

separate mixed-effect models to estimate shifts in peak dates, daily biomass, and median body

size, with Julian date as a random slope and random intercept, using the *lmer* function (package

'lme4', Bates et al. 2015) in the R programming environment (v4.0.3, R Core Team 2020).

To identify potential changes in the composition of the invertebrate assemblage, we

repeated the above analyses with each of the six Orders that comprised 79.9% of all observed

invertebrates – Acari (8.0%), Araneae (19.9%), Coleoptera (5.6%), Diptera (34.8%), Hemiptera

(2.4%), and Hymenoptera (9.2%; Supporting information, Fig. S2). We excluded Collembola

(19%) from these analyses as they were imperfectly recorded from 2009 – 2012. We

standardized response variables according to Gelman (2008), but report coefficients in their

original units throughout the text. We considered variables whose 95% confidence intervals did

not include zero as biologically relevant.

Chick growth and body condition

We modelled chick growth with a logistic growth function using the 'nlme' package (Pinhiero et

al., 2019) to predict the age-specific mass of chicks (Ricklefs, 1968; Senner et al., 2017). We set

the asymptotic mass to the population's mean adult mass (249 g; Senner et al., 2017). Next, we

developed separate growth models with chick ID as a random intercept, and constant or yearly

growth coefficient and inflection points (Pinhiero & Bates, 2019). We performed 100 iterations

for each model and included site-specific estimates from Senner et al. (2017) as starting values.

We compared 12 candidate models using Akaike's Information Criterion scores (AIC_c) corrected

for small sample sizes (Burnham & Anderson, 2002). We used the model with the lowest AIC_c

score to calculate the body-condition index (BCI) for each recaptured individual by dividing the

observed weight gain since last capture by the curve-predicted weight gain over the same time.

To investigate how resource characteristics influenced chick growth, we modelled BCI

from resource abundance and quality in all years with godwit monitoring except 2014, which

lacked recaptures. We first determined the timescale over which predictors influenced BCI – day

of, 1-day, 3-day, 7-day average – using AIC_c scores. We then built a generalized additive model

(GAM) with a gaussian error term that included (1) daily invertebrate biomass and (2) median

invertebrate body size as fixed effects (package 'gamlss'; Rigby and Stasinopoulos 2005). We

also included (3) hatch date as a blocking variable, random intercepts for (4) study year and (5)

brood, and a cubic spline for (6) chick age. Again, we compared models by AIC_c scores. When

no model had a weight (w_i) > 0.90, we used model averaging within the 'MuMIn' package and

report conditional average coefficients (Bartoń, 2015).

Effect of resources on survival: constant or age-varying?

To determine how invertebrate biomass or body size affected daily chick survival, we built a

Bayesian hierarchical survival model. We constructed daily encounter histories for all

individuals, beginning with an individual's hatch date and ending with their expected fledging

date. Because we assumed chicks not located for three consecutive days were dead, we included

two days of unknown fate to allow for Markov chain Monte Carlo (MCMC) prediction. We

modelled encounter histories as a Bernoulli variable and assumed fates were known.

In the second portion of our model, we incorporated parameters hypothesized to

influence chick survival. We constructed a logit-linear mixed model to estimate the additive

effects of daily invertebrate biomass, median invertebrate body size, hatch date, and chick age,

along with random intercepts for each brood ID ($n = 98$), study year ($n = 7$), and study plot ($n =$

2). We averaged our continuous parameters across 3-day periods and standardized all variables.

To test whether the effects of invertebrate body size or daily biomass varied with chick age — a

proxy for metabolism — we built separate models with interactive terms between chick age and

either median invertebrate body size or daily invertebrate biomass. We compared age-interaction

models using deviance information criterion (DIC) and included the interaction from the model

with the lower DIC score in all further tests. We chose diffuse priors for all our predictors

(Normal(0, τ)) and constrained random intercepts close to 0 (mean = N(0, 1000), SD =

Uniform(0, 25)).

To identify the top model, we performed model selection using the indicator-variable

approach (Converse et al., 2013; Link & Barker, 2006). We assigned a Bernoulli variable

(weights) with a 0.5 prior to each predictor to model its inclusion (1) or absence (0) from each

MCMC sample. We maintained an equal number of parameters across samples by fixing the

model variance, $\tau = K *$ Gamma(3.29, 7.8), for all parameters, where K is the number of

parameters (Link & Barker, 2006). The posterior mean of the weight indicator is evidence for

inclusion in the model. We calculated Bayes factors (BF) from predictor weights (Link &

Barker, 2006). We included predictors with BF > 3 in our top model along with their random

intercepts. When an interaction term was chosen, both additive terms were also included.

We constructed models of daily chick survival in R with the 'runjags' and 'rjags'

packages (JAGS 4.1.0; Plummer 2012, 2013; Denwood 2016). Models accessed three parallel

chains to perform 5,000 iterations. We removed 600 and 1,000 iterations for adaptation and

burn-in, respectively, with a one-third thinning factor. We assessed model performance based on

the values of the Gelman-Rubin statistic < 1.1 and chain mixing (Gelman 1996). For all tests, we

report the beta coefficients in logit-form, 95% credible interval, and Bayesian p-value

(probability of slope $\neq$ 0).

Population match and reproductive success

To quantify population-level mismatching, we built resource and consumer demand curves for

each season. Additionally, we built competing demand curves from the 'peak demand' and

'whole demand' conceptual models (Fig. 2) to test the for the interaction of dynamic consumer

demand and resource availability. (1) *Peak demand*: Following Kwon et al. (2019), we calculated

the number of all hatched godwit chicks expected to be 11-days old (i.e., age of peak growth

rate; Senner et al. 2017) for each day of the season, and converted both the daily values of

invertebrate biomass (hereafter, 'resource curve') and counts of 11-day old chicks to their

seasonal proportions. (2) *Whole demand*: For this curve, we multiplied the maximum number of

chicks of each age per day of the season by age-specific estimates of resting metabolic rate in

godwit chicks taken from Williams et al. (2007). Resting metabolic rate approximates the

amount of energy individuals use to maintain homeostasis and therefore represents an

individual's minimum energetic requirement independent of other factors (i.e., thermal

environment). We then estimated the cumulative energetic requirements (kJ d^{-1}, kilojoules per

day) of all chicks per day of the season and converted these to seasonal proportions to produce

the whole demand curve.

309   We modelled the shape of the peak demand, whole demand, and resource curves using

separate GAMs with a quadratic time function – day + day^2 (Kwon et al. 2019). We restricted the

analyses to 10 May – 10 July for comparison among study years. We approximated error terms

as a gaussian distribution ($\sim N[\mu,\sigma]$) and zero-inflated beta distributions ($\sim zBeta[z|\alpha, \beta]$) for the

peak demand and whole demand curves, respectively, and a beta distribution ($\sim Beta[\alpha, \beta]$) for

the resource curve, all with logit-link functions. We fit the resource curve with a penalized spline

(k = 10) to estimate mean predicted values for each day of the season while capturing the

modality of the resource curve (Vatka et al., 2016). We then estimated the degree of overlap

between the peak demand or whole demand curves and the resource curve by calculating the

proportional area overlap using the *integrate.xy* function ('sfsmisc', Maechler 2020). We also

estimated the (3) 'curve height' in each season (i.e., cumulative resource availability) from the

area under the resource curve. Lastly, we calculated the (4) 'difference in dates' (i.e., synchrony)

between the resource and peak demand curves in each season from the point at which each

curve's derivative was zero.

To determine how mismatching affected godwit reproductive success, we built four

univariate linear models relating the different measures of mismatching to fledging rates – (1)

peak demand, (2) whole demand, (3) difference in dates, and (4) curve height. We extrapolated

daily survival rate (DSR) estimates from our global Bayesian model to 28 days with the

associated error using the Delta method (Powell 2007). We compared among the four models by

calculating (1) model weights from their AIC_c scores and (2) the proportion of the variation in

fledging rates they explained (i.e., R^2).

Results

We located 142 godwit nests from 2009 – 2019, of which 128 survived to hatch. We individually

marked 349 chicks (2009 – 2011, $n = 195$; 2014 – 2016, $n = 106$; 2019, $n = 48$) and attached

radios to 128 chicks from 102 distinct broods. We monitored radioed chicks for 11.1 ± 10.9 d

and recaptured them 1.5 times ± 0.83 ($n = 103$).

Interannual changes in resources

We recorded the body-lengths of 69,598 adult invertebrates across 14 orders, 41,298 of which

were potential godwit prey (i.e., 1.5 – 9 mm in length). Sample days showed wide variation in

biomass ($\bar{x} = 132.9$ mg, range: 0 – 948.4 mg) and median invertebrate body size ($\bar{x} = 1.5$ mg,

range: 0.2 – 13.4 mg). We found no interannual shift in the predicted peak dates of all

invertebrates ($\beta = -1.68 \pm 3.08$ d, 95% Confidence Interval: -3.34, 5.50 d) or individual orders

(Fig. 3, left). However, both daily invertebrate biomass ($\beta = -2.49 \pm 0.50$ mg, 95% CI: -3.49, -

1.51; Fig. 3, center) and median invertebrate body size (β = -0.33 ± 0.03 mg, 95% CI: -0.028, -

0.37; Fig. 3, right) decreased over the course of the study. At the order level, only Acari became

more abundant over time (β = 0.20 ± 0.02 mg, 95% CI = 0.15, 0.25). Meanwhile, Araneae (β = -

0.67 ± 0.09 mg, 95% CI = -0.49,-0.85), Diptera (β = -0.24 ± 0.02 mg, 95% CI = -0.20, -0.29),

and Hemiptera (β = -0.23 ± 0.06 mg, 95% CI = -0.10, -0.35) showed consistent shrinkage.

368   Figure 3. Interannual changes of within season peak timing (left), observed daily invertebrate

369   biomass (center), and median invertebrate body size (right) of six common Orders and the

370   invertebrate assemblage overall. Linear regression estimates are shown as hollow circles, with

371   95% confidence intervals shown as horizontal lines. Variables with no consistent effect had

372   intervals that crossed zero (grey line).

Chick growth and body condition

We modeled godwit chick growth from 179 mass-at-capture estimates. Chick growth did not

differ among years, and our top-performing growth function included both a constant logistic

coefficient ($K = 0.13 \pm 4.2 \times 10^{-3}$) and inflection point ($T_i = 17.5 \pm 0.5$ days; Supporting

information, Table S1).

The fit of our global model was highest with 7-day averaged covariates and no random

effects (Supplementary Materials Appendix A, Table S2). Our top model explaining chick BCI

($n = 89$) included invertebrate biomass and our blocking variable, hatch date, with a smoothed

age effect ($w_i = 0.75$; Supplementary Materials Appendix A, Table S3). Chick growth improved

with higher invertebrate biomasses ($\beta = 1.8 \times 10^{-4} \pm 3.8 \times 10^{-5}$ mg^{-1}, CI: 1.2×10^{-4}, 2.8×10^{-4};

Fig. 4a), but decreased with later hatch dates ($\beta = -0.013 \pm 0.003$ d^{-1}, CI: -0.0053, -0.019; Fig.

4b). Invertebrate body size had no consistent effect (Supporting information, Fig. S3). Chicks

therefore grew as well as or better than expected (e.g., BCI ≥ 1) following weeks with

invertebrate biomasses in the upper 15% of those observed or if they hatched before 5 June.

Effect of resources on survival: constant or age-varying?

Of the 128 godwit chicks in our study, we excluded 6 due to human-caused mortality or

equipment failure at deployment. The mean DSR of the remaining 122 chicks was $86 \pm 24\%$,

meaning that $19.2 \pm 33\%$ survived to fledge, although this varied among years and broods

(Supporting information, Table S4).

The model with an age-varying effect of invertebrate body size (DIC = 283.0)

outperformed the model with an age-varying effect of invertebrate biomass (DIC = 287.9). We

therefore used the former in our subsequent tests. The constant effect of invertebrate biomass and

the age-varying invertebrate size effect had 79% and 85% posterior inclusion probabilities,

respectively (Table 1). We also included constant effects of age and invertebrate size to

accompany the interaction term.

Chick survival improved with higher invertebrate biomasses and larger invertebrate body

size, and the latter effect increased throughout development (Table 2). Each 1% increase in daily

invertebrate biomass (+ 1.5 mg) improved daily chick survival by 0.66% (Fig. 5a), while each

1% increase in median invertebrate body size (+ 0.06 mg) led to a 1.02% increase in daily chick

survival. This 'size' effect then grew by 2.2% with each day that a chick survived (Fig. 5b). Age

itself, however, had no consistent effect on chick survival.

Population match and reproductive success

The model fit for the whole demand curve (AIC_c = -300.1) was 25.7-times better than the peak

demand curve (AIC_c = –248.7). Godwits had, on average, 51.9 ± 9.2% overlap with resource

phenology according to the peak demand model, but 44.7 ± 11.6% overlap according to the

whole demand model. Years also differed in curve height ($\bar{x}$ = 8800 ± 3668 mg) and the

difference between the peak dates of the resource and demand curves ($\bar{x}$ = 14.7 ± 16.36 d).

Godwit fledging rates varied among study years (Supporting information, Table S5) but

were lowest in 2014 and 2015. Models differed in their ability to explain population-level

reproductive success but the whole demand model was best supported (Supporting information,

Table S6). The whole demand model explained 55% of the variation in godwit fledging rates (β

= 1.19 ± 0.41; $R^2_{adj.}$ = 0.55; w_i = 0.43; Fig. 6a; Supporting information, Fig. S4). The difference

in dates model performed similarly well (β = -0.68 ± 0.27; $R^2_{adj.}$ = 0.48; w_i = 0.36; Fig. 6b) but

was 7% less likely to be the top model. Both the curve height (β = 2.49 ± 1.44; $R^2_{adj.}$ = 0.25; w_i =

0.10; Fig. 6c) and peak demand overlap models ($\beta = 1.00 \pm 0.56$; $R^2_{adj.} = 0.26$; $w_i = 0.11$; Fig. 6d;

Supporting information, Fig. S5) were unlikely to be the top model given their low model

weights and the low amounts of interannual variation in fledging rates explained by either.

Discussion

The disconnect between empirical studies and the theoretical predictions of the match-mismatch

hypothesis casts doubt upon the risks climate change-induced phenological mismatches pose to

consumer populations (Visser & Gienapp, 2019; Keogan et al., 2020). To remedy this gap and

connect mismatches to demographic processes, Kharouba & Wolkovich (2020) urged

researchers to define pre-climate change baselines, collect per-capita data on resources and

consumers, and test competing biological mechanisms. We developed mismatch models aimed at

fulfilling these recommendations while adopting an ontogenetic view of consumer demand.

Using this approach, we built upon the findings of Senner et al. (2017) and identified heretofore

undetected individual- and population-level fitness effects of mismatching in the Alaskan

breeding population of Hudsonian godwits. Our study joins the growing literature suggesting that

mismatches do not fall neatly into a 'matched' or 'mismatched' paradigm (Keogan et al., 2020;

Simmonds et al., 2020). Instead, models built around the underlying biological mechanisms

connecting consumers and resources are key to clarifying how mismatching affects consumer

fitness (Takimoto & Sato, 2020).

More than mistiming: the tandem drivers of resource availability

We found that resources affected godwit chick survival in two distinct ways: first, periods with

reduced resource abundance resulted in poorer growth and lower survival and, second, access to

larger invertebrates was increasingly important to the survival of older chicks. Our findings

differ from those of previous godwit studies, which found no effects of limited resource

availability in the Alaskan godwit breeding population (Senner et al., 2017; Wilde et al., in

revision). While these studies did not investigate the influence of invertebrate body size on

godwit chicks, our contradictory conclusions likely stem from our use here of hierarchical

models that easily approximate time-varying effects on survival (Royle & Dorazio, 2009).

Increasing energetic demands throughout ontogeny mean that the effects of resource limitation

are unlikely to be constant over an individual's lifetime (Yang & Rudolf, 2010; Takimoto &

Sato, 2020). Therefore, models that accommodate variable predictor effects may be key to

clarifying how resource characteristics affect consumer fitness.

Godwit chicks had improved growth and survival following periods with high resource

abundance. Having adequate resources during energetically stressful periods is a major driver of

animal fitness (Bastille-Rousseau et al., 2015), especially in seasonal environments (McKinnon

et al., 2012). Given their high energetic demands and rapid development, chicks of shorebird

species across the Arctic have exhibited survival costs following reduced resource abundance

(Schekkerman et al., 2003; Saalfeld et al., 2019). Godwit chicks in this study had 3 – 75% higher

body condition indices and 17% higher daily survival probabilities, on average, during periods of

higher-than-average invertebrate abundance. Importantly, while we also detected effects of hatch

date (i.e., phenology) on chick growth, these did not translate into an effect on survival. Our

results therefore suggest that relating fitness measures to resource availability captures the effects

of mismatching while defining its specific costs in biological terms (Dunn et al., 2011).

In addition to the effects of resource abundance, the quality (i.e., median body size) of

invertebrates became increasingly important as godwit chicks aged. Optimal foraging theory

predicts that consumers should select resources with the most energy content relative to foraging

effort (Krebs et al., 1977). Chicks of black-tailed godwits (*Limosa limosa*), for instance,

prioritize the rapid intake of small prey early in life, but switch to the slower intake of larger prey

as they grow older (Schekkerman & Boele, 2009). While we did not observe foraging behaviors

directly, we hypothesize that Hudsonian godwit chicks may make a similar transition. In fact,

increasing selection of larger prey could explain the especially high costs of poor resource

quality for older chicks. We found that periods of below-average prey size resulted in 29% lower

survival for chicks below 5-days of age, but 50% lower survival for chicks older than 11-days.

Changes in resource quality, though rarely explored in the context of mismatches, can enact

strong selection on consumer populations (Keogan et al., 2020; Yang et al., 2020). Because some

individuals will encounter high-quality conditions in years when they are 'mismatched' (Kerby

et al., 2012), accounting for the effects of multiple factors of resource availability could improve

our ability to document the true effects of mismatching.

Taken together, the additive effects of resource quantity and quality are likely to worsen

in Beluga given the changes we observed in the invertebrate community. Climate-induced

reductions in resource availability are common across terrestrial and marine systems (Bowden et

al., 2015; Weterings et al., 2018). Arctic invertebrates, in particular, are simultaneously emerging

earlier (Høye et al., 2007), becoming less abundant (van Klink et al., 2020), and smaller in size

(Bowden et al., 2015; Jonsson et al., 2015) with increasing spring temperatures. Here, we found

a linear decrease in the daily abundance (-2%) and body size (-5%) of invertebrates, but no

change in the date of peak occurrence of invertebrates over the course of our study. Although we

did not detect a linear shift in the timing of the resource peak, this may relate to the occurrence of

opposing trends in abundance during the early and late portions of the godwit breeding season. In

a post-hoc test, we found that days during the godwit nest incubation period (16 May – 6 June)

from 2014 – 2019 had 83% higher invertebrate biomass than those from 2009 – 2012, but 41%

lower biomasses on days during the chick-rearing period (6 June – 4 July). Meanwhile,

invertebrate body size was 42 – 72% smaller in the later period. Therefore, should these trends

continue, developing godwit chicks may face increasingly untenable conditions as food becomes

both less abundant and of poorer quality (i.e., smaller size). More broadly, our results suggest

that resource timing, quality, and quantity can act as concomitant drivers of phenological

mismatches (Rollins & Benard, 2020), and that their effects may be most apparent when placed

in the context of the consumer life cycle (Yang et al., 2020).

Modelling the demand-resource interaction clarifies the population effects of mismatching

Variation in godwit reproductive success at the population level was best explained by our whole

demand model of mismatching, although the simpler difference in dates model also performed

well. Estimates from overlap and dates models do often correlate (Ramakers et al., 2020), but

may perform differently depending on a species' life history and trophic specialization (Miller-

Rushing et al., 2010). Thus, while difference in dates models may suffice for godwits and other

species with narrow, synchronous, breeding phenologies or those that rely on singular resource

pulses (Miller-Rushing et al., 2010), they would likely perform poorly in species with highly

variable nest initiation dates or those capable of multiple nesting events (Phillimore et al., 2016).

Because overlap models account for both synchrony and the magnitude of interacting consumer-

resource pairs, they are more likely to capture mismatching as a disrupted interaction (Kerby et

al., 2012). Overlap models are therefore likely more generalizable, but using both overlap and

difference in dates models could help when exploring how mismatching occurs on a case-by-

case basis (Kellermann & van Riper, 2015).

Not all overlap models are equivalent, however. Overlap models have received mixed

support (Ramakers et al., 2020), but their ability to accurately quantify mismatching at the tails

of the consumer curve has been suggested as an important component of their effectiveness

(Kerby et al., 2012). Accordingly, whereas our peak demand model performed relatively poorly

and explained only 21% variation in godwit reproductive success, our whole demand model had

25-fold better model fit than the peak demand model and explained 55% of the variation in

fledging rates. This difference likely stems from the inability of the peak demand model to

accurately capture consumer demand at the upper (i.e., right-hand) tail of the consumer curve,

corresponding to the period when individual-level consumer demand is greatest. Our results

therefore show that incorporating additional nuance into the statistical concept of consumer

phenologies can greatly improve overlap models (Lindén, 2018).

The need to accurately identify mismatches is made most clear by the accumulating

evidence for variable and non-linear responses by consumer populations to mismatching (Visser

& Both, 2005; Phillimore et al., 2016). So called 'tipping points' – thresholds past which an

effect abruptly changes (Latty & Dakos, 2019) – buffer consumer populations from the negative

impacts of moderate mismatching and may contribute to the lack of consistent responses to

mismatching across consumer populations (Simmonds et al., 2020). In godwits, we found that

greater population-level mismatching consistently drove poorer fledging success, but that there

may be thresholds past which the effects are most severe. For instance, godwits experienced

~19% poorer overlap and ~28 days greater mismatching in 2014 and 2015 compared to the long-

term average. Mismatching on this scale resulted in 24% lower fledging rates and near complete

reproductive failure for the population. Similarly low fledging rates for Hudson Bay breeding

godwits, which are mismatched by 11-days on-average (Senner et al., 2017), suggests that for

godwits, this tipping point may exist when populations are mismatched by more than ~10 days or

have less than 40% overlap with the resource curve.

Importantly, though, the 2014 and 2015 seasons in Beluga coincided with a period of

anomalous and prolonged near-surface warming in the northeastern Pacific called the 'blob'

(Cavole et al., 2016). Thus, while the conditions in these atypical years may provide useful

insights into potential outcomes of a warming climate on coastal communities in the region

(Auth et al., 2018), mismatches of this magnitude are unlikely to become the norm. Beluga

godwits have been able to advance their timing of migration and reproduction in response to

recent long-term, linear, warming trends (Senner 2012; Senner et al., 2017) and may therefore be

able to do so in the future. Nonetheless, significant spring warming and earlier snow

disappearance dates projected for the North American sub-Arctic mean that godwits and other

migratory populations may soon face accelerating, potentially non-linear, warming (Littell et al.,

2018; Lader et al., 2020). Quantifying the strength and effects of mismatching in real time will

thus be crucial for conservation going forward (Simmonds et al., 2020).

Conclusions

By modelling the unspoken assumptions of the match-mismatch hypothesis, we stand to adopt a

more powerful definition of mismatching in biological terms and, in so doing, be better able to

identify the circumstances under which consumer populations perform poorly. Our work also

illustrates the role of ontogeny in shaping an individual's changing response to resource

availability over time (Yang & Rudolf, 2010), and helps explain the empirical-theoretical

disconnect in phenological studies. Importantly, our models are transferrable to other systems,

whereby remotely sensed indices and knowledge of a population's age-structure could

approximate resource availability and metabolic rate, respectively, when these data are otherwise

unavailable (Lumbierres et al., 2017). Finally, we show how treating mismatches as an outcome

of both demand and resource dynamics provides insight into the structure of individual-level

effects and the mechanism behind population-level responses (Takimoto & Sato, 2020).

Replacing the categorical 'matched or mismatched' view of mismatching with one that explicitly

recognizes the underlying mechanism may be critical to monitoring and conserving animal

populations in an uncertain future.

Acknowledgements

We thank the Cook Inlet Region Inc. for permitting us access to their lands. We thank Drs. Sarah

Converse and Emily Weiser for analytical guidance. Funding was provided by the Association of

Field Ornithologists, Wilson Ornithological Society, American Ornithological Society, Arctic

Audubon Society, Cornell Lab of Ornithology, Athena Fund at the Cornell Lab of Ornithology,

University of South Carolina, Faucett Family Foundation, the David and Lucile Packard

Foundation, National Science Foundation (PCE-1110444 and DGE-1144153), and the U.S. Fish

and Wildlife Service (4074, 5147). All procedures met the ethical standards of the University of

South Carolina (2449-101417-042219), Alaska Department of Fish and Game (20-024), and

USGS (24191). Any use of trade, product, or firm names is for descriptive purposes only and

does not imply endorsement by the U.S. Government. The authors declare no conflict of interest.

Author Contributions

LRW and NRS conceived of the study. All authors collected field data. LRW analyzed the data

and wrote the manuscript. All authors contributed to revisions.

Data Availability Statement

Data are deposited at Dryad (https://doi.org/10.5061/dryad.x69p8czh0) and computer code for all

analyses are available at github (http://doi.org/10.5281/zenodo.4298755).

References

Auth, T. D., Daly, E. A., Brodeur, R. D., & Fisher, J. L. (2018). Phenological and distributional

shifts in ichthyoplankton associated with recent warming in the northeast Pacific Ocean.

Global Change Biology, *24*(1), 259–272. https://doi.org/10.1111/gcb.13872

Bartoń, K. (2015). Multiple-Model Inference (Package 'MuMIn'). *R v. 3.3.2*.

Bastille-Rousseau, G., Potts, J. R., Schaefer, J. A., Lewis, M. A., Ellington, E. H., Rayl, N. D.,

Mahoney, S. P., & Murray, D. L. (2015). Unveiling trade-offs in resource selection of

migratory caribou using a mechanistic movement model of availability. *Ecography*,

38(10), 1049–1059. https://doi.org/10.1111/ecog.01305

Bates D. M. Maechler B. Bolker and S. Walker. (2015). Fitting linear mixed-effects models

using lme4. *Journal of Statistical Software*, *67*: 1–48.

Both, C., Asch, M. V., Bijlsma, R. G., Burg, A. B. V. D., & Visser, M. E. (2009). Climate

change and unequal phenological changes across four trophic levels: Constraints or

adaptations? *Journal of Animal Ecology*, *78*(1), 73–83. https://doi.org/10.1111/j.1365-

2656.2008.01458.x

Both, C., & Visser, M. E. (2001). Adjustment to climate change is constrained by arrival date in

a long-distance migrant bird. *Nature*, *411*(6835), 296–298.

https://doi.org/10.1038/35077063

Bowden, J. J., Eskildsen, A., Hansen, R. R., Olsen, K., Kurle, C. M., & Høye, T. T. (2015).

High-Arctic butterflies become smaller with rising temperatures. *Biology Letters*, *11*(10),

20150574. https://doi.org/10.1098/rsbl.2015.0574

Burnham, K. P., & Anderson, D. R. (2002). *Model selection and multimodel inference: A*

practical information–theoretic approach. Springer Science and Business Media.

Cavole, L. M., Demko, A. M., Diner, R. E., Giddings, A., Koester, I., Pagniello, C. M. L. S.,

Paulsen, M.-L., Ramirez-Valdez, A., Schwenck, S. M., Yen, N. K., Zill, M. E., & Franks,

P. J. S. (2016). Biological Impacts of the 2013–2015 Warm-Water Anomaly in the

Northeast Pacific: Winners, Losers, and the Future. *Oceanography*, *29*(2), 273–285.

Chmura, H. E., Kharouba, H. M., Ashander, J., Ehlman, S. M., Rivest, E. B., & Yang, L. H.

(2019). The mechanisms of phenology: The patterns and processes of phenological shifts.

Ecological Monographs, *89*(1), e01337. https://doi.org/10.1002/ecm.1337

Converse, S. J., Royle, J. A., Adler, P. H., Urbanek, R. P., & Barzen, J. A. (2013). A hierarchical

nest survival model integrating incomplete temporally varying covariates. *Ecology and*

Evolution, *3*(13), 4439–4447. https://doi.org/10.1002/ece3.822

Corkery, C. A., Nol, E., & Mckinnon, L. (2019). No effects of asynchrony between hatching and

peak food availability on chick growth in Semipalmated Plovers (Charadrius

semipalmatus) near Churchill, Manitoba. *Polar Biology*, *42*(3), 593–601.

https://doi.org/10.1007/s00300-019-02456-w

Cushing, D. H. (1974). The natural regulation of fish populations. In *Sea Fisheries Research* (pp.

399–412). F. R. Harden Jones, ed. https://ci.nii.ac.jp/naid/10022014307/#cit

Cushing, D. H. (1990). Plankton Production and Year-class Strength in Fish Populations: An

Update of the Match/Mismatch Hypothesis. In J. H. S. Blaxter & A. J. Southward (Eds.),

Advances in Marine Biology (Vol. 26, pp. 249–293). Academic Press.

https://doi.org/10.1016/S0065-2881(08)60202-3

Denwood, M. J. 2016. runjags: An R Package Providing Interface Utilities, Model Templates,

Parallel Computing Methods and Additional Distributions for MCMC Models in JAGS.

Journal of Statistical Software, 71(9), 1-25. doi:10.18637/jss.v071.i09

Dunn, P. O., Winkler, D. W., Whittingham, L. A., Hannon, S. J., & Robertson, R. J. (2011). A

test of the mismatch hypothesis: How is timing of reproduction related to food abundance

in an aerial insectivore? *Ecology*, *92*(2), 450–461. https://doi.org/10.1890/10-0478.1

Ganihar, S. R. (1997). Biomass estimates of terrestrial arthropods based on body length. *Journal*

of Biosciences, *22*(2), 219–224. https://doi.org/10.1007/BF02704734

Gelman, A. (2008). Scaling regression inputs by dividing by two standard deviations. *Statistics*

in Medicine, *27*(15), 2865–2873. https://doi.org/10.1002/sim.3107

Høye, T. T., Post, E., Meltofte, H., Schmidt, N. M., & Forchhammer, M. C. (2007). Rapid

advancement of spring in the High Arctic. *Current Biology*, *17*(12), R449–R451.

https://doi.org/10.1016/j.cub.2007.04.047

Jonsson, M., Hedström, P., Stenroth, K., Hotchkiss, E. R., Vasconcelos, F. R., Karlsson, J., &

Byström, P. (2015). Climate change modifies the size structure of assemblages of

emerging aquatic insects. *Freshwater Biology*, *60*(1), 78–88.

https://doi.org/10.1111/fwb.12468

Kellermann, J. L., & van Riper, C. (2015). Detecting mismatches of bird migration stopover and

tree phenology in response to changing climate. *Oecologia*, *178*(4), 1227–1238.

https://doi.org/10.1007/s00442-015-3293-7

Keogan, K., Lewis, S., Howells, R. J., Newell, M. A., Harris, M. P., Burthe, S., Phillips, R. A.,

Wanless, S., Phillimore, A. B., & Daunt, F. (2020). No evidence for fitness signatures

consistent with increasing trophic mismatch over 30 years in a population of European

shag Phalacrocorax aristotelis. *Journal of Animal Ecology*, *n/a*(n/a).

https://doi.org/10.1111/1365-2656.13376

Kerby, J. T., Wilmers, C. C., & Post, E. (2012). Climate change, phenology and the nature of

consumer–resource interactions: Advancing the match/mismatch hypothesis. In *Trait-*

mediated indirect interactions: Ecological and evolutionary perspectives (eds Ohgushi T,

Schmitz O, Holt R) (pp. 508–525). Cambridge University Press.

Kharouba, H. M., & Wolkovich, E. M. (2020). Disconnects between ecological theory and data

in phenological mismatch research. *Nature Climate Change*, *10*(5), 406–415.

https://doi.org/10.1038/s41558-020-0752-x

Krebs, C. J., Erichson, J. T., Weber, M. I., & Charnov, E. I. (1977). Optimal prey selection in the

great tit (Parus major). *Animal Behaviour*, *25*(1), 30–38. https://doi.org/10.1016/0003-

3472(77)90064-1

Kwon, E., Weiser, E. L., Lanctot, R. B., Brown, S. C., Gates, H. R., Gilchrist, G., Kendall, S. J.,

Lank, D. B., Liebezeit, J. R., McKinnon, L., Nol, E., Payer, D. C., Rausch, J., Rinella, D.

J., Saalfeld, S. T., Senner, N. R., Smith, P. A., Ward, D., Wisseman, R. W., &

Sandercock, B. K. (2019). Geographic variation in the intensity of warming and

675   phenological mismatch between Arctic shorebirds and invertebrates. *Ecological*

676   *Monographs*, *89*(4), e01383. https://doi.org/10.1002/ecm.1383

677 Lader, R., Walsh, J. E., Bhatt, U. S., & Bieniek., P. A. (2020). Anticipated changes to the snow

678   season in Alaska: Elevation dependency, timing and extremes. International Journal of

679   Climatology 40(1):169-187. *Journal of Climatology*, *40*(1), 169–187.

680 Lameris, T. K., Scholten, I., Bauer, S., Cobben, M. M. P., Ens, B. J., & Nolet, B. A. (2017).

681   Potential for an Arctic-breeding migratory bird to adjust spring migration phenology to

682   Arctic amplification. *Global Change Biology*, *23*(10), 4058–4067.

683   https://doi.org/10.1111/gcb.13684

684 Latty, T., & Dakos, V. (2019). The risk of threshold responses, tipping points, and cascading

685   failures in pollination systems. *Biodiversity and Conservation*, *28*(13), 3389–3406.

686   https://doi.org/10.1007/s10531-019-01844-2

687 Leung, M. C.-Y., Bolduc, E., Doyle, F. I., Reid, D. G., Gilbert, B. S., Kenney, A. J., Krebs, C. J.,

688   & Bêty, J. (2018). Phenology of hatching and food in low Arctic passerines and

689   shorebirds: Is there a mismatch? *Arctic Science*, *4*(4), 538–556.

690   https://doi.org/10.1139/as-2017-0054

691 Lindén, A. (2018). Adaptive and nonadaptive changes in phenological synchrony. *Proceedings*

692   *of the National Academy of Sciences*, *115*(20), 5057–5059.

693   https://doi.org/10.1073/pnas.1805698115

694 Link, W. A., & Barker, R. J. (2006). Model Weights and the Foundations of Multimodel

695   Inference. *Ecology*, *87*(10), 2626–2635. https://doi.org/10.1890/0012-

696   9658(2006)87[2626:MWATFO]2.0.CO;2

Littell, J. S., McAfee, S. A., & Hayward, G. D. (2018). Alaska snowpack response to climate

change: Statewide snowfall equivalent and snowpack water scenarios. Water 10(5):668.

Water, *10*(5), 668.

Lumbierres, M., Méndez, P., Bustamante, J., Soriguer, R., & Santamaría, L. (2017). Modeling

Biomass Production in Seasonal Wetlands Using MODIS NDVI Land Surface

Phenology. *Remote Sensing*, *9*(4), 392. https://doi.org/10.3390/rs9040392

Maechler M. (2020). sfsmisc: Utilities from 'Seminar fuer Statistik' ETH Zurich. R package

version 1.1-7.

McKinnon, L., M. Picotin, E. Bolduc, C. Juliet, & J. Bety. (2012). Timing of breeding, peak food

availability, and effects of mismatch on chick growth in birds nesting in the High Arctic.

Canadian Journal of Zoology, *90*, 961–971.

Miller-Rushing, A. J., Høye, T. T., Inouye, D. W., & Post, E. (2010). The effects of phenological

mismatches on demography. *Philosophical Transactions of the Royal Society B:*

Biological Sciences, *365*(1555), 3177–3186. https://doi.org/10.1098/rstb.2010.0148

Parmesan, C., & Yohe, G. (2003). A globally coherent fingerprint of climate change impacts

across natural systems. *Nature*, *421*(6918), 37–42. https://doi.org/10.1038/nature01286

Pinheiro J., D., Bates, S. DebRoy, D. Sarkar, and R Core Team. (2019). nlme: Linear and

Nonlinear Mixed Effects Models. R package version 3.1-140.

Phillimore, A. B., Leech, D. I., Pearce-Higgins, J. W., & Hadfield, J. D. (2016). Passerines may

be sufficiently plastic to track temperature-mediated shifts in optimum lay date. *Global*

Change Biology, *22*(10), 3259–3272. https://doi.org/10.1111/gcb.13302

Plummer, M. (2012). JAGS: just another Gibbs sampler. Version 3.3.0.

Plummer, M. (2013). Package rjags: Bayesian graphical models using MCMC. Version 3.10.

Powell L. A. (2007). Approximating variance of demographic parameters using the delta

method: a reference for avian biologists. *Condor, 109*: 949–954.

R Core Team (2020) R: A Language and Environment for Statistical Computing. RFoundation

for Statistical Computing, Vienna, Austria.

Ramakers, J. J. C., Gienapp, P., & Visser, M. E. (2020). Comparing two measures of

phenological synchrony in a predator–prey interaction: Simpler works better. *Journal of*

Animal Ecology, *89*(3), 745–756. https://doi.org/10.1111/1365-2656.13143

Reed, T. E., Jenouvrier, S., & Visser, M. E. (2013). Phenological mismatch strongly affects

individual fitness but not population demography in a woodland passerine. *Journal of*

Animal Ecology, *82*(1), 131–144. https://doi.org/10.1111/j.1365-2656.2012.02020.x

Rigby, R. A., & Stasinopoulos, D. M. (2005). Generalized additive models for location, scale

and shape. *Journal of the Royal Statistical Society*, *54*, 507–554.

Rogers, L. E., Buschbom, R. L., & Watson, C. R. (1977). Length-Weight Relationships of

Shrub-Steppe Invertebrates1. *Annals of the Entomological Society of America*, *70*(1), 51–

53. https://doi.org/10.1093/aesa/70.1.51

Rollins, H. B., & Benard, M. F. (2020). Challenges in predicting the outcome of competition

based on climate change-induced phenological and body size shifts. *Oecologia*, *193*(3),

749–759. https://doi.org/10.1007/s00442-020-04705-w

Royle, J. A., & Dorazio, R. M. (2009). *Hierarchical Modeling and Inference in Ecology, The*

Analysis of Data from Populations, Metapopulations and Communities.

Saalfeld, S. T., McEwen, D. C., Kesler, D. C., Butler, M. G., Cunningham, J. A., Doll, A. C.,

English, W. B., Gerik, D. E., Grond, K., Herzog, P., Hill, B. L., Lagassé, B. J., & Lanctot,

R. B. (2019). Phenological mismatch in Arctic-breeding shorebirds: Impact of snowmelt

and unpredictable weather conditions on food availability and chick growth. *Ecology and*

Evolution, *9*(11), 6693–6707. https://doi.org/10.1002/ece3.5248

Samplonius, J. M., Bartošová, L., Burgess, M. D., Bushuev, A. V., Eeva, T., Ivankina, E. V.,

Kerimov, A. B., Krams, I., Laaksonen, T., Mägi, M., Mänd, R., Potti, J., Török, J., Trnka,

M., Visser, M. E., Zang, H., & Both, C. (2018). Phenological sensitivity to climate

change is higher in resident than in migrant bird populations among European cavity

breeders. *Global Change Biology*, *24*(8), 3780–3790. https://doi.org/10.1111/gcb.14160

Schekkerman, H., & Boele, A. (2009). Foraging in precocial chicks of the black-tailed godwit

Limosa limosa: Vulnerability to weather and prey size. *Journal of Avian Biology*, *40*(4),

369–379. https://doi.org/10.1111/j.1600-048X.2008.04330.x

Schekkerman, H., Tulp, I., Piersma, T., & Visser, G. H. (2003). Mechanisms promoting higher

growth rate in arctic than in temperate shorebirds. *Oecologia*, *134*(3), 332–342.

https://doi.org/10.1007/s00442-002-1124-0

Schekkerman, H., & Visser, G. H. (2001). Prefledging Energy Requirements in Shorebirds:

Energetic Implications of Self-Feeding Precocial Development. *The Auk*, *118*(4), 944–

957. https://doi.org/10.1093/auk/118.4.944

Senner, N. R. (2012). One species but two patterns: Populations of the Hudsonian Godwit (

Limosa haemastica) differ in spring migration timing. *The Auk*, *129*(4), 670–682.

https://doi.org/10.1525/auk.2012.12029

Senner, N. R., Stager, M., & Sandercock, B. K. (2017). Ecological mismatches are moderated by

local conditions for two populations of a long-distance migratory bird. *Oikos*, *126*(1), 61–

72. https://doi.org/10.1111/oik.03325

Simmonds, E. G., Cole, E. F., Sheldon, B. C., & Coulson, T. (2020). Phenological asynchrony: A

ticking time-bomb for seemingly stable populations? *Ecology Letters*.

https://doi.org/10.1111/ele.13603

Swift, R. J., Rodewald, A. D., & Senner, N. R. (2017a). Environmental heterogeneity and biotic

interactions as potential drivers of spatial patterning of shorebird nests. *Landscape*

Ecology, *32*(8), 1689–1703. https://doi.org/10.1007/s10980-017-0536-5

Swift, R. J., Rodewald, A. D., & Senner, N. R. (2017b). Breeding habitat of a declining shorebird

in a changing environment. *Polar Biology*, *40*(9), 1777–1786.

https://doi.org/10.1007/s00300-017-2101-8

Swift, R. J., Rodewald, A. D., & Senner, N. R. (2018). Context-dependent costs and benefits of a

heterospecific nesting association. *Behavioral Ecology*, *29*(4), 974–983.

https://doi.org/10.1093/beheco/ary042

Takimoto, G., & Sato, T. (2020). Timing and duration of phenological resources: Toward a

mechanistic understanding of their impacts on community structure and ecosystem

processes in stream food chains. *Ecological Research*, *35*(3), 463–473.

https://doi.org/10.1111/1440-1703.12098

Thackeray, S. J., Henrys, P. A., Hemming, D., Bell, J. R., Botham, M. S., Burthe, S., Helaouet,

P., Johns, D. G., Jones, I. D., Leech, D. I., Mackay, E. B., Massimino, D., Atkinson, S.,

Bacon, P. J., Brereton, T. M., Carvalho, L., Clutton-Brock, T. H., Duck, C., Edwards, M.,

… Wanless, S. (2016). Phenological sensitivity to climate across taxa and trophic levels.

Nature, *535*(7611), 241–245. https://doi.org/10.1038/nature18608

van Klink, R., Bowler, D. E., Gongalsky, K. B., Swengel, A. B., Gentile, A., & Chase, J. M.

(2020). Meta-analysis reveals declines in terrestrial but increases in freshwater insect

abundances. *Science*, *368*, 417 – 420. https://doi.org/10.1126/science.aax9931

Vatka, E., Orell, M., & Rytkönen, S. (2016). The relevance of food peak architecture in trophic

interactions. *Global Change Biology*, *22*(4), 1585–1594.

https://doi.org/10.1111/gcb.13144

Visser, M. E., & Both, C. (2005). Shifts in phenology due to global climate change: The need for

a yardstick. *Proceedings of the Royal Society B: Biological Sciences*, *272*(1581), 2561–

2569. https://doi.org/10.1098/rspb.2005.3356

Visser, M. E., & Gienapp, P. (2019). Evolutionary and demographic consequences of

phenological mismatches. *Nature Ecology & Evolution*, *3*(6), 879–885.

https://doi.org/10.1038/s41559-019-0880-8

Visser, M. E., Noordwijk, A. J. van, Tinbergen, J. M., & Lessells, C. M. (1998). Warmer springs

lead to mistimed reproduction in great tits (Parus major). *Proceedings of the Royal*

Society of London. Series B: Biological Sciences, *265*(1408), 1867–1870.

https://doi.org/10.1098/rspb.1998.0514

Visser, M. E., te Marvelde, L., & Lof, M. E. (2012). Adaptive phenological mismatches of birds

and their food in a warming world. *Journal of Ornithology*, *153*(S1), 75–84.

https://doi.org/10.1007/s10336-011-0770-6

Walker, B. M., Senner, N. R., Elphick, C. S., & Klima, J. (2011). Hudsonian Godwit (Limosa

haemastica), version 2.0. In A. F. Poole (Ed.), *The Birds of North America*.

Weterings, M. J. A., Moonen, S., Prins, H. H. T., Wieren, S. E. van, & Langevelde, F. van.

(2018). Food quality and quantity are more important in explaining foraging of an

intermediate-sized mammalian herbivore than predation risk or competition. *Ecology and*

Evolution, *8*(16), 8419–8432. https://doi.org/10.1002/ece3.4372

Wilde, L. R., Swift, R. J., & Senner, N. R. (in revision). Flexible space use and density-

dependent heterospecific interactions determine fledging success in a precocial bird.

Journal of Animal Ecology.

Williams, J. B., Tieleman, B. I., Visser, G. H., & Ricklefs, R. E. (2007). Does Growth Rate

Determine the Rate of Metabolism in Shorebird Chicks Living in the Arctic?

Physiological and Biochemical Zoology, *80*(5), 500–513. https://doi.org/10.1086/520126

Yang, L. H., Cenzer, M. L., Morgan, L. J., & Hall, G. W. (2020). Species-specific, age-varying

plant traits affect herbivore growth and survival. *Ecology*, *101*(7), e03029.

https://doi.org/10.1002/ecy.3029

Yang, L. H., & Rudolf, V. H. W. (2010). Phenology, ontogeny and the effects of climate change

on the timing of species interactions. *Ecology Letters*, *13*(1), 1–10.

https://doi.org/10.1111/j.1461-0248.2009.01402.x

Tables

Table 1. Bayesian model selection on variables in a global logistic model predicting daily

survival rate in godwit chicks near Beluga River, AK from 2009 – 2019. Predictors were selected

using the indicator-variable approach, in which posterior inclusion probabilities (weights) and

Bayes Factors (BF) were estimated from a Bernoulli variable associated with each predictor.

Variables of the global model with BF > 3 and their component parts (i.e., interaction terms)

were included in the top model.

Variable	Weight, global	BF, global	Weight, top	BF, top
Age	0.53	1.125	0.52	1.081
Size	0.54	1.177	0.52	1.068
Hatch	0.54	1.175	-	-
Biomass	0.79	3.730	0.77	3.340
Age × Size	0.85	5.614	0.80	4.042

841   *Age: number of days since chick hatched; Hatch: individual chick's hatch date; Size: daily*

842   *median invertebrate body size; Biomass: daily invertebrate biomass*

853   Table 2. Standardized effect of variables on the survival rates of godwit chicks near Beluga

854   River, AK from 2009 – 2019. Posterior probabilities were estimated from a hierarchical model (n

855   = 122, posterior samples = 5000) with both survival and stochastic model components.

Predictor	Mean (SD)	95% Credible Interval	Pr $\neq$ 0
Intercept	-2.388 (5.253)	-13.10, 7.17	0.35
Age	-0.001 (0.201)	-0.41, 0.38	0.48
Size	0.331 (0.185)	-0.03, 0.71	0.96
Biomass	0.26 (0.169)	-0.08, 0.58	0.87
Age × Size	0.679 (0.103)	0.48, 0.88	1.00

861   *Age: number of days since hatch; Hatch: individual's hatch date; Size: daily median invertebrate*

862   *body size; Biomass: daily invertebrate biomass*

Figures

Figure 4. Effect of daily invertebrate biomass (a) and hatch date (b) on godwit chick body

condition index (BCI). BCI (hollow points) is the ratio of the observed to expected weight gain

since an individual's last measurement. BCI > 1 correspond with above average growth and BCI

< 1 below average growth. Regression line (black) and 95% confidence interval (grey) are

shown.

Figure 5. Effects of daily invertebrate biomass (a) and invertebrate body size (b) on the daily

survival of godwit chicks from the posterior mean estimates of a Bayesian hierarchical model

(credible intervals not shown). Biomass (dashed) had a constant effect, but the effect of size

varied with age (shade of grey, in days).

Figure 6. Correlation of seasonal fledging rates with measures of difference in dates (upper, left),

curve height (upper, right), whole demand percent overlap (lower, left), and peak demand

percent overlap (lower, right). Yearly fledging rates were extrapolated from daily survival rates

with accompanying 95% credible intervals. Regression lines (black) with 95% confidence

interval (grey) are shown.