Atomic Force Microscopy Reveals Distinct Protofilament-scale Structural Dynamics in Depolymerizing Microtubule Arrays

Authors: S. S. Wijeratne[1,2], M. Marchan[1†], J. S. Tresback[3†], R. Subramanian[1,2*]

Affiliations:

[1]Department of Molecular Biology, Massachusetts General Hospital, Boston, MA 02114, USA.

[2]Department of Genetics, Harvard Medical School, Boston, MA 02115, USA.

[3]Center for Nanoscale Systems, Harvard University, Cambridge, MA 02139, USA.

*Correspondence: radhika@molbio.mgh.harvard.edu.

† These authors contributed equally to this work.

ABSTRACT

The dynamic reorganization of microtubule-based cellular structures, such as the spindle and the axoneme, fundamentally depends on the dynamics of individual polymers within multi-microtubule arrays. A major class of enzymes, implicated in both the complete demolition and fine size control of microtubule-based arrays, are depolymerizing kinesins. How different depolymerases differently remodel microtubule arrays is poorly understood. A major technical challenge in addressing this question is that existing optical or electron-microscopy methods lack the spatial-temporal resolution to observe the dynamics of individual microtubules within larger arrays. Here we use Atomic Force Microscopy (AFM) to image depolymerizing arrays at single microtubule and protofilament resolution. We discover previously unseen modes of microtubule array destabilization by conserved depolymerases. We find that the kinesin-13 MCAK mediates asynchronous protofilament depolymerization and lattice-defect propagation, whereas the kinesin-8 Kip3p promotes synchronous protofilament depolymerization. Unexpectedly, MCAK can depolymerize the highly stable axonemal doublets, but Kip3p cannot. We propose that distinct protofilament-level activities underlie the functional dichotomy of depolymerases, resulting in either large-scale destabilization or length regulation of microtubule arrays. Our work establishes AFM as a powerful strategy to visualize microtubule dynamics within arrays and reveals how nanometer-scale substrate specificity leads to differential remodeling of micron-sized cytoskeletal structures.

20   **MAIN TEXT**

The dynamic formation and dismantling of protein arrays underlie a broad range of cellular functions in both prokaryotes and eukaryotes. A prototypical example of dynamic polymeric protein structures are micron-sized arrays of microtubules, which assemble into essential cellular machines and tracks such as the mitotic spindle in dividing cells, axonal arrays in neurons and

25   axonemes in cilia and flagella. Microtubules themselves are complex cylindrical macromolecular assembly of, most commonly, 13-15 protofilaments that are composed of repeating α,β-tubulin heterodimers. The intrinsic dynamic instability of microtubules and its regulation by a host of different Microtubule Associated Proteins (MAPs) are critical for the assembly and disassembly of microtubule arrays (1). How nanometer-scale dynamics of protofilaments (~4 nm) and

30   microtubules (~25 nm) result in the organization and remodeling of micron-sized multi-microtubule arrays remains poorly understood.

In vitro reconstitution and visualization by optical microscopy have provided tremendous insights into microtubule dynamic instability and its regulation by MAPs. However, these studies have been limited to single or pairs of microtubules as light microscopy does not have the

35   resolution to identify individual microtubules within a complex array of multiple microtubules. In addition, it is challenging to image individual protofilaments within each microtubule by this method. Structural intermediates of microtubule-remodeling reactions have been inferred from electron microscopy studies, but the single snapshot nature of the technique lacks temporal resolution to follow reaction dynamics in real time. To address these technical limitations and offer

40   insights into microtubule array remodeling at single microtubule and protofilament resolution in real time, we employed Atomic Force Microscopy (AFM) imaging (2-8). This technique allowed the direct visualization of microtubule depolymerization in two different arrays, antiparallel microtubule bundles that are found in the mitotic spindle and doublet microtubule arrays that form axonemes in cilia and flagella (9-11).

45   A critical reaction that governs the size and stability of microtubule arrays is microtubule depolymerization, which is catalyzed by a class of enzymes known as microtubule depolymerases. This reaction is required for rapid large-scale reorganization of the cytoplasm. For example, the mitotic spindle is built and disassembled every time a cell divides and the cilium is constructed and deconstructed each cell cycle (12-14). In addition to large scale reorganization of microtubule

50   networks and arrays, microtubule dynamics and its regulation are important for fine tuning the size of microtubule arrays (15-18). A fundamental conundrum is how the same reaction, the removal of tubulin from microtubules, results in different outcomes ranging from large-scale remodeling or fine-length regulation of microtubule arrays.

      Prototypical depolymerases are members of the two major family of kinesins, the

55   vertebrate kinesin-13 family, such as MCAK, and the budding yeast kinesin-8 family, such as Kip3p (19-21). While the non-motile MCAK and processive Kip3p proteins have different mechanisms for arriving at the microtubule ends, enzymatically MCAK and Kip3p are both catastrophe factors and catalyze the removal of tubulin from microtubule ends (22, 23) (24, 25). Structural studies suggests that at the microtubule end, both enzymes recognize the curved

60   conformation of tubulin in a similar manner (26-28). Despite these similarities, these proteins differently regulate dynamic instability such that Kip3p limits the distribution of maximum microtubule lengths whereas MCAK promotes rapid filament shortening (29-31). These differences are reflected in their distinct functions; kinesin-8 proteins are largely involved in length control of structures such as the spindle and the cilium, while kinesin-13s are additionally

65   implicated in large-scale cytoskeleton remodeling such as the depolymerization of interphase microtubules during entry into mitosis and suppression of cilium biogenesis (32-36). However, what underlies the differences in activity of these prototypical kinesin-family depolymerases and how differences in depolymerase activity at the single microtubule level translate to distinct remodeling of complex arrays remain unknown.

70   The AFM imaging reported here reveals the structural dynamics that underlie microtubule array destabilization and provide a framework for linking the action of enzymes on the nanometer-scale protofilaments to the remodeling of micron-sized arrays. The study sheds light on the long-standing question of how different depolymerases are tuned for distinct cellular activities, such as rapid remodeling or length control of microtubule arrays. Our findings highlight differences in

75   enzyme activity on the protofilament scale as a critical parameter that governs the fate of microtubules within complex structures, thereby dictating how such arrays are remodeled.

Imaging the depolymerization of PRC1-crosslinked microtubule arrays by Atomic Force Microscopy

80   As a first step towards imaging the depolymerization of microtubule arrays by MCAK, we reconstituted microtubule bundles using the antiparallel crosslinking protein PRC1 (Protein Regulator of Cytokinesis-1) on a mica surface (Fig. 1A) (see Methods and SI Appendix, Extended Methods). AFM images of the bundles reveal that PRC1 crosslinking results in 2D microtubule arrays (Fig. 1B-C). This is advantageous as every microtubule in the bundle can be clearly

85   distinguished visually (see Methods and SI Appendix, Extended Methods). This is reflected in a height plot as a series of peaks with an average height of ~30 nm (Fig. 1D). The 3D rendition of the images, obtained from the surface topography data of Fig. 1B, shows dense linkages connecting the overlapping microtubules in an array, consistent with the cooperative binding of PRC1 to overlapping microtubules (Fig. 1C) (37, 38).

90   We used two main criteria to set the imaging conditions for investigating microtubule dynamics within PRC1-crosslinked bundles (3-4 mins/frame with 256 x 256 pixels): (1) the spatial and temporal resolution is suitable for imaging the entire array as well as individual microtubules within the bundle, and (2) no sample damage is visible in the time frame of the experiment (15-30 mins). Experiments were performed with both GMPCPP- and GMPCPP + taxol-stabilized

95   microtubules. After locating a microtubule bundle by AFM, MCAK was added into the sample

chamber at the indicated concentrations and time lapse image series was acquired (*note:* solution

concentrations are reported throughout the manuscript; local concentrations on the mica surface

may be different). The rate of microtubule depolymerization by MCAK in the absence of PRC1 is

similar to the rates observed by TIRF (SI Appendix, Fig. S1A-C), which shows that the activity of

100   the enzyme is intact in this assay (SI Appendix, Fig. S1D-E) (24, 29, 39). The time-lapse AFM

imaging of PRC1-crosslinked bundles reveals that depolymerization of individual microtubules

within an array by MCAK can be visualized in real time (Fig. 1E, SI Appendix, Fig. S2, Videos 1-

2). Depolymerization begins shortly after MCAK addition, as we observed depolymerization of

microtubules within the arrays in the first image taken after MCAK addition. This is visualized in

105   sections of microtubule that lack complete tubules (Fig. 1E, time=0 min). In control experiments

without MCAK, PRC1-crosslinked microtubules remain stable and do not depolymerize

significantly during the experiment (SI Appendix, Fig. S3).

These observations reveal AFM imaging as a powerful method to examine dynamic

changes in a multi-microtubule array with nanometer-scale spatial resolution, thus offering a view

110   of structural changes that are extremely difficult to detect by other imaging methods (40).

MCAK depolymerizes microtubule protofilaments asynchronously and propagates defects within crosslinked bundles.

Close examination of the time lapse AFM images revealed two striking features of the

115   reaction intermediates. First, the loss of microtubules was observed to be associated with the

appearance of stripe-like features in the AFM images, which correspond to protofilaments. For

example, in the magnified time-lapse montages, partial and bidirectional depolymerization is

observed at the microtubule ends (Fig. 1F-K, SI Appendix, Fig. S2A). The stripes arise due to

protofilaments at different heights from the surface, which are resolved because the high

120   resolution of AFM in the axis perpendicular to the mica surface (see SI Appendix, Extended Methods). These observations show that in the presence of MCAK, the depolymerization of protofilaments within a microtubule is asynchronous (Fig. 1F, 1I, SI Appendix, Fig. S2A). The stripiness is observed on non-crosslinked as well as crosslinked microtubules with one or two neighbors. These observations suggest that microtubule protofilaments depolymerize at different

125   rates in the presence of MCAK, resulting in an asynchronous loss of protofilaments from the ends (Fig. 1L). To ensure that the features observed do not arise from the frequency of AFM scanning, we increased the time interval between scans to 5-10 mins after MCAK addition and found that the features were unaltered and independent of time interval (SI Appendix, Fig. S2D-E). Similar results were observed with microtubules stabilized by either GMPCPP or GMPCPP + taxol (Fig.

130   1F, SI Appendix, Fig. S2). EM studies have described curved ram horn shape of protofilaments during depolymerization. In the AFM experiments, these transient structures can only be captured if they lie in the plane of the mica surface as shown in SI Appendix, Fig. S2B.

       Second, in addition to depolymerization from the ends, we observed that breaks can also appear in the middle of microtubule arrays. These are likely to be defects formed by the loss of

135   tubulin from the microtubule lattice. We find that defects can be propagated by MCAK, again accompanied by the protofilament-associated stripiness (Fig. 2A-C, SI Appendix, Fig. S2B). Defects propagate both along the diameter and length of microtubules, with different rates (Fig. 2D-E), suggesting that destabilization of inter-protofilament interactions is slower than depolymerization along the length of an individual protofilaments in the presence of MCAK. The

140   observed depolymerization rates in the two opposite directions along the microtubule length are consistent with the unequal rates of plus and minus end microtubule depolymerization by MCAK (Fig. 2D) (23). More defects were observed in our taxol stabilized samples compared to GMPCPP, suggesting that polymerization conditions contribute to defects (41). Defect propagation was also observed in Total Internal Reflection Fluorescence (TIRF) microscopy-based assays with

145   GMPCPP microtubules in the presence of MCAK (300-500 nM) (SI Appendix, Fig. S1A-C). Thus, while the number of defects observed in AFM may be a combination of pre-existing lattice defects and any additional ones induced by the AFM tip, the TIRF and AFM data show that MCAK can propagate these defects to induce destabilization of arrays.

How does PRC1-mediated bundling of microtubules affect array destabilization by MCAK?
150   To address this, we quantitatively examined the effect of neighboring microtubules in bundles on the depolymerization reaction. Quantitative measurement of the depolymerization rates of microtubules that have zero, one or two neighbors (see Methods) showed that bundling has a protective effect on microtubules against MCAK-mediated depolymerization (SI Appendix, Fig. S1E). We find that the depolymerization rates of microtubules with two neighbors are 3-fold lower
155   than microtubules with zero or one neighbor. Second, microtubule depolymerization rates also depend on PRC1 concentration (SI Appendix, Fig. S1E). For instance, the depolymerization rate decreased when the solution concentration of PRC1 was increased by 10-fold from 10 nM to 100 nM. This reduction could be alleviated by increasing the MCAK concentration by ~10-fold. Under all conditions, the presence of two neighboring microtubules has a significant effect on the
160   protection of bundles. This is likely to arise from the highly dense pattern of PRC1 occupancy in overlap regions. Consequently, microtubules with two neighbors, one on either side, are likely to have the least number of exposed plus/minus protofilament ends (hereafter referred to as "exposed protofilament ends").

The features observed during AFM imaging of microtubule array depolymerization by
165   MCAK suggest that a single or few protofilaments with exposed ends can be an effective substrate for MCAK, and that these protofilaments are asynchronously removed by the enzyme. Altogether, we demonstrate that microtubule depolymerization can be visualized by AFM in real time at the single microtubule and protofilament resolution within arrays. By using this imaging modality, we provide the first view of protofilament level depolymerization by a microtubule depolymerase,

170   nearly two decades after such possibility was hypothesized (42, 43). Therefore, we formally show that protofilament-level depolymerization occurs on microtubules within a larger array in the presence of MCAK. The observed asynchronous loss of protofilaments and defect propagation suggests that MCAK can use an exposed protofilament ends as a substrate as would be advantageous for large-scale destabilization of the microtubule arrays.

Structural dynamics of microtubule depolymerization by Kip3p are distinct from MCAK

       To investigate whether different depolymerases exhibit distinct structural dynamics, we examined the depolymerization of PRC1-crosslinked microtubule bundles by Kip3p. We mainly focused on GMPCPP stabilized microtubules since depolymerization of doubly-stabilized

180   GMPCPP-taxol microtubules by Kip3p is extremely slow (23, 24). We first examined Kip3p activity on single microtubules in the absence of PRC1 (SI Appendix, Fig. S4A). The depolymerization rate increases with Kip3p concentration (SI Appendix, Fig. S4B) and are similar to the rates observed by TIRF microscopy (SI Appendix, Fig. S4C-D). Depolymerization is primarily seen at one end of the microtubules although we also observe some slow minus-end depolymerization

185   (SI Appendix, Fig. S4B inset; 0.34±0.07 nm/s on the plus end compared to 0.08±0.02 nm/s on the minus end at 1 nM solution concentration of Kip3p). Low levels of minus depolymerization are also observed in TIRF assays at concentrations > 50 nM (SI Appendix, Fig. S4C-D). Enhanced local concentration of Kip3p (on the mica substrate or on the cantilever tip) may contribute to Kip3p-induced depolymerization activity from both ends of the microtubules as has also been

190   observed with a non-motile Kip3p mutant (26).   Control experiments confirm that the depolymerization in this experiment is enzyme mediated (SI Appendix, Fig. S4E). We next examined Kip3p activity on PRC1-crosslinked microtubules. We selected a field where the ends of several microtubules within the PRC1 bundle were in view, added Kip3p, and collected time-lapse AFM images (Fig. 3A, SI Appendix, Fig. S5A-B, Video 3). We observed that the structural

195   features of depolymerization of PRC1-bound crosslinked and individual microtubules by Kip3p
were distinct from MCAK. First, in contrast to MCAK, no stripes were observed at the ends of
depolymerizing microtubules and the protofilaments were lost in unison, leaving only one or few
protofilament remnants (< 5 nm in height) that were adhered on the mica surface (Fig. 3B-D). This
is reflected in the shifting of the entire edge of the corresponding height profiles with
200   depolymerization without features of intermediate heights. Second, unlike with MCAK, we rarely
observed defect propagation events using the same batch of microtubules. In the rare instances
where we saw depolymerization from the middle of a doubly stabilized GMPCPP-taxol
microtubule, we again observed no stripes suggesting that defects are propagated by Kip3p in
unison, i.e. synchronously (SI Appendix, Fig. S5B). Third, in contrast to MCAK, we did not see a
205   significant effect of PRC1 concentration or neighbors on depolymerization rate (SI Appendix, Fig.
S5C). Control experiments in the presence of ATP alone, confirm that the observed
depolymerization of PRC1-crossslinked bundles in the presence of Kip3 is specific to the
presence of depolymerases in the assay (SI Appendix, Fig. S3).

In addition, we observed that Kip3p destabilizes highly curved microtubules, consistent
210   with the reported accumulation of Kip3 at these sites (26) (Fig. 3E-F, SI Appendix, Fig. S6A).
Destabilization of the curved region was neither observed in the presence of the motile
conventional kinesin (K401) or ATP alone (SI Appendix, Fig. S6B-C). We used time-resolved AFM
to follow the structural intermediates of the process by which the curved segment disintegrates in
real time in the presence of Kip3p (Fig. 3E-F, SI Appendix, Video 4) (26). As seen in the zoomed
215   3D view, the curved section destabilized and depolymerized faster in one direction (Fig. 3F).
Again, we observed that depolymerization of the entire microtubule occurs when most of the
protofilaments in the curved region are lost. Overall, the features of microtubule depolymerization,
such as synchronous depolymerization of protofilaments, lack of defect propagation, and
accelerated breaking at curved microtubule segments by Kip3p in the absence of PRC1 (SI

220   Appendix, Fig. S4A, S6A), are similar to that observed in experiments with PRC1 (Fig. 3A-F, SI

Appendix, Fig. S5A-B). Together these data suggest that unlike MCAK, Kip3p depolymerizes

protofilament ends synchronously. This is likely due to the accumulation of Kip3p at microtubule

ends and a preference to stall at ends, rather than at sections of partially exposed protofilament

ends such as those in defects.

225   Altogether, this suggests that the two depolymerases, Kip3p and MCAK, exhibit distinct

preferences in terms of substrate specificity at microtubule ends and defects at the protofilament

level, and in the context of an array of bundled microtubules. These findings shed light on how

the two depolymerases may be tuned for distinct functions. While the properties of MCAK make

it well suited for remodeling of arrays and depolymerization at defects, Kip3p activity seems better

230   aligned with a role of a length regulator at microtubule ends.

Visualizing the depolymerization of doublet-microtubules using AFM.

As a step towards examining how other microtubule arrays are remodeled by depolymerases, we

focused on the axoneme, an array of nine outer doublet and two central singlet microtubules that

235   forms the backbone of the cilia and flagella (Fig. 4A). The doublets are composed of the A-tubule,

which contains 13 protofilaments, and the B-tubule, which is an incomplete microtubule containing

ten protofilaments (44, 45). At the distal cilium tip, the doublets transition into an array composed

of singlets (44). Axoneme are one of the most stable arrays of microtubules in cells and

dissociating them into soluble tubulin requires fairly harsh treatments like sonication and

240   detergent, or specific ionic conditions (46, 47). Depolymerases of the kinesin-13 and kinesin-8

families are proposed to act on axonemes to control cilium length and stability (32-34). However,

the activity of depolymerases on doublet microtubules has not been visualized, and it is unknown

if and how these enzymes depolymerize doublets and impact axoneme stability.

We purified axonemes from *Lytechinus pictus* sea urchin sperm. We first focused on

245   individual doublets present in this sample. In high spatial resolution mode, the AFM images clearly

show two joined tubules, with heights of 25 and 35 nm respectively (Fig. 4B-C). Another feature

of these microtubule doublets were the periodic repeats separated by ~30 nm, which are likely to

be the outer dynein arms on the A tubule (Fig. 4B-C). Both of these structural features of

microtubules are consistent with cryo-EM structures of the axoneme and previous AFM images

250   of doublet microtubules (48, 49). First, we examined whether depolymerases act on doublets

microtubules by performing AFM experiments with 2D doublet sheets from partially dissociated

axonemes, composed of two or more doublets linked together (SI Appendix, Fig. S7A-D). In the

presence of MCAK, we observed that one of the tubules depolymerizes from both ends first,

followed by the depolymerization of the second tubule (Fig. 4D, SI Appendix, Figs. S7E-F and

255   Video 5). This can be visualized in the height plots as a deepening of the minima between

adjacent doublets (dotted lines) and the conversion of an asymmetric broad peak to a single sharp

peak (indicated by black arrow) (Fig. 4D). The AFM time-lapse data of a microtubule doublet and

a singlet in the same field of view in the presence of Kip3p revealed that the doublet

depolymerizes very slowly (rate= 0.07 nm/s), but the single microtubule is depolymerized on the

260   same time scale (rate= 0.43 nm/s) (Fig. 4E and SI Appendix, Video 6). This is consistent with

previous fluorescent studies with the kinesin-8 protein Kif19A, where single microtubules

nucleated from an axoneme were depolymerized, but the axoneme itself appears intact (32).

Within the doublet, the average depolymerization rate measured for the A- and B-tubule in the

presence of Kip3p was close to zero (0.02$\pm$0.02 nm/s) (Fig. 4F), while we estimated that MCAK

265   leads to faster B-tubule depolymerization than the A-tubule (Fig. 4F; A rate= 0.09$\pm$0.06 nm/s; B

rate= 0.38$\pm$0.12 nm/s). Control experiments with the doublets without enzyme showed no

significant depolymerization over the same time scale (SI Appendix, Fig. S7G). Taken together,

these data suggest clear differences in how MCAK and Kip3p process microtubule doublets.

Axonemes are heavily decorated with several proteins. Prior reports have shown that

270 treatment with a solution of Tris-EDTA-DTT (TED) can dissociate a subset of axonemal proteins, particularly the dyneins (50). Our AFM imaging revealed that these TED-treated doublets have a smoother surface in comparison to untreated doublets (SI Appendix, Fig. S8A-B). We examined if TED treatment alters the depolymerization of doublets. Similar to the doublet samples without TED treatment (Fig. 4), we observed bidirectional depolymerization of one tubule of an isolated

275 doublet with MCAK (SI Appendix, Fig. S8C). TED treatment resulted in faster depolymerization of both tubules (compared to non-TED with MCAK) with the B-tubule being lost at a higher rate than the A-tubule (SI Appendix, Fig. S8D; A rate= 0.26 ± 0.10 nm/s; B rate= 0.63 ± 0.11 nm/s). In the presence of Kip3p, TED treatment also permitted slow depolymerization of the B-tubule, but the A tubule remained intact (SI Appendix, Fig. S8E-F; A rate= 0.04 ± 0.02 nm/s; B rate= 0.17 ± 0.05

280 nm/s). To ensure that depolymerization is not arising from frequent AFM scanning, we have imaged the doublets every 10 minutes upon adding a depolymerase (SI Appendix, Fig. S8G). The results show that the depolymerization activity of the doublet with MCAK is independent of AFM scanning. In addition, in control experiments, without the enzyme, TED-treated doublets showed no significant depolymerization (SI Appendix, Fig. S8H).

285 Taken together, our results show for the first time that doublet microtubules can be enzymatically depolymerized with MCAK, and the rate of depolymerization of the B-tubule in a doublet is faster than the A-tubule. In contrast, doublet microtubules are poor substrates for Kip3p even when the doublet is stripped of associated proteins. Thus, proteins that have Kip3p-like properties may be selectively functional at the distal cilium tip, where the axoneme is composed

290 mostly of singlet microtubule, to fine-tune cilium length. In contrast, the properties of depolymerases like MCAK make them better suited to depolymerize both singlets and doublets for processes such as cilia disassembly or inhibition of ciliogenesis.

Destabilization of axonemal structures by MCAK

295   Our observations with the doublet microtubule depolymerization raise the question of how

preferential depolymerization of B-tubule by MCAK impacts stability of the entire axoneme. We

first imaged intact axonemes adsorbed onto the mica surface. Axonemes were ~200 nm in height,

which is consistent with the diameter of the axoneme from cryo-EM measurements (Fig. 5A-D

and SI Appendix, Fig. S9A-F) (48). The AFM amplitude image showed longitudinal striations,

300   which likely arise from the nine outer doublets that are around ~20-30 nm apart (Fig. 5B and SI

Appendix, Fig. S9B). In AFM imaging experiments with ATP alone, we observed no substantial

change in the overall size and integrity of the axoneme over time (Fig. 5E).

Next, we selected an axoneme, added depolymerase (t = 0), and monitored the changes

in the axoneme structure with time. With the addition of MCAK, we observed a rapid loss of

305   microtubules from the axoneme. For example, we documented that an intact axoneme (~200 nm

in max height) loses most of its tubules upon MCAK addition (SI Appendix, Fig. S9G).

Occasionally, we were able to capture intermediates in this reaction due to adsorption of the

dissociated microtubules on mica (Fig. 5F, SI Appendix, Fig. S9H, Video 7). In the example shown

in Fig. 5F-G, MCAK addition resulted in the rapid unfurling of the axoneme and the scattering of

310   microtubules on the surface.

These data suggest that the preferential and rapid depolymerization of one tubule in a

doublet by enzymes such as MCAK may result in disintegration of the structure by breaking the

links between doublets in an axoneme.

315   **Discussion**

Collectively, this study demonstrates the power of AFM imaging in visualizing dynamic

processes within dense microtubule arrays in real time and at spatial resolution that, for the first

time, allowed observations of individual protofilaments. AFM imaging of depolymerizing microtubule arrays revealed previously unseen structural dynamics and provided new mechanistic insights into how differences in the activities of microtubule remodeling factors at the level of protofilaments and microtubules can lead to differential fate of complex multi-microtubule arrays.

Our data show that the prototypical members of two major depolymerases of the kinesin superfamily, the kinesin-13 MCAK and the kinesin-8 Kip3p, depolymerize complex microtubule arrays in distinct ways. How is this achieved at the mechanistic level? The answer to this question has not been clear despite greater than two decades of enzymatic and structural studies. From the perspectives of enzymology and substrate preference, Kip3 and MCAK are similar. First, both enzymes use family specific loops in the motor domains to target curved microtubules ends (26, 31). Second, depolymerization catalyzed by both motors is not coupled to ATP hydrolysis. Instead, ATPase is inhibited at the microtubule end, and the role of hydrolysis is primarily to dissociate the enzymes from tubulin dimers after depolymerization, recycling for the next depolymerization reaction (26, 51). While an obvious difference lies in the diffusive scanning of MCAK and the processive motility of Kip3 on microtubules, these differences reflect the mechanism by which the enzymes arrive at microtubule ends and not their depolymerase activity (22-25). Indeed, it is observed that a non-motile Kip3 can depolymerize microtubules (26). Our data suggest that the observed differences stem from distinct substrate specificity of the depolymerases at the protofilament level (Fig. 6). We propose that MCAK molecules can effectively depolymerize an exposed protofilament end segment that it encounters at the middle or end of the lattice. This is reflected in the observation that MCAK can propagate defects and protofilaments can be asynchronously depolymerized from the ends. In contrast, a short exposed protofilament end like in a defect, is not sufficient for Kip3-mediated depolymerization from that site. Instead, Kip3 molecules synchronously depolymerize from the multi-protofilament ends of

microtubules, where they accumulate. This is consistent with the previously observed co-operativity in Kip3 activity (52, 53). This difference in protofilament-level substrate preference of the two enzymes lead to distinct action of crosslinked microtubules and doublets despite similarities in tubulin binding and ATPase activity at microtubule ends.

Our results also provide insights into the structural mechanism by which these depolymerases differentially regulate "microtubule aging". Aging refers to the observation that catastrophe results from a multi-step process, and its probability increases with microtubule lifetime (29). MCAK reduces this process to a first order reaction while Kip3 simply accelerates the rate of steps leading to catastrophe, and consequently MCAK is able to disassemble arrays faster than Kip3 (29). Current models propose that the catastrophe vulnerable state arises from the accumulation of defects and flayed protofilament ends that have fewer laterally stabilized protofilaments (40, 54, 55). How might MCAK and Kip3p differently affect lateral stabilization? The protofilament-level substrate specificity and efficient depolymerization of exposed protofilament ends by MCAK would prevent or greatly reduce the inherent lateral stabilization of protofilaments, thus effectively eliminating the need for accumulation of laterally un-stabilized protofilaments over time or aging. On the other hand, the stabilization of curved tubulin at microtubule ends by Kip3 will serve to increase the rate of aging but not the multi-step nature of the process. Together these observations provide a structural mechanism that can serve as a framework for understanding the molecular underpinnings of aging process, and it's regulation by depolymerizing enzymes.

The difference in substrate preference at the protofilament level has multiple implications for structural dynamics and regulation of microtubules and the arrays, and the role depolymerases play in these processes. First, lattice defects, which are sites with only a few exposed protofilament ends, are propagated by MCAK but not by Kip3p, suggesting that microtubule arrays with higher level of lattice defects may be especially sensitive to MCAK activity. This may be

advantageous for disassembly of microtubule arrays by MCAK in conjunction with proteins such as microtubule severing enzymes (56, 57). On the other hand, the inability to act on defects would be advantageous for depolymerases like Kip3 and other homologs of Kip3, such as Kif18A, that suppress microtubule dynamics for length control (58). Second, the restriction of Kip3p activity to microtubule ends makes it less sensitive to the crosslinking of microtubules by PRC1 compared to MCAK. This may arise from lower density of PRC1 crosslinks at microtubule ends, the primary site of Kip3p action, either because of crowding of Kip3p at ends or due to disruption of microtubule geometry at ends. This feature of Kip3p would allow it to act as an effective length-regulator even in the context densely crosslinked microtubule bundles. It is noteworthy that despite protection by PRC1 crosslinks, MCAK remains an overall faster remodeler due to rapid kinetics of tubulin removal and ability to access exposed protofilaments at both ends of microtubules and at defects. The properties of MCAK also allow it to enzymatically depolymerize one of the most stable microtubule structures in the cell, the doublet microtubules (46, 49, 59). We find that axonemal doublet microtubules can be depolymerized by MCAK, with B-tubule preferentially depolymerized over the A-tubule. The lower stability of the B-tubule due to fewer MIPs and incomplete tubule structure (48, 60, 61), likely exposes protofilaments that are effectively depolymerized by MCAK. Consequently, MCAK mediates the rapid destabilization of axonemal arrays. We think that the fast depolymerization of one tubule compromises the stability of the axonemal array by breaking the links between doublets. In contrast, proteins like Kip3p, which preferentially depolymerize singlets over doublets can act as length regulators at the distal cilium tip, which is predominantly comprised of singlet microtubules (62).

As demonstrated here, previously unknown structural dynamics of individual microtubules and protofilaments within complex arrays can be clearly visualized in the AFM time lapse images. In the case of the two major family of depolymerases, we find that functional dichotomy in the action of depolymerases, which determines how micron-scale arrays are remodeled, can arise

from differences in enzyme activity on the nanometer-sized protofilaments. These differences in the observed structural dynamics enable a greater diversity of microtubule array remodeling outcomes as would be beneficial in different cellular contexts when either large scale reorganization or fine-tuning of the architecture of cellular structures in required.

ACKNOWLEDGEMENTS

We thank S. Jiang for purifying MCAK and R. Ohi and D. Pellman for generously sharing Kip3p plasmids and proteins. R.S. was supported by the Pew Biomedical Foundation, the Smith Foundation and the NIH Director's New Innovator Award. This work was performed in part at the Center for Nanoscale Systems (CNS), a member of the National Nanotechnology Coordinated Infrastructure Network (NNCI), which is supported by the National Science Foundation under NSF award no. 1541959. CNS is part of Harvard University.

DATA AVAILABILITY: All data needed to evaluate the conclusions in the paper are present in the paper or the supplementary materials.

AUTHOR CONTRIBUTIONS

S.S.W. designed and performed the experiments, analyzed data and cowrote paper; M.M. performed TIRF experiments and analyzed data; J.S.T. assisted with AFM experiments; R.S. conceived of and designed the experiments, cowrote the paper and supervised the project. All authors discussed the results and commented on the manuscript.

COMPETING INTERESTS

The authors declare no competing interests.

415 **FIGURE TITLES AND LEGENDS**

Fig. 1. MCAK depolymerizes microtubule protofilaments asynchronously.

(A).   Schematic of an antiparallel microtubule array crosslinked by PRC1 (dotted lines).

(B).   AFM image of a microtubule bundle crosslinked by 100 nM PRC1. Each microtubule within the flat 2D array can be clearly distinguished. The x-y scale bar is 70 nm. The z-scale is 0

420   to 40 nm (dark to light brown). The AFM image is colored according to height from the surface.

(C).   The 3D rendition of a zoomed in region from B.

(D).   The corresponding height profile from dotted line in B.

(E).   Successive AFM images show depolymerization of individual microtubules within a PRC1-

425   crosslinked bundle by MCAK from 0-39 min. The image at 0 min represents the first frame taken after adding MCAK (GMPCPP + Taxol microtubules, PRC1:100 nM; MCAK: 70 nM). The x-y scale bar is 100 nm. The z-scale is 0 to 40 nm (dark to light brown).

(F)&(I).   Two examples of depolymerization from the experiment in E showing stripe-like appearance of depolymerizing protofilaments (boxes 1-2).

430   (G)&(J). The 3D rendition of the AFM time-lapse images from F and I.

(H)&(K). Corresponding height profiles from the white dotted line in panels F and I show that the stripiness results from protofilaments at different heights relative to the surface.

(L).   Schematic of a microtubule undergoing asynchronous protofilament-level depolymerization.

435   The scanning rate is 4 min/frame in B and ~3 min/frame in E with 256 x 256 pixels. The arrow on panels B) and E) indicates the scanning direction in the fast axis. See also SI Appendix, Figs. S1-3.

Fig. 2. MCAK propagates defects within crosslinked bundles.

440   (A).   Successive AFM images showing depolymerization of individual microtubules within a microtubule bundle by MCAK at 0-12 min. The image at 0 min represents the first frame taken after adding MCAK (GMPCPP + Taxol microtubules, PRC1:100 nM; MCAK: 70 nM). The x-y scale bar is 50 nm.

(B).   Successive time-lapse montages of a zoomed-in region of a section of microtubule from

445   the experiment in A (box 1) showing defect propagation on three microtubules within the array at the indicated times (arrows). 'S' indicates the edge with slow depolymerization, and 'F' indicates the edge with fast depolymerization. The x-y scale bar is 30 nm.

(C).   The corresponding average height profiles along the length of the microtubule from the dotted line in B.

450   (D).   Depolymerization rates of 'slow (S)' and 'fast (F)' events from defect propagation with MCAK (PRC1: 100 nM; MCAK: 70 nM; GMPCPP + Taxol microtubules: slow= 0.02±0.003 nm/s, n=19, fast=0.06±0.01 nm/s, n=17; GMPCPP microtubules: slow= 0.05±0.007 nm/s, n=18, fast=0.11±0.02 nm/s, n=18).

(E).   Depolymerization rates of 'diameter (D)' and 'length (L)' events from defect propagation

455   with MCAK (PRC1: 100 nM; MCAK: 70 nM; GMPCPP + Taxol microtubules: diameter= 0.01±0.002 nm/s, n=20, length=0.06±0.01 nm/s, n=20; GMPCPP microtubules: diameter= 0.03±0.004 nm/s, n=22, length=0.17±0.03 nm/s, n=22).

The scanning rate is ~3 min/frame with 256 x 256 pixels. The arrow on panel A) indicates the scanning direction in the fast axis. See also SI Appendix, Fig. S2.

Fig. 3. Structural dynamics of microtubule depolymerization by Kip3p are distinct from MCAK.

(A). Successive AFM time lapse images of a PRC1-crosslinked microtubule bundle in the presence of Kip3p at the indicated times (GMPCPP microtubules, PRC1:100 nM; Kip3p: 4 nM). The blue circles are fiduciary marks which show that the microtubules are depolymerizing and not sliding. The x-y scale bar is 100 nm.

(B)-(D). Zoomed-in regions from the experiment in A (boxes) showing bluntness at the depolymerizing microtubule end for N=0 (B), N=1 (C), N=2 (D) events. The height profiles corresponding to the dotted lines show that protofilaments at the ends of the microtubules are lost synchronously (white arrows). The x-y scale bar is 40 nm. For A-D, the z-scale is 0 to 40 nm (dark to light brown).

(E). Successive AFM time-lapse images show the destabilization of two highly curved regions (white arrows at 0 min) at the indicated times. Over time, the microtubule starts to depolymerize faster from one end with synchronous loss of protofilaments (blue arrows) (GMPCPP microtubules, PRC1:10 nM; Kip3p: 1 nM). The x-y scale bar is 100 nm. The z-scale is 0 to 30 nm (dark to light brown).

(F). The 3D rendition shows a magnified view of this depolymerization activity (dotted box at 0 min in E).

The scanning rate is ~3 min/frame with 256 x 256 pixels. The arrow on panels A) and E) indicates the scanning direction in the fast axis. See also SI Appendix, Figs. S4-S6.

Fig. 4. Visualizing the activity of MCAK and Kip3p on doublet-microtubules using AFM.

(A). Schematic of the axoneme structure. Axonemes consists of 9 outer microtubule doublets (MTD). Each doublet contains an A and B tubule. Outer dynein arms (ODA) present on the A tubule form a repeating pattern.

485   (B). AFM height image of an MTD. The A and the B tubule in an MTD were distinguished by the height of the tubules in the joined doublet (25 and 35 nm) and the periodic striations, which are separated by 30 nm (arrows). The x-y scale bar is 50 nm.

(C). The 3D representation of B and the corresponding height profile of dotted line in B.

(D). Successive AFM images show a 2D microtubule doublet sheet with MCAK at the indicated

490   times. The zoomed in region (boxed region) shows the depolymerization activity of alternate tubule in an array (arrows) and its corresponding height profiles over time (dotted line). The height profiles show the deepening of the minima and changing of the asymmetric peak into a single sharp peak (dotted line, arrow) (-TED sample, MCAK: 7 nM). The x-y scale bar of the zoomed-out and zoomed-in image is 80 nm and 40 nm, respectively. The z-scale is 0 to

495   40 nm (dark to light brown).

(E). Successive AFM height and amplitude images show a microtubule doublet (D) and a singlet (S) in the presence Kip3p at the indicated times. The corresponding height profiles from dotted line show that the height of the doublet doesn't change over time, but the height of the singlet reduces with time (dotted lines) (-TED sample, Kip3p: 5 nM). The x-y scale bar

500   is 100 nm. The z-scale is 0 to 50 nm (dark to light brown).

(F).   Depolymerization rates of A and B tubules in the -TED sample with MCAK and Kip3p.

MCAK: A rate=0.03±0.08 nm/s, n=12; B rate=0.38±0.12 nm/s, n=10; Kip3p: A

rate=0.003±0.015 nm/s, n=22; B rate=0.05±0.09 nm/s, n=22. Data were pooled from

experiments with protein concentrations less than 10 nM. The line on the boxes indicate

505   the mean of the sample. Statistical calculations used an unpaired t-test with Kolmogorov-

Smirnov correction for non-Gaussian distribution. * indicates a P-value of <0.05.

The scanning rate is ~4 min/frame in B and ~3 min/frame in E-F with 256 x 256 pixels. The arrow

on panels B), D) and E) indicates the scanning direction in the fast axis. See also SI Appendix,

Figs. S7 and S8.

Fig. 5. Destabilization of axonemal structures by depolymerases.

(A)-(C). AFM height (A), amplitude (B) and 3D (C) images of an intact axoneme from Lytechinus pictus sea urchin sperm. The AFM amplitude (B) and the 3D (C) images show longitudinal striations, likely from the nine outer doublets, which are around ~20-30 nm apart. The x-y scale bar is 300 nm. The z-scale is 0–200 nm (dark to light brown).

(D). The height profile of the selected dotted line from height image in A is ~200 nm.

(E). Successive AFM images of an intact axoneme with 1 mM ATP at the indicated times. With ATP alone, no significant change was observed in the axoneme structure. The x-y scale bar is 300 nm. The z-scale is 0–300 nm (dark to light brown).

(F). Successive AFM images of an axoneme in the presence of MCAK (1 nM) at the indicated times. At t=0, the axoneme has been partially frayed. The x-y scale bar is 500 nm. The z-scale is 0–90 nm (dark to light brown).

(G). Schematic of proposed intermediate that results in unfurling of the axoneme with MCAK. The scanning rate in A-C is ~4 min/frame in B and ~3 min/frame in E-F with 256 x 256 pixels. The arrow on panels A), E) and F) indicates the scanning direction in the fast axis. See also SI Appendix, Fig. S9.

Fig. 6. Summary of AFM findings

Schematic of AFM imaging illustrates the distinct structural dynamics and intermediates during microtubule depolymerization with different enzymes and its impact on microtubule arrays. In the context of an individual microtubule: (1) the depolymerization of protofilaments is asynchronous with MCAK and synchronous with Kip3p, and (2) MCAK propagates lattice defects whereas Kip3p does not. In the context of microtubule arrays: (1) Crosslinking by PRC1 protects microtubules against depolymerization by MCAK and does not significantly influence the depolymerization by Kip3p, and (2) MCAK

550   depolymerizes doublet microtubules and results in the destabilization of axonemal arrays.

This arises from fast depolymerization of one tubule which compromises the stability of

the cylindrical doublet array.

REFERENCES

1. N. B. Gudimchuk, J. R. McIntosh, Regulation of microtubule dynamics, mechanics and function through the growing tip. *Nat Rev Mol Cell Biol* 10.1038/s41580-021-00399-x (2021).

2. N. Kodera, D. Yamamoto, R. Ishikawa, T. Ando, Video imaging of walking myosin V by high-speed atomic force microscopy. *Nature* **468**, 72-76 (2010).

3. T. Ando *et al.*, A high-speed atomic force microscope for studying biological macromolecules. *Proc Natl Acad Sci U S A* **98**, 12468-12472 (2001).

4. I. A. Schaap, C. Carrasco, P. J. de Pablo, C. F. Schmidt, Kinesin walks the line: single motors observed by atomic force microscopy. *Biophys J* **100**, 2450-2456 (2011).

5. A. Vinckier *et al.*, Immobilizing and imaging microtubules by atomic force microscopy. *Ultramicroscopy* **57**, 337-343 (1995).

6. L. Hamon, P. A. Curmi, D. Pastre, High-resolution imaging of microtubules and cytoskeleton structures by atomic force microscopy. *Methods Cell Biol* **95**, 157-174 (2010).

7. Y. F. Dufrene *et al.*, Imaging modes of atomic force microscopy for application in molecular and cell biology. *Nat Nanotechnol* **12**, 295-307 (2017).

8. C. Elie-Caille *et al.*, Straight GDP-tubulin protofilaments form in the presence of taxol. *Curr Biol* **17**, 1765-1770 (2007).

9. C. Duellberg, F. J. Fourniol, S. P. Maurer, J. Roostalu, T. Surrey, End-binding proteins and Ase1/PRC1 define local functionality of structurally distinct parts of the microtubule cytoskeleton. *Trends Cell Biol* **23**, 54-63 (2013).

10. S. C. Goetz, K. V. Anderson, The primary cilium: a signalling centre during vertebrate development. *Nature reviews. Genetics* **11**, 331-344 (2010).

11. R. Subramanian, T. M. Kapoor, Building complexity: insights into self-organized assembly of microtubule-based architectures. *Developmental cell* **23**, 874-885 (2012).

12. J. B. Woodruff, D. G. Drubin, G. Barnes, Spindle assembly requires complete disassembly of spindle remnants from the previous cell cycle. *Mol Biol Cell* **23**, 258-267 (2012).

13. I. Sanchez, B. D. Dynlacht, Cilium assembly and disassembly. *Nat Cell Biol* **18**, 711-717 (2016).

14. N. McHedlishvili, H. K. Matthews, A. Corrigan, B. Baum, Two-step interphase microtubule disassembly aids spindle morphogenesis. *BMC Biol* **16**, 14 (2018).

15. G. Goshima, J. M. Scholey, Control of mitotic spindle length. *Annu Rev Cell Dev Biol* **26**, 21-57 (2010).

16. H. Ishikawa, W. F. Marshall, Ciliogenesis: building the cell's antenna. *Nat Rev Mol Cell Biol* **12**, 222-234 (2011).

17. Z. Hu, Y. Liang, D. Meng, L. Wang, J. Pan, Microtubule-depolymerizing kinesins in the regulation of assembly, disassembly, and length of cilia and flagella. *Int Rev Cell Mol Biol* **317**, 241-265 (2015).

18. S. Shrestha, M. Hazelbaker, A. L. Yount, C. E. Walczak, Emerging Insights into the Function of Kinesin-8 Proteins in Microtubule Length Regulation. *Biomolecules* **9** (2018).

19. C. E. Walczak, S. Gayek, R. Ohi, Microtubule-depolymerizing kinesins. *Annu Rev Cell Dev Biol* **29**, 417-441 (2013).

20. M. L. Reilly, A. Benmerah, Ciliary kinesins beyond IFT: Cilium length, disassembly, cargo transport and signalling. *Biol Cell* **111**, 79-94 (2019).

21. J. Howard, A. A. Hyman, Microtubule polymerases and depolymerases. *Curr Opin Cell Biol* **19**, 31-35 (2007).

22. M. L. Gupta, Jr., P. Carvalho, D. M. Roof, D. Pellman, Plus end-specific depolymerase activity of Kip3, a kinesin-8 protein, explains its role in positioning the yeast mitotic spindle. *Nat Cell Biol* **8**, 913-923 (2006).

23. V. Varga *et al.*, Yeast kinesin-8 depolymerizes microtubules in a length-dependent manner. *Nat Cell Biol* **8**, 957-962 (2006).

24. J. Helenius, G. Brouhard, Y. Kalaidzidis, S. Diez, J. Howard, The depolymerizing kinesin MCAK uses lattice diffusion to rapidly target microtubule ends. *Nature* **441**, 115-119 (2006).

25. A. Desai, S. Verma, T. J. Mitchison, C. E. Walczak, Kin I kinesins are microtubule-destabilizing enzymes. *Cell* **96**, 69-78 (1999).

26. H. Arellano-Santoyo *et al.*, A Tubulin Binding Switch Underlies Kip3/Kinesin-8 Depolymerase Activity. *Developmental cell* **42**, 37-51 e38 (2017).

27. T. Ogawa, R. Nitta, Y. Okada, N. Hirokawa, A common mechanism for microtubule destabilizers-M type kinesins stabilize curling of the protofilament using the class-specific neck and loops. *Cell* **116**, 591-602 (2004).

28. A. B. Asenjo *et al.*, Structural model for tubulin recognition and deformation by kinesin-13 microtubule depolymerases. *Cell Rep* **3**, 759-768 (2013).

29.   M. K. Gardner, M. Zanic, C. Gell, V. Bormuth, J. Howard, Depolymerizing kinesins Kip3 and MCAK shape cellular microtubule architecture by differential control of catastrophe. *Cell* **147**, 1092-1103 (2011).

30.   D. Wang *et al.*, Motility and microtubule depolymerization mechanisms of the Kinesin-8 motor, KIF19A. *Elife* **5** (2016).

31.   A. M. Mulder *et al.*, A new model for binding of kinesin 13 to curved microtubule protofilaments. *J Cell Biol* **185**, 51-57 (2009).

32.   S. Niwa *et al.*, KIF19A is a microtubule-depolymerizing kinesin for ciliary length control. *Dev Cell* **23**, 1167-1175 (2012).

33.   T. Kobayashi, W. Y. Tsang, J. Li, W. Lane, B. D. Dynlacht, Centriolar kinesin Kif24 interacts with CP110 to remodel microtubules and regulate ciliogenesis. *Cell* **145**, 914-925 (2011).

34.   T. Piao *et al.*, A microtubule depolymerizing kinesin functions during both flagellar disassembly and flagellar assembly in Chlamydomonas. *Proc Natl Acad Sci U S A* **106**, 4713-4718 (2009).

35.   C. E. Walczak, T. J. Mitchison, A. Desai, XKCM1: a Xenopus kinesin-related protein that regulates microtubule dynamics during mitotic spindle assembly. *Cell* **84**, 37-47 (1996).

36.   T. Maney, A. W. Hunter, M. Wagenbach, L. Wordeman, Mitotic centromere-associated kinesin is important for anaphase chromosome segregation. *J Cell Biol* **142**, 787-801 (1998).

37.   P. Bieling, I. A. Telley, T. Surrey, A minimal midzone protein module controls formation and length of antiparallel microtubule overlaps. *Cell* **142**, 420-432 (2010).

38.   R. Subramanian *et al.*, Insights into antiparallel microtubule crosslinking by PRC1, a conserved nonmotor microtubule binding protein. *Cell* **142**, 433-443 (2010).

39.   J. R. Cooper, M. Wagenbach, C. L. Asbury, L. Wordeman, Catalysis of the microtubule on-rate is the major parameter regulating the depolymerase activity of MCAK. *Nat Struct Mol Biol* **17**, 77-82 (2010).

40.   C. E. Coombes, A. Yamamoto, M. R. Kenzie, D. J. Odde, M. K. Gardner, Evolving tip structures can explain age-dependent microtubule catastrophe. *Curr Biol* **23**, 1342-1348 (2013).

41.   A. Rai *et al.*, Taxanes convert regions of perturbed microtubule growth into rescue sites. *Nat Mater* **19**, 355-365 (2020).

42.   H. Niederstrasser, H. Salehi-Had, E. C. Gan, C. Walczak, E. Nogales, XKCM1 acts on a single protofilament and requires the C terminus of tubulin. *J Mol Biol* **316**, 817-828 (2002).

43.   C. A. Moores *et al.*, A mechanism for microtubule depolymerization by KinI kinesins. *Mol Cell* **9**, 903-909 (2002).

44. P. Satir, Studies on cilia. 3. Further studies on the cilium tip and a "sliding filament" model of ciliary motility. *J Cell Biol* **39**, 77-94 (1968).

45. D. Nicastro *et al.*, Cryo-electron tomography reveals conserved features of doublet microtubules in flagella. *Proc Natl Acad Sci U S A* **108**, E845-853 (2011).

46. R. Orbach, J. Howard, The dynamic and structural properties of axonemal tubulins support the high length stability of cilia. *Nat Commun* **10**, 1838 (2019).

47. L. I. Binder, J. L. Rosenbaum, The in vitro assembly of flagellar outer doublet tubulin. *J Cell Biol* **79**, 500-515 (1978).

48. D. Nicastro *et al.*, The molecular architecture of axonemes revealed by cryoelectron tomography. *Science* **313**, 944-948 (2006).

49. M. Owa *et al.*, Inner lumen proteins stabilize doublet microtubules in cilia and flagella. *Nat Commun* **10**, 1143 (2019).

50. R. W. Linck, Flagellar doublet microtubules: fractionation of minor components and alpha-tubulin from specific regions of the A-tubule. *J Cell Sci* **20**, 405-439 (1976).

51. M. Wagenbach, S. Domnitz, L. Wordeman, J. Cooper, A kinesin-13 mutant catalytically depolymerizes microtubules in ADP. *J Cell Biol* **183**, 617-623 (2008).

52. M. Bugiel, M. Chugh, T. J. Jachowski, E. Schaffer, A. Jannasch, The Kinesin-8 Kip3 Depolymerizes Microtubules with a Collective Force-Dependent Mechanism. *Biophys J* **118**, 1958-1967 (2020).

53. V. Varga, C. Leduc, V. Bormuth, S. Diez, J. Howard, Kinesin-8 motors act cooperatively to mediate length-dependent microtubule depolymerization. *Cell* **138**, 1174-1183 (2009).

54. C. Duellberg, N. I. Cade, T. Surrey, Microtubule aging probed by microfluidics-assisted tubulin washout. *Mol Biol Cell* **27**, 3563-3573 (2016).

55. P. Zakharov *et al.*, Molecular and Mechanical Causes of Microtubule Catastrophe and Aging. *Biophys J* **109**, 2574-2591 (2015).

56. D. J. Sharp, J. L. Ross, Microtubule-severing enzymes at the cutting edge. *J Cell Sci* **125**, 2561-2569 (2012).

57. F. J. McNally, A. Roll-Mecak, Microtubule-severing enzymes: From cellular functions to molecular mechanism. *J Cell Biol* **217**, 4057-4069 (2018).

58. Y. Du, C. A. English, R. Ohi, The kinesin-8 Kif18A dampens microtubule plus-end dynamics. *Curr Biol* **20**, 374-380 (2010).

59. G. B. Witman, K. Carlson, J. L. Rosenbaum, Chlamydomonas flagella. II. The distribution of tubulins 1 and 2 in the outer doublet microtubules. *J Cell Biol* **54**, 540-555 (1972).

60. M. Ma *et al.*, Structure of the Decorated Ciliary Doublet Microtubule. *Cell* **179**, 909-922 e912 (2019).

61. M. Ichikawa *et al.*, Tubulin lattice in cilia is in a stressed form regulated by microtubule inner proteins. *Proc Natl Acad Sci U S A* **116**, 19930-19938 (2019).

62. P. Kiesel *et al.*, The molecular structure of mammalian primary cilia revealed by cryo-electron tomography. *Nat Struct Mol Biol* **27**, 1115-1124 (2020).

63. J. S. Bonifacino (1998) Current protocols in cell biology. (John Wiley, New York), pp volumes (loose-leaf).

64. H. G. Hansma, D. E. Laney, DNA binding to mica correlates with cationic radius: assay by atomic force microscopy. *Biophys J* **70**, 1933-1939 (1996).

65. D. Pastre *et al.*, Adsorption of DNA to mica mediated by divalent counterions: a theoretical and experimental study. *Biophys J* **85**, 2507-2518 (2003).

66. P. R. Heenan, T. T. Perkins, Imaging DNA Equilibrated onto Mica in Liquid Using Biochemically Relevant Deposition Conditions. *ACS Nano* **13**, 4220-4229 (2019).

METHODS

Protein purification

PRC1 was expressed and purified as described previously (38). Fluorescent MCAK plasmid was synthesized and the protein expressed and purified as described (24). Kip3p plasmid

560 and protein were a gift from R. Ohi (University of Michigan). Both proteins were purified using standard Ni-NTA purification protocol followed by Superose-6 size exclusion chromatography.

Microtubule polymerization

GMPCPP polymerized and taxol-stabilized microtubules were prepared as described previously (38). Briefly, GMPCPP seeds were prepared from a mixture of unlabeled bovine tubulin,

565 X-rhodamine-tubulin, and biotin tubulin, which were diluted in BRB80 buffer (80 mM PIPES pH 6.8, 1.5 mM $MgCl_2$, 0.5 mM EGTA, pH 6.8) and mixed by tapping lightly. The diluted seeds were transferred to a 37°C heating block and covered with foil to minimize light exposure. After approximately 1 hour, 100 µL of warm BRB80 buffer was added to the microtubules and spun at 75,000 rpm, 10 min and 37°C to remove free unpolymerized tubulin. Following the centrifugation

570 step, the supernatant was discarded, and the pellet was washed by round of centrifugation with

100 µL BRB80. The pellet was resuspended in 16 µL of BRB80 and stored at room temperature covered in foil. 20 µM taxol was added for the preparation of doubly stabilized microtubules. Rhodamine-labeled microtubules were used for both TIRF and AFM experiments.

Isolation of axoneme

Sperm flagellar axonemes were isolated and purified from sea urchin Lytechinus pictus (Marinus Scientific, LLC, Newport Beach, CA), according to the procedures of Salmon et al. (63). Briefly, sea urchin sperm were collected by inducing them to spawn by injecting the animal with 0.5 M KCl. The sperm were diluted with artificial sea water and put on ice for 20 mins. The sperm suspensions were then centrifuged at 500 rpm. Afterwards, the supernatant was collected, and the sperm were pelleted by centrifuging at 5000 rpm. The sperm pellets were resuspended by trituration in 20% sucrose to osmotically remove the plasma membranes, and the demembranated sperm were homogenized to break the sperm heads from the tails. The suspensions were centrifuged at 10,000 rpm to pellet whole sperm and sperm heads, and the supernatants recentrifuged at 13,000 rpm to pellet the detached sperm tails. The pellet was stratified into a top white layer that contains the demembranated tails and a bottom yellow layer that contains heads and debris. The white layer was collected and resuspended by trituration in isolation buffer. The resuspended tails break the tails into fragments and were further centrifuged at 10,000 rpm. The top white layer was resuspended by trituration in isolation buffer centrifuged to completely separate the tail fragments from heads and debris until the pellet is a single layer of pure white. The white pellet was resuspended in extraction buffer and homogenized and incubated on ice for 45 min to extract dyneins and central pair MTs from the tail fragments. The extracted axonemes were separated from soluble proteins by centrifuging them at 13,000 rpm. The axoneme pellet was resuspended by trituration in extraction buffer, and the axoneme fragments were re-extracted by incubation. The extracted axonemes were pelleted by centrifugation at 13,000 rpm. The extracted axoneme pellet was resuspended by trituration in

isolation buffer containing 50% glycerol. Unless specified, all steps were performed at 4°C, and all centrifugation times were 5-10 min.

600   Doublet fractionation

Axoneme pellets were taken up in Tris-EDTA-DTT solution (TED, 2 mM Tris, 0.2 mM EDTA and 0.5 mM DTT, pH 7.8). As described in the literature, the TED treatment has shown to remove 40-50% of the protein in the sea urchin doublets, which remain associated in sheets of 9 or fewer (50). The treatment time is empirically determined to obtain a mixture of sheets and
605   isolated doublets.

Atomic Force Microscope experiments

Sample preparation

Microtubule adsorption on mica is achieved by increasing the multivalent cation
610   concentration of the buffer as described previously (6). To prepare microtubule bundles, taxol-stabilized or GMPCPP microtubules, PRC1 and BRB80 buffer with an additional 5 mM $MgCl_2$ were combined in a tube. The microtubule and protein mixture was quickly spun down for a few minutes and ~20 µL of this mixture was deposited on a mica substrate freshly cleaved by scotch tape. Single microtubule, doublet, and axoneme samples were diluted with BRB80 and 5 mM
615    $MgCl_2$ and deposited on mica as described above. After a few minutes of incubation, ~10 µL of BRB80 buffer was added to the mica and to the AFM tip before imaging the sample by AFM.

AFM imaging

To acquire static AFM images of the sample, clean regions showing flat microtubule arrays were chosen. To get a high-resolution image of the sample, the scan size was decreased to ~ <1
620   µm. For time-lapse experiments, after finding a flat microtubule bundle on mica, we zoomed into a 1-2 µm region preferably with a few microtubule ends in the scan area. After acquiring this t=0 image, the scanning was briefly paused, the distance between tip and sample was increased by

100-200 nm and 10-20 µL of the depolymerase was injected on to the surface with a pipette without touching the mica disc to prevent losing the scanned area. The amount of liquid on the mica was monitored to ensure that the sample did not dehydrate, and additional buffer was added to the mica to maintain solution volume during the experiments. A range of enzyme concentrations was tested, and final conditions were empirically determined because some protein is lost to the surface and cantilever in these experiments.

All AFM experiments were carried out by tapping mode in liquid with the Asylum Cypher S with a silicon nitride tip (BL-AC40TS, radius: 8 nm; spring constant: 0.09 N/m; Oxford Instruments). After adding the depolymerase, imaging was started at a frame rate of ~3 mins/frame (256 × 256 pixels at ~1.5 Hz). In liquid, the drive frequency of the tip was ~ 25 kHz. We kept the scan rate at or below 1.5 Hz and scanned at 50-100 pN tip forces to minimize sample damage. Data were collected on microtubules in various orientations due to random binding to the surface, and the features observed were not specific to microtubule orientation with respect to the scan direction. Image acquisition took place over ~30 mins while constantly maintaining the drive amplitude of the tip throughout the experiment. The drive amplitude was maintained slightly above a point where the tip starts to make contact with the surface. To exclude the possibility that depolymerization is arising from the AFM tip, microtubule bundles or doublets were imaged every 5-10 minutes upon adding depolymerase (SI Appendix Fig. 2D-E, 7G). Because of sample contamination and tip damage, using the same tip for all the experiments is not possible. For each new experiment, a new tip was used, and we ensured that we obtained the same quality of AFM images as determined by the spatial resolution or the width of the sample.

Data Analysis

Raw AFM data were processed with the Asylum Research (version 16.14.216) and Gwiddyon software (http://gwyddion.net/). All AFM images (height, phase and amplitude) were flattened and any horizontal scars from scanning artifacts were removed prior to analysis. Only

the clearest AFM images were used for analysis. After data processing, the height profiles of the microtubules were obtained from the AFM height image.

The number of neighbors was determined by the number of microtubules which were physically contacting an individual microtubule in parallel in a large bundle. For example, a microtubule with no neighbors nearby has N=0, a microtubule in contact with another microtubule has N=1, and a microtubule in contact with two microtubules has N=2.

The depolymerization rates from AFM images were determined by calculating the average length change along the microtubule between the first frame and the last frame from the height time-lapse images (Fig. 2D-E & 4F, SI Appendix, Fig. S1D-E, S4B, S5C & S8D).

For the rate of defect propagation, after the appearance of a defect, the average length change over time from both edges of a defect was measured. In Fig. 2D, 'slow' refers the edge with the smallest change in length relative to the 'fast' edge. In Fig. 2E, 'diameter' refers to change in length over time around the diameter of the microtubule and 'length' refers to the change in length over time along the length of the microtubule. The units were converted from nm/min to nm/secs for easier comparison to our TIRF assay measurements and previous literature.

In vitro fluorescence microscopy assay

The microscope slides (Gold Seal Cover Glass, 24 × 60 mm, thickness No.1.5) and coverslips (Gold Seal Cover Glass, 18 × 18 mm, thickness No.1.5) were cleaned and functionalized with biotinylated PEG and non-biotinylated PEG, respectively, to prevent nonspecific surface sticking, according to standard protocols (38). Flow chambers were built by applying three strips of double-sided tape to a slide and attaching the coverslip. Sample chamber volumes were approximately 6–8 μL.

Experiments were performed as described previously (38). Biotinylated GMPCPP microtubules, labeled with rhodamine, were immobilized in a flow chamber by first coating the surface with neutravidin (0.2 mg/ml). To visualize microtubule depolymerization, MCAK or Kip3p

675 and 1 mM ATP were flowed into the chamber in assay buffer (BRB80 buffer supplemented with 5 mM TCEP, 2 mM MgCl$_2$, 0.2 mg/ml k-casein, 4 mg/ml glucose oxidase, 0.35 mg/ml glucose catalase, 1% b-mercaptoethanol, and 5% sucrose), and a time-lapse sequence of images was immediately acquired at a rate of 6-12 frames/min. Data were collected for 5–25 min.

All experiments were performed on Nikon Ti-E inverted microscope with a Ti-ND6-PFS 680 perfect focus system equipped with an APO TIRF 100x oil/1.49 DIC objective (Nikon). The microscope was outfitted with a Nikon-encoded x-y motorized stage and a piezo z-stage, an sCMOS camera (Andor Zyla 4.2), and two-color TIRF imaging optics (Lasers: 488 nm and 561 nm; Filters: Dual Band 488/561 TIRF exciter).

ImageJ (NIH) was used to process the image files. Briefly, raw time-lapse images were 685 converted to tiff files. From these images, individual microtubule depolymerization events were identified and converted to kymographs by the MultipleOverlay and MultipleKymograph plug-ins (J. Reitdorf and A. Seitz; https://www.embl.de/eamnet/html/body_kymograph.html).

The following criteria were used to select microtubules for quantitative analysis: (1) firmly attached to the surface throughout depolymerization; (2) both ends visible in the initial frame; (3) 690 not overlapping with another microtubule or tubulin aggregate. A threshold was applied to each kymograph to distinguish microtubule from background. To calculate depolymerization rates, the derivative of the position versus time coordinates of each external edge of the microtubule in the rhodamine channel was measured using Velocity_Measurement_Tool macro and then converted from pixels/frame to nm/s. Breaks within microtubules were visually identified from kymographs 695 with a triangular area of low (background) intensity within the bright microtubule region, which indicates a break in the microtubule that continues to be depolymerized from both sides. Total microtubule length for each condition was calculated as a sum of the widths of all quantified kymographs, converted from pixels to microns.

Figure 1

Figure 2

Figure 3

A GMPCPP MTs, 100 nM PRC1 + 4 nM Kip3p

9 min · *N*=0 · *N*=1 · *N*=2

12 min

15 min

18 min

0 — 40 nm

B *N*=0

C *N*=1

D *N*=2

Distance (nm)

E GMPCPP MTs, 10 nM PRC1 + 1 nM Kip3p

0 min · 6 min · 12 min · 18 min · 24 min · 30 min

0 — 30 nm

F

0 — 30 nm

Figure 4

Figure 5

Figure 6