The evolutionary assembly of forest communities along environmental gradients: recent diversification or sorting of pre-adapted clades?

Running title: Andean community assembly across elevation

Alexander G. Linan[1*†], Jonathan A. Myers[2], Christine E. Edwards[1], Amy E. Zanne[3], Stephen A. Smith[4], Gabriel Arellano[4], Leslie Cayola[1,5], William Farfan-Ríos[1,2], Alfredo F. Fuentes[1,5], Karina Garcia-Cabrera[6], Sebastián Gonzales-Caro[7], M. Isabel Loza[1,5,8], Manuel J. Macía[9,10], Yadvinder Malhi[11], Beatriz Nieto-Ariza[12], Norma Salinas Revilla[13], Miles Silman[14], and J. Sebastián Tello[1†]

1. Center for Conservation and Sustainable Development, Missouri Botanical Garden, St. Louis, Missouri, USA
2. Department of Biology, Washington University in St. Louis, St. Louis, Missouri, USA
3. Department of Biological Sciences, The George Washington University, Washington DC, USA
4. Department of Ecology and Evolutionary Biology, University of Michigan, Ann Arbor, Michigan, USA
5. Herbario Nacional de Bolivia, Universidad Mayor de San Andrés, La Paz, Bolivia
6. Escuela Profesional de Biología, Universidad Nacional de San Antonio Abad del Cusco, Cusco, Peru
7. Departamento de Ciencias Forestales, Universidad Nacional de Colombia Sede Medellín, Universidad Nacional de Colombia, Medellín, Colombia
8. Department of Biology, University of Missouri-St Louis, St. Louis, Missouri, USA
9. Departamento de Biología, Área de Botánica, Universidad Autónoma de Madrid, Madrid, Spain
10. Centro de Investigación en Biodiversidad y Cambio Global (CIBC-UAM), Universidad Autónoma de Madrid, Madrid, Spain.
11. Environmental Change Institute, School of Geography and the Environment, University of Oxford, Oxford, England, United Kingdom
12. Hospital Central de Ivirgarzama, Puerto Villarroel, Bolivia
13. Institute for Nature Earth and Energy, Pontificia Universidad Catolica del Peru, Lima, Peru
14. Center for Energy, Environment and Sustainability, Winston-Salem, North Carolina, USA

† These authors contributed equally to this study

E-mails:
alinan@mobot.org*
jamyers@wustl.edu

Christine.Edwards@mobot.org
JuanSebastian.Tello@mobot.org
wfarfan@gmail.com
belarivera23@gmail.com
alfrefuentes@gmail.com
lcayola@gmail.com
manuel.macia@uam.es
sebastian.gonzalez.caro@gmail.com
yadvinder.malhi@ouce.ox.ac.uk
gabriel.arellano.torres@gmail.com
azanne@gwu.edu
eebsmith@umich.edu
nosare@gmail.com
silmanmr@wfu.edu
karigac@gmail.com
beatriz.nieto.ariza@gmail.com
55

*Correspondence phone number: 1(314)-577-9473 ext.77264

Statement of authorship: JST, JAM, AEZ and CEE developed and designed the study. JST, CL,
AFF, MIL, GA and MJM collected the Madidi Project dataset; MS, WFR, KGC, NSR, and YM
collected the ABERG dataset. SAS produced the phylogenetic data. AGL and JST performed
data analyses. AGL and JST wrote the manuscript, and all authors contributed significantly to
revisions.

Data accessibility statement: The Madidi Project's dataset used in our analyses correspond to
version 4.1, which is deposited in Zenodo (https://doi.org/10.5281/zenodo.4276558).
Additionally, raw data of the Madidi Project are stored and managed in Tropicos
(https://tropicos.org/home), the botanical database of the Missouri Botanical Garden. These data
can be viewed and accessed via the Madidi Project's module at
http://legacy.tropicos.org/Project/MDI. The Andes Biodiversity and Ecosystem Research Group
(ABERG) is a team of 38 researchers from 12 universities dedicated to understanding
biodiversity distribution and ecosystem function in the Peruvian Andes. ABERG is committed to
data exchange within the scientific community and promoting collaboration among other tropical
ecosystem scientists. For more information and to request data contact Miles Silman or
Yadvinder Malhi (http://www.andesconservation.org/). The R code created for analyses is
available at https://github.com/Linan552/Madidi-project.

Keywords: Andes; community assembly; phylogenetics; turnover; neotropics; elevational

gradient; adaptive diversification; dispersal; species sorting

Abstract

Historical biogeographic events such as mountain orogeny are associated with the creation of environmental gradients, giving rise to the assembly of communities of species observed today. However, key gaps remain in our understanding of the relative importance of different eco-evolutionary processes acting as drivers of community assembly across environmental gradients. In this study, we test two non-exclusive hypotheses of the eco-evolutionary processes that shape tree communities across the Central Andean elevational gradient: Communities are assembled via 1) immigration and ecological sorting of pre-adapted clades, and 2) recent adaptive diversification along the elevational gradient. We used species surveys in the Bolivian and Peruvian Andes and a novel phylogenetic framework to test the relative importance of these hypotheses. Although adaptive diversification has previously been observed in specific clades, immigration and sorting of clades pre-adapted to montane habitats is the primary mechanism shaping communities across elevations.

Introduction

Large-scale biogeographic events—such as the emergence of novel environmental conditions, biotic interchanges, or the evolution of key innovations—can have lasting consequences for biodiversity, community assembly, and species distributions (Ricklefs 2006; Fussmann *et al.* 2007; Pelletier *et al.* 2009; Claramunt & Cracraft 2015; Uribe-Convers & Tank 2015). Although theory and empirical evidence suggest that processes occurring in the deep past can contribute to the modern structure of local ecological communities, most research in community ecology during the last few decades has been dominated by a focus on mechanisms at small spatial and temporal scales (Ricklefs 1987). Studies largely overlook the broader biogeographic context in which communities of co-occurring species are embedded (Chesson 2000; Adler *et al.* 2007). Only recently have ecologists begun bridging this gap by developing ecological theory and empirical tests that truly integrate community assembly across eco-evolutionary scales (Emerson & Gillespie 2008; McGill *et al.* 2019; Bañares-de-Dios *et al.* 2020; Segovia *et al.* 2020). The extent to which community assembly is contingent upon regional context and biogeographic history has broad implications for ecological and evolutionary theory and for understanding how and why communities respond to environmental change (Chase 2003; Fukami 2015; Vellend 2016; McPeek 2017).

Recent studies provide important insights into how ongoing ecological processes change along environmental gradients (Bricca *et al.* 2019; Bañares-de-Dios *et al.* 2020; Neves *et al.* 2020). However, much less is known about how the emergence of the gradients themselves shape the evolution of species and phenotypes that underlie community assembly. Two non-mutually exclusive processes may explain how communities assemble along gradients following the

emergence of novel environmental conditions (Fig. 1). First, the emergence of new

environments—e.g., due to climate change, island formation, or mountain orogeny— may create

opportunities for immigration and ecological sorting of pre-adapted clades (ISPC hypothesis;

Box 1; Fig. 1A; Donoghue 2008). According to this hypothesis, when environmental conditions

change within a region and new gradients are created, community assembly across these new

habitats is dominated by the immigration of species that are pre-adapted because they occupy

similar habitats in a different region. This means that the combination of traits needed to colonize

new habitats evolved before the origin of the environmental gradient. Diversification following

colonization would not involve adaption to novel environments (i.e. niche conservatism) owing,

for example, to competition with species pre-adapted to other environments (Fukami 2015;

Tanentzap *et al.* 2015). Thus, even though diversification might occur after the origin of the

gradient, new species would be restricted mainly to environments to which their ancestors were

already pre-adapted. In this way, community assembly across environmental gradients would

result in the ecological sorting of species within clades that predate the new environments in the

region. This scenario of community assembly is consistent with the idea that "it is easier to move

than to evolve" (Donoghue 2008).

Second, the emergence of new environments may create opportunities for recent adaptive

diversification across environments (the RAD hypothesis; Box 1; Fig. 1A). According to this

hypothesis, when new environmental gradients are created, community assembly across habitats

is dominated by adaptation in response to the emerging environmental conditions, resulting in

the diversification of clades across the environmental gradient (Stroud & Losos 2016). Thus, the

traits needed to colonize new habitats evolve after the origin of the environmental gradient. In

this scenario, niche conservatism is minimal or non-existent, and community assembly results

from the diversification of one or more clades that were originally adapted to a subset of

environmental conditions, but that diversify to occupy emerging novel environmental space. This

scenario for community assembly following the emergence of environmental gradients is

consistent with the classic ideas of biome shifts and adaptive radiation driven by ecological

opportunity (Schluter 2000; Losos 2010; Donoghue & Edwards 2014).

Here we present and test a novel community-phylogenetic framework and method to

disentangle the relative importance of ISPC and RAD in determining the assembly of

communities along large-scale environmental gradients. These effects on community assembly

can be inferred from unique patterns in the phylogenetic structure of compositional turnover. In

particular, signatures of these two processes can be traced when species turnover is decomposed

into components that correspond to *within-* and *among-clade turnover*, where clades correspond

to independent lineages that originated before the emergence of the gradient. These within- and

among-clade turnover components, in turn, reflect the effects of diversification after and before

the emergence of the gradient on community composition across environments. Here, we

illustrate these patterns using a hypothetical elevational gradient of a mountain depicted in Fig.

1A. The ISPC hypothesis predicts that for communities *at the same elevation*, variation in

community composition should be dominated by *within-clade turnover*, reflecting strong niche

conservatism of a few clades that are pre-adapted to the environments at that specific elevation

(Fig. 1B). As communities are *farther apart along the elevational (i.e. environmental) gradient*,

variation in community composition should become increasingly dominated by *among-clade*

turnover, reflecting the shift in dominance from species in one pre-adapted clade to another.

Alternatively, the RAD hypothesis predicts that for communities at *similar or contrasting*

elevations, variation in community composition should be dominated by turnover *within clades,*

reflecting how multiple clades evolved niche differences in response to new environmental

conditions that allow them to have broad elevational (i.e., environmental) distributions (Fig. 1C).

Although we developed and tested this conceptual framework in the context of mountain uplift,

our approach is applicable to study community assembly after the emergence of any type of

environmental gradient at any spatial or temporal scale.

In the Neotropics, the geologically recent uplift of the Andean mountains created a striking

elevational and environmental gradient that had profound consequences for global climate and

biodiversity (Rahbek & Graves 2001; Antonelli *et al.* 2009; Ehlers & Poulsen 2009; Graham

2009; Jiménez *et al.* 2009; Hoorn *et al.* 2010). Indeed, the tropical Andes are considered the most

species-rich biodiversity hotspot, containing 15% of all plant species (>45,000 species) in only

1% of the world's land area (Myers *et al.* 2000; Rahbek & Graves 2001; Jiménez *et al.* 2009).

However, our current understanding of the eco-evolutionary forces that shape community

assembly across elevations in the hyper-diverse Andean biotas is limited. First, many studies

focus on the evolution and distribution of relatively small clades compared to entire

communities; these studies have provided evidence for an important role of adaptive

diversification (Antonelli *et al.* 2009; Pérez-Escobar *et al.* 2017) in some cases and immigration

and colonization of pre-adapted clades in others (Hughes & Eastwood 2006; Jin *et al.* 2015;

Lagomarsino *et al.* 2016). Such studies demonstrate that both processes have occurred, but

provide limited insights into how evolutionary history of individual clades contribute to the

assembly of entire ecological communities and regional biotas. Second, studies that focus on the

phylogenetic structure of Andean communities are relatively few and often fail to differentiate

the effects of diversification before and after the emergence of the gradient (Graham *et al.* 2009;

Parra *et al.* 2011; Bacon *et al.* 2018; Montaño-Centellas *et al.* 2019; Ramírez *et al.* 2019). To

date, no study has sought to disentangle the relative importance of immigration and sorting of

pre-adapted clades versus post-Andean uplift adaptive radiation in shaping the enormous

variation in plant community composition across elevational gradients.

In this study, we combined tree-species distribution data and phylogenetic information from

two large networks of Andean-forest plots to test how RAD and ISPC contribute to the assembly

of Andean tree communities. We test these hypotheses in the context of the uplift of the Central

Andes, which is associated with the formation of the Altiplano plateau during the last 30 my.

(Fig. 1). Moreover, we developed a novel method to decompose measures of species turnover

among plots distributed across the elevational gradient into among- and within- pre-Andean

clade components (Fig. 1 and Box 1; Legendre & Cáceres 2013). These components measure the

relative contributions of ISPC and RAD, respectively. This work provides both a novel

framework for examining phylogenetic community turnover and expands our current

understanding of how historical processes contribute to community assembly.

Methods

Community composition data across elevations

We utilized data from two large-scale forest plot networks in the Central Andes of Bolivia

(the Madidi Project; 30,165 km^2) and Peru (the Andes Biodiversity and Ecosystem Research

Group [ABERG]; 1,765 km^2). Both datasets contain information on tree community composition

spanning the entire elevational range of forests in this region of the Andes from lowland

Amazonia to the tree line. Our datasets include information on species composition across 73 1-

ha plots (large plots hereafter; 50 in Bolivia and 23 in Peru), as well as 494 0.1-ha plots (small

plots hereafter; 458 in Bolivia and 36 in Peru; Fig. 2). Within these plots, all woody plants with a

diameter at breast height (DBH) $\geq$ 10 cm in large plots and DBH $\geq$ 2.5 cm in small plots were

tagged, measured and identified to species or morpho-species. Large and small plots characterize

different plant communities; while large plots consider only adults of large tree species, small

plots include younger individuals and also species that do not reach 10 cm DBH, including many

shrubs. Thus, we separated out data by plot size and conducted analyses independently.

Additionally, we excluded high elevation plots (> 3,800 m) and plots with $\leq$ 3 species. Most of

these plots represent *Polylepis*-dominated forests fragments within a matrix of Paramo

grasslands/shrublands. The ecology and composition of these Paramo forests is clearly distinct

from the continuous forest cover along the elevational gradient.

Within the Bolivian and Peruvian datasets, we conducted extensive taxonomic work to

standardize species and morpho-species names across plots. Morpho-species, however, could not

be standardized between the Bolivian and Peruvian data. To test the effect of morphospecies on

results, analyses were repeated with and without morphospecies. Both analyses produced nearly

identical results (Fig. S1); for simplicity, we present only analyses including morphospecies.

Representative specimens at each site were collected and deposited in herbaria, mainly at the

Herbario Nacional de La Paz (LPB), the Missouri Botanical Garden (MO) and Universidad

Nacional de San Antonio Abad del Cusco (CUZ) in Peru. The final dataset contains 494 small

plots and 73 large plots, distributed from 175 m to 4,365 m in elevation. The small plots

contained 2,731 species, whereas the large plots contained 1,904 species (Table 1).

Phylogenetic reconstruction and defining clades of pre-Andean origin

To test our hypotheses, we needed a phylogenetic framework that grouped species into clades

that diverged from one-another before the origin of the elevational gradient (i.e. clades that pre-

date the uplift of the Central Andes). To do this, we based our analyses on Smith and Brown's (

2018) global mega-phylogeny of seed plants, which combined NCBI sequence data, results from

the Open Tree of Life project (Hinchliff *et al.* 2015), and advances in bioinformatic methods

(PyPHLAWD; Smith & Walker 2019) to produce the most comprehensive time-calibrated

species-level phylogeny to date. To include species and morphospecies in our dataset that were

not in the original phylogeny, we used the R package V.PhyloMaker (Jin & Qian 2019). Using

genus and family level taxonomic information, missing taxa not included in the mega-phylogeny

were joined to the halfway point of the family/genus branch (V.PhyloMaker scenario= "S3").

For genera not represented in the mega-phylogeny, we joined species to sister genera in the

phylogeny based on support in the literature (when possible) using the 'bind.relative' option of

V.PhyloMaker. Finally, we pruned from the phylogeny, all species that were absent in our forest

plots. The resulting phylogeny included 3,143 species.

The formation of the Andean cordillera has been a complex and heterogeneous process. In

the Central Andes, the history of mountain formation is closely tied to the development of the

Altiplano plateau, currently located at nearly 3,800 m in elevation. While the traditional view of

mountain uplift invokes a slow and gradual process, recent evidence suggests that the uplift of

the Altiplano was dominated by spurs of rapid rise with intervening periods of stasis (Garzione *et*

al. 2008, 2017). Although the northern Andes is considered much younger, the best available

evidence suggests that most of the uplift in the Central Andes occurred within the last 30 million

years. Thus, our analyses used this age as a main reference for the origin of the elevational

gradient and to delimit pre-Andean clades.

Pre-Andean clades in the time-calibrated regional phylogeny were defined as those whose

stems intersect the 30 my reference. In this way, each pre-Andean clade in our study diverged

from others before the uplift of the central Andes, whereas all species within pre-Andean clades

resulted from diversification that occurred after mountain uplift had started. We used the

function treeSlice in the R package "Phytools" (Revell 2012) to fragment the regional phylogeny

into these clades. Species present in small plots formed 473 pre-Andean clades with an average

of 5.77 species per clade, whereas species in the large plots formed 355 clades, averaging 5.36

species per clade (Table 1 and Fig. S2). Finally, we sought see understand how our results varied

by defining different ages for pre-Andean clades. Thus, in addition to creating a dataset with pre-

Andean clades defined as 30 million years old, we made a second dataset which defined clades

as 60 million years old. This represents a much more conservative estimate of the timing of

Andean uplift (Hoorn *et al.* 2010). The results from these alternative analyses were nearly

identical, and thus are presented only in the Supplementary Material (Fig. S3).

Decomposing total turnover into among- and within-clade turnover

To test hypotheses about the relative importance of ISPC and RAD, we developed a method

to decompose variation in species turnover between two communities into additive components

representing the contribution of turnover among-groups and within-groups (Legendre & Cáceres

2013). For our analyses, groups are defined by clades of pre-Andean origin, but this

decomposition method is broadly applicable to species groupings based on any criteria. Analyses

were based on the Sørensen pair-wise dissimilarity index (S; Sørensen 1948), which uses

presence/absence data:

273
$$S = \frac{b+c}{2a+b+c}$$

Here, a represents the number of shared species between two communities, b is the number

of species present only in the first community, and c is the number of species present only in the

second community. Since species are aggregated into clades, species in b can be further divided

into two components: b_{WG} is the fraction of b that correspond to species in groups present in both

communities, while b_{AG} is the fraction of b corresponding to species in groups present only in the

first community. The same process can be done for c, producing the corresponding components

c_{WG} and c_{AG}. In this way, the additive within-group (S_{WG}) and among-groups (S_{AG}) components

of Sørensen dissimilarity are defined as:

$$S_{WG} = \frac{b_{WG} + c_{WG}}{2 \times a + b + c}$$

$$S_{AG} = \frac{b_{AG} + c_{AG}}{2 \times a + b + c}$$

Further details of the decomposition method can be found in the Supplementary Material,

where we also show that this approach could be applied to other turnover metrics, such as Bray-

Curtis distances. The R code that performs this decomposition is available at

https://github.com/Linan552/Madidi-project. When within- and among-clade dissimilarities are

transformed into components of total turnover (S_{WG}/S and S_{AG}/S, respectively), these values

correspond to the contribution of diversification after (S_{WG}/S) and before (S_{AG}/S) the uplift of the

central Andes to community species turnover (Fig. 1, S4). Thus, a high among-clade component

indicates that turnover is mainly dominated by species that diverged from one another before the

uplift of the Central Andes (Fig. 1A, left). In contrast, high within-clade component indicates

that turnover is dominated by species that diverged from one another after the uplift of the

Central Andes (Fig. 1A, right).

293   As described in the introduction, the immigration and sorting of pre-adapted clades

(ISPC) and the recent adaptive diversification (RAD) hypotheses make predictions about how

these components of turnover will be related to environmental (i.e. elevational) distances. Thus,

we plotted the components of turnover for each pair of plots against their elevational distance.

The ISPC hypothesis predicts that as communities are farther apart along the elevational

gradient, variation in community composition should become increasingly dominated by among-

clade turnover (Fig. 1B). Alternatively, the RAD hypothesis predicts that variation in community

composition should be dominated by turnover within clades regardless of elevational distance

(Fig. 1C).

Assessing significance of empirical data using null models and ruling out effects of

geographic distance

To test whether observed patterns are different from those expected by chance, we compared

the components of turnover in the empirical data with components produced by a null model that

eliminated any phylogenetic structure in the distribution of species, but retained other important

elements of the data that might shape turnover patters. We ran a "tip-randomization null model"

in which species were randomly re-assigned to tips in the phylogeny, such that species were

randomly reshuffled among pre-Andean clades. This randomization algorithm maintained the

number of species per clade, the diversity gradient across elevations, the average range size in

each community, and importantly, the empirical turnover observed between pairs of plots. The

only aspect of the data that was randomized was the empirical membership of species in clades

of pre-Andean origin. We randomized the data and re-calculated components of turnover for

each pair of plots 999 times. From these null expectations, we calculated standardized effect

sizes as the empirical value minus the mean of null distribution divided by the standard deviation

of the null distribution. These values represent the magnitude of the difference between the

empirical components of turnover and the null expectation, where there is no phylogenetic

structure in species distributions. As was done for the empirical components of turnover, we

related these standardized effect values against the difference in elevation for each pair of plots.

Additionally, we compared the rate of change in turnover components with difference in

elevation between the empirical data and null model expectations. To do this, we used slopes

from a linear regression between components of turnover and elevational distance. Because these

relationships are non-linear, we used a logit transformation on species turnover prior to

regression analyses (Cleveland 1981). These transformations produced a reasonable linearization

of the relationships in large and small plot datasets, allowing us to capture the rate of change in a

single parameter (see Fig. S5). The empirical slopes were then compared with the distribution of

999 slopes generated by the null model. If empirical slopes were significantly greater or smaller

($p<0.05$) than null slopes, we concluded that phylogenetic structure exists in the community

composition changes along the elevational gradient.

Finally, to control for the effects of geographic distance on the analysis of turnover across

elevations, only a subset of pair-wise plot comparisons for each dataset were used. This subset of

plot pairs minimized variation in geographic distances, but maximized the elevational range

represented in the data (Fig. S6). For the large-plot dataset, we selected plot pairs only between

50 and 90 km apart (8% of the total range in geographic distances), and for the small-plot

dataset, we selected plot pairs between 110 and 160 km apart (10% of geographic range). In both

datasets, the elevational distances between plots ranged from 175 to 3765 m.

Additionally, we analyzed how the within- and among-clade components of turnover changes

as a function of geographic distance. For these analyses, we used pairs of plots spanning the

entire geographic range of the study (max distance between plots; 495 km), but were at similar

elevations (0 to 200 m of elevational distance, Fig. S6). Like our analyses along elevational

distances, we compared patterns of variation in the within- and among-clade turnover against

null model expectations. These analyses show how turnover and its components change across

space but within the same environmental conditions (results presented in the Supplementary

Material).

Results

Species composition changed dramatically across elevations. Species turnover (Sørensen

dissimilarity) among forest plots showed a saturating relationship with elevational distance,

increasing rapidly as elevational distance increases and then reaching an asymptote at complete

turnover (Fig. 3). Indeed, plots separated by more than 2,000 to 2,500 m of elevation never

shared species. We found a similar relationship between species composition and geographic

distance (Fig. S7). However, Sørensen dissimilarity did not increase as dramatically with

increasing geographic distance and it never reached complete turnover; for example, we found

that plots can still share species when they are in similar environments even if plots are 400 km

away from one another (one in Peru the other in Bolivia; Fig. S7).

We also found strong elevational gradients in the within- and among-clade components of

species turnover (Fig. 4A&D). For forest plots occurring at the same elevation (zero meters in

elevational difference), among- and within-clade components were equal in magnitude (Fig.

4A&D). This result indicates that communities in the same environment shared species in the

same pre-Andean clades, but also that multiple different clades contributed to community

composition among these plots. As elevational difference increased, among-clade turnover rose

rapidly, while within-clade turnover decreased (Fig. 4A&D). Although the increase in the

among-clade component was monotonic in the small-plots dataset, it saturated at around 2,000 m

of elevational difference for the large-plots dataset. In both datasets, however, when plots were

separated by more than 1,000 to 1,500 m in elevation, pairs of communities were found where

100% of the turnover corresponded to the among-clade component. This result means that some

pairs of plots at opposite ends of the elevational gradient shared neither species nor clades 30 my

old, which originated before the uplift of the Central Andes. These results support the idea that

the immigration and ecological sorting of pre-Andean clades had a major effect in shaping

community composition across elevations - the ISPC hypothesis.

The predictions of the ISPC hypotheses were also supported using standardized effect sizes –

as measured using our null model (Fig. 4 C&F). Indeed, standardized effect sizes for within- and

among-clade components were both close to zero for plots at the same elevation. As elevational

differences increased, standardized effect sizes increased for among-clade turnover and

decreased for within-clade turnover (Fig. 4 C&F). Moreover, when plots were separated by

>1,500 m in elevation, the empirical values differed by more than two standard deviations from

null expectations (i.e. standardized effect sizes greater than 2; Fig. 4C and 4D). The comparison

of the empirical and null regression slopes also showed that the change in the empirical

components was much more pronounced than the change expected by the null model (Fig. 5).

Finally, we found that geographic distance did not have the same effect on components of

turnover as elevational difference. Among large plots, among- and within-clade turnover

remained constant and of similar magnitude with increasing geographic distance (Fig. 4 vs. S8).

For small plots, on the other hand, the magnitude of the within-clade turnover component

increases with geographic distance. This pattern remained when using standardized effect sizes

(Fig. 4 vs. S8).

Discussion

__Community assembly across contrasting elevations is dominated by the immigration and__

__ecological sorting of clades that pre-date mountain uplift__

Our results showed clearly that changes in species composition across elevations were driven

primarily by a replacement of clades of pre-Andean origin. These results were robust to analyses

using different age estimates of pre-Andean clades (30 vs. 60 mya), inclusion or exclusion of

morpho-species, or delimitations of forest communities (trees $\geq$ 10 cm DBH in large plots vs.

trees $\geq$ 2.5 cm DBH in small plots). While adaptive diversification is likely to have occurred in

our study system, our results suggest that this process has had a reduced influence on patterns of

community assembly. In contrast, we found strong evidence for a high relative importance of the

ISPC hypothesis. The new environments created by the uplift of the Central Andes during the

last 30 my were colonized primarily by clades of species that were pre-adapted to the emerging

environmental conditions. Diversification within these clades resulted in new tree species that

had elevational distributions similar to those occupied by the immigrating species. In this way,

the ecological sorting of pre-Adapted clades according to their pre-adaptations is the eco-

evolutionary process that dominates the regional assembly of tree communities across the

elevational gradient.

Our study focuses on the structure of species assemblages, and how biogeographic processes

shape patterns of diversity. The assembly of communities, however, integrates the evolutionary

history of multiple independent clades of species. Several previous studies that have taken this

approach, focusing on the evolution of clades after the Andean uplift. This research shows that

groups of animals and plants across the Andes have diversified in ways that are consistent with

our results (Bell & Donoghue 2005; Hughes & Eastwood 2006; Chaves *et al.* 2011; Nürk *et al.*

2013). One of the best studied biogeographic histories in the Andes is that of the plants in the

genus *Lupinus*, which colonized the Andes from temperate North America (Hughes & Eastwood

2006), and were likely pre-adapted to the cold conditions of alpine environments (Nevado *et al.*

2016). This clade experienced an explosive diversification in the Andes, but most of the resulting

species occupy only high-elevation habitats. Their diversification was likely fueled by the

interaction between insularity of high-mountain habitats and climatic fluctuations during

Quaternary (Nevado *et al.* 2018). Adaptive diversification also played an important role in the

radiation of the Andean lupins (Nevado *et al.* 2016). Indeed, species in the clade show a huge

diversity of phenotypes, life forms and micro-habitat use (Hughes & Eastwood 2006). Their

adaptive diversification, however, did not involve large numbers of species colonizing the new

environments at different elevations created by mountain uplift. Several clades of plants

distributed at the highest elevations in the Andes seem to show similar patterns of diversification

(Madriñán *et al.* 2013). A general pattern of conservatism in elevational distribution was also

documented for several clades of trees by Griffiths et al. (2020). Clades with a biogeographic

history similar those of the Andean lupins would contribute little to changes in species

composition along the elevational gradient. Instead, this pattern of diversification, when

experienced by numerous clades pre-adapted to different elevations can lead to the observed

patterns of clade turnover found in our study. Indeed, turnover among communities across the

elevational gradient have an evolutionary origin that is rooted deep in the past, and that mostly

pre-dates the emergence of the environmental gradient.

Studies of particular clades, like those highlighted above, are insightful and have helped us

advance our understanding of the patterns and mechanisms of diversification. However, this

approach does not address directly the eco-evolutionary forces behind the assembly of diverse

communities, which is the focus of our analyses. To the best of our knowledge, our study is the

first effort to explicitly test the role that diversification before and after the origin of the

environmental gradient (i.e., the uplift of the Central Andes) had on community structure across

elevations. While previous studies have not tested the role of mountain uplift directly, our results

are supported by previous research of Andean communities, which have suggested an important

role for niche conservatism in community assembly across elevations (Graham *et al.* 2009;

Hardy *et al.* 2012; Jin *et al.* 2015; Ramírez *et al.* 2019; Worthy *et al.* 2019; Bañares-de-Dios *et*

al. 2020). A recent important study in this respect is that by Segovia et al. (2020), who

demonstrated a clear link in the phylogenetic composition of Andean tree communities to

temperate regions of North and South America. In particular, they highlight the role that freezing

conditions at high elevations play in creating environments that are invaded by temperate clades.

Similarly, niche conservatism has been implied in the eco-evolutionary assembly of seasonally

dry forest communities, which occur in rain-shadowed valleys along the Andes. Our study,

however, goes further than simply demonstrating niche conservatism or phylogenetic clustering

of communities. Instead, we provide evidence that the assembly of communities across

elevations is primarily driven by the immigration and sorting of clades that evolved appropriate

adaptations even before the emergence of the environmental gradient (Hardy *et al.* 2012; Chi *et*

al. 2014; Kubota *et al.* 2018).

Mountain uplift might create opportunities for adaptive radiation, but this process has a

limited effect on community assembly along elevational gradients

Adaptive radiations have played a critical role in the formation of biodiversity, giving rise to

an often-stunning array of morphological and species diversity (Gillespie *et al.* 2020). Previous

studies have suggested that ecological opportunity is an important determinant, maybe a

prerequisite, of adaptive radiations (Stroud & Losos 2016; Gillespie *et al.* 2020), allowing

species to diversify rapidly to fill available niche space. The uplift of the Central Andes created

environments that were previously unavailable in the region, likely opening up new unoccupied

niche space for species. Moreover, as we discussed earlier, numerous rapid radiations have been

documented in the Andes (Madriñán *et al.* 2013); some of them, like that of *Lupinus* or *Espeletia*

(Hughes & Eastwood 2006; Pouchon *et al.* 2018) are as dramatic as those in clades that

epitomize adaptive radiation (e.g. stickleback fish or African Great Lake cichlids; Gillespie *et al.*

2020). If ecological opportunity existed and rapid diversification in the mountains is well

documented, then why did we not find a strong signal for recent adaptive radiation in the

assembly of communities across elevations?

There are several reasons that could explain our lack of evidence for recent adaptive

radiations across the elevational gradient. First, recent and rapid radiations in the Andes may not

involve adaptive diversification. Instead, high rates of species accumulation could be fueled

solely by allopatric speciation resulting from repeated cycles of habitat isolation and re-

connection driven by climatic oscillations (Nevado *et al.* 2018; Flantua *et al.* 2019). This process

would produce a large number of species that replace one another across geography but within

the same environment (Hughes & Eastwood 2006; Chaves *et al.* 2011). Furthermore, nearly all

cases of recent montane radiations occur in clades occupying the highest elevations (Bell &

Donoghue 2005; Hughes & Eastwood 2006; Nürk *et al.* 2013; Hughes & Atchison 2015), and

more studies are needed to know if these patterns of diversification also occur at mid or low

elevations (but see Lagomarsino *et al.* 2016). Second, adaptive radiations may have occurred

along environmental dimensions other than those of the elevational gradient. Indeed, some of the

classic examples of Andean diversification involve fast evolution of phenotypes, even if the

elevational distribution of the clade is highly conserved (Hughes & Eastwood 2006; Nürk *et al.*

2018; Pouchon *et al.* 2018). Finally, some clades may have adaptively radiated across the

elevational gradient, but these clades are rare and contribute little to overall assembly patterns.

Indeed, biogeographic studies have documented significant shifts in elevational distribution

during the evolutionary history of several groups of plants and animals (Elias *et al.* 2009; Bacon

et al. 2018). Some of these shift in elevational distribution might be accompanied by shifts in life

form (as exemplified by *Espeletia*; Pouchon *et al.* 2018; but see Zanne *et al.* 2013) which do not

contribute to the assembly of tree communities that are the focus of our study. The relative

frequency of adaptive diversification across elevations versus niche conservatism has not been

evaluated; however, Griffith et al. (2020) found that in Peru, most clades of trees have narrow

elevational distributions, and only a few show wider elevational distributions than expected by

chance. The overall lack of clades that appear to have adaptively radiated across elevations could

be due to evolutionary priority effect, whereby different elevational niches may have been

preempted by immigrating pre-adapted clades (Fukami 2015). While the role of adaptive

radiation in community assembly deserves further study, our results suggest that Andean

community assembly is mainly the result of different pre-adapted clades that originated before

Andean uplift, which colonized available niches before other clades could adaptively radiate to

occupy a broad elevational gradient (Tanentzap *et al.* 2015).

Conclusions: future directions and implications for conservation

In this study, we developed a novel conceptual framework (Fig. 1), as well as new methods

of decomposing species turnover (Fig. 1 & S4), to investigate the biogeographic origins of

community assembly along environmental gradients. We use this approach to study how the

uplift of the Central Andes led to the variation in community composition along iconic

elevational gradients. Our approach, however, can be applied to any system in which the timing

of the emergence of an environmental gradient is known and time-calibrated phylogenies can be

generated. We envision future studies using this method to understand the eco-evolutionary

assembly in systems beyond Andean forests such as along precipitation gradients (Parolari *et al.*

2020), across contrasting soil conditions (Capurucho *et al.* 2020), or even under different

disturbance regimes (Cavender-Bares & Reich 2012). Methods such as these can be used to test

hypotheses about specific process of community assembly, going beyond documenting niche

conservatism or phylogenetic aggregation. Our approach will facilitate deeper insights into how

the emergence of environmental gradients shape modern natural ecosystems.

Our analyses demonstrate that species turnover across elevations in the Central Andes is

driven primarily by the turnover of clades that are at least 30 my old. These results suggest a

strong role for immigration and ecological sorting of pre-adapted clades to the novel

environments across elevations created by the uplift of the Central Andes. Adaptive

diversification following the emergence of the elevational gradient is likely restricted to a few

clades or to narrow elevational bands, having little impact on the assembly of communities along

such a large environmental gradient. Our results add to a growing body of evidence suggesting

that present day communities are strongly influenced by the ability of lineages to track

environmental conditions through space and geological time (Donoghue 2008; Emerson &

Gillespie 2008; Carvajal-Endara *et al.* 2017; Griffiths *et al.* 2020; Segovia *et al.* 2020).

This finding has important implications for the long-term persistence of communities facing

the effects of human-mediated global change. Increases in atmospheric temperatures are

predicted to cause elevational shifts in environmental conditions, such that climates that

currently occur at specific elevations will occur at higher elevations in the future (Harsch *et al.*

2009; Ruiz-Labourdette *et al.* 2012; Freeman *et al.* 2018; O'Sullivan *et al.* 2020). Our work on

historical patterns of community assembly suggests that ecosystems are more likely to track

shifting habitats rather than adapt to novel conditions (Sheldon *et al.* 2011; Ruiz-Labourdette *et*

al. 2012; Freeman *et al.* 2018; Feeley *et al.* 2020). Communities and species at the highest

elevations might be specially threatened by climate change since their environments will

disappear at the top of mountains and new pre-adapted competitors will move in from lower

elevations (Colwell *et al.* 2008). Thus, communities occupying the highest-elevation sites in the

Andes should be prioritized for monitoring and conservation efforts. Because their habitat may

not persist over the long term, ex situ conservation (either through conservation seed banking or

living collections) of the species endemic to the highest elevations should be a specific priority.

Acknowledgements

We thank the Dirección General de Biodiversidad, the Bolivian Park Service (SERNAP),

the Madidi National Park and local communities for permits, access, and collaboration in

Bolivia, where fieldwork was supported by the National Science Foundation (DEB 0101775,

DEB 0743457, DEB 1836353). Additional financial support to The Madidi Project has been

provided by the Missouri Botanical Garden, the National Geographic Society (NGS 7754-04 and

NGS 8047-06), the Comunidad de Madrid (Spain), Consejo Superior de Investigaciones

Científicas (Spain), Centro de Estudios de América Latina (Banco Santander and Universidad

Autónoma de Madrid, Spain), and the Taylor and Davidson families. Fieldwork in the ABERG

transect was supported by NSF, the Gordon and Betty Moore Foundation and the UK Natural

Environment Research Council. We thank all the researchers, students and local guides that were

involved in the collection of the data, particularly Carla Maldonado, Maritza Cornejo, Alejandro

Araujo, Javier Quisbert, Narel Paniagua and Peter Jørgenson. Finally, we thank Iván Jiménez for

helpful discussions, ideas and comments.

References

Adler, P.B., HilleRislambers, J. & Levine, J.M. (2007). A niche for neutrality. *Ecol. Lett.*, 10, 95–104.

Antonelli, A., Nylander, J.A.A., Persson, C. & Sanmartin, I. (2009). Tracing the impact of the Andean uplift on Neotropical plant evolution. *Proc. Natl. Acad. Sci.*, 106, 9749–9754.

Bacon, C.D., Velásquez-Puentes, F.J., Hoorn, C. & Antonelli, A. (2018). Iriarteeae palms tracked the uplift of Andean Cordilleras. *J. Biogeogr.*, 45, 1653–1663.

Bañares-de-Dios, G., Macía, M.J., Granzow-de la Cerda, Í., Arnelas, I., Martins de Carvalho, G., Espinosa, C.I., *et al.* (2020). Linking patterns and processes of tree community assembly across spatial scales in tropical montane forests. *Ecology*, 101.

Bell, C. & Donoghue, M.J. (2005). Phylogeny and biogeography of Valerianaceae (Dipsacales) with special reference to the South American valerians. *Org. Divers. Evol.*, 5, 147–159.

Bricca, A., Conti, L., Tardella, M.F., Catorci, A., Iocchi, M., Theurillat, J.P., *et al.* (2019). Community assembly processes along a sub-Mediterranean elevation gradient: analyzing the interdependence of trait community weighted mean and functional diversity. *Plant Ecol.*, 220, 1139–1151.

Capurucho, J.M.G., Borges, S.H., Cornelius, C., Vicentini, A., Prata, E.M.B., Costa, F.M., *et al.* (2020). Patterns and Processes of Diversification in Amazonian White Sand Ecosystems: Insights from Birds and Plants. In: *Neotropical diversification: patterns and processes* (eds. Rull, V. & Carnaval, A.C.). Springer Nature, Cham, Switzerland, pp. 245–270.

Carvajal-Endara, S., Hendry, A.P., Emery, N.C. & Davies, T.J. (2017). Habitat filtering not dispersal limitation shapes oceanic island floras: species assembly of the Galápagos archipelago. *Ecol. Lett.*, 20, 495–504.

Cavender-Bares, J. & Reich, P.B. (2012). Shocks to the system: community assembly of the oak savanna in a 40-year fire frequency experiment. *Ecology*, 93, S52--S69.

Chase, J.M. (2003). Community assembly: when should history matter? *Oecologia*, 136, 489–498.

Chaves, J.A., Weir, J.T. & Smith, T.B. (2011). Diversification in Adelomyia hummingbirds follows Andean uplift. *Mol. Ecol.*, 20, 4564–4576.

Chesson, P. (2000). Mechanisms of Maintenance of Species Diversity. *Annu. Rev. Ecol. Syst.*, 31, 343–366.

Chi, X., Tang, Z. & Fang, J. (2014). Patterns of phylogenetic beta diversity in China's grasslands in relation to geographical and environmental distance. *Basic Appl. Ecol.*, 15, 416–425.

Claramunt, S. & Cracraft, J. (2015). A new time tree reveals Earth history's imprint on the evolution of modern birds. *Sci. Adv.*, 1, e1501005.

Cleveland, W.S. (1981). LOWESS: A program for smoothing scatterplots by robust locally weighted regression. *Am. Stat.*, 35, 54.

Colwell, R.K., Brehm, G., Cardelús, C.L., Gilman, A.C., Longino, J.T., Cardelus, C.L., *et al.*
(2008). Global Warming, Elevational Range Shifts, and Lowland Biotic Attrition in the Wet
Tropics. *Science (80-.).*, 322, 258–261.

Donoghue, M.J. (2008). A phylogenetic perspective on the distribution of plant diversity. *Proc.*
Natl. Acad. Sci., 105, 11549–11555.

Donoghue, M.J. & Edwards, E.J. (2014). Biome shifts and niche evolution in plants. *Annu. Rev.*
Ecol. Evol. Syst., 45, 547–572.

Ehlers, T.A. & Poulsen, C.J. (2009). Influence of Andean uplift on climate and paleoaltimetry
estimates. *Earth Planet. Sci. Lett.*, 281, 238–248.

Elias, M., Joron, M., Willmott, K., Silvia-Brandao, K.L., Kaiser, V., ARIAS, C.F., *et al.* (2009).
Out of the Andes: patterns of diversification in clearwing butterflies. *Mol. Ecol.*, 18, 1716–
1729.

Emerson, B.C. & Gillespie, R.G. (2008). Phylogenetic analysis of community assembly and
structure over space and time. *Trends Ecol. Evol.*, 23, 619–630.

Feeley, K.J., Bravo-Avila, C., Fadrique, B., Perez, T.M. & Zuleta, D. (2020). Climate-driven
changes in the composition of New World plant communities. *Nat. Clim. Chang.*, 10, 965–
970.

Flantua, S.G.A., O'Dea, A., Onstein, R.E., Giraldo, C. & Hooghiemstra, H. (2019). The
flickering connectivity system of the north Andean páramos. *J. Biogeogr.*, 46, 1808–1825.

Freeman, B.G., Scholer, M.N., Ruiz-Gutierrez, V. & Fitzpatrick, J.W. (2018). Climate change
causes upslope shifts and mountaintop extirpations in a tropical bird community. *Proc. Natl.*
Acad. Sci. U. S. A., 115, 11982–11987.

Fukami, T. (2015). Historical Contingency in Community Assembly: Integrating Niches, Species
Pools, and Priority Effects. *Annu. Rev. Ecol. Evol. Syst.*, 46, 1–23.

Fussmann, G.F., Loreau, M. & Abrams, P.A. (2007). Eco-evolutionary dynamics of communities
and ecosystems. *Funct. Ecol.*, 21, 465–477.

Garzione, C.N., Hoke, G.D., Libarkin, J.C., Withers, S., MacFadden, B., Eiler, J., *et al.* (2008).
Rise of the Andes. *Science*, 320, 1304–1307.

Garzione, C.N., McQuarrie, N., Perez, N.D., Ehlers, T.A., Beck, S.L., Kar, N., *et al.* (2017).
Tectonic Evolution of the Central Andean Plateau and Implications for the Growth of
Plateaus. *Annu. Rev. Earth Planet. Sci.*, 45, 529–559.

Gillespie, R.G., Bennett, G.M., Meester, L. De, Feder, J.L., Fleischer, R.C., Harmon, L.J., *et al.*
(2020). Comparing Adaptive Radiations Across Space, Time, and Taxa. *J. Hered.*, 1–20.

Graham, A. (2009). The Andes: A geological overview from a biological perspective. *Ann.*
Missouri Bot. Gard., 96, 371–385.

Graham, C.H., Parra, J.L., Rahbek, C. & McGuire, J. a. (2009). Phylogenetic structure in tropical
hummingbird communities. *Proc. Natl. Acad. Sci.*, 106, 19673–19678.

Griffiths, A.R., Silman, M.R., Farfán Rios, W., Feeley, K.J., Cabrera, G., Meir, P., *et al.* (2020).
Evolutionary heritage shapes tree distributions along an Amazon-to-Andes elevation
gradient. *Biotropica*, 1–13.
Hardy, O.J., Couteron, P., Munoz, F., Ramesh, B.R. & Pélissier, R. (2012). Phylogenetic
turnover in tropical tree communities: Impact of environmental filtering, biogeography and
mesoclimatic niche conservatism. *Glob. Ecol. Biogeogr.*, 21, 1007–1016.
Harsch, M.A., Hulme, P.E., McGlone, M.S. & Duncan, R.P. (2009). Are treelines advancing? A
global meta-analysis of treeline response to climate warming. *Ecol. Lett.*, 12, 1040–1049.
Hinchliff, C.E., Smith, S.A., Allman, J.F., Burleigh, J.G., Chaudhary, R., Coghill, L.M., *et al.*
(2015). Synthesis of phylogeny and taxonomy into a comprehensive tree of life. *Proc. Natl.*
Acad. Sci., 112, 12764–12769.
Hoorn, W., Steeg, H., Bermudez, M.A., Mora, A., Sevink, J., Sanmartin, I., *et al.* (2010).
Amazonia Through Time: Andean Uplift, Climate Change, Landscape Evolution, and
Biodiversity. *Science (80-.).*, 330, 927–931.
Hughes, C. & Eastwood, R. (2006). Island radiation on a continental scale: Exceptional rates of
plant diversification after uplift of the Andes. *Proc. Natl. Acad. Sci.*, 103, 10334–10339.
Hughes, C.E. & Atchison, G.W. (2015). The ubiquity of alpine plant radiations: From the Andes
to the Hengduan Mountains. *New Phytol.*, 207, 275–282.
Jiménez, I., Distler, T. & Jørgensen, P.M. (2009). Estimated plant richness pattern across
northwest South America provides similar support for the species-energy and spatial
heterogeneity hypotheses. *Ecography (Cop.).*, 32, 433–448.
Jin, L.S., Cadotte, M.W. & Fortin, M.J. (2015). Phylogenetic turnover patterns consistent with
niche conservatism in montane plant species. *J. Ecol.*, 103, 742–749.
Jin, Y. & Qian, H. (2019). V.PhyloMaker: an R package that can generate very large phylogenies
for vascular plants. *Ecography (Cop.).*, 42, 1353–1359.
Kubota, Y., Kusumoto, B., Shiono, T. & Ulrich, W. (2018). Environmental filters shaping
angiosperm tree assembly along climatic and geographic gradients. *J. Veg. Sci.*, 29, 607–
618.
Lagomarsino, L.P., Condamine, F.L., Antonelli, A., Mulch, A. & Davis, C.C. (2016). The abiotic
and biotic drivers of rapid diversification in Andean bellflowers (Campanulaceae). *New*
Phytol., 210, 1430–1442.
Legendre, P. & Cáceres, M. De. (2013). Beta diversity as the variance of community data:
dissimilarity coefficients and partitioning. *Ecol. Lett.*, 16, 951–963.
Losos, J.B. (2010). Adaptive radiation, ecological opportunity, and evolutionary determinism.
Am. Nat., 175, 623–39.
Madriñán, S., Cortés, A.J. & Richardson, J.E. (2013). Páramo is the world's fastest evolving and
coolest biodiversity hotspot. *Front. Genet.*, 4, 1–7.

McGill, B.J., Chase, J.M., Hortal, J., Overcast, I., Rominger, A.J., Rosindell, J., *et al.* (2019). Unifying macroecology and macroevolution to answer fundamental questions about biodiversity. *Glob. Ecol. Biogeogr.*, 28, 1925–1936.

McPeek, M.A. (2017). *Evolutionary Community Ecology*. Princeton University Press, Princeton.

Montaño-Centellas, F.A., McCain, C. & Loiselle, B.A. (2019). Using functional and phylogenetic diversity to infer avian community assembly along elevational gradients. *Glob. Ecol. Biogeogr.*, 1–14.

Myers, N., Mittermeier, R.A., Mittermeier, C.G., da Fonseca, G.A.B. & Kent, J. (2000). Biodiversity hotspots for conservation priorities. *Nature*, 403, 853–858.

Nevado, B., Atchison, G.W., Hughes, C.E. & Filatov, D.A. (2016). Widespread adaptive evolution during repeated evolutionary radiations in New World lupins. *Nat. Commun.*, 7, 1–9.

Nevado, B., Contreras-Ortiz, N., Hughes, C. & Filatov, D.A. (2018). Pleistocene glacial cycles drive isolation, gene flow and speciation in the high-elevation Andes. *New Phytol.*, 219, 779–793.

Neves, D.M., Dexter, K.G., Baker, T.R., Coelho de Souza, F., Oliveira-Filho, A.T., Queiroz, L.P., *et al.* (2020). Evolutionary diversity in tropical tree communities peaks at intermediate precipitation. *Sci. Rep.*, 10, 1–7.

Nürk, N.M., Michling, F. & Linder, H.P. (2018). Are the radiations of temperate lineages in tropical alpine ecosystems pre-adapted? *Glob. Ecol. Biogeogr.*, 27, 334–345.

Nürk, N.M., Scheriau, C. & Madriñán, S. (2013). Explosive radiation in high Andean Hypericum-rates of diversification among New World lineages. *Front. Genet.*, 4.

O'Sullivan, K.S.W., Ruiz-Benito, P., Chen, J.C. & Jump, A.S. (2020). Onward but not always upward: individualistic elevational shifts of tree species in subtropical montane forests. *Ecography (Cop.).*, 1–12.

Parolari, A.J., Paul, K., Griffing, A., Condit, R., Perez, R., Aguilar, S., *et al.* (2020). Liana abundance and diversity increase with rainfall seasonality along a precipitation gradient in Panama. *Ecography (Cop.).*, 43, 25–33.

Parra, J.L., Rahbek, C., McGuire, J.A. & Graham, C.H. (2011). Contrasting patterns of phylogenetic assemblage structure along the elevational gradient for major hummingbird clades. *J. Biogeogr.*, 38, 2350–2361.

Pelletier, F., Garant, D. & Hendry, A.P. (2009). Eco-evolutionary dynamics. *Philos. Trans. R. Soc. B Biol. Sci.*, 364, 1483–1489.

Pérez-Escobar, O.A., Chomicki, G., Condamine, F.L., Karremans, A.P., Bogarín, D., Matzke, N.J., *et al.* (2017). Recent origin and rapid speciation of Neotropical orchids in the world's richest plant biodiversity hotspot. *New Phytol.*, 215, 891–905.

Pouchon, C., Fernández, A., Nassar, J.M., Boyer, F., Aubert, S., Lavergne, S., *et al.* (2018).

Phylogenomic analysis of the explosive adaptive radiation of the Espeletia complex (Asteraceae) in the tropical Andes. *Syst. Biol.*, 67, 1041–1060.

Rahbek, C. & Graves, G.R. (2001). Multiscale assessment of patterns of avian species richness. *Proc. Natl. Acad. Sci.*, 98, 4534–4539.

Ramírez, S., González-Caro, S., Phillips, J., Cabrera, E., Feeley, K.J. & Duque, Á. (2019). The influence of historical dispersal on the phylogenetic structure of tree communities in the tropical Andes. *Biotropica*, 51, 500–508.

Revell, L.J. (2012). phytools: An R package for phylogenetic comparative biology (and other things). *Methods Ecol. Evol.*, 3, 217–223.

Ricklefs, R.E. (1987). Community Diversity: Relative Roles of Local and Regional Processes. *Science (80).*, 235, 167–171.

Ricklefs, R.E. (2006). Evolutionary diversification and the origin of the diversity-environment relationship. *Ecology*, 87, 3–13.

Ruiz-Labourdette, D., Nogués-Bravo, D., Ollero, H.S., Schmitz, M.F. & Pineda, F.D. (2012). Forest composition in Mediterranean mountains is projected to shift along the entire elevational gradient under climate change. *J. Biogeogr.*, 39, 162–176.

Schluter, D. (2000). *The ecology of adaptive radiation*. Oxford University Press.

Segovia, R.A., Pennington, R.T., Baker, T.R., de Souza, F.C., Neves, D.M., Davis, C.C., *et al.* (2020). Freezing and water availability structure the evolutionary diversity of trees across the Americas. *Sci. Adv.*, 6.

Sheldon, K.S., Yang, S. & Tewksbury, J.J. (2011). Climate change and community disassembly: Impacts of warming on tropical and temperate montane community structure. *Ecol. Lett.*, 14, 1191–1200.

Smith, S.A. & Brown, J.W. (2018). Constructing a broadly inclusive seed plant phylogeny. *Am. J. Bot.*, 105, 302–314.

Smith, S.A. & Walker, J.F. (2019). Py phlawd: A python tool for phylogenetic dataset construction. *Methods Ecol. Evol.*, 10, 104–108.

Sorensen, T.A. (1948). A method of establishing groups of equal amplitude in plant sociology based on similarity of species content and its application to analyses of the vegetation on Danish commons. *Biol. Skar.*, 5, 1–34.

Stroud, J.T. & Losos, J.B. (2016). Ecological Opportunity and Adaptive Radiation. *Annu. Rev. Ecol. Evol. Syst.*, 47, 507–532.

Tanentzap, A.J., Brandt, A.J., Smissen, R.D., Heenan, P.B., Fukami, T. & Lee, W.G. (2015). When do plant radiations influence community assembly? The importance of historical contingency in the race for niche space. *New Phytol.*, 207, 468–479.

Uribe-Convers, S. & Tank, D.C. (2015). Shifts in diversification rates linked to biogeographic movement into new areas: An example of a recent radiation in the Andes. *Am. J. Bot.*, 102,

1854–1869.

Vellend, M. (2016). *The Theory of Ecological Communities*. *Theory Ecol. Communities*.
Princeton University Press, Princeton.

Worthy, S.J., Jiménez Paz, R.A., Pérez, Á.J., Reynolds, A., Cruse-Sanders, J., Valencia, R., *et al.*
(2019). Distribution and community assembly of trees along an andean elevational gradient.
Plants, 8, 7–10.

Zanne, A.E., Tank, D.C., Cornwell, W.K., Eastman, J.M., Smith, S.A., FitzJohn, R.G., *et al.*
(2013). Three keys to the radiation of angiosperms into freezing environments. *Nature*, 506,
89–92.

746
747

BOX 1 - Glossary

Pre-Andean clade: A clade that diverged from others before the uplift of the Central Andes. Fig. 1 shows predicted elevational distributions of three pre-Andean clades (colors) based on our hypotheses.

Pre-adapted clade: A pre-Andean clade that had, before its immigration to the Central Andes, already evolved adaptations to the novel environmental conditions created by mountain uplift.

Turnover: Observed variation in species or functional-trait composition among forest plots or biogeographic regions. For example, in two species assemblages [A, B] & [A, C], turnover is generated by the replacement of species B in the first assemblage with species C in the second.

Within-clade turnover: Proportion of total turnover that corresponds to shifts in species composition within a pre-Andean clade. For example, in two assemblages [A, B] & [A, C], within-clade turnover would be high if species B and C belong to the same pre-Andean clade.

Among-clade turnover: Proportion of total turnover that corresponds to shifts in species composition among multiple pre-Andean clades. For example, in two assemblages [A, B] & [A, C], among-clade turnover would be high if species B and C belong to different pre-Andean clades.

748
749

750 **Table 1.** Summary of datasets used for analyses and p-values assessing significance of empirical gradients in among-clade and
751 within-clade turnover across elevational (elev) and geographic (geo) distances.
752

Morpho-species status	No. species	Clade age	No. clades	No. clades with one sp.	Mean species per clade	P (among-clade vs. elev)	P (within-clade vs. elev)	P (among-clade vs. geo)	P (within-clade vs. geo)
Large plots (73)									
Included	1889	30	352	202	5.37	0.001	0.001	0.553	0.553
Included	1889	60	142	34	13.30	0.001	0.001	0.424	0.424
Excluded	1349	30	309	168	4.37	0.001	0.001	0.223	0.223
Excluded	1349	60	139	34	9.71	0.001	0.001	0.87	0.87
Small plots (404)									
Included	2326	30	425	239	5.47	0.001	0.001	0.025	0.025
Included	2326	60	159	34	14.63	0.001	0.001	0.005	0.005
Excluded	1741	30	386	213	4.51	0.001	0.001	0.002	0.002
Excluded	1741	60	154	36	11.31	0.001	0.001	0.001	0.001

Figures

Figure 1. Conceptual models to explain the assembly of regional biotas after the emergence of new environments. (A) shows the distributions of species (symbols), traits (sizes) and clades (colors) along an elevational gradient as expected by the ISPC (left) and RAD (right) hypotheses. The gray broken line marks the emergence of the novel environmental conditions due to mountain uplift. The phylogeny describes the evolutionary relationships among species in the target communities, and the colors indicate different clades of pre-Andean origin (clades that diverged before the uplift of the Central Andes). ISPC and RAD predict contrasting spatial patterns in how species turnover is partitioned into within- and among-clade components. ISPC (A, left) – If the uplift of the Central Andes created opportunities for immigration of pre-adapted clades, communities at different elevations will be assembled by species of different pre-Andean clades. (B) If ISPC is the dominant scenario of community assembly, the among-clade component will increase rapidly as differences in elevation between plots increase, while the within-clade component will decrease. These two components will intersect, and the among-

clade component will become the most important. Within a specific elevation, ISPC also predicts
that the within-clade component will be consistently more important as geographic distance
increases. RAD (A, right) – If mountain uplift created opportunities for adaptive diversification,
environments at different elevations will be occupied by species of the same pre-Andean clades.
(C) If RAD is the dominant scenario, the within-clade component will dominate the turnover of
species both as elevational differences or geographic distances increase.

Figure 2. Regional network of forest plots used in this study. The study region is in the
Central Andes, particularly the Madidi National Park in Bolivia (the Madidi Project) and the
Manu National Park in Peru (ABERG). In both of these regions, we have sampled forest
communities across extensive elevational gradients that go from the Amazon to the tree line
(4,190 m difference in elevation). In Madidi and Manu, we have data from 73 permanent 1-ha

plots (large plots). Within these plots, all woody plants with a diameter at breast height (DBH)
equal or greater than 10 cm have been tagged, measured, and identified. Additionally, in Madidi
and Manu, we have data from 494 temporary 0.1-ha plots (small plots), where woody plants with
a DBH $\geq$ 2.5 cm have been surveyed. Our analyses have been repeated for large and small plot
separately.

Figure 3. Species turnover across elevations. Sørensen dissimilarity plotted against difference
in elevation for each pair of plots in our two datasets. These patterns are presented separately for
(A) large 1-ha plots and (B) small 0.1-ha plots. These analyses demonstrate that as difference in
elevation increases, dissimilarity in species composition between plot also increases. This
dissimilarity is exemplified by the lack of shared species between Amazonian forests (C; Laguna
Chalalan in Bolivia at 400 m elevation) and upper montane cloud forests (D; Trocha Union in
Peru at 3,260 m). Pictures by Christopher Davidson, Sharon Christoph and William Farfan-Rios.
799

Figure 4. Decomposition of species turnover across elevational gradients into among-clade and within-clade components – 30 MY clades from small and large plots. Sorensen dissimilarities between each pair of plots were decomposed into within-clades (blue lines) and among-clades (yellow lines) components. We then plotted these components of turnover against difference in elevation in large plots (first row) and small plots (second row). Finally, we compared spatial patterns in variation of these components with a tip-randomization null model that removes any phylogenetic structure in the distribution of species across elevation. (A. & D.) empirical patterns; (B. & E.) patterns for the mean of the expectations in the null model; (C & F) patterns for standardized effect sizes showing the deviation of the empirical values from null expectations.

813
814
815 **Figure 5. Comparison of linear slopes.** Empirical slopes of logit transformed proportional
816 turnover across elevational distance (triangles) compared to the distribution of 999 slopes
817 produced by the null model (boxplots) in (A) large plots and (B) small plots.
818
819
820