The single *Marchantia polymorpha FERONIA* homolog reveals an

ancestral role in regulating cellular expansion and integrity

Martin A. Mecchia[1#], Moritz Rövekamp[1#+], Alejandro Giraldo-Fonseca[1], Dario Meier[1],

Philippe Gadient[1], Hannes Vogler[1], Daria Limacher[1], John L. Bowman[2], and Ueli

Grossniklaus[1]*

[1]Department of Plant and Microbial Biology and Zurich-Basel Plant Science Center, University of

Zurich, 8008 Zurich, Switzerland

[2]School of Biological Sciences, Monash University, Clayton, Melbourne, Victoria 3800, Australia

[+]Current Affiliation: Kantonsschule Uetikon am See, 8708 Uetikon am See, Switzerland

#These authors contributed equally to the manuscript.

*Author for correspondence: grossnik@botinst.uzh.ch

Running tittle: Ancestral role of *Cr*RLK1L receptors in plants

Summary statement

The *Cr*RLK1L family arose in land plants and the *Marchantia polymorpha genome*

contains a single copy*,* Mp*FER,* which is broadly expressed and regulates cell expansion

and cell wall integrity.

Abstract

Plant cells are surrounded by a cell wall, a rigid structure that is not only important for cell

and organ shape, but crucial for intercellular communication and interactions with the

environment. In the flowering plant *Arabidopsis thaliana*, the 17 members of the

Catharanthus roseus RLK1-like (*Cr*RLK1L) receptor kinase family are involved in a multitude of physiological and developmental processes, making it difficult to assess their primary or ancestral function. To reduce genetic complexity, we characterized the single *Cr*RLK1L gene of *Marchantia polymorpha*, Mp*FERONIA* (Mp*FER*). Plants with reduced Mp*FER* levels show defects in vegetative development, i.e., rhizoid formation and cell expansion, and have reduced male fertility. In contrast, Mp*fer* null mutants and overexpression lines severely affect cell integrity and morphogenesis of the gametophyte. Thus, the *Cr*RLK1L gene family originated from a single gene with an ancestral function in cell expansion and the maintenance of cellular integrity. During land plant evolution, this ancestral gene diversified to fulfil a multitude of specialized physiological and developmental and roles in the formation of both gametophytic and sporophytic structures essential to the life cycle of flowering plants.

Key Words

cell wall integrity, cell elongation, *Cr*RLK1L receptor kinases, *Marchantia polymorpha*, signaling pathway evolution

Introduction

The functioning of plant cells is contingent upon the cell wall acting as a barrier between the cell and its environment. This complex matrix of polysaccharides, glycoproteins, and other organic compounds defines the growth and shape of a cell and provides protection against biotic and abiotic stresses [1], [2], [3]. Moreover, the cell wall has to resist turgor pressure while also allowing cell expansion during growth. Thus, sensing and controlling cell wall integrity (CWI) is crucial for plant cells, and receptor kinases (RKs) play an

important role in sensing extracellular cues that activate intracellular pathways. Different RK subfamilies are defined by their extracellular domain (ECD), which is more variable than their transmembrane and kinase domains [4]. Over the last decade, the *Cr*RLK1L subfamily emerged as important CWI sensors [5], [6].

Functions have been assigned to 15 of the 17 *Cr*RLK1L members encoded in the *A. thaliana* genome: they are required for sensing environmental cues and cell-cell communication in diverse contexts, including reproduction, hormone signaling, cell expansion, innate immunity, and various stress responses [7]–[23]. For instance, *FERONIA* (At*FER*), *THESEUS1* (At*THE1*), *HERCULES1* (At*HERK1*), and At*HERK2* are required for cell elongation as At*fer* single, At*the1*At*herk1* double, and At*the1*At*herk1*At*herk2* triple mutants exhibit stunted growth [8], [10]–[12]. Root hairs burst in mutants affecting At*FER* and *[Ca^{2+}]$_{cyt}$-ASSOCIATED PROTEIN KINASE1* (At*CAP1*)/*ERULUS* (At*ERU*) [13], [17]. Moreover, At*FER* plays a role in powdery mildew resistance [12], innate immunity [16], [23], calcium signaling [24], phytohormone signaling [11], [25]–[27], and mechano- and heavy metal sensing [21], [28]. In most contexts, the *Cr*RLK1Ls tend to promote cellular growth and cell elongation. In contrast, At*THE1* actively inhibits cell elongation in hypocotyls when cell wall perturbations occur, *e.g.*, due to mutations in the cell wall biosynthesis machinery or upon treatment with isoxaben, a cellulose synthesis inhibitor [8], [29].

In addition to regulating cellular growth during the vegetative phase of the life cycle, several *Cr*RLK1Ls play a role during fertilization by controlling the growth and reception of the pollen tube. In the female gametophyte, At*FER* is highly expressed in the two synergid cells that flank the egg cell and are important for double fertilization [7], [30]. At*FER* is crucial for pollen tube reception and the release of the two sperm cells [7], [11], [24], [31], [32], a process also involving two other, redundantly acting synergid-expressed

*Cr*RLK1Ls, *AtHERK1* and *ANJEA* (*AtANJ*) [18]. Moreover, *AtFER* and *AtANJ* also play a

role in pollen-stigma recognition during pollen germination [1]. In contrast to these

synergid-expressed *Cr*RLK1Ls, four pollen-expressed *Cr*RLK1Ls, named *ANXUR1*

(At*ANX1*), At*ANX2, BUDDHA'S PAPER SEAL1* (At*BUPS1*) and At*BUPS2* [9], [15], [19],

[33], are redundantly required for pollen tube integrity and tip growth.

To transduce extracellular cues, *Cr*RLK1Ls bind secreted peptides of the RAPID

ALKALINIZATION FACTOR (RALF) family [23], [33]–[36], which induce complex

formation with members of LORELEI (LRE) family of GPI-anchored proteins [33], [36]–

[38]. Hence, it was suggested that at least some *Cr*RLK1Ls act as CWI sensors that

coordinate cell elongation in response to changes in the cell wall [39]–[44]. A role as CWI

sensors is also consistent with recent findings that AtFER interacts with pectin [43], [45],

[46].

The 17 *A. thaliana Cr*RLK1Ls act partially redundantly and sometimes even have

opposite effects on cellular growth (reviewed in [20], [42], [44], [47]). To investigate the

original function of *Cr*RLK1Ls, we characterized them in a system with reduced genetic

complexity: the liverwort *Marchantia polymorpha* encodes a single *Cr*RLK1L [48], [49].

Liverworts represent an early diverging land plant lineage and have been hypothesized

to retain, at least in part, characteristics of the earliest land plants [48], [50]–[55]. The

presence of similar types of gene families in all plant lineages suggests that the

differences in development evolved by co-opting and modifying existing developmental

programs and genetic networks, rather than through the evolution of novel genes [50],

[51], [56]. *M. polymorpha* is thus an ideal system to study the ancestral role of genes and

how it was modulated and diversified during land plant evolution [48], [57], [58].

We refer to the single *Cr*RLK1L encoded in the *M. polymorpha* genome, Mp4g15890

(Mapoly0869s0001), as Mp*FERONIA* (Mp*FER*) [47]–[49] (aka Mp*THESEUS,* Mp*THE*)

[59]. An Mp*fer/the* mutant was identified in a large T-DNA screen for defective rhizoid

formation, developing short and irregularly shaped rhizoids with brown tips, indicating

rhizoid rupture [59]. In the same screen, a mutant affecting the ortholog of the *A. thaliana*

receptor-like cytoplasmic kinase *MARIS* (At*MRI*), known to act downstream of At*FER*,

was identified with a similar phenotype (named Mp*mri* aka Mp*pti*, Mp7g17560

[Mapoly0051s0094]) [59], [60]. As in *A. thaliana*, a dominant active version of Mp*MRI* can

partially rescue the Mp*fer* rhizoid phenotype [60]. This suggests that at least some of the

machinery associated with cell elongation and CWI sensing is conserved between *A.*

thaliana and *M. polymorpha*.

Here, we report that Mp*FER* function is required for a variety of aspects of *M. polymorpha*

gametophyte development. Phylogenetic analyses suggest that *Cr*RLK1L genes first

appeared in the common ancestor of land plants. The characterization of lines with

reduced Mp*FER* levels indicates that, in addition to its function in rhizoid formation,

Mp*FER* controls cell size and organ growth and is involved in male gametogenesis and

fertility. Analysis of the Mp*FER* expression pattern points to an involvement in female

sexual organogenesis and sporophyte development. Our data suggests that the broad

involvement of *Cr*RLK1L function in the development and physiology of land plants is an

ancestral and conserved characteristic, although the functions of *Cr*RLK1Ls were

extended and adapted to control additional developmental processes in the sporophyte

during the course of land plant evolution.

Results

***Cr*RLK1L is conserved in land plants and arose in this lineage**

The *M. polymorpha* genome encodes a single *Cr*RLK1L homolog (Mp4g15890

[Mapoly0869s0001]) [47]–[49]. Like other *Cr*RLK1L members, this gene encodes an

intracellular serine/threonine kinase domain, a transmembrane domain, and an

extracellular malectin-like domain with two malectin domains (Fig. 1A). Given that the

ECD of *Cr*RLK1Ls in *A. thaliana* is crucial for their function and, in contrast to the kinase

domain, not interchangeable [61], we performed a phylogenetic analysis with the amino

acid sequences of the predicted malectin-like domain of Mp4g15890, the *Cr*RLK1Ls of

the mosses *Physcomitrium patens* (5 members), *Sphagnum fallax* (7 members),

Ceratodon purpureus (3 members), the hornworts *Anthoceros agrestis Bonn* (one

member)*, Anthoceros agrestis* Oxford (one member)*, Anthoceros punctatus* (one

member)*, the lycophyte *Selaginella moellendorffii* (two members), the basal angiosperm

Amborella trichopoda (nine members)*, and the 17 *Cr*RLK1Ls of *A. thaliana* (Fig. 1B). The

*M. polymorpha Cr*RLK1L member grouped together with the *Cr*RLK1Ls of basal land

plants and was in the same clade as, e.g., AtFER, AtHERK2, and AtANX1/2, while

AtTHE1 was in a different one (Fig. 1B). Based on this phylogenetic position and the fact

that the malectin-like domain of the *M. polymorpha Cr*RLK1L shared the highest amino

acid identity with AtFER (Fig. S1A), we named it MpFER [48], [49], [60].

Interestingly, no significant similarity to the MpFER malectin-like domain was found when

compared to available genomic data of six Chlorophyte algae (*Chlamydomonas*

reinhardtii, Dunaliella salina, Volvox carteri, Coccomyxa subellipsoidea, Micromonas

pusilla, *Ostreococcus lucimarinus*) and expression data of six Charophycean algae

(*Klebsormidium nitens, Nitella mirabilis, Coleochaete orbicularis, Spirogyra pratensis,*

Mesostigma viride, Closterium peracerosum–strigosum–littorale complex). In *Closterium*

complex, an RK named CpRLK1 expressed during sexual reproduction was suggested

to be a *Cr*RLK1L homolog [62]. However, comparing the full sequence or the ECD of

CpRLK1 with the predicted *M. polymorpha* proteome, higher identity to another RK was

found (Mp1g17720 [Mapoly0001s0111]) (Figs S1B,C). Moreover, no RALF orthologs

have been reported in algae, suggesting the origin of the *Cr*RLK1L pathway in land plants

[48]. A similar conclusion was recently reached by analyzing the RALF family in *P. patens*

[63]. Moreover, the best candidate obtained by BLAST analysis using the MpFER

malectin-like domain against the genome of the Charophycean alga *Chara braunii*

clustered outside the *Cr*RLK1L phylogenetic tree (Figs. 1B, S1B).

To further support MpFER as a member of the *Cr*RLK1L subfamily, the three-dimensional

structure of its ECD was modelled [64] using the structure of the AtFER ECD (PDB

6a5b.1A, [36]) as a template (Fig. 1C). Overall, the predicted structure of the MpFER ECD

is highly similar to that of AtFER, keeping an almost identical number of secondary

structures (7 alpha-helices and 33 beta-sheets in the MpFER ECD [36] and 8 alpha-helix

and 34 beta-sheets in the AtFER ECD, respectively), and a similar spatial disposition for

32 of the beta-sheet structures and 6 of the alpha-helices (Fig. 1C). When comparing the

structural superposition of the AtFER and MpFER ECDs (Figs. 1D and S1D), we

observed that most conserved parts of the malectin-like domain are in the core of the

protein, while the more variable parts are located peripherally. Moreover, the 3D structure

of the CpRLK1 ECD could not be properly modelled using the AtFER ECD (Fig. S1D,E),

neither the one from *C. braunii* (Fig. S1D,F). Taken together, these results suggest that

the malectin-like domain of MpFER could interact with similar proteins as AtFER does in

A. thaliana and may thus be involved in similar pathways. Sequence comparison of the

MpFER kinase domain showed a conservation of the activation loop and a Lys important

for catalytic activity (Fig. 1E). Furthermore, several determined phosphorylation sites

found in AtFER, AtERU, and CrRLK1 were conserved in MpFER (Fig. 1E) [14], [34], [65],

[66]. Given that the genomes of *M. polymorpha* and *Anthoceros* spp. encode a single

*Cr*RLK1L and this RK family is not present in Charophycean algae, we propose MpFER

to be basal and orthologous to all other land plant *Cr*RLK1Ls, and that the *Cr*RLK1L family

arose as plants conquered land.

Mp*FER* is broadly expressed throughout the *M. polymorpha* life cycle

In *A. thaliana*, the expression patterns of the 17 *Cr*RLK1L members span vegetative and

reproductive stages and both generations of the life cycle (reviewed in [44]), pointing to

the importance of *Cr*RLK1Ls for basic cellular functions. To assess whether this pattern

is reflected in the land plant with the most ancestral characteristics, a promoter fragment

of 3.2 kb (_{pro}Mp*FER*) was used to drive expression of a triple yellow fluorescent protein

(trpVNS) with a nuclear localization signal (_{pro}Mp*FER*:*trpVNS*) and transformed into *M.*

polymorpha sporelings. All transformants expressing the trpVNS fluorescent protein (45

of 48 independent lines) exhibited an indistinguishable expression pattern (Fig. 2). During

vegetative stages of gametophyte development, expression was observed in most cells

of the thallus (Fig. 2A). In concordance with the importance of Mp*FER* for rhizoids [59],

high trpVNS expression was observed in these tip-growing cells (Fig. 2B). While trpVNS

was expressed in gemma cups, no expression was observed in mature gemmae, possibly

due to their dormancy (Fig. 2C).

In mature female sexual organs, strong expression was observed in the entire

archegoniophore (Fig. 2D). The archegonia expressed trpVNS in most cells, except for

the egg cell (Fig. 2E). However, after fertilization, _{pro}Mp*FER*:*trpVNS* became active in the

zygote (Fig. 2F). In male sexual organs, trpVNS was broadly detected in the antheridial

splash platform (Fig. 2G) and, specifically, in the spermatogenous tissue of the antheridia

and the non-reproductive jacket cells surrounding them at different developmental stages

(Fig. 2H). Inside the sporogonium, the sporogenous cells and subsequently the

developing spores, but not the elater cells, strongly expressed trpVNS (Fig. 2I).

Taken together, Mp*FER* is expressed in most tissues during vegetative stages of

gametophyte development as well as in the antheridia and archegonia during the sexual

reproduction. Interestingly, the ₚᵣₒMp*FER* promoter is not active in mature gemmae and

the unfertilized, quiescent egg cell, although expression is initiated after dormancy and in

the zygote. In concordance with the collectively ubiquitous expression of *Cr*RLK1Ls in *A.*

thaliana, the Mp*FER* expression pattern reinforces the importance of this family for basic

cellular functions in land plants.

Mp*FER* controls cell expansion during vegetative gametophyte development

To analyze the function of Mp*FER*, three independent artificial microRNA constructs

targeting Mp*FER* were designed (amiR-Mp*FER*), based on the endogenous microRNA

Mp*miR160* (Fig. S2, [67]) and driven by the ubiquitous ₚᵣₒMp*EF1* promoter [68]. While

amiR-Mp*FER*1 and amiR-Mp*FER*2 target sites in the ECD coding sequence, amiR-

Mp*FER*3 targets the sequence encoding the kinase domain (Fig. 3A,B). For each amiR-

Mp*FER* construct, several independent transgenic lines were obtained that showed a

reduction in thallus size (Figs. 3C,D and S3A). Mp*FER* levels were quantified by qRT-

PCR in two independent amiR-Mp*FER*2 and amiR-Mp*FER*3 lines each, which showed a

similar phenotype (Fig. 3E). Mp*FER* expression ranged from 10% to 20% of the wild-type

level in all amiR-Mp*FER* lines analyzed (Fig. 3E).

Previously, the T-DNA insertion mutant Mp*fer-1* was identified in a screen for mutants

with defective rhizoids ([59], referred to as Mp*the*). As the T-DNA inserted into the 3' UTR

of the Mp*FER* gene (Fig. 3A; [59]), Mp*FER* expression was analyzed by qRT-PCR but it

showed no reduction compared to the wild type (Fig. S3B), suggesting that the T-DNA

insertion does not affect Mp*FER* transcription. Sequencing of the Mp*FER* coding

sequence in the Mp*fer-1* mutant did not show any mutation, while 3' RACE-PCR showed

that transcript terminated about 1.3 kb inside the T-DNA and was not polyadenylated (Fig.

S3C). Because thallus size of Mp*fer-1* mutants is similar to that of amiR-Mp*FER* lines

(Figs. 3C,D and S3A) and efficient translation depends on the presence of a poly(A)-tail

[69] , we propose that MpFER protein levels are reduced in the Mp*fer-1* mutant despite

normal transcript levels.

Rhizoids are tip-growing cells analogous to the root hairs in flowering plants

(angiosperms). Reduced Mp*FER* activity strongly impaired rhizoid formation and the

rhizoids collapsed (Fig. S3D). This had also been reported for the Mp*fer-1* mutant [59]

and is in concordance with the function of At*FER* and At*ANX1/2* in tip-growing root hairs

[13], [17], [34], [70] and pollen tubes [9], [15], respectively. As *Cr*RLK1L members are

important for cell expansion in angiosperms [8], [10], [12], [20], [35], [71], [72], cell size of

epidermal thallus cells in fully differentiated regions with minimal growth was measured

in Mp*fer-1* and amiR-Mp*FER* lines, showing a reduction in cell area as compared to the

wild type (Figs. 3F and S3E,F). The average epidermal cell area was similarly reduced,

although the distribution of cell areas differed between amiR-Mp*FER* lines and Mp*fer-1*

(Figs. 3F and S3E,F).

In conclusion, Mp*FER* has a fundamental role in rhizoid formation and growth during

vegetative development. These results suggest an ancestral and conserved function of

the *Cr*RLK1Ls in polar cell growth and cell expansion.

Mp*FER* is important for the morphological integrity of the gametophyte

Given that Mp*fer-1* and amiR-Mp*FER* lines retain some MpFER activity, we generated

Mp*fer* knock-out mutants with the goal to unveil potentially hidden *Cr*RLK1L functions by

generating a plant without any *Cr*RLK1L activity. Using CRISPR/Cas9, two sites in the

ECD-coding sequence of Mp*FER* were targeted (Fig. 4A,B). At least 10 independent lines

were selected and analyzed for each target site. All plants with severely affected thallus

development contained a frame shift mutation at the respective target site (Fig. 4B,C)

while Mp*FER* was not affected in transgenic plants with normal development.

The thallus area of the newly generated, amorphic knock-out mutants was more strongly

reduced compared to Mp*fer-1* [59], confirming that, like the amiR-Mp*FER* lines, Mp*fer-1*

is a hypomorphic mutant (Fig. 4C,D). In contrast, rhizoid integrity was similarly affected

in hypo- and amorphic Mp*fer* alleles, suggesting that rhizoid formation is more sensitive

to reduced Mp*FER* activity than thallus growth (Fig. 4E,F). Despite of the strong impact

on thallus development, both hypo- and amorphic Mp*fer* mutants form gemma cups.

The strong disruption of thallus growth in amorphic Mp*fer* mutants prompted us to

investigate whether cells of the thallus died in the absence of Mp*FER* activity. Indeed,

using tryphan blue staining, we detected dead cells in these mutants, indicating a loss of

morphological integrity (Fig. 4G). To investigate the mechanical properties of thallus cells,

we measured their apparent stiffness using cellular force microscopy on 3-day old

gemmalings and found that amorphic Mp*fer-2* mutants had a lower apparent stiffness

than the wild type (Fig. 4H). The apparent stiffness of the thalli depends on both the

stiffness of the cell wall and the turgor pressure of the cells. Since the apparent stiffness

is lower in Mp*fer-2* mutants, it is likely that a reduction in the stiffness their cell wall causes

the cells to burst and die.

Some reproductive but not all CWI sensing functions of *Cr*RLK1Ls are conserved

in land plants

In *A. thaliana*, at least seven *Cr*RLK1L family members play a role in reproduction, with

At*ANX1/2* and At*BUPS1/2* being important for pollen tube growth [9], [15], [19], [33] and

At*FER*, At*HERK1*, and At*ANJ* for pollen tube reception by the synergid cells [7], [18]. To

characterize the function of Mp*FER* in reproductive development, amiR-Mp*FER3* lines

were transferred to sexual organ-inducing conditions [73]. The number of

antheridiophores produced was significantly less in amiR-Mp*FER* lines compared to the

wild type (Figs. 5A and S3G) and the antheridiophore splash platforms were also smaller

(Fig. 5B). The reduction in Mp*FER* expression in antheridiophores of amiR-Mp*FER* lines

was confirmed by qRT-PCR and results were comparable to those using thallus tissue

(Fig. 5C). A pronounced reduction in spermatogenous tissue was observed in many

antheridia isolated from amiR-Mp*FER*3-3 lines (Fig. 5D,E). Whether this phenotype is

caused by reduced cell expansion as was observed in thalli or by a lower cell division rate

is currently unclear. Moreover, although the plants were grown under optimal conditions

in axenic culture boxes, the effects on reproductive structures could be partly indirect due

to the reduction in plant size and/or rhizoid function.

To determine male fertility of these amiR-Mp*FER* lines with reduced spermatogenous

tissue, we performed crosses with wild-type female plants. The spermatocyte

concentration harvested from wild-type antheridia was almost 3-fold higher than from

amiR-Mp*FER* lines and was adjusted using a hemocytometer. In just one of the crosses

(n=20), a single sporophyte was formed, resulting in 0.05 ± 0.05 (mean $\pm$ SE) sporophytes

per archegoniophore, whereas the same females crossed with wild-type males yielded

9.95 ± 2.33 (mean $\pm$ SE) sporophytes per archegoniophore. In summary, Mp*FER* plays

a role in antheridiophore development and spermatocytes of plants with reduced Mp*FER*

activity are poorly fertile. Thus, *Cr*RLK1Ls have a conserved role in reproduction in

addition to their role in cell expansion and integrity during vegetative development.

Some *A. thaliana Cr*RLK1Ls are involved in surveying CWI and in inhibiting growth when

CWI is impaired. Cell wall defects caused either by mutations affecting cellulose

biosynthesis or by treatment with isoxaben, a cellulose synthesis inhibitor, can be

suppressed by a mutation in At*THE1* [8], [29]. To investigate whether Mp*FER* also acts

as CWI sensor, we treated wild-type, Mp*fer-1*, and Mp*fer-2* gemmae with isoxaben (Fig.

5F,G). As in the wild type, isoxaben affected the development of gemmae with reduced

Mp*FER* activity, all growing progressively less with increasing isoxaben concentration

(Fig. 5F,G). Moreover, isoxaben treatment was lethal for Mp*fer*-2 knock out mutants (Fig.

5F,G). Similar observations were reported for At*fer-4*, where isoxaben treatment

increased the number of dead cells in hypocotyls [74]. These results suggest that the

function of At*THE1* in repressing cellular growth when CWI is impaired appeared later in

the course of land plant evolution or was lost in the *Marchantia* lineage.

The *Cr*RLK1L signaling pathway is conserved in land plants

As downregulation of Mp*FER* or At*FER* produce similar phenotypes concerning polarized

growth and cell expansion in *M. polymorpha* and *A. thaliana*, respectively, interspecific

complementation was attempted. First, the coding sequence of Mp*FER* fused to the

Green Fluorescent Protein (GFP) gene under the control of the viral *35S* promoter

(*pro35S*:Mp*FER-GFP*) was transformed into heterozygous At*fer-1* mutants. Then, GFP

expression at the cell periphery and complementation of the bursting root hair and

reduced rosette size phenotypes were assessed in transgenic plants (Fig. 6A). At*fer-1*

homozygotes expressing the MpFER-GFP protein look similar to At*fer-1* mutants,

indicating that Mp*FER* does not complement these At*fer-1* loss-of-function phenotypes

(Fig. 6A). While we cannot exclude that this is because we used a heterologous promoter,

we consider this unlikely as expression the *35S* promoter is routinely used to complement

root hair phenotypes [75], [76]. In a complementary experiment, the coding sequence of

At*FER* fused to the Citrine gene under the control of the Mp*EF1* promoter

(*pro*Mp*EF1:*At*FER-Cit*) was transformed into amiR-Mp*FER*3-2 plants. Ten independent

lines with Citrine expression at the cell periphery were phenotypically characterized (Fig.

6B,C). All had bursting rhizoids and a reduced thallus size, similar to the parental amiR-

Mp*FER*3-2 line, suggesting that At*FER* does not rescue the vegetative phenotypes

produced by down-regulation of Mp*FER*. A sequence alignment of At*FER* with amiR-

Mp*FER3* showed not complementarity (Fig. 3B), confirming that the failure to complement

is not cause by a potential downregulation of At*FER* by amiR-Mp*FER3*. Taken together,

these results show that MpFER and AtFER cannot substitute each other with respect to

their function in cell expansion and tip growth.

In *A. thaliana*, it was shown that AtFER forms a complex with AtRALF peptides and the

LRE/LLG co-receptors [36]. As no interspecific complementation was observed, the

conservation of binding surfaces between *Cr*RLK1L, RALF, and LRE homologs were

analyzed. The *M. polymorpha* genome encodes three RALF peptides (MpRALF:

Mp1g27120   [Mapoly0002s0166];   MpRALF2:   Mp2g21670   [Mapoly0040s0047];

MpRALF3: Mp7g07270 [Mapoly0076s0067]) [47], [48]. We performed a phylogenetic

analysis using the amino acid sequences of the predicted mature peptides of *A. thaliana*

and *M. polymorpha* RALFs (Fig. S4A). All MpRALF peptides clustered together with

AtRALFs known to interact with *Cr*RLK1L receptors, including AtRALF1 and AtRALF23,

which serve as AtFER ligands [23], [34] (Fig. S4A). The MpRALFs shared the four

conserved Cys residues as well as the YXXY and YY motifs with the AtRALF1 subgroup

(Fig. S4B). However, only MpRALF1 and MpRALF3 had an RRXL motif important for S1P

cleavage (Fig. S4B,C)[77], consistent with the presence of one S1P ortholog in basal

plants (Mp8g07990 [Mapoly0155s0018], Fig. S4D). There are two MpLRE/LLG proteins

encoded in the *M. polymorpha,* genome (Fig. S5A,B). Both the MpLRE1 (Mp5g09600

[Mapoly0048s0110]) and MpLRE2 (Mp4g22100 [Mapoly0090s0020]) proteins showed

conservation of the eight Cys and the ND motif distinctive of the family (Fig. S5B);

however, MpLRE1 did not contain a GPI anchoring site (Fig. S5B). Structure prediction using AtLLG1 as a template [36] showed a conserved general structure (Fig. S5C,D). Some of the Mp*RALF* and Mp*LRE* family members show a similar expression pattern as Mp*FER* (Fig. S5F), suggesting that the corresponding proteins could form a complex similar to that described in *A. thaliana*.

To gain insights into the formation of a putative *Cr*RLK1L signaling complex, the MpFER/MpRALF/MpLRE complex was modelled. The general structure appeared conserved (Fig. S5E); however, analysis of conserved sites between *M. polymorpha* and *A. thaliana* showed a low amino acid conservation in the interacting surfaces of the complex subunits (Fig. 6D). This suggests that the lack of interspecific complementation may be due to differences in amino acids that are important for complex formation, and that the proteins forming the complex have diverged, following distinct routes of co-evolution in the two lineages.

Because interspecific complementation of the amiR-Mp*FER* phenotypes was unsuccessful, we asked whether suppression using a downstream component of *Cr*RLK1L signaling identified in *A. thaliana* was possible. AtMRI, a receptor-like cytoplasmic kinase, acts downstream of At*FER* or At*ANX1/2* in the regulation of polar tip growth [78], [79]. The dominant AtMRI[R240C] mutation can suppress pollen tube rupture in the At*anx1*/At*anx2* double mutant and the root hair defects of At*fer-4* [78], [79]. A mutation in Mp*MRI* (aka Mp*PTI*) with a similar bursting rhizoid phenotype as observed in Mp*fer-1*, was identified in the same genetic screen for *M. polymorpha* mutants with defective rhizoids [59]. To test whether Mp*MRI* is a conserved downstream component of the Mp*FER* signal transduction pathway, a dominant-active form of the protein equivalent to AtMRI[R240C] was transformed into the amiR-Mp*FER*3-2 line, driven by 2 kb of the endogenous promoter ($_{pro}$Mp*MRI*:Mp*MRI*[R240C]). Several independent lines showed a

partial restoration of rhizoid growth; however, we also observed defects in thallus

development of lines with higher levels of Mp*MRI* expression, mainly abnormalities in the

epidermis (Figs. 6E and S6A-E). This suggest that Mp*MRI* acts downstream of Mp*FER*

in the signal transduction pathway controlling polarized growth since the origin of land

plants, in agreement with a recent report on the functional characterization of Mp*MRI* [60].

Overexpression of Mp*FER* affects morphological integrity

Because overexpressing At*FER* in *M. polymorpha* did neither lead to any obvious

phenotypes nor complementation of amiR-Mp*FER* lines (Fig. 6B,C), we also

overexpressed At*FER* and Mp*FER* in the wild type. As in the amiR-Mp*FER* background,

overexpression of AtFER ($_{pro}$Mp*EF1:*At*FER-Cit*) in a wild-type background had no effect,

consistent with the idea that At*FER* is unable to form a complex with the corresponding

M. polymorpha proteins. However, when expressing an MpFER-Citrine fusion protein

($_{pro}$Mp*EF1:*Mp*FER-Cit*) in wild-type plants, thallus development was strongly affected

(Fig. 7A). As expected, MpFER-Cit localized to the membrane (Fig. 7A inset), and higher

protein levels correlated with more severe phenotype (Fig. S6F,G). Scanning electron

microscopy of epidermis of plant expressing high levels of Mp*FER* showed defects in the

formation of air chambers and air pores (Fig. 7B), similar to what we observed when

expressing the dominant-active Mp*MRI*^{R240C} (Fig. S6A-C). This phenotype is reminiscent

of the *quasimodo1* mutant in *A. thaliana*, in which cell walls have a reduced pectin content

and cell adhesion is compromised [80]. Moreover, overexpression of Mp*FER* affects the

number of lobes in the antheridial receptacle, producing only 4 instead of 8 (Fig. S6H),

supporting the importance of Mp*FER* during sexual development. Considering the

importance of Mp*FER* for normal rhizoid formation, we also analyzed rhizoids in the

$_{pro}$Mp*EF1:*Mp*FER-Cit* lines: while rhizoid morphology looked normal, a decrease in rhizoid

number was observed (Fig. 7C,D). To analyze cell wall properties of *MpFER*

overexpressor lines, chlorophyll efflux rates were measured, as an indicator for epidermal

permeability (Fig. 7E). Overexpression of *MpFER* clearly reduced the chlorophyll efflux

compared to the wild type, while the opposite was observed in Mp*fer-2* mutants (Fig. 7E).

Taken together, the results obtained by overexpressing Mp*FER* support a role of Mp*FER*

in cell wall formation and the maintenance of tissue integrity during plant morphogenesis.

Discussion

During land plant evolution, many developmental aspects have changed in order to adapt

to new environments, producing the enormous diversity observed in the plant kingdom.

However, the control and maintenance of CWI remained a key aspect for the biology of

a plant cell. CWI sensing is not only important for vegetative growth by cell expansion,

but it is also essential for sexual reproduction and defense responses.

The *Cr*RLK1L family was first characterized for its importance during angiosperm

fertilization, a process comprising aspects of polar cell elongation, CWI control, and cell-

cell communication [7], [31], [32]. Lately, different members of this family were found to

carry out diametrically opposite roles in diverse aspects of plant development, which has

made it difficult to define the core or ancestral function of this gene family comprising 17-

members in *A. thaliana* [7]–[13], [15]–[21], [33].

Here, we report the characterization of the single *Cr*RLK1L gene encoded in the genome

of *M. polymorpha*. Structurally, the MpFER ECD and kinase domain share similarities

with previous characterized *Cr*RLK1L members, including the malectin-like domain and

conserved phosphorylation sites. Conservation of RALF and LRE members in *M.*

polymorpha suggests that MpFER also forms a complex at the plasma membrane [36].

While modeling predicted that the general structure of the complex was conserved, the

interaction surfaces seem to have diverged and co-evolved independently in the respective lineages.

We observed a broad involvement of Mp*FER* at both vegetative and reproductive stages of development: During vegetative growth of the gametophyte, Mp*FER* is required for rhizoid formation and cell expansion but it also plays a role in male gametogenesis and is expressed in female reproductive organs. These findings suggest that the *Cr*RLK1Ls have, also in ancestral land plants, held roles in various aspects of both vegetative and reproductive development. Thus, the role of *Cr*RLK1Ls in angiosperm reproduction does not represent a derived feature from a purely vegetative function in cell elongation in bryophytes but constitutes a conserved feature of this gene family.

Moreover, the almost ubiquitous expression of Mp*FER* in *M. polymorpha* is consistent with the collective expression of different *Cr*RLK1L members in essentially all tissues and organs of *A. thaliana* [44]. Thus, gene duplication allowed the expansion of the *Cr*RLK1L family in angiosperms, diversifying their expression patterns and functions but also leading to genetic redundancy [9], [18], [33], [44].

*Cr*RLK1L members are important sensors of CWI during polarized growth, both in pollen tubes [9], [33] and root hairs [17]. In early divergent land plants, rhizoids are tip-growing cell with a function analogous to that of root hairs, important for taking up nutrients and water [59], [81]. Based on the conservation of many genes controlling the development and growth of tip-growing cells forming rooting systems, it was suggested that these pathways were active in the earliest land plants that existed about 470 million years ago [59]. That Mp*FER* also controls rhizoid integrity points to the importance of the *Cr*RLK1L pathway for polarized growth since the origin of land plants. Moreover, the fact that amorphic Mp*fer* mutants contain many dead cells demonstrates the importance of the *Cr*RLK1L pathway for cellular integrity during vegetative growth. As the occurrence of

dead cells was recently also reported in At*fer-4* mutant seedlings [74], the function of this pathway in maintaining cellular integrity may also be ancestral. Whether Mp*FER* also plays a central role in plant innate immunity, as it does in angiosperms [16], [23], remains to be determined.

The signaling mechanisms downstream of Mp*FER* in *M. polymorpha* development are still unclear. Mp*MRI*, the *M. polymorpha* ortholog of At*MRI*, which acts downstream of AtFER and AtANX1/2 in *A. thaliana* during polarized growth [78], does have a similar rhizoid phenotype as Mp*FER* [59], [60]. The suppression of the rhizoid phenotype in amiR-Mp*FER* transformants with the constitutively active form MpMRI^{R240C} corroborates a conserved *Cr*RLK1L signaling cascade during polarized growth in *M. polymorpha* but whether this pathway also relies on Ca^{2+} and reactive oxygen species (ROS) as second messengers as it does in *A. thaliana* remains to be determined.

In *A. thaliana*, loss-of-function mutations in the motor protein AtKINESIN-13a, a microtubule-based motor, cause cell elongation in petals, leaves, and hypocotyls concomitant with changes in cell wall composition in leaves [82]. This effect depends on functional AtTHE1 and does not occur in At*kinesin-13-a*/At*the1* double mutants. Interestingly, in this case, AtTHE1 stimulates cell elongation in response to defects in cell wall deposition [82]. Based on our results, we have no indication that Mp*FER* acts as a CWI sensor and regulates growth similar to At*THE1* in response to cell wall disturbances. We have not observed a context in *M. polymorpha* development in which Mp*FER* inhibits cell expansion. Therefore, the ability to restrict growth in response to disturbances in CWI could represent an evolutionary derived feature, as gene duplication allowed for the diversification and specialization of *Cr*RLK1Ls to regulate cellular growth in a more complex and context-dependent way. The capacity to flexibly regulate growth appears to be fundamentally context-dependent, as a similar discrepancy in promoting or inhibiting

growth is also known for some well-studied growth promoting agents, like the phytohormones auxin and brassinosteroid, which can have growth-inhibitory effects dependent on tissue and concentration. The reverse was noted for the phytohormones ethylene and abscisic acid, which are usually considered growth inhibitors but can also promote growth, dependent on the context [83]–[85].

Taken together, our results suggest an ancestral and conserved function of the *Cr*RLK1Ls in polar cell growth and cell expansion, but also during sexual reproduction. In addition, we probe the relevance of *Cr*RLK1L members for cell integrity and morphogenesis. In angiosperms, *Cr*RLK1Ls occupy an important position at the interface of developmental and environmental inputs, which are integrated by a downstream signaling machinery that controls cell shape and polar growth to ensure normal development, e.g., by preventing cell rupture upon CWI defects. Whether the same downstream machinery is utilized in a similar fashion in *M. polymorpha* remains to be shown, even though the similarity in function points to a related mechanism. Even less is known about upstream aspects of *Cr*RLK1L signaling. Future studies on the transcriptional regulation and the upstream components, such as the putative RALF ligand(s), in the highly simplified *M. polymorpha* system could provide fundamental insights into the molecular mechanism of the *Cr*RLK1L pathway, which is so central to land plant physiology, growth, and development.

Materials and methods

Experimental model

M. polymorpha subsp. *ruderalis* (Bischl. & Boissel.-Dub.) plants of the Tak-1 accession were grown on sterile culture on half-strength Gamborg's B5 basal medium (PhytoTechnology Laboratories), supplemented with 1% phytoagar in Petri dishes sealed

with air-permeable tape. The plants were kept under fluorescent light and long-day

conditions (16 h light at 22°C, 8 h dark at 20 °C) in Percival growth cabinets (models AR-

41L3 and AR-41/L2). Alternatively, to induce sexual reproduction, plants were cultivated

in jiffy pots filled with a 1:1 mixture of soil ("Einheitserde D73 + Bims", Universalerde) and

sand or in culture boxes in sterile culture on half-strength Gamborg's B5 basal medium

(Duchefa Eco2 Box, #E1654), under fluorescent light supplemented with far-red light

(GreenPower LED module HF far-red, #929000464503 and #929000632103 Philips).

Wild-type plants in all experiments were *Arabidopsis thaliana* L. (Heyn) accession L*er*-0.

The At*fer-2* mutant was obtained from the Signal Collection at the Salk Institute. Plants

were grown in soil (ED73; Universalerde), covered with a thin quartz sand layer, under

long photoperiods (16 hs light/8 hs dark) at 23°C and 60% relative humidity.

Phylogenetic analysis

Protein sequences were identified in plant genomes via BLASTp searches in

https://phytozome.jgi.doe.gov/pz/portal.html. To elucidate the evolutionary relationship

across land plant evolutionary history, we focused on *A. thaliana, Selaginella*

moellendorffii, Physcomitrium patens and *M. polymorpha*, as representative species for

angiosperms, lycophytes, mosses and liverworts, respectively. Complete or partial coding

protein sequences were aligned using the ClustalW parameters and were conducted in

MEGA7. Phylogenetic trees were constructed with Neighbor-joining method, with a

bootstrap test of 1000 replicates. The evolutionary distances were computed using the

Poisson correction method.

Vector construction

Expression constructs: a BJ36 plasmid [86] containing a tripleVENUS-NLS (trpVNS)

fragment was modified by adding a ligation-independent cloning (LiC) adapter site [87].

A promoter fragment of Mp*FER* of 3.2 kb was amplified using primers specified below

and cloned into the BJ36 vector via LiC cloning. *pro*Mp*FER* promoter fragment fused to

trpVNS was then shuttled via *Not*I restriction sites into the HART01 [86] expression

vector. The expression vector was also directly modified to contain the LiC sites, so

promoter fragments can now be directly cloned in front of the trpVNS in the HART01

vector via LiC-cloning. This vector was called VHL (trpVNS-containing HART01 vector

with LiC sites).

amiRNA constructs: For functional analyses three independent artificial microRNA

(amiRNA) constructs were made (Fig. S2). The endogenous microRNA precursor

Mp*miR160* was used as both backbone and template to design the constructs which

generate single species small RNA molecules of 21 nt length, complementary to the

target gene transcript [67], [88]. Three miR160 backbones containing independent amiR-

Mp*FER*/amiR-Mp*FER** duplexes were synthesized by GenScript and fused to the

*pro*Mp*EF1* promoter in the BJ36 shuttle vector and shuttled to the HART01 expression

vector as described previously [88]. The folding structure and design procedures were

explained in detail in [67], [88]. The sequences for the amiRNAs were designed to retain

the exact physical properties of the endogenous Mp*miR160* template. The folding

patterns were analyzed in the Mfold software [89] and the sequences synthesized (Fig.

S2).

CRISPR/Cas gRNA: Selection of target sites was done using CasFinder [90].

Complementary oligos were designed and annealed for ligation into gRNA, in the

pMpGE_En03 vector, previously digested with BsaI restriction enzyme. Resulting gRNAs

were incorporated into binary vector pMpGE011 through gateway recombination.

Over-expression of Mp*FER* and At*FER*: The full-length Mp*FER* sequence was amplified

from genomic *M. polymorpha* DNA with attB sites and Gateway-cloned via pDONR207.

For *A. thaliana* expression, Mp*FER* was introduced into the expression vectors

pMDC201, which contains a *2X35S* promoter fragment and an in-frame C-terminal

mGFP6 [91]. For *M. polymorpha* expression, Mp*FER* was introduced intro the binary

vector pMpGWB308, which contains a MpEF1 promoter and an in-frame C-terminal

citrine [92]. Similarly, for expression of At*FER* in *M. polymorpha*, coding sequence of

AtFER was introduced into pMpGWB308 binary vector, for the generation of

$_{pro}$Mp*EF1:*At*FER-Cit* construct.

Quantitative real-time PCR (qRT-PCR) and droplet digital PCR (ddPCR)

RNA extraction from *M. polymorpha*: RNA extraction from thallus tissue (ca. 100mg

including apical notch) was done using the Rneasy plant mini kit (#74904, Qiagen). To

remove contaminating DNA, the TURBO DNAfree™ Kit (AM1907 Ambion) was used

according to manufacturer's recommendations. To extract RNA from the gametophores

(2 antheridiophores > 5mm diameter were pooled, respectively, for each replicate), the

Direct(-zol) RNA MiniPrep (#R2050, ZymoResearch) was used with TRIzol Reagent

(#15596026, Ambion) according to the manufacturer's protocol, including on-column

DNase digestion and subsequent TURBO DNA-free™ Kit (AM1907 Ambion) treatment.

RNA samples were quantified using the Qubit® RNA HS Assay Kit (#Q32852, Life

Technologies), a Nanodrop ND-1000 Spectrometer or an Agilent 2100 BioAnalyzer.

cDNA synthesis: cDNA was synthesized from 0.5 μg or 1.0 μg of total RNA. Reverse

transcription was performed in 25 μl with 20μg/ml Oligo(dT) 12-18 primers and 200 units

of SuperScript® II Reverse Transcriptase (#18064-014, Invitrogen) according to the

manufacturer's protocol. The resulting cDNA was diluted 1:9 by adding nuclease-free

water. To control for genomic DNA contamination, control replicates of all samples were

incubated without SuperScript® II reverse transcriptase.

Primer tests and quantitative RT-PCR: Primer efficiency and concentration tests were

carried out for Mp*FER* and suitable reference genes as described previously (Rövekamp

et al., 2016, Althoff et al., 2014). Amplification experiments were carried out using a 7500

Applied Biosystem Fast Real-Time PCR System. Reactions were performed in 20 μl

volumes containing 10 μl 2X SYBR-green mastermix (SsoAdvanced™ Universal

SYBRGreen Supermix), 200 nM (Mp*EF1* and Mp*FER*) or 250 nM (Mp*ACTIN*) forward and

reverse primers, and 1 μl diluted cDNA. Where possible, three technical and biological

replicates were performed for each reaction. The primers used for the qRT-PCR are

summarized in Table S1.

For ddPCR analysis, individual PCR reactions were performed in a total volume of 25 μl,

using 1X ddPCR EvaGreen Supermix, with droplets generated according to

manufacturer's recommendations. Reading of the PCR-amplified droplets was carried out

by the QX200 Droplet Reader (Bio-Rad) and analysed by the QuantaSoft TM Software

(v1.4, Bio-Rad). The counts from ddPCR were normalised through a log2 transformation.

Afterwards, relative expression was calculated against the geometric mean of the counts

for all three reference genes (Mp*ACT1*, Mp*ACT7*, and Mp*APT3*) [93]: (log2(gene tested

+ 1) - log2(geom. mean of all references + 1)). The primers used for the ddPCR are

summarized in Table S1.

3' RACE-PCR

RNA from wild-type and Mp*fer-1* mutant plants were extracted as previously described.

A poly(G)-tail was added at the 3' end of the RNA according to a Poly(A) Polymerase

protocol (Thermo Scientific), using GTP instead of ATP. Reverse transcription was

performed using anchored Oligo(dC) instead of Oligo(dT). Nested PCR on the cDNA was performed using two forward primers that matched the 3' UTR sequence of Mp*FER* (3'RACE-Fwd1 and 3'RACE-Fwd2) and Oligo(dC). PCR fragments were purified and sequenced.

Transformation of *M. polymorpha* and *A. thaliana*

Agrobacterium-mediated transformation using regenerating thalli of *M. polymorpha* was done by co-cultivation 15-day old plant fragments with transformed *Agrobacterium tumefaciens* (GV3101) cells [94], [95]. After three days of incubation, positive transformants were selected on Gamborg's B5 plates supplemented with 10 *µ*g/ml Hygromicin B (Invitrogen), 0.5 µM Chlorosulfuron or Kanamycin, and 100 µg/ml of Cefotaxime sodium. Isogenic lines were obtained by using plants derived from gemmae of the T1 generation (G1 generation). G1 or subsequent gemmae generations were used for all experiments, as they are derived from single cells [96], [97]. Transformation of *A. thaliana* via *A. tumefaciens* (GV3101) was performed by floral dipping according to [98]. Primers used for amplification of promoter fragments of Mp*FER* are listed in Table S2.

Microscopy

Plants were observed in a Lumar.V12 (Zeiss) or a Leica MZFLIII dissection microscope and photographed with an AxioCam HRc (Zeiss) or Leica DFC 420C camera. Clearings were observed using a Leica DMR microscope and photographed with an Axiocam 105 color camera. Fluorescence reporter expression were analyzed using a Leica SP5 confocal microscope. Fiji [99] and GIMP (version 2.8.10) software were used for adjustments of brightness, contrast, channel selection, and image size.

Tissue clearing for bright field microscopy: *M. polymorpha* tissues were fixed in Carnoy's

solution and incubated at 4°C overnight, followed by rehydration in an ethanol (EtOH)

series of 85%, 70%, 50%, and 30%. Samples were incubated for 1 hour at 4°C for each

step and then incubated in chloral hydrate solution at 4°C overnight. Samples were

mounted in chloral hydrate.

Calcofluor-white staining for cell size measurements: Pieces of fresh thallus tissue

located in fully differentiated zones [100] of plants grown on plates were dissected out

and put into water. The pieces were transferred to PBS pH 6.1 containing 100 μg/ml

Calcofluor-white and vacuum-infiltrated for 1 hour before being mounted on slides. The

area of the epidermal cells was measured using Imaris 8.3.1. software. Cells of air pores

and the two adjacent cell layers were excluded from the analysis, as well as the spiralling

cells surrounding newly developing air pores.

For cell-death staining, seven-day old gemmalings were stained with lactophenol-

trypanblue (10 mL of lactic acid, 10 mL of glycerol, 10 g of phenol, 10 mg of trypan blue,

dissolved in 10 mL of distilled water) and boiled for approximately 1 min in the stain

solution and then decolorized in chloral hydrate (2.5 g of chloral hydrate dissolved in 1

mL of distilled water) for at least 30 min. They were mounted in chloral hydrate and

imaged in dissecting microscope.

For scanning electron microscopy (SEM), 20-day old plants of WT and $_{pro}$Mp*EF1*:Mp*FER-*

Cit lines *#5* and #9 were harvested and fixed in using Glutaraldehyde (0.25% in 20mM

HEPES) over night at 4 °C. After fixation, dehydratation with Acetone series (from 50%

to 100% acetone "ultradry") was done. The samples were then critical point dried, without

the following sputter coated step. Microscopic imaging was done on a JSM-6010 (Jeol,

Freising, Germany) at the Center for Microscopy and Image Analysis at the University of

Zurich.

To evaluate the stiffness of the gemmaling tissue, we used cellular force microscopy (CFM; [101], [102]). A microelectromechanical systems (MEMS)-based FT-S100 force sensor (FemtoTools AG, Buchs, Switzerland) with a glued-on tungsten probe (T-4-22; Picoprobe by GGB Industries INC, Naples, FL, USA) was used to indent the gemmaling tissue. The sensor has a force range of ±100 µN and a resolution of 15 nN measured at 500 Hz. The sensor was attached with a custom-made acrylic arm to a xyz positioner (SLC-2460-M; SmarAct), which was fixed on a xyz piezo stage (P-563.3CD PIMars; Physik Instrumente (PI) GmbH & Co., Karlsruhe, Germany). The xyz positioner was used for the rough positioning of the sensor on a part of a gemma that was as much as possible in contact with the supporting glass slide and away from the meristematic zones. Nine evenly spaced measurements were then taken in a 3-by-3 grid, again using the xyz positioner to place the sensor, while the indentation at each position was driven by the piezo stage. This procedure was repeated a second time on each gemmaling, whenever possible on the opposite lobe.

The stiffness analysis was achieved with Matlab (Mathworks, Natick, MA, USA) based on the slope fitted to the force-displacement curves, which were acquired by CFM. In a first round of data curation, each force-indentation curve was visually inspected, and curves that didn't have a clear contact point or showed a clear displacement of the gemmaling, indicated by an irregular slope, were discarded. The remaining data exhibited a bimodal distribution with a high density at very soft values around 1 µN, indicating the presence of an artefact due to gemmalings that were not in contact with the slide and, therefore, were displaced rather than indented. To identify such failed measurements, we pooled all data and fitted a Gaussian mixture model with two components. We then removed all measurements that were closer than 3 standard deviations to the mean of the distribution

with the smallest mean (mean = 1.16 µN and sd = 0.56 µN). Hence, all measurements

with a value below 2.9 µN were removed.

Boxplots and statistical analysis were performed in Matlab. The statistical tests used to

assess the significance of differences between groups are indicated in the figure legends.

Western blot analysis

M. polymorpha explants were ground in liquid nitrogen and resuspended in 300 µL of

Laemmli buffer [103]. After centrifugation at 6,000 rpm for 5 min at 4˚C, the supernatant

was recovered. Protein accumulation was confirmed by Western blotting, using the anti-

GFP antibody (1:3000, Torrey Pines Biolabs). As secondary antibodies, the Agrisera

S09602 goat anti-rabbit antibody was used with the ECL-chemistry of FUSION FX -

Western Blot & Chemi Imaging (Vilber Lourmat).

Growth rate experiments

Gemmae of wild-type, Mp*fer-1*, and independent transformation lines of amiR-Mp*FER*

lines were grown on plates and the thallus area was measured over time. 10 or 30

gemmae of four independent transformation lines were grown on half-strength Gamborg's

B5 and scanned on an Epson 2450 Photo scanner at 600dpi. Plant area was measured

using color threshold settings and particle analysis in the Fiji software [99].

Expression map

Gene expression of *Cr*RLK1L signaling pathway components was calculated from

publicly available RNA-seq data of 39 samples of *M. polymorpha* Tak-1 and Tak.2.

Samples were grouped by tissue and analyzed under the same pipeline: Nextera paired-

end adapters were trimmed from sequencing reads using the bbduk tool embedded in

the BBMap software package [104]. Read ends with quality below 20 were also trimmed

and the minimum read length was set to 25. The rest of parameters of bbduk were set as

default. Reads were mapped to *M. polymorpha* reference genome (v3.1) [48] using the

Tophat software (v2.1.1) [105], designed for RNA-sequencing read mapping. Tophat

contains Bowtie (v2.3.2.0) [106] as the aligner software. All parameters were set to default

values except for the RAM memory and number of used Cores. Mapped reads were

sorted using samtools (v1.3.1) [107] and counted with the software HTSeq-count"

(v.0.9.1) [108]. Gene expression values were calculated using the package DESeq2

(Release 3.6) [109] for R software (v3.4.4) [110]. Another R package, ggplot2 (release

2.2.1) [111] was used for producing the figures. A summary of the samples that were

used in this study is provided in Table S3.

Chlorophyll efflux rate

To measure the chlorophyll efflux rate, 10-day-old gemmalings were used [112]. Sixteen

replicates of MpFER-overexpressor, Mp*fer*-2 mutant, and wild-type lines were immersed

in an 80% EtOH solution in 5-mL tubes. The tubes were agitated gently on a shaker

platform. Aliquots of 2 µL were removed every min, during the first 10 minutes. The

amount of chlorophyll extracted by the EtOH solution was quantified with a

spectrophotometer (Nanodrop 8000, Thermo Scientific) and calculated from UV light

absorption at 647 and 664 nm. The micromolar concentration of total chlorophyll per gram

of fresh weight of tissue per ml of EtOH was calculated using the equation Total

micromoles chlorophyll = $7.93(A_{664}) + 19.53(A_{647})$.

Three-dimensional (3D) protein structure modeling

Comparative modeling of protein 3D structures was performed using the SWISS-MODEL

online-tool [64]. The software was fed with the three protein sequences of MpFER,

MpRALF1, and MpLRE1 as input to model, and one PDB file as a 3D template, which

contains the published crystal structure of the protein complex formed by AtFER,

AtRALF23, and AtLLG1 [36]. Quality control of the model was assessed using the same

tool and the results are summarized in Table S4. Spatial comparisons of the *M.*

*polymorph*a predicted structures with the crystal structures of *A. thaliana* were performed

in PyMol (Table S4) [113].

728 QUANTIFICATION AND STATISTICAL ANALYSIS

729 Analyses were performed in InfoStat (http://www.infostat.com.ar) or Matlab. Tests are

730 indicated in the corresponding figure legend.

732 Acknowledgments

We thank Liam Dolan for kindly providing the Mp*fer-1*/Mp*the* mutant, Valeria Gagliardini

for help with qRT-PCR and ddPCR, Anja Grossniklaus for help in measuring epidermal

cell area, Célia Baroux and Ethel Mendocillo-Sato for instructions on confocal microscopy

and the use of Imaris Software, Tiago Meier for helping with sample preparation for SEM

microscopy, Isabel Monte and Cyril Zipfel for comments on the manuscript, and Christof

Eichenberger, Frédérique Pasquer, Arturo Bolaños, Daniela Guthörl, Daniel Prata, and

Peter Kopf for general lab support.

741 Author contributions

MAM: Conceptualization, Formal analysis, Investigation, Methodology, Validation,

Visualization, Writing – original draft, Writing – review & editing

MR: Conceptualization, Formal analysis, Investigation, Methodology, Validation,

Visualization, Funding acquisition, Writing – original draft, Writing – review & editing

AG-F: Data curation, Formal analysis, Software, Validation, Visualization, Writing –

original draft, Writing – review & editing

DL: Investigation, Validation

DM: Formal analysis, Investigation, Validation, Writing – review & editing

PG: Formal analysis, Investigation, Validation, Writing – review & editing

HV: Formal analysis, Investigation, Validation, Writing – review & editing

JLB: Funding acquisition, Resources, Supervision, Validation, Writing – review & editing

UG: Conceptualization, Formal analysis, Funding acquisition, Project administration,

Supervision, Validation, Writing – original draft, Writing – review & editing

Funding disclosure

This work was supported by the University of Zurich, Monash University, and grants from

the Forschungskredit of the University of Zurich (FK-16-090,

http://www.research.uzh.ch/en/funding/phd/fkcandoc) to MR, the Australian Research

Council (http://www.arc.gov.au; DP130100177) to JLB, and the Swiss National Science

Foundation (http://www.snf.ch; 310030B_160336 and 31003A_179553) to UG. The

funders had no role in study design, data collection and analysis, decision to publish, or

preparation of the manuscript.

Competing interests

766 The authors have declared that no competing interests exist.

Main figure titles and legends

Figure 1. *Cr*RLK1Ls Are Conserved among Land Plants

(A) Representation of AtFER and MpFER proteins. The malectin domains are represented in blue and kinase domains in green. SP, signal peptide; TM, transmembrane domain (yellow). Amino acids important for regulation and activity are marked in red, violet, and light blue as described in Fig. 1E.

(B) A rooted neighbor-joining tree of the amino acid sequence of the predicted malectin-like domain (corresponding to aa 76-431 of MpFER). *Cr*RLK1L members from *Marchantia polymorpha* (Mp, green), *Physcomitrium patens* (Pp), *Sphagnum fallax* (Sphfalx), *Ceratodon purpureus* (Cpur)*, Anthoceros agrestis* Bonn (AagrBONN)*, Anthoceros agrestis* Oxford (AagrOXR)*, Anthoceros punctatus* (Apun)*, Selaginella moellendorffii* (Smo), *Amborella trichopoda* (AmTr)*,* and *Arabidopsis thaliana* (At, blue) were used. The best hit from *Chara braunii* (CHBRA) was also included. The numbers indicate the bootstrap values (%) from 1000 replications. The given scale represents a substitution frequency of 0.1 amino acids per site.

(C) Cartoon representation of the predicted three-dimensional structure of the MpFER malectin-like domain, showing predicted alpha-helix and beta-sheet structures.

(D) Structural superposition of the malectin-like domains of AtFER (blue) and MpFER (green).

(E) Alignment of the cytoplasmic domains of MpFER, AtERU, AtFER, and *Cr*RLK. Kinase domains are in green, putative phosphorylation sites in light blue (Ser) and violet (Thr), and the conserved catalytic Lys in red.

See also Fig. S1

Figure 2. Mp*FER* Is Broadly Expressed in most Tissues throughout the *M.*

***polymorpha* Life Cycle**

Expression of $_{pro}$Mp*FER*:*trpVNS* in different organs of male and female *M. polymorpha*

plants. Fluorescence is visualized as either green (epifluorescence microscope) or yellow

(confocal microscope) color, depending on the panel.

(A to D) Bright field (left) and epifluorescence (right) images of a thallus, (A) rhizoid (C),

gemmae cups (C), and an archegoniophore (D). Red arrows indicate meristematic zones,

the yellow arrow gemmae. Scale bars, 0.5 mm.

(E and F) Confocal images of archegonia before (E) and 2 days after fertilization (F).

Scale bar, 50 μm.

(G) Bright field (left) and epifluorescence (right) images of an antheridiophore. Scale bar,

1 mm.

(H) Confocal image of antheridia. Scale bar, 100 μm.

(I) Confocal image of sporophyte and spores. Scale bar, 25μm.

Figure 3. Mp*FER* Controls Cell Size during Gametophyte Development

(A) Representation of the mature Mp*FER* mRNA. The coding sequence is in green,

locations of the amiRNA target sites in blue. The location of the T-DNA insertion in Mp*fer-*

1 is shown.

(B) Base complementary of mature amiR-Mp*FER*1, amiR-Mp*FER*2, and amiR-Mp*FER*3

with the Mp*FER* transcript. Very low complementary was found between amiR-Mp*FER*3

and At*FER*.

(C) Representative pictures of 10-day old gemmalings of wild-type (WT), Mp*fer-1*, and

amiR-Mp*FER*3-2 lines. Scale bar, 1 mm.

(D) Violin plot of thallus area of 14-day old gemmalings of WT, Mp*fer-1*, and amiR-

Mp*FER*3-2 lines. n = 30. Different letters indicate significant differences according to the

non-parametric Kruskal-Wallis test and Dunn's test for multiple comparisons (P < 0.001).

Orange circles indicate the group mean and the corresponding vertical bars the standard

deviation for each group.

(E) Relative expression level of Mp*FER* in thallus tissue from WT and two independent

insertion lines of amiR-Mp*FER*2 and amiR-Mp*FER*3, as measured by qRT-PCR. Mp*EF1*

was used as internal control. Shown are means ± standard errors of the means (SEM) of

three biological replicates. Statistical analysis was performed by one-way analysis of

variance (ANOVA) followed by a post-hoc Duncan test (*P < 0.01).

(F) Violin plot of cell size in WT, *Mpfer-1*, and amiR-Mp*FER*3-2 lines. Only areas with fully

expanded cells in approximately 3-week old plants were used for measurements.

Different letters indicate significant differences according to the non-parametric Kruskal-

Wallis test and Dunn's test for multiple comparisons (P < 0.001). Orange circles indicate

the group mean and the corresponding vertical bars the standard deviation for each

group.

See also Figs. S2,S3

Figure 4. The Integrity of Thalli from Mp*fer* Knock-out Lines Is Severely Affected

(A) Representation of the mature Mp*FER* mRNA with the coding sequence in green, and

the location of the gRNA target sites in red.

(B) Sequences of gRNA target sites in the confirmed Mp*fer* knock-out mutants. Deletions

and insertions are highlighted in red.

(C) Representative pictures of 14-day old gemmalings of the wild type (WT), Mp*fer-1*, and two independent Mp*fer* knock-out mutants. Scale bar, 0.5 cm.

(D) Violin plot of thallus area of 7-day old gemmalings of WT, Mp*fer-1*, and two independent Mp*fer* knock-out mutants. n = 25. Different letters indicate significant differences in a one-way analysis of variance (ANOVA) follow by a post-hoc Duncan test ($P < 0.01$). Orange circles indicate the group mean and the corresponding vertical bars the standard deviation for each group.

(E) Representative pictures of 7-day old gemmalings growing in upside-down plates of WT, Mp*fer-1*, and two independent Mp*fer* knock-out mutants. Scale bar, 500 μm.

(F) Violin plot of rhizoid number in 3-day old gemmalings of WT, Mp*fer-1*, and two independent Mp*fer* knock-out mutants. n = 16. Statistical analysis was performed by one-way analysis of variance (ANOVA) followed by a post-hoc Duncan test (*$P < 0.01$).

(G) Trypan blue staining of 7-day old gemmalings of WT, Mp*fer-1*, and two independent Mp*fer* knock-out mutants. Scale bar, 1 mm.

(H) Violin plot of the reduction in the apparent stiffness of gemmaling tissue in Mp*fer-2* compared to the WT. Statistical analysis was performed using a Kruskal-Wallis test followed by a Tukey-Kramer post-hoc test (*$P < 0.001$). Orange circles indicate the group mean and the corresponding vertical bars the standard deviation for each group.

Figure 5. Conservation of *Cr*RLK1L Function in Land Plants

(A) Violin plot of the number of antheridiophores produced per plant for wild-type (WT, n=9) and two independent amiR-Mp*FER*3 lines (amiR-Mp*FER*3-2, n = 9; amiR-Mp*FER*3-3, n = 12). Different letters indicate a significant difference in a one-way analysis of variance (ANOVA) followed by a post-hoc Duncan test ($P < 0.01$).

(B) Antheridiophore splash platform size distribution of WT (n = 32), amiR-Mp*FER*2-3 (n = 34), and amiR-Mp*FER*3-3 (n = 11) lines. For all antheridia with a stalk >8 mm, length and width of the platform were recorded.

(C) qRT-PCR of Mp*FER* levels in antheridiophores in a WT and amiR-Mp*FER*3-3 line. Mp*EF1* was used as internal control. Shown are means ± SEM of three biological replicates. Statistical analysis was performed by one-way analysis of variance (ANOVA) followed by a post-hoc Duncan test (*P < 0.01).

(D and E) Mature antheridia of a WT (D) and amiR-Mp*FER*3-3 line (E). Bright field images (left) and in cross-section (right). The spermatogenous areas are indicated in red. Scale bar, 100*µ*m.

(F) Representative pictures of 6-day old gemmalings of WT and Mp*fer-2* mutants at different isoxaben concentrations. Scale bar, 1 mm.

(G) Thallus area of gemmalings of WT, Mp*fer-1*, and Mp*fer-2* lines growing on media containing different isoxaben concentrations. n = 35. Shown are means ± SEM of two biological replicates. Different letters indicate a significantly difference in a one-way analysis of variance (ANOVA) followed by a post-hoc Duncan test (P < 0.05).

See also Fig. S3

Figure 6. Interspecific Complementation of *A. thaliana* and *M. polymorpha* Plants with Reduced Levels of *FER* Activity

(A) Roots of independent *A. thaliana* lines expressing 35S:Mp*FER-GFP* in L*er* wild-type (WT) and At*fer-1* plants. Scale bar, 200 µm.

(B) 10-day old gemmalings of WT and amiR-Mp*FER*3-2 *M. polymorpha* plants with and without expression of At*FER-Cit.* Scale bar, 0.5 mm.

(C) Citrine expression in 7-day old gemmalings of WT and amiR-Mp*FER*3-2 plants with

and without expression of At*FER-Cit*. Scale bar, 100 μm.

(D) Cartoon representation of the *Cr*RLK1L[ECD] polar interactions sites with LRE/RALF

when forming a protein complex. Upper panels correspond to contact areas between

AtFER (purple), AtLLG2 (blue-grey), and AtRALF23 (yellow), and lower panels to the

modeled MpFER structure (green) superimposed onto the same complex. Polar

interactions are depicted as doted yellow lines. Left panels show the predicted polar

interactions of the complex that are conserved between *A. thaliana* and *M. polymorpha*,

whereas the right pannels show the polar interaction between AtFER and AtLLG2 that is

predicted to be absent between MpFER and AtLLG2.

(E) Suppression of the amiR-Mp*FER* phenotype by a dominant-active version of Mp*MRI*

under its own promoter (*pro*Mp*MRI:MRI*[R240C]). Scale bar, 1 mm.

See also Figs. S4-S6

Figure 7: Overexpression of Mp*FER* Affects Morphological Integrity

(A) Representative pictures of 10-day old gemmalings of wild-type (WT) and two different

lines overexpressing Mp*FER* (*pro*Mp*EF1:*Mp*FER-Cit)*. Scale bar, 1 mm. Inset: Citrine

expression at the plasma membrane in *pro*Mp*EF1:*Mp*FER-Cit* lines *#5* and *#9* as observed

under confocal microscopy. Scale bar, 20 μm.

(B) Scanning electron microscopical images of thalli from 20-day old plants of WT and

*pro*Mp*EF1:*Mp*FER-Cit* lines *#5* and *#9*. The blue region demarks an air chamber, while air

pore cells are colored in red. Scale bar, 500 μm.

(C) Representative pictures of 3-day old gemmalings growing in upside-down plates of

WT and *pro*Mp*EF1:*Mp*FER-Cit #5* lines. Scale bar, 1 mm.

(D) Violin plot of the number of rhizoids in 3-day old gemmalings of WT and

$_{pro}$Mp*EF1:*Mp*FER-Cit* lines *#5* and #9. n = 20. Statistical analysis was performed by one-

way analysis of variance (ANOVA) followed by a post-hoc Duncan test (*P < 0.01).

(E) Graphs showing amount of effluxed chlorophyll as a function of time from 15-day old

gemmalings of WT, Mp*fer-2* and $_{pro}$Mp*EF1:*Mp*FER-Cit #5* lines.

See also Fig. S6

918 **References**

[1]   Zhu S, Fu Q, Xu F, Zheng H, Yu F., 'New paradigms in cell adaptation: decades of discoveries on
the CrRLK1L receptor kinase signalling network.', *New Phytol*, vol. 232, no. 3, pp. 1168–1183,
2021.
[2]   Showalter AM., 'Structure and function of plant cell wall proteins.', *Plant Cell*, vol. 5, pp. 9–23,
1993.
[3]   Franck CM, Westermann J, Boisson-Dernier A., 'Plant Malectin-Like Receptor Kinases: From Cell
Wall Integrity to Immunity and Beyond', *Annu Rev Plant Biol*, vol. 69, pp. 301–328, 2018.
[4]   Shiu SH, Bleecker AB., 'Plant receptor-like kinase gene family: diversity, function, and signaling',
Sci STKE, vol. 2001, no. 113, p. re22, 2001.
[5]   Steinwand BJ, Kieber JJ., 'The role of receptor-like kinases in regulating cell wall function', *Plant
Physiol.*, vol. 153, pp. 479–84, 2010.
[6]   Wolf S, Hématy K, Hofte H., 'Growth control and cell wall signaling in plants', *Annu Rev Plant Biol*,
vol. 63, pp. 381–407, 2012.
[7]   Escobar-Restrepo J-M, Huck N, Kessler S, Gagliardini V, Gheyselinck J, et al., 'The FERONIA
receptor-like kinase mediates male-female interactions during pollen tube reception', *Science*,
vol. 317, pp. 656–660, 2007.
[8]   Hématy K, Sado PE, Van Tuinen A, Rochange S, Desnos T, et al., 'A receptor-like kinase mediates
the response of Arabidopsis cells to the inhibition of cellulose synthesis', *Curr Biol.*, vol. 17, pp.
922–931, 2007.
[9]   Boisson-Dernier A, Roy S, Kritsas K, Grobei MA, Jaciubek M, Schroeder, J.I., Grossniklaus, U.,
'Disruption of the pollen-expressed FERONIA homologs ANXUR1 and ANXUR2 triggers pollen tube
discharge', *Development*, vol. 136, pp. 3279–3288, 2009.
[10] Guo H, Li L, Ye H, Yu X, Algreen A, Yin Y., 'Three related receptor-like kinases are required for
optimal cell elongation in Arabidopsis thaliana', *PNAS*, vol. 106, pp. 7648–7653, 2009.
[11] Guo H, Ye H, Li L, Yin Y., 'A family of receptor-like kinases are regulated by BES1 and involved in
plant growth in Arabidopsis thaliana', *Plant Signal Behav*, vol. 4, no. 8, pp. 784–786, 2009.
[12] Kessler SA, Shimosato-Asano H, Keinath NF, Wuest SE, Ingram G, et al., 'Conserved molecular
components for pollen tube reception and fungal invasion', *Science*, vol. 330, pp. 968–971, 2010.
[13] Bai L, Ma X, Zhang G, Song S, Zhou Y, et al., 'A receptor-like kinase mediates ammonium
homeostasis and is important for the polar growth of root hairs in Arabidopsis', *Plant Cell*, vol. 26,
pp. 1497–511, 2014.
[14] Schoenaers S, Balcerowicz D, Breen G, Hill K, Zdanio M, Mouille G, Holman TJ, Oh J, Wilson MH,
et al., 'The Auxin-Regulated CrRLK1L Kinase ERULUS Controls Cell Wall Composition during Root
Hair Tip Growth.', *Curr. Biol.*, vol. 28, pp. 722–732, 2018.

[15] Miyazaki S, Murata T, Sakurai-Ozato N, Kubo M, Demura T, et al., 'ANXUR1 and 2, sister genes to
FERONIA/SIRENE, are male factors for coordinated fertilization', *Curr Biol.*, vol. 19, pp. 1327–
1331, 2009.
[16] Keinath NF, Kierszniowska S, Lorek J, Bourdais G, Kessler SA, Shimosato-Asano H, Grossniklaus U,
Schulze WX, Robatzek S, Panstruga RJ, 'PAMP (pathogen-associated molecular pattern)-induced
changes in plasma membrane compartmentalization reveal novel components of plant
immunity.', *J Biol Chem*, vol. 285, pp. 39140–39149, 2010.
[17] Duan Q, Kita D, Li C, Cheung AY, Wu HM., 'FERONIA receptor-like kinase regulates RHO GTPase
signaling of root hair development', *PNAS*, vol. 107, pp. 17821–17826, 2010.
[18] Galindo-Trigo S, Blanco-Touriñán N, DeFalco TA, Wells ES, Gray JE, Zipfel C, Smith LM., 'CrRLK1L
receptor-like kinases HERK1 and ANJEA are female determinants of pollen tube reception', *EMBO*
Rep, vol. 21, no. 2, p. e48466, 2020.
[19] Zhu L, Chu LC, Liang Y, Zhang XQ, Chen LQ, Ye D., 'The Arabidopsis CrRLK1L protein kinases BUPS1
and BUPS2 are required for normal growth of pollen tubes in the pistil', *Plant J*, vol. 95, no. 3, pp.
474–486, 2018.
[20] Li C, Yeh F-L,Cheung AY, Duan Q, KitaD, et al., 'Glycosylphosphatidylinositol-anchored proteins as
chaperones and co-receptors for FERONIA receptor kinase signaling in Arabidopsis', *eLife*, vol. 4,
p. e06587, 2015.
[21] Richter J, Ploderer M, Mongelard G, Gutierrez L, Hauser MT., 'Role of CrRLK1L Cell Wall Sensors
HERCULES1 and 2, THESEUS1, and FERONIA in Growth Adaptation Triggered by Heavy Metals and
Trace Elements', *Front Plant Sci*, vol. 8, p. 1554, 2017.
[22] Richter J, Watson JM, Stasnik P, Borowska M, Neuhold J, Berger M, Stolt-Bergner P, Schoft V,
Hauser MT., 'Multiplex mutagenesis of four clustered CrRLK1L with CRISPR/Cas9 exposes their
growth regulatory roles in response to metal ions', *Sci Rep*, vol. 8, no. 1, p. 12182, 2018.
[23] Stegmann M, Monaghan J, Smakowska-Luzan E, Rovenich H, Lehner A, et al., 'The receptor kinase
FER is a RALF-regulated scaffold controlling plant immune signaling', *Science*, vol. 355, pp. 287–
289, 2017.
[24] Ngo QA, Vogler H, Lituiev DS, Nestorova A, Grossniklaus U., 'A calcium dialog mediated by the
FERONIA signal transduction pathway controls plant sperm delivery', *Dev Cell*, vol. 29, pp. 491–
500, 2014.
[25] Yu F, Qian L, Nibau C, Duan Q, Kita D, et al., 'FERONIA receptor kinase pathway suppresses
abscisic acid signaling in Arabidopsis by activating ABI2 phosphatase', *PNAS*, vol. 109, pp. 14693–
14698, 2012.
[26] Mao D, Yu F, Li J, Van de Poel B, Tan D, et al., 'FERONIA receptor kinase interacts with S-
adenosylmethionine synthetase and suppresses S-adenosylmethionine production and ethylene
biosynthesis in Arabidopsis', *Plant Cell Env.*, vol. 38, pp. 2566–2574, 2015.
[27] Deslauriers SD, Larsen PB., 'FERONIA is a keymodulator of brassinosteroid and ethylene
responsiveness in Arabidopsis hypocotyls', *Mol Plant*, vol. 3, pp. 626–640, 2010.
[28] Shih H-W, Miller ND, Dai C, Spalding EP, Monshausen GB., 'The receptor-like kinase FERONIA is
required for mechanical signal transduction in Arabidopsis seedlings', *Curr Biol.*, vol. 24, pp.
1887–1892, 2014.
[29] Denness L, McKenna JF, Segonzac C, Wormit A, Madhou P, et al., 'Cell wall damage-induced lignin
biosynthesis is regulated by a reactive oxygen species- and jasmonic acid-dependent process in
Arabidopsis', *Plant Physiol*, vol. 156, pp. 1367–1374, 2011.
[30] Okuda S, Tsutsui H, Shiina K, Sprunck S, Takeuchi H, Yui R, Kasahara RD, Hamamura Y, Mizukami
A, Susaki D, Kawano N, Sakakibara T, Namiki S, Itoh K, Otsuka K, Matsuzaki M, Nozaki H, Kuroiwa
T, Nakano A, Kanaoka MM, Dresselhaus T, Sasaki N, Higashiyama T., 'Defensin-like polypeptide
LUREs are pollen tube attractants secreted from synergid cells', *Nature*, vol. 458, no. 7236, pp.
357–361, 2009.

[31]   Huck N, Moore JM, Federer M, Grossniklaus U., 'The Arabidopsis mutant feronia disrupts the
1003   female gametophytic control of pollen tube reception', *Development*, vol. 130, pp. 2149–2159,
1004   2003.
1005   [32]   Rotman N, Rozier F, Boavida L, Dumas C, Berger F, Faure J-E., 'Female control of male gamete
1006   delivery during fertilization in Arabidopsis thaliana', *Curr Biol.*, vol. 13, pp. 432–436, 2003.
1007   [33]   Ge Z, Bergonci T, Zhao Y, Zou Y, Du S, et al., 'Arabidopsis pollen tube integrity and sperm release
1008   are regulated by RALF-mediated signaling', *Science*, vol. 358, pp. 1596–1600, 2017.
1009   [34]   Haruta M, Sabat G, Stecker K, Minkoff BB, Sussman MR., 'A peptide hormone and its receptor
1010   protein kinase regulate plant cell expansion', *Science*, vol. 343, pp. 408–411, 2014.
1011   [35]   Dünser K, Gupta S, Herger A, Feraru MI, Ringli C, Kleine-Vehn J., 'Extracellular matrix sensing by
1012   FERONIA and Leucine-Rich Repeat Extensins controls vacuolar expansion during cellular
1013   elongation in Arabidopsis thaliana', *EMBO J*, vol. 38, no. 7, p. pii: e100353, 2019.
1014   [36]   Xiao Y, Stegmann M, Han Z, DeFalco TA, Parys K, Xu L, Belkhadir Y, Zipfel C, Chai J., 'Mechanisms
1015   of RALF peptide perception by a heterotypic receptor complex', *Nature*, vol. 572, no. 7768, pp.
1016   270–274, 2019.
1017   [37]   Mecchia MA, Santos-Fernandez G, Duss NN, Somoza SC, Boisson-Dernier A, et al., 'RALF4/19
1018   peptides interact with LRX proteins to control pollen tube growth in Arabidopsis', *Science*, vol.
1019   358, pp. 1600–1603, 2017.
1020   [38]   Capron A, Gourgues M, Neiva LS, Faure J-E, Berger F, Pagnussat G, Krishnan A, Alvarez-Mejia C,
1021   Vielle-Calzada JP, Lee YR, Liu B, Sundaresan V., 'Maternal control of male-gamete delivery in
1022   Arabidopsis involves a putative GPI-anchored protein encoded by the LORELEI gene', *Plant Cell*,
1023   vol. 20, pp. 3038–3049, 2008.
1024   [39]   Boisson-Dernier A, Kessler SA, Grossniklaus U., 'The walls have ears: the role of plant CrRLK1Ls in
1025   sensing and transducing extracellular signals', *J Exp Bot*, vol. 62, 2011.
1026   [40]   Hématy K, Höfte H., 'Novel receptor kinases involved in growth regulation', *Curr Opin Plant Biol.*,
1027   vol. 11, no. 3, pp. 321–328, 2008.
1028   [41]   Cheung AY, Wu HM., 'THESEUS 1, FERONIA and relatives: a family of cell wall-sensing receptor
1029   kinases?', *Curr Opin Plant Biol*, vol. 14, no. 6, pp. 632–641, 2011.
1030   [42]   Nissen KS, Willats WGT, Malinovsky FG., 'Understanding CrRLK1L Function: Cell Walls and Growth
1031   Control', *Trends Plant Sci*, vol. 21, no. 6, pp. 516–527, 2016.
1032   [43]   Feng W, Kita D, Peaucelle A, Cartwright HN, Doan V, Duan Q, Liu MC, Maman J, Steinhorst L,
1033   Schmitz-Thom I, Yvon R, Kudla J, Wu HM, Cheung AY, Dinneny JR., 'The FERONIA Receptor Kinase
1034   Maintains Cell-Wall Integrity during Salt Stress through Ca2+ Signaling', *Curr Biol.*, vol. 28, no. 5,
1035   pp. 666–675, 2018.
1036   [44]   Lindner H, Müller LM, Boisson-Dernier A, Grossniklaus U., 'CrRLK1L receptor-like kinases: not just
1037   another brick in the wall', *Curr Opin Plant Biol*, vol. 15, no. 6, pp. 656–669, 2012.
1038   [45]   Tang W, Lin W, Zhou X, Guo J, Dang X, Li B, Lin D, Yang Z., 'Mechano-transduction via the pectin-
1039   FERONIA complex activates ROP6 GTPase signaling in Arabidopsis pavement cell morphogenesis',
1040   *Curr Biol*, vol. S0960-9822, no. 21, pp. 01590–6, 2021.
1041   [46]   Lin W, Tang W, Pan X, Huang A, Gao X, Anderson CT, Yang Z., 'Arabidopsis pavement cell
1042   morphogenesis requires FERONIA binding to pectin for activation of ROP GTPase signaling', *Curr
1043   Biol*, vol. S0960-9822, no. 21, pp. 01589–X, 2021.
1044   [47]   Galindo-Trigo S, Gray JE, Smith LM., 'Conserved Roles of CrRLK1L Receptor-Like Kinases in Cell
1045   Expansion and Reproduction from Algae to Angiosperms', *Front Plant Sci*, vol. 7, p. 1269, 2016.
1046   [48]   Bowman JL, Kohchi T, Yamato KT, Jenkins J, Shu S, Ishizaki K, et al., 'Insights into Land Plant
1047   Evolution Garnered from the Marchantia polymorpha Genome', *Cell*, vol. 171, no. 2, pp. 287-
1048   304.e15, 2017.
1049   [49]   Rövekamp M, Bowman JL, Grossniklaus U., 'Back to the Rhizoids of Plant Sexual Development.',
1050   *Abstr. 23rd Int. Congr. Sex. Plant Reprod. Porto Port.*, p.
1051   https://www.fc.up.pt/sprporto2014/doc/23rd%20International%20congress%20on%20sexual%2
1052   0plant%20reproduction%20ABSBook.pdf, 2014.

[50] Bowman JL, Floyd SK, Sakakibara K, 'Green genes-comparative genomics of the green branch of
life', *Cell*, vol. 129, no. 2, pp. 229–234, 2007.
[51] Nishiyama T, Fujita T, Shin-I T, Seki M, Nishide H, Uchiyama I, Kamiya A, Carninci P, Hayashizaki Y,
Shinozaki K, et al., 'Comparative genomics of Physcomitrella patens gametophytic transcriptome
and Arabidopsis thaliana: implication for land plant evolution', *PNAS*, vol. 100, pp. 8007–8012,
2003.
[52] Qiu YL, Li L, Wang B, Chen Z, Knoop V, Groth-Malonek M, Dombrovska O, Lee J, Kent L, Rest J, et
al., 'The deepest divergences in land plants inferred from phylogenomic evidence', *PNAS*, vol.
103, pp. 15511–15516, 2006.
[53] Wickett NJ, Mirarab S, Nguyen N, Warnow T, Carpenter E, Matasci N, Ayyampalayam S, Barker
MS, Burleigh JG, Gitzendanner MA, et al., 'Phylotranscriptomic analysis of the origin and early
diversification of land plants', *PNAS*, vol. 1111, pp. E4859–E4868, 2014.
[54] Mishler BD, Churchill SP, 'A cladistic approach to the phylogeny of the "bryophytes"', *Brittonia*,
vol. 36, no. 4, pp. 406–424, 1984.
[55] Kenrick P, Crane PR, 'The origin and early evolution of plants on land', *Nature*, vol. 389, pp. 33–
39, 1997.
[56] Floyd SK, Bowman JL., 'The ancestral developmental tool kit of land plants', *Int J Plant Sci*, vol.
168, pp. 1–35, 2007.
[57] Bowman JL, Araki T, Arteaga-Vazquez MA, Berger F, Dolan L, Haseloff J, Ishizaki K, Kyozuka J, Lin
SS, Nagasaki H, Nakagami H, Nakajima K, Nakamura Y, Ohashi-Ito K, Sawa S, Shimamura M,
Solano R, Tsukaya H, Ueda T, Watanabe Y, Yamato KT, Zachgo S, Kohchi T, 'The Naming of Names:
Guidelines for Gene Nomenclature in Marchantia', *Plant Cell Physiol*, vol. 57, no. 2, pp. 257–261,
2016.
[58] Ishizaki K, 'Evolution of land plants: insights from molecular studies on basal lineages', *Biosci
Biotechnol Biochem*, vol. 81, no. 1, pp. 73–80, 2017.
[59] Honkanen S, Jones VAS, Morieri G, Champion C, Hetherington AJ, Kelly S, Proust H, Saint-Marcoux
D, Prescott H, Dolan L., 'The Mechanism Forming the Cell Surface of Tip-Growing Rooting Cells Is
Conserved among Land Plants', *Curr Biol.*, vol. 26, no. 23, pp. 3238–3244, 2016.
[60] Westermann J, Streubel S, Franck CM, Lentz R, Dolan L, Boisson-Dernier A., 'An Evolutionarily
Conserved Receptor-like Kinases Signaling Module Controls Cell Wall Integrity During Tip
Growth', *Curr Biol.*, vol. 29, no. 23, p. 4153, 2019.
[61] Kessler SA, Lindner H, Jones DS, Grossniklaus U., 'Functional analysis of related CrRLK1L receptor-
like kinases in pollen tube reception', *EMBO Rep*, vol. 16, pp. 107–115, 2014.
[62] Hirano N, Marukawa Y, Abe J, Hashiba S, Ichikawa M, Tanabe Y, 'A Receptor-Like Kinase, Related
to Cell Wall Sensor of Higher Plants, is Required for Sexual Reproduction in the Unicellular
Charophycean Alga, Closterium peracerosum–strigosum– littorale Complex', *Plant Cell Physiol*,
vol. 56, no. 7, pp. 1456–1462, 2015.
[63] Ginanjar EF, Teh OK, Fujita T., 'Characterisation of rapid alkalinisation factors in Physcomitrium
patens reveals functional conservation in tip growth', *New Phytol*, p. doi: 10.1111/nph.17942,
2021.
[64] Schwede T, Kopp J, Guex N, Peitsch MC, 'SWISS-MODEL: an automated protein homology-
modeling server', *Nucleic Acids Res.*, vol. 31, no. 13, pp. 3381–3385, 2003.
[65] Liu L, Zheng C, Kuang B, Wei L, Yan L, Wang T, 'Receptor-Like Kinase RUPO Interacts with
Potassium Transporters to Regulate Pollen Tube Growth and Integrity in Rice', *PLoS Genet*, vol.
12, no. 7, p. e1006085, 2016.
[66] Schulze-Muth P, Irmler S, SchröderG, Schröder J, 'Novel type of receptor-like protein kinase from
a higher plant (Catharanthus roseus): cDNA, gene, intramolecular autophosphorylation, and
identification of a threonine important for auto- and substrate phosphorylation', *J Biol Chem*, vol.
271, pp. 26684–26689, 1996.
[67] Flores-Sandoval E, Dierschke T, Fisher TJ, Bowman JL., 'Efficient and Inducible Use of Artificial
MicroRNAs in Marchantia polymorpha.', *Plant Cell Physiol*, vol. 57, pp. 281–290, 2016.

[68] Althoff F, Kopischke S, Zobell O, Ide K, Ishizaki K, Kohchi T, Zachgo S., 'Comparison of the MpEF1α
and CaMV35 promoters for application in Marchantia polymorpha overexpression studies',
Transgenic Res., vol. 23, no. 2, pp. 235–244, 2014.
[69] Sachs AB, Varani G., 'Eukaryotic translation initiation: there are (at least) two sides to every
story.', *Nat Struct Biol*, vol. 7, no. 5, pp. 356–61, 2000.
[70] Gonneau M, Desprez T, Martin M, Doblas VG, Bacete L, Miart F, Sormani R, Hématy K, Renou J,
Landrein B et al., 'Receptor kinase THESEUS1 is a RAPID ALKALINIZATION FACTOR34 receptor in
Arabidopsis', *Curr Biol.*, vol. 28, pp. 2452–2458, 2018.
[71] Gachomo EW, Jno Baptiste L, Kefela T, Saidel WM, Kotchoni SO, 'The Arabidopsis CURVY1 (CVY1)
gene encoding a novel receptor-like protein kinase regulates cell morphogenesis, flowering time
and seed production', *BMC Plant Biol*, vol. 14, p. 221, 2014.
[72] Merz D, Richter J, Gonneau M, Sanchez-Rodriguez C, Eder T, Sormani R, Martin M, Hématy K,
Höfte H, Hauser MT, 'T-DNA alleles of the receptor kinase THESEUS1 with opposing effects on cell
wall integrity signaling', *J Exp Bot*, vol. 68, pp. 4583–4593, 2017.
[73] Chiyoda S, Ishizaki K, Kataoka H, Yamato KT, Kohchi T, 'Direct transformation of the liverwort
Marchantia polymorpha L. by particle bombardment using immature thalli developing from
spores', *Plant Cell Rep*, vol. 27, no. 9, pp. 1467–1473, 2008.
[74] Malivert A, Erguvan Ö, Chevallier A, Dehem A, Friaud R, Liu M, Martin M, Peyraud T, Hamant O,
Verger S., 'FERONIA and microtubules independently contribute to mechanical integrity in the
Arabidopsis shoot', *PLoS Biol 2021 Nov 121911*, vol. 19, no. 11, p. e3001454, 2021.
[75] Baumberger N, Ringli C, Keller B., 'The chimeric leucine-rich repeat/extensin cell wall protein
LRX1 is required for root hair morphogenesis in Arabidopsis thaliana', *Genes Dev*, vol. 15, no. 9,
pp. 1128–39, 2001.
[76] Kim CM, Dolan L., 'ROOT HAIR DEFECTIVE SIX-LIKE Class I Genes Promote Root Hair Development
in the Grass Brachypodium distachyon.', *PLoS Genet*, vol. 12, no. 8, p. e1006211, 2016.
[77] Srivastava R, Liu JX, Guo H, Yin Y, Howell SH, 'Regulation and processing of a plant peptide
hormone, AtRALF23, in Arabidopsis', *Plant J*, vol. 59, no. 6, pp. 930–939, 2009.
[78] Boisson-Dernier A, Franck CM, Lituiev DS, Grossniklaus U., 'Receptor-like cytoplasmic kinase
MARIS functions downstream of CrRLK1L-dependent signaling during tip growth', *PNAS*, vol. 112,
pp. 12211–12216, 2015.
[79] Liao H-Z, Zhu M-M, Cui H-H, Du X-Y, Tang Y, et al., 'MARIS plays important roles in Arabidopsis
pollen tube and root hair growth', *J Integr Plant Biol*, vol. 58, pp. 927–940, 2016.
[80] Bouton S, Leboeuf E, Mouille G, Leydecker MT, Talbotec J, Granier F, Lahaye M, Höfte H, Truong
HN., 'QUASIMODO1 encodes a putative membrane-bound glycosyltransferase required for
normal pectin synthesis and cell adhesion in Arabidopsis.', *Plant Cell*, vol. 14, no. 10, pp. 2577–90,
2002.
[81] Bowen EJ, 'A NOTE ON THE CONDUCTION OF WATER IN FIMBRIARIA BLEUMEANA', *Ann. Bot.*, vol.
49, no. 196, pp. 844–848, 1935.
[82] Fujikura U, Elsaesser L, Breuninger H, Sánchez-Rodríguez C, Ivakov A, Laux T, Findlay K, Persson S,
Lenhard M., 'Atkinesin-13A modulates cell-wall synthesis and cell expansion in Arabidopsis
thaliana via the THESEUS1 pathway', *PLoS Genet*, vol. 10, no. 9, p. e1004627, 2014.
[83] Pierik R, Tholen D, Poorter H, Visser EJ, Voesenek LA, 'The Janus face of ethylene: growth
inhibition and stimulation', *Trends Plant Sci*, vol. 11, no. 4, pp. 176–183, 2006.
[84] Wolf S, Höfte H., 'Growth Control: A Saga of Cell Walls, ROS, and Peptide Receptors', *Plant Cell*,
vol. 26, no. 5, pp. 1848–1856, 2014.
[85] Humplík JF, Bergougnoux V, Van Volkenburgh E, 'To Stimulate or Inhibit? That Is the Question for
the Function of Abscisic Acid', *Trends Plant Sci*, vol. 22, no. 10, pp. 830–841, 2017.
[86] Flores-Sandoval E, Eklund DM, Bowman, JL, 'A Simple Auxin Transcriptional Response System
Regulates Multiple Morphogenetic Processes in the Liverwort Marchantia polymorpha', *PLoS
Genet*, vol. 11, p. e1005207, 2015.

[87] De Rybel B, van den Berg W, Lokerse AS., Liao C-Y, van Mourik H, Moller B, Llavata-Peris CI, Weijers D, 'A Versatile Set of Ligation-Independent Cloning Vectors for Functional Studies in Plants', *Plant Physiol.*, vol. 156, no. 3, pp. 1292–1299, 2011.

[88] Rövekamp M, Bowman JL, Grossniklaus U, 'Marchantia MpRKD Regulates the Gametophyte-Sporophyte Transition by Keeping Egg Cells Quiescent in the Absence of Fertilization', *Curr. Biol.*, vol. 26, no. 13, pp. 1782–1789, 2016.

[89] Zuker M., 'Mfold Web Server for Nucleic Acid Folding and Hybridization Prediction', *Nucleic Acids Res*, vol. 31, no. 13, pp. 3406–3415, 2003.

[90] Aach J, Mali P, Church GM, 'CasFinder: Flexible algorithm for identifying specific Cas9 targets in genomes', *BioRxiv Doi 101101005074*, 2014.

[91] Curtis MD, Grossniklaus U, 'A gateway cloning vector set for high-throughput functional analysis of genes in planta', *Plant Physiol.*, vol. 133, no. 2, pp. 462–469, 2003.

[92] Ishizaki K, Nishihama R, Ueda M, Inoue K, Ishida S, Nishimura Y, Shikanai T, Kohchi T, 'Development of Gateway Binary Vector Series with Four Different Selection Markers for the Liverwort Marchantia polymorpha', *PLoS One*, vol. 10, no. 9, p. e0138876, 2015.

[93] Vandesompele J, De Preter K, Pattyn F, Poppe B, Van Roy N, De Paepe A, Speleman F., 'Accurate normalization of real-time quantitative RT-PCR data by geometric averaging of multiple internal control genes', *Genome Biol*, vol. 3, no. 7, p. RESEARCH0034, 2002.

[94] Kubota A, Ishizaki K, Hosaka M, Kohchi T, 'Efficient Agrobacterium-mediated transformation of the liverwort Marchantia polymorpha using regenerating thalli.', *Biosci Biotechnol Biochem*, vol. 77, no. 1, pp. 167–172, 2013.

[95] Ishizaki K, Chiyoda S, Yamato KT, Kohchi, T, 'Agrobacterium-Mediated Transformation of the Haploid Liverwort Marchantia polymorpha L., an Emerging Model for Plant Biology', *Plant Cell Physiol.*, vol. 49, no. 7, pp. 1084–1091, 2008.

[96] Barnes CR, Land WJG., 'Bryological papers. I. The origin of air chambers', *Bot Gaz*, vol. 44, pp. 197–213, 1907.

[97] Hughes SJ, 'On conidia of fungi, and gemmae of algae, bryophytes, and pteridophytes', *Can J Bot*, vol. 49, pp. 1319–1339, 1971.

[98] Bent A., 'Agrobacterium Protocols', *Totowa NJ Humana Press*, vol. Volume I, no. Second edition, 2006.

[99] Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S, et al., 'Fiji: An open-source platform for biological-image analysis', *Nat. Methods*, vol. 9, no. 7, pp. 676–682, 2012.

[100] Solly JE, Cunniffe NJ Harrison CJ, 'Regional Growth Rate Differences Specified by Apical Notch Activities Regulate Liverwort Thallus Shape', *Curr. Biol.*, vol. 27, no. 1, pp. 16–26, 2017.

[101] Felekis D, Muntwyler, S., Vogler H, Beyeler F, Grossniklaus U and Nelson BJ, 'Quantifying growth mechanics of living, growing plant cells in situ using microrobotics.', *Micro Nano Lett*, vol. 6, pp. 311–6, 2011.

[102] Vogler H, Draeger C, Weber A, Felekis D, Eichenberger C, Routier-Kierzkowska AL, Boisson-Dernier A, Ringli C, Nelson BJ, Smith RS, Grossniklaus U., 'The pollen tube: a soft shell with a hard core. 2013 Feb;73(4):617-27.', *Plant J*, vol. 73, no. 4, pp. 617–27, 2013.

[103] Laemmli UK, 'Cleavage of structural proteins during the assembly of the head of bacteriophage T4', *Nature*, vol. 227, no. 5259, pp. 680–685, 1970.

[104] Bushnell B, 'BBMap: a fast, accurate, splice-aware aligner (No. LBNL-7065E)', *Lawrence Berkeley Natl. LabLBNL Berkeley CA U. S.*, 2014.

[105] Trapnell C, Pachter L, Salzberg SL, 'TopHat: discovering splice junctions with RNA-Seq', *Bioinformatics*, vol. 25, no. 9, pp. 1105–1111, 2009.

[106] Langdon WB, 'Performance of genetic programming optimised Bowtie2 on genome comparison and analytic testing (GCAT) benchmarks', *BioData Min.*, vol. 8, no. 1, p. 1, 2015.

[107] Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al, 'The sequence alignment/map format and SAMtools', *Bioinformatics*, vol. 25, no. 16, pp. 2078–2079, 2009.

[108] Anders S, Pyl PT, Huber W., 'HTSeq—a Python framework to work with high-throughput
sequencing data', *Bioinformatics*, vol. 31, no. 2, pp. 166–169, 2015.
[109] Love MI, Huber W, Anders S, 'Moderated estimation of fold change and dispersion for RNA-seq
data with DESeq2', *Genome Biol.*, vol. 15, no. 12, p. 550, 2014.
[110] Team RC, 'R: A language and environment for statistical computing', 2013.
[111] Wickham H, 'ggplot2: elegant graphics for data analysis', vol. Springer, 2016.
[112] Lolle SJ, Berlyn GP, Engstrom EM, Krolikowski KA, Reiter W-D , Pruitt RE, 'Developmental
Regulation of Cell Interactions in the Arabidopsis fiddlehead-1 Mutant: A Role for the Epidermal
Cell Wall and Cuticle', *Dev. Biol.*, vol. 189, pp. 311–321, 1997.
[113] DeLano WL, 'Pymol: An open-source molecular graphics tool', *CCP4 Newsl. Protein Crystallogr.*,
vol. 40, no. 1, pp. 82–92, 2002.

1216
1217

1218

Figure 1. *Cr*RLK1Ls Are Conserved among Land Plants

(A) Representation of AtFER and MpFER proteins. The malectin domains are

represented in blue and kinase domains in green. SP, signal peptide; TM, transmembrane

domain (yellow). Amino acids important for regulation and activity are marked in red,

violet, and light blue as described in Fig. 1E.

(B) A rooted neighbor-joining tree of the amino acid sequence of the predicted malectin-

like domain (corresponding to aa 76-431 of MpFER). *Cr*RLK1L members from *Marchantia*

polymorpha (Mp, green), *Physcomitrium patens* (Pp), *Sphagnum fallax* (Sphfalx),

Ceratodon purpureus (Cpur)*, Anthoceros agrestis* Bonn (AagrBONN)*, Anthoceros*

agrestis Oxford (AagrOXR)*, Anthoceros punctatus* (Apun)*, Selaginella moellendorffii* (Smo), *Amborella trichopoda* (AmTr)*,* and *Arabidopsis thaliana* (At, blue) were used. The best hit from *Chara braunii* (CHBRA) was also included. The numbers indicate the bootstrap values (%) from 1000 replications. The given scale represents a substitution frequency of 0.1 amino acids per site.

(C) Cartoon representation of the predicted three-dimensional structure of the MpFER malectin-like domain, showing predicted alpha-helix and beta-sheet structures.

(D) Structural superposition of the malectin-like domains of AtFER (blue) and MpFER (green).

(E) Alignment of the cytoplasmic domains of MpFER, AtERU, AtFER, and *Cr*RLK. Kinase domains are in green, putative phosphorylation sites in light blue (Ser) and violet (Thr), and the conserved catalytic Lys in red.

See also Fig. S1

1242

Figure 2. Mp*FER* Is Broadly Expressed in most Tissues throughout the *M.*

***polymorpha* Life Cycle**

Expression of $_{pro}$Mp*FER*:*trpVNS* in different organs of male and female *M. polymorpha*

plants. Fluorescence is visualized as either green (epifluorescence microscope) or yellow

(confocal microscope) color, depending on the panel.

(A to D) Bright field (left) and epifluorescence (right) images of a thallus, (A) rhizoid (C),

gemmae cups (C), and an archegoniophore (D). Red arrows indicate meristematic zones,

the yellow arrow gemmae. Scale bars, 0.5 mm.

(E and F) Confocal images of archegonia before (E) and 2 days after fertilization (F).

Scale bar, 50 μm.

(G) Bright field (left) and epifluorescence (right) images of an antheridiophore. Scale bar,

1 mm.

(H) Confocal image of antheridia. Scale bar, 100 μm.

(I) Confocal image of sporophyte and spores. Scale bar, 25μm.

1257

Figure 3. Mp*FER* Controls Cell Size during Gametophyte Development

(A) Representation of the mature Mp*FER* mRNA. The coding sequence is in green, locations of the amiRNA target sites in blue. The location of the T-DNA insertion in Mp*fer-1* is shown.

(B) Base complementary of mature amiR-Mp*FER*1, amiR-Mp*FER*2, and amiR-Mp*FER*3 with the Mp*FER* transcript. Very low complementary was found between amiR-Mp*FER*3 and At*FER*.

(C) Representative pictures of 10-day old gemmalings of wild-type (WT), Mp*fer-1*, and amiR-Mp*FER*3-2 lines. Scale bar, 1 mm.

(D) Violin plot of thallus area of 14-day old gemmalings of WT, Mp*fer-1*, and amiR-Mp*FER*3-2 lines. n = 30. Different letters indicate significant differences according to the non-parametric Kruskal-Wallis test and Dunn's test for multiple comparisons (P < 0.001). Orange circles indicate the group mean and the corresponding vertical bars the standard deviation for each group.

(E) Relative expression level of Mp*FER* in thallus tissue from WT and two independent

insertion lines of amiR-Mp*FER*2 and amiR-Mp*FER*3, as measured by qRT-PCR. Mp*EF1*

was used as internal control. Shown are means ± standard errors of the means (SEM) of

three biological replicates. Statistical analysis was performed by one-way analysis of

variance (ANOVA) followed by a post-hoc Duncan test (*P < 0.01).

(F) Violin plot of cell size in WT, *Mpfer-1*, and amiR-Mp*FER*3-2 lines. Only areas with fully

expanded cells in approximately 3-week old plants were used for measurements.

Different letters indicate significant differences according to the non-parametric Kruskal-

Wallis test and Dunn's test for multiple comparisons (P < 0.001). Orange circles indicate

the group mean and the corresponding vertical bars the standard deviation for each

group.

See also Figs. S2,S3

1284

1285

B

MpFER 5'...CCATGTACAGGCTGAATGT-CGGAGGCGGAA...3'
Mpfer-2 5'...CCATGTACAGGCTGAATGTACGGAGGCGGAA...3'
Mpfer-3 5'...CCATGTACAGGCTGAATGTTCGGAGGCGGAA...3'
Mpfer-4 5'...CCATGTACAGGCTGA--------GGCGGAA...3'
Mpfer-5 5'...CCATGTACAGGCTGAAT-T-CGGAGGCGGAA...3'

MpFER 5'...TGTACCCGAACGATGTGCCGAGGTGGATGGC...3'
Mpfer-6 5'...TGTACCCGAACGATGTGCC-AGGTGGATGGC...3'
Mpfer-7 5'...TGTACCCGAACGAGGTG--------GATGGC...3'
Mpfer-8 5'...TGTACCCGAACGATGTGCC-AGGTGGATGGC...3'

1286

Figure 4. The Integrity of Thalli from Mp*fer* Knock-out Lines Is Severely Affected

(A) Representation of the mature Mp*FER* mRNA with the coding sequence in green, and the location of the gRNA target sites in red.

(B) Sequences of gRNA target sites in the confirmed Mp*fer* knock-out mutants. Deletions and insertions are highlighted in red.

(C) Representative pictures of 14-day old gemmalings of the wild type (WT), Mp*fer-1*, and two independent Mp*fer* knock-out mutants. Scale bar, 0.5 cm.

(D) Violin plot of thallus area of 7-day old gemmalings of WT, Mp*fer-1*, and two independent Mp*fer* knock-out mutants. n = 25. Different letters indicate significant differences in a one-way analysis of variance (ANOVA) follow by a post-hoc Duncan test (P < 0.01). Orange circles indicate the group mean and the corresponding vertical bars the standard deviation for each group.

(E) Representative pictures of 7-day old gemmalings growing in upside-down plates of WT, Mp*fer-1*, and two independent Mp*fer* knock-out mutants. Scale bar, 500 μm.

(F) Violin plot of rhizoid number in 3-day old gemmalings of WT, Mp*fer-1*, and two independent Mp*fer* knock-out mutants. n = 16. Statistical analysis was performed by one-way analysis of variance (ANOVA) followed by a post-hoc Duncan test (*P < 0.01).

(G) Trypan blue staining of 7-day old gemmalings of WT, Mp*fer-1*, and two independent Mp*fer* knock-out mutants. Scale bar, 1 mm.

(H) Violin plot of the reduction in the apparent stiffness of gemmaling tissue in Mp*fer-2* compared to the WT. Statistical analysis was performed using a Kruskal-Wallis test followed by a Tukey-Kramer post-hoc test (*P < 0.001). Orange circles indicate the group mean and the corresponding vertical bars the standard deviation for each group.

Figure 5. Conservation of *Cr*RLK1L Function in Land Plants

(A) Violin plot of the number of antheridiophores produced per plant for wild-type (WT, n=9) and two independent amiR-Mp*FER*3 lines (amiR-Mp*FER*3-2, n = 9; amiR-Mp*FER*3-3, n = 12). Different letters indicate a significant difference in a one-way analysis of variance (ANOVA) followed by a post-hoc Duncan test (P < 0.01).

(B) Antheridiophore splash platform size distribution of WT (n = 32), amiR-Mp*FER*2-3 (n

= 34), and amiR-Mp*FER*3-3 (n = 11) lines. For all antheridia with a stalk >8 mm, length

and width of the platform were recorded.

(C) qRT-PCR of Mp*FER* levels in antheridiophores in a WT and amiR-Mp*FER*3-3 line.

Mp*EF1* was used as internal control. Shown are means ± SEM of three biological

replicates. Statistical analysis was performed by one-way analysis of variance (ANOVA)

followed by a post-hoc Duncan test (*P < 0.01).

(D and E) Mature antheridia of a WT (D) and amiR-Mp*FER*3-3 line (E). Bright field images

(left) and in cross-section (right). The spermatogenous areas are indicated in red. Scale

bar, 100*μ*m.

(F) Representative pictures of 6-day old gemmalings of WT and Mp*fer-2* mutants at

different isoxaben concentrations. Scale bar, 1 mm.

(G) Thallus area of gemmalings of WT, Mp*fer-1*, and Mp*fer-2* lines growing on media

containing different isoxaben concentrations. n = 35. Shown are means ± SEM of two

biological replicates. Different letters indicate a significantly difference in a one-way

analysis of variance (ANOVA) followed by a post-hoc Duncan test (P < 0.05).

See also Fig. S3

1336

1337

Figure 6. Interspecific Complementation of *A. thaliana* and *M. polymorpha* Plants

with Reduced Levels of *FER* Activity

(A) Roots of independent *A. thaliana* lines expressing 35S*:*Mp*FER-GFP* in L*er* wild-type

(WT) and At*fer-1* plants. Scale bar, 200 μm.

(B) 10-day old gemmalings of WT and amiR-Mp*FER*3-2 *M. polymorpha* plants with and

without expression of At*FER-Cit*. Scale bar, 0.5 mm.

(C) Citrine expression in 7-day old gemmalings of WT and amiR-Mp*FER*3-2 plants with

and without expression of At*FER-Cit*. Scale bar, 100 μm.

(D) Cartoon representation of the *Cr*RLK1L^{ECD} polar interactions sites with LRE/RALF

when forming a protein complex. Upper panels correspond to contact areas between

AtFER (purple), AtLLG2 (blue-grey), and AtRALF23 (yellow), and lower panels to the

modeled MpFER structure (green) superimposed onto the same complex. Polar

interactions are depicted as doted yellow lines. Left panels show the predicted polar

interactions of the complex that are conserved between *A. thaliana* and *M. polymorpha*,

whereas the right pannels show the polar interaction between AtFER and AtLLG2 that is

predicted to be absent between MpFER and AtLLG2.

(E) Suppression of the amiR-Mp*FER* phenotype by a dominant-active version of Mp*MRI*

under its own promoter ($_{pro}$Mp*MRI*:*MRI*^{R240C}). Scale bar, 1 mm.

See also Figs. S4-S6

1357

1358

1359 Figure 7: Overexpression of Mp*FER* Affects Morphological Integrity

(A) Representative pictures of 10-day old gemmalings of wild-type (WT) and two different

lines overexpressing Mp*FER* (*pro*Mp*EF1*:Mp*FER-Cit)*. Scale bar, 1 mm. Inset: Citrine

expression at the plasma membrane in *pro*Mp*EF1:*Mp*FER-Cit* lines *#5* and #9 as observed

under confocal microscopy. Scale bar, 20 μm.

(B) Scanning electron microscopical images of thalli from 20-day old plants of WT and

*pro*Mp*EF1:*Mp*FER-Cit* lines *#5* and #9. The blue region demarks an air chamber, while air

pore cells are colored in red. Scale bar, 500 μm.

(C) Representative pictures of 3-day old gemmalings growing in upside-down plates of

WT and *pro*Mp*EF1:*Mp*FER-Cit #5* lines. Scale bar, 1 mm.

(D) Violin plot of the number of rhizoids in 3-day old gemmalings of WT and

$_{pro}$Mp*EF1:*Mp*FER-Cit* lines *#5* and #9. n = 20. Statistical analysis was performed by one-

way analysis of variance (ANOVA) followed by a post-hoc Duncan test (*P < 0.01).

(E) Graphs showing amount of effluxed chlorophyll as a function of time from 15-day old

gemmalings of WT, Mp*fer-2* and $_{pro}$Mp*EF1:*Mp*FER-Cit #5* lines.

See also Fig. S6