Isolation and characterization of *Pseudomonas chlororaphis* strain ST9 and its potential as a bioinoculant for agriculture

Iris Bertani [♦,1], Elisa Zampieri[♦,2,3], Cristina Bez[1], Andrea Volante[2,4], Vittorio Venturi[1*] and Stefano Monaco[2*]

♦ Equal Contribution

[1] International Centre for Genetic Engineering and Biotechnology, 34149 Trieste, Italy

[2] Council for Agricultural Research and Economics-Research Centre for Cereal and Industrial Crops, s.s. 11 to Torino, km 2.5, 13100 Vercelli, Italy

[3] Institute for Sustainable Plant Protection, National Research Council, Turin, Italy

[4] Council for Agricultural Research and Economics-Research Centre for Vegetable and Ornamental Crops, Corso Inglesi 508, 18038 Sanremo, IM, Italy (present address)

Keywords: rice, plant growth promoting bacteria, qPCR, microbiome, inoculum persistence, genome analysis

* Correspondence: Vittorio Venturi (venturi@icgeb.org) and Stefano Monaco (stefano.monaco@crea.gov.it)

Abstract

The development of biotechnologies based on beneficial microorganisms for improving soil fertility and crop yields could help addressing many current agriculture challenges, such as food security, climate change, pests control, soil depletion while decreasing the use of chemical fertilizers and pesticides. Plant Growth Promoting (PGP) microbes can be used as probiotics in order to increase plant tolerance/resistance to abiotic/biotic stresses and in this context strains belonging to the *Pseudomonas chlororaphis* group have shown to have potential as PGP candidates. In this work a new *P. chlororaphis* isolate is reported and tested for (i) *in vitro* PGP features, (ii) whole genome sequence analysis, and (iii) its effects on root microbiome, plant growth and on the expression of different plant genes in greenhouse experiments. The potential use of this *P. chlororaphis* strain as a plant probiotic is discussed.

Introduction

Agriculture of the 21st century has several challenges to face. Among them are the increase in population and the growing demand for food in the context of climate change, soil depletion, competition between different land uses and the need to reduce chemical fertilizers and pesticides. The use of beneficial microorganisms in crop production is an attractive solution as they can have beneficial effects on soil and plant by improving soil fertility, plant yields and reducing the use of agrochemicals (Hayat *et al*., 2010). The development of next-generation sequencing methodologies has led to numerous studies on plant microbiomes documenting that plants are colonized and live in association with a large number of microorganisms. Many of these, indicated as PGP (Plant Growth Promoting) microbes, play important roles in plant health and resistance to biotic and abiotic stresses (Berendsen *et al*., 2012; Bulgarelli *et al*., 2013; Backer *et al*., 2018; Orozco-Mosqueda *et al*., 2018; Khatoon *et al*., 2020). PGP microbes allow the reduction of fertilizers requirements, improving crop nutrient use efficiency (nitrogen, phosphate, etc.), and protecting the host plant by pathogens invasion through niche exclusion mechanisms and antibacterial/antifungal compound production (Bender *et al*., 2016; Pii *et al*., 2015). PGP microorganisms can also elicit transcriptional changes in hormone-, defence- and cell wall-related genes (Spaepen *et al*., 2014), increase root length (Hong *et al*., 1991), and activate auxin-response genes that reinforce plant growth (Ruzzi and Aroca, 2015). In the last decades the interest in developing PGP probiotic microorganisms has increased with the goal to reduce chemical fertilization (Mehnaz, 2016) and pesticide use, both in conventional and organic farming, with the aim of offering healthier food and improving sustainability of crop production (Hazra *et al*., 2018). One class of microorganisms that are being studied for many years for their application are Plant Growth Promoting Rhizobacteria (PGPR) (Oleńska *et al*., 2020).

Among the group of PGPR, strains belonging to the *Pseudomonas chlororaphis* species have been found in association with a wide range of plants, both mono- and dicotyledonous, and both wild and cultivated (Biessy *et al*., 2019). *Pseudomonas chlororaphis* is currently classified as four subspecies, namely *chlororaphis*, *aureofaciens*, *aurantiaca*, and *piscium* (Peix *et al*., 2007; Burr *et al*., 2010). Several strains of *P. chlororaphis* have shown good potential for application as plant probiotics (Zhao *et al*., 2013; Chen *et al*., 2015; Arrebola *et al*, 2019; Anderson *et al*., 2020) due to their rhizosphere colonization abilities and plant associated beneficial phenotypes such as chemotaxis and motility (Arrebola *et al*., 2020), biofilm formation (Calderon *et a*l., 2013, 2014), P solubilization (Ahemad, 2015), ACC deaminase (Glick, 2014), IAA production (Kang *et al*., 2006; Oh *et al*., 2013) and biocontrol. *P. chlororaphis* strains produce different antifungal compounds such as Prn (pyrrolnitrin), PCN (phenazine-1-carboxamide), PCA (phenazine-1-carboxylic acid), 2-OH-PHZ (2-hydroxyphenazine), HPR (2-hexyl-5-propyl-alkylresorcinol) and HCN (hydrogen cyanide). These molecules inhibit the growth of various phytopathogens belonging to the *Fusarium* group (Chin-A-Woeng *et al*., 1998; Hu *et al*., 2014) and to different species of *Colletotrichum,*

Phytophthora, *Pythium*, *Sclerotinia* and *Rhizoctonia* (Liu *et al.*, 2007), protecting plants such as maize (Tagele *et al.*, 2019), tomato (Zhang *et al.*, 2015). In addition, good product formulation protocols have been developed and commercialized products having *P. chlororaphis* strains have been produced (Nam *et al.*, 2018).

In this work, we report the isolation and characterization of *P. chlororaphis* strain ST9. This strain was able to colonize and persist in rice roots for the entire rice vegetative cycle and studies are presented of its effect on the root microbiome, on plant growth and on the expression of several plant genes. The potential use of this *P. chlororaphis* strain as a plant probiotic for rice cultivation is discussed.

Material and Methods

Strain isolation, growth and identification

One gram of bulk soil from Padriciano, Trieste, Italy (45° 39' 32" North, 13° 50' 28" East) was taken and resuspended in 5 ml of PBS (phosphate buffer solution). Serial dilutions were performed and plated on 1/6 Tryptic Soy medium (BD, Maryland, 21152. USA), solidified with 1.5% agar. Growth was allowed for 72 hrs at 28°C and single colonies were isolated and purified through further streaking. Several isolates displaying varying morphologies, were selected and identified. Identification occurred through 16S rRNA sequencing: briefly, primers fD1 and rR2 were used for the amplification of the 16S rRNA gene and primers 518F and 800R (Supplementary Table (ST) ST1) were used for its sequencing (Eurofins, Ebersberg, Germany). A small bacterial collection was established and *P. chlororaphis* ST9 was one of the isolates stored at -80°C for further studies.

In vitro **phenotypic characterization**

Presence of lipolytic and proteolytic activities was determined by streaking the bacterial isolates on 1/6 TSA (Tryptic Soy Agar) medium amended with 1% Glyceryl tributyrin (Smeltzer *et a*l., 1992) and 2% of powder milk (Huber *et al.*, 2001), respectively. Exopolysaccharide (EPS) production was tested by streaking the bacterial isolates on Yeast Extract Mannitol medium (Zlosnik *et al.*, 2008) and indole acetic acid (IAA) production was verified as described by Bric *et al.* (1991) using the Salkowski reagent. The 1-aminocyclopropane-1-carboxylic acid (ACC) deaminase activity was detected by means of M9 minimal medium with ACC as unique N source (Penrose *et al.*, 2003), while the ability to solubilize P was verified using the NPRBB growth medium (Nautiyal, 1999). N-acyl homoserine lactone (AHL) production was assessed by T-streak technique, using the biosensor *Chromobacterium violaceum* CV026 after incubation for 1-2 days (Steindler and Venturi, 2007). Motility was checked on M8 medium plates with 0.3% (swimming) or 0.5% (swarming) agar (Kohler *et al.*, 2000). Anti-bacterial and anti-fungal activities were tested by plating different plant pathogens (*Dickeya zeae, Pseudomonas fuscovaginae, Magnaporthe oryzae, Fusarium* sp*., Aspergillus* sp*.* and *Phytophora infestans*) adjacent to a 16 hrs old streak of ST9 on TSA medium: pathogens growth was checked after 72-96 hrs. Plant growth promotion traits were tested on 300 surface sterilized rice seeds. In detail, 16 hrs pre-germinated seeds were submerged for 2 hrs in a ST9 bacterial suspension (OD_{600} 0.5) or PBS, as control; after inoculum application seeds were rinsed with sterile water and allowed to germinate in the dark at 30°C. One hundred seeds were evaluated for emergence after 3 days, 100 seeds were grown for 10 days and then used to measure coleoptile length and the growth of the last 100 seeds was interrupted after 12 days for dry mass weight.

Generation of a rifampicin resistant spontaneous mutants

The *P. chlororaphis* ST9 strain was grown in 1/6 TS medium for 16 hrs at 30°C with 100 rpm shaking. A 1/100 dilution of the overnight culture was put in 1/6 TS medium with 15 µg/ml rifampicin and the culture was grown again in the same growth conditions. The procedure was repeated, increasing concentration of Rifampicin (25, 50 and 100 µg/ml), and the final culture was plated on 1/6 TS. One colony was chosen and streaked on TSArif100; its identity, *P. chlororaphis* ST9, was confirmed again by 16S rRNA sequencing.

Bacterial genomic DNA extraction and sequencing

Genomic DNA extraction for genome sequencing was performed with the pronase Sarkosyl lysis method (Better *et al.*, 1983*)*. 2 µg of DNA, quantified through Nanodrop (Thermo Scientific, Waltham, Massachusetts, USA) and checked by gel electrophoresis, were used for genome sequencing by the Exeter Sequencing Service (University of Exeter, Exeter, UK). The sequence was carried out using the Illumina technique with the Hiseq 2500 platform with the 125 base pair paired end system. Reads were assembled using SPAdes 3.9.03 (Bankevich *et al.*, 2012). The assembled *P. chlororaphis* ST9 genome was uploaded in the RAST Annotation Server (Overbeek *et al.*, 2014) and was automatically annotated using the RASTtk annotation scheme (Brettin *et al.*, 2015). The genome sequence is presented as a unique contig and it is available on the RAST server (http://rast.nmpdr.org; "guest" as login and password) under Genome ID 286.2086.

P. chlororaphis ST9 taxonomy

Multi-locus Sequence Analysis (MLSA) was performed using 5 housekeeping genes: *recA* (recombinase A), *gyrB* (DNA gyrase subunit B), *rpoD* (RNA polymerase sigma factor), *carA* (Carbamoyl-phosphate synthase small chain) and *atpD* (ATP synthase subunit beta), plus the 16S rRNA locus. Sequences of these loci were obtained from the ST9 genome sequence. For the phylogenetic analysis, we concatenated the gene sequences in the following order: 16S-*recA*-*gyrB*-*rpoD*-*carA*-*atpD* resulting in a single sequence. For the same genes, orthologue sequences from 16 *P. chlororaphis* species were obtained from the NCBI database and chained. The used sequences are reported in Supplementary Materials SM1. As outgroup, we used the orthologue concatenate from *Pseudomonas putida KT2440 and P. fluorescens Pf-01*. The phylogenetic analysis was performed using the NGPhylogeny.fr public platform (Lemoine *et al.*, 2019).

Plant inoculation

Seeds of *O. sativa* L. cv. Baldo were surface-sterilized with 50% sodium hypochlorite solution (commercial bleach) for 60 min, rinsed with sterile water and incubated in a wet dark environment for seven days at 30°C for germination. Two growth conditions (treatments) were considered: with and without the ST9 inoculation. The roots of 70 seedlings were soaked for 60 min in a ST9 RifR bacterial solution at 0.5 of OD_{600} (treated, "T" plants), while the control was performed soaking the

roots of 70 seedlings in a sterile PBS solution (untreated, "UT" plants). The inoculated and the control seedlings were transferred to plastic tubes containing 0.4% water-agar and Hoagland solution for 72 hrs and then transplanted into pots in greenhouse.

Greenhouse experiments

For each of the two treatments, fourteen plastic pots (23 cm x 21 cm) were used, filled with non-sterile paddy field soil (47.8% sand, 9.4% clay, 42.8% silt, pH 6.4, organic matter 1.45%) taken from the experimental rice field of CREA-CI in Vercelli (VC, Italy). Pots were placed in greenhouse under uncontrolled temperature, light and humidity parameters. In each pot, five seedlings, undergoing the same treatment, were sown. Plants were watered every day with tap water and kept in a greenhouse for 90 days, following the natural photoperiod. Four time points were considered: T1, 10 days post inoculation (dpi); T2, 28 dpi; T3, 40 dpi and T4, 90 dpi. At T1, T2 and T3 ten plants per treatment were sampled, washed and stored at -80°C for gene expression analysis. At T1, T2 and T4 ten plants per treatment were sampled and utilized for bacterial counting and microbiome analysis. A time line of the experiment is presented in SF1. The remaining 20 plants were used for physiological and morphological evaluations. In particular, Nitrogen Balance Index (NBI), indicator of the plant nitrogen status at the beginning of flowering stage (Tremblay *et al.*, 2012) was calculated as ratio between chlorophyll (CHL) and flavonoid (FLA) concentration recorded by the DUALEX 4 Scientific (Dx4) chlorophyll meter (Force-A, Paris, France) (Goulas *et al.*, 2004). Measurements were carried out on both adaxial and abaxial faces of the panicle leaf for each plant. The total height was also measured for each plant as well as the dry weight of shoots (obtained after 48 h at 65°C).

Colonization counts

The roots of 10 rice plants for each treatment at each time point were cleaned from soil and washed thoroughly under tap water before processing them. For T1, complete roots were macerated, while for T2 and T4, 600 mg of root samples were macerated and resuspended in 2 ml of PBS. Serial dilutions, up to 10^{-5}, were made and 100 μL of each dilution was plated in triplicate on TSA (Tryptic soy agar) and TSA with rifampicin (50 μg/L). Plates were incubated at 28°C for 48 hrs and the emerged colonies were counted and presented as function of plated volume, dilution factor, macerated volume and starting material weight in order to determine the number of total cultivable CFUs (Colony Forming Units) and of rifampicin resistant CFUs present in 1 g of roots.

Bacterial genomic DNA extraction and 16S rRNA gene amplicon library preparation and sequencing

Five hundred mg of root samples from 10 ST9-treated and untreated plants at T2 and T4 time points were used to extract DNA and perform microbiome analysis. DNA was extracted using the PowerSoil Kit (Qiagen, Hilden, Germany) following supplier instructions. For the 16S amplicon

libraries preparation, 12.5 ng of DNA, quantified using Nanodrop, were used for each sample to prepare the 16S rRNA amplicon libraries. Library preparation was done using the Illumina methodology following the 16S Metagenomic Sequencing Library Preparation protocol (https://emea.support.illumina.com/content/dam/illumina-support/documents/documentation/chemistry_documentation/16s/16s-metagenomic-library-prep-guide-15044223-b.pdf). Sequencing was performed at CBM scrl (Trieste, Italy) with MiSeq methodology.

Microbiome sequence analysis

FASTQ files were demultiplexed using the QIIME 1.9.1 split_libraries_fastq.py script and then analysed using DADA2 v1.4.0 (Callahan *et al.*, 2016) adapting the methods from the DADA2 Pipeline Tutorial (1.4) and including dereplication, singletons and chimera removing (sample inference was performed using the inferred error model, while chimeric sequences were removed using the removeBimeraDenovo function). The Greengenes (GG) database (McDonald *et al.*, 2012), giving a final Operational Taxonomic Unit (OTU) table, was employed to assign bacterial taxonomy by means of the assign Taxonomy function with a 97% sequence similarity. The resulting OTUs were clustered at Genus taxonomic level obtaining the final profile and abundance of bacterial taxa in the different samples. Statistical analysis was performed using the vegan package version 2.5-4 (Oksanen *et al.*, 2013) and phyloseq package (McMurdie and Holmes, 2013) in R version 3.5.2 (R CoreTeam, 2014). Relative abundances of OTUs between samples were calculated. To test the differential representation of microbial taxa in diverse samples the Deseq2 package (Love *et al.*, 2014) was used.

RNA extraction and cDNA conversion

The RNA from three biological replicates for each thesis at each time point was extracted using the RNeasy Plant Mini Kit (Qiagen, Hilden, Germany), according to the manufacturer's instructions. After the RNA extraction, any amount of DNA was removed using DNase (RQ1 RNase-Free DNase, Promega, Madison, Wisconsin, USA) and measured using Qubit (Invitrogen, ThermoFisher Scientific, Massachusetts, USA). The absence of genomic DNA was verified through PCR on RNA with the primers for the reference gene (*OsACT* Lee *et al.*, 2010). Total RNA was used for each sample to synthesize the cDNA, according to the SuperScript II Reverse Transcriptase (Invitrogen, ThermoFisher Scientific, Massachusetts, USA) procedure using random primers.

Primer selection

The genes analysed in this study (Table 1) were selected based on their regulation and role during the interaction between PGPR and rice. In particular, in addition to genes coding for a

metallothionein – like protein and a multiple stress-responsive zinc-finger protein (Brusamarello-Santos *et al.*, 2012), genes involved in ethylene and auxin pathways were considered (Song *et al.*, 2009; Brusamarello-Santos *et al.*, 2012; Vargas *et al.*, 2012). Before RT-qPCR, all primers (Table ST1) were tested *in silico* on PRIMERBLAST and in PCR reactions on genomic DNA extracted from Baldo rice. The DNA extraction was performed using the DNeasy Plant Mini Kit (Qiagen, Hilden, Germania) according to the manufacturer's instructions.

Gene expression analysis

Quantitative RT-PCR was carried out with 7500 Fast Real Time Systems (Applied Biosystem, ThermoFisher Scientific, Massachusetts, USA). Each PCR reaction was conducted on a total volume of 10 µl, containing 1 µl cDNA, 5 µl SYBR Green Reaction Mix and 0.3 µl of each primer (10 µM) using a 96-well plate. The used primers are listed in ST1. The following PCR programme, which includes the calculation of a melting curve, was used: 95°C for 10 min, 40 cycles of 95°C for 15 s, 60°C for 1 min, 95°C for 15 s, 60°C for 1 min, and 95°C for 15 s. All the reactions were performed for three biological and technical replicates. The baseline range and CT (cycle threshold) values were automatically calculated using the 7500 Fast Real Time Systems software. In order to compare data from different PCR runs or cDNA samples, the CT values of all the genes were normalized to the CT value of *OsACT1*, the reference gene. The candidate gene expression was normalized to that of the reference gene by subtracting the Ct value of the reference gene from the Ct value of the candidate gene efficiency correction, from the equation $2^{-\Delta\Delta CT}$ (Livak and Schmittgen, 2001), where ΔΔCT represents the ΔCT sample–ΔCT control.

Data analysis

The statistical analysis of the phenotyping data was done using JMP7 (SAS Institute Inc., Cary, NC, 1989-2019). Concerning gene expression, statistical analyses were carried out using REST 2009, version 2.0.13 (Qiagen, Hilden, Germany) (Pfaffl *et al.*, 2002), considering 0.05 as the *p*-value. Only significant expression values were considered and visualized as heat maps by a custom R (version 3.6.3) script (command "heatmap.2"). In order to reduce the data set dimension, a PCA (Principal Component Analysis) was carried out on ΔCT data for each biological replicate through R (version 3.6.3) (CRAN package "ggfortify"). The input files for R analysis were tabular file in .csv including ID# genes and ID# samples.

Results

Identification and characterization of P. chlororaphis ST9

It was of interest to isolate and characterize new *P. chlororaphis* strains from the soil in order to test their potential use as plant probiotics. Strain ST9 was isolated form bulk soil as described in the Materials and Methods section. The genome sequence of *P. chlororaphis* ST9 was determined and established that its chromosome was 6.7 Mb long with a GC content of 63%; according to the RASTtk annotation scheme it contains 6185 predicted protein-coding sequences (CDSs) and 83 RNAs. Phylogenetic analyses based on MSLA using 6 loci (16S rRNA, *recA*, *gyrB*, *rpoD*, *carA* and *atpD;* Hilario *et al.,* 2004-) and 16 sequences of *P. chlororaphis* isolates revealed that strain ST9 belongs to the *Pseudomonas chlororaphis* subsp. *aurantiaca* group (Fig. SF2). As other *P. chlororaphis* strains, its genome possesses several loci encoding for anti-bacterial/anti-fungal compounds including operons for the biosynthesis of phenazine, of two R-tailocins bacteriocines and of pyrrolnitrin. Other loci potentially involved in microbial pathogen antagonism, cell-cell signal interferences, niche colonization, plant protection and plant-beneficial traits were found and reported in Table 3. In summary, *P. chlororaphis* ST9 contains many of the features that can potentially make this strain a plant probiotic and biocontrol agent.

In order to begin assessing possible PGP potential of *Pseudomonas chlororaphis* ST9, several *in vitro* phenotypes were determined. A summary of the tested phenotypes is summarized in Table 2. Among the performed tests, the ability of ST9 to inhibit the growth of a wide variety of pathogens, including rice pathogens such as *Dickeya zeae*, *Pseudomonas fuscovaginae* and *Magnaporthe oryzae* was observed. Other PGP related *in vitro* phenotypes such as lipolytic and proteolytic activities and motility were observed. In summary, this isolate possesses several phenotypic abilities many of which can be of importance in plant colonization and PGP.

***P. chlororaphis* ST9 root colonization ability, persistence and effect on the microbiome in the rhizosphere**

It was of interest to evaluate the ability of *P. chlororaphis* ST9 to colonize and persist in the rhizosphere of rice plants. Rice seeds were inoculated with *P. chlororaphis* ST9, grown in greenhouse and roots were then collected at T1 (10 dpi), T2 (28 dpi) and T4 (90 dpi) together with un-inoculated controls for ST9 strain CFU counts. Results are summarized in Figure 1 which show that the level of colonization after 10 days was significantly high (10^7 cfu/g of root, -wet weight) then it decreased and stabilized around a value of 10^4 cfu/g at 90 dpi. Colonies grown on TSA rifampicin plates were randomly chosen for identification through 16S rRNA amplification and sequencing to confirm their identity as *P. chlororaphis* ST9. Rifampicin resistant colonies present in the untreated plants resulted to be a mixture of different bacteria: some of them were part of an analogue experiment carried out in concomitance (data not shown). These results evidenced that under the conditions tested, *P. chlororaphis* ST9 was an efficient root colonizer and able to persist

in the rhizosphere.

It was of interest to determine the possible effects of strain ST9 colonization on the rhizosphere microbiome. A 16S rDNA community profiling approach was performed, characterizing and comparing the microbial root community at 28 and 90 dpi in ST9 treated and untreated plants. The richness and diversity values of the bacterial communities after normalization are shown in Figure 2.

The *Pseudomonas* genus, was significantly more abundant (*p*-value<0.001) in the *P. chlororaphis* ST9 inoculated plants as presented in Figure 2A. The presence of *P. chlororaphis* ST9 was confirmed by finding its 16S V3-V4 region sequence among the reads obtained by the NGS experiment. Significant differences (*p*-value = 8.66e-05) in Shannon alpha diversity (Figure 2B) were found between inoculated and un-inoculated samples both at 28 and 90 dpi. Beta diversity analysis based on Bray Curtis distance (Figure 2C) was performed to compare the microbial compositions of the two different tested conditions. The results obtained highlighted differences between the microbial populations in relation to the treatment. As result of the *P. chlororaphis* treatment, several genera resulted differently distributed as evidenced by the heatmap of Figure 3 and in SF3 and SF4. Among the genera more abundant in untreated samples were *Duganella*, *Clostridium*, *Aeromonas, Enterobacter and Vogesella*; the majority of them are also defined as animal/human pathogens and just few species, such as *Clostridium puniceum* (Lund *et al*., 1981) and *Enterobacter cloacae* (García-González *et al*., 2018), are correlated with plant diseases. On the other hand, genera such as *Janthinobacterium*, known for its antifungal features (Haack *et al*., 2016), *Flavobacterium*, often reported as a plant growth promoting rhizobacterium (Kolton *et al.,* 2016), *Bacillus, Paenibacillus*, *Bradyrhizobium* and others known for their PGP potential (Sansinenea, 2019; Radhapriya *et al*., 2018; Greetatorn, *et al*., 2019; Ferreira *et al*., 2019; Berg, 2009; Souza *et al*., 2015) were significantly more abundant in the *P. chlororaphis* ST9 treated communities than in untreated ones.

Plant gene expression and phenotypic analysis of *P. chlororaphis* inoculated plants

In order to determine the effects of *P. chlororaphis* ST9 on plant growth, several phenotypic parameters were assayed. Statistical analyses were carried out on chlorophyll and flavonoid content and on the nitrogen balance index (NBI) as physiological parameters. In addition, plant height and dry shoot biomass at 90 dpi (after flowering stage) was also established. Results did not show any statistical differences between control and ST9 inoculated plants [p(F)<0.05] (Figure 4), although a tendency on higher NBI and lower flavonoid contents was measured in inoculated plants compared with the control. It was concluded that under the conditions tested, no significant plant beneficial effect was observed upon seed inoculation of *P. chlororaphis* ST9.

In order to investigate the effect on PGPR beneficial plant-bacteria interactions, RT-qPCR was performed on 14 genes selected for their role in ethylene and auxin pathways (Table 1). In Supplementary Table 2 (ST2) the fold change was shown for genes that were significantly and not

significantly differentially expressed, while in Figure 5 the heat map representation of the transcript levels coupled to a hierarchical clustering in ST9 inoculated plants is presented. At T2, significant differences in the expression of some loci was evidenced when compared to the other two time points. The most up- and down-regulated genes at T2 were *OsETR3* and *Osmetallotionein*, respectively. At T1, with the exception of *Osmetallotionein,* there were no significantly differentially regulated genes. At T3 the most up-regulated gene was *OsIAA14*, while *OsERF3* was the most down-regulated. The only genes that were never significantly differentially expressed were *OsIAA11* and *OsIAA4.* The PCA on ΔCT data of control (un-inoculated) and ST9 inoculated samples for each replicate and time point showed, without outlier observations, a clear distinction between control and inoculated samples at T2, where we have found the highest number of differently regulated genes (Figure 6). On the other hand, there was no separation between data at T1 and at T3, where we have found a reduced number of regulated genes. In summary, most loci displayed some level of variation in gene expression upon inoculation with strain ST9; one gene was regulated at T1, while 12 and five genes were regulated at T2 and T3, respectively.

Discussion

This study presents the identification of a *P. chlororaphis* strain and characterisation studies in relation to plant growth promotion (PGP). The work performed included root colonisation and persistence, effect on the root bacterial microbiome, effect on targeted plant gene expression and a greenhouse plant growth promotion test. Results showed that *P. chlororaphis* ST9 is a good rice root colonizer integrating well in the microbiome affecting plant gene expression however in our greenhouse experiment it did not result in a plant growth promotion effect.

P. chlororaphis was able to persist in the root compartment for 90 days without decreasing its abundance below a concentration of 10^4 cfu/g of root. An initial good root colonization rate then slowly decreasing in the later stages of plant growth has also been observed for other PGPR strains. For example in Chaudhary *et al*. (2013) *Azotobacter* strain ST24 colonizes well the wheat rhizosphere decreasing during growth and reaching a density of 10^4cfu/ml at 90 days after sowing. Similarly Solanky and Garg (2014) performed the inoculation of *Azotobacter* on *Brassica campestris* and again detected its persistence at 30 and 60 day post inoculation being approximately 2×10^4 cfu per plant at 60 dpi. Mosquito *et al*. (2020) inoculated a *Kosakonia* sp. strain in rice in the field and it was persistent at 30 dpi, while it was undetectable at 60 and 90 dpi. The effect of ST9 inoculation on the total bacterial population evidenced an increase in biodiversity with an enrichment of certain bacterial genera (e. g. *Janthinobacterium*, *Flavobacterium*, *Bacillus, Paenibacillus*, *Bradyrhizobium*) which are known to establish beneficial association with plants (Berg, 2009; Souza *et al*., 2015; Radhapriya *et al*., 2018; Ferreira *et al*., 2019; Greetatorn, *et al*., 2019; Sansinenea, 2019).

No plant beneficial effect on the rice inoculated with strain ST9 was observed. The reason is not known regardless that the strain was present and persistent in the rhizosphere. It would be appropriate to challenge rice with biotic and abiotic stresses and determine whether the presence of strain ST9 would have resulted in a tolerance effect. On the bases of physiological parameters however ST9 inoculated plants displayed less stress symptoms than controls. A similar trend was also observed Andreozzi *et al*. (2019) where NBI was significantly lower in the uninoculated control with respect to the *Herbaspirillum huttiense* RCA24 + *Enterobacter cloacae* RCA25 inoculated Baldo rice plants.

Gene expression studies were performed with 14 rice loci related to ethylene and auxin pathways together with genes coding for a metallothionein–like protein and a multiple stress-responsive zinc-finger protein; a role for these genes during rice-PGPR interaction has been demonstrated (Song *et al*., 2009; Brusamarello-Santos *et al*., 2012; Vargas *et al*., 2012; Ambreetha *et al*., 2018). Genes *OsERS1, OsERS2, OsETR2, OsETR3*, homologs to *Arabidopsis thaliana* ethylene receptors, were transcribed at higher level in the *P. chlororaphis* ST9 inoculated plants. Similarly, Vargas *et al*

(2012) also observed increase in expression of these loci when inoculated with *Azospirillum brasilense* and *Burkholderia kururiensis.* In their study rice varieties having more BNF (Biological Nitrogen Fixation) capacities showed higher bacterial colonization as well as up-regulation of ethylene receptors genes. In this study, these loci were significantly up- regulated at 28 dpi indicating that the Baldo rice variety displays good nitrogen fixing capacity as also reported by Andreozzi *et al.* (2019). The *OsERF2* and *OsERF3* genes, encoding for transcriptional factors related to ethylene, were also differently expressed in ST9 inoculated plants indicating that this strain could be involved in ethylene hormone levels. Indole-3-acetic acid (IAA) pathway modulation is important during an effective plant colonization, both by pathogenic and by nonpathogenic organisms (Brusamarello-Santos *et al.*, 2012). In *P. chlororaphis* inoculated plants it was established that loci *OsIAA1, OsIAA13* and *OsIAA14* were up-regulated at T2, *OsIAA1* was down-regulated at T3, while *OsIAA11* and *OsIAA4* were not differently transcribed in the three considered time points. Some of these trends are in accordance and others in contrast with previous observations (Brusamarello-Santos *et al.*, 2012; Ambreetha *et al.*, 2018); this could be due to the different timing and experimental applied conditions. Expression of genes associated to defense can be also affected by bacterial presence in the rhizosphere (Brusamarello-Santos *et al.,* 2012). *OsISAP1*, encoding for a multiple stress-responsive zinc-finger protein, was up-regulated at T2, suggesting an activation of the plant defense mechanisms. The gene coding for a metallothionein, which is a metal binding protein involved in metal homeostasis, was always down-regulated at each time point in agreement with previous data (Brusamarello-Santos *et al.,* 2012). The plant defense system could be primed by *P. chlororaphis* ST9 maintaining a low level of stress; future studies on the response to biotic stress of strain ST9 inoculated plant will determine whether the change of expression of these loci provides immunity to microbial pathogens.

Inoculated PGPR must interact or compete with other microorganisms in the rhizosphere microbiome and this can cause a short persistence of the inoculated bacteria possibly affecting its probiotic effects (Backer *et al.*, 2018). The ability of *P. chlororaphis* ST9 to successfully colonize and persist in the rhizosphere is an important trait for its possible use as bioinoculant for agriculture. Furthermore, the plethora of antimicrobial activities encoded in its genome makes *P. chlororaphis* ST9 a good candidate for biotic stress tolerance tests upon its inoculation. Plant gene expression analysis demonstrated some positive effects of the presence of strain ST9 on the expression of plant hormonal pathways such as IAA and ethylene; further investigation should be carried out on rice defence genes in order to better clarify the type of interaction. *P. chlororaphis* strains are considered safe for the environment and human health (EPA, 2009) and their use in agriculture is has been permitted through the application of live microorganism formulations (Rosas 2017; Anderson *et al.*, 2018) and via the production and purification of metabolites (Liu *et al.*, 2016; Peng *et al.*, 2018). Future experiments with *P. chlororaphis* ST9 will reveal its full potential as a

bioinoculant for PGP, tolerance to abiotic and most importantly, biotic stresses.

Reference

Ahemad M. (2015). Phosphate-solubilizing bacteria-assisted phytoremediation of metalliferous soils: a review. 3 Biotech 5:111-121

Ambreetha S, Chinnadurai C, Marimuthu P, Balachandar D. (2018). Plant-associated *Bacillus* modulates the expression of auxin-responsive genes of rice and modifies the root architecture. Rhizosphere 5:57-66.

Anderson AJ, Kim YC. (2020). Insights into plant-beneficial traits of probiotic *Pseudomonas chlororaphis* isolates. J Med Microbiol 69:361-371

Anderson JA, Staley J, Challender M, Heuton J. (2018). Safety of *Pseudomonas chlororaphis* as a gene source for genetically modified crops. Transgenic Res 27:103-113.

Andreozzi A, Prieto P, Mercado-Blanco J, Monaco S, Zampieri E, Romano S, Valè G, Defez R, Bianco C. (2019). Efficient colonization of the endophytes *Herbaspirillum huttiense* RCA24 and *Enterobacter cloacae* RCA25 influences the physiological parameters of *Oryza sativa* L. cv. Baldo rice. Environ Microbiol doi:10.1111/1462-2920.14688.

Arrebola E, Cazorla FM. (2020). Aer Receptors Influence the *Pseudomonas chlororaphis* PCL1606 Lifestyle. Front Microbiol 11:1560.

Arrebola E, Tienda S, Vida C, de Vicente A, Cazorla FM. (2019). Fitness Features Involved in the Biocontrol Interaction of *Pseudomonas chlororaphis* With Host Plants: The Case Study of PcPCL1606. Front Microbiol 10:719.

Backer R, Rokem JS, Ilangumaran G, Lamont J, Praslickova D, Ricci E, Subramanian S, Smith DL. (2018). Plant growth-promoting rhizobacteria: context, mechanisms of action, and roadmap to commercialization of biostimulants for sustainable agriculture. Front Plant Sci 9:1473.

Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Prjibelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. (2012). SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. J Comput Biol 19:455-477.

Bender SF, Wagg C, Van Der Heijden MG. (2016). An underground revolution: biodiversity and soil ecological engineering for agricultural sustainability. Trends Ecol Evol 31:440-452.

Berendsen RL, Pieterse CMJ, Bakker P. (2012). The rhizosphere microbiome and plant health. Trends Plant Sci 17:478-486.

Berg G. (2009). Plant-microbe interactions promoting plant growth and health: perspectives for controlled use of microorganisms in agriculture. Appl Microbiol Biotechnol 84:11-18.

Better M, Lewis B, Corbin D, Ditta G, Helinski DR. (1983). Structural relationships among *Rhizobium meliloti* symbiotic promoters. Cell 35:479-485.

Biessy A, Novinscak A, Blom J, Leger G, Thomashow LS, Cazorla FM, Josic D, Filion M. (2019). Diversity of phytobeneficial traits revealed by whole-genome analysis of worldwide-isolated phenazine-producing Pseudomonas spp. Environ Microbiol 21:437-455.

Brettin T, Davis JJ, Disz T, Edwards RA, Gerdes S, Olsen GJ, Olson R, Overbeek R, Parrello B, Pusch GD, Shukla M, Thomason JA 3rd, Stevens R, Vonstein V, Wattam AR, Xia F. (2015). RASTtk: a modular and extensible implementation of the RAST algorithm for building custom annotation pipelines and annotating batches of genomes. Sci Rep 5:8365.

Bric JM, Bostock RM, Silverstone SE. (1991). Rapid *in situ* assay for indoleacetic acid production by bacteria immobilized on a nitrocellulose membrane. Appl Environ Microbiol 57:535-538.

Brusamarello-Santos LCC, Pacheco F, Aljanabi SMM, Monteiro RA, Cruz LM, Baura VA, Pedrosa FO, Souza EM, Wassem R. (2012). Differential gene expression of rice roots inoculated with the diazotroph Herbaspirillum seropedicae. Plant Soil 356:113.

Bulgarelli D, Schlaeppi K, Spaepen S, Ver Loren van Themaat E, Schulze-Lefert P. (2013). Structure and functions of the bacterial microbiota of plants. Annu Rev Plant Biol 64:807-838.

Burr SE, Gobeli S, Kuhnert P, Goldschmidt-Clermont E, Frey J. (2010). *Pseudomonas chlororaphis* subsp. *piscium* subsp. *nov.*, isolated from freshwater fish. Int J Syst Evol Microbiol 60:2753-2757.

Calderón CE, de Vicente A, Cazorla FM. (2014). Role of 2-hexyl, 5-propyl resorcinol production by *Pseudomonas chlororaphis* PCL1606 in the multitrophic interaction the avocado rhizosphere during the biocontrol process. FEMS Microb. Ecol 89:20-31.

Calderon CE, Perez-Garcia A, de Vicente A., Cazorla FM. (2013). The *dar* genes of *Pseudomonas chlororaphis* PCL1606 are crucial for biocontrol activity via production of the antifungal compound 2-hexyl, 5-propyl resorcinol. Mol Plant Microbe Interact 26:554-565.

Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP. (2016). DADA2: high resolution sample inference from Illumina amplicon data. Nat Methods 13:581.

Cazorla FM, Duckett SD, Bergström ET, Odijk R, Lugtenberg BJJ, Thomas-Oates JE, Bloemberg GV. (2006). Biocontrol of avocado dematophora root rot by antagonistic *Pseudomonas fluorescens* PCL1606 correlates with the production of 2-hexyl, 5-propyl resorcinol. Mol Plant Microbe Interact 19:418-428.

Chaudhary D, Narula N, Sindhu SS, Behl RK. (2013). Plant growth stimulation of wheat (*Triticum aestivum* L.) by inoculation of salinity tolerant *Azotobacter* strains. Physiol Mol Biol Plants 4:515-519.

Chen Y, Shen X, Peng H, Hu H, Wang W, Zhang X. (2015). Comparative genomic analysis and phenazine production of *Pseudomonas chlororaphis*, a plant growth-promoting rhizobacterium. Genomics Data 4:33-42.

Chin-A- PA, Woeng TF, Bloemberg GV, van der Bij AJ, van der Drift KM, Schripsema J, Kroon B, Scheffer RJ, Keel C, Bakker Tichy HV. (1998). Biocontrol by phenazine-1-carboxamide-producing *Pseudomonas chlororaphis* PCL1391 of tomato root rot caused by *Fusarium oxysporum* f. sp. *radicis-lycopersici*. Mol Plant–Microbe Interact 11:1069-1077.

Cho S-T, Chang H-H, Egamberdieva D, Kamilova F, Lugtenberg B, Kuo C-H (2015) Genome Analysis of *Pseudomonas fluorescens* PCL1751: A Rhizobacterium that Controls Root Diseases and Alleviates Salt Stress for Its Plant Host. PLoS ONE 10: e0140231.

Compant S, Samad A, Faist H, Sessitsch A. (2019). A review on the plant microbiome: Ecology, functions, and emerging trends in microbial application. J Adv Res. 19:29-37.

Dorosky RJ, Yu JM, Pierson LS, III, Pierson EA. (2017). *Pseudomonas chlororaphis* produces two distinct R-tailocins that contribute to bacterial competition in biofilms and on roots. Appl Environ Microbiol 83:e00706-17.

Flury P, Aellen N, Ruffner B, Péchy-Tarr M, Fataar S, Metla Z, Dominguez-Ferreras A, Bloemberg G, Frey J, Goesmann A, Raaijmakers JM, Duffy B, Höfte M, Blom J, Smits TH, Keel C, Maurhofer M. (2016). Insect pathogenicity in plant-beneficial pseudomonads: phylogenetic distribution and comparative genomics. ISME J 10:2527-2542.

Ferreira CMH, Soares HMVM, Soares EV. (2019). Promising bacterial genera for agricultural practices: An insight on plant growth-promoting properties and microbial safety aspects. Sci Total Environ 682:779-799.

Flury P, Aellen N, Ruffner B, Péchy-Tarr M, Fataar S, Metla Z, Dominguez-Ferreras A, Bloemberg G, Frey J, Goesmann A, Raaijmakers JM, Duffy B, Höfte M, Blom J, Smits TH, Keel C, Maurhofer M. (2016). Insect pathogenicity in plant-beneficial pseudomonads: phylogenetic distribution and comparative genomics. ISME J. 10:2527-2542.

García-González T, Sáenz-Hidalgo HK, Silva-Rojas HV, Morales-Nieto C, Vancheva T, Koebnik R, Ávila-Quezada GD. (2018). *Enterobacter cloacae*, an Emerging Plant-Pathogenic Bacterium Affecting Chili Pepper Seedlings. Plant Pathol J 34:1-10.

Glick BR. (2014). Bacteria with ACC deaminase can promote plant growth and help to feed the world. Microbiol Res 169:30-39.

Goulas Y, Cerovic ZG, Cartelat A, Moya I. (2004). Dualex: a new instrument for field measurements of epidermal ultraviolet absorbance by chlorophyll fluorescence. Appl Opt 43:4488-4496.

Greetatorn T, Hashimoto S, Sarapat S, Tittabutr P, Boonkerd N, Uchiumi T, Teaumroong N. (2019). Empowering rice seedling growth by endophytic *Bradyrhizobium* sp. SUTN9-2. Lett Appl Microbiol 68:258-266.

Haack FS, Poehlein A, Kröger C, Voigt CA, Piepenbring M, Bode HB, Daniel R, Schäfer W, Streit WR. (2016). Molecular Keys to the *Janthinobacterium* and *Duganella* spp. Interaction with the Plant Pathogen *Fusarium graminearum*. Front Microbiol 7:1668.

Hayat R, Ali S, Amara U, Khalid R, Ahmed I. (2010). Soil beneficial bacteria and their role in plant growth promotion: a review. Ann Microbiol 60:579-598.

Hazra KK, Swain DK, Bohra A, Singh SS, Kumar N, Nath CP. (2018). Organic rice: potential production strategies, challenges and prospects. Organic Agriculture. 8:39-56.

Hilario E, Buckley TR, Young JM. (2004). Improved resolution on the phylogenetic relationships among *Pseudomonas* by the combined analysis of *atp D*, *car A*, *rec A* and 16S rDNA. Antonie van Leeuwenhoek 86:51-64.

Hong YW, Glick B R, Pasternak JJ. (1991). Plant microbial interaction under gnotobiotic conditions - a scanning electron-microscop estudy. Curr Microbiol 23:111-114.

Hu W, Gao Q, Hamada MS, Dawood DH, Zheng J, Chen Y, Ma Z. (2014). Potential of *Pseudomonas chlororaphis* subsp. *aurantiaca* Strain Pcho10 as a Biocontrol Agent Against *Fusarium graminearum*. Phytopathol 104:1289-1297.

Huber B, Riedel K, Hentzer M, Heydorn A, Gotschlich A, Givskov M, Molin S, Eberl L. (2001). The cep quorum-sensing system of *Burkholderia cepacia* H111 controls biofilm formation and swarming motility. Microbiol 147:2517-2528.

Kang BR, Yang KY, Cho BH, Han TH, Kim IS, Lee MC, Anderson AJ, Kim YC. (2006). Production of indole-3-acetic acid in the plant-beneficial strain *Pseudomonas chlororaphis* O6 is negatively regulated by the global sensor kinase GacS. Curr Microbiol 52:473-476.

Khatoon Z, Huang S, Rafique M, Fakhar A, Kamran MA, Santoyo G. (2020). Unlocking the potential of plant growth-promoting rhizobacteria on soil health and the sustainability of agricultural systems. J Environ Manage 273:111118.

Kohler T, Curty LK, Barja F, van Delden C, Pechere JC. (2000). Swarming of *Pseudomonas aeruginosa* is dependent on cell-to-cell signaling and requires flagella and pili. J Bacteriol 182:5990-5996.

Kolton M., Erlacher A., Berg G., Cytryn E. (2016). The *Flavobacterium* Genus in the Plant Holobiont: Ecological, Physiological, and Applicative Insights. In: Castro-Sowinski S. (eds) Microbial Models: From Environmental to Industrial Sustainability. Microorganisms for Sustainability, vol 1. Springer, Singapore.

Lane DJ. (1991). 16S/23S rRNA sequencing. In: Nucleic acid techniques in bacterial systematics. Stackebrandt E, Goodfellow M, eds., John Wiley and Sons, New York, NY, pp. 115-175.

Lee HB, Hong JP, Kim SB. (2010). First Report of Leaf Blight Caused by *Pantoea agglomerans* on Rice in Korea. Plant Dis 94:1372.

Lemoine F, Correia D, Lefort V, Doppelt-Azeroual O, Mareuil F, Cohen-Boulakia S, Gascuel O. (2019). NGPhylogeny.fr: new generation phylogenetic services for non-specialists. Nucleic Acids Res 47:W260-W265.

Liu H, He Y., Jiang H, Peng H, Huang X, Zhang X, Thomashow LS, Xu Y. (2007). Characterization of a phenazine-producing strain *Pseudomonas chlororaphis* GP72 with broad-spectrum antifungal activity from green pepper rhizosphere. Curr Microbiol 54:302-306.

Liu K, Hu H, Wang W, Zhang X. (2016). Genetic engineering of *Pseudomonas chlororaphis* GP72 for the enhanced production of 2-Hydroxyphenazine. Microb Cell Fact 15:131.

Livak KJ, Schmittgen TD. (2001). Analysis of relative gene expression data using real-time quantitative PCR and the 2(–Delta Delta C(T)) Method. Methods 25:402-408.

Love MI, Huber W, Anders S. (2014). Moderated estimation of fold change and dispersion for RNA seq data with DESeq2. Genome Biol 15:550.

Lund BM, Brocklehurst TF, Wyatt GM. (1981). Characterization of Strains of *Clostridium puniceum* sp. nov., a Pink-pigmented, Pectolytic Bacterium. J Gen Microbiol 122:17-26.

McDonald D, Price MN, Goodrich J, Nawrocki EP, DeSantis TZ, Probst A, Andersen GL, Knight R, Hugenholtz P. (2012). An improved Greengenes taxonomy with explicit ranks for ecological and evolutionary analyses of bacteria and archaea. ISME J 6:610-618.

McMurdie PJ, Holmes S. (2013). Phyloseq: an R package for reproducible interactive analysis and graphics of microbiome census data. PloS ONE 8**:**e61217.

Mehnaz S. (2016). "An overview of globally available bioformulations," in Bioformulations: For Sustainable Agriculture, eds NK Arora, S Mehnaz, R. Balestrini (Berlin: Springer), 267-281.

Mitter B, Pfaffenbichler N, Flavell R, Compant S, Antonielli L, Petric A, Berninger T, Naveed M, Sheibani-Tezerji R, von Maltzahn G, Sessitsch A. (2017). A New Approach to Modify Plant Microbiomes and Traits by Introducing Beneficial Bacteria at Flowering into Progeny Seeds. Front Microbiol *8*:11.

Mosquito S, Bertani I, Licastro D, Compant S, Myers MP, Hinarejos E, Levy A, Venturi V. (2020). In Planta Colonization and Role of T6SS in Two Rice Kosakonia Endophytes. Mol Plant Microb Interact 33:349-363.

Nam HS, Anderson AJ, Kim YC. (2018). Biocontrol Efficacy of Formulated *Pseudomonas chlororaphis* O6 against Plant Diseases and Root-Knot Nematodes. Plant Pathol J 34:241-249.

Nandi M, Selin C, Brassinga AKC, Belmonte MF, Fernando WGD, Loewen PC, de Kievit TR. (2015). Pyrronitrin and hydrogen cyanide production by *Pseudomonas chlororaphis* PA23 exhibits nematicidal and repellent activity against *Caenorhabditis elegans*. PLoS ONE 10:e0123184.

Nautiyal CS. (1999). An efficient microbiological growth medium for screening phosphate solubilizing microorganisms. FEMS Microbiol Lett 170:265-270.

Oh SA, Kim JS, Park JY, Han SH, Dimkpa C, Anderson AJ, Kim YC. (2013). The RpoS Sigma Factor Negatively Regulates Production of IAA and Siderophore in a Biocontrol Rhizobacterium, *Pseudomonas chlororaphis* O6. Plant Pathol J. 29:323-329.

Oksanen J, Blanchet FG, Kindt R, Legendre P, Minchin PR, O'hara R, Simpson GL, Solymos P, Stevens MHH, Wagner H. (2013). Package 'vegan'. Community ecology package, version, 2:1-295.

Oleńska E, Małek W, Wójcik M, Swiecicka I, Thijs S, Vangronsveld J. (2020). Beneficial features of plant growth-promoting rhizobacteria for improving plant growth and health in challenging conditions: A methodical review. Sci Total Environ 743:140682.

Orozco-Mosqueda MDC, Rocha-Granados MDC, Glick BR, Santoyo G. (2018). Microbiome engineering to improve biocontrol and plant growth-promoting mechanisms. Microbiol Res 208:25-31.

Overbeek R, Olson R, Pusch GD, Olsen GJ, Davis JJ, Disz T, Edwards RA, Gerdes S, Parrello B, Shukla M, Vonstein V, Wattam AR, Xia F, Stevens R. (2014). The SEED and the Rapid Annotation of microbial genomes using Subsystems Technology (RAST). Nucleic Acids Res 42:D206–D214.

Peix A, Valverde A, Rivas R, Igual JM, Ramírez-Bahena MH, Mateos PF, Santa-Regina I, Rodríguez-Barrueco C, Martínez-Molina E, Velázquez E. (2007). Reclassification of *Pseudomonas aurantiaca* as a synonym of *Pseudomonas chlororaphis* and proposal of three subspecies, *P. chlororaphis* subsp. *chlororaphis* subsp. *nov.*, *P. chlororaphis* subsp. *aureofaciens* subsp. *nov.*, comb. *nov.* and *P. chlororaphis* subsp. *aurantiaca* subsp. *nov.*, comb. *nov.* Int J Syst Evol Microbiol 57:1286-1290.

Peng H, Zhang P, Bilal M, Wang W, Hu H, Zhang X. (2018). Enhanced biosynthesis of phenazine-1-carboxamide by engineered *Pseudomonas chlororaphis* HT66. Microb Cell Fact 17:117.

Penrose DM, Glick BR. (2003). Methods for isolating and characterizing ACC deaminase-containing plant growth-promoting rhizobacteria. Physiol Plant 118:10-15.

Pfaffl MW, Horgan GW, Dempfle L. (2002). Relative expression software tool (REST[©]) for group-wise comparison and statistical analysis of relative expression results in real-time PCR. Nucleic Acid Res 30:e36.

Pii Y, Mimmo T, Tomasi N, Terzano R, Cesco S, Crecchio C. (2015). Microbial interactions in the rhizosphere: beneficial influences of plant growth promoting rhizobacteria on nutrient acquisition process. A review. Biol Fertil Soils 51:403-415.

Radhapriya P, Ramachandran A, Palani P. (2018). Indigenous plant growth-promoting bacteria enhance plant growth, biomass, and nutrient uptake in degraded forest plants. 3 Biotech 8:154.

R Core Team (2014). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. http://www.R-project.org/.

Rosas SB. (2017). *Pseudomonas chlororaphis* subsp. *aurantiaca* SR1: isolated from rhizosphere and its return as inoculant. A review. Int Biol Rev 1:1-19.

Ruzzi M, Aroca R. (2015). Plant growth-promoting rhizobacteria act as biostimulants in horticulture. Sci Hortic 196:124-134.

Sansinenea E. (2019) *Bacillus* spp.: As Plant Growth-Promoting Bacteria. In: Singh H, Keswani C, Reddy M, Sansinenea E, García-Estrada C. (eds) Secondary Metabolites of Plant Growth Promoting Rhizomicroorganisms. Springer, Singapore.

Smeltzer MS, Hart ME, Iandolo JJ. (1992). Quantitative spectrophotometric assay for staphylococcal lipase. Appl Environ Microbiol 58:2815-2819.

Solanki M, Garg FC. (2014). The use of *lacZ* marker in enumeration of *Azotobacter chroococcum* in carrier based inoculants. Braz J Microbiol 45:595-601.

Song Y, Wang L, Xiong L. (2009). Comprehensive expression profiling analysis of *OsIAA* gene family in developmental processes and in response to phytohormone and stress treatments. Planta 229:577-591.

Souza RD, Ambrosini A, Passaglia LM. (2015). Plant growth-promoting bacteria as inoculants in agricultural soils. Genetics Mol Biol 38:401-419.

Spaepen S, Bossuyt S, Engelen K, Marchal K, Vanderleyden J. (2014). Phenotypical and molecular responses of *Arabidopsis thaliana* roots as a result of inoculation with the auxin-producing bacterium *Azospirillum brasilense*. New Phytol 201:850-861.

Steindler L, Venturi V. (2007). Detection of quorum-sensing N-acyl homoserine lactone signal molecules by bacterial biosensors. FEMS Microbiol Lett 266:1-9.

Tagele SB, Lee HG, Kim SW, Lee YS. (2019). Phenazine and 1-Undecene Producing **Pseudomonas chlororaphis** subsp. **aurantiaca** Strain KNU17Pc1 for Growth Promotion and Disease Suppression in Korean Maize Cultivars**. J Microbiol Biotechnol 29:66-78.

Tieman DM, Taylor MG, Ciadi JA, Klee HJ. (2000). The tomato ethylene receptors NR and LeETR4 are negative regulators of ethylene response and exhibit functional compensation within a multigene family. Proc Nat Acad Sci 97:5663-5668.

Tremblay N, Wang Z, Cerovic ZG. (2012). Sensing crop nitrogen status with fluorescence indicators. A review. Agron Sustain Dev 32:451-464

Vargas L, de Carvalho TLG, Ferreira PCG, Divan Baldani VL, Baldani JI, Silva Hemerly A. (2012). Early responses of rice (*Oryza sativa* L.) seedlings to inoculation with beneficial diazotrophic bacteria are dependent on plant and bacterial genotypes. Plant Soil 356:127.

Weisburg WG, Barns SM, Pelletier DA, Lane DJ. (1991). 16S ribosomal DNA amplification for phylogenetic study. J Bacteriol 173:697-703.

Zhang L, Khabbaz SE, Wang A, Li H, Abbasi PA. (2015). Detection and characterization of broad-spectrum antipathogen activity of novel rhizobacterial isolates and suppression of *Fusarium* crown and root rot disease of tomato. J Appl Microniol 118:685-703.

Zhao LF, Xu YJ, Ma ZQ, Deng ZS, Shan CJ, Wei GH. (2013). Colonization and plant growth promoting characterization of endophytic *Pseudomonas chlororaphis* strain Zong1 isolated from *Sophora alopecuroides* root nodules. Braz J Microbiol. 44:623-631.

Zhu X, van Pee KH, Naismith JH. (2010). The Ternary Complex of PrnB (the Second Enzyme in the Pyrrolnitrin Biosynthesis Pathway), Tryptophan, and Cyanide Yields New Mechanistic Insights into the Indolamine Dioxygenase Superfamily. J Biol Chem 285:21126-21133

Zlosnik JE, Hird TJ, Fraenkel MC, Moreira LM, Henry DA, Speert DP. (2008). Differential mucoid exopolysaccharide production by members of the *Burkholderia cepacia* complex. J Clin Microbiol 46:1470-1473.

Table 1: List of the genes considered for plant gene expression

Gene name	Putative function	Reference
OsERS1	Ethylene response sensor 1	Vargas *et al.,* 2012
OsERS2	Ethylene response sensor 2	Vargas *et al.,* 2012
OsETR2	Ethylene responsive 2	Vargas *et al.,* 2012
OsETR3	Ethylene responsive 3	Vargas *et al.,* 2012
OsIAA1	auxin-responsive protein IAA1-like	Song *et al.,* 2009
OsIAA4	auxin-responsive protein IAA4	Song *et al.,* 2009
OsIAA11	auxin-responsive protein IAA11	Song *et al.,* 2009
OsIAA13	auxin-responsive protein IAA13-like	Song *et al.,* 2009
OsIAA14	auxin-responsive protein IAA14-like	Song *et al.,* 2009
OsACT1	Actin 1	Lee *et al.,* 2010
OsARF2	Similar to auxin response factor 2	Brusamarello-Santos *et al.,* 2012
OsERF2	Similar to ethylene response factor 2	Brusamarello-Santos *et al.,* 2012
OsERF3	Similar to ethylene response binding factor 3	Brusamarello-Santos et al., 2012
OsISAP1	Multiple stress-responsive zinc-finger protein	Brusamarello-Santos *et al.,* 2012
Osmetallo-tionein	metallothionein-like protein type 1	Brusamarello-Santos *et al.,* 2012

Table 2. Phenotypic characterization of PGP features of *P. chlororaphis* ST9

	In vitro PGP activity							Antagonistic activity					*In vivo* PGP activity		
	Protease	Lipase	IAA production	P solubilization	ACC deaminase	EPS production	Motility	*Dickeya zeae*	*Pseudomonas fuscovaginae*	*Aspergillus sp*	*Fusarium sp*	*Magnaporthe grisea*	Emergence 4 dpi	Coleoptile length 10 dpi	Dry weight biomass 12 dpi
P. chlororaphis ST9	+	+	-	+	-	-	+	+	+	+	+	+	-	+*	-

* Non parametric Mann Whitney t test; *p*-value <0.0001

Table 3. PGP and protection features in the genome of *P. chlororaphis*

SYSTEM / COMPOUND	TARGET / BENEFICIAL EFFECT	REFERENCES	ST9 GENES
ANTAGONISM			
Phenazine 1-carboxylic acid (PCA)	Antifungal redox-active antibiotic	Chin-A-Woeng et al., 1998	7 genes cluster fig|286.2086.peg.5005 to fig|286.2086.peg.5011 phzABCDEFG
2-hydroxy phenazine	Fungistatic and bacteriostatic	Liu et al., 2016	fig|286.2086.peg.5012; phzO
Tailocine R	Antibacterial Persistence within the rhizosphere microbiome	Dorosky R et al 2018	40 genes cluster fig|286.2086.peg.1042- fig|286.2086.peg.1082
Pirrolnitrin	Antifungal compound	Nandi et al., 2015 Zhu X et al, 2010	4 genes cluster fig|286.2086.peg.3071 to fig|286.2086.peg.3074; prnA, prnB, prnC, prnD
2-hexyl, 5-propyl resorcinol (HRP)	Antifungal compound and signal molecule	Cazorla et al., 2006 Calderon C et al 2013	5 genes cluster fig|286.2086.peg.4110 to fig|286.2086.peg.4114; darA, darB, darC, darR, darS
Hydrogen cyanide	Metalloenzymes inhibitor and antifungal	Nandi et al., 2015	2 genes cluster fig|286.2086.peg.2368 to fig|286.2086.peg.2369
SIGNAL INTERFERENCES			
Acyl-homoserine lactone acylase	Quorum Quenching		fig|286.2086.peg.2769
N-acyl-L-homoserine lactone synthetase	Quorum sensing	Shah N. et al 2020	3 systems fig|286.2086.peg.5003 to fig|286.2086.peg.5004 phzI, phzR fig|286.2086.peg.4930 and fig|286.2086.peg.4931 aurI, aurR fig|286.2086.peg.2434 to fig|286.2086.peg.2435 luxI, luxR
NICHE COLONIZATION			
poly-beta-1,6-N-acetyl-D-glucosamine	biofilm adhesin	Cho S-T et al 2015	4 genes cluster fig|286.2086.peg.20 - fig|286.2086.peg.24 pgaA, pgaB, pgaC, pgaD
Motility & chemotaxis	flagella	Cho S-T et al 2015	6 genes cluster fig|286.2086.peg.1427 to fig|286.2086.peg.1433 flgF, flgG, flgH, flgI, flgK, flgL 36 gene cluster fig|286.2086.peg.1451 to fig|286.2086.peg.1486 9 gene cluster fig|286.2086.peg.4398 to fig|286.2086.peg.4406
PLANT PROTECTION			
Chitinase	insecticidal	Flury P et al 2016	5 genes fig|286.2086.peg.1868 and fig|286.2086.peg.1869 fig|286.2086.peg.3224 and fig|286.2086.peg.3225 fig|286.2086.peg.1868 4987 chitin binding protein and chitinases
Proteases	insecticidal	Flury P et al 2016	fig|286.2086.peg.3194 aprX/prtA and other related genes
Lipases	insecticidal	Flury P et al 2016	fig|286.2086.peg.3118 phospholipase C and others related genes
rebB	insecticidal	Flury P et al 2016	fig|286.2086.peg.35 to fig|286.2086.peg.36 rebB like protein (Refractile Inclusion Bodies)
BENEFICIAL ACTIVITIES TO PLANT			
Phosphate solubilization	Phosphorus solubilization enzyme		fig|286.2086.peg.702 alkD
Achromobactine	Fe chelation and competition		13 genes cluster From fig|286.2086.peg.3101 to fig|286.2086.peg.3113 Achromobactin synthesis, receptor, transport

Figure 1. Rice root colonization by *P. chlororaphis* ST9 was evaluated at three time points: 10, 28 and 90 dpi (days post inoculation). The average of the cfu/g of fresh weight root of 10 plants for each treatment and time point is presented. Each sample was plated on TSA to count all cultivable bacteria, and on TSArif to count just the rifampicin resistant CFUs, mainly *P. chlororaphis* ST9. Statistical analysis: same letter means no significant difference (t-test, non-parametric, Mann Whitney, *p*-value < 0.05).

A

B

C

Figure 2. Microbiome analysis. (A) *Pseudomonas* genus abundance according to the plant treatment and time point; the Welch two Sample t-test was used to compare untreated *versus* treated samples at 28 dpi (*p*-value = 0.0006793) and at 90 dpi (*p*-value = 0.0003417). (B) Alpha Diversity (Shannon index) at community level in accordance to the treatment and the time point; the Wilcoxon rank sum test was used, *p*-value for untreated *versus* treated samples both at 28 and 90 dpi results < 0.01. (C) Principal component analysis of the samples in accordance to the treatment and the time point, similarity between the different communities was evaluated using the Bray-Curtis test; the bacterial communities of the untreated plants cluster together and differently from the bacterial communities colonizing the ST9 treated samples (*p*-value < 0.001).

UT28: untreated samples at 28 dpi; T28: ST9 treated samples at 28 dpi; UT90: untreated samples at 90 dpi; T90: ST9 treated samples at 90 dpi.

Figure 3. Heat map showing the relative abundance (% of sequencing reads) of the 50 predominant genera. Rows are bacterial genera. Columns are samples. Colors indicate taxa with a higher (blue) or lower (light blue) relative abundance in each sample. Genera showing a different distribution and abundance level between the samples are highlighted with *: green if enriched in the treated samples, red if enriched in the untreated ones. UT28: untreated samples at 28 dpi; T28: ST9 treated samples at 28 dpi; UT90: untreated samples at 90 dpi; T90: ST9 treated samples at 90 dpi.

Figure 4: Total height (A), dry shoot biomass (B), chlorophyll (C) and flavonoid content (D), and NBI (E) in control un-inoculated and in ST9 inoculated plants.

Figure 5: Heat map representation of the transcript levels coupled to a hierarchical clustering in ST9 inoculated plants. Each column represents a time point, while each row represents a gene. Expression levels are coloured green for low intensities and red for high intensities (see scale at the top left corner). The black cells represent genes not significantly different from those of the untreated samples.

Figure 6: PCA carried out on ΔCT of each biological replicate of untreated control (K) and ST9 inoculated plants (S) at each time point (T1=10 dpi; T2=28 dpi; T3=40dpi). The cumulative variability percentage is given by the sum of 67.42% for PC1 and 26.31% for PC2, resulting in 93.73%. KT2 and ST2 are clearly separated in the space, differently by the other time points.