Drosophila glue protects from predation

Flora Borne[1], Stéphane R. Prigent[2], Mathieu Molet[3] and Virginie Courtier-Orgogozo[1]

[1]Institut Jacques Monod, CNRS, UMR7592, Université de Paris, 15 rue Hélène Brion, 75013, Paris, France

[2]Institut de Systématique, Évolution, Biodiversité, ISYEB, Muséum national d'Histoire naturelle, CNRS, Sorbonne Université, EPHE, Université des Antilles, 75005 Paris, France

[3]Sorbonne Université, UPEC, CNRS, IRD, INRA, Institute of Ecology and Environmental Sciences of Paris (iEES Paris), 75005 Paris, France

FB: flora.borne@ens-paris-saclay.fr ORCID: 0000-0002-6660-8559

SRP: srpfly@gmail.com ORCID: 0000-0002-8117-3848

MM: mathieu.molet@sorbonne-universite.fr ORCID: 0000-0002-1247-8904

VCO: virginie.courtier@normalesup.org, ORCID: 0000-0002-9297-9230

Abstract

Animals can be permanently attached to a substrate for several days, weeks or months in aerial environments at certain stages of their development such as eggs and pupae. Pupa adhesion has evolved multiple times in insects and is thought to maintain the animal in a place where it is not detectable by predators. Here, we investigate whether pupa adhesion in Drosophila could also protect the animal by preventing potential predators from detaching the pupa. We measured the adhesion of Drosophila species originating from the same area and found that pupa adhesion varies among species, which can be explained by different glue production strategies. Then, we compared attached and manually detached pupae in both field and laboratory assays to investigate the role of pupa adhesion to prevent predation. First, we found that attached pupae remain on site 30 % more than detached pupae in the field after three days, probably because they are less predated. Second, we observed that attached pupae are less efficiently predated by ants in the laboratory, because they are not carried back to the ant nest and because more ants are needed to consume them onsite. Our results show that pupa adhesion is a crucial mechanical trait for Drosophila fly survival that can prevent the animal from being taken away by predators.

Keywords

bioadhesion, bioadhesive, predation, insect, Drosophila, ant, pupa

Introduction

Multiple animals are immobile and permanently attached to a substrate. Most of them are found in aquatic environments (Wahl, 1997), and also within living hosts for parasites. In aerial environments, animals can also be attached to a substrate during certain stages of development where no feeding from outside nutrients is required. Eggs from multiple invertebrate species are fixed on leaves, wood or on the host tissue in some parasitic species (Fordyce and Nice, 2002; Hinton, 1981; Voigt and Gorb, 2010). This is also the case of pupae, a non-feeding life stage of holometabolous insects occurring between the larval and the adult stages, and undergoing a complete metamorphosis (Heming, 2003).

Modes of attachment and the ability to stick to the substrate appear to change rapidly during animal evolution. In stick and leaf insects gluing eggs to a substrate has evolved independently seven times (Robertson et al., 2018). Wasp parasitoids have evolved different pupation strategies consisting in hanging their cocoon to a leaf (*Meteorus pulchricomis*) or attaching their cocoon on a leaf (Microplitis sp, Harvey et al., 2011). In flies, pupae of certain species such as *Drosophila melanogaster* and *Phormia regina* are glued to a substrate whereas others such as *Musca domestica*, *Calliphora erythrocephala* or *Sarcophaga falculata* are not (Fraenkel and Brookes, 1953).

Permanent attachment of eggs and pupae has been associated with several functions. First, attachment can allow the organisms to remain in a favorable environment. Females of many butterfly species choose to lay and attach their eggs directly on the host plant on which their larvae will start feeding (Hinton, 1981; Wiklund, 1984). Butterflies laying their eggs during winter have evolved different strategies to avoid their eggs to be blown away if the dead host leaf falls far away from the plant. They lay their eggs near the host plant on lichens, small herbs or on tree trunks close by, or on the woody part of the host plant, or they glue their eggs less strongly to the host leaf so that the eggs would detach from the dead leaf and fall close to the host plant in case of strong wind (Fordyce & Nice, 2002; Hayes, 1982; Wiklund, 1984). Second, hanging chrysalis may facilitate adult emergence (Chapman, 1982). Third, attachment may protect immobile animals from predation. The defensive role of

adhesion towards predators can involve multiple aspects. Permanent attachment, when associated with clumping behaviors, in which individuals of a particular species group closely to one another, can confer a better protection from predators. For example, in a freshwater caddisfly, pupal grouping behavior with conspecifics confers protection against a planarian flatworm predator (Wrona and Dixon, 1991). Attachment can also prevent predators from accessing the immobile animal. In *Mecynogea lemniscata* spiders, females make a cocoon which is hanging from a leaf to protect their eggs. Cocoons are more attacked by predators such as ants when they are on the ground than when they are hanging (Hieber, 1992). Similarly, cocoons of the parasitoid *Meteorus pulchricornis* are less predated when they are hanging than when they are artificially attached to leaves (Shirai & Maeto, 2009). To our knowledge, nothing is known about the function of pupal attachment in Diptera. In this study, we investigated whether the glue attaching Drosophila pupae can protect them from predation.

Drosophila larvae produce a glue right before pupariation which allows the pupa to stay attached to a substrate during metamorphosis. After expectoration, the glue spreads between the body and the substrate and dries within a few minutes (Beňová-Liszeková et al., 2019). This glue is made of a few proteins called salivary gland secreted proteins (Sgs) which have evolved rapidly across and within species (Beckendorf and Kafatos, 1976; Da Lage et al., 2019; Korge, 1975; Korge, 1977).

In nature, pupae have been observed in various sites. In Drosophila, pupae are found on rotten fruits (Beltramí et al., 2012), on or below the soil surface (Ballman et al., 2017; Grossfield, 1978; Woltz and Lee, 2017) and even on beer glass bottles (Vouidibio et al., 1989). Aerial pupation sites are usually close to the ground, thus accessible to ground dwelling species. Multiple cases of predation on fruit fly pupae have been reported; the most common predators were ants, rove beetles and spiders (Hennessey, 1997; Thomas, 1995; Urbaneja et al., 2006). Birds and small mammals were also found to prey on fruit fly pupae (Bigler et al., 1986; Thomas, 1993). To our knowledge, all studies of pupa predation in Drosophila were performed on *D. suzukii* (Ballman et al., 2017; Gabarra et al., 2015; Woltz and Lee, 2017). In these analyses, ants and spiders were the most common predators of pupae. Ground beetles, earwigs and crickets were also identified as potential natural predators of pupae. As ants were previously observed to dig up and carry pupae out of the soil (Woltz and Lee, 2017), we hypothesized that pupa adhesion to a substrate might have another, yet

unexplored, effect against predation: preventing potential predators from detaching the
animal. Here, we first compared the adhesion strength of pupae from different drosophila
species from the same ecological community to explore whether different species have
evolved different attachment strategies. Then, we compared the ability of attached and
detached pupae of one of these species to remain on site in a natural environment. Finally, we
compared the ability of attached and detached pupae of two of these species to resist
predation in the laboratory using the most common natural predator found in the field, an ant
species.

Materials and Methods

Fly culture

105 Flies were cultured at 25 °C in plastic vials on standard medium [4 liters: 83.5 g yeast, 335.0
106 g cornmeal, 40.0 g agar, 233.5 g saccharose, 67.0 mL Moldex, 6.0 ml propionic acid]. For *D.*
107 *suzukii*, this medium was supplemented with 200 g of D-glucose.

Fly collection

Drosophila flies were collected at the Bois de Vincennes in Paris, France. On July 3 2020,
five traps made from 0.5-L plastic bottles were settled in the ornithological reserve
(48°50'05.5"N; 2°26'11.4"E). Small holes were made in the sides of the bottles that allowed
drosophilid flies to enter but prevented entry by larger insects. Traps were baited with pieces
of banana and hung to tree branches or within the understorey vegetation. They were
distantly placed, three of them in the forest, one at the margin of a meadow and another close
to a pond (Table S1). Flies in the bottles were collected two days and five days later. On July
16 2020, drosophilid flies were also collected with a sweeping net over a compost at the
forest services facilities. Collected flies were transferred with an aspirator into plastic vials
containing a piece of humid tissue paper for the time of transportation. In the laboratory, flies
were checked under a binocular stereomicroscope and isolated by species in culture vials
with standard cornmeal. When species could not be precisely determined, females were
isolated in small culture vials with instant Drosophila medium (Formula 4-24, Carolina
Biological Supply Company, Burlington, NC,USA) and species were then identified based on
key morphological characters in the male progeny. Results of the fly collections are presented
in Table S1. Combining isofemale lines of the same species when necessary, we managed to

obtain mass culture for most species, including *D. hydei*, *D. simulans* and *D. suzukii*. These

three stocks were raised in the laboratory for 3-4 months at 25°C before being used in the

experiments described above.

Adhesion assays

Third instar wandering larvae were transferred on glass slides (Menzel Superfrost microscope

glass slide from ThermoScientific™ #AGAB000080) with soft forceps, kept in a box with

wet cotton. 15 to 21 h after transfer, pupae naturally attached to the glass slides (with their

ventral part adhering to the glass slide) were used for the adhesion tests. The pull-off force

necessary to detach the pupa from the glass slide was measured using a universal test

machine (LS1S/H/230V Lloyd Instruments) with a 5N force sensor (YLC-0005-A1 Lloyd

Instruments), in a set up similar to the one we published earlier (Borne et al., 2020). Double-

sided adhesive tape (tesa, extra strong, #05681-00018) was attached to a cylindrical metal

part in contact with the force sensor. The force was set to 0 before each run. The force sensor

was moved down with a constant speed of 1 mm/min until it pressed the pupa with a force of

0.07 N (0.25 N for *D. hydei*) then let still at a pressure of 0.03 N (0.21 N for *D. hydei*) for 10s

and finally moved up with a constant speed of 0.2 mm/s until the pupa was detached. Force-

by-time curves were recorded using NEXYGENPlus software (Lloyd Instruments). We used

the maximal force reached during the experiment, corresponding to the force at which the

pupa was detached, as the adhesion strength of the individual. Pupae whose pupal case broke

during the assay (*D. suzukii*: 1/27, *D. simulans*: 2/37; *D. hydei*: 7/50) or pupae which were

not detached (*D. suzukii*: 0/27, *D. simulans*: 5/37, *D. hydei*: 12/50) were excluded from the

analysis. After pupa detachment, images of glue prints remaining on glass substrates were

taken with a Keyence VHX-2000 Z20 x20 or x100. Contours of prints areas were digitized

manually by the same person using imageJ (1.50d, java 1.8.0_212, 64-bit) (Schneider et al.,

2012). Pictures were anonymized for manual contour acquisition so that the digitizer did not

know the genotype. We measured the surface of the print corresponding to the pupa-substrate

interface as defined previously (Borne et al., 2020). Three prints for *D. suzukii* and one print

for *D. simulans* were not detectable on the slides and were not used in the analysis.

To assess pupal size, we used pupae that were raised in the same condition as for the

adhesion tests but that were not used for the tests. We imaged attached pupae from the dorsal

view and measured the area of the pupal case as described to measure the glue area.

Predation assay in the field

In the laboratory, *D. simulans* third instar wandering larvae were collected with soft forceps and let to pupate in Petri dishes (55-mm diameter) in a closed plastic box containing wet paper for 17 to 24 hr at room temperature. 15 larvae were put per dish. All Petri dishes were then brought to the field. For the condition 'attached pupae', a few pupae were removed in order to have exactly 10 pupae glued in the lid of the Petri dish. For the condition 'detached pupae', pupae were all detached from the dish with featherweight entomology forceps and exactly 10 detached pupae were kept in the lid. To distinguish between the two conditions ("non collées" versus "collées" in French), the letters 'NC' or 'C' were written on pieces of paper that were taped on the external surface of the lids to facilitate visualisation and counting of the pupae. The lids of the Petri dishes were put in the center of buckets previously installed in the ornithological reserve of Bois de Vincennes. These 40×35 cm buckets contain local soil, leaf litter and vegetation; they are pierced at the bottom for water draining and are semi-buried (10 cm deep) (Honorio et al., 2020). In total, 28 buckets were used and contained both one dish with 'attached pupae' and one dish with 'detached pupae' (56 Petri dishes in total). We randomly alternated the West / East orientation of the two conditions inside the buckets. Pupae were counted at 0h (September 8 2020, day 1 morning, at 11 am), 6h30 (day 1 afternoon), 22h30 (day 2 morning), 31h (day 2 afternoon), 47h (day 3 morning), and 54h (day 3 afternoon) after the start of the experiment without being touched. Animals present in the dishes at counting times were photographed and later identified based on the pictures (Table S3). Insects present in the dishes at the end of recording (54h) were collected and kept in 90% ethanol for identification.

Ant predation assay in the laboratory

Seven colonies of *T. nylanderi* were collected on September 17 2020 in Bois de Vincennes (48°50'20.0"N 2°26'57.2"E), brought to the lab and allowed to move into artificial nests consisting of two microscope glass slides separated by a 1-mm auto-adhesive plastic foam harboring three chambers, covered with a black plastic film to maintain darkness. Each artificial nest was placed in a foraging area consisting of a plastic box ($11.5 \times 11.5 \times 5.5$ cm) as described in Honorio et al (2020). Water was always available in a tube plugged with cotton. Colonies were fed frozen Drosophila and diluted honey and then they were starved for

10 days until the beginning of the experiment, on October 13 2020. Prior and during the
experiment, colonies were kept at room temperature on the bench and under indirect sunlight.
Each day at around 9h30 am, and for 6 days (between October 13 and October 21 2020), one
glass slide presenting two pupae was put into the foraging area of each colony, at about 10
cm from the entrances of the artificial nest. On each day, half of the colonies were given one
slide with 2 *D. simulans* pupae and the other half one slide with 2 *D. suzukii* pupae, and we
alternated species every day. We used the same two Drosophila lines as for our adhesion
assays described above.

The glass slides were prepared as follows. Six third instar wandering larvae were
transferred on glass slides (Menzel Superfrost microscope glass slide from
ThermoScientific™ #AGAB000080) placed in a Petri dish kept in a box containing wet
cotton and let to pupate for 14 to 19 hours. On each slide, only one pupa was kept attached,
the other ones were detached slightly with soft forceps and one detached pupa was left on the
slide. Pupae were about 1 cm apart from each other on the slide. Left / right positions of the
two pupae were randomly assigned per colony and changed every day. The initial locations
of the pupae were identified by a mark under the slide.

Ant foraging areas were checked by eye every 5 minutes for 3 hours after the glass
slide with the 2 pupae was added. The number of ants in contact with the pupae were
counted. If the pupa was brought to the nest, we noted the time when the pupa was present
inside the nest for the first time. If the pupa was not brought to the nest, we noted the time
when the pupa was fully consumed by the ants (no Drosophila body remnants visible by eye).
211

Quantification and Statistical Analysis
We used R v3.6.1 (R Core Team 2015) to conduct our statistical analyses. Adhesion forces
were not normally distributed and statistical differences in forces between species were tested
by Kruskal-Wallis tests followed by multiple pairwise Wilcoxon tests. To test whether
adhesion forces were correlated to pupa - substrate contact areas for each species, we
performed standardized major axis regressions using *sma* function from the rsmatr-package
in R (Warton et al., 2012). Slopes were compared among species. For predation assays in
Bois de Vincennes and in the laboratory, means were compared using Wilcoxon tests because
data were not normally distributed. We did not correct P-values for multiple testing, as
suggested by Nakagawa (2004).

Fig 1. Pupa adhesion varies between species originating from the same location . (A)

Adhesion strength of three Drosophila lines collected in Vincennes. Force indicates the force

required to detach a pupa naturally attached to a glass slide. Each dot corresponds to a single

pupa and n indicates the total number of pupae tested for each species. Ends of the boxes

define the first and third quartiles. The black horizontal line represents the median. The

vertical line on the top of the box extends to the largest value no further than 1.5 * IQR from

the upper hinge of the box. The vertical line on the bottom of the box extends to the smallest

value at most 1.5 * IQR of the hinge. (IQR: inter-quartile range is the distance between the

first and the third quartiles). Data beyond the end of these lines are "outlying" points. *

indicates significant differences between *D. suzukii* and *D. hydei*, *D. suzukii* and *D. simulans*

and *D. simulans* and *D. hydei* (Kruskal-Wallis chi-squared = 63.77, *df* = 2, *p* = 1e-14,

followed by all pairwise comparison Wilcoxon test, *p* <0.05). (B) Pupa adhesion is correlated

with the glue print area corresponding to pupa-substrate contact area for each species

(standardised major axis regressions: *D. suzukii*: R^2 = 0.41, *p* = 0.0009, *D. simulans*: R^2 =

0.55, *p* = 5e-06, *D. hydei*, R^2 = 0.17, *p* = 0.02). (C) Adhesion strength corrected by the pupa-

substrate contact area. Boxplots as in 1A. n.s. indicates not significant (*p* > 0.05).

Results

Glue adhesion strength varies between species from a same location

We collected drosophilid flies in the forest near Paris in June 2020 and noted the presence of

9 *Drosophila* species (Table S1). We established fly stocks of the most common species.

Three of them, *Drosophila simulans*, *D. hydei* and *D. suzukii*, were assayed for pupal

adhesion. We found that *D. simulans* detached at a median strength of 234.2 mN (Fig. 1A),

similar to what has been found previously for its sister species *D. melanogaster* (Borne et al.,

2020). We measured a lower adhesion for *D. suzukii* pupae with a median strength of 78.7

mN and higher adhesion for *D. hydei* with a median strength of 482.6 mN. Adhesion strength

was significatively different between the three species (Kruskal-Wallis chi-squared = 63.77,

df = 2, *p* = 1.4e-14, followed by all pairwise comparison Wilcoxon test, *p* < 0.001).

By examining the glue prints left by the pupae on glass slides after detachment, we

found that adhesion forces correlated with the surface of the glue print delimiting the contact

between the pupa and the glass slide for each species (*D. suzukii*: R^2 = 0.41, *p* = 0.0009, *D.*

simulans: R^2 = 0.55, *p* = 5e-06, *D. hydei*, R^2 = 0.17, *p* = 0.02). There was no difference in

slope among the three species (*p* = 0.2) and the common slope was about 491, meaning that

adhesion force increases by 491 mN for 1 mm^2 increase of the pupa - substrate contact area

for each species. After dividing the adhesion force by the surface of contact, we found a

difference in adhesion between *D. simulans* and *D. hydei* (Fig. 1C, Kruskal-Wallis *chi-*

squared = 9.61, *df* = 2, *p* = 0.008, followed by all pairwise comparison Wilcoxon test, *p* =

0.0008) but not between *D. simulans* and *D. suzukii* (*p* = 0.1) and *D. suzukii* and *D. hydei* (*p* =

0.6).

To test whether the production of the glue was related to the size of the pupa in each

species, we imaged attached pupae raised in the same condition as the pupae used for the

adhesion tests and measured the area of the pupal case. We found that the three species had

different pupal size (Kruskal-Wallis *chi-squared* = 73.876, *df* = 2, *p* < 2e-16, followed by all

pairwise comparison Wilcoxon test, *p* < 0.001) with *D. hydei* pupae presenting the biggest

size (4.63 mm^2), then *D. suzukii* (2.78 mm^2) and *D. simulans* (2.12 mm^2). The ratio of glue

print area over pupal case area was 0.21 for *D. simulans*, 0.30 for *D. hydei* and 0.06 for *D.*

suzukii, suggesting that *D. suzukii* pupae produce less glue relative to their size than *D.*

simulans and *D. hydei*.

Fig 2. Attached pupae are taken away less frequently than detached pupae in nature.

(A) Picture of the half-buried buckets installed in Bois de Vincennes. (B) Picture of two dishes placed in the center of a bucket and containing attached ("C") or detached ("NC") pupae. (C-E) Predators observed in the dishes during the experiments: a red mite spider (C), two *Temnothorax nylanderi* ants (D) and a cockroach (E). (F) Boxplot represents the number of pupae present in one dish at the counting time. Pupae were counted twice a day in the morning (AM) and in the afternoon (PM). Each dot represents the count for one dish. White boxes represent dishes with detached pupae and grey boxes dishes with attached pupae. Boxplots are defined as previously (Fig. 1A). * represents a significant difference between the number of remaining pupae between the attached and detached conditions ($p < 0.05$, Wilcoxon test). Scale bars: (B) 1 cm, (C-E) 1 mm.

Attached pupae are taken away less frequently than detached pupae in a semi-natural environment

To test whether glue attachment may protect pupae from predation in a semi-natural environment, we chose to use *D. simulans*, as we could obtain a large number of pupae from our fly strain. We compared the disappearance of pupae naturally attached to the plastic lid of Petri dishes with pupae mechanically detached from the lid. We placed two dishes containing respectively 10 attached and 10 detached pupae in 28 open buckets in Bois de Vincennes (two Petri dishes in each bucket) for 54 h and monitored the number of remaining pupae twice a day (Fig. 2A,B). At the end of the experiment, 10 pupae (median = 10) remained in the dish with attached pupae (all still attached) compared to 6-7 pupae (median = 6.5) in the dish with detached pupae. We found that attached pupae stayed significantly more in the Petri dishes than detached ones, with differences becoming significant from day 2 AM until the end of the experiment (Fig. 2F; paired Wilcoxon rank tests with continuity correction: day 1 PM: $V = 9$, $p = 0.2$; day 2 AM: $V = 89.5$, $p = 0.02$; day 2 PM: $V = 92.5$, $p = 0.01$; day 3 AM: $V = 91$, $p = 0.01$; day 3 PM: $V = 110$, $p = 0.005$). After three days, attached pupae remained on site 30 % more than detached pupae. During the countings, a few species were observed in the Petri dishes: *Temnothorax nylanderi* ants, red spider mites, cockroaches and woodlice (Table S5). Red spider mites were seen to fix themselves to both attached and detached pupae (Fig. 2C). *Temnothorax nylanderi* was the only species that was clearly seen consuming both attached and detached pupae in the dishes (Fig. 2D). Cockroaches were also observed, but only on the first day (Fig. 2E). Woodlice were also found in the dishes but never in contact with the pupae.

Fig 3. Attached pupae require more time and more ants to go away. (A) Number of ants in direct contact with the pupa over the duration of the experiment. Each cell represents one experiment with X_Y corresponding to the experiment with the colony X during the trial Y. Grey cells represent experiments using *D. suzukii* and white cells using *D. simulans*. Red lines represent the number of ants over time on attached pupa and blue lines on detached pupa. Vertical lines represent the time when the attached pupa (in red) or the detached pupa (in blue) arrives in the nest (dashed line) or is fully consumed outside the nest (full line) (B)

Time during which ants are in contact with attached pupa (grey box) and detached pupa (white box) outside the nest (until the pupa is completely consumed outside the nest or enters into the nest). Each dot represents one experiment. (C) Maximum number of ants observed in contact with detached pupa (white box) and attached pupa (grey box) during the duration of the experiment. Each dot represents one experiment. Boxplots are defined as previously (Fig. 1A). * represents significant differences ($p < 0.05$, Wilcoxon tests).

Attached pupae are predated less efficiently by ants

To understand further how predators may act when they encounter strongly attached, loosely attached or detached pupae, we decided to monitor in the laboratory pupae predation by the ant *Temnothorax nylanderi*, which was the most commonly found predator of *D. simulans* pupae in our field assay. Seven ant colonies were collected in Bois de Vincennes. After 10-day starvation, each ant colony was given on each day one glass slide with two pupae, an attached and a detached one (either two pupae of *D. simulans*, or two pupae of *D. suzukii*, we alternated colonies each day). We examined the ant-Drosophila interactions every 5 minutes for 3 hours after adding the glass slide with pupae.

We found that in all replicates all pupae were consumed by the ants. For both fly species, we observed a difference between detached and attached pupae: detached pupae were mostly taken to the nest and eaten there while most of the attached pupae were eaten on site (Fig. 3A, number of pupae taken to the nest in *D. simulans:* detached 21/21, attached 3/21, *chi2* = 28.10, *df* = 1, $p < 10^{-6}$; in *D. suzukii:* detached 19/21, attached 7/21, *chi2* = 12.22, *df* = 1, $p = 0.0005$). In 9 cases (6 for *D. suzukii* and 3 for *D. simulans*), both the attached and the detached pupa were taken to the nest; in all those cases the detached pupa was always taken to the nest first (about 37 min earlier for *D. suzukii* and 50 min earlier for *D. simulans* in median, Fig. 3A).

Ants spent 15 min and 45 min (median) outside the nest in contact with *D. simulans* and *D. suzukii* detached pupae, respectively, while they spent respectively six and two times longer in contact with attached pupae (Fig. 3B, respectively 100 min and 110 min in *D. simulans* and *D. suzukii*, paired Wilcoxon rank tests with continuity correction *D. simulans: V* = 4.5, $p = 0.0001$; *D. suzukii: V* = 18, $p = 0.0007$). The maximum number of ants observed in direct contact with the fly pupa over the duration of the experiment was significantly higher

for attached pupae than for detached pupae (Fig. 3A,C, median of 9 for *D. simulans* and 8 for

D. suzukii for attached pupae compared to median of 2 for detached pupae in both species,

paired Wilcoxon rank tests with continuity correction *D. simulans: V* = 2.5, *p* = 0.0002; *D.*

suzukii: V = 10.5, p = 0.0007). The difference was still significant after correcting for the

amount of time, by comparing the maximum number of ants in contact with the pupa until the

first pupa is brought to the nest or fully consumed (paired Wilcoxon rank tests with

continuity correction, *D. simulans*: 3 vs 2, *V* = 24, *p* = 0.02; *D. suzukii* 6 vs 2, *V* = 7, *p* =

0.004).

The time for the first ant to touch a pupa was slightly significantly different between

attached and detached pupae only in *D. suzukii* (*D. suzukii*: 15 min vs 20 min, *V* = 31, *p* =

0.03; *D. simulans*: 15 min vs 20 min, *V* = 62.5, *p* = 0.3) but not different between *D. simulans*

and *D. suzukii* (Wilcoxon rank tests with continuity correction, attached pupae: *W* = 195.5, *p*

= 0.5; detached pupae: *W* = 200, *p* = 0.6). No differences were found between *D. simulans*

and *D. suzukii* regarding the time until the detached pupae is brought to the nest from the start

of the experiment (45 min vs. 65 min, *W* = 160.5*, p* = 0.296), the time that ants spent in

contact with attached pupa (*W* = 194, *p* = 0.5) and detached pupa (*W* = 161, *p* = 0.1), the

maximum number of ants over the duration of the experiment on detached pupae (*W* = 196.5,

p = 0.5) or on attached pupae (*W* = 190.5, *p* = 0.5), the maximum number of ants until the

first pupa disappeared on detached pupae (*W* = 231.5, *p* = 0.8) and the maximum number of

ants on attached pupae (*W* = 265, *p* = 0.3).

Discussion

In the same environment, pupa adhesion strength varies among species.

Using our previously published pull-force measurement assay (Borne et al., 2020), we

provide here the first evidence that pupa adhesion varies between Drosophila species. Our

result is in agreement with the rapid evolution of glue genes (Da Lage et al., 2019). Our

analysis unravels at least two mechanisms leading to changes in the quantity of glue produced

and resulting in changes in adhesion strength among species: (1) a change in body size

(probably linked with a change in salivary gland size), as observed between *D. hydei* and *D.*

simulans, (2) a change in the amount of glue production independently of body size, as

observed between *D. suzukii* and *D. simulans*.

The variation that we observed in adhesion strength among individuals within a given

species is much higher than measurement error (our universal test machine has an accuracy of

±0.5%) and could be due to individual variation in size, shape, weight, glue production, or

position of the pupa relative to the substrate. We note that our experiment might not reflect

natural conditions as we have not tested adhesion on natural substrates and in natural

conditions. Drosophila pupation behavior and pupation sites have been thoroughly

investigated in the lab (Erezyilmaz and Stern, 2013; Schnebel and Grossfield, 1992; Vandal

et al., 2008) and more rarely in nature (Beltramí et al., 2010). Pupation behavior depends on

abiotic factors such as temperature (Schnebel and Grossfield, 1992), darkness (Rizki and

Davis, 1953) or the nature of the substrates (Godoy-Herrera and Luis, 1998). In particular, *D.*

simulans prefers to pupate on rough and humid surfaces while *D. hydei* prefers dry and

smooth surfaces. *D. simulans* was often reported as pupating in fruits in the lab (Erezyilmaz

and Stern, 2013; Vandal et al., 2008) and from field sampling (Beltramí et al., 2010) but other

natural sites have not been investigated. In *D. suzukii*, recent studies in the field have found

that pupae are present in the soil rather than in fruits (Ballman et al., 2017; Woltz and Lee,

2017). Additionally, pupation behavior depends on biotic factors and particularly on the

presence of conspecifics and alien species. In *D. simulans* and *D. hydei* and in other

Drosophila species, pupae are aggregated with conspecifics (Beltramí et al., 2010; Ringo and

Dowse, 2012). In *D. simulans* and *D. buzzatii*, larvae change their site choice in presence of

heterospecific larval cues (Beltramí et al., 2012). As many parameters seem to influence

pupation behavior, it is hard to know how differences in pupa ecology lead to differences in

pupa adhesion between species.

Fixation of the pupa prevents predation

Comparing the disappearance of attached and detached *D. simulans* pupae in the field,

we found that being attached allows pupae to stay on site more efficiently. Because pupae

lied within the lid of petri dishes, we can infer that pupae were not blown away by small wind

but we cannot know whether pupae that disappeared were indeed predated or not. The ant

Temnothorax nylanderi was the main predator that we observed in the field consuming

pupae. This observation is in agreement with previous studies which found that ants prey on

fruit fly pupae (Urbaneja et al., 2006; Ballman et al., 2017; Woltz and Lee, 2017). During our

laboratory assay, *T. nylandei* ants predated attached and detached pupae with distinct

behaviours: they brought most of the detached pupae to the nest while they ate the attached ones directly on site. The latter strategy requires ants to spend more time outside the nest and to recruit more foragers. The presence of parasites, predators and competitors in the wild would certainly make this strategy costly. Additionally, *T. nylanderi* is a solitary foraging species that recruits nestmates one by one (Glaser and Grueter, 2018). Under natural conditions, it would take a relatively long time to gather many foragers around the pupae. In our field assay, no more than two ants were observed together in a lid (Table S5). We found that ants act similarly on *D. suzukii* and *D. simulans* pupae, suggesting that there is no difference in strategy to predate loosely attached pupae such as *D. suzukii* or more strongly attached ones such as *D. simulans*, and that both species are equally attractive as preys.

To our knowledge, this study is the first to show that fly pupa adhesion can protect from predation. Our experiments are simple and can be easily applicable to other species, and not only for pupae but also for eggs, to check if this phenomenon is general in flies and insects.

Other strategies can protect pupae from predation

Pupal adhesion is only one of several strategies for pupae to escape predators. A common strategy is cryptic coloration to hide from visual predators (Gaitonde et al., 2018; Lindstedt et al., 2019). The brownish color of Drosophila pupae could contribute to hiding the animal when pupating in the leaf litter or in the soil. In some cases, pupae mimic non-living things such as leaves or sticks like the Common Maplet butterfly chrysalis (Lindstedt et al., 2019). To avoid non visual predators, pupa has also evolved chemical defences either to chemically hide from predators (Mizuno et al., 2018) or to make the pupa toxic (Deyrup et al., 2014). Pupae have also evolved different types of physical defence (spin, hard pupal case, urticating hairs...). In Drosophila, the pupa is covered by a relatively thick (about 20 μm, Mitchell et al., 1971) and hard cuticle which has been hypothesized to protect the animal from predator attacks. Additionally, pupae have evolved different behaviors such as interacting with conspecifics during larval stage particularly to form aggregation. Aggregated pupae may be more visible to visual predators but predation risk is diluted in group living (Wrona and Dixon, 1991). This last strategy is found in Drosophila species (Beltramí et al., 2010; Ringo and Dowse, 2012) but its contribution to protection from predators has not been tested.

443   For the first time, we report that pupa adhesion varies among Drosophila species and

444 that pupa attachment can protect from predation. Our results unravel a previously unknown

445 important trait for Drosophila survival in the wild, the ability of pupae to firmly adhere to a

446 substrate. Further studies of Drosophila glue combining genetic and phenotypic approaches

447 should provide insight on the molecular basis for diverse bioadhesive properties, adapted to

448 various habitats and climates.

Data Availability Statement

450 Raw data and scripts have been deposited in DRYAD (temporary link for private access:

451 https://datadryad.org/stash/share/SnN69p739EA9Q0KW-

452 Tz6B7AZwEDkfRTZNIEedf6sZV8).

Acknowledgements

We are grateful to Michel Neff and the Division du Bois de Vincennes de la Ville de Paris for

allowing us to use an area of wood land of the Réserve Ornithologique du Bois de Vincennes

as an experimental plot. We thank Céline Bocquet for her help regarding collection and

rearing of ant colonies, Manon Monier and Gauthier Beuque for their help with experiment

installation in Vincennes and Aurélie Surtel for drawing the outlines of all the glue prints.

The research leading to this paper has received funding from CNRS (MITI project

GLUEADAPT) and from the European Research Council under the European Community's

Seventh Framework Program (FP7/2007–2013 Grant Agreement no. 337579) to V.C.-O. F.B.

was supported by a PhD fellowship from École normale supérieure Paris-Saclay. MM and the

Vincennes setup were supported by the Institute of Ecology and Environmental Sciences -

Paris and the Institut de la Transition Environnementale de Sorbonne Université.

Author Contributions

468 FB, MM and VCO designed the experiments, VCO supervised the project, SRP collected,

469 identified and raised wild-caught flies, FB performed all the other experiments, FB, MM and

470 VCO analyzed the results, FB wrote the original draft and all authors edited and contributed

471 to the final version of this paper.

Conflict of Interest Statement

The authors have no conflict of interest to declare.

References

Ballman, E. S., Collins, J. A., & Drummond, F. A. (2017). Pupation Behavior and Predation on Drosophila suzukii (Diptera : Drosophilidae) Pupae in Maine Wild Blueberry Fields. *Journal of Economic Entomology*, *110*(6), 2308-2317. https://doi.org/10.1093/jee/tox233

Beckendorf, S. K., & Kafatos, F. C. (1976). Differentiation in the salivary glands of Drosophila melanogaster : Characterizaion of the glue proteins and their developmental appearance. *Cell*, *9*(3), 365-373. https://doi.org/10.1016/0092-8674(76)90081-7

Beltramí, M., Medina-Muñoz, M., Pino, F., Ferveur, J.-F., & Godoy-Herrera, R. (2012). Chemical Cues Influence Pupation Behavior of Drosophila simulans and Drosophila buzzatii in Nature and in the Laboratory. *PloS one*, *7*, e39393. https://doi.org/10.1371/journal.pone.0039393

Beltramí, M., Muñoz, M., Arce, D., & Godoy-Herrera, R. (2010). Drosophila pupation behavior in the wild. *Evolutionary Ecology*, *24*. https://doi.org/10.1007/s10682-009-9310-8

Beňová-Liszeková, D., Beňo, M., & Farkaš, R. (2019). Fine infrastructure of released and solidified *Drosophila* larval salivary secretory glue using SEM. *Bioinspiration & Biomimetics*, *14*(5), 055002. https://doi.org/10.1088/1748-3190/ab2b2b

Bigler, F., Neuenschwander, P., Delucchi, V., & Michelakis, S. (1986). Natural enemies of preimaginal stages of Dacus oleae Gmel.(Dipt., Tephritidae) in Western Crete. II.

Impact on olive fly populations. *Bollettino del Laboratorio di Entomologia Agraria"*

Filippo Silvestri", Italy, *43*, 79–96.

Borne, F., Kovalev, A., Gorb, S., & Courtier-Orgogozo, V. (2020). The glue produced by

Drosophila melanogaster for pupa adhesion is universal. *The Journal of Experimental*

Biology, *223*(8), jeb220608. https://doi.org/10.1242/jeb.220608

Chapman, R. F. (1982). *The Insects : Structure and function*. Harvard University Press.

Da Lage, J.-L., Thomas, G. W. C., Bonneau, M., & Courtier-Orgogozo, V. (2019). Evolution

of salivary glue genes in Drosophila species. *BMC Evolutionary Biology*, *19*(1), 36.

https://doi.org/10.1186/s12862-019-1364-9

Deyrup, S. T., Eckman, L. E., Lucadamo, E. E., McCarthy, P. H., Knapp, J. C., & Smedley,

S. R. (2014). Antipredator activity and endogenous biosynthesis of defensive

secretion in larval and pupal Delphastus catalinae (Horn) (Coleoptera :

Coccinellidae). *Chemoecology*, *24*(4), 145-157. https://doi.org/10.1007/s00049-014-

0156-3

Erezyilmaz, D. F., & Stern, D. L. (2013). Pupariation Site Preference Within and Between

Drosophila Sibling Species. *Evolution*, *67*(9), 2714-2727.

https://doi.org/10.1111/evo.12146

Fordyce, J. A., & Nice, C. C. (2002). Variation in butterfly egg adhesion : Adaptation to local

host plant senescence characteristics?: Variation in butterfly egg adhesion. *Ecology*

Letters, *6*(1), 23-27. https://doi.org/10.1046/j.1461-0248.2003.00389.x

Fraenkel, G., & Brookes, V. J. (1953). THE PROCESS BY WHICH THE PUPARIA OF

MANY SPECIES OF FLIES BECOME FIXED TO A SUBSTRATE. *The Biological*

Bulletin, *105*(3), 442-449. https://doi.org/10.2307/1538461

Gabarra, R., Riudavets, J., Rodríguez, G. A., Pujade-Villar, J., & Arnó, J. (2015). Prospects

for the biological control of Drosophila suzukii. *BioControl*, *60*(3), 331-339.

https://doi.org/10.1007/s10526-014-9646-z

Gaitonde, N., Joshi, J., & Kunte, K. (2018). Evolution of ontogenic change in color defenses

of swallowtail butterflies. *Ecology and Evolution*, *8*(19), 9751-9763.

https://doi.org/10.1002/ece3.4426

Glaser, S., & Grueter, C. (2018). Ants (Temnothorax nylanderi) adjust tandem running when

food source distance exposes them to greater risks. *Behavioral Ecology and*

Sociobiology, *72*, 40. https://doi.org/10.1007/s00265-018-2453-2

Godoy-Herrera, R., & Luis, S.-C. (1998). The behavior of sympatric Chilean populations of

Drosophila larvae during pupation. *Genetics and Molecular Biology*, *21*.

https://doi.org/10.1590/S1415-47571998000100007

Grossfield. (1978). Non-sexual behavior of Drosophila. In *The genetics and biology of*

Drosophila (Ashburner M, Wright TRF, Vol. 2b, p. 3-126). Academic Press.

Harvey, J. A., Gols, R., & Tanaka, T. (2011). Differing Success of Defense Strategies in Two

Parasitoid Wasps in Protecting Their Pupae Against a Secondary Hyperparasitoid.

Annals of the Entomological Society of America, *104*(5), 1005-1011.

https://doi.org/10.1603/AN10192

Hayes, J. L. (1982). A Study of the Relationships of Diapause Phenomena and Other Life

History Characters in Temperate Butterflies. *The American Naturalist*, *120*(2),

160-170. https://doi.org/10.1086/283979

Heming, B. S. (2003). *Insect Development and Evolution*. Cornell University Press.

https://www.jstor.org/stable/10.7591/j.ctv75d5sv

Hennessey, M. K. (1997). Predation on wandering larvae and pupae of caribbean fruit fly

(diptera : Tephritidae) in guava and carambola grove soils. *Journal of Agricultural and*

Urban Entomology, *14*(2), 129-138. Scopus.

Hieber, C. S. (1992). Spider cocoons and their suspension systems as barriers to generalist

and specialist predators. *Oecologia*, *91*(4), 530-535.

https://doi.org/10.1007/BF00650327

Hinton, H. E. (1981). *Biology of insect eggs* (1st ed). Pergamon Press.

Honorio, R., Doums, C., & Molet, M. (2020). Manipulation of worker size diversity does not

affect colony fitness under natural conditions in the ant Temnothorax nylanderi.

Behavioral Ecology and Sociobiology, *74*(8), 104. https://doi.org/10.1007/s00265-

020-02885-2

Korge, G. (1975). Chromosome puff activity and protein synthesis in larval salivary glands of

Drosophila melanogaster. *Proceedings of the National Academy of Sciences*, *72*(11),

4550-4554. https://doi.org/10.1073/pnas.72.11.4550

Korge, Günter. (1977). Larval saliva in Drosophila melanogaster : Production, composition,

and relationship to chromosome puffs. *Developmental Biology*, *58*(2), 339-355.

https://doi.org/10.1016/0012-1606(77)90096-3

Lindstedt, C., Murphy, L., & Mappes, J. (2019). Antipredator strategies of pupae : How to

avoid predation in an immobile life stage? *Philosophical Transactions of the Royal*

Society B: Biological Sciences, *374*(1783), 20190069.

https://doi.org/10.1098/rstb.2019.0069

Mitchell, H. K., Weber-Tracy, U. M., & Schaar, G. (1971). Aspects of cuticle formation in

Drosophila melanogaster. *Journal of Experimental Zoology*, *176*(4), 429-443.

https://doi.org/10.1002/jez.1401760406

Mizuno, T., Hagiwara, Y., & Akino, T. (2018). Chemical tactic of facultative myrmecophilous

lycaenid pupa to suppress ant aggression. *Chemoecology*, *28*(6), 173-182.

https://doi.org/10.1007/s00049-018-0270-8

Nakagawa, S. (2004). A farewell to Bonferroni : The problems of low statistical power and

publication bias. *Behavioral Ecology*, *15*(6), 1044-1045.

https://doi.org/10.1093/beheco/arh107

Ringo, J., & Dowse, H. (2012). Pupation Site Selection in Four Drosophilid Species :

Aggregation and Contact. *Journal of Insect Behavior*, *25*.

https://doi.org/10.1007/s10905-012-9324-x

Rizki, M. T. M., & Davis, Charles G. (1953). Light as an Ecological Determinant of

Interspecific Competition between Drosophila willistoni and Drosophila melanogaster.

The American Naturalist, *87*(837), 389-392. https://doi.org/10.1086/281799

Robertson, J. A., Bradler, S., & Whiting, M. F. (2018). Evolution of Oviposition Techniques in

Stick and Leaf Insects (Phasmatodea). *Frontiers in Ecology and Evolution*, *6*, 216.

https://doi.org/10.3389/fevo.2018.00216

Schmidt-Nielsen, K. (1997). *Animal physiology : Adaptation and environment*. Cambridge

University Press.

Schnebel, E. M., & Grossfield, J. (1992). Temperature effects on pupation-height response

in four Drosophila species group triads. *Journal of Insect Physiology*, *38*(10),

727-732. https://doi.org/10.1016/0022-1910(92)90024-8

Schneider, C. A., Rasband, W. S., & Eliceiri, K. W. (2012). NIH Image to ImageJ : 25 years

of image analysis. *Nature Methods*, *9*(7), 671-675.

https://doi.org/10.1038/nmeth.2089

Shirai, S., & Maeto, K. (2009). Suspending cocoons to evade ant predation in *Meteorus*

pulchricornis , a braconid parasitoid of exposed-living lepidopteran larvae.

Entomological Science, *12*(1), 107-109. https://doi.org/10.1111/j.1479-

8298.2009.00301.x

Thomas, D. B. (1995). Predation on the soil inhabiting stages of the Mexican fruit fly.

Southwestern Entomologist, *20*(1), 61-71.

Thomas, Donald B. (1993). Survivorship of the Pupal Stages of the Mexican Fruit Fly

Anastrepha ludens (Loew) (Diptera : Tephritidae) in an Agricultural and a

Nonagricultural Situation. *Journal of Entomological Science*, *28*(4), 350-362.

https://doi.org/10.18474/0749-8004-28.4.350

Urbaneja, A., Marí, F. G., Tortosa, D., Navarro, C., Vanaclocha, P., Bargues, L., &

Castañera, P. (2006). Influence of Ground Predators on the Survival of the

Mediterranean Fruit Fly Pupae, Ceratitis capitata, in Spanish Citrus Orchards.

Biocontrol, *51*(5), 611-626. https://doi.org/10.1007/s10526-005-2938-6

Vandal, N. B., Siddalingamurthy, G. S., & Shivanna, N. (2008). Larval pupation site

preference on fruit in different species of *Drosophila*. *Entomological Research*, *38*(3),

188-194. https://doi.org/10.1111/j.1748-5967.2008.00163.x

Voigt, D., & Gorb, S. (2010). Egg attachment of the asparagus beetle Crioceris asparagi to

the crystalline waxy surface of Asparagus officinalis. *Proceedings of the Royal*

Society B: Biological Sciences, *277*(1683), 895-903.

https://doi.org/10.1098/rspb.2009.1706

Wahl, M. (1997). Living attached : Aufwuchs, fouling, epibiosis. Fouling organisms of the

Indian Ocean: biology and control technology. *Oxford & IBH Publ. Co. Put Ltd., New*

Delhi, 31-83.

Warton, D. I., Duursma, R. A., Falster, D. S., & Taskinen, S. (2012). Smatr 3– an R package

for estimation and inference about allometric lines. *Methods in Ecology and*

Evolution, *3*(2), 257-259. https://doi.org/10.1111/j.2041-210X.2011.00153.x

Wiklund, C. (1984). Egg-laying patterns in butterflies in relation to their phenology and the

visual apparency and abundance of their host plants. *Oecologia*, *63*(1), 23-29.

https://doi.org/10.1007/BF00379780

Woltz, J. M., & Lee, J. C. (2017). Pupation behavior and larval and pupal biocontrol of

Drosophila suzukii in the field. *Biological Control*, *110*, 62-69.

https://doi.org/10.1016/j.biocontrol.2017.04.007

Wrona, F. J., & Dixon, R. W. J. (1991). Group Size and Predation Risk : A Field Analysis of

Encounter and Dilution Effects. *The American Naturalist*, *137*(2), 186-201.

Supplementary materials (available in DRYAD)

Table S1. List of Drosophila flies collected in Bois de Vincennes. The number corresponds
to the total number of individuals collected at each site. Traps were emptied twice, two days
and five days after traps were set up in the field. GPS coordinates are shown for each trap and
for the compost.

Table S2. Adhesion force and pupa-substrate interface measurements. Results of the

adhesion assay performed on pupae from *D. suzukii*, *D. simulans* and *D. hydei* strain originating from Vincennes. Sample_ID column corresponds to a unique identification for each pupa, Temperature_assay, Humidity, Pressure_mba indicate respectively the room temperature (°C), the room humidity (%) and the atmospheric pressure (mba) at the moment of the assay. date_measurement and time_measurement correspond to the day and hours of the assay. date_substrate and time_substrate correspond to the day and hours at which larvae were put on the substrate. force_detachment_mN corresponds to the maximum force reached during the experiment in mN. Comment_on_this_sample reports particular observations: the pupa did not detach from the substrate ("not_detached"), the pupal case broke during the assay ("cuticle_broke") or nothing special happened during the assay ("ok"). Area_px corresponds to the pupa-substrate contact surface measured in pixel and scale_px, scale_mm correspond respectively to the scale present on the picture in pixel and mm.

Table S3. Pupal size measurement. Measurements of the size of pupae of *D. simulans*, *D. suzukii*, *D. hydei*. ID column corresponds to a unique identification for each pupa. area corresponds to the area obtained by measuring the contour of the pupa in pixel. scale_px and scale_mm are defined as in Table S2.

Table S4. Pupal count in the field experiment. Results of the experiment performed in the field. bucket_ID column corresponds to a unique identification for each bucket. Orientation_C and Orientation_NC give, respectively, the orientation of attached and detached pupa within the bucket. Count_C and Count_NC indicate, respectively, the number of attached and detached pupa in a dish. Time indicates the time at which pupae were counted: t0 after 0h (8 September 2020, day 1 morning, at 11 am) t1 after 6h30 (day 1 PM), t2 after 22h30 (day 2 AM), t3 after 31h (day 2 PM), t4 after 47h (day 3 AM), and t5 after 54h (day 3 PM).

Table S5. Animals observed in the dishes during the experiment with half-buried buckets in Bois de Vincennes. The number indicates the number of dishes where the respective animals were observed. In total, 56 dishes were examined at each time point.

Table S6. Ant count during predation assay in the laboratory. The number indicates the number of ants observed in direct contact with the pupa at each time point. The column "X_Y" corresponds to the count over one experiment with the ant colony X at the trial Y.

Table S7. Results table of the predation assay in the laboratory. Each row corresponds to the description of the experiment for one pupa. condition column indicates the initial state of the pupa ("attached" or "detached"), strategies indicates whether the pupa was brought to the nest over the duration of the experiment ("to_nest") or was never brought to the nest and consumed outside the nest ("on_site"). Time corresponds to the time at which the pupa was brought to the nest or fully consumed outside the nest in min. max_ant corresponds to the maximum number of ants observed in contact with the pupa over the duration of the

experiment. orientation indicates whether the pupa was initially on the left or on the right side of the glass slide at the beginning of the experiment. first_ant corresponds to the time in min at which the first interaction between the pupa and an ant is observed. max_ant_at_first_done corresponds to the maximum number of ants observed in direct interaction with the pupa between the beginning and the time when the first pupa is brought to the nest or fully consumed outside the nest.

R script. Drosophila_glue_predation.R R script used to prepare the figures and run the statistical tests.

pupa_prints Folder containing pictures of the prints left by the pupae on glass slides after detachment during our adhesion assay force measurements.

pupal_size Folder containing the pictures of pupae used to measure the size of the pupae of the different species.

Pictures_Vincennes Folder containing pictures taken during and after the experiments in the field.

Pictures_predation_lab Folder containing pictures and video of the predation assays in the laboratory.