Massive colonization of protein-coding exons by selfish genetic elements in *Paramecium* germline genomes

Short title: Massive invasion of *Paramecium* genes by selfish genetic elements

Diamantis Sellis[1#a¶], Frédéric Guérin[2#b¶], Olivier Arnaiz[3], Walker Pett[1], Emmanuelle Lerat[1],

Nicole Boggetto[2], Sascha Krenek[4], Thomas Berendonk[4], Arnaud Couloux[5], Jean-Marc Aury[5],

Karine Labadie[6], Sophie Malinsky[7,8], Simran Bhullar[7], Eric Meyer[7], Linda Sperling[3], Laurent

Duret[1&*], Sandra Duharcourt[2&*]

[1] Université de Lyon, CNRS, Laboratoire de Biométrie et Biologie Evolutive UMR 5558, F-69100, Villeurbanne, France

[2] Université de Paris, CNRS, Institut Jacques Monod, F-75006 Paris, France

[3] Université Paris-Saclay, CEA, CNRS, Institute for Integrative Biology of the Cell (I2BC), 91198, Gif-sur-Yvette, France.

[4] TU Dresden, Institute of Hydrobiology, Dresden, Germany

[5] Génomique Métabolique, Genoscope, Institut de biologie François Jacob, CEA, CNRS, Université d'Évry, Université Paris-Saclay, F-91000 Evry, France.

[6] Genoscope, Institut de biologie François-Jacob, Commissariat à l'Energie Atomique (CEA), Université Paris-Saclay, F-91000 Evry, France.

[7] Institut de Biologie de l'Ecole Normale Supérieure (IBENS), Ecole Normale Supérieure, CNRS, INSERM, Université PSL, Paris, France.

[8] Université de Paris, F-75006 Paris, France

[#a]Current Address: CosmoTech, 5 Passage du Vercors 69007 Lyon, France
[#b]Current Address: Scipio bioscience, 92120 Montrouge, France

* Corresponding authors
Emails: sandra.duharcourt@ijm.fr; Laurent.Duret@univ-lyon1.fr

[¶]These authors contributed equally to this work.
[&]These authors also contributed equally to this work.

Abstract

Ciliates are unicellular eukaryotes with both a germline genome and a somatic genome in the same cytoplasm. The somatic macronucleus (MAC), responsible for gene expression, is not sexually transmitted but develops from a copy of the germline micronucleus (MIC) at each sexual generation. In the MIC genome of *Paramecium tetraurelia*, genes are interrupted by tens of thousands of unique intervening sequences, called Internal Eliminated Sequences (IESs), that have to be precisely excised during the development of the new MAC to restore functional genes. To understand the evolutionary origin of this peculiar genomic architecture, we sequenced the MIC genomes of nine *Paramecium* species (from ~100 Mb in *P. aurelia* species to > 1.5 Gb in *P. caudatum*). We detected several waves of IES gains, both in ancestral and in more recent lineages. While the vast majority of IESs are single-copy in present-day genomes, we identified several families of mobile IESs, including non-autonomous elements acquired via horizontal transfer, that generated tens to thousands of new copies. These observations provide the first direct evidence that transposable elements can account for the massive proliferation of IESs in *Paramecium*. The comparison of IESs of different evolutionary ages indicates that, over time, IESs shorten and diverge rapidly in sequence while they acquire features that allow them to be more efficiently excised. We nevertheless identified rare cases of IESs that are under strong purifying selection across the *aurelia* clade. The cases examined contain or overlap cellular genes that are inactivated by excision during development, suggesting conserved regulatory mechanisms. Similar to the evolution of introns in eukaryotes, the evolution of *Paramecium* IESs highlights the major role played by selfish genetic elements in shaping the complexity of genome architecture and gene expression.

Introduction

In multicellular organisms, the division of labor between transmission and expression of the genome is achieved by separation of germline and somatic cells. Such a division is also observed in some unicellular eukaryotes, including ciliates [1]. The ciliate *Paramecium tetraurelia* separates germline and somatic functions into distinct nuclei in the same cell. Somatic functions are supported by the highly polyploid macronucleus (MAC) that is streamlined for gene expression and destroyed at each sexual cycle. Germline functions are ensured by two small, diploid micronuclei (MIC) that are transcriptionally silent during vegetative growth. During sexual events, the MICs undergo meiosis and transmit the germline genome to the zygotic nucleus. New MICs and new MACs differentiate from mitotic copies of the zygotic nucleus. MAC differentiation involves massive and reproducible DNA elimination events (for review: [2,3]). In addition to the variable elimination of large regions containing repeats, ~45,000 unique, short, interspersed Internal Eliminated Sequences (IESs) are precisely removed from intergenic and coding regions [4,5]. Precise excision of IESs at the nucleotide level is essential to restore functional cellular genes, since 80% of the IESs are inserted within protein-coding genes, and about half of the ~40,000 genes are interrupted by IESs. IESs are invariably bounded by two 5'-TA-3' dinucleotides, one of which is left at the junction in the MAC genome after excision. IES excision in the developing MAC is initiated by DNA double-strand breaks at IES ends by the endonuclease PiggyMac (Pgm) assisted by other proteins, which are likely part of the excision machinery or interact with it [6–9].

Despite significant progress in characterization of the mechanisms underlying IES elimination, the evolutionary origin of IESs remains mysterious. On the basis of sequence similarities between the consensus found adjacent to the TA dinucleotide at IES ends and the extremities

of DNA transposons from the IS630-Tc1-mariner (ITm) superfamily, Klobutcher and Herrick

hypothesized that IESs might be degenerated remnants of transposable elements (TEs)[10,11].

This hypothesis was further substantiated by the discovery that the endonuclease responsible

for IES excision in *P. tetraurelia* is encoded by a domesticated PiggyBac transposase [6],

assisted by a related family of catalytically inactive transposases [7]. All-by-all sequence

comparison of the *P. tetraurelia* 45,000 IESs and of their flanking sequences identified eight

families of "mobile IESs" (2 to 6 copies), *i.e.* homologous IESs inserted at non-homologous

sites in the genome [4]. One such family (with 6 copies) was found similar to the Terminal

Inverted Repeats of *Thon*, a DNA transposon of the ITm superfamily, indicating that some IESs

derive from TEs [4]. These cases provided support to the notion that at least some IESs have

derived from recently mobilized elements. However, the rather small number of mobile IESs

detected (23 copies out of 45,000 IESs) suggested a limited activity of transposable IESs in the

recent evolutionary history of the *P. tetraurelia* lineage [4]. There is also evidence that some

IESs originated from MAC sequences, as described for instance for the IESs involved in mating

type determination in several species [12,13]. The extent to which the 45,000 IESs detected in

P. tetraurelia derive from TEs or from MAC sequences therefore remained unclear.

In order to gain insight into the evolutionary origin of IESs in the *Paramecium* lineage, we

adopted a comparative genomic approach. *P. tetraurelia* belongs to the *Paramecium aurelia*

group of species that comprises a score of morphologically similar yet genetically isolated

species [14–17] (see [13] for a detailed description of the species). Here, we sequenced the

germline MIC genomes of eight *P. aurelia* species and one outgroup (*P. caudatum*), compared

the IES repertoire across these nine species, and analyzed the evolutionary trajectories of IESs

in the *Paramecium* lineage.

Results

Sequencing of somatic and germline genomes in nine *Paramecium* species: gigantic germline genome in *P. caudatum*

In order to determine the origin and evolution of IESs in the *Paramecium* lineage, we sequenced the germline MIC genome and the somatic MAC genome of several *Paramecium* species. We selected 8 species from the *Paramecium aurelia* complex and one outgroup species, *P. caudatum*, which diverged from the *aurelia* complex before the two most recent *Paramecium* whole genome duplications [15]. To sequence the germline MIC genome, we purified the germline nuclei (MICs) of each species using a flow cytometry procedure that we previously developed for *P. tetraurelia* [5] (S1 Fig), and used paired-end Illumina sequencing (see Materials and Methods and S1 Table).

Species (strain)	MIC genome		MAC-destined regions			
	Size (Mb) (a)	Sequencing Depth (b)	Size (Mb) (c)	Nb. of protein genes	Nb. of IESs (d)	IES density (per kb) (e)
P. tetraurelia (51)	108	31	72	40,460	44,128	0.62
P. octaurelia (138)	108	56	72	44,398	44,509	0.61
P. biaurelia (V1-4)	119	58	74	40,261	45,384	0.65
P. tredecaurelia (209)	127	71	65	36,179	42,275	0.66
P. pentaurelia (87)	112	194	72	41,676	42,686	0.57
P. primaurelia (AZ9-3)	114	65	73	42,619	43,766	0.59
P. sonneborni (ATCC 30995)	286	13	82	49,951	60,198 (~85,000)	1.05
P. sexaurelia (AZ8-4)	123	141	68	36,094	47,002	0.70

					8,762	
P. caudatum (My43c3d)	1,300	13	30	18,673	(~15,000)	0.47

Table 1. Characteristics of analyzed genomes.

Species are ordered according to the phylogeny (Fig 1). (a) MIC genome size (in Mb) was estimated based on k-mer counts (see S1 Table for additional estimates based on flow cytometry). (b) Sequencing depth in MIC-specific regions estimated from the distribution of k-mer counts (S2 Fig). (c) Size of constitutive MAC genome assembly. (d) The sensitivity of IES detection was limited in *P. caudatum* and *P. sonneborni* because of the relatively low sequencing depth of their MIC genome. Estimates of total number of IESs, based the IES density observed in regions with sufficient read depth (see e), are indicated in parenthesis for these two species. (e) IES density measured in MAC-destined sequences, after exclusion of regions with insufficient MIC read depth (< 15X).

To estimate the size of the MIC genomes, we employed two distinct approaches. First, we used the MIC-enriched preparations from *Paramecium* cultures to yield values for DNA quantity in the MICs by flow cytometry (see Materials and Methods and S1 Data). The estimated MIC genome sizes are within a similar range (140-173 Mb) for the *P. aurelia* species, except for *P. sonneborni* (S1 Table), which was estimated to be roughly the double of the others. The second, independent approach for genome size estimations was based on the sequence reads themselves and used the k-mer method described in [18,19]. The estimated MIC genome sizes were comprised between 108 Mb to 123 Mb for the *aurelia* species, with a considerably larger MIC genome (283 Mb) for *P. sonneborni* (Table 1). While the values obtained using the flow cytometry method were greater than those with the k-mer method, the estimated MIC genome sizes were within a similar range for both methods (S3 Fig).

With both methods, the estimated MIC genome size of *P. caudatum* strain My43c3d was the

largest among the species analyzed (app. 1,300-1,600 Mb). To confirm this observation, we

estimated the genome size of 9 additional strains belonging to the two major clades A and B

described in the *caudatum* lineage, as well as another divergent strain [20]. The data confirmed

that the MIC genome size in the *caudatum* lineage is far bigger than that in the *aurelia* lineage

and revealed great variations of genome size among the different strains (from 1,600 Mb to

5,500 Mb), even within the same clade (S1 Data). To investigate the composition of the gigantic

caudatum genomes, we searched for the presence of repeats in the MIC sequence reads of strain

My43c3d. We identified two major satellite repeats, Sat1 and Sat2 (332 bp and 449 bp,

respectively), which represent 42% and 29% respectively of the MIC genome (S4 Fig). Both

Sat1 and Sat2 repeats were detected in the *P. caudatum* strains of the clade B, to which the

strain My43c3d belongs, but not in the other *P. caudatum* strains (S4 Fig). Thus, the repeats are

not shared by all *P. caudatum* strains and most likely invaded the MIC genome after the

divergence between clades A and B.

The MAC genome was sequenced for four *aurelia* species that had not been sequenced

previously (S2 Table). We defined the '*constitutive*' MAC genome as the DNA sequences

retained in all MAC copies (see Materials and Methods and S2 Table). The size of the

constitutive MAC genome assembly was similar among *P. aurelia* species (66-73 Mb) with a

noticeably larger size for *P. sonneborni* (83 Mb) (Table 1). The number of protein coding genes

follows a similar distribution (36,179 to 42,619) in *aurelia* species, with a larger number of

genes in *P. sonneborni* (Table 1). This contrasts with the much smaller MAC genome size and

number of genes of the outgroup *P. caudatum* [15].

In conclusion, the eight species of the *aurelia* complex that we analyzed share similar genome

characteristics, with a MIC genome of ~110-160 Mb, 50-70% of which is retained during MAC

development (~70-80 Mb). The only notable exception is *P. sonneborni*, with a 300-400 Mb

MIC genome, of which about 25% is retained in its MAC. The MIC genome of the outgroup

P. caudatum is much larger (~1,300-1,600 Mb). Only 2% of MIC sequences are retained in the

MAC of *P. caudatum* strain My43c3d and 83% of the MIC-specific sequences consist of

repeated DNA (S4 Fig).

IES repertoire

IESs were identified by comparing MIC sequence reads to the MAC genome assembly (see

Materials and Methods; [4,21]). Overall, the number of detected IESs in MAC-destined

sequences is similar across *Paramecium* species (~42,000-47,000 IESs), with the exception of

P. sonneborni (~60,000 IESs) and *P. caudatum* (~9,000 IESs). It should be noted that the

sensitivity of IES detection depends on sequencing depth (S5 Fig). For most species, we expect

that >99% of IESs have been detected, except in *P. caudatum*, and *P. sonneborni*, where the

sequencing depth was limited, due to the unexpected large size of their MIC genome (Table 1).

To circumvent this issue, we compared the IES density across species by taking into account

only IESs annotated in regions with at least 15X depth of MIC sequence reads mapped onto the

MAC assembled genome (Table 1): in *P. caudatum*, the density of detected IES sites in MAC-

destined regions (0.5 IESs per kb) is only slightly lower than in other species (~0.6 IESs per

kb). This suggests that the genome of *P. caudatum* probably contains about 15,000 IESs in its

MAC-destined regions.

Our approach is designed to identify IESs only if they are present within loci retained in the

MAC. Hence, IESs located in MIC-specific regions (e.g., IESs nested within other IESs

[22,23]) remain undetected. Interestingly, in four species whose MAC genome was sequenced

at very high depth *(P. octaurelia, P. primaurelia, P. pentaurelia* and *P. sonneborni*), the initial

MAC genome assemblies included 10 to 16 Mb of MAC-variable regions, which correspond

to DNA sequences that are not completely eliminated and instead, are retained in a small

fraction of MAC copies (see Materials and Methods, S6 Fig and S2 Table). We identified many

IESs in these regions, at a density (0.4 to 0.5 IESs per kb, S2 Table) nearly as high as in MAC-

destined regions (Table 1). This suggests that in addition to IESs located in MAC-destined

regions, many other IESs are present within MIC-specific regions.

In all species, the vast majority of IESs in MAC-destined regions (73% to 81%) are located in

protein-coding exons and ~5% are located in introns. Overall, there is a slight enrichment of

IESs within genes (S3 Table). This enrichment is not true for all gene categories. In particular,

we observed a depletion of IESs in highly expressed genes: on average, the IES density in the

top 10% most expressed genes is 37% lower than in the bottom 10% (S7 Fig). This pattern,

consistent with previous observations in *P. tetraurelia*, suggests that IES insertions are counter-

selected in highly expressed genes [4].

Age distribution of IESs

In order to explore the origin and evolution of IESs, we resolved the phylogenetic relationship

among the sequenced species. To do so, we classified all protein sequences into families

(N=13,617 gene families) and inferred the species phylogeny using the subset of 1,061 gene

families containing one single sequence from each species. In agreement with previous reports

[16,24], we found strong support for a division of the *aurelia* complex in two subclades

(hereafter referred to as subclades A and B), separating *P. sonneborni* and *P. sexaurelia* from

the other *aurelia* species (Fig 1). We then used this species phylogeny to identify gene

duplications and speciation events in each of the 13,617 gene families, using the PHYLDOG

tree reconciliation method [25].

Fig 1. Dynamics of IES insertion/loss in *Paramecium*.

The species phylogeny was reconstructed from a concatenated alignment of 1,061 single-copy

genes. All internal branches are supported by 100% bootstrap values (except branch *:

bootstrap support = 83%). The age of IESs located within coding regions was inferred from the

pattern of presence/absence within gene family alignments (N=13,617 gene families). Only

IESs present within well-aligned regions were included in this analysis. The number of dated

IESs and the fraction predicted to be old (predating the divergence between *P. caudatum* and

the *P. aurelia* lineages), intermediate (before the radiation of the *P. aurelia* complex) or recent

are reported for each species. Rates of IES gain (in red) and loss (in blue) were estimated along

each branch using a Bayesian approach. Gain rates are expressed per kb per unit of time (using

the branch length – in substitutions per site – as a proxy for time). Loss rates are expressed per

IES per unit of time (see S8 Fig for more details on the measure of gain/loss rates). NB:

estimates of loss rate along terminal branches of the phylogeny also include false negatives (i.e.

IESs that are present but that have not been detected), and hence may be overestimated.

In order to date events of IES gain or loss, we mapped the position of IES excision sites in

multiple alignments of each gene family (nucleic sequence alignments based on protein

alignments): IESs located at the exact same position within a codon were assumed to derive

from a single ancestral insertion event, whatever their present level of sequence similarity (S9

Fig). IESs located at homologous sites are hereafter referred to as 'co-orthologous' IESs. We

use the term 'co-orthologous' rather than 'orthologous', because these sets frequently include

paralogs, notably as a result of whole genome duplications. To avoid ambiguities due to low

quality alignments, we only analyzed IESs present within well-conserved protein-coding

regions (which represent from 45% to 51% of IESs located in coding regions; Fig 1). We then

used the reconciled gene tree to map events on the species phylogeny and estimate rates of IES

gain and loss along each branch of the species tree using a Bayesian approach accounting for

IES losses and missing data (see Materials and Methods). In the absence of fossil records, it is

impossible to date speciation events (in million years). We therefore used sequence divergence

(number of amino-acid substitutions per site) along branches of the phylogeny as a proxy for

time.

Overall, 10.8% of IESs detected in *aurelia* species predate the divergence from *P. caudatum*

(referred to as 'Old' IESs in Fig 1), 79% were gained after the divergence of *P. caudatum*, but

before the radiation of the *aurelia* complex ('Intermediate' in Fig 1) and 10.2% are more recent.

The rate of IES gain varied widely over time: a burst of insertions occurred in the ancestral

branch leading to the *aurelia* clade, followed by a progressive slowdown in most lineages,

except in *P. sonneborni* where the rate of IES gain strongly increased again in the recent period

(18.8% of IESs detected in *P. sonneborni* are specific to that species). The IES gain rate has

remained substantial in *P. sexaurelia* and *P. tredecaurelia*, but has dropped to very low levels

in *P. tetraurelia/P.octaurelia* and in *P. pentaurelia/P. primaurelia* lineages, about 20 times

lower than in *P. sonneborni* or in the ancestral *aurelia* lineage (Fig 1). The rate of IES loss

appears to be more uniform along the phylogeny, with only 2 to 3-fold variation (Fig 1).

Recent waves of mobilization of IESs

The episodic bursts of IES gains that we observed in the phylogeny are reminiscent of the

dynamics of invasion by TEs. To test the hypothesis that IESs might correspond to TEs, we

searched for evidence of mobile IESs, i.e. IES sequences sharing significant sequence similarity

but inserted at different (non-homologous) loci. In a first step, we compared all IESs against

each other with BLASTN to identify clusters of similar IESs. Consistent with previous

observations made in *P. tetraurelia* [4], we found that the vast majority of IESs correspond to unique sequences in all MIC genomes, but a fraction of IESs are present in multiple copies (Fig 2A). In a second step, all clusters with ≥ 10 interspersed copies were manually inspected, to precisely delineate the boundaries of the repeated element and create a multiple alignment of full-length copies. We then used these representative multiple alignments to perform an exhaustive sequence similarity search based on HMM profiles over the entire IES dataset (see Materials and Methods). Among the hits, we distinguished two categories: 1) cases where the detected copy is located within the IES but does not include the extremities of the IES, and 2) cases where the extremities of the copy correspond precisely to the extremities of the IES. The first category probably corresponds to TEs that were inserted within a pre-existing IES (i.e. nested repeats). The second category corresponds to cases where the transposed element is the IES itself (i.e. mobile IESs). Overall, we detected 24 families with at least 10 copies of mobile IESs, totaling 7,443 copies of mobile IESs (Table 2).

Repeat family	Length (bp)	Nb. of mobile IESs	Number of mobile IESs per species								
			pca	pse	pso	ptr	ppe	ppr	pbi	poc	pte
FAM_2183	233	4153	0	4	3252	897	0	0	0	0	0
FAM_3	290	1783	0	0	0	15	344	321	766	146	191
FAM_2938	765	378	0	7	370	1	0	0	0	0	0
FAM_2317	768	331	0	82	140	109	0	0	0	0	0
FAM_2942	211	110	0	0	110	0	0	0	0	0	0
FAM_2334	214	106	0	17	89	0	0	0	0	0	0
FAM_2321	471	84	1	11	58	10	1	1	2	0	0
FAM_78	50	56	0	0	0	0	34	22	0	0	0
FAM_1402 (TIR *Thon)*	693	53	0	6	8	5	0	0	3	16	15
FAM_1257 (TIR *Merou)*	522	49	0	0	5	5	3	2	7	12	15

FAM_670	46	44	0	0	9	1	2	5	2	24	1
FAM_2649	762	40	0	0	16	24	0	0	0	0	0
FAM_1473	98	33	0	0	4	7	1	0	5	10	6
FAM_51	231	32	0	0	0	0	12	9	11	0	0
FAM_692	93	26	0	0	4	13	5	4	0	0	0
FAM_1294 (*Baudroie*)	1706	26	0	0	0	0	1	2	18	4	1
FAM_2314 (DDE)	3421	24	0	20	2	2	0	0	0	0	0
FAM_2802	32	22	0	0	2	20	0	0	0	0	0
FAM_3194	230	20	20	0	0	0	0	0	0	0	0
FAM_837	50	18	0	0	0	0	14	4	0	0	0
FAM_1165	77	15	0	0	0	0	1	1	0	12	1
FAM_2936	223	15	0	0	15	0	0	0	0	0	0
FAM_1259	231	14	0	0	0	0	1	0	13	0	0
FAM_3023	350	11	0	0	11	0	0	0	0	0	0
Total		7443	21	147	4095	1109	419	371	827	224	230

Table 2. Taxonomic distribution of mobile IESs.

Copies of mobile IESs were searched among IES sequences (see Materials and Methods). Detected copies were divided in two categories: nested copies (i.e. copies inserted within an IES, but not including the extremities of the IES) and *bona fide* mobile IESs (i.e. copies whose extremities correspond to the extremities of the IES). This table lists all families for which at least one species contains ≥ 10 copies of *bona fide* mobile IESs in its genome (see S4 Table for information on nested copies). Species codes: pso: *P. sonneborni*, ptr: *P. tredecaurelia*, pte: *P. tetraurelia*, pbi: *P. biaurelia*, poc: *P. octaurelia*, pse: *P. sexaurelia*, ppr: *P. primaurelia*, ppe: *P. pentaurelia*, pca: *P. caudatum*.

Four of these mobile IESs present homology with DNA transposons of the ITm superfamily previously identified in *P. tetraurelia* [4,5] (Table 2). FAM_2314 (3.4 kb) includes an intact open reading frame (ORF) encoding a DDE transposase. FAM_1294 (1.7 kb) is homologous

to *Baudroie*, a composite Tc1-mariner element, and includes an ORF with similarity to tyrosine-type recombinases. FAM_1402 (0.7 kb) and FAM_1257 (0.5 kb) correspond to non-autonomous elements, homologous to the terminal inverted repeats (TIR) of *Thon* and *Merou* respectively. The other families of mobile IESs do not match with any known TEs. Their relatively short lengths (32 bp to 765 bp) and the absence of similarity with any known protein, indicate that they most probably correspond to non-autonomous elements, mobilized by transposases expressed from active TEs.

The genomic distribution of mobile IESs within MAC-destined regions is similar to that of other IESs: most of the families are predominantly located within protein-coding regions (which represent ~70% of the MAC genome) (S4 Table). The only notable exceptions are FAM_1257, FAM_1402 and FAM_78 elements, which are under-represented within genes (S4 Table). In particular, FAM_78 elements are exclusively found in intergenic regions.

As explained previously, it is possible to date insertions for the subset of IESs located within well-conserved protein-coding regions. The vast majority (97.5%) of mobile IES copies that can be dated correspond to recent insertions (as compared to only 9.5% of recent insertions for the other IESs). FAM_3 is present in all genomes of the subclade A (Table 2), and 94% of dated insertions are shared by at least two species, which indicates that this element has been very active at the beginning of the radiation of this clade. For the other families of mobile IESs, more than 97% of insertion loci are species-specific. Thus, all the families of mobile IESs that we detected are relatively recent, posterior to the radiation of the *aurelia* clade. This most probably reflects the fact that more ancient families are difficult to recognize, because of the rapid divergence of IES sequences.

In agreement with previous analyses [4], we detected few cases of recent mobilization of IESs

in *P. tetraurelia* (although we detected more copies than reported by [4], owing to the higher

sensitivity of the sequence similarity search protocol used here). However, several other species

contain IES families with a high number of recently inserted copies (Table 2). The largest

family (FAM_2183) corresponds to a 233 bp-long non-autonomous element, for which we

detected a total of 4,126 copies in *P. sonneborni*, and 1,089 copies in *P. tredecaurelia*, among

which 20% are nested within IESs and 80% correspond to mobile IESs (Table 2, Fig 2B-D).

The high level of sequence similarity between all copies (average pairwise identity=72%; Fig

2) suggests that both species have been invaded recently by this mobile element. To assess

whether some insertions are ancestral to these two species, we analyzed copies inserted in well-

conserved coding regions (i.e. for which it is possible to check whether they are co-orthologous

or not; N=1,973 copies). We identified only two FAM_2183 elements inserted at co-

orthologous sites in both species. Thus, the quasi-totality of FAM_2183 copies result from

independent waves of insertion in the two lineages. Interestingly, FAM_2183 is absent from

other *Paramecium* species (except 4 copies in *P. sexaurelia*). It is important to note that *P.*

sonneborni and *P. tredecaurelia* belong to two distantly related subclades of the *aurelia*

complex (Fig 1). This patchy phylogenetic distribution therefore suggests that FAM_2183 has

been subject to horizontal transfers between *P. sonneborni* and *P. tredecaurelia* lineages. The

alternative hypothesis of vertical inheritance of the family since the last common ancestor of

the *aurelia* complex would imply that all copies of the family have been lost independently in

at least three lineages of subclade A. Furthermore, the phylogeny of this family is mainly star-

like, with *P. sonneborni* and *P. tredecaurelia* copies interspersed across the entire tree (Fig 2B).

This topology is incompatible with a scenario of vertical inheritance, in which copies from each

species would have been expected to form two distant clades. Interestingly, the tree topology

indicates that several events of horizontal transfer, not just one, occurred between these two

lineages. This finding is corroborated by four other families (FAM_2317, FAM_2321,

FAM_2649, FAM_2802) that are shared specifically by *P. tredecaurelia* and the *P.*

sonneborni/P. sexaurelia clade. These multiple events of horizontal transfer suggest an

important genetic flux between these two lineages. And indeed, besides these mobile IESs, we

also found many large genomic segments (up to 200 kb) that appear to have been introgressed

into MIC-specific regions of *P. sonneborni* genome, from a lineage closely related to *P.*

tredecaurelia (these results will be described in detail in another article). The hypothesis that

we propose is that these introgressed segments brought some active transposable elements (both

autonomous or non-autonomous), including some mobile IESs, which then were able to

proliferate within the *P. sonneborni* genome. The finding of two FAM_2183 copies present at

co-orthologous sites is *a priori* unexpected under this scenario. However, they both correspond

to nested copies. Thus, one possible explanation is that, by chance, these copies were inserted

independently in both lineages within pre-existing co-orthologous IESs. An alternative

hypothesis is that both nested insertions might have been transferred between species through

the introgression of larger genomic segments. Like most species of the *aurelia* complex, *P.*

sonneborni and *P. tredecaurelia* have a worldwide geographic distribution [26,27]. Our

observations indicate that despite the strong reproductive isolation between extant species [26],

horizontal transfers did occur recently between these genetically very distant lineages.

Fig 2. Proportion of IESs corresponding to repeated sequences, and phylogenetic

analysis of the largest family of mobile IESs (A) The vast majority of IESs correspond to

unique sequences, but a fraction of IESs are present in multicopies. For each species, all IESs

were compared against each other with BLASTN (with an E-value threshold of 10^{-5}). The

distribution of the number of BLAST hits per IES (excluding self-hits) is displayed for each

species. (B) The largest family of multicopy IESs (FAM_2183) includes 4,153 copies. We present here a phylogenetic tree of a subset of sequences (200 IESs from *P. tredecaurelia* in black, and 200 from *P. sonneborni* in red), randomly sampled from the entire FAM_2183 alignment (computed with PhyML [28]). The tree topology is mainly star-like, which indicates that most copies derive from several bursts of insertions. (C) Length distribution of the 4,153 FAM_2183 mobile IESs. (D) Sequence logo [29], based on the alignment of the entire FAM_2183 family. The information content reported in the logo was computed relative to the base composition of IESs, to take into account their high AT-content. All copies present a high level of sequence similarity (average pair-wise identity 72%) throughout their entire length, not just at their ends.

IES excision mechanism varies with IES age

Like any biological process, the excision of IESs during new MAC development is not 100% efficient [4,30]. For instance, the IES retention rate in *P. tetraurelia* MAC chromosomes is on average 0.8% in wild-type cells [31]. We observed that a substantial fraction of IESs have a much lower excision efficiency. In all *Paramecium* species, the proportion of 'weak' IESs (defined as IESs with more than 10% retention in wild-type cells) differs strongly among genomic compartments: from 0.7% on average for IESs located within genes (introns or exons), to 5.4% for IESs in intergenic regions (S10 Fig). This difference probably results from the fact that IESs with low excision efficiency are more deleterious, and therefore more strongly counter-selected, in genes than in intergenic regions. Interestingly, we also observed that within coding regions, the proportion of weak IESs is much higher for newly gained IESs (2.1% on average) than older ones (0.3%) (S10 Fig). This suggests that after a wave of insertions,

genomes accumulate changes in cis (within IESs) and/or in trans (in the IES excision

machinery) that progressively make these new IESs more efficiently excised, presumably in

response to the selective pressure against retention of IESs within coding regions.

In *P. tetraurelia*, functional analyses have revealed that different classes of IESs rely on

different excision pathways [31–33]. A large subset of IESs (70%) requires the histone H3

methyltransferase Ezl1 for excision [31]. A much smaller subset (7%), all among the of *EZL1*-

dependent IESs, also requires the Dcl2/3 proteins, which are necessary for the biogenesis of 25

nt long scnRNAs [31,34]. The remaining 30% of IESs require neither Ezl1 nor Dcl2/3 to

complete excision. Using published IES excision efficiency datasets upon silencing of *EZL1* or

DCL2/3 [31], we found that 92% of newly inserted *P. tetraurelia* IESs are sensitive to Ezl1, as

compared to 39% for old ones (Fig 3). Similarly, the proportion of Dcl2/3-dependent IESs

varies from 17% for new IESs to 3% for old ones. These observations suggest that newly

inserted IESs, like TEs themselves [35], initially depend on histone marks deposited by Ezl1

(and to some extent on the scnRNA pathway). Over time, histone marks and scnRNAs become

dispensable as IESs gradually acquire features that allow them to be efficiently excised.

Fig 3. Old and young IESs rely on different excision pathways.

The subset of IESs whose excision relies on histone marks or scnRNAs have been identified in

P. tetraurelia by measuring their IES retention score (IRS) upon silencing of *EZL1* and

DCL2/DCL3 [31]. Barplots represent the proportion of sensitive IESs (IRS>10%) according to

their age, for each silencing experiment. The age of an IES insertion is defined by the

phylogenetic position of the last common ancestor (LCA) of species sharing an IES at the same

site (New: *P. tetraurelia*-specific IES; Node *n*: the LCA corresponds to node number *n* in the

species phylogeny; Old: the LCA predates the *P. aurelia*/*P. caudatum* divergence). The

We also compared the length of IESs according to their age. As previously observed for *P. tetraurelia* [4], IESs have a characteristic length distribution with the same ~10 bp periodicity in all *aurelia* species (Fig 4A), likely reflecting structural constraints on the excision process [4,7]. We observed that the length distribution of IESs changes drastically over evolutionary time. For instance, in *P. sonneborni*, *P. tredecaurelia* and *P. tetraurelia*, the proportion of IESs in the first peak of the length distribution (< 35 bp) ranges from 1-10% for new IESs to 81-84% for old ones (Fig 4B), and similar patterns are observed in all other *aurelia* species (S11 Fig). In *P. caudatum*, the overall length distribution is shifted towards shorter IESs (71% in the first peak, compared to 35% in *aurelia*; Fig 4A). This suggests that this lineage has not been subject to IES insertion waves for a long period of time, in agreement with the paucity of recognizable mobile IESs in that genome (Table 2).

Fig 4. Length distribution of IESs according to their age.

(A) Comparison of the length distribution of IESs in *P. caudatum* (N=8,172 IESs) and in species from the *aurelia* clade (N=392,082 IESs). The fraction of IESs present within each peak of the distribution is indicated for the first 10 peaks. (B) Comparison of the length distribution of IESs according to their age (for the subset of datable IESs located in coding regions). The age of IES insertions is defined as in Fig 3. Results from other species are presented in S11 Fig.

Genomic distribution of IESs according to their age

Because of the rapid divergence of non-coding sequences, it is generally not possible to assess homology among IES insertion sites located in intergenic regions, and hence it is not possible

to date them directly. We therefore used the length of IESs as a rough proxy for their age, to

investigate their genomic distribution over time. We observed that in all *aurelia* species, long

IESs (> 100 bp, presumably young) are uniformly distributed across genomic compartments

(introns, coding regions and intergenic regions) (S12 Fig). Conversely, short IESs (<35 bp,

presumably older) are enriched in coding regions (on average, 81% of short IESs in coding

regions, vs. 70% expected; S12 Fig). This suggests that IESs located within intergenic regions

have a shorter lifespan than those located in coding regions.

Evidence that some IESs are functional

A large majority of detected IESs predate the divergence of the *aurelia* clade (Fig 1). Because

of the rapid evolution of non-coding sequences, co-orthologous IESs from different species are

generally too divergent to be recognized by sequence similarity search. Yet the comparison of

all sequences against each other revealed several interesting exceptions. Overall, we identified

69 clusters of co-orthologous IESs conserved across at least 5 of the 8 species of the *aurelia*

clade, and showing at least 70% identity between sequences from subclades A and B (87%

identity on average). Each of these 69 IESs correspond to unique sequences (we did not detect

any interspersed homologs within genomes, only co-orthologs). These highly conserved IESs

are similar to other IESs in terms of length (mean=75 bp) or genomic distribution (79% within

protein-coding genes, 21% in intergenic regions). Their high levels of sequence conservation

indicate that they are subject to strong selective constraints, and hence that they have a function,

beneficial for *Paramecium*. By definition, IESs are absent from the MAC genome so they

cannot be expressed in vegetative cells. However, they can potentially be transcribed during

the early development of the new MAC, before IES excision occurs [32,36]. To gain insight

into their possible functions, we analyzed the transcription of conserved IESs using

polyadenylated RNAseq data from autogamy time course experiments in *P. tetraurelia* [37].

Among the 56 families of highly conserved IESs present in *P. tetraurelia*, 10 (18%) are

transcribed at substantial levels (>1 RPKM) during autogamy (as compared to 0.8% for other

IESs) (S5 Table). One of these IESs (~800 bp-long) contains a gene encoding a putative DNA-

binding protein, well conserved in all species of the *aurelia* clade and expressed at high levels

during the early stages of autogamy (Fig 5A,C). Several paralogs of this gene are present in the

MAC genome, but it is the only member of this family to be located within an IES. We

hypothesize that this copy might have originated by retrotransposition of one of the MAC

paralogs, and that its insertion within a pre-existing IES might have been exapted, as a way to

restrict its expression specifically to early MAC development.

Fig 5. Examples of highly conserved co-orthologous IESs. (A) Cluster 4968 corresponds to

a set of co-orthologous IESs that contain a gene (blue), oriented in the opposite direction with

respect to the gene (blue) in which the IES is inserted (yellow). The IES-embedded gene

encodes a protein belonging to a family of CENP-B DNA-binding proteins, and is likely of

cellular origin. RNA-Seq data (shown here for the *Paramecium tetraurelia* member of this

cluster: IES.PTET.51.1.16.324097) show that this gene is expressed at meiosis, with mRNA

steady state levels strongly decreasing before most genome-wide IES excision occurs [37]. **(B)**

Cluster 9405 corresponds to a set of co-orthologous IESs that contain the promoter and the start

codon of a gene (blue) encoding a protein of unknown function. The IES is drawn in yellow.

RNA-Seq data (shown here for one *Paramecium tetraurelia* member of this cluster:

IES.PTET.51.1.28.104046) indicate that this gene is expressed at the time genome

rearrangements occur, with a peak in steady-state mRNA level at T11, when IES excision peaks

[37]. (D) Nucleotide alignment of co-orthologous IESs from cluster 9405. The ATG start codon

and first few amino acids encoded by the IES are shown below the alignment. (C) Alignment

of the proteins encoded by IESs from cluster 4968. [IES IDs in the alignments, from top to

bottom (C): IES.PPENT.87.1.080.374281, IES.PPRIM.AZ9-3.1.072.352290, IES.PBIA.V1_4.1.0053.93462.rc, IES.POCT.138.1.076.348969, IES.PTET.51.1.61.348482, IES.PTRED.209.1.7180000129326.89420.rc, IES.PTRED.209.1.7180000129337.349035, IES.PBIA.V1_4.1.0022.293968, IES.PPENT.87.1.047.444829, IES.PPRIM.AZ9-3.1.042.468625, IES.PSEX.AZ8_4.1.047.365473, IES.PSEX.AZ8_4.1.067.78433.rc, IES.PSON.ATCC_30995.1.070.107189.rc, IES.POCT.138.1.047.443628, IES.PTET.51.1.28.104046.rc. (D): IES.PSON.ATCC_30995.1.014.460171, IES.PSEX.AZ8_4.1.004.493121, IES.PTRED.209.1.7180000129358.256806, IES.PTET.51.1.16.324097, IES.POCT.138.1.015.320413, IES.PBIA.V1_4.1.0089.86544, IES.PPENT.87.1.022.397764, IES.PPRIM.AZ9-3.1.019.378988.]

All other highly conserved IESs are much shorter (< 300 bp), most probably too short to encode proteins. But we found examples suggesting that some of them contribute to the regulation of the expression of their host gene. For instance, we identified a conserved IES located at the 5' end of a gene of unknown function, encompassing the transcription start site and the beginning of the first exon (including the 5'UTR and the first 9 codons). The excision of the IES during MAC development leads to the loss of the initiation codon and of the promoter region, and thereby to the silencing of this gene in vegetative cells (Fig 5B,D). These examples illustrate that recruitment of the IES excision machinery during evolution to contribute new functions beneficial for *Paramecium*.

Discussion

 A majority of *Paramecium* IES insertions result from the transposition of mobile IESs

To explore the evolutionary origin of IESs, we analyzed the MIC genomes of eight species of

the *P. aurelia* complex, and of an outgroup species, *P. caudatum*. Unexpectedly, we discovered

that the MIC genomes of *P. caudatum* strains are at least one order of magnitude larger than

those of *P. aurelia* species (~1,600 to 5,500 Mb vs ~110-160 Mb). The sequencing of *P.*

caudatum My43c3d revealed that its huge MIC genome size is caused by the amplification of

two major satellite repeats, which represent 71% of its MIC-limited genome (S4 Fig). The high

variability of genome sizes across the *P. caudatum* lineage makes this clade an attractive model

system to study the possible phenotypic consequences of genome size variations within a

species.

All the *Paramecium* MIC genomes we sequenced present a high density of IESs in MAC-

destined sequences: from 0.5 IES per kb in *P. caudatum* up to 1 IES per kb in *P. sonneborni*

(Table 1). The vast majority of these IESs (83% on average) are located within genes, as

expected given the very high gene density in MAC genomes (S3 Table). In *aurelia* species,

there are on average 0.95 IESs per protein-coding gene. The IES density varies among genes,

but overall, ~50% of the ~40,000 genes contain at least one IES. Moreover, the analysis of

MIC-specific regions that are occasionally retained in the MAC (MAC-variable regions)

revealed similar IES densities (S2 Table), which suggests that, in addition to IESs located in

MAC-destined regions, many other IESs are located within MIC-specific regions.

To explore the origin and evolution of these tens of thousands of IESs, we sought to identify

homologous IESs across the 9 *Paramecium* species. We distinguished two categories of

homologs: interspersed homologs (i.e. similar IES sequences located at non-homologous loci)

and co-orthologous IESs (i.e. IESs deriving from a single ancestral insertion event at a given

genomic site). We first compared all IES against each other to identify sets of homologs based

on sequence similarity, within genomes or across species. The vast majority of IESs correspond

to sequences that are unique within each genome (Fig 2). We identified only a handful of co-

orthologous IES sequences that are conserved in sequence across the *aurelia* complex (see

below). Apart from these rare cases, most IES sequences evolve rapidly, as expected for non-

coding sequences. It should be noted that species from the *aurelia* complex are genetically quite

divergent (S6 Table). For instance, the average synonymous divergence (measured in

orthologous protein-coding genes) is 0.95 substitutions/site between species of the A and B

subclades. At this evolutionary scale, in the absence of selective pressure, IES homologs are

expected to be far too diverged to be recognizable by sequence similarity. Thus, homologs

(interspersed or co-orthologs) that can be detected by sequence similarity essentially

correspond to IESs whose last common ancestor is relatively recent (more recent than the A

and B subclades).

However, it is also possible to identify co-orthologous IESs based on their shared position

within multiple alignments of homologous genes. Thus, for the subset of IESs located in coding

regions, we were able to infer rates of IES gain and loss across the species phylogeny (Fig 1).

Overall, about 90% of IESs detected in *aurelia* species predate the radiation of that clade, but

fewer than 10% are shared with *P. caudatum*. Thus, the vast majority of *aurelia* IESs result

from a major wave of IES gains that occurred after the divergence of *P. caudatum,* but before

the radiation of the *aurelia* complex. Similarly, 80% of IESs detected in *P. caudatum* are

specific to that lineage, which implies that multiple independent events of massive IES

invasions occurred during evolution.

The burst of IES gains at the base of the *aurelia* clade was followed by a progressive slowdown

in most species, except in the *P. sonneborni* lineage, which has been subject to a second wave

of IES insertions (Fig 1). Interestingly, the comparison of IES sequences revealed thousands of

interspersed homologous copies, resulting from the recent and massive mobilization of a small

number of IESs. Several families of mobile IESs present homology with known ITm

transposons, and some of them encode transposases. But most mobile IESs do not appear to

have any protein-coding potential, and therefore likely correspond to non-autonomous

elements, whose mobility depends on the expression of active transposons. The number of

detectable (recently mobilized) IES copies varies widely across species (Table 2). For instance,

mobile IESs have been very active in the *P. sonneborni* lineage (4,095 copies), much more than

in its sister lineage, *P. sexaurelia* (147 copies). Thus, mobile IESs account for at least 20% of

the difference in IES number between these two species (Fig 1). Given the rapid evolution of

IESs, interspersed homologs that can be detected probably represent only the tip of the iceberg

of mobile IESs. Overall, we found a strong correlation ($R^2=0.86$, $p=8 \times 10^{-4}$) between the number

of mobile IES copies detected in each species (Table 2) and the rate of IES gain along

corresponding branches of the phylogeny (Fig 1), which suggests that most gains result from

transposition.

Interestingly, the five most active families in *P. tredecaurelia* all show the signature of

horizontal transfer with the distantly related *P. sonneborni* lineage (Table 2). This is the case

of the largest family that we identified (FAM_2183: 3,252 and 897 copies in each species,

respectively; Fig 2). This pattern is reminiscent of the typical life cycle of many DNA

transposons: when a new element enters a genome, it is initially very active and produces a

wave of insertions. Its activity then progressively slows down, largely because defense

mechanisms become more efficient in the host genome. In the long-term, DNA transposons

escape extinction only if they can occasionally be transmitted to a new host [38]. Thus, the

variation of IES insertion rates that we observed in the *Paramecium* phylogeny fits very well

with the dynamics of TEs: rare episodes of massive proliferation (promoted by horizontal

transfer of an active copy to a new naive host), followed by progressive slowdown of

transposition activity.

TEs are not the unique source of IES gains. Mutations in MAC-destined regions can generate

sequence motifs that are recognized by the IES excision machinery, and thereby create new

IESs. There is indeed evidence that cryptic IES signals occasionally trigger the excision of

MAC-destined sequences [4,30], and that some IESs originated from MAC-destined sequences

[12,13]. However, our results suggest that the vast majority of IESs correspond to

unrecognizable fossils of mobile elements – as initially proposed by Klobutcher and Herrick

[10,11].

Indirect evidence of mutational burden caused by IES invasions

In all *Paramecium* species, we observed a deficit of IESs in highly expressed genes (S7 Fig).

As previously reported in *P. tetraurelia* [4], this pattern most probably reflects selective

pressure against IES insertions within genes. Indeed, the IES excision machinery (like any other

biological machinery) is not 100% efficient: a small fraction of IES copies are retained in the

MAC or subject to imprecise excision [30]. Typically, the average IES retention rate in MAC

chromosomes is 0.8% in *P. tetraurelia* [31]. For IESs located within genes, such excision errors

are expected to have deleterious consequences on fitness, in particular for genes that have to be

expressed at high levels [4]. And indeed, in agreement with the hypothesized selective pressure

against IESs within genes, we observed that the proportion of 'weak' IESs (i.e. IESs with a

relatively high retention frequency) is much lower in genes than in intergenic regions (S10 Fig).

Despite their selective cost, weakly deleterious IES insertions can eventually become fixed by

random genetic drift. Once fixed, the fitness of the organism will depend on its ability to

properly excise the IES during MAC development. Over time, selection should favor the

accumulation of substitutions and indels that increase the efficiency of IES excision. Indeed,

we did observe that the proportion of weak IESs decreases with their age (S10 Fig).

Interestingly, older IESs, which are also shorter, are less dependent on the Ezl1 and Dcl2/3

proteins (Fig 3). This suggests that after their insertion, IESs progressively acquire features that

make them more efficiently excised, by a pathway that requires neither scnRNAs nor histone

marks [31].

Exaptation of the IES excision machinery

As mentioned previously, orthologous sequences that evolve neutrally are not expected to

display any significant similarity between species of subclades A and B (synonymous

divergence ~0.95 substitution per site; S6 Table). Yet, we identified 69 clusters of co-

orthologous IESs displaying a high level of sequence conservation across the entire *aurelia*

complex (on average 87% identity between the closest pair of homologs from subclades A and

B). Their level of conservation implies that these IESs are subject to strong purifying selection,

and hence, that they fulfill a function that contributes to the fitness of *Paramecium*. Notably,

we identified one IES that contains a protein-coding gene (Fig 5A,C). This gene is expressed

during the early stages of autogamy, likely from the new developing MAC, before IES excision

(Fig 5A). Interestingly, 18% of the conserved IESs are transcribed during autogamy (as

compared to 0.8% for other IESs). Most conserved IESs are too short to encode proteins, but

they may contribute to gene regulation (e.g. Fig 5B,D). Given the enrichment of conserved IESs

in genes expressed during early autogamy, it is tempting to speculate that these IESs may play

a role in controlling the IES excision machinery itself. Indeed, this machinery must be tightly

regulated to ensure that all IESs are efficiently excised, while limiting off-target excision of

MAC-destined regions, which occurs occasionally in MAC chromosomes [4,30]. Thus,

developmental disruption of genes encoding IES excision factors by the excision machinery

may provide a simple regulatory feedback loop to decrease the activity of the IES excision

machinery as soon as a large fraction of IESs have been excised: if a given IES drives the

expression of a protein factor that is essential for IES excision, then this process is progressively

interrupted by the removal of this IES during MAC development. More generally, such IESs

may provide an exquisite developmental process to regulate DNA elimination events and /or

MAC differentiation.

Given that each of these 69 IESs is conserved across species and unique within genomes, it

seems *a priori* unlikely that they derive from transposable elements. We propose that these

conserved IESs may in fact originate from MAC-destined segments. Indeed, mutations within

genes can create sequence motifs recognized by the IES excision machinery, and thereby trigger

the deletion of functional elements during MAC development. Such mutations are expected to

be generally deleterious, and hence to be counter-selected. But occasionally, they might have

been positively selected in some genes because they provide a mean to modulate their

expression during MAC development or to generate phenotypic diversity among clones, as

previously observed for IESs overlapping mating-type genes in many species [12,13]. This

scenario would notably explain the presence of conserved IESs overlapping with cellular genes

(e.g. Fig 5B). Thus, we propose that the IES excision machinery, which evolved initially to

ensure the efficient removal of selfish genetic elements from the MAC genome, has been

exapted during evolution as a new way to regulate the expression of some genes.

IES losses

In all species, we observed that the length of IESs is negatively correlated with their age (Fig 4, S11 Fig). This pattern is similar to that observed in other eukaryotes, where fixed copies of TEs tend to shrink over time, due to the accumulation of small deletions [39]. For IESs located in non-coding region, this progressive shrinking can ultimately lead to their disappearance. Furthermore, these IESs can also be lost by transformation into a MAC-destined sequence (e.g. via mutations the TA dinucleotides). However, for IESs located within exons, losses can only occur by precise complete deletions that leave the open reading frame intact. We did observe such cases of precise loss (S9 Fig). One possible mechanism is that the IES excision machinery, which is normally at work during MAC development, might occasionally operate within the MIC or zygotic nucleus. An alternative hypothesis is that IESs might be lost from the MIC lineage by gene conversion, through homologous recombination with MAC-derived DNA fragments. Interestingly, this scenario might explain cases where we observed concomitant losses of neighboring IESs (see e.g. IES 5 and 6 in S9 Fig). Further studies will be needed to determine the mechanisms underlying precise IES loss. The rate of IES loss has remained quite stable and relatively low across the phylogeny (Fig 1). Conversely, the rate of IES gains has been much more erratic, characterized by episodic waves of insertions, during which the IES gain rate largely exceeded the loss rate (Fig 1). In the end, the number of IESs reflects the balance between gain and loss rates. Thus, the large number of IESs in *Paramecium* can simply be explained by massive invasions of mobile IESs, followed by periods of lower activity, during which IES copies progressively diverge, and occasionally get lost by deletion from the MIC.

Scenario for the evolution of IESs

In most organisms, gene regulatory elements and coding regions constitute a no man's land for TEs, because insertions that disrupt gene function are strongly counter-selected. But in some

ciliates, it is possible for mobile elements to proliferate within genes in the MIC genome, as long as they are efficiently and precisely excised during the development of the MAC genome, before genes start to be expressed. DNA transposons encode transposases that allow their mobilization by a 'cut-and-paste' process. Generally, the excision step leaves a few nucleotides at the original insertion site, but one peculiarity of piggyBac transposases is that they can excise copies precisely, without leaving any scar [40]. This feature may have predisposed piggyBac to extend its niche to genic regions in ciliates. We speculate that the very first proto-IESs corresponded to piggyBac elements whose transposase was domesticated to target the developing MAC. As soon as several copies of these proto-IESs have been fixed within genes, then the host organism has become dependent on the activity of the piggyBac transposase to ensure that all these copies are precisely excised from its MAC. This selective pressure would have perfected the domestication of the piggyBac transposase by its host, and then, progressively, the evolution of the other components that contribute to the efficient excision of proto-IESs. Once the IES excision machinery in place in the ancestral *Paramecium* lineage, other families of TEs (including non-autonomous elements) could hijack the machinery and in turn exploit this intragenic niche, eventually creating the tens of thousands of IESs found in present-day *Paramecium* genes. The first steps of this scenario remain speculative, since there are no recognizable traces of piggyBac-related IESs in present-day genomes. But the discovery of thousands of mobile IESs directly demonstrates the major contribution of TEs to the expansion of the IES repertoire.

The coexistence of MAC and MIC is a common feature of all ciliates, yet they do not all contain such a high density of IESs in coding regions. Notably, there are ~12,000 IESs in the germline genome of *Tetrahymena thermophila* (~0.1 IES per kb of MAC-destined sequence), but only 11 of them are located within coding regions [41]. These exonic IESs differ from other IESs by

their strongly conserved terminal inverted repeats ending with 5'-TTAA-3', the target site of

piggyBac transposons. They are excised precisely (restoring a single TTAA) by two

domesticated piggyBac transposases, Tpb1 and Tpb6, which may thus have retained the

cleavage specificity of their transposon ancestor [42,43]. We analyzed these 11 exonic IESs: 8

of them are inserted in protein-coding regions that are not conserved in *Paramecium*, and the

other 3 are inserted at sites that do not contain IESs in *Paramecium*. There is therefore no

evidence for shared exonic IESs between *T. thermophila* and *Paramecium*. The vast majority

of the ~12,000 *T. thermophila* IESs are excised by another domesticated piggyBac transposase,

Tpb2 [44]. Although Tpb2 retains the cleavage geometry of piggyBac transposases, producing

staggered double-strand breaks with 4-nt 5' overhangs [44], it has lost almost all sequence

specificity and is thought to be recruited at IES ends by chromatin marks [45]. As a result,

several possible cleavage sites are usually present at IES ends and the rejoining of flanking

sequences generates microheterogeneity in the MAC sequence [41], which explains why Tpb2-

dependent IESs are restricted to introns and intergenic regions [41]. It is important to note that

Tpb2 is an essential gene in *T. thermophila* [44], suggesting that genome-wide retention of IESs

in the MAC is still highly detrimental. Interestingly, phylogenetic analyses indicate that the

Paramecium endonuclease PiggyMac (Pgm) and Tpb2 are more closely related to each other

than to Tpb1 or Tpb6, and may even be orthologs [7]. In the case of Pgm, however, sequence

specificity was relaxed only for the two distal positions of the 4-nt cleavage sites, and the central

TAs remain a strict requirement for IES excision in *Paramecium*. Although piggyBac

transposons are completely absent from the present-day *Paramecium* germline, this

evolutionary solution may have been favored because it also allowed for precise excision of

Tc1/mariner insertions, which in turn would have allowed continuous accumulation of

insertions within exons [4].

Importantly, the fact that a mechanism of precise excision exists in *T. thermophila* (via Tpb1

and Tpb6) raises the question of why intragenic IESs are not more abundant in its genome. A

similar question arises from the distribution of introns in eukaryotes: why are introns very

abundant in some lineages but rare in others (e.g. ~7 introns per gene in vertebrates vs ~0.04 in

hemiascomycetous yeast)? Part of the explanation may reside in the fact that, because of

population genetic forces, some lineages are more subject to random genetic drift than others,

and therefore are more permissive to invasion by weakly deleterious genetic elements [46,47].

It is also possible that the abundance of intragenomic parasites is strongly affected by

contingency – rare events of massive invasion, followed by long periods during which copies

are lost at a slow rate.

Parallel scenario for the evolution of IESs and spliceosomal introns

The above scenario for the origin of IESs in *Paramecium* presents an interesting parallel with

the one proposed for the evolution of spliceosomal introns in eukaryotes. Indeed, it had long

been postulated, based on similarities in biochemical processes, that spliceosomal introns

derive from mobile elements (group II self-splicing introns) [48]. In eukaryotes, the spread of

introns in protein-coding genes has been facilitated by the fact that transcription and translation

occur in separate compartments, thus offering the opportunity for these mobile elements to be

excised from the mRNA in the nucleus without interfering with its translation in the cytoplasm

[48] – like IESs, which are excised from genes before they get expressed in the MAC. Once the

first introns were established, selection drove the emergence of host factors contributing to the

efficiency of the splicing process, which progressively led to the evolution of the modern

spliceosome [49]. In turn, the existence of the spliceosome released the requirement for introns

to maintain their self-splicing activity [50], and allowed other TEs to hijack this machinery.

The recent discovery of non-autonomous DNA transposons that generated thousands of introns

in genomes of algae directly demonstrated that mobile elements are a major source of new

introns [51]. Like IESs, introns represent a burden for their host, because of errors of the

splicing machinery [46,47,52–54], but they also contributed to innovations. Indeed, the

spliceosome has been exapted during evolution to produce alternative splicing, which not only

contributed to diversify the protein repertoire [55] but also allowed new modes of post-

transcriptional regulation gene expression [56], in particular in genes that encode splicing

factors [57,58]. This pattern is reminiscent of highly conserved IESs that we uncovered in

Paramecium lineages, which appear to be particularly enriched within genes that are expressed

during early MAC development.

In summary, the evolution of the nuclear envelope opened the way for introns to invade genes

in eukaryotes [48], and likewise, the separation of somatic and germline functions between the

MIC and the MAC offered the possibility for selfish genetic elements to invade genes in ciliates.

Genetic conflicts between these selfish elements and their host genome resulted in the evolution

of complex cellular machineries (the spliceosome, the IES excision machinery), which, in the

short term, reduced excision errors, but in the long term facilitated proliferation of these

elements within genes. The paradigm of intragenomic parasites [59–61] provides a simple and

powerful explanation for the "raison d'être" of these mysterious pieces of non-coding DNA

that interrupt genes.

Materials and Methods

Cells and cultivation

All experiments were carried out with the *Paramecium* strains listed in Table 1. *Paramecium aurelia* cells were grown in a wheat grass powder (WGP, Pines International, USA) infusion medium bacterized the day before use with *Klebsiella pneumoniae* and supplemented with 0.8 mg/L of β-sitosterol (Merck). Cultivation and autogamy were carried out at 27 °C. Monoclonal cultures of the *P. caudatum* cells were grown in a 0.25% Cerophyl infusion inoculated with *Enterobacter aerogenes* at 22°C [62].

Micronucleus-enriched preparation

To purify the MICs from vegetative cells, we used the same strategy as the one previously published [5,19], with some optimization for the sorting steps. For *Paramecium aurelia*, transgenic cells expressing a micronuclear (MIC)-localized version of the Green Fluorescent Protein (GFP) were obtained by microinjection of the vegetative macronucleus with the *P. tetraurelia* CenH3a-GFP plasmid, described in [63]. In the transformed clones, GFP was exclusively found in the MICs and the transformed clones were selected for their GFP signal/noise ratio. Viability of the sexual progeny after autogamy of the transformed clones was systematically monitored to make sure that the presence of the transgene did not impair the functionality of the MICs. A MIC-enriched preparation was obtained from approximately 3 L of exponentially growing vegetative cells after fractionation and Percoll density gradient centrifugation as described in [5] and kept at -80°C until further use.

A slightly different procedure was used for *Paramecium caudatum* cells, which were not transformed with the CenH3a-GFP transgene. The MICs of *P. caudatum* strain My43c3d (used for genome sequencing) were purified with a protocol modified from [64]. Briefly, 3L of a starved culture (~600 cells/mL) were filtered through 8 layers of gauze and concentrated by

centrifugation in pear-shaped centrifuge tubes. Packed cells were transferred to a 250 mL cell

culture flask, resuspended in 150 mL sterile Eau de Volvic and incubated over night at 22°C.

All subsequent steps were performed at 4°C or on ice. The overnight culture was again

concentrated by centrifugation and the cell pellet was resuspended and washed in 0.25 M TSCM

buffer (10 mM Tris-HCl, pH 6.8, 0.25 M sucrose, 3 mM CaCl2, 8mM MgCl2) [65]. After

centrifugation for 3 min at 100 g, pelleted cells were resuspended and incubated for 5 min in

10 mL 0.25M sucrose-lysis buffer (10 mM Tris-HCl, pH 6.8, 0.25 M sucrose, 3 mM CaCl2,

1mM MgCl2, 0.1% Nonidet-P40, 0.1% Na-deoxycholate). The cell suspension was centrifuged

for 2 min at 500 g and the packed cells were lysed in 1 mL of 0.25M sucrose-lysis buffer by

about 10-20 strokes on a vortex machine. Lysed cells were washed in 14 mL of 0.25 M TSCM

buffer and centrifuged for 1 min at 100 g. The supernatant (containing the MICs) was

centrifuged for 10 min at 1,500 g and the pellet was resuspended in 8 mL of 60% Percoll. This

suspension was centrifuged for 15 min at 24,000 g in a fixed-angle rotor and the micronuclei

formed a diffuse band near the middle of the centrifuge tube. This MIC containing layer was

carefully removed with a pipette in about 2 mL, diluted with 10 mL of 0.25 M TSCM buffer

and pelleted by centrifugation for 10 min at 1,500 g. The MIC pellet was resuspended in 100

µL of 0.25 M TSCM buffer, carefully mixed with 50 µL of 50% glycerol and kept at -80°C

until further use.

The MICs of the other *P. caudatum* strains were purified with a similar protocol, but omitting

the Percoll step and replacing it with centrifugation across a sucrose cushion. Lysed cells were

resuspended in 9 mL of 0.25 M TSCM buffer and this suspension was carefully layered on top

of a sucrose cushion consisting of 2 mL of 1.6 M TSCM buffer and 2 mL of 0.9 M TSCM

buffer and centrifuged in a swinging bucket rotor for 10 min at 300 g with lowest acceleration

and braking levels. Depending on the strain, the micronuclei accumulated at the bottom of the

0.25 M or 0.9 M TSCM cushion and were removed by careful pipetting of the respective phases

to new 15-mL tubes. MIC- containing suspensions were diluted with 0.25 M TSCM buffer,

centrifuged for 10 min at 1,500 g and the MIC pellets were subsequently treated as described

above.

Quantification of MIC DNA content by flow cytometry

We measured the absolute DNA content in the nuclei with propidium iodide, a fluorophore that

is insensitive to differences in base composition, and compared DNA content of MIC-enriched

preparations to a standard (tomato nuclei) of known genome size (see S1 Data). MIC-enriched

samples were thawed on ice, diluted 1/5 to 1/10 in washing buffer (0.25 M sucrose; 10 mM Tris

pH 7.4; 5 mM MgCl2; 15 mM NaCl; 60 mM KCl; 0.5 mM EGTA) and stained on ice with

propidium iodide at 100 µg/mL final concentration. We used Tomato nuclei obtained from

Montfavet 63-5 hybrid F1 seeds as internal standards of known genome size. Tomato nuclei

were obtained from 1 cm^2 of young leaves chopped in a Petri dish with a scalpel. 800 µL of a

modified Galbraith buffer [66], containing 45 mM MgCl2, 30 mM Sodium-Citrate and 20 mM

MOPS pH 7.0, 40 µg/mL RNAse A, 0.1% Triton X-100, 5 mM sodium metabisulfite ($S_2O_5Na_2$)

was added. The nuclei were collected by pipetting, filtered on 70 µm mesh, and stained on ice

with propidium iodide at 100 µg/mL final concentration.

The samples were analyzed on a CyanADP Cytomation analyzer from Beckman-Coulter

equipped with 3 lasers: 405 nm, 488 nm and 635 nm. Fluorescence intensity (PE signal in pulse-

height) of the nuclei was measured at 575/25 nm, after excitation with the 488 nm laser. Results

are deduced from 2C nuclei in individuals considered diploid and are given as C-values [67].

The ratio of fluorescence intensity of 2C-nuclei from sample and standard allows calculation

of genome size. C corresponds to the nuclear genome size (the whole chromosome complement

with chromosome number n), 1C and 2C being, respectively, the DNA contents of the haploid

(n) and diploid (2n) sets of chromosomes. The haploid nuclear DNA content is expressed in

picograms or million base pairs, where 1 pg = 978 Mbp [68], considering Tomato 2C DNA

(pg) =1.99, according to [69]. The raw data and calculations are provided in S1 Data.

Micronucleus sorting by flow cytometry and flow imaging

To sort the MICs, the MIC-enriched samples were submitted to flow cytometry. Sorting by

flow cytometry allowed the separation of the small, diploid MICs from the highly polyploid

MAC and the bacteria abundant in *Paramecium* cultures. *P. aurelia* MICs were sorted based

on the SSC, FSC, DAPI (DNA staining), and GFP signals. *P. caudatum* MICs, which are bigger

than *aurelia* MICs, could be sorted based on their SSC, FSC, and DAPI signals, without the

use of a MIC-specific GFP fluorophore. Quality control was performed by flow cell imaging,

using the ImageStreamX (Amnis/Merck Millipore) imaging flow cytometer, as previously

described [5]. The MICs represented >99% of the sorted sample, except for *P. sonneborni*

(97%). An example of sorting is shown in Fig S1.

Genomic DNA extraction and sequencing

For MAC DNA sequencing, genomic DNA was extracted from vegetative *Paramecium* cell

culture after centrifugation and washes with Tris 10mM pH 7.4. For MIC DNA sequencing,

DNA was extracted from the sorted MIC samples. The cell or nuclei pellet was treated with 3

volumes of proteinase K solution (0.5 M EDTA pH 9; 1% N-lauroylsarcosine; 1% SDS; 1

mg/mL proteinase K) at 55 °C overnight. Genomic DNA was extracted with Tris–HCl-phenol

pH 8 with gentle agitation followed by dialysis against TE (10 mM Tris–HCl; 1 mM EDTA,

pH 8) 25% ethanol then against Tris 1 mM pH 8. An RNAse A treatment was performed on

MAC DNA, followed by phenol extraction and dialysis as described above. DNA concentration

was quantified using QuBit High sensibility kit (Invitrogen) and stored at 4 °C.

As the amounts of DNA extracted from the MIC are too low (30-50 ng), only an overlapping paired-end library could be prepared for *de novo* sequencing. Briefly, 30-50 ng of MIC DNA were sonicated using the E210 Covaris instrument (Covaris, Inc., USA) in order to generate fragments mostly around 500bp. Illumina libraries were then prepared using the NEBNext DNA Sample Prep Master Mix Set (New England Biolabs, MA, USA) and DNA fragments were PCR-amplified using Platinum Pfx DNA polymerase (Invitrogen) and P5 and P7 primers. Amplified library fragments of roughly 500 – 600 bp were size selected on 2% agarose gel. Libraries traces were validated on a Agilent 2100 Bioanalyzer (Agilent Technologies, USA) and quantified by qPCR using the KAPA Library Quantification Kit (KapaBiosystems) on a MxPro instrument (Agilent Technologies, USA). The libraries were sequenced using 251 base-length read chemistry in a paired-end flow cell on the Illumina HiSeq2500 sequencer (Illumina, USA).

For the MAC genomes, an overlapping paired-end library as described above and four additional mate-pair libraries (about 5 kb, 8 kb, 11 kb and 13 kb) were prepared following Nextera protocol (Nextera Mate Pair sample preparation kit, Illumina). Each library was sequenced using 100 base-length read chemistry on a paired-end flow cell on the Illumina HiSeq2000 (Illumina, USA).

Information about the sequencing data generated for this study is available in S7 Table.

RNA extraction and sequencing

For the purpose of gene annotation, we sequenced mRNAs from vegetative cells (S7 Table). 400 mL cultures of exponentially growing cells at 1000 cells/mL were centrifuged and flash-frozen in liquid nitrogen prior to TRIzol (Invitrogen) treatment, modified by the addition of glass beads for the initial lysis step.

RNA-Seq library preparation was carried out from 1 μg total RNA using the TruSeq Stranded

mRNA kit (Illumina, San Diego, CA, USA), which allows mRNA strand orientation (sequence

reads occur in the same orientation as anti-sense RNA). Briefly, poly(A)+ RNA was selected

with oligo(dT) beads, chemically fragmented and converted into single-stranded cDNA using

random hexamer priming. Then, the second strand was generated to create double-stranded

cDNA. cDNAs were then 3'-adenylated, and Illumina adapters were added. Ligation products

were PCR-amplified. Ready-to-sequence Illumina libraries were then quantified by qPCR using

the KAPA Library Quantification Kit for Illumina Libraries (KapaBiosystems, Wilmington,

MA, USA), and library profiles evaluated with an Agilent 2100 Bioanalyzer (Agilent

Technologies, Santa Clara, CA, USA). Each library was sequenced using 101 bp paired end

read chemistry on a HiSeq2000 Illumina sequencer.

MAC Genome assembly

The MAC genomes sequenced for this project were all assembled according to the following

steps.

First, long Illumina reads were obtained from 250 bp overlapping paired-end reads sequenced

from ~450 bp fragments. The reads were fused with fastx_mergepairs, an in-house tool

developed at Genoscope using the fastx library (http://hannonlab.cshl.edu/fastx_toolkit/). An

alignment of at least 15 bp with at least 90% identity and fewer than 4 errors was required to

fuse two reads into one longer read. The set of fused reads, completed with any reads that could

not be fused, was assembled into contigs by the Newbler version 2.9 overlap-layout-consensus

assembler, with a minimal alignment identity of 99% and a minimal alignment size of 99 bp.

Scaffolds were built from the contigs using 4 Illumina mate-pair libraries with respective insert

sizes of 5 kb, 8 kb, 11 kb and 13 kb. The scaffolder SSpace [70] was used, with default

parameters and an acceptable variation in mate-pair insert size of 25%. Gap closing was a two

step process with SOAPdenovo2 GapCloser software [71]. The first step used the Illumina

paired-end reads, the second step used the Illumina mate-pair libraries. Finally, Kraken software

[72] and the NR nucleotide database were used to detect and remove non-eukaryotic scaffolds,

owing mainly to bacterial contaminants (see below).

Filtering

Scaffolds with a length inferior to 2 kb or with a G+C content greater than 40% were filtered.

Contaminant scaffolds were identified and removed from the assembly provided the Kraken

Kmer score was superior to 10 or a BLASTN match (-evalue 1e-40 –perc_identity 70) against

RefSeq database (excluding *Paramecium* sequences) covered at least 20% of the scaffold

length. If the mitochondrial genome (more or less fragmented) could be identified by a

BLASTN (-evalue 1e-1 –perc_identity 70) against the *P. tetraurelia* mitochondrial genome, the

scaffold(s) were tagged as mitochondrial. A handful of chimeric scaffolds were detected and

corrected in the *P. octaurelia*, *P. primaurelia* and *P. sexaurelia* assemblies by visual inspection

of available long-range sequencing data (remapped mate-pairs) (see S2 Data)).

The constitutive MAC

Paired-end MAC DNA sequencing data were mapped on the MAC genome assembly using

Bowtie2 (v.2.2.3 –local, otherwise default parameters) [73]. We defined the constitutive MAC

as consisting of all regions of the assemblies with the expected average read coverage.We

defined the regions of low coverage at scaffold extremities as MAC-variable regions. These

regions result from the variability of programmed genome rearrangement patterns during MAC

development [74]. While most MIC loci are either fully eliminated during MAC development

(MIC-limited sequences) or fully retained (MAC-destined sequences), MAC-variable regions

correspond to DNA sequences that are not completely eliminated and instead, are retained in a

small fraction of MAC copies. MAC-variable regions represent ~15% of the initial MAC

genome assembly (S2 Table). In relation to the MAC DNA-seq depth, a minimum expected

coverage (v1.9 of samtools depth –q 10 –Q 10) was defined for each assembly (*P. octaurelia*

50X, *P. pentaurelia* 35X, *P. primaurelia* 20X, and *P. sonneborni* 35X). For each scaffold

extremity, a Perl script analyzed the coverage in sliding 2 kb windows. The first window from

the end of the scaffold with a coverage above the minimum expected coverage delimitated the

end of the MAC-variable regions. Only regions of minimum size 4 kb were kept. The script

adjusted region ends using the MAC telomerisation sites and the ends of coding genes. After

this automatic pipeline, each scaffold and mask was adjusted by eye using Circos drawings [75]

(see example S6 Fig, representing DNA and RNA coverage, density in non-coding genes and

positions of the MAC telomerisation sites). The positions of the regions used to reconstruct the

constitutive MAC for each MAC assembly are provided in S2 Data.

IES annotation

Annotation of IESs was performed using the ParTIES toolkit [21] with default parameters.

Briefly, this involves (i) alignment of MIC paired-end reads with a reference MAC genome to

establish a catalog of potential IES insertion sites and to exclude reads that match perfectly

across these sites hence do not contain IESs; (ii) assembly of the remaining reads with Velvet

to obtain contigs that may contain IESs; (iii) alignment of the contigs with the MAC reference

genome to determine the position and the sequence of the IESs.

Gene annotation

Gene annotation for the 9 species was carried out using a pipeline specifically tuned for the

high gene density and tiny intron size (20 − 30 nt) characteristic of *Paramecium* somatic

genomes. RNA-Seq transcriptome data was used to predict transcription units with the TrUC

v1.0 software (https://github.com/oarnaiz/TrUC), as detailed in [37]. EuGene v4.1 software

[76] configured with curated *Paramecium tetraurelia* genes [37] was used for *ab initio*

predictions and to combine annotation evidence (the transcription units, the *ab initio* predictions

and comparative genomics evidence). Gene annotation completeness was calculated using

BUSCO (v4.1.4 --mode prot --lineage_dataset alveolate; [77]) through the Galaxy Portal

(https://usegalaxy.org/).

Assembly-free Genome Size Estimation

Illumina paired-end sequencing reads were used to estimate genome size based on counting all

substrings of 17 nt in the reads, using jellyfish software version 2.2.10 [78]:

`jellyfish count -t 12 -C -m 17 -s 5G -o <sample.jf> <sample_paired_end_reads.fastq>`

`jellyfish histo -o <sample.histo> <sample.jf>`

The method for genome size estimation, described in [18,19], assumes that the total number

of *k*-mers (in this case 17-mers) divided by the sequencing depth is a good approximation for

genome size.

As discussed in [19] for *Paramecium* genomes, the histogram of *k*-mer depth for a perfect,

homozygous genome with no repeated sequences (and no sequencing errors) is fit by a Poisson

distribution, the peak corresponding to sequencing depth. The estimate of genome size is

obtained by dividing the total *k*-mer count (excluding the peak near the origin that results from

k-mers with sequencing errors) by the sequencing depth. This is straightforward for MAC

genomes. As shown in S2 Table, the estimated MAC genome sizes were in good agreement

with the size of the constitutive MAC genome assemblies.

For MIC genomes, variable amounts of contamination from MAC DNA lead to a second peak

at higher *k*-mer depth corresponding to the sum of MAC-destined k-mers in the MIC DNA and

the MAC *k*-mers in the contaminating MAC DNA. This was only a significant problem for the

P. tetraurelia, *P. sexaurelia* and *P. sonneborni* MIC DNA samples, which were approximately

corrected by assuming that a proportion of the *k*-mers counted from this second peak up to a

depth of 500 were contributed by the contaminating MAC reads, while all the *k*-mers with a

depth greater than 500, corresponding to highly repeated sequences, are of MIC origin (S2 Fig).

The proportion of contaminating MAC DNA needed for this calculation was confirmed using

IES retention scores (IRS) calculated with the MIC sequencing reads [21]. The position of the

peak in the IRS distribution indicates the proportion of MIC (IRS ~ 1) and MAC (IRS ~ 0)

DNA in the sample, as illustrated in S2 Fig.

1015

Identification of gene families

We performed an all against all BLASTP (ncbi-blast+ v. 2.2.30+) [79] search using the

predicted protein sequences from each genome including also the proteins of *Tetrahymena*

thermophila (June 2014 assembly http://ciliate.org) as an outgroup. From the resulting output

we determined gene families with SiLiX v. 1.2.9 [80]. The resulting gene families were aligned

with MAFFT v7.305b (2016/Aug/16) [81] using the --auto option. Gene families with fewer

than 3 genes or average pairwise identity less than 50% were excluded from downstream

analyses. From the protein alignments we reconstructed the nucleotide coding sequence

alignments.

1025

Paramecium species phylogeny

To reconstruct the species phylogeny, we first selected single-copy gene families present in all

nine *Paramecium* species (N=1,061 genes). When available, the *T. thermophila* homolog was

also included as an outgroup. We estimated the maximum likelihood phylogeny using IQtree

v.1.4.2 [82], considering each gene as a separate partition. We performed model testing on each

partition and chose the best codon model (determined by the largest BIC). We evaluated the

results by 1,000 bootstrap replicates. All internal branches but one are supported by 100%

bootstrap values (Fig 1). We will hereafter refer to this species tree inferred from single-copy

gene families as *Tree1*.

1035

The rationale for analyzing single-copy gene families is that these sets of homologous

sequences are *a priori* expected to correspond to orthologs. However, given that *Paramecium*

have been subject to three rounds of whole genome duplications followed by massive gene

losses [83], it is possible that some single-copy gene families include paralogs. To check

whether hidden paralogs might have biased the estimation of the species tree, we used

PHYLDOG v.2.0beta (build 10/10/2016), a maximum likelihood method to jointly infer rooted

species and gene trees, accounting for gene duplications and losses [25]. The analysis was

performed using all gene families (N=13,617). The default program options were used with

additionally setting a random starting species tree and BIONJ starting gene trees. The

duplication and loss parameters were optimized with the average then branchwise option and

the genomes were not assumed to have the same number of genes. We also ran PHYLDOG

considering *Tree1* as the fixed species tree, and keeping the remaining options identical. The

topology of the most likely species tree inferred with PHYLDOG is almost identical to *Tree1*

(it only slightly differs in the positions of *P. biaurelia* and *P. tredecaurelia*), and its likelihood

is not significantly different from that obtained when running PHYLDOG with *Tree1* as a

species tree. Thus, the species tree inferred by PHYLDOG using all gene families (N=13,617)

shows no significant disagreement with the phylogeny based on single-copy gene families

(*Tree1*). We therefore hereafter considered *Tree1* as the reference species tree for all our

analyses. To identify duplication and speciation nodes in gene phylogenies, we computed

reconciled trees for each gene family with PHYLDOG, using *Tree1* as a species tree.

1056

1057

1058

To identify events of IES gain and loss along the species phylogeny, it is necessary to analyze

the pattern of presence/absence of IESs at homologous loci across species. One difficulty is that

some IESs may remain undetected (false negatives). In particular, the sensitivity of ParTIEs

depends on the local read coverage [21]. To take into account the uncertainty arising from the

variable local read coverage along scaffolds of each species we calculated the coverage of MIC

reads mapped against the MAC genome. We identified genes with extreme values of coverage

(less than the 10th percentile or more than the 90th over all genes) or with an absolute read

coverage of less than 15 reads. These genes correspond to regions with possible assembly errors

or to regions of low power to detect IESs, and we marked them as problematic for IES

annotation. IESs in these genes were considered to have an uncertain status of presence, and if

no IES was annotated the genes were marked as potentially containing IESs. To avoid issues

due to genome assembly errors, we excluded from our analyses all IESs identified on small

scaffolds (< 10 kb).

1072

Taking into account the uncertainty of IES location (floating IESs).

To identify co-orthologous IES loci, i.e. that result from a single ancestral insertion event, we

searched for IESs located at a same site across homologous sequences. It should be noted that

the location of IESs, inferred from the comparison of MIC and MAC sequences, is sometimes

ambiguous. This occurs when the IES boundaries overlap a motif repeated in tandem (S13 Fig).

Such cases, hereafter called "floating IESs", represent 7% of all IESs. In the vast majority of

cases (86%) the alternative locations of floating IESs differ by only two bp (as in the example

shown in S13 Fig), and there are less than 1% of floating IESs for which the uncertainty in IES

position exceeds 5 bp. To determine the exact location of IESs and capture the inherent

ambiguity due to possible floating IESs we used a 10 bp window around each annotated IES

location to determine if the IES was classified as floating. If so, the alternative locations were

added to the IES annotation.

1085

Co-orthologous IES insertion sites

To detect co-orthologous IES loci, we compared the position of IESs within homologous genes.

To do so, we analyzed gene families with more than 3 sequences and average pairwise identity

(at the protein sequence level) of more than 50%. To avoid ambiguity in the identification of

homologous sites, we filtered protein sequence alignments with GBlocks v0.91b [84] and we

only retained IESs located within conserved alignment blocks. An IES insertion site spans two

nucleotides (TA). In a multiple sequence alignment including gaps, an IES locus can be larger

(e.g. T--A). Two IES loci were considered as homologous if they have at least one shared site

within the alignment (taking into account all potential locations in the case of floating IESs). In

the case of floating IESs overlapping the boundaries of conserved alignment blocks, the

presence or absence of homologous IES loci in other sequences cannot be reliably inferred. We

therefore only retained IESs for which all homologs (if any) are entirely located within the

conserved alignment blocks (i.e. we discarded sets of co-orthologous IES loci that included

some floating IESs for which some of the possible alternative positions were located outside of

the conserved alignment blocks).

1101

Ancestral state reconstruction and inference of IES insertion and loss rates

To explore the dynamics of IES gain and loss we used a Bayesian approach to reconstruct the

ancestral states of presence and absence of IESs using revBayes 1.0.0 beta 3 (2015-10-02) [85].

We constructed binary character matrices (presence/absence) for each gene family containing

at least one IES unambiguously located within a conserved alignment block (see above). We

assumed a model of character evolution with one rate of gain and one rate of loss sampled from

the same exponential distribution with parameter α and a hyperprior sampled from an

exponential with parameter 1. We excluded from the analysis 5 gene families for which

revBayes could not compute a starting probability due to very small numbers. We used

PHYLDOG reconciled gene trees (see above) to fix gene tree topologies and branch lengths.

We ran 5×10^5 iterations. The search parameters were optimized in an initial phase of 10,000

iterations with tuning interval 1,000. Good sampling of the parameter space was verified by

inspecting the time series and autocorrelation plots of the parameters. The convergence was

validated by inspecting the multivariate Gelman and Rubin's diagnostic plots for different

iterations.

1117

Thus, for a given IES locus in a given gene family, revBayes provides an estimate of the

probability of presence of an IES at each node of the gene phylogeny. We used these

probabilities of presence along the gene phylogeny to estimate rates of IES gains or losses in

each branch of the species tree. Because of gene duplications, a given branch in the species tree

can be represented by several paths in the gene tree. Thus, we considered all paths in the gene

tree that connect the corresponding speciation nodes (see S8 Fig for a simplified example). To

measure the IES gain rate at a given IES locus (*c*), in a given gene family (*g*), we define p^+_{cgij}

as the sum of increase in probability of presence of an IES at this locus along all paths of gene

family *g* connecting speciation nodes *i* and *j* (where *i* is a direct ancestor of *j*). Let *ng* be the

length in kilobase pairs of gene family *g* alignment (counting only well aligned sites, where the

presence of IESs can be assessed). Let *Ig* be the number of IES loci in family *g*. Let *kgij* be the

number of paths connecting speciation nodes *i* and *j* in family *g*. Let *bij* be the branch length

connecting nodes *i* and *j* in the species tree (*bij* is taken here as a proxy for time). Let p^+_{gij} be

the sum of increase in probability of presence of an IES, cumulated over all IES loci in family g. We define p^+_{ij} as the sum of increase in probability of presence of an IES, cumulated over all IES loci along the path i to j of family g. We define G_{ij} as the rate of IES gain over all gene families (f) along path ij expressed in number of IES gains per kilobase pairs of alignment per unit of time:

$$G_{ij} = \frac{\sum_{g=1}^{g=f} p^+_{gij}}{\sum_{g=1}^{g=f} n_g k_{gij} \cdot b_{ij}}$$

We define in a similar manner the rate of IES loss. For a given gene family g, let p^-_{cgij} be the sum of decreases in probability of presence of an IES in IES locus c along a lineage in gene family g connecting speciation nodes i and j. Let I_g be the number of IES loci in family g. Let p^-_{gij} be the sum of decrease in probability of presence of an IES, cumulated over all IES loci in family g. We define as L_{ij} the rate of IES loss over all gene families (f) along path ij expressed in number of IES losses per IES, per unit of time.

$$L_{ij} = \frac{\sum_{g=1}^{g=f} p^-_{gij}}{\sum_{g=1}^{g=f} I_g \cdot kg_{ij} \cdot b_{ij}}$$

IES age of insertion

The age of first insertion for each group of co-orthologous IES locations is defined as the age of the most recent common ancestor of all nodes in which an ancestral IES was present with probability larger than 99%.

Identification of homologous IES sequences and characterization of mobile IESs

To characterize families of homologous IES sequences, we first compared all IESs (from all species) against each other with *blastn* (ncbi blast+ v2.5.0, [79]):

1154

```
blastn -evalue 1e-8 -query IES.fa -db IES -dust yes -task blastn -
max_target_seqs 10000
```

1157

We retained all pairs of homologous IESs for which BLAST alignments encompass the first and last 20 nt of the query and subject sequences. This ensures that the detected sequence homology includes the boundaries of the IESs, and is not merely due to the presence of repeated sequences inserted within a pre-existing IES.

To identify potentially mobile IESs, we searched for homologous IES sequences present at different (non-homologous) genomic loci. For this, we extracted 100 nt on each side of the IES location, and compared all these flanking regions against each other with *blastn* (using the same parameters as above). Pairs of homologous IES sequences with strong hits in flanking regions ($\geq$75% identity over 150 nt or more) were classified as 'homologous IESs at homologous loci'. The other pairs were classified as 'candidate mobile IESs'. We clustered each group based on sequence similarity using SiLiX [80] with default parameters.

We further analyzed all clusters of candidate mobile IESs having at least 10 sequences (N=57 clusters). For each cluster, we constructed multiple sequence alignments with MAFFT v7.305b (with `--adjustdirection` and `--auto` options). We manually inspected these alignments to select full-length copies and create a multiple alignment covering the entire repeated element. At this stage, we excluded 11 clusters corresponding to very AT-rich sequences, for which it was not clear whether the detected sequence similarities were due to homology or to their highly biased sequence composition. Furthermore, two clusters were split into subfamilies, to include only sequences that are homologous over their entire length. We

1179 then used these seed alignments to build an HMM profile for each repeat family and search for

1180 homologous copies among the entire IES dataset with NHMMER version 3.1b2 [86].

1181

In total, NHMMER identified 12,184 IESs having a significant hit (E-value $< 10^{-3}$) in the dataset

of HMM profiles. Among detected hits, we distinguished two categories: 1) cases where the

detected repeated element is located within the IES but does not overlap with the extremities of

the IES (i.e. nested repeats), and 2) cases where the extremities of the HMM profile align with

the extremities of the IES (with a tolerance of 3 bp to allow alignment uncertainties). IESs

belonging to this latter category were hereafter considered as 'mobile IESs'. For subsequent

analyses, we selected all families with more than 10 mobile IESs in at least one genome (N=24

families of mobile IESs). Multiple alignments, HMM profiles and the list of matching IESs are

available (https://doi.org/10.5281/zenodo.4726682).

1191

Acknowledgments

We thank Damien de Vienne for his help in phylogenetic analyses, Alexey Potekhin (Saint

Petersburg State University, St Petersburg, Russia) and Ewa Przybos (Institute of Systematics

and Evolution of Animals, Polish Academy of Sciences, Cracow, Poland) for the maintenance

of *Paramecium* stock collections and providing some *Paramecium aurelia* strains. This work

was performed using the computing facilities of the CC LBBE/PRABI.

1198

1199

References

1. Cheng C-Y, Orias E, Leu J-Y, Turkewitz AP. The evolution of germ-soma nuclear differentiation in eukaryotic unicells. Curr Biol. 2020;30: R502–R510. doi:10.1016/j.cub.2020.02.026

2. Betermier M, Duharcourt S. Programmed Rearrangement in Ciliates: Paramecium. Microbiol Spectr. 2014;2. doi:10.1128/microbiolspec.MDNA3-0035-2014

3. Rzeszutek I, Maurer-Alcalá XX, Nowacki M. Programmed genome rearrangements in ciliates. Cell Mol Life Sci. 2020. doi:10.1007/s00018-020-03555-2

4. Arnaiz O, Mathy N, Baudry C, Malinsky S, Aury J-M, Denby Wilkes C, et al. The Paramecium Germline Genome Provides a Niche for Intragenic Parasitic DNA: Evolutionary Dynamics of Internal Eliminated Sequences. PLoS Genetics. 2012;8: e1002984. doi:10.1371/journal.pgen.1002984

5. Guérin F, Arnaiz O, Boggetto N, Denby Wilkes C, Meyer E, Sperling L, et al. Flow cytometry sorting of nuclei enables the first global characterization of Paramecium germline DNA and transposable elements. BMC Genomics. 2017;18. doi:10.1186/s12864-017-3713-7

6. Baudry C, Malinsky S, Restituito M, Kapusta A, Rosa S, Meyer E, et al. PiggyMac, a domesticated piggyBac transposase involved in programmed genome rearrangements in the ciliate Paramecium tetraurelia. Genes Dev. 2009;23: 2478–2483. doi:10.1101/gad.547309

7. Bischerour J, Bhullar S, Denby Wilkes C, Régnier V, Mathy N, Dubois E, et al. Six domesticated PiggyBac transposases together carry out programmed DNA elimination in Paramecium. Elife. 2018;7. doi:10.7554/eLife.37927

8. Vanssay A de, Touzeau A, Arnaiz O, Frapporti A, Phipps J, Duharcourt S. The Paramecium histone chaperone Spt16-1 is required for Pgm endonuclease function in programmed genome rearrangements. PLOS Genetics. 2020;16: e1008949.

doi:10.1371/journal.pgen.1008949

9.   Abello A, Régnier V, Arnaiz O, Le Bars R, Bétermier M, Bischerour J. Functional

diversification of Paramecium Ku80 paralogs safeguards genome integrity during precise

programmed DNA elimination. PLoS Genet. 2020;16: e1008723.

doi:10.1371/journal.pgen.1008723

10.   Klobutcher LA, Herrick G. Consensus inverted terminal repeat sequence of

Paramecium IESs: resemblance to termini of Tc1-related and Euplotes Tec transposons.

Nucleic Acids Res. 1995;23: 2006–2013. doi:10.1093/nar/23.11.2006

11.   Klobutcher LA, Herrick G. Developmental genome reorganization in ciliated

protozoa: the transposon link. Prog Nucleic Acid Res Mol Biol. 1997;56: 1–62.

doi:10.1016/s0079-6603(08)61001-6

12.   Singh DP, Saudemont B, Guglielmi G, Arnaiz O, Goût J-F, Prajer M, et al. Genome-

defence small RNAs exapted for epigenetic mating-type inheritance. Nature. 2014;509: 447–

452. doi:10.1038/nature13318

13.   Sawka-Gądek N, Potekhin A, Singh DP, Grevtseva I, Arnaiz O, Penel S, et al.

Evolutionary Plasticity of Mating-Type Determination Mechanisms in Paramecium aurelia

Sibling Species. Genome Biol Evol. 2020;13. doi:10.1093/gbe/evaa258

14.   McGrath CL, Gout J-F, Johri P, Doak TG, Lynch M. Differential retention and

divergent resolution of duplicate genes following whole-genome duplication. Genome

Research. 2014;24: 1665–1675. doi:10.1101/gr.173740.114

15.   McGrath CL, Gout J-F, Doak TG, Yanagi A, Lynch M. Insights into Three Whole-

Genome Duplications Gleaned from the *Paramecium caudatum* Genome Sequence. Genetics.

2014;197: 1417–1428. doi:10.1534/genetics.114.163287

16.   Gout J-F, Johri P, Arnaiz O, Doak TG, Bhullar S, Couloux A, et al. Universal trends

of post-duplication evolution revealed by the genomes of 13 Paramecium species sharing an

ancestral whole-genome duplication. bioRxiv. 2019; 573576. doi:10.1101/573576

17.   Sonneborn TM. Paramecium aurelia. In: King RC, editor. Handbook of Genetics:

Plants, Plant Viruses, and Protists. Boston, MA: Springer US; 1974. pp. 469–594.

doi:10.1007/978-1-4684-2994-7_20

18.   Li R, Fan W, Tian G, Zhu H, He L, Cai J, et al. The sequence and de novo assembly of

the giant panda genome. Nature. 2010;463: 311–317. doi:10.1038/nature08696

19.   Duharcourt S, Sperling L. The Challenges of Genome-Wide Studies in a Unicellular

Eukaryote With Two Nuclear Genomes. Meth Enzymol. 2018;612: 101–126.

doi:10.1016/bs.mie.2018.08.012

20.   Johri P, Krenek S, Marinov GK, Doak TG, Berendonk TU, Lynch M. Population

Genomics of Paramecium Species. Molecular Biology and Evolution. 2017;34: 1194–1216.

doi:10.1093/molbev/msx074

21.   Denby Wilkes C, Arnaiz O, Sperling L. ParTIES: a toolbox for Paramecium

interspersed DNA elimination studies. Bioinformatics. 2016;32: 599–601.

doi:10.1093/bioinformatics/btv691

22.   Mayer KM, Mikami K, Forney JD. A mutation in Paramecium tetraurelia reveals

functional and structural features of developmentally excised DNA elements. Genetics.

1998;148: 139–149.

23.   Duharcourt S, Keller AM, Meyer E. Homology-dependent maternal inhibition of

developmental excision of internal eliminated sequences in Paramecium tetraurelia. Mol Cell

Biol. 1998;18: 7075–7085. doi:10.1128/mcb.18.12.7075

24.   Przyboś E, Tarcz S, Rautian M, Sawka N. Delimiting Species Boundaries within a

Paraphyletic Species Complex: Insights from Morphological, Genetic, and Molecular Data on

Paramecium sonneborni (Paramecium aurelia species complex, Ciliophora, Protozoa). Protist.

2015;166: 438–456. doi:10.1016/j.protis.2015.07.001

25.   Boussau B, Szollosi GJ, Duret L, Gouy M, Tannier E, Daubin V. Genome-scale

coestimation of species and gene trees. Genome Research. 2013;23: 323–330.

doi:10.1101/gr.141978.112

26.   Przyboś E, Tarcz S. New Stands of the Paramecium aurelia spp. Complex (Protista,

Oligohymenophorea) in Ethiopia, Madagascar, Taiwan, and Romania. Folia Biologica.

2018;66: 111–119. doi:10.3409/fb_66-3.12

27.   Przyboś E, Tarcz S, Rautian M, Sawka N. Delimiting Species Boundaries within a

Paraphyletic Species Complex: Insights from Morphological, Genetic, and Molecular Data on

Paramecium sonneborni (Paramecium aurelia species complex, Ciliophora, Protozoa). Protist.

2015;166: 438–456. doi:10.1016/j.protis.2015.07.001

28.   Guindon S, Dufayard J-F, Lefort V, Anisimova M, Hordijk W, Gascuel O. New

algorithms and methods to estimate maximum-likelihood phylogenies: assessing the

performance of PhyML 3.0. Syst Biol. 2010;59: 307–321. doi:10.1093/sysbio/syq010

29.   Crooks GE, Hon G, Chandonia J-M, Brenner SE. WebLogo: a sequence logo

generator. Genome Res. 2004;14: 1188–1190. doi:10.1101/gr.849004

30.   Duret L, Cohen J, Jubin C, Dessen P, Gout J-F, Mousset S, et al. Analysis of sequence

variability in the macronuclear DNA of Paramecium tetraurelia: A somatic view of the

germline. Genome Research. 2008;18: 585–596. doi:10.1101/gr.074534.107

31.   Lhuillier-Akakpo M, Frapporti A, Denby Wilkes C, Matelot M, Vervoort M, Sperling

L, et al. Local Effect of Enhancer of Zeste-Like Reveals Cooperation of Epigenetic and cis-

Acting Determinants for Zygotic Genome Rearrangements. PLoS Genetics. 2014;10:

e1004665. doi:10.1371/journal.pgen.1004665

32.   Maliszewska-Olejniczak K, Gruchota J, Gromadka R, Denby Wilkes C, Arnaiz O,

Mathy N, et al. TFIIS-Dependent Non-coding Transcription Regulates Developmental

Genome Rearrangements. PLoS Genet. 2015;11: e1005383.

doi:10.1371/journal.pgen.1005383

33.   Gruchota J, Denby Wilkes C, Arnaiz O, Sperling L, Nowak JK. A meiosis-specific

Spt5 homolog involved in non-coding transcription. Nucleic Acids Res. 2017;45: 4722–4732.

doi:10.1093/nar/gkw1318

34.   Sandoval PY, Swart EC, Arambasic M, Nowacki M. Functional diversification of

Dicer-like proteins and small RNAs required for genome sculpting. Dev Cell. 2014;28: 174–

188. doi:10.1016/j.devcel.2013.12.010

35.   Frapporti A, Miró Pina C, Arnaiz O, Holoch D, Kawaguchi T, Humbert A, et al. The

Polycomb protein Ezl1 mediates H3K9 and H3K27 methylation to repress transposable

elements in Paramecium. Nature Communications. 2019;10: 2710. doi:10.1038/s41467-019-

10648-5

36.   Lepère G, Bétermier M, Meyer E, Duharcourt S. Maternal noncoding transcripts

antagonize the targeting of DNA elimination by scanRNAs in Paramecium tetraurelia. Genes

Dev. 2008;22: 1501–1512. doi:10.1101/gad.473008

37.   Arnaiz O, Van Dijk E, Bétermier M, Lhuillier-Akakpo M, de Vanssay A, Duharcourt

S, et al. Improved methods and resources for paramecium genomics: transcription units, gene

annotation and gene expression. BMC Genomics. 2017;18: 483. doi:10.1186/s12864-017-

3887-z

38.   Blumenstiel JP. Birth, School, Work, Death, and Resurrection: The Life Stages and

Dynamics of Transposable Element Proliferation. Genes. 2019;10: 336.

doi:10.3390/genes10050336

39.   Petrov DA, Sangster TA, Johnston JS, Hartl DL, Shaw KL. Evidence for DNA Loss as

a Determinant of Genome Size. Science. 2000;287: 1060–1062.

doi:10.1126/science.287.5455.1060

40.   Mitra R, Fain-Thornton J, Craig NL. piggyBac can bypass DNA synthesis during cut

and paste transposition. EMBO J. 2008;27: 1097–1109. doi:10.1038/emboj.2008.41

41.   Hamilton EP, Kapusta A, Huvos PE, Bidwell SL, Zafar N, Tang H, et al. Structure of

the germline genome of Tetrahymena thermophila and relationship to the massively

rearranged somatic genome. eLife. 2016;5. doi:10.7554/eLife.19090

42.   Cheng C-Y, Young JM, Lin C-YG, Chao J-L, Malik HS, Yao M-C. The piggyBac

transposon-derived genes TPB1 and TPB6 mediate essential transposon-like excision during

the developmental rearrangement of key genes in Tetrahymena thermophila. Genes Dev.

2016;30: 2724–2736. doi:10.1101/gad.290460.116

43.   Feng L, Wang G, Hamilton EP, Xiong J, Yan G, Chen K, et al. A germline-limited

piggyBac transposase gene is required for precise excision in Tetrahymena genome

rearrangement. Nucleic Acids Research. 2017;45: 9481–9502. doi:10.1093/nar/gkx652

44.   Cheng C-Y, Vogt A, Mochizuki K, Yao M-C. A Domesticated piggyBac Transposase

Plays Key Roles in Heterochromatin Dynamics and DNA Cleavage during Programmed DNA

Deletion in Tetrahymena thermophila. MBoC. 2010;21: 1753–1762. doi:10.1091/mbc.e09-

12-1079

45.   Vogt A, Mochizuki K. A domesticated PiggyBac transposase interacts with

heterochromatin and catalyzes reproducible DNA elimination in Tetrahymena. PLoS Genet.

2013;9: e1004032. doi:10.1371/journal.pgen.1004032

46.   Lynch M. The Origins of Eukaryotic Gene Structure. Molecular Biology and

Evolution. 2006;23: 450–468. doi:10.1093/molbev/msj050

47.   Lynch M. Intron evolution as a population-genetic process. Proceedings of the

National Academy of Sciences. 2002;99: 6118–6123. doi:10.1073/pnas.092595699

48.   Cavalier-Smith T. Intron phylogeny: a new hypothesis. Trends in Genetics. 1991;7:

145–148. doi:10.1016/0168-9525(91)90377-3

49.   Wahl MC, Will CL, Lührmann R. The Spliceosome: Design Principles of a Dynamic

RNP Machine. Cell. 2009;136: 701–718. doi:10.1016/j.cell.2009.02.009

50.   Doolittle WF. Is junk DNA bunk? A critique of ENCODE. Proceedings of the

National Academy of Sciences. 2013;110: 5294–5300. doi:10.1073/pnas.1221376110

51.   Huff JT, Zilberman D, Roy SW. Mechanism for DNA transposons to generate introns

on genomic scales. Nature. 2016;538: 533–536. doi:10.1038/nature20110

52.   Pickrell JK, Pai AA, Gilad Y, Pritchard JK. Noisy Splicing Drives mRNA Isoform

Diversity in Human Cells. PLoS Genetics. 2010;6: e1001236.

doi:10.1371/journal.pgen.1001236

53.   Saudemont B, Popa A, Parmley JL, Rocher V, Blugeon C, Necsulea A, et al. The

fitness cost of mis-splicing is the main determinant of alternative splicing patterns. Genome

Biology. 2017;18: 208. doi:10.1186/s13059-017-1344-6

54.   Tress ML, Abascal F, Valencia A. Most Alternative Isoforms Are Not Functionally

Important. Trends in Biochemical Sciences. 2017;42: 408–410.

doi:10.1016/j.tibs.2017.04.002

55.   Nilsen TW, Graveley BR. Expansion of the eukaryotic proteome by alternative

splicing. Nature. 2010;463: 457–463. doi:10.1038/nature08909

56.   McGlincy NJ, Smith CWJ. Alternative splicing resulting in nonsense-mediated mRNA

decay: what is the meaning of nonsense? Trends in Biochemical Sciences. 2008;33: 385–393.

doi:10.1016/j.tibs.2008.06.001

57.   Lareau LF, Inada M, Green RE, Wengrod JC, Brenner SE. Unproductive splicing of

SR genes associated with highly conserved and ultraconserved DNA elements. Nature.

2007;446: 926–929. doi:10.1038/nature05676

58.   Lareau LF, Brenner SE. Regulation of Splicing Factors by Alternative Splicing and

NMD Is Conserved between Kingdoms Yet Evolutionarily Flexible. Molecular Biology and

Evolution. 2015;32: 1072–1079. doi:10.1093/molbev/msv002

59.   Doolittle WF, Sapienza C. Selfish genes, the phenotype paradigm and genome

evolution. Science. 1980;284: 601–603.

60.   Orgel LE, Crick FHC. Selfish DNA: the ultimate parasite. Science. 1980;284: 604–

607.

61.   Werren JH. Selfish genetic elements, genetic conflict, and evolutionary innovation.

Proceedings of the National Academy of Sciences. 2011;108: 10863–10870.

doi:10.1073/pnas.1102343108

62.   Krenek S, Berendonk TU, Petzoldt T. Thermal performance curves of Paramecium

caudatum: a model selection approach. Eur J Protistol. 2011;47: 124–137.

doi:10.1016/j.ejop.2010.12.001

63.   Lhuillier-Akakpo M, Guérin F, Frapporti A, Duharcourt S. DNA deletion as a

mechanism for developmentally programmed centromere loss. Nucleic Acids Res. 2016;44:

1553–1565. doi:10.1093/nar/gkv1110

64.   Cummings DJ. Isolation and partial characterization of macro- and micronuclei from

Paramecium aurella. J Cell Biol. 1972;53: 105–115. doi:10.1083/jcb.53.1.105

65.   Freiburg M. Isolation and characterization of macronuclei of Paramecium caudatum

infected with the macronucleus-specific bacterium Holospora obtusa. J Cell Sci. 1985;73:

389–398.

66.   Galbraith DW, Harkins KR, Maddox JM, Ayres NM, Sharma DP, Firoozabady E.

Rapid Flow Cytometric Analysis of the Cell Cycle in Intact Plant Tissues. Science. 1983;220:

1049–1051. doi:10.1126/science.220.4601.1049

67.   Bourge M, Brown SC, Siljak-Yakovlev S. Flow cytometry as tool in plant sciences,

with emphasis on genome size and ploidy level assessment. Genetics & Applications. 2018;2:

1–12. doi:10.31383/ga.vol2iss2pp1-12

68.   Dolezel J, Bartos J, Voglmayr H, Greilhuber J. Nuclear DNA content and genome size

of trout and human. Cytometry A. 2003;51: 127–128. doi:10.1002/cyto.a.10013

69.   Marie D, Brown SC. A cytometric exercise in plant DNA histograms, with 2C values for 70 species. Biol Cell. 1993;78: 41–51. doi:10.1016/0248-4900(93)90113-s

70.   Boetzer M, Henkel CV, Jansen HJ, Butler D, Pirovano W. Scaffolding pre-assembled contigs using SSPACE. Bioinformatics. 2011;27: 578–579. doi:10.1093/bioinformatics/btq683

71.   Luo R, Liu B, Xie Y, Li Z, Huang W, Yuan J, et al. SOAPdenovo2: an empirically improved memory-efficient short-read de novo assembler. Gigascience. 2012;1: 18. doi:10.1186/2047-217X-1-18

72.   Wood DE, Salzberg SL. Kraken: ultrafast metagenomic sequence classification using exact alignments. Genome Biol. 2014;15: R46. doi:10.1186/gb-2014-15-3-r46

73.   Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. Nat Methods. 2012;9: 357–359. doi:10.1038/nmeth.1923

74.   Le Mouël A, Butler A, Caron F, Meyer E. Developmentally regulated chromosome fragmentation linked to imprecise elimination of repeated sequences in paramecia. Eukaryotic Cell. 2003;2: 1076–1090. doi:10.1128/ec.2.5.1076-1090.2003

75.   Krzywinski M, Schein J, Birol I, Connors J, Gascoyne R, Horsman D, et al. Circos: an information aesthetic for comparative genomics. Genome Res. 2009;19: 1639–1645. doi:10.1101/gr.092759.109

76.   Foissac S, Gouzy J, Rombauts S, Mathe C, Amselem J, Sterck L, et al. Genome Annotation in Plants and Fungi: EuGene as a Model Platform. Current Bioinformatics. 2008;3: 87–97.

77.   Simão FA, Waterhouse RM, Ioannidis P, Kriventseva EV, Zdobnov EM. BUSCO: assessing genome assembly and annotation completeness with single-copy orthologs. Bioinformatics. 2015;31: 3210–3212. doi:10.1093/bioinformatics/btv351

78.   Marçais G, Kingsford C. A fast, lock-free approach for efficient parallel counting of

occurrences of k-mers. Bioinformatics. 2011;27: 764–770. doi:10.1093/bioinformatics/btr011

79.   Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, et al.

BLAST+: architecture and applications. BMC Bioinformatics. 2009;10: 421.

doi:10.1186/1471-2105-10-421

80.   Miele V, Penel S, Duret L. Ultra-fast sequence clustering from similarity networks

with SiLiX. BMC Bioinformatics. 2011;12: 116. doi:10.1186/1471-2105-12-116

81.   Katoh K, Standley DM. MAFFT multiple sequence alignment software version 7:

improvements in performance and usability. Mol Biol Evol. 2013;30: 772–780.

doi:10.1093/molbev/mst010

82.   Nguyen L-T, Schmidt HA, von Haeseler A, Minh BQ. IQ-TREE: a fast and effective

stochastic algorithm for estimating maximum-likelihood phylogenies. Mol Biol Evol.

2015;32: 268–274. doi:10.1093/molbev/msu300

83.   Aury J-M, Jaillon O, Duret L, Noel B, Jubin C, Porcel BM, et al. Global trends of

whole-genome duplications revealed by the ciliate Paramecium tetraurelia. Nature. 2006;444:

171–178. doi:10.1038/nature05230

84.   Talavera G, Castresana J. Improvement of phylogenies after removing divergent and

ambiguously aligned blocks from protein sequence alignments. Syst Biol. 2007;56: 564–577.

doi:10.1080/10635150701472164

85.   Höhna S, Landis MJ, Heath TA, Boussau B, Lartillot N, Moore BR, et al. RevBayes:

Bayesian Phylogenetic Inference Using Graphical Models and an Interactive Model-

Specification Language. Syst Biol. 2016;65: 726–736. doi:10.1093/sysbio/syw021

86.   Wheeler TJ, Eddy SR. nhmmer: DNA homology search with profile HMMs.

Bioinformatics. 2013;29: 2487–2489. doi:10.1093/bioinformatics/btt403

87.   Goubert C, Modolo L, Vieira C, ValienteMoro C, Mavingui P, Boulesteix M. De novo

assembly and annotation of the Asian tiger mosquito (Aedes albopictus) repeatome with

dnaPipeTE from raw genomic reads and comparative analysis with the yellow fever mosquito

(Aedes aegypti). Genome Biol Evol. 2015;7: 1192–1205. doi:10.1093/gbe/evv050

1453

1454

Supporting information

S1 Fig. Multi-gate flow cytometry strategy for sorting the MICs.

GFP, DAPI-positive MICs from *P. sonneborni* vegetative cells transformed with the *P. tetraurelia CENH3a-GFP* transgene [63] were sorted based on size, granularity, DAPI staining and GFP signal (see Materials and Methods). P4 and P8 were sorted separately. Based on quality control by flow imaging (Imagestream) indicating 97% purity, the two samples P4 and P8, which represent 1.91% of total events, were combined for DNA extraction and sequencing. Two populations are visualized and likely correspond to 2n and 4n MICs.

S2 Fig. Estimating the proportion of MIC and MAC DNA in the sample based on IES retention score.

The histograms on the left show the k-mer depth profiles. The peak at the origin can be attributed to sequencing errors (k-mers that occur only once or a few times). The position of the largest peak beyond the origin corresponds to k-mers present once in the genome and provides the sequencing depth. As *Paramecium aurelia* genomes have undergone whole genome duplications, there are a significant number of k-mers at 2X and even 4X the sequencing depth arising from genes (or regions of genes) present in 2 or 4 copies, clearly visible for *P. octaurelia* and *P. primaurelia*. The profile for *P. tetraurelia* however has a first peak (MIC sequences that occur once) at 31X followed by a larger peak that is not at the 2X position as it arises because of MAC DNA contamination. The column on the right shows histograms of IES retention scores. Only the *P. tetraurelia* sample is significantly contaminated by MAC DNA: the average IES retention score of 0.4 indicates 40% MIC and 60% MAC DNA in this sample.

S3 Fig. Comparison of cytometry and k-mer MIC genome size estimates.

Flow cytometry estimates of DNA content of micronuclei and k-mer counting estimates of

genome size are described in Materials and Methods. In order to show all of the data, both axes

of the graph are log-transformed. Simple linear regression was carried out on the untransformed

data with R. The linear model that fits the data is presented as a dashed blue line; $R^2 = 0.99$, p-

value = 1.3×10^{-09}.

1486

S4 Fig. Repeat content of *P. caudatum* MIC genome.

(A) Abundance of repeat families identified by DNAPipeTE in *P. caudatum* strain My43c3d.

The repeat content of the *P. caudatum* MIC genome was analyzed with DNAPipeTE [87], using

a sample of 3,500,000 sequence reads (corresponding to a read depth of ~0.5X). DNAPipeTE

identified 67 repeat families that collectively constitute 83% of the MIC genome. Among them,

there are two major satellite repeats Sat1 and Sat2, which represent respectively 42% and 29%

of the MIC genome.

(B) Sequences of the two major satellite repeats Sat1 and Sat2 in *P. caudataum* My43c3d (332

bp and 449 bp long). These two satellite repeats share homology over a ~200 bp-long region.

Primer sequences used for specific PCR amplification of each repeat are indicated in bold.

(C) Detection of Sat1 and Sat2 *in P. caudatum* strains. Whole cell genomic DNA was used to

perform duplex PCR with a set of primers located within each repeat (Sat1 or Sat2, in bold

panel B) and another set of primers within the 18S ribosomal DNA as a loading control. The

expected size of the 18SrDNA PCR product was 301 bp using primers 18S_F953:

AGACGATCAGATACCGTCGTAG and 18S_R1300: CACCAACTAAGAACGGCCATGC.

L: 1-kb NEB ladder. Neg.: negative control (no DNA). Sat1 was amplified with primers

comp2975_F1: TTGTGCTGTAGGGCTCAATAAT and comp2975_R1:

CTCAAAATTCGACGCTGACAA at the expected size (198 bp) in *the P. caudatum* clade B

strains tested (My43c3d; C033; C083; C131; C147). The repeat could not be amplified in *P.*

caudatum DNA from clade A strains (C023; C065; C104; C119), from strain C026 or from

strain Indo_1.6I.

Sat2 was amplified with primers comp5240_F1: TGCTGCTGATTTTGGATCTCG and

comp5240_R1: CCGAGAACGGCCATTACAAG at the expected size (168 bp) in the *P.*

caudatum clade B strains tested (My43c3d; C033; C083; C131; C147). The repeat could not

be amplified in *P. caudatum* DNA from clade A strains (C023; C065; C104; C119), from strain

C026 or from strain Indo_1.6I.

1513

S5 Fig. Impact of sequencing depth on IES detection. To assess the impact of sequencing

depth on the sensitivity of IES detection, we subsampled sequence reads from the *P. tetraurelia*

MIC dataset so as to obtain subsets of lower depth (from 2x to 30x), on which we applied the

same IES detection procedure. (A) Number of detected IESs vs. sequencing depth. (B-H)

Length distributions of IESs detected within each subset (the percentage of IESs in each peak

is indicated). Sequencing depth affects the number of detected IESs (for depths <15x), but not

their length distribution.

1521

S6 Fig. Example of a MAC-variable region

Circular representation of one scaffold of ~730 kb. The tracks from the exterior to the interior

of the circle: G+C content of 100nt sliding windows (black), MAC DNA-seq depth (purple),

the density in predicted non-coding genes (orange), RNA-seq depth (red) and the density of

detected telomerisation sites (green). The external blue arc shows the region identified as being

MAC-variable. These regions were determined by an automatic pipeline (see Materials and

Methods), then adjusted by eye for each scaffold.

1529

S7 Fig. Intragenic IES density vs. gene expression level.

Expression levels (RPKM) were measured with RNAseq datasets from vegetative cells. For each species, expressed genes were classified into 10 bins of equal sample size according to their expression level, and IES density computed within each bin. Non-expressed genes (6.6% of the entire dataset) were excluded. (A) *Paramecium aurelia* species. (B) *P. caudatum*.

S8 Fig. Measuring the rate of IES gain or loss along the species phylogeny.

To illustrate our methodology, we show here an example of a gene family with 3 genes, two from *P. sonneborni* (pson1, pson2) and one from *P. sexaurelia* (psex1). Two IES loci are found in this family (A, B). The probability of presence of an IES (estimated by Bayesian ancestral state reconstruction - see Materials and Methods) is indicated by shaded circles for each locus at each node of the gene phylogeny. We focus here on the branch of the species tree leading from the common ancestor of *P. sexaurelia* and *P. sonneborni* to the leaf node of *P. sonneborni* (the red branch in the species tree, shown in the insert). The length of this branch (b), is taken as a proxy for time. Because of a duplication event, this branch of the species tree corresponds to two paths in the gene tree ($k=2$). To estimate the IES gain rate, we calculate for each path the sum of increase in the probability of presence of an IES, for all IES loci (p^+). Along the first path (from the root to pson1), we have $p^+A1=0.5$ and $p^+B1=0$. Along the second path (from the root to pson2), we have $p^+A2=0.5$ and $p^+B2=0$. The average gain rate along all paths, per unit of time and per bp, is thus given by $G=(p^+A1 + p^+B1 + p^+A2 + p^+B2)/(k \times b \times n_g)$, where n_g is the number of well aligned sites in the gene family alignment (i.e. the number of sites where the presence of co-orthologous IESs can be assessed). Similarly, to estimate the IES loss rate, we calculate for each path the sum of decrease in the probability of presence of an IES, for all IES loci (p^-). Along the first path (from the root to pson1), we have $p^-A1=0$ and $p^-B1=0.4$. Along the second path (from the root to pson2), we have $p^-A2=0$ and $p^-B2=0.4$. The average gain rate

1555   along all paths, per unit of time and per bp, is thus given by $L=(p^-A1 + p^-B1 + p^-A2 + p^-B2)/(k$

1556 $\times b \times I)$, where I is the number of IES loci in the gene family (here $I=2$).

1557

S9 Fig. Dating events of IES insertion/loss.

(A) To date events of IES loss or gain, it is first necessary to identify IESs that derive from a

single ancestral insertion event (co-orthologous IESs). For this, we aligned coding sequences

(based on the protein alignment) and mapped the position of IESs: IESs located at the exact

same position within a codon were assumed to be co-orthologous. We then used the reconciled

gene tree to map events in the species phylogeny, using a maximum likelihood approach (see

Materials and Methods). The example shown here corresponds to a gene family encoding a

putative RNA 3'-terminal phosphate cyclase (PTET.51.1.P0920097, POCTA.138.1.P0960088,

PBIA.V1_4.1.P01950012,   PTRED.209.2.P71800001293600070,   PPENT.87.1.P1090087,

PPRIM.AZ9-3.1.P0020612,   PSON.ATCC_30995.1.P0860097,   PSEX.AZ8_4.1.P0910047,

PCAU.43c3d.1.P00760109). The positions of IESs are indicated by red rectangles. (B) The

presence of IESs (red bars) within each of these genes is indicated with regard to the species

phylogeny. Six distinct IESs were identified in this gene family: IES2 is shared by all species

and therefore predates the divergence between *P. caudatum* and the *aurelia* clade; IES4 most

probably corresponds to a gain in the *P. sexaurelia* lineage; IES5 and IES6 predate the

divergence of the *aurelia* clade and have been subsequently lost in the *P. tetraurelia/P.*

octaurelia lineage; IES1 might correspond to a gain at the base of the *aurelia* clade or a loss in

the *P. caudatum* lineage (and vice versa for IES3).

1576

S10 Fig. Prevalence of weak IESs.

(A) Proportion of weak IESs (i.e. IESs with a retention frequency $\geq 10\%$ in WT vegetative cells)

among IESs located in different genomic compartments. The proportion of weak IESs among

intergenic IESs was compared to that of IESs located in introns or coding regions, by a Chi-

squared test. (B) Proportion of weak IESs according to the age of IESs (for IESs located in

coding regions): New = species-specific IES; Old = IES predating the divergence between *P.*

caudatum and the *aurelia* lineage. The number of new IESs is indicated for each species. The

proportion of weak IESs among new IESs was compared to that of older ones, by a Chi-squared

test. Species codes: pso: *P. sonneborni*, ptr: *P. tredecaurelia*, pte: *P. tetraurelia*, pbi: *P.*

biaurelia, poc: *P. octaurelia*, pse: *P. sexaurelia*, ppr: *P. primaurelia*, ppe: *P. pentaurelia*, pca:

P. caudatum. (*: p-value < 0.05; **: p-value < 1e-3; ***: p-value < 1e-6).

1588

1589 **S11 Fig. Length distribution of IESs according to their age**.

1590 Comparison of the length distribution of IESs according to their age (for the subset of datable

1591 IESs located in coding regions). The age of an IES site is defined as in Fig 3. Results for other

1592 species are shown in Fig 4.

1593

S12 Fig. Genomic distribution of IESs according to their length.

Green bars indicate the percentage of IESs located in each compartment of the MAC genome

(introns, protein-coding regions, intergenic regions) for each species. Grey bars indicate the

percentage of the MAC genome in each compartment. (A) Long IESs (>100 bp). (B) Short IESs

(<35 bp). For each species, the relative proportion of IESs in the three compartments (intron,

intergenic, coding regions) was compared for short IESs (<35 bp) vs long IESs (> 100 bp) by

a Chi-squared test (p-value $<10^{-16}$ in all species).

1601

S13 Fig. Example of floating IES.

Comparison of MIC and MAC sequences indicates the presence of an IES at this locus.

However, because of the presence of a repeated motif at the boundaries of the IES (blue arrows),

it is not possible to determine which of the two possible segments (IES-1 in black or IES-2 in

red) is actually excised *in vivo*. Such IESs that cannot be unambiguously positioned are called

'floating IESs'. They represent 6.8% of the 400,254 IESs detected across all species. In the vast

majority of cases (86%) the alternative locations of floating IESs differ by only two bp (as in

the example shown here), and there are less than 1% of floating IESs for which the uncertainty

in IES position exceeds 5 bp.

1611

S1 Table. MIC genomes: sequencing data and size estimates. (a) Highly enriched sorted

MIC (97-99%) are contaminated with the highly polyploid MAC. For instance, 97% of MIC

purity led to 60% of MAC DNA contamination in the case of P. tetraurelia (see S2 Fig). MIC

genome size (in Mb) was estimated based on (b) flow cytometry analysis and (c) k-mer counts.

Size estimation before correction, based on MAC contamination, is indicated in parentheses

(see Material and Methods).

1618

S2 Table. MAC genome assemblies used in this study.

The assemblies of the four MAC genomes sequenced in the course of this project include both

'constitutive MAC' regions (i.e. regions that are always retained in the MAC) and 'MAC-

variable regions' (i.e. regions that are mostly restricted to the MIC, but that are retained at low

frequency in MAC nuclei). The size and content of these two types of regions are indicated.

The completeness of MAC genome assemblies was assessed using BUSCO [77].

1625

1626   **S3 Table. Distribution of IESs in different genomic compartments.**

1627   Values in parentheses indicate the proportions expected under the hypothesis of uniform IES

1628   distribution along MAC-destined regions.

1629

S4 Table. Genomic distribution of copies of mobile IESs.

Copies of mobile IESs were searched within IES sequences (see Materials and Methods). Detected copies are divided in two categories: nested copies (i.e. copies inserted within an IES, but not including the extremities of the IES) and *bona fide* mobile IESs (i.e. copies whose extremities correspond to the extremities of the IES). This table lists all families for which at least one species contains ≥ 10 copies of *bona fide* mobile IESs in its genome. The genomic distribution (in coding regions, introns and intergenic regions) of IESs containing these copies (nested copies or *bona fide* mobile IESs) is indicated.

S5 Table. Highly conserved IESs that are transcribed during MAC development and/or associated to genes that are upregulated during MAC development. The transcription level of IESs is indicated for different stages during MAC development (S, T0 to T45) and in vegetative cells (V) [37]. Only *P. tetraurelia* IESs are shown in the table, because this is the only species for which developmental transcriptome data is available [37]. (*) The IES pte.MICA.16.324097 (FAM_4968) contains a complete protein-coding gene, which is expressed during development (HTH CenpB-type DNA-binding domain see Fig 5A). The gene in which this IES is inserted (PTET.51.1.P0160202) is not specifically expressed during development.

S6 Table. Genetic distances between species of the *aurelia* complex. This matrix reports for each pair of species, the number of orthologous genes analyzed (above the diagonal), and their median synonymous divergence (below the diagonal).

S7 Table. Sequencing data generated for this study.

1655 **S1 Data. Flow cytometry-based estimations of MIC genome size in *Paramecium*.**

1656

1657 **S2 Data. Locations of MAC-variable regions and MAC assembly curation**

1658 This file provides the positions of MAC-variable regions identified in the MAC assemblies of

1659 *P. octaurelia, P. pentaurelia, P. primaurelia,* and *P. sonneborni*. In addition, it indicates the

1660 positions of putative assembly chimeras that have been identified in *P. octaurelia*, *P.*

1661 *primaurelia* and *P. sexaurelia*.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure S1

Figure S2

Figure S3

B
```
>Sat1: Caudatum_satellite_1 (332 bp)
CATCAAACTTTCTTTTCCTTATGACAAGTCAATTGACATTTTTCGTCAATTTTATGAGGA
GTCATAGGTACCTAATATTTGTGCTGTAGGGCTCAATAATAGCTCATTTTAATGCTGGTA
CAGAGGGCTTTAAAATAGCCATCTCAGTTTTTCTTAATGGAATCTAGATCGCCAGAACGG
GCATTTCATGACTTTTTTAATTATATCATATCGATGGTCATTTCATCTCCAAAGTGTCTA
AAAACACGTTATTCGTTGTCAGCGTCGAATTTTGAGTAAAACTGTCATCTCACTTTTCCT
TAATGGAATTTATTTGAGGAGATCCAATTTCC

>Sat2: Caudatum_satellite_2 (449 bp)
TTTTTGAGATAAAGAACTCGGGTTTTTTTCTATATAATACTGGTACCAAACGCTACAAAT
CGTATGCCTCAGATTTTCTTATTGTTAGCTAAATCCCTGAGAACGCTCATTTTCATCTGG
AAAGTCATCGGGTTTGGTATTTCGCCTGAAAAAATATAAAAAATTTTTGAGATAAAGAAC
TCGGATCTTTTTCTATATAATACTGGTACTAAAAGCTGCAAATCGTATAATCTCGATTTT
CTTAATGGATGCTAAATCCCGAGAACGGCCATTACAAGGGAATTTTAATTTTGCAAAATC
GATGGTTTTTTGATGTCCTACGTTGCTAAAAACATGTTTTATGATATCAGGAGCCCATTT
TACCCCTAATCCACTCTTTTATAGACCTTAATGGAATTTAGTACCCGAGATCCAAAATCA
GCAGCAAATAGCATTTTTCATAGGGAATT
```

Figure S4

Figure S5

Figure S6

Figure S7

Figure S8

Figure S9

Figure S10

Figure S11

Figure S12

MAC: TCCTAATTCAATTGT**TATA**ATATATTATTAAA

MIC: TCCTAATTCAATTGT**TATA**GAATGAATATTCAATTATAATTA**TATA**ATATATTATTAAA

IES-1

IES-2

Figure S13