Variation and process of life history evolution in insular dwarfism as revealed by a natural experiment

Shoji Hayashi[1,2,*,**], Mugino O. Kubo[3,*,**], Marcelo R. Sánchez-Villagra[4], Hiroyuki Taruno[5], Masako Izawa[6,7], Tsunehiro Shiroma[6], Takayoshi Nakano[2], and Masaki Fujita[8]

[1] Faculty of Biosphere-Geosphere Science, Okayama University of Science, Kita-ku Ridaicho 1-1, Okayama, 700-0005, Japan

[2] Division of Materials and Manufacturing Science, Graduate School of Engineering, Osaka University, Yamada-Oka, Suita, Osaka, 565-0871, Japan

[3] Department of Natural Environmental Studies, Graduate School of Frontier Sciences, The University of Tokyo, 5-1-5 Kashiwanoha, Kashiwa, Chiba 277-8563, Japan

[4] Paläontologisches Institut und Museum der Universität Zürich, Karl Schmid-Strasse 4, CH-8006 Zürich, Switzerland

[5] Osaka Museum of Natural History, Nagai Park 1-23, Higashi-Sumiyoshi-ku, Osaka, 546-0034, Japan

[6] Faculty of Science, University of the Ryukyus, Okinawa 903-0213, Japan

[7] Kitakyushu Museum of Natural History & Human History, 2-4-1 Higashida, Yahatahigashi-ku, Kitakyushu 805-0071, Japan

[8] Department of Anthropology, National Museum of Nature and Science, 4-1-1 Amakubo, Tsukuba-shi, Ibaraki 305-0005, Japan

* Correspondence:

Shoji Hayashi and Mugino O. Kubo

** Both authors contributed equally to this work and share the first authorship.

Email: shoji-hayashi@ous.ac.jp (S.H.), mugino@k.u-tokyo.ac.jp (M.O.K)

Keywords: bone histology, deer, demography, growth rate, life history

Abstract

Islands are a classic focus for evolutionary studies. One subject of much interest has been the evolution of "dwarfs", significantly smaller island mammals relative to their

continental counterparts. Although a consensus has been achieved that multivariate ecological causes are behind body size changes, the processes involved remain largely

unexplored. Life history variables, including the age of first reproduction, growth rate, and longevity, are probably key to understanding the process of insular dwarfism. The

Japanese Archipelago with numerous islands offers a unique natural experiment of evolution into different sizes within the same group of organisms, deer. Thus, we

investigated eight deer populations with a total number of 52 individuals exhibiting body size variation, both extant and fossil, to clarify the effect of insularity on life

history traits. We applied several methods to both extant and extinct populations to resolve life history changes among deer populations. Skeletochronology, using lines

of arrested growth formed in long bones (femur and tibia), successfully reconstructed body growth curves and revealed a gradual change in growth trajectories reflecting

the degree of insularity. Slower growth rates with prolonged growth periods in more isolated deer populations were revealed. An extensive examination of bone

microstructure further corroborated this fact, with much slower growth and later somatic maturity evident in fossil insular deer isolated for more than 1.5 Myr. Finally,

mortality patterns assessed by demographic analysis showed variation among deer populations, with a life history of insular populations shifting toward the "slow life".

Introduction

Among the most iconic examples of visible evolutionary changes are island mammals that dramatically changed in body size [1-3]. While elephants, hippos, and

deer on islands across the world became small, rodents and other small mammals became larger [2]. These changes are which led to the suggestion of the "island rule"

[1]. Classic examples are found in fossil species: the pygmy elephant (*Elephas falconeri*) of Sicily, which has a body mass only 1% of its mainland ancestor [4], and

a gigantic gymnure (*Deinogalerix koenigswaldi*) of Miocene Gargano having a body mass about 200 times heavier than its ancestor [5]. This leads to a hypothesis of the

optimal body size of mammals on isolated islands, although the particular optimum of an insular population varies with the characteristics of the islands and the species, and

with interactions among species [5, 6].

Moreover, since body size has a close relationship with various aspects of

68   organisms' life, recent studies have focused on the relationship between body size and

life history, including growth rate, age of reproduction, and longevity on islands [7-9].

70   A reduction in the body size of large mammals has also been associated with a

modification of the growth trajectory [10, 11], which in turn affects the lifespan, e.g.,

72   the timing of sexual maturity or longevity. Life history theory, which extended from

the r/K selection theory in ecology [12], proposed the idea that mammalian

74   populations can be placed along a fast-slow continuum [13]. Species that adapted to

unstable environments, under which high intrinsic rates of natural increase (r) were

76   adaptive, evolved a set of life history traits, such as rapid body growth, early sexual

maturity, many offspring for a clutch, and short life span, occupying the "fast" end of

78   the continuum (formerly referred as "r-selected"). On the other hand, in stable

environments, the populations are maintained near their carrying capacity (K), where

80   resultant intraspecific competitions favor those which have slow body growth, late

sexual maturity, a small number of offspring for a clutch, and a long life span,

82   occupying the "slow" end of the continuum ("K-selected"). Based on this theoretical

background, life history traits were estimated for fossil insular species. Such studies

84   have been conducted by applying histological analysis of bones developed to

reconstruct the paleoecology of dinosaurs and other Mesozoic vertebrates [7, 14-16],

86   the method being further recognized by mammalian paleontologists as a useful tool to

infer life history traits in mammals [8, 17-22]. Köhler and Moyà-Solà [8] showed that

88   in a fossil dwarfed bovine *Myotragus balearicus* from the Balearic Islands of the

Mediterranean, its long bone histology showed lamellar-zonal tissue throughout the

90   cortex, which is characteristic of ectothermic reptiles. The bone microstructures

indicate that *M. balearicus* is estimated to have had a low metabolism and a very slow

92   growth rate. This life history shift toward a slow life in *M. balearicus* was further

corroborated by analyses of tooth enamel microstructure, tooth eruption and mortality

94   pattern, which implied delayed somatic maturity, lower mortality and extended

longevity of this species [9, 23, 24]. Followed these paleohistological analyses were

96   fossil mammals from other Mediterranean islands: e.g. a fossil cervid (*Candiacervus*)

from Crete showing a slight shift toward a slower life history [18, 25], and a fossil

98   dwarf hippo (*Hippopotamus minor*) from Cyprus with no evident modification in

bone histology [19]. The bone histological observations and life history implications

100   of fossil dwarfs have been accumulated in the last decade, nonetheless, these studies

were based on different species from single islands, therefore it was hard to generalize

102   the evolution of body size and life history on islands. Multiple comparisons of closely

related species or populations from both mainlands and islands are crucial to drawing

104   general conclusions.

Further important issues awaiting to be explored are the process of body size and

106   life history evolution [26], including how ecological factors affect processes, such as

the lack of predators and competitors, resource limitation on islands, founder effects

108   of island immigrants, and the remoteness and duration of isolation [2, 5, 6, 27, 28].

Drastic body size changes have been reported for fossil species or populations [2].

110   Nevertheless, as far as we focused only on fossil evidence, ecological processes

undergone during these changes remain unclear because of the lack of detailed

112   ecological information. To achieve this change, again, studies on the mainland and

insular populations of a single extant species are particularly useful, as they provide

114   ecological contrasts between focal populations and allow confounding factors other

than living on islands to be controlled. Variation in isolation durations among the

116   populations can therefore mimic the process of insular settlement.

In this study, we focused on deer in the Japanese Archipelago, both extant and

118   extinct populations from both islands and the mainland (Fig. 1). Our targets cover an

extant sika deer (*Cervus nippon*) population that was introduced recently (c.a. 400

120   years ago) on a small island, four extant insular and mainland deer populations [sika

deer, Reeves's muntjac (*Muntiacus reevesi*)] with variable body sizes, and extinct

122   insular deer populations (*Cervus astylodon* and Muntiacini gen. et. sp. indet., referred

to as "the Ryukyu muntjac") that were isolated for more than 1.5 Myr on the Okinawa

124   Island [29], representing different degrees of insular effect. We also included an

extinct gigantic mainland deer (*Sinomegaceros yabei*) from the Late Pleistocene

126   deposits on the Honshu mainland, having a comparable body size to the largest extant

deer; the moose (*Alces alces*) [30]. The immense variation of the island's size and

128   duration of isolation observed in the eight understudied deer populations enabled us to

infer the process of body size and life history changes on islands.

130   This study aims to test whether the different degree of insular effect results in the

change of life history traits among Japanese deer populations. Specifically, we

132   hypothesize that a gradual change in life history traits of deer populations is expected

according to the intensity of insular effects, with longer isolation on smaller islands,

134   resulting in a shift toward the more pronounced slow life. For this objective, we first

applied skeletochronology to long bones, using lines of arrested growth (LAGs)

136   within the primary periosteal bone tissue, to depict body growth rates. Obtained

growth curve parameters were statistically compared among deer populations to test

138   the hypothesis. Subsequently, detailed qualitative descriptions of bone histology were

conducted aiming to corroborate the skeletochronological findings on growth patterns

140   and somatic maturity. We focused on periosteal bone tissue and vascularization

pattern as an indicator of the relative bone growth rate [8, 31]. Lastly, we conducted

142   demographic analyses of age data, which were collected either through ecological

surveys or analysis of tooth wear obtained from fossil deer assemblages, to

144   quantitatively compare mortality patterns among deer populations.

146   **Materials and Methods**

    1. Locality and specimen information for the sampled deer used in the
148   **histological analyses**

    For histological investigation and estimation of growth curves from LAGs, we used

150   femora and tibiae of extant and fossil deer. All extant deer were wild individuals and

their skeletal specimens were stored in museums. Both tibia and femur were sampled

152   from the same individual for the extant deer. On the other hand, all fossil specimens

were excavated in disassociated conditions. Therefore, we considered that each of the

154   femora or tibiae originated from a different individual. A summary of the sampled

populations together with the sample size is provided in Table 1.

    1-1. Extant deer populations

158   Sika deer from Hokkaido mainland: all individuals (N = 7) were hunted in Hokkaido

Prefecture (43°N, 143°E), northern Japan (Fig. 1). Hokkaido is one of the four main

160   islands constituting the Japanese archipelago and was considered "mainland" for this

study. Hokkaido sika deer inhabit subarctic coniferous forests mixed with cool

162   temperate deciduous trees and are known to be the largest subspecies (*C. nippon*

yesoensis [32-34]). The complete skeletons are housed in the Hokkaido University

164   Museum (HOUMVC), Sapporo, Japan. The age at death was recorded in the museum

inventory list, which was assessed based on tooth eruption for fawns and yearlings

166 and the tooth cementum annuli formed at the roots of the incisors for adults.

168   Sika deer (N = 8) and Reeves's muntjac (N = 5) from Honshu mainland: all

individuals were hunted or collected in Chiba Prefecture (35°N, 140°E, Fig. 1), which

170 is in the central region of Honshu. Honshu is one of the four main islands constituting

the Japanese archipelago and was considered "mainland" for this study. Both of these

172 species inhabit evergreen broadleaved forests in the temperate zone. Reeves's muntjac

was introduced to Chiba Prefecture ca. 1960–1980 following an escape from a private

174 park [35]. The source population is unknown, but the body size of these deer is

smaller than the continental population but closer to that of the Taiwanese population

176 [35]. It is not clear whether Reeves's muntjac studied here or those in Taiwan are

dwarfed populations compared with their counterpart in the continent. The complete

178 skeletons of these individuals are stored at the Natural History Museum and Institute,

Chiba (CBM), Japan. The age at death was recorded in the museum inventory list,

180 which was assessed based on tooth eruption or the cementum annuli for the sika deer

and from tooth eruption and the molar wear condition for Reeves's muntjac.

   Sika deer from Yakushima Island: all individuals (N = 4) were hunted or collected

184 on Yakushima Island (30°18′ N, 130°30′ E, Fig. 1) in Kagoshima Prefecture, Japan,

where they inhabit subtropical evergreen broadleaved forests. Yakushima Island deer

186 are considered to be the subspecies *C. nippon yakushimae* [33] and exhibit

considerable phenotypic differences from the mainland populations, such as small

188 body size, simplified antlers, and shortened limbs [33, 36]. Yakushima deer diverged

from the population on Kyushu Island (one of the four main islands constituting the

190 Japanese archipelago) ca. 0.1 Mya when the Ohsumi Strait was formed between the

two islands [37]. Yakushima Island was possibly connected to Kyushu Island by a

192 land bridge during the Last Glacial Maximum (ca. 25,000 ya), but the genetic

differentiation between the two populations implies a much longer period of isolation

194 for Yakushima deer [38, 39]. The complete skeletons are stored at Tochigi Prefectural

Museum (TPM), Utsunomiya, Japan. The age at death was recorded in the museum

196 inventory list, which was assessed based on tooth eruption for fawns and yearlings

and the tooth cementum annuli of the incisors and molars for adults.

Sika deer from Kerama Islands: all individuals (N = 7) were collected as carcasses
resulting from natural deaths on Kerama Islands (26°12′ N, 127°24′ E, Fig. 1) in

Okinawa Prefecture, Japan, by two of the authors (MI and TS) during their fieldwork.
Sika deer were introduced to Kerama Islands from Kyushu ca. 400 years ago possibly

from Kagoshima Prefecture, Kyushu mainland, Japan [40], and inhabit subtropical

evergreen broadleaved forests. These deer exhibit the phenotypic characteristics of
insular dwarf deer (similar to Yakushima Island deer) having smaller body size than

the possible source population of Kagoshima [40] and so are also regarded as a
subspecies of sika deer (*C. nippon keramae*) [33]. Most individuals were complete

skeletons, but some bone elements were lost at the time of collection in the field for
some individuals. All of the specimens are stored at the Faculty of Science, University

of the Ryukyus (URB), Nishihara, Japan. The age at death was recorded in the
inventory list, which was assessed based on tooth eruption for fawns and yearlings

and the tooth cementum annuli for adults. However, no reliable age at death was
obtained for an individual URB-MAM-34.

1-2. Fossil deer populations

Cervus astylodon (femur N = 6, tibia N = 5) and Muntiacini gen. et sp. indet. (the
Ryukyu muntjac, femur N = 4, tibia N = 2): both of these fossil deer species were

excavated from the Hananda-Gama Cave site (26° 8′ 7″ N, 127° 45′ 32″ E, Fig.
1) in the southern part of Okinawa Island [41]. These animal remains are mainly

isolated materials consisting of limb bones and teeth and are considered to have fallen
into the cave through fissures or sinkholes following the decomposition of the

carcasses [41]. The detailed geological age of these fossils is not clear due to a lack of
suitable specimens for radiocarbon dating. However, based on radiocarbon dating at

other sites that have yielded fossils of these species, the geological age of both species
has been estimated as the Latest Pleistocene (>20 ka, [41]). *C. astylodon* has been

recognized as a typical insular dwarf due to its very small body size and shortened
limbs [2]. The Ryukyu muntjac used to be referred to as *Dicrocerus* sp. [2], which is a

genus that is known mostly from Miocene Europe. However, the finding that the
antlers have a clear burr and grow from the posterior parts of the frontals oriented

backward clearly indicates that it does not belong to this genus (Gertrud Rößner, personal communication to MOK). Rather, it appears to be closely related to

Muntiacus based on the presence of tusk-like canines and a lack of antlers in the females, neither of which is found in *Dicrocerus* [42]. A full revision of this species is

required, including a detailed description and comparisons with other fossil and extant taxa. At this moment, we did not have any reliable information on its ancestry and the

degree of dwarfism, though the estimated body mass for this species implied that its body size seemed to be comparable to the continental extant muntjac species. In this

paper, we present it as Muntiacini gen. et sp. indet. and refer to it as "the Ryukyu muntjac." All of the specimens that were examined are stored at Okinawa Prefectural

Museum and Art Museum (OPM), Naha, Japan.

Sinomegaceros yabei (femur N = 2, tibia N = 2): this fossil deer had a gigantic body size that was comparable to that of Irish elk (*Megaloceros giganteus*) [30, 43]. The

skeletal remains were excavated from the Kumaishi-do Cave site (35°45″ N, 137°′ 00″ E, Fig. 1) in Hachiman-cho, Gujo city, Gifu Prefecture, central Japan. Most of

the animal remains obtained from this site are isolated materials, but an incomplete skeleton that includes the skull has also been discovered [43]. The geological age of

these fossils is the Late Pleistocene based on radiocarbon dating of the bones of cervids and proboscideans [44]. All of the specimens we sampled are stored at the

Osaka Museum of Natural History (OMNH), Osaka, Japan.

Detailed information on the extant and fossil specimens is provided in Supplementary Information (SI) Tables S1–S3.

2. Thin sectioning and X-ray computed tomography (CT) scanning

256 We applied histological analyses of long bones to infer life history traits [8, 17-22]. We used 62 extant and 21 fossil long bones for histological analyses (Table 1). All

258 specimens were then photographed and standard measurements were taken, following which thin sections of the midshaft of long bones were prepared based on the

260 methodology described in previous studies [45, 46]. We produced 83 thin sections (62 sections for extant species, and 21 sections for extinct ones; see also detail in SI Table

262 S1–S3). The thin sections were photographed using a digital film scanner (Pixus Mp

800, Canon) and analyzed using an Optiphot2-pol microscope (Nikon). Microscopic photographs were finally taken with a Nikon Df camera.

Neonatal specimens of sika deer (Honshu mainland: CBM-ZZ-757) and Reeves's muntjac (CBM-ZZ-4974) were scanned using an experimental animal X-ray CT scanner (Latheta LCT-200, Aloka; 24-μm resolution, 80 kV, 0.2 mA) at Okayama University of Science, Okayama, Japan, to determine the initial diameter of the cortical bone and open medullary cavity in these species. Afterward, image segmentation and visualization were conducted using VG-Studio Max (Volume Graphics) version 3.1.

3. Age assessment

To allow for the comparison of growth patterns, we estimated ages as follows: for most of the extant deer samples, age at death was recorded in the museum inventory lists, which were assessed using tooth eruption or the number of dental cementum annuli in the tooth root (SI Tables S3 and S4). However, with fossil deer samples, long bones were unassociated with skulls and mandibles. Therefore, it was required to estimate the age at death from the number of LAGs found in the long bones. Several previous studies have reported a good correlation between the number of LAGs and the age at death in several vertebrate groups [47], while other studies indicated the limitation of LAGs as reliable ages due to loss of LAGs by bone remodeling and/or the expansion of the medullary cavity [e.g. 25, 48]. A study examined this correlation in extant deer species [red deer (*Cervus elaphus*)], which reported that although the number of LAGs in the tibia corresponded to the actual age, the number in the femur corresponded to the age before the deposition of the EFS [49]. Therefore, the relationship was examined first using extant deer between the actual age at death and the number of LAGs in bones (see SI for details). Results showed that the estimated age of the extant deer based on the number of LAGs, including those that appeared in EFS in both the femur and tibia was highly correlated with the actual age determined using the tooth cementum annuli (SI Figure S6 and Table S4). Furthermore, while some old adults who were over 8 years (HOUMVC-00037, CBM-ZZ-412, and URB-MAM-193) had fewer LAGs in their long bones than their actual age, diameters of their expanded medullary cavities were identical or even larger than bone diameters

observed in neonates and fawns (<1 year old), proposing that the first (and also

296   second in URB-MAM-193) LAG was eliminated in old individuals (SI Table S4).

Thus, since the number of LAGs matched the actual age in extant sika deer, by

298   counting the number of LAGs in their long bones, the age of individuals of the fossil

deer was also estimated (SI Table S5). Secondary bone remodeling, which had erased

300   any evidence of the primary bone tissue including LAGs, were seen only at the inner

medullary surface and part of the cortex where ligaments were strongly attached (e.g.,

302   the labium laterale in the femur) in adults, indicating a good preservation of the

primary bone tissues in most parts of the cortex.

4. Body mass estimation

306   The body mass during ontogeny was estimated from bone diameters at the LAGs

[50]. Body mass estimation formulas for both the femora and tibiae were obtained and

308   the formula that uses the mediolateral diameter (MLD) of the femora and tibiae was

selected for use in this study (see SI for details). In addition to diameters at the LAGs,

310   external diaphysial measurements [anteroposterior diameter (APD) and MLD] were

collected and used to estimate body mass at death. Raw measurement data of

312   diameters at the LAGs, with the estimated body mass used for growth curve

modeling, are presented in Supplementary Table S13 (separately provided in excel

314   format).

316   **5. Comparison of growth patterns**

Growth curves of body mass were constructed for each specimen that was older than

318   yearling (i.e., those that had more than one LAG), which had at least four data points,

including starting and terminal points. The exception to this was the extant Reeves's

320   muntjac, for which a fawn and yearling were included in the growth curve fitting

because this species attained maturity early in life [35]. Before fitting the growth

322   curves, neonatal body mass data were obtained for both the extant and fossil taxa as

the starting point (SI Table S6 and Figure S7).

324   To fit growth curves to the obtained data, the number of LAGs was first transformed

into the age in years. For each specimen, the body mass (W, in kg) was then

326   formulated with age (x, in years) using the Gompertz curve, which is described below:

$$W = aExp(-Exp(-b(x - c))$$

328   where a, b, and c represent the asymptote (i.e., the body mass when body growth

ceases), growth rate, and inflection point, respectively. The Gompertz curve fitting was rationalized as this model showed the highest value of goodness of fit represented by the lowest AICc among the fitted growth curves (SI Table S7). Furthermore, the growth rate (b) and inflection point (c) were statistically compared among species/populations. First of all, the normality of the parameter distribution was confirmed by the Shapio-Wilk tests (for each species/population, $P > 0.05$). We then compared the growth rate and inflection point between the femur and tibia using extant deer samples by paired t-test. We found that the null hypothesis of no difference in these parameters between the femur and tibia was not rejected (for the growth rate, N = 15, $t = 0.10$, d.f. = 14, $P = 0.92$; for the inflection point, N = 15, $t = 0.60$, d.f. = 14, $P = 0.56$). Therefore, to increase the sample size we combined data from the femur and tibia for fossil samples, though we admit that further data are needed to clarify the difference in growth curves between the femur and tibia. We conducted one-way ANOVA and subsequent multiple comparisons using the Tukey-Kramer method to examine differences in the growth parameters among the eight species/populations, averaging the growth curve parameters from the femur and tibia for the extant species to avoid repeated sampling from the same individual. Additionally, two-way ANOVA using bone type (i.e., femur or tibia) and species/populations as the factors was conducted. Growth curve parameters were averaged for each population/species to produce a representative growth curve for each group. All statistical analyses were conducted in JMP Pro 16.0 (SAS Institute Inc.).

6. Qualitative assessment of bone histology and external fundamental system (EFS) observation

To corroborate findings derived from estimated growth curves, we additionally investigated bone histology in detail. We observed the same histological sections used for the growth curve modeling. The nomenclature and definitions of bone tissues were based on the study by Francillon-Vieillot *et al*. [51] and Castanet *et al*. [52]. We focused on the following bone tissue characteristics: 1) the type of primary periosteal bone tissue and vascularization pattern as an indicator of the relative bone growth rate and 2) the presence of an external fundamental system (EFS). A primary periosteal bone tissue known as a fibro-lamellar (FBL) bone, which has a high level of laminar to plexiform

362 vascularization (PV), is an indicator of fast-growth, whereas a parallel-fibered (PF)
bone tissue is formed during periods of slower growth [e.g. 31]. Growth rate correlates

364 with intensity and vascularization patterns [16, 53, 54], where lower levels of
vascularization associated with the longitudinal orientation of canals indicated a slower

366 bone depositional rate. Additionally, EFS is characterized by the PF bone tissue
constituting without vascularization, thereby forming an edge of slow-growing bone

368 tissues as a sign of somatic maturity [e.g. 48, 55, 56]. By counting LAGs within EFS,
we estimated the ages at which EFS formation started (see SI for details). Somatic

370 maturity was additionally investigated by the status of fusion of both proximal and
distal epiphyses. Lastly, we conducted correlation tests between the age at which EFS

372 development started and the inflection point (c), which marks the transition from the
initial phase of exponential growth to the subsequent phase of asymptotic growth and

374 is known to be correlated with the timing of sexual maturity in laboratory-reared mice
[57]. For this purpose, we used the mid-values of EFS developmental age for each

376 specimen. For example, OPM-HAN06-588 (*Cervus astylodon*) had the range of 9 – 10
years, therefore we used the value of 9.5 years as the age at which EFS development

378 started. As for the extant specimens, we averaged the inflection point obtained from the
femur and tibia growth curves. Non-parametric Spearman's rank correlation coefficient

380 (ρ) was calculated in three datasets (extant only, fossil only, and all individuals
combined).

384 **7. Comparison of survivorship curves**
     Since a change in mortality patterns also represents life history changes, we

386 estimated survivorship curves of both extinct and extant deer populations. We
obtained age data for the three extant sika deer populations that were used in the

388 histological investigations either from published papers or through investigation of the
cementum annuli of the incisor roots: Hokkaido mainland (N = 1,060) [58], Honshu

390 mainland (N = 594) [59], and Yakushima Island (N = 74, cementum annuli of
museum specimens). Histological analysis of cementum layers of the lower first

392 incisors was conducted on Yakushima Island deer specimens housed in the University
Museum of the University of Tokyo, the Tochigi Prefectural Museum, and the

394 Hokkaido University Museum to identify their age at death. This analysis was done
by a specialized laboratory (Matson Laboratory, Montana, U.S.A.) and we received

age data with high reliability. The age data from three populations were based on randomly culled individuals and so were considered to be indicative of the age structure of actual extant populations. Because the hunting of female deer had been legally prohibited until 1994 and these deer populations were increasing in number at the time of culling (approximately from the late 1990s to early 2000s), the deer populations were less affected by the hunting pressure [59]. Additionally, we obtained mortality data of sika deer from Kinkazan Island from the published literature (N = 264) [60]. The Kinkazan population is protected for religious reasons and thus free from hunting pressure, which was also the case for *C. astylodon* and the Ryukyu muntjac inhabiting the Pleistocene Okinawa Island without human and other predators [60, 61]. The age data for extant Reeves's muntjac from England was obtained from a published reference (N = 84)[62].

   The age data of two fossil insular deer from Okinawa Island were also obtained from published references (*C. astylodon*: N = 45 [60]; the Ryukyu muntjac: N = 65 [63]). Kubo et al. [60] estimated the age at death for *C. astylodon* based on the height of the lower third molar by applying the molar wear rate of extant sika deer. Variability of age estimation based on the molar wear rate was tested and discussed in previous studies [59, 60, 64], after which we adopted the most appropriate age estimates. Ozaki [63] estimated the age at death of the Ryukyu muntjac by applying the molar wear scoring method developed by Chapman *et al.* [62] for extant Reeves's muntjac. Fossil *S. yabei* and extant Kerama Islands deer were excluded from these survivorship curve comparisons due to a lack of reliable age data.

   To depict the survivorship curves of three culled sika deer populations, we constructed life tables following the method described in a reference [65], assuming stationary age distributions, and applied probit regression to smooth the age distributions. The age data for the other populations (Kinkazan sika deer, Reeves's muntjac, and two fossil insular deer) were based on natural deaths, therefore the age class frequency was divided by the sample size to obtain a multiple of the frequency of mortality (dx). Life tables were then derived from this mortality frequency [65]. The survivorship curves were presented with an initial value of 1,000 individuals.

   The observed age class frequencies and derived life tables for each deer population are shown in SI Tables S11 and S12.

For more details on the materials and methods used in this study, see the SI and

430   Supplementary Figures and Tables therein.

432   **Results**

Growth curve models and growth rate comparison using skeletochronology

434   The LAGs in the long bones of both extant and fossil deer were well preserved (SI

Figure S1–4), allowing us to measure bone diameters at the LAGs. A growth curve of

436   each species/population is presented in Figure 2A and SI Figure S10. Among the sika

deer populations, deer from the two mainland areas (Honshu and Hokkaido mainland)

438   showed faster growth than those from the two insular populations (Yakushima and

Kerama Islands). Growth trajectories of body masses also corresponded with those

440   obtained from investigations of culled individuals [66, 67] and through long-term field

observations of wild deer (Agetsuma and Agetsuma-Yanagihara, personal

442   communication), thereby validating our estimation method of body growth from

skeletochronology. The extant Reeves's muntjac showed the fastest growth, which was

444   reported as well in the analysis of culled individuals that showed early sexual maturity

at ca. 6 months [35].

446   Compared to the extant references, two fossil insular deer species (*C. astylodon* and

the Ryukyu muntjac) showed remarkably slow and prolonged growth, whereas the

448   Pleistocene giant deer (*S. yabei*), had fast growth, which suggested a growth pattern

similar to other extant large deer species, such as moose [68].

450   Statistical comparisons of growth parameters revealed that both fossil and extant

insular deer had significantly slower growth rates than Reeves's muntjac and *S. yabei*

452   (both $P < 0.05$, SI Tables S9, S10). However, growth rates of the four extant sika deer

populations and fossil insular deer were not significantly different from each other (P

454   > 0.05, SI Tables S9, S10), due to the transitional position of extant insular deer,

although the limited numbers of possible samples may also play a role.

Variation in bone tissue among deer populations

458   Histological characteristics of long bones corroborated the above statistical

comparisons of growth curves. Figure 3 summarizes the main histological features of

460   examined deer (also see SI Tables S1–S3).

As observed, all extant deer populations (i.e., the four extant sika deer populations,

462   the extant Reeves's muntjac), including the gigantic fossil *S. yabei*, exhibited a similar

bone tissue structure to continental cervids [18, 48, 49], consisting of a fast-growing FBL bone with PV in most parts of the cortex (Figure 3G–L; SI Figures S2A–I, S3G–I, and S4A–O). Additionally, slow-growing PF bone tissue was observed in the outer cortex. This PF bone tissue was more extensively and notably developed in the two insular populations (sika deer from Yakushima and Kerama Islands), with LAGs at the inner to the middle cortex being tighter than in the mainland deer (Figure 3D–F, SI Figures S2D–F, and S3G–I, J–L), suggesting a slower growth rate among insular populations. Furthermore, the cortical bone tissues of two fossil insular deer (*C. astylodon* and Ryukyu muntjac) were different from those of extant deer and gigantic fossil deer (*S. yabei*), in that they were primarily PF bone tissue with multiple LAGs, thereby implying a much slower growth rate than other deer populations. Their bone tissues were also similar to those of the extinct dwarf bovid (*Myotragus balearicus*) and typical extant reptiles [e.g. 69]. Moreover, although LAGs were spaced evenly throughout the cortex (Figure 3A–C; SI Figures S1A–C and S3A–F), they became closer as they approached the periosteal surface in older individuals, which indicated a decrease in growth rate with age. Only a few areas of the innermost cortex showed a prevalence of the FBL bone tissue with primary osteons, which alternated with nonvascular PF bone tissue with LAGs, suggesting that the bone was deposited slowly during most periods of the lifetime. The slow growth rate was further supported by the observed vascularization pattern. Our fossil insular deer samples showed less vascularization and their vascular canals oriented longitudinally, which is not observed in other cervids and large mammals [18, 22]. In summary, the bone histological features of the fossil insular deer supported their slow growth rates.

Results also showed that all extant specimens with EFS had fused epiphyses both in the femora and tibiae (SI Figure S5, Table S3). As observed, the proximal epiphysis of the tibiae and femora fused later than the distal epiphysis. Moreover, EFS developed contemporaneously with proximal fusions. In two mainland sika deer populations, the timing of EFS development and proximal epiphyseal fusion showed a good correspondence with the age when the rate of growth decelerated (SI Figures S8, S9, Table S3), including the reported age of somatic maturity in the analyses of culled individuals [66, 67]. Notably, the extant sika deer from Kerama Islands also showed delayed EFS development and epiphyseal fusions compared with other investigated deer populations. Based on the observation of extant specimens, we further investigated

496   the relationship between EFS development and epiphyseal fusion in fossil specimens
(SI Figure S5, Tables S1, 2). Though most fossil specimens did not preserve both

498   proximal and distal epiphyses, all specimens with fused proximal epiphysis had EFS
and vice versa. Therefore, the correspondence of epiphyseal fusion and EFS was also

500   held in fossil species as well. Findings revealed that two insular extinct deer showed
much delayed EFS development (Min. 9 years, Max. 16 years for *C. astylodon*; Min. 5

502   years, Max. 11 years for the Ryukyu muntjac, see SI Tables S1, S2) than other
investigated deer populations, suggesting a delay of somatic maturity.

504   Histological observation of delayed somatic maturity was further corroborated by the
growth curve parameter. We found a significant positive correlation between the

506   inflection point and the age at which EFS development started (extant only, $\rho = 0.87$, *P*
$= 0.0012$; fossil only, $\rho = 0.84$, $P = 0.0003$; all data combined, $\rho = 0.91$, $P < 0.0001$; SI

508   Figure S11). Though the inflection point was younger than the actual age when the body
growth deaccelerated and the age at which EFS development started, the significant

510   positive correlation indicated that the higher values of the inflection point can be used
as an indicator of delayed somatic maturity. Statistical comparison of inflection points

512   among studied deer showed that the inflection point was significantly greater in *C.*
astylodon than in other deer populations ($P < 0.05$, SI Tables S9, S10), implying again

514   that sexual maturity can be delayed in this species.

516   **Survivorship curve comparison among extant and fossil deer**
     To clarify life history differences in mortality patterns, we compared the survivorship

518   curves of fossil *C. astylodon* and four other extant sika deer populations from the
Hokkaido mainland, Honshu mainland, Yakushima Island, and Kinkazan Island (Figure

520   2B). Since the age estimation of *C. astylodon* was based on the lower third molar height,
we confined survivorship curves to juvenile and onward for *C. astylodon* and the sika

522   deer. A low mortality from the juvenile through to the prime age period characterized
the survivorship curve of *C. astylodon*, followed by an increase in mortality during

524   senescence, which appeared like an inversed L, and was categorized as a type I
survivorship curve [70], including a characteristic of slow life species. At the other

526   extreme, extant sika deer from the Hokkaido mainland showed relatively high mortality
in the younger age class, which was followed by a more gradual decrease in the number

528   of survivors from the prime age to senescence, thereby serving as a characteristic of
type II survivorship curves. Results also showed that survival curves of sika deer from

the Honshu mainland and Yakushima Island were located between those of *C. astylodon* and the Hokkaido sika deer, with a steady decrease in survivorship until the

prime age, though the Honshu population showed an increased mortality rate after the prime age. Also, although the survivorship curve of the Kinkazan Island deer was

clustered with Honshu and Yakushima Island populations, a lower mortality rate in the initial phase and a higher rate in the later phase was shown. This slight difference in the

pattern is proposed to be associated with the lack of hunting pressure in the Kinkazan population. However, the intensity of hunting in the sika deer does not strongly affect

overall survival curves as described in the Methods section. Furthermore, though the Kinkazan deer inhabited the predator-free island, their mortality pattern was different

from that of the *C. astylodon*, implying a further advanced insular effect in the latter. A previous study reported that the maximum observed age of *C. astylodon* was 25 years

[60], which was greater than that of the extant sika deer (17, 19, 17, and 21 years for the Honshu, Hokkaido, Yakushima Island, and Kinkazan Island populations,

respectively; SI Tables S11, 12).

    We additionally compared survivorship curves between the Ryukyu muntjac and the

546   extant Reeves's muntjac (Figure 2C). As observed, the extant muntjac had a type II survivorship curve as in the extant sika deer from the Honshu mainland. Similarly, the

548   Ryukyu muntjac had a type I survivorship curve, indicating that its mortality rate did not also increase until senescence, thereby implying a strong insular effect on life

550   history traits in Okinawa Island again.

552   **Discussion**

    All interdisciplinary analysis results from growth rate comparison, histological

554   investigation, and analysis of survivorship curves of both extant and extinct deer species supported our expectations that the evolution of a new combination of life

556   history traits amounts to repeated establishments of a "slow life" in insular environments. Though the sample size of deer for each population was not very large,

558   our findings are among the first to clarify the gradual shift to slow life histories for insular forms, placing the two fossil insular deer populations from Okinawa Island at

560   the slow end of the fast-slow continuum.

    It is possible to infer ecological processes of life history change from the current

562   dataset obtained from deer populations. As observed in mainland deer populations,

any modification of life history traits happened in mainland populations (Hokkaido and Honshu mainland sika deer, Reeves's muntjac, and *S. yabei*), similar to the condition of continental relatives [18]. The Japanese Archipelago is considered "an island" in literature [e.g. 2]. Nevertheless, from the current dataset, the Japanese mainland deer is proposed not to be under a strong insular effect on life history traits. However, further investigation of sika deer populations from the Asian continent will shed light on the degree of insularity for the Japanese mainland sika deer.

We hypothesize a process of life history evolution on Japanese island deer as follows: the first step can be exemplified by sika deer on Kerama Islands. As observed, bone histological features showing signs of slow growth accompanied the smallest body size and slowest growth rate among understudied extant deer populations. These deer were introduced to Kerama Islands from the Kyushu mainland ca. 400 years ago by humans (Okinawa Prefectural Board of Education 1996). Fast changes in body size in less than 6,000 years have also been documented previously in red deer from Jersey [71]. Furthermore, domestic cattle that were introduced to Amsterdam Island experienced a rapid shrinking of body size in ca. 110 years [72, 73]. Though changes in growth trajectories might exist, these changes were not revealed in these interesting populations. With the Kerama deer, this drastic change in growth trajectory is proposed to have occurred within 400 years, which equates to ca. 80 generations (assuming a generation time of 5 years for the sika deer). The change in the growth trajectory of the Kerama deer is therefore proposed to have been brought about by phenotypic plasticity rather than a genetic modification in response to the less abundant understory vegetation in the subtropical evergreen broadleaved forest, accounting for the poor nutritional status of the Kerama deer [74], which was supported by the observation of a possible malnourished individual (URB-MAM-55; see SI for details). Lister [26] originally hypothesized the importance of phenotypic plasticity in the initial stage of insular dwarfism, and behind this would be epigenetic mechanisms controlling body growth, which respond to habitat environments [75, 76]. These mechanisms should be a challenging theme in the study of insular evolution. Another topic to be clarified in the Kerama deer would be the life history shift observed in its demographic aspect. It is therefore relevant to obtain reliable mortality data for the Kerama deer for this purpose.

The Yakushima Island deer illustrates the next step in life history evolution, which also showed a slower growth rate and intermediate histological features, coupled with

a moderate shift in survivorship curves toward those of slow life species. Contrary to the situation of the Kerama deer, the slower growth rate of Yakushima deer is proposed to have a genetic basis because; 1) Yakushima deer is not malnourished and have enjoyed a recent population increase and 2) some of their macromorphological features have a genetic basis [36, 38]. Long-term field observations of the Yakushima deer also supported our finding that life history traits of this population were shifting to slow life (Agetsuma and Agetsuma-Yanagihara, personal communication).

The clearest evolutionary transformation was shown in the two fossil deer species isolated from Okinawa Island over 1.5 Ma. Their growth patterns, bone histological characteristics, and mortality patterns were different from other deer populations. As with the Kerama deer, it was proposed that phenotypic plasticity partially influenced their growth trajectories. However, we do not currently have reliable sources of information on the nutritional status of the fossil deer, except for their dietary habits [61, 64]. Nevertheless, reports of healed bone fractures on several leg bones from a museum collection of fossil insular deer species on the Okinawa Island have been recorded, which implies that this deer survived long enough to recover from serious leg injuries. These findings refute the possibility that fossil Okinawa deer were malnourished and their body growth was suppressed due to a lack of resources. It was also notable that the same trend was observed for the fossil deer and muntjac, further supporting the shift toward slow life as an adaptation to the insular environment. This process is proposed to have started after their initial settlement on the Okinawa Island in part through phenotypic plasticity of growth trajectory due to a lack of sufficient foods during growing periods. A life history change would then have become genetically fixed through the natural selection of more slow life individuals, which occurred in the Yakushima Island deer. The outcome will have been the development of extremely slow life animals, despite their small body size. Unfortunately, this life history trait will have made these species vulnerable to human exploitation because animals having slow life histories have overall lower population recruitment than fast-life animals and their population recruitment is most declined by the increase of adult mortality rate [13, 70]. Consequently, the two Okinawa deer species became extinct at the time of or soon after the Paleolithic human arrival to the Okinawa Islands [77, 78], probably due to hunting. This extinction applies to other island dwarfs that became extinct during the Pleistocene era [79, 80, but see 81]. Recently, Rozzi et al. [82]

630 analyzed body size changes and extinction risk of over 1,500 island mammals

worldwide and clarified that insular dwarfs and giants are susceptible to extinction by

632 the arrival of modern humans. It is plausible that behind the extinction there could be

a life history shift toward the slow life in these insular extreme forms.

634   Our findings also agreed with worldwide observations of life history changes in

insular mammals previously documented (Figure 4). A living cervid (*Odocoileus*

636 *hemionus*) from Blakely Island showed slower growth rates than those from the

mainland population [10]. As the tooth eruption was delayed in the Blakely Island

638 deer, this life history change is proposed to have a genetic basis, similar to the case of

the Yakushima Island deer. With longer isolation but variable island sizes, the life

640 history response varied among mammals from Mediterranean islands. A fossil dwarf

hippo (*Hippopotamus minor*) from Cyprus showed no evident modification in bone

642 histology [19], though their life history traits remained unclarified. In contrast, a fossil

cervid (*Candiacervus*) from Crete showed slight modifications toward a slower life

644 history [18, 25]. The Sicilian pygmy elephant (*Elephas falconeri*) also showed a

modification in life history. This species was observed to have a fast life history as the

646 age distribution of its fossil assemblage was skewed to calves and immature

individuals [83]. Nevertheless, a histological study suggested that this pygmy elephant

648 had a much slower growth rate than its mainland relatives, and sexual maturity was

within the range of extant elephants [84], thus this contradiction requires a further

650 synthetic study on this species. The differences in the life history response among

Mediterranean mammals, however, are unsurprising, given that these Mediterranean

652 islands were much bigger than those for which we report life history changes (i.e., the

Blakely Island, Kerama Islands, Yakushima Island, Okinawa Island, and Mallorca

654 Island). The most explicit change in life history was observed in *Myotragus*

balearicus from the Mallorca Island, which underwent an exceptionally long time of

656 isolation (5.2 Ma) in predator-free environments. *Myotragus* showed a dramatic

decrease in bone growth rate and an evolution toward a slow life history [8, 21, 23],

658 implying the commonality of this evolutionary change with the Okinawa fossil deer.

These results propose that both island size and duration of isolation, with smaller

660 islands and longer isolation, strongly affected the degree of modification in life

history, thereby resulting in life history changes toward the slow life.

Conclusions

664 The extant Japanese sika deer can modify their life history traits through a combination of phenotypic plasticity and natural selection processes. This ability

666 underlies the process of insular dwarfism also recorded in fossil deer species (*C. astylodon* and the Ryukyu muntjac) from the Okinawa Island, which were estimated

668 to have a slow life history. After combining results from the present and previous studies on the life histories of insular mammals, we conclude that both island size and

670 duration of isolation strongly affect the degree of modification in life history traits, with longer isolation on smaller islands resulting in more prominent life history

672 changes. We also demonstrated that the evolution of a gradual and transitional shift toward the slow life history in large mammals from the mainland to insular

674 populations was the main life history change accompanying insular dwarfism, which makes the insular dwarfs vulnerable to human exploitation.

Data Availability Statement

678 All data are available in the main text and electronic supplementary information.

680 **Ethic statement**

 Ethical review and approval were not required for this study used museum

682 specimens and the investigations for the specimens were officially approved by the curators.

Author contributions

686 SH, MOK, and MF designed the research. SH, HT, and TN, prepared bone histological sections, took measurements and created visualizations. MOK, MI, and

688 TS collected ecological data. MOK conducted statistical analyses. SH, MOK, MSV, and MF drafted the original manuscript. All authors also contributed to reviews and

690 edits of the manuscript and approved the final manuscript.

692 **Funding**

 This work was supported by grants from the Sanyo Broadcasting Foundation (to SH

694 and MOK) and JSPS KAKENHI Grant Numbers JP16K18615 (to MOK), JP18H05254 (to TN) and JP19K04060 (to SH and MOK). MSV is supported by

696   Swiss SNF 31003A_169395. This work was partially supported JST-CREST Grant
        Number JPMJCR2194 (to TN).

Acknowledgments

700   We thank the following curators for access to museum collections under their care:
        Shimoinaba, S. (Natural History Museum and Institute, Chiba, Japan), Hayashi, T.

702   (Tochigi Prefectural Museum), Eda, M. (The Hokkaido University Museum), Usami,
        K. (Okinawa Prefectural Museum and Art Museum). Oshiro I. is also acknowledged

704   for his advice on the fossil Okinawa deer and support during this study. Furthermore,
        we thank Ochiai, K. and Shimoinaba, S. for providing ecological data on sika deer

706   and Reeves's muntjac in Honshu. Agetsuma, N. and Agetsuma-Yanagihara, Y. are
        appreciated for sharing data from their long-term field survey of the Yakushima deer.

708   We are grateful to Nakamura, K. and Nomura, H. as well for technical assistance in
        thin-sectioning, and Kodaira, S. and Kodama, R. for CT scanning. We appreciate

710   Kawada, S. (National Museum of Nature and Science, Tokyo), van der Geer, A.
        (Naturalis Biodiversity Center), Köhler, M. (ICP), Takakuwa, Y. (Gunma Museum of

712   Natural History), Kikuchi, H. and Umemura, Y. for providing photos of *Odocoileus
        hemionus, Hippopotamus minor, Candiacervus* spp.*, Myotragus balearicus,*

714   *Sinomegaceros yabei, Elephas falconeri,* and sika deer (Yakushima Island and
        Hokkaido mainland, respectively). We are grateful for the comments of Weisbecker,

716   V. and Köhler, M. on the earlier version of this manuscript.

Conflict of interest

718
        We have no competing interests to declare.

References

722   1.   Foster JB: **Evolution of Mammals on Islands.** *Nature* 1964, **202:**234-235.
        2.   van der Geer A, Lyras G, de Vos J, Dermitzakis M: *Evolution of Island
724   Mammals: Adaptation and Extinction of Placental Mammals on Island.* West
              Sussex: John Wiley & Sons Ltd.; 2010.
726   3.   Benitez-Lopez A, Santini L, Gallego-Zamorano J, Mila B, Walkden P,
              Huijbregts MAJ, Tobias JA: **The island rule explains consistent patterns of
728   body size evolution in terrestrial vertebrates.** *Nat Ecol Evol* 2021, **5:**768-
              786.
730   4.   Larramendi A, Palombo MR: **Body Size, biology and encephalization
              quotient of *Palaeoloxodon* ex gr. *P. falconeri* from Spinagallo Cave**

732   **(Hyblean plateau, Sicily).** *Hystrix-Italian Journal of Mammalogy* 2015, **26:**102-109.

734   5.   Lomolino MV, van der Geer AA, Lyras GA, Palombo MR, Sax DF, Rozzi R: **Of mice and mammoths: generality and antiquity of the island rule.** *J*
736   *Biogeogr* 2013, **40:**1427-1439.

  6.   Lomolino MV, Sax DF, Palombo MR, van der Geer AA: **Of mice and**
738   **mammoths: evaluations of causal explanations for body size evolution in insular mammals.** *J Biogeogr* 2012, **39:**842-854.

740   7.   Sander PM, Mateus O, Laven T, Knotschke N: **Bone histology indicates insular dwarfism in a new Late Jurassic sauropod dinosaur.** *Nature* 2006,
742   **441:**739-741.

  8.   Köhler M, Moyà-Solà S: **Physiological and life history strategies of a fossil**
744   **large mammal in a resource-limited environment.** *Proc Natl Acad Sci U S A* 2009, **106:**20354-20358.

746   9.   Marín-Moratalla N, Jordana X, García-Martínez R, Köhler M: **Tracing the evolution of fitness components in fossil bovids under different selective**
748   **regimes.** *Comptes Rendus Palevol* 2011, **10:**469-478.

  10.   Long ES, Courtney KL, Lippert JC, Wall-Scheffler CM: **Reduced body size**
of insular black-tailed deer is caused by slowed development. *Oecologia* 2019, **189:**675-685.

11.   Palkovacs EP: **Explaining adaptive shifts in body size on islands: a life history approach.** *Oikos* 2003, **103:**37-44.

12.   Begon M, Harper JL, Townsend CR: *Ecology: From Individuals to Ecosystems, 4*th *Edition.* Oxford: Blackwell Publishing Ltd.; 2006.

13.   Oli MK: **The fast–slow continuum and mammalian life-history patterns: an empirical evaluation.** *Basic Appl Ecol* 2004, **5:**449-463.

14.   Erickson GM: **On Dinosaur Growth.** *Annual Review of Earth and Planetary Sciences* 2014, **42:**675-697.

15.   Erickson GM: **Growth patterns in brooding dinosaurs reveals the timing of sexual maturity in non-avian dinosaurs and genesis of the avian**
condition. *Biol Lett* 2007, **3:**558-561.

  16.   Castanet J, Rogers KC, Cubo J, Boisard J-J: **Periosteal bone growth rates in**
764   **extant ratites (ostriche and emu). Implications for assessing growth in dinosaurs.** *CR Acad Sci Paris, Sciences de la vie / Life Sciences* 2000,
766   **323:**543-550.

  17.   Amson E, Kolb C, Scheyer TM, Sánchez-Villagra MR: **Growth and life**
history of Middle Miocene deer (Mammalia, Cervidae) based on bone histology. *Comptes Rendus Palevol* 2015, **14:**637-645.

18.   Kolb C, Scheyer TM, Lister AM, Azorit C, de Vos J, Schlingemann MA, Rossner GE, Monaghan NT, Sanchez-Villagra MR: **Growth in fossil and**
extant deer and implications for body size and life history evolution. *BMC Evol Biol* 2015, **15:**19.

19.   Kolb C, Scheyer TM, Veitschegger K, Forasiepi AM, Amson E, Van der Geer AA, Van den Hoek Ostende LW, Hayashi S, Sanchez-Villagra MR:
Mammalian bone palaeohistology: a survey and new data with emphasis on island forms. *PeerJ* 2015, **3:**e1358.

20.   Köhler M, Marín-Moratalla N, Jordana X, Aanes R: **Seasonal bone growth and physiology in endotherms shed light on dinosaur physiology.** *Nature*
2012, **487:**358-361.

21.   Marín-Moratalla N, Jordana X, Köhler M: **Bone histology as an approach to providing data on certain key life history traits in mammals: implications for conservation biology.** *Mammalian Biology* 2013, **78:**422-429.

22.   Klevezal GA: *Recording Structures of Mammals. Determination of Age and Reconstruction of Life History.* Rotterdam: A. A. Balkema; 1996.

23.   Jordana X, Marín-Moratalla N, DeMiguel D, Kaiser TM, Köhler M: **Evidence of correlated evolution of hypsodonty and exceptional longevity in endemic insular mammals.** *Proc Biol Sci* 2012, **279:**3339-3346.

24.   Jordana X, Köhler M: **Enamel microstructure in the fossil bovid Myotragus balearicus (Majorca, Spain): Implications for life-history evolution of dwarf mammals in insular ecosystems.** *Palaeogeogr Palaeoclimatol Palaeoecol* 2011, **300:**59-66.

25.   Miszkiewicz JJ, Van Der Geer AAE: **Inferring longevity from advanced rib remodelling in insular dwarf deer.** *Biological Journal of the Linnean Society* 2022, **136:**41-58.

26.   Lister AM: **Dwarfing in island elephants and deer: processes in relation to time of isolation.** *Symp Zool Soc* 1996, **69:**277-292.

27.   Raia P, Meiri S: **The island rule in large mammals: paleontology meets ecology.** *Evolution* 2006, **60:**1731-1742.

28.   van der Geer AA, Lyras GA, Lomolino MV, Palombo MR, Sax DF: **Body size evolution of palaeo-insular mammals: temporal variations and interspecific interactions.** *J Biogeogr* 2013, **40:**1440-1450.

29.   Otsuka H, Takahashi A: **Pleistocene vertebrate faunas in the Ryukyu Islands: their migration and extinction.** *Tropics* 2000, **10:**25-40.

30.   Taruno H, Kawamura Y, Ishida S, Okumura K: **Yabe's giant deer and elk remains from the Late Pleistocene of Kumaishi-do Cave, Gifu Prefecture, central Japan (Part 2): Postcranial bones.** *Bulletin of the Osaka Museum of Natural History* 2017, **71:**17-142.

31.   de Margerie E, J. Cubo, Castanet J: **Bone typology and growth rate: testing and quantifying 'Amprino's rule' in the mallard (*Anas platyrhynchos*).** *Comptes Rendus Biologies* 2002, **325:**221-230.

32.   Groves C: **The genus Cervus in eastern Eurasia.** *European Journal of Wildlife Research* 2006, **52:**14-22.

33.   Ohtaishi N: **Preliminary memorandum of classification, distribution and geographic variation on Sika deer.** *Honyurui Kagaku (Mammalian Science)* 1986, **53:**13-17 (in Japanese with English abstract).

34.   Kubo MO, Takatsuki S: **Geographical body size clines in sika deer: path analysis to discern amongst environmental influences.** *Evol Biol* 2015, **42:**115-127.

35.   Chiba Prefecture, Deer Research Group on Boso: *Science report on emergent survey of invasive alien species (Reeves's muntjac) in Boso.* Chiba: Chiba Prefecture; 2007.

36.   Terada C, Tatsuzawa S, Saitoh T: **Ecological correlates and determinants in the geographical variation of deer morphology.** *Oecologia* 2012, **169:**981-994.

37.   Ohshima K: **The history of straits around the Japanese Islands in the Late-Quaternary.** *The Quaternary Research* 1990, **29:**193-208 (in Japanese with English abstract).

38. Terada C, Saitoh T: **Phenotypic and genetic divergence among island populations of sika deer (*Cervus nippon*) in southern Japan: a test of the local adaptation hypothesis.** *Population Ecology* 2018, **60:**211-221.

39. Nagata J, Masuda R, Tamate HB, Hamasaki S, Ochiai K, Asada M, Tatsuzawa S, Suda K, Tado H, Yoshida MC: **Two genetically distinct lineages of the sika deer, *Cervus nippon*, in Japanese islands: comparison of mitochondrial D-loop region sequences.** *Mol Phylogenet Evol* 1999, **13:**511-519.

40. Okinawa Prefectural Board of Education: *A report of conservation and management of sika deer in Kerama Islands.* Okinawa: Okinawa Prefectural Board of Education; 1996.

41. Okinawa Prefectural Museum and Art Museum (Ed.). **Excavation of the Hananda-Gama Cave site, Okinawa**. Naha: Kokusai-insatsu Co.; 2009.

42. Azanza B, DeMiguel D, DeMiguel D, Andrés M, Andrés M: **The antler-like appendages of the primitive deer *Dicrocerus elegans*: morphology, growth cycle, ontogeny, and sexual dimorphism.** *Estudios Geológicos* 2011, **67:**579-602.

43. Okumura K, Ishida S, Taruno H, Kawamura Y: **Yabe's giant deer and elk remains from the Late Pleistocene of Kumaishi-do Cave, Gifu Prefecture (Part 1): Antlers, a skull, mandibles, and teeth.** *Bulletin of the Osaka Museum of Natural History* 2016, **70:**1-82 (in Japanese with English abstract).

44. Okumura K, Ishida S, Kawamura Y, Kumada M, Tamiya S: **Latest Pleistocene Mammalian assemblage of Kumaishi-do Cave, Gifu Prefecture and the significance of its ^{14}C age.** *Earth Science (Chikyu Kagaku)* 1982, **36:**214-218 (in Japanese with English abstract).

45. Chinsamy A, Raath MA: **Preparation of fossil bone for histological examination.** *Palaeontologia Africana* 1992, **29:**39-44.

46. Sander PM: **Longbone histology of the Tendaguru sauropods: implications for growth and biology.** *Paleobiology* 2000, **26:**466-488.

47. Castanet J, Croci S, Aujard F, Perret M, Cubo J, de Margerie E: **Lines of arrested growth in bone and age estimation in a small primate: *Microcebus murinus*.** *J Zool* 2004, **263:**31-39.

48. Jordana X, Marín-Moratalla N, Moncunill-Solè B, Nacarino-Meneses C, Köhler M: **Ontogenetic changes in the histological features of zonal bone tissue of ruminants: A quantitative approach.** *Comptes Rendus Palevol* 2016, **15:**255-266.

49. Calderon T, DeMiguel D, Arnold W, Stalder G, Köhler M: **Calibration of life history traits with epiphyseal closure, dental eruption and bone histology in captive and wild red deer.** *J Anat* 2019, **235:**205-216.

868 50. Lehman TM, Woodward HN: **Modeling growth rates for sauropod dinosaurs.** *Paleobiology* 2008, **34:**264-281.

870 51. Francillon-Vieillot H, Buffrénil Vd, Castanet J, Géraudie J, Meunier FJ, Sire JY, Zylberberg L, Ricqlès Ad: **Microstructure and Mineralization of Vertebrate Skeletal Tissues.** In *Skeletal Biomineralization: Patterns, Processes and Evolutionary Trends.* Edited by Carter JG. New York: Van Nostrand Reinhold; 1989: 471-530

52. Castanet J, Francillon-Vieillot H, Meunier FJ, de Ricqlès A: **Bone and individual aging.** In *Bone Vol 7: Bone Growth.* Edited by Hall BK. Boca Raton: CRC Press; 1993: 245–283

878   53.   Orlandi-Oliveras G, Nacarino-Meneses C, Koufos GD, Köhler M: **Bone histology provides insights into the life history mechanisms underlying**
880   **dwarfing in hipparionins.** *Sci Rep* 2018, **8:**17203.

54.   Lee AH, Huttenlocker AK, Padian K, Woodward HN: **Analysis of growth**
882   **rates.** In *Bone Histology of Fossil Tetrapods: Advancing Methods, Analysis, and Interpretation.* Edited by Padian K, Lamm ET. Berkeley: University of
884   California Press; 2013: 217-251

55.   Woodward HN, Padian K, Lee AH: **Skeletochronology.** In *Bone Histology of*
886   *Fossil Tetrapods: Advancing Methods, Analysis, and Interpretation.* Edited by Padian K, Lamm ET. Berkeley: University of California Press; 2013: 195 –
888   216

56.   Andrade RCLP, Bantim RA, Lima FJ, dos Santos Campos L, Eleutério LHS,
890   Sayão J: **New data about the presence and absence of the external fundamental system in archosaurs.** *Cadernos de Cultura e Ciência* 2015,
892   **14:**200-211.

57.   Monteiro LS, Falconer DS: **Compensatory growth and sexual maturity in**
894   **mice.** *Animal Science* 1966, **8:**179-192.

58.   Hokkaido Institute of Environmental Sciences: **Results of a survey related to**
sika deer and brown bear sighting on Hokkaido. Sapporo: Hokkaido Institute of Environmental Sciences (in Japanese); 1997.

59.   Ozaki M, Kaji K, Matsuda N, Ochiai K, Asada M, Ohba T, Hosoi E, Tado H, Koizumi T, Suwa G, Takatsuki S: **The relationship between food habits,**
molar wear and life expectancy in wild sika deer populations. *J Zool* 2010, **280:**202-212.

60.   Kubo MO, Fujita M, Matsu'ura S, Kondo M, Suwa G: **Mortality profiles of late Pleistocene deer remains of Okinawa Island: evidence from the**
Hananda-Gama cave and Yamashita-cho cave I sites. *Anthropological Science* 2011, **119:**183-201.

61.   Kubo MO, Yamada E, Fujita M, Oshiro I: **Paleoecological reconstruction of Late Pleistocene deer from the Ryukyu Islands, Japan: combined**
evidence of mesowear and stable isotope analyses. *Palaeogeogr Palaeoclimatol Palaeoecol* 2015, **435:**159-166.

62.   Chapman NG, Brown WAB, Rothery P: **Assessing the age of Reeves' muntjac (*Muntiacus reevesi*) by scoring wear of the mandibular molars.** *J*
Zool 2005, **267**.

63.   Ozaki M: **Estimation of age at death from tooth wear of lower molars of**
914   **excavated deer and comparison of age structures with modern deer populations.** In *Excavation of the Hananda-Gama Cave site, Okinawa.* Edited
916   by Okinawa Prefectural Museum and Art Museum. Naha: Kokusai-insatsu Co.; 2009: 24-34 (in Japanese)

918   64.   Kubo MO, Fujita M: **Diets of Pleistocene insular dwarf deer revealed by dental microwear texture analysis.** *Palaeogeogr Palaeoclimatol Palaeoecol*
920   2021, **562:**110098.

65.   Caughley G: *Analysis of Vertebrate Populations.* New York: John Wiley &
922   Sons Inc.; 1977.

66.   Ochiai K, Asada M: **Growth in the body size of sika deer (*Cervus nippon*)**
924   **on the Boso Peninsula, central Japan.** *Journal of the Natural History Museum and Institute, Chiba* 1995, **3:**223-232 (in Japanese with English
926   abstract).

928 67. Suzuki M, Onuma M, Yokoyama M, Kaji K, Yamanaka M, Ohtaishi N: **Body size, sexual dimorphism, and seasonal mass fluctuations in a larger sika deer subspecies, the Hokkaido sika deer (*Cervus nippon yesoensis* Heude,**
930 **1884).** *Can J Zool* 2001, **79:**154-159.

68. Sand H, Cederlund G, Danell K: **Geographical and latitudinal variation in**
932 **growth patterns and adult body size of Swedish moose (*Alces alces*).** *Oecologia* 1995, **102:**433-442.

934 69. Woodward HN, Horner JR, Farlow JO: **Quantification of intraskeletal histovariability in *Alligator mississippiensis* and implications for**
936 **vertebrate osteohistology.** *PeerJ* 2014, **2:**e422.

70. Pianka ER: *Evolutionary Ecology.* New York: Harper & Row; 1978.
938 71. Lister AM: **Rapid dwarfing of red deer on Jersey in the Last Interglacial.** *Nature* 1989, **342:**539-542.

940 72. Rozzi R, Lomolino MV: **Rapid Dwarfing of an Insular Mammal - The Feral Cattle of Amsterdam Island.** *Sci Rep* 2017, **7:**8820.

942 73. Berteaux D, Micol T: **Population studies and reproduction of the feral cattle (*Bos taurus*) of Amsterdam Island, Indian Ocean.** *J Zool* 1992,
944 **228:**265–276.

74. Shiroma T, Ohta H: **Geographical variation in cranial morphology in sika**
deer with a special reference on Kerama deer In *A report of conservation and management of sika deer in Kerama Islands.* Edited by Okinawa
Prefectural Board of Education. Okinawa: Okinawa Prefectural Board of Education; 1996: 13-55 (in Japanese)

75. Price TD, Qvarnstrom A, Irwin DE: **The role of phenotypic plasticity in driving genetic evolution.** *Proc Biol Sci* 2003, **270:**1433-1440.

76. Lafuente E, Beldade P: **Genomics of Developmental Plasticity in Animals.** *Front Genet* 2019, **10:**720.

77. Fujita M, Yamasaki S, Katagiri C, Oshiro I, Sano K, Kurozumi T, Sugawara H, Kunikita D, Matsuzaki H, Kano A, et al: **Advanced maritime adaptation**
in the western Pacific coastal region extends back to 35,000-30,000 years before present. *Proc Natl Acad Sci U S A* 2016, **113:**11184-11189.

78. Fujita M, Yamasaki S, Sugawara H, Eda M: **Body size reduction in wild boar (*Sus scrofa*) from the late Pleistocene Maehira Fissure Site in**
Okinawa-jima Island, Japan, with relevance to human arrival. *Quaternary International* 2014, **339:**289-299.

79. Sondaar PY: **Early human exploration and exploitation of islands.** *Tropics* 2000, **10:**203-230.

80. Alcover JA, Sans A, Palmer M: **The Extent of Extinctions of Mammals on Islands.** *J Biogeogr* 1998, **25:**913-918.

81. Louys J, Braje TJ, Chang CH, Cosgrove R, Fitzpatrick SM, Fujita M, Hawkins S, Ingicco T, Kawamura A, MacPhee RDE, et al: **No evidence for**
widespread island extinctions after Pleistocene hominin arrival. *Proc Natl Acad Sci U S A* 2021, **118**.

82. Rozzi R, Lomolino MV, van der Geer A, Silvestro D, Lyons SK, Bover P, Alcover JA, Benítez-López A, Tsai C-H, Fujita M, et al: **Dwarfism and**
gigantism drive human-mediated extinctions on islands. *Science* 2023, **in press**.

83. Raia P, Barbera C, Conte M: **The fast life of a dwarfed giant.** *Evol Ecol* 2003, **17:**293-312.

84. Köhler M, Herridge V, Nacarino-Meneses C, Fortuny J, Moncunill-Solè B, Rosso A, Sanfilippo R, Palombo MR, Moyà-Solà S: **Palaeohistology reveals**
a slow pace of life for the dwarfed Sicilian elephant. *Sci Rep* 2021, **11:**22862.
85. Palombo MR: **Endemic elephants of the Mediterranean Islands: knowledge, problems and perspectives.** In *The World of Elephants -*
International Congress, Rome 2001. pp. 486-491. Rome; 2001:486-491.
86. Kawamura Y: **Fossil record of sika deer in Japan.** In *Sika Deer: Biology and*
984 *Management of Native and Introduced Populations.* Edited by McCullough DR, Takatsuki S, Kaji K. Tokyo: Springer; 2009: 11-25

988 **Figures and Tables**

FIGURE 1

992 **Locations of the sampled deer populations in Japan.** Dagger (†) indicates fossils

excavated from Late Pleistocene deposits. Localities of extant taxa are shown in

994 closed stars, whereas those of extinct taxa are in open stars. The approximate body

height (BH) was also indicated for each species/population. Photography of fossil

996   cervids was courtesy of the Gunma Museum of Natural History (*Sinomegaceros yabei*) and Okinawa Prefectural Museum & Art Museum (*Cervus astylodon* and

998   Muntiacini gen. et sp. indet.).

1000   **FIGURE 2**

Growth (A) and survivorship curves (B-C) for studied Japanese deer. The insular

1002   deer have a significantly slower growth rate and a delayed growth plateau (somatic

maturity) compared with the mainland deer (A), and an ecological shift exists in the

1004   mortality pattern between the insular and mainland deer (B, C). As the age estimation of

C. astylodon was based on the lower third molar height, we confined the survivorship

1006   curves to juvenile and onward for *C. astylodon* and the sika deer (B).

FIGURE 3

Bone histological features of studied Japanese deer. (A–C) Midshaft cross-section (A),
the cortex under normal light (B), and the cortex under polarized light (C) of a femur of
fossil *Cervus astylodon* (OPM-HAN07-1603). (D–F) Midshaft cross sections (D), the
cortex under normal light (E), and the cortex under polarized light (F) of a femur of
extant sika deer (*Cervus nippon*) from Yakushima Island (TPM-M-313). (G–I) Midshaft
cross section (G), the cortex under normal light (H), and the cortex under polarized light
(I) of a femur of extant sika deer from the Honshu mainland (CBM-ZZ-412). (J–L)
Midshaft cross section (J), the cortex under normal light (K), and the cortex under
polarized light (L) of a femur of fossil *Sinomegaceros yabei* (OMNH-QV-4062). Red
lines in A, D, G, and J, and yellow arrows in B, C, E, F, H, I, and K indicate lines of
arrested growth. The outer cortex is at the upper right of B, C, E, and F and on the right in

1020   H, I, and K. BM, body mass; EFS, external fundamental system; FBL, fibro-lamellar

bone tissue; LAG, line of arrested growth; LV, longitudinal vascular canal; MC,

1022   medullary cavity; PFB, parallel-fibered bone tissue; PV, plexiform to laminar vascular

canal. [†] Fossil cervids. Note: only fossil deer from Okinawa Islands had a distinctly

1024   longer-term geographical isolation history and exhibited bone tissue, indicating a slower

growth rate [31].

FIGURE 4

The Diagram summarizes evolutionary processes, the relationship between insularity, body size, and life history observed in extant and fossil large mammals. Cervid species/populations of the Japanese Archipelago were investigated in this study and information on animals from other islands we obtained from references: *Odocoileus hemionus* [10], *Hippopotamus minor* [2], *Candiacervus* spp. [2],

1031   *Elephas falconeri* [2, 85], *Myotragus balearicus* [8]. Photographs of fossil mammals in Mediterranean islands courtesy of Gunma Museum

1032   of Natural History (*Elephas falconeri*), Alexandra van der Geer (*Hippopotamus minor* and *Candiacervus* spp.), and Meike Köhler

1033   (*Myotragus balearicus*).

1034   [†]Fossil taxa. [*] Life history of *Elephas falconeri* was disputed [83, 84]. [**] *Muntiacus reevesi* was an artificially introduced animal in the

1035   1980s. Therefore, the isolation period of >0.43 Myr is inapplicable to it [35].

1036

1037 **TABLE 1 Summary of the general information of sampled Japanese deer populations.**

Species/Population	Locality (island size)	Timing of immigration/isolation [reference]	Habitat type[*]	EBM (kg)[‡]	N of individuals used in histological analyses	N of demographic analysis samples [reference of data]
Extant species/population						
Cervus nippon (Hokkaido mainland)	Hokkaido (83,424 km^2)	Separation from Honshu mainland population: Late Pleistocene (ca. 0.02 Ma) [86]	Subarctic coniferous forests or mixed forests	105.2	7	1060 [58]
Cervus nippon (Honshu mainland)	Honshu (227,962 km^2)	Middle Pleistocene (0.43 Ma) [86]	Temperate evergreen broadleaved forests	45.5	8	594 [60]
Muntiacus reevesi		Recent introduction (1960–1980) [35]		9.2	5	84 [62]
Cervus nippon (Yakushima Island)	Yakushima Island (505 km^2)	Isolation from Kyushu mainland: Late Pleistocene (ca. 0.1–0.02 Ma; see SI Appendix for details and references therein)	Subtropical evergreen broadleaved forests	37.2	4	74 [this study]
Cervus nippon (Kerama Islands)	Kerama Islands (36 km^2)	Historic introduction (ca. 400 years ago) [74]	Subtropical evergreen broadleaved forests	27.1	7	NA
Fossil species						

Species/Population	Locality (island size)	Timing of immigration/isolation [reference]	Habitat type[*]	EBM (kg) [‡]	N of individuals used in histological analyses	N of demographic analysis samples [reference of data]
[†]*Sinomegaceros yabei*	Honshu (227,962 km^2)	Middle Pleistocene (>0.43 Ma) [30]	NA	564.5	4 (Femur: 2, Tibia: 2)	NA
[†]*Cervus astylodon*	Okinawa Island (1,207 km^2)	Early Pleistocene (ca. 1.5 Ma) [29]	NA	25.4	11 (Femur: 6, Tibia: 5)	45 [60]
[†]Muntiacini gen. et sp. indet.				16.5	6 (Femur: 4, Tibia: 2)	65 [63]

1038   [*] Habitat was obtained from a reference [34]. [†] Fossil cervids. [‡] The estimated body mass (EBM) was obtained from fitted growth curves (SI

1039   Figures S8, S9) by averaging asymptotes for each species/population (SI Table S8).

1040