Variation and process of life history evolution in insular dwarfism as revealed by a

natural experiment

Author affiliation:

Shoji Hayashi[1,2,*,**], Mugino O. Kubo[3,*,**], Marcelo R. Sánchez-Villagra[4], Hiroyuki

Taruno[5], Masako Izawa[6,7], Tsunehiro Shiroma[6], Takayoshi Nakano[2], Masaki Fujita[8]

[1] Faculty of Biosphere-Geosphere Science, Okayama University of Science, Kita-ku

Ridaicho 1-1, Okayama, 700-0005, Japan; [2] Division of Materials and Manufacturing

Science, Graduate School of Engineering, Osaka University, Yamada-Oka, Suita, Osaka,

565-0871, Japan; [3] Department of Natural Environmental Studies, Graduate School of

Frontier Sciences, The University of Tokyo, 5-1-5 Kashiwanoha, Kashiwa, Chiba 277-

8563, Japan; [4] Paläontologisches Institut und Museum der Universität Zürich, Karl

Schmid-Strasse 4, CH-8006 Zürich, Switzerland; [5] Osaka Museum of Natural History,

Nagai Park 1-23, Higashi-Sumiyoshi-ku, Osaka, 546-0034, Japan; [6] Faculty of Science,

University of the Ryukyus, Okinawa 903-0213, Japan; [7] Kitakyushu Museum of Natural

History & Human History, 2-4-1 Higashida, Yahatahigashi-ku, Kitakyushu 805-0071,

Japan; [8] Department of Anthropology, National Museum of Nature and Science, 4-1-1

Amakubo, Tsukuba-shi, Ibaraki 305-0005, Japan

[*] Both authors contributed equally to this work.

[**] *Authors for correspondence*: Shoji Hayashi e-mail: hayashi@big.ous.ac.jp, Mugino O.

Kubo e-mail: mugino@k.u-tokyo.ac.jp

Keywords: bone histology, deer, demography, growth rate, islands, Japan, *K*-strategy,

life history, skeletochronology

Abstract

Background

Islands offer a classic topic for evolutionary studies. Few other subjects have

historically raised as much fascination as the island large mammals that having evolved

into dwarfs. Consensus has been gained that multivariate ecological causes are behind

those changes, but what remains largely unexplored are the processes involved. Recent

studies focused on associated change of body size and life history (age of reproduction,

growth rate, longevity) – a key to understand the process of insular dwarfism. The

Japanese Archipelago offers worldwide a unique natural experiment, as in its numerous

islands of different sizes the same group of organisms, deer, have evolved into different

sizes. We investigated eight deer populations exhibiting body size variation, both extant

and fossil, to clarify the effect of insularity on life history traits.

Results

We applied several methods to both extant and extinct populations for resolving life

history change among deer populations. Skeletochronology using lines of arrested

growth successfully reconstructed body growth curves and revealed a gradual change in

the growth trajectories reflecting degree of insularity: slower growth with prolonged

growth periods in more isolated deer populations. An extensive examination of bone

microstructure further corroborated it, and clearly indicated much slower growth and

later somatic maturity in fossil insular deer, which had been isolated for more than 1.5

Myr. Finally, mortality patterns revealed by demographic analysis varied among deer

populations and life history of insular populations shifted toward "slow-life" of *K*-

strategists.

Conclusion

We clarified the evolutionary and ecological process behind insular dwarfism, which occurred in conjunction with life history shifts. The process initiated with phenotypic plasticity responding to the resource-limitation on predator-free islands after the settlements of islanders. Under the insular environments, natural selection favored K-selected animals and the life history traits were genetically fixed. Extreme K-strategy found in fossil dwarfs on islands would make them vulnerable to anthropogenic changes, that would explain termination of insular dwarfs after human arrival on islands.

Background

Among the most iconic examples of visible evolutionary change are island mammals
that dramatically changed in body size [1-3]. Elephants, hippos and deer in islands
across the world became very small, whereas small mammals became larger [2], what
led to the suggestion of an 'island rule' [1]. A classic example is the pygmy elephant
(*Elephas falconeri*) of Sicily, which had a body mass of only 1% of its mainland
ancestor [4].

Body size has a close relationship with various aspects of organism's life. Recent
studies are focusing on the relationship between body size and life history – growth rate,
age of reproduction, longevity – on islands [5-7]. The reduction of body size in large
mammals appears to be associated with modification of the growth trajectory [8, 9], and
this in turn affects the life schedule, e.g. timing of sexual maturity or longevity. Life
history theory, which has been extended from r/K selection theory in ecology [10],
predicts that species adapted to unstable environments, under which high intrinsic rates
of natural increase (r) are adaptive, suit a set of life history traits such as rapid body
growth, early sexual maturity, large number of offspring for a clutch and a short life
span, being referred as "r-selected". On the other hand, in stable environments the
populations are maintained near their carrying capacity (K) and resultant intraspecific
competitions favor those which have slow body growth, late sexual maturity, small
number of offspring for a clutch and a long life span, referred as "K-selected". Based on
this theoretical background, life history traits were estimated for fossil insular species.
Some dwarfed species show a distinctly slower growth rate and delayed somatic
maturity compared with continental taxa [6], whereas others do not show extreme
modifications in their life histories [11, 12]. Since these studies were based on different

species from single islands, it is hard to generalize body size and life history evolution

on islands.

Further important issue awaiting to be explored is the process of body size and life

history evolution, and how ecological factors affect the process, e.g. a lack of predators

and competitors, resource limitation on islands, founder effects of island immigrants,

and the remoteness and duration of isolation [2, 13-16]. Drastic body size changes have

been reported for fossil species or populations [2]. Nevertheless, as far as we focused

only on the fossil evidence, the ecological process of the changes remains unclear. To

achieve this, studies of mainland and insular populations of a single extant species are

particularly useful because they allow confounding factors other than living on islands

to be controlled. Variation in isolation durations among the populations can mimic the

process of insular settlement.

In the present study, we focused on deer in the Japanese Archipelago, both extant and

extinct populations from both islands and mainland (Fig. 1). Our targets cover an extant

sika deer (*Cervus nippon*) population recently introduced (c.a. 400 years ago) on a small

island, four extant insular and mainland deer populations [sika deer, Reeves's muntjac

(*Muntiacus reevesi*)] with variable body sizes, and extinct insular deer (*Cervus*

astylodon and Muntiacini gen. et. sp. indet., hereafter referred to as "the Ryukyu

muntjac") isolated for more than 1.5 Myr on the Okinawa Island [17], representing

different degrees of insular effect. We also included an extinct gigantic mainland deer

(*Sinomegaceros yabei*) from the Late Pleistocene deposits on Honshu mainland, having

a comparable body size to the largest extant deer, the moose (*Alces alces*) [18]. The

immense variation of the island size and duration of isolation found in the eight deer

populations enables us to infer the process of body size and life history changes on

islands.

We applied long bone histological analyses established for reconstruction of

palaeobiology of fossil species [6, 7, 11, 12, 19-21], which could be applicable to both

extant and extinct taxa to clarify variation of body growth rates and life histories of the

deer in Japan. Skeletochronology using lines of arrested growth (LAGs) was applied to

depict body growth rates, which was further corroborated by detailed qualitative

descriptions of bone histology aiming to infer growth pattern and somatic maturity. We

also conducted demographic analyses of age data, which were collected either

ecological surveys or analysis of tooth wear of fossil deer assemblages, to quantitatively

compare mortality patterns among the deer populations. By making the most of these

interdisciplinary analytical methods, we expected a gradual change in life history traits

in the deer populations according to the intensity of insular effects, with longer isolation

on smaller islands resulting in the shift toward *K*-strategy.

Results

Growth curve models and growth rate comparisons using skeletochronology

The lines of arrested growth (LAGs) in the long bones of both the extant and fossil deer

were well preserved (SI Appendix, Figure S1 – 4), allowing us to measure the bone

diameters at the LAGs. We estimated ontogenetic change in body mass and depicted a

growth curve for each individual (SI Appendix, Figure S8, 9, Table S8). We then

averaged each of three growth curve parameters (the asymptote, growth rate, and

inflection point) generated by the Gompertz curve fitting for each species/population to

deduce a representative growth curve for each (Fig. 2A, SI Appendix, Figure S10).

Among the sika deer populations, deer from the two mainland areas (Honshu and

Hokkaido mainland) showed faster growth than those from the two insular populations

(Yakushima and Kerama Islands). The growth trajectories of body mass corresponded

to those obtained by investigations of culled individuals [22, 23] and by long-term field

observations of wild deer (Agetsuma and Agetsuma-Yanagihara, personal

communication), validating our estimation method of body growth from

skeletochronology. The extant Reeves's muntjac showed the fastest growth, which was

also reported in analysis of culled individuals, corresponded with its early sexual

maturity at ca. 6 months [24].

Compared to the extant references, the two fossil insular deer (*C. astylodon* and the

Ryukyu muntjac) showed remarkably slow and prolonged growth, whereas the

Pleistocene giant deer *S. yabei* had fast growth, suggesting a similar growth pattern to

other extant large deer, such as moose [25]. Statistical comparisons of the growth

parameters revealed that both the fossil and extant insular deer had significantly slower

growth rates than Reeves's muntjac and *S. yabei* (SI Appendix, Tables S10, S11).

However, the growth rates of the four extant sika deer populations and the fossil insular

deer were not significantly different from each other (SI Appendix, Tables S10, S11),

due to the transitional position of the extant insular deer, although the limited numbers

of possible samples may also play a role. The inflection point, which marks the

transition from the initial phase of exponential growth to the subsequent phase of

asymptotic growth, is known to be associated with the timing of sexual maturity [26]

and was greatest in *C. astylodon* (SI Appendix, Tables S10, S12), implying that sexual

maturity may be delayed in this species. However, the differences were not always

statistically significant, reflecting high levels of variation within species/populations.

Variation in bone tissue among deer populations in Japan

The nomenclature and definitions of the bone microstructures in the following

descriptions are based on Francillon-Vieillot et al. [27] and Castanet et al. [28].

Histological characteristics of long bones corroborated the above statistical

comparisons of growth curves. We focused on the following bone tissue characters: 1)

type of primary periosteal bone tissue and vascularization pattern as an indicator of

relative bone growth rate [e.g. 29] and 2) the presence of the external fundamental

system (EFS), which is constituted by parallel-fibred bone tissue without

vascularization and forming an edge of slow-growing bone tissue, as a sign of somatic

maturity [e.g. 30, 31] . Fig. 3 summarizes the main histological features of the cervids

we examined (also see SI Appendix, Tables S1–S3).

All of the extant deer populations (i.e., the four extant sika deer populations, the extant

Reeves's muntjac) and the gigantic fossil *S. yabei* exhibited a similar bone tissue

structure to continental cervids [11, 32, 33], consisting of fast-growing periosteal bone

tissue known as a fibro-lamellar bone with a high level of laminar to plexiform

vascularization in most of the cortex (Fig. 3G–L; SI Appendix, Figures S2A–I, S3G–I,

and S4A–O). Slow-growing parallel-fibered bone tissue was observed in the outer

cortex. Notably, the parallel-fibered bone tissue was more extensively developed in the

two insular populations (sika deer from Yakushima and Kerama Islands) with the LAGs

at the inner to middle cortex being tighter than in the mainland deer (Fig. 3D–F, SI

Appendix, Figures S2D–F, and S3G-I, J-L), suggesting a slower growth rate among the

insular populations. The cortical bone tissues of two fossil insular deer, *C. astylodon*

and the Ryukyu muntjac, were different from those of the extant deer and the gigantic

fossil *S. yabei*, in that they were primary parallel-fibered bone tissue with multiple

LAGs, implying much slower growth rate than other deer. Their bone tissues were

similar to those of the extinct dwarf bovid *Myotragus* [6] and of typical extant reptiles

[e.g. 34]. The LAGs were spaced evenly throughout the cortex (Fig. 3A–C; SI

Appendix, Figures S1A–C and S3A–F) but became closer together as they approached

the periosteal surface in older individuals, indicating a decrease in growth rate with age.

Only a few areas of the innermost cortex showed a prevalence of fibro-lamellar bone

tissue with primary osteons, which alternated with nonvascular parallel-fibered bone

tissue with LAGs, suggesting bone deposited slowly during most of the lifetime. There

was relatively little vascularization and the vascular canals oriented longitudinally

compared with other cervids and large mammals [11, 21]. Since the growth rate

correlates with intensity and pattern of vascularization [35-37], where lower level of

vascularization associated with longitudinal orientation of canals indicated slower bone

depositional rate, the bone histological features of the fossil insular deer further

supported their slow growth rates.

All extant specimens with EFS had fused epiphyses both in femora and tibiae (SI

Appendix, Figure S5, Table S3). The proximal epiphysis of the tibiae and femora fused

later than the distal epiphysis, and EFS developed contemporaneously with the proximal

fusions. In two mainland sika deer populations, the timing of EFS development and

proximal epiphyseal fusion showed good correspondence with the age when the rate of

growth decelerated (SI Appendix, Figures S8, S9, Table S3) and the reported age of

somatic maturity in the analyses of culled individuals [22, 23] . Notably, the extant sika

deer from the Kerama Islands showed delayed EFS development and epiphyseal fusions

compared with other investigated deer. Based on the observation of extant specimens,

we further investigated the relationship between EFS development and epiphyseal

fusion in fossil specimens (SI Appendix, Figure S5, Tables S1, 2). Though most of the

fossil specimens did not preserve both proximal and distal epiphyses, all the specimens

with fused proximal epiphysis had EFS and vice versa. Therefore, the correspondence

of epiphyseal fusion and EFS was also held in fossil species. Two insular extinct deer

from Okinawa Island showed much delayed EFS development (Min. 9 years, Max. 16

years for *C. astylodon*; Min. 5 years, Max. 11 years for the Ryukyu muntjac, see SI

Appendix, Table S1, S2) than other investigated deer, suggesting the delay of somatic

maturity.

Survivorship curve comparisons among extant and fossil cervids.

To clarify life history differences in mortality patterns, we compared the survivorship

curves of fossil *C. astylodon* (from age data of n = 45) and the four extant sika deer

populations on the Hokkaido mainland (n = 1060), Honshu mainland (n = 594),

Yakushima Island (n = 74), and Kinkazan Island (n = 264) (Fig. 2B). The deer mortality

data of the Kinkazan Island population was additionally included as it provided the

mortality pattern free from hunting pressure [38], that is also the case for *C. astylodon*

and the Ryukyu muntjac inhabiting on Pleistocene Okinawa Island without human and

other predators [38, 39]. As the age estimation of *C. astylodon* was based on the lower

third molar height, we confined the survivorship curves to juvenile and onwards for *C.*

astylodon and the sika deer. The survivorship curve of *C. astylodon* was characterized

by very low mortality from the juvenile through to the prime age period, followed by an

increase in mortality during senescence, appearing like an inversed L, which is

categorized as a type I survivorship curve [40] and characteristic to *K*-selected species.

At the other extreme, extant sika deer from Hokkaido mainland showed relatively high

mortality in the younger age class, followed by a more gradual decrease in the number

of survivors from the prime age to senescence, which is characteristic of a type II

survivorship curve. The survivorship curves of sika deer from Honshu mainland and

Yakushima Island lay between those of *C. astylodon* and Hokkaido sika deer, being

characterized by a steady decrease in survivorship until the prime age, though the

Honshu population showed an increased mortality rate after the prime age. The

survivorship curve of Kinkazan Island was also clustered with Honshu and Yakushima

Island populations, but showed lower mortality rate in the initial phase and higher rate

in the later phase. This slight difference in pattern might be associated with the lack of

hunting pressure in the Kinkazan population, however, overall, the survivorship curves

were not strongly affected by the intensity of hunting in the sika deer (see the details in

SI Appendix). Though the Kinkazan deer inhabited in the predator-free island, still their

mortality pattern was different from that of *C. astylodon*, implying further advanced

insular effect in the latter. The maximum observed age of *C. astylodon* was 25 years

[38], which was greater than that of the extant sika deer (17, 19, 17, and 21 years for the

Honshu, Hokkaido, Yakushima Island, and Kinkazan Island populations, respectively;

SI Appendix, Table S13, 14).

We additionally compared the survivorship curves between the Ryukyu muntjac (n =

65) and the extant Reeves's muntjac (n = 84) (Fig. 2C). The extant muntjac had a type II

survivorship curve as in the extant sika deer from Honshu mainland. Similarly, to *C.*

astylodon, the Ryukyu muntjac had a type I survivorship curve, indicating that its

mortality rate also did not increase until senescence, again implying strong insular effect

on life history traits in Okinawa Island.

Discussion

All of the interdisciplinary analyses of growth rate comparison, histological

investigation, and analysis of survivorship curves of both extant and extinct deer species

supported our expectation that the evolution of a new combination of life history traits

amounts to repeated establishment of a *K*-strategy in insular environments. Our findings

are amongst the first to clarify the gradual shift to *K*-selected life history for insular

forms, placing the two fossil insular deer from Okinawa Island on the extreme *K*-

strategists.

It is possible to infer the ecological process of the life history change from the current

dataset of the deer populations. As we have seen in the deer populations in mainlands,

any modifications of life history traits happened in the mainland populations (Hokkaido

and Honshu mainland sika deer, Reeves's muntjac, and *S. yabei*), similar to the

condition of the continental relatives [11]. The Japanese Archipelago is considered as

"an island" in the literature [e.g. 2], nevertheless from the current dataset, the Japanese

mainland deer may not be under the strong insular effect in the aspect of life history

traits. Further investigation of sika deer populations from Asian Continent will shed

light on the degree of insularity for Japanese mainland sika deer.

The first step of the life history change on islands could be exemplified in sika deer on

Kerama Islands, which showed the smallest body size and the slowest growth rate

among the studied extant deer populations accompanied by the bone histological

features showing the signs of slow-growth. These deer were introduced to Kerama

Islands from the Kyushu mainland ca. 400 years ago by humans (Okinawa Prefectural

Board of Education 1996). Although fast changes in body size in less than 6,000 years

have been previously documented in red deer in other islands [41], it is surprising that

this drastic change in growth trajectory could have occurred over such a short period of

time, which equates to ca. 80 generations (assuming a generation time of 5 years for the

sika deer). This change in growth trajectory might have been brought about by

phenotypic plasticity rather than genetic modification in response to the less abundant

understory vegetation in the subtropical evergreen-broadleaved forest and thus the poor

nutritional status of the Kerama deer [42], which was supported by the observation of a

possible malnourished individual (URB-MAM-55; see SI Appendix for details). It

would be relevant to obtain reliable mortality data for the Kerama deer in order to

clarify the life history shift being observed in its demographic aspect.

The next step of life history evolution would be illustrated by the Yakushima Island

deer, which also showed slower growth rate and intermediate histological features,

coupled by a moderate shift of survivorship curve toward those of *K*-selected species.

Contrary to the situation of the Kerama deer, the slower growth rate of Yakushima deer

is considered to have a genetic basis because 1) Yakushima deer are not malnourished

and have enjoyed a recent population increase [43] and 2) some of their

macromorphological features have a genetic basis [44, 45]. A long-term field

observations of Yakushima deer also supported our finding that the life history traits of

this population is shifting to *K*-strategist (Agetsuma and Agetsuma-Yanagihara,

personal communication).

The most apparent evolutionary transformation is shown in the two fossil deer isolated

on Okinawa Island over 1.5 Ma. Their growth pattern, bone histological characters, and

mortality pattern were totally different from other deer populations. As with Kerama

deer, it is possible that phenotypic plasticity partially influenced the growth trajectories

of them. We do not currently have any reliable sources of information on the nutritional

status of fossil deer, except for their dietary habits [39, 46]. However, reports of healed

bone fractures on a number of leg bones from a museum collection of fossil insular deer

of Okinawa Island [47] implies that these deer survived long enough to recover from

serious leg injuries, refuting the possibility that fossil Okinawa deer were malnourished

and their body growth was suppressed due to a lack of resources. It is also notable that

exactly the same trend was observed for the fossil deer and muntjac, further supporting

the shift toward *K*-strategy as an adaptation to the insular environment. This process

may have started after their initial settlement on Okinawa Island in part through

phenotypic plasticity of growth trajectory due to a lack of sufficient foods during

growing periods, as possibly occurred in Kerama deer. The life history change would

then have become genetically fixed through the natural selection of more *K*-shifted

individuals, as appears to have occurred in Yakushima Island deer. The final outcome

will have been the development of extreme *K*-strategists, despite their small body size.

Unfortunately, this life history trait will have made these species vulnerable to human

exploitation, because animals with a *K*-strategy have lower population recruitment [40].

Consequently, the two Okinawa deer became extinct at the time of or soon after

Paleolithic human arrival to the Okinawa Islands [48, 49], possibly due to hunting. This

might be also applicable to other island dwarfs extinct during Pleistocene [50, 51],

though the definitive role of Pleistocene *Homo* on their extinction has been recently

questioned [52].

Our findings were also in good correspondence with the worldwide observations of

life history change in insular mammals previously documented (Fig. 4). A living cervid

Odocoileus hemionus from Blakely Island showed slower growth rate than a mainland

population [8]. As the tooth eruption was also delayed in the Blakely Island deer, this

life history change may have a genetic basis, similar to our case of Yakushima Island

deer. With longer isolation but variable island sizes, the life history response varied

among the mammals from Mediterranean islands. A fossil hippo *Hippopotamus minor*

from Cyprus did not show apparent modification in bone histology and life history [12],

whereas a fossil cervid *Candiacervus* from Crete showed slight modifications toward

slower life history [11]. Sicilian pygmy elephant *Elephas falconeri* provided a

contradictory case: in the demographic aspect it was interpret to have *r*-selected life

history as the age distribution of the fossil assemblage was skewed to calves and

immature individuals [53], nevertheless a histological study suggested that this pygmy

elephant had a much slower growth rate than its mainland relatives, and sexual maturity

was within the range of extant elephants [54]. These differences in the life history

response, however, are not surprising considering the fact that these Mediterranean

islands were much bigger than those we have observed apparent life history changes (i.e.

Blakely Island, Kerama Islands, Yakushima Island, Okinawa Island, and Mallorca

Island). The most explicit change in life history was observed in *Myotragus balearicus*

from Mallorca Island, which underwent an exceptionally long time of isolation (5.2 Ma)

in predator-free environments. *Myotragus* showed a dramatic decrease in bone growth

rate and an evolution towards a slow life history [6, 7, 55], implying the commonality of

this evolutionary change with the Okinawa fossil deer. These results suggest that the

degree of modification in life history is strongly affected by both island size and

duration of isolation, with smaller islands and longer isolations resulting in life history

changes into *K*-strategists.

Conclusion

Extant Japanese sika deer are able to modify their life history traits through a

combination of phenotypic plasticity and natural selection. This ability likely underlies

the process of insular dwarfism also recorded in fossil deer (*C. astylodon* and the

Ryukyu muntjac) from Okinawa Island, which were estimated to have a *K*-strategy life

history. Combining the results of the present and previous studies on life histories of

insular mammals, we conclude that the degree of modification in life history is strongly

affected by both island size and duration of isolation, with longer isolations on smaller

islands resulting in more prominent life history changes. We demonstrate the evolution

of a gradual, transitional shift toward a 'slow life history' in large mammals from

mainland to insular populations as the main life history change accompanying insular

dwarfism.

Material and methods

Materials

Samples of extant sika deer were obtained from two mainland populations (Hokkaido

and Honshu) and two insular populations (Yakushima Island and Kerama Islands) (Fig.

1). The four populations had different properties of their habitat environments and

showed considerable differences in body size [56](Table 1). Samples of extant Reeve's

muntjac were obtained from the Honshu mainland (Fig. 1). All extant samples were

obtained from wild individuals.

Three Pleistocene fossil deer species were examined: *Cervus astylodon* and Muntiacini

gen. et sp. indet. (the Ryukyu muntjac), collected from the Hananda-Gama Cave site of

the Late Pleistocene (>20 ka, [47]) on Okinawa Island, Japan; and *Sinomegaceros yabei*,

collected from the Kumaishi-do Cave of the Late Pleistocene (16–23 ka [57]) on

Honshu mainland, Japan.

Detailed information on the extant and fossil specimens is provided in the SI Appendix

and is listed in Table 1 and SI Appendix, Tables S1–S3.

Thin sectioning and X-ray computed tomography (CT) scanning

We applied histological analyses of long bones to infer life history traits [6, 7, 11, 12,

19-21]. We used in total 62 extant and 21 fossil long bones for histological analyses

(Table 1). All of the specimens were photographed and standard measurements were

taken, following which thin sections of the midshaft of the long bones were made based

on the methodology described in previous studies [58, 59]. For the extant species, we

sampled both femur and tibia from the same individual, whereas we used either femur

or tibia for fossil specimens, as their skeletal remains were not associated. The thin

sections were photographed with a digital film scanner (Pixus Mp 800, Canon) and

analyzed using an Optiphot2-pol microscope (Nikon). Microscopic photographs were

taken with a Nikon Df camera.

Neonatal specimens of sika deer (Honshu mainland: CBM-ZZ-757) and Reeves's

muntjac (CBM-ZZ-4974) were scanned using an experimental animal X-ray CT scanner

(Latheta LCT-200, Aloka; 24-µm resolution, 80 kV, 0.2 mA) at the Okayama

University of Science, Okayama, Japan, to determine the initial diameter of the cortical

bone and an open medullary cavity in these species. Image segmentation and

visualization were performed using VG-Studio Max (Volume Graphics) version 3.1.

Age assessment

To allow comparison of growth patterns, we estimated ages as follows. For most of the

extant deer samples, age at death was assessed by tooth eruption or the number of dental

cementum annuli in the tooth root (SI Appendix, Tables S3 and S4). In the case of fossil

deer samples, the long bones were not associated with skulls and mandibles, therefore it

was required to estimate the age at death from the number of LAGs found in the long

bones. Several previous studies have reported a good correlation between the number of

LAGs and the age at death in a range of vertebrate groups [60]. However, only one

study has examined this correlation in an extant deer species [red deer (*Cervus*

elaphus)], which reported that the number of LAGs in the tibia corresponds to the actual

age but the number in the femur corresponds to the age before the deposition of the

external fundamental system (EFS) [32]. Therefore, the relationship between the actual

age at death and the number of LAGs in the bones of the extant cervids sampled was

examined first. This showed that the estimated age of the extant deer based on the

number of LAGs including those appeared in EFS in both the femur and tibia was

highly correlated with the actual age determined by tooth cementum annuli (SI

Appendix, Figure S6 and Table S4). While some old adults that were over 8 years of

age (HOUMVC-00037, CBM-ZZ-412, and URB-MAM-193) had fewer LAGs in their

long bones than their actual age, the diameters of their expanded medullary cavities

were identical or even larger than the bone diameters observed in neonates and fawns

(<1 year old), indicating that the first (and also second in URB-MAM-193) LAG is

likely eliminated in old individuals (SI Appendix, Table S4). Thus, since the number of

LAGs matched the actual age in extant Japanese cervids, the age of individuals of fossil

taxa was also estimated by counting the number of LAGs in their long bones (SI

Appendix, Table S5). Haversian bones were seen only at the inner medullary surface

and the part of the cortex where the ligaments are strongly attached (e.g., the labium

laterale in the femur) in adults, indicating the good preservations of the primary bone

tissues in most parts of the cortex.

Body mass estimation

The body mass during ontogeny was estimated from the bone diameters at the LAGs

[61]. Body mass estimation formulas for both the femora and tibiae were obtained from

[62] and the formula that uses the mediolateral diameter (MLD) of the femora and tibiae

was selected for use in this study (see SI Appendix for details). In addition to the

diameters at the LAGs, the external diaphyseal measurements [anteroposterior diameter

(APD) and MLD] were collected and used to estimate the body mass at death. Raw

measurement data of diameters at the LAGs together with the estimated body mass used

for growth curve modeling are presented in Supplementary Table S15.

Comparison of growth patterns

Growth curves of body mass were constructed for each individual specimen that were

older than yearlings (i.e., those that had more than one LAG), which had at least four

data points including starting and terminal points. The exception to this was the extant

Reeves's muntjac, for which a fawn and yearling were included in the growth curve

fitting because this species is known to have rapid growth [24]. Before fitting the

growth curves, neonatal body mass data were obtained for both the extant and fossil

taxa as the starting point (SI Appendix, Table S6, and Figure S7).

To fit growth curves to the obtained data, the number of LAGs was first transformed

into age in years. For each specimen, the body mass (W, in kg) was then formulated

with age (x, in years) using the Gompertz curve, which is described as:

$$W = aExp\left(-Exp(-b(x-c))\right)$$

where *a*, *b*, and *c* represent the asymptote (i.e., the body mass when body growth

ceases), growth rate, and inflection point, respectively. The Gompertz curve fitting was

rationalized as this model showed the highest value of goodness of fit represented by

the lowest AICc among the fitted growth curves (SI Appendix Table S7). The growth

curve parameters were then averaged for each population/species to produce a

representative growth curve for each group, and the growth rate (*b*) and inflection point

(*c*) were statistically compared among species/populations using the Tukey–Kramer

method. These analyses were conducted separately for the femora and tibiae, and all

statistical analyses were conducted in JMP Pro 16.0 (SAS Institute Inc.).

Comparison of survivorship curves

As life history change is also represented by the change in mortality pattern, we

estimated survivorship curves of both extinct and extant deer populations. Kubo et al.

[38] estimated the age at death for *C. astylodon* based on the height of the lower third

molar by applying the molar wear rate of extant sika deer. Here, the same age data (n =

45) were used to depict survivorship curves following the method of Caughley [63].

Variability of age estimation based on the molar wear rate was tested and discussed in

previous studies [38, 46, 64], and we used the most appropriate age estimates. Age data

of three sika deer populations that were used to investigate the long bone histology

(Hokkaido, Honshu, and Yakushima Island populations) and an insular population free

from hunting (Kinkazan Island) were collected from the literature [38, 65]. No reliable

published age data were available for the Yakushima Island population, so instead

histological analysis of the cementum layers of the lower first incisors was performed

on museum specimens housed in the University Museum of the University of Tokyo,

the Tochigi Prefectural Museum, and the Hokkaido University Museum to identify their

age at death. Age data were collected from 1060, 594, 74, and 264 individual deer in the

Hokkaido, Honshu, Yakushima Island, and Kinkazan Island populations, respectively.

The age at death of the fossil Ryukyu muntjac was also estimated by evaluating the

wear score of the lower third molars (n = 65). Ozaki [66] previously estimated the age at

death of the Ryukyu muntjac by applying the molar wear scoring method developed by

Chapman et al. [67] for extant Reeves's muntjac. While no age data of extant Reeves's

muntjac from Honshu mainland have been reported to date, despite their ecology being

well investigated for management purposes [24], age data have been reported for extant

Reeves's muntjac from England [67]. Therefore, these age data were used to depict the

survivorship curve of extant Reeves's muntjac (n = 84). Fossil *S. yabei* and extant

Kerama Islands deer were not included in these survivorship curve comparisons due to a

lack of reliable age data.

For more details on the materials and methods used in this study, see the SI Appendix.

Acknowledgments

We thank the following curators for the access to the museum collections under their

care: Shimoinaba, S. (Natural History Museum and Institute, Chiba, Japan), Hayashi, T.

(Tochigi Prefectural Museum), Eda, M. (The Hokkaido University Museum), Usami, K.

(Okinawa Prefectural Museum and Art Museum). Oshiro I. is acknowledged for his

advice on the fossil Okinawa deer and support for this study. We also thank Ochiai, K.

and Shimoinaba, S. for providing ecological data of sika deer and Reeves's muntjac in

Honshu. Agetsuma, N. and Agetsuma-Yanagihara, Y. are appreciated for sharing data

of their long-term field survey of Yakushima deer. We are grateful to Nakamura, K. and

Nomura, H. for technical assistance for thin-sectioning; Kodaira, S. and Kodama, R. for

CT scanning; Kawada, S. (National Museum of Nature and Science, Tokyo), van der

Geer, A. (Naturalis Biodiversity Center), Köhler, M. (ICP), Takakuwa, Y. (Gunma

Museum of Natural History), Kikuchi, H. and Umemura, Y. for providing photos of

Odocoileus hemionus, Hippopotamus minor and *Candiacervus* spp., *Myotragus*

baleanicus, Sinomegaceros yabei, Elephas falconeri, and sika deer (Yakushima Island

and Hokkaido mainland, respectively). We are grateful to V. Weisbecker and M. Köhler

for their comments on the earlier version of this manuscript.

Additional information

Funding

This work was supported by grants from the Sanyo Broadcasting Foundation (to SH and

MOK) and JSPS KAKENHI Grant Numbers JP16K18615 (to MOK) and JP19K04060

(to SH and MOK). MRSV is supported by Swiss SNF 31003A_169395.

Competing interest

We have no competing interests to declare.

Additional files

Supplementary files

⬚ Supplementary file 1. Supplementary information including additional

methodological descriptions, references, figures and tables.

Supplementary file 2. Raw data of measurements in excel format (Table S15).

Data accessibility

519 All data not presented in the manuscript are available as electronic supplementary

520 information and uploaded to Dryad Digital Repository

521 (https://doi.org/10.5061/dryad.wdbrv15n8).

References

1. Foster JB: **Evolution of Mammals on Islands**. *Nature* 1964, **202**(492):234-235.

2. van der Geer A, Lyras G, de Vos J, Dermitzakis M: **Evolution of Island**

Mammals: Adaptation and Extinction of Placental Mammals on Island.

West Sussex: John Wiley & Sons Ltd.; 2010.

3. Benitez-Lopez A, Santini L, Gallego-Zamorano J, Mila B, Walkden P,

Huijbregts MAJ, Tobias JA: **The island rule explains consistent patterns of**

body size evolution in terrestrial vertebrates. *Nat Ecol Evol* 2021, **5**(6):768-

786.

4. Larramendi A, Palombo MR: **Body Size, biology and encephalization quotient**

of *Palaeoloxodon* ex gr. *P. falconeri* from Spinagallo Cave (Hyblean plateau,

Sicily). *Hystrix* 2015, **26**(2):102-109.

5. Sander PM, Mateus O, Laven T, Knotschke N: **Bone histology indicates**

insular dwarfism in a new Late Jurassic sauropod dinosaur. *Nature* 2006,

441(7094):739-741.

6. Köhler M, Moyà-Solà S: **Physiological and life history strategies of a fossil**

large mammal in a resource-limited environment. *Proc Natl Acad Sci U S A*

540   2009, **106**(48):20354-20358.

7.   Marin-Moratalla N, Jordana X, Köhler M: **Bone histology as an approach to**

providing data on certain key life history traits in mammals: implications

for conservation biology. *Mammalian Biology* 2013, **78**(6):422-429.

8.   Long ES, Courtney KL, Lippert JC, Wall-Scheffler CM: **Reduced body size of**

insular black-tailed deer is caused by slowed development. *Oecologia* 2019,

189(3):675-685.

9.   Palkovacs EP: **Explaining adaptive shifts in body size on islands: a life**

history approach. *Oikos* 2003, **103**(1):37-44.

10.   Begon M, Harper JL, Townsend CR: **Ecology: From Individuals to**

Ecosystems, 4th Edition. Oxford: Blackwell Publishing Ltd.; 2006.

11.   Kolb C, Scheyer TM, Lister AM, Azorit C, de Vos J, Schlingemann MA,

Rossner GE, Monaghan NT, Sanchez-Villagra MR: **Growth in fossil and**

extant deer and implications for body size and life history evolution. *BMC*

Evol Biol 2015, **15**:19.

12.   Kolb C, Scheyer TM, Veitschegger K, Forasiepi AM, Amson E, Van der Geer

AA, Van den Hoek Ostende LW, Hayashi S, Sanchez-Villagra MR:

Mammalian bone palaeohistology: a survey and new data with emphasis on

island forms. *PeerJ* 2015, **3**:e1358.

13.   Lomolino MV, Sax DF, Palombo MR, van der Geer AA: **Of mice and**

mammoths: evaluations of causal explanations for body size evolution in

insular mammals. *J Biogeogr* 2012, **39**(5):842-854.

14.   Lomolino MV, van der Geer AA, Lyras GA, Palombo MR, Sax DF, Rozzi R: **Of**

mice and mammoths: generality and antiquity of the island rule. *J Biogeogr*

2013, **40**(8):1427-1439.

15.   Raia P, Meiri S: **The island rule in large mammals: paleontology meets**

ecology. *Evolution* 2006, **60**(8):1731-1742.

16.   van der Geer AA, Lyras GA, Lomolino MV, Palombo MR, Sax DF: **Body size**

evolution of palaeo-insular mammals: temporal variations and interspecific

interactions. *J Biogeogr* 2013, **40**(8):1440-1450.

17.   Otsuka H, Takahashi A: **Pleistocene vertebrate faunas in the Ryukyu Islands:**

their migration and extinction. *Tropics* 2000, **10**:25-40.

18.   Taruno H, Kawamura Y, Ishida S, Okumura K: **Yabe's giant deer and elk**

remains from the Late Pleistocene of Kumaishi-do Cave, Gifu Prefecture,

central Japan (Part 2): Postcranial bones. *Bulletin of the Osaka Museum of*

Natural History 2017, **71**:17-142.

19.   Amson E, Kolb C, Scheyer TM, Sánchez-Villagra MR: **Growth and life**

history of Middle Miocene deer (Mammalia, Cervidae) based on bone

histology. *Comptes Rendus Palevol* 2015, **14**(8):637-645.

20.   Köhler M, Marin-Moratalla N, Jordana X, Aanes R: **Seasonal bone growth and**

physiology in endotherms shed light on dinosaur physiology. *Nature* 2012,

487(7407):358-361.

21.   Klevezal GA: **Recording Structures of Mammals. Determination of Age and**

Reconstruction of Life History. Rotterdam: A. A. Balkema; 1996.

22.   Ochiai K, Asada M: **Growth in the body size of sika deer (*Cervus nippon*) on**

the Boso Peninsula, central Japan. *Journal of the Natural History Museum*

and Institute, Chiba 1995, **3**(2):223-232 (in Japanese with English abstract).

23.   Suzuki M, Onuma M, Yokoyama M, Kaji K, Yamanaka M, Ohtaishi N: **Body**

588   **size, sexual dimorphism, and seasonal mass fluctuations in a larger sika**

589   **deer subspecies, the Hokkaido sika deer (*Cervus nippon yesoensis* Heude,**

590   **1884)**. *Can J Zool* 2001, **79**(1):154-159.

591 24. Chiba Prefecture, Deer Research Group on Boso: **Science report on emergent**

592   **survey of invasive alien species (Reeves's muntjac) in Boso**. Chiba: Chiba

593   Prefecture; 2007.

594 25. Sand H, Cederlund G, Danell K: **Geographical and latitudinal variation in**

595   **growth patterns and adult body size of Swedish moose (*Alces alces*)**.

596   *Oecologia* 1995, **102**:433-442.

597 26. Monteiro LS, Falconer DS: **Compensatory growth and sexual maturity in**

598   **mice**. *Animal Science* 1966, **8**(2):179-192.

599 27. Francillon-Vieillot H, Buffrénil Vd, Castanet J, Géraudie J, Meunier FJ, Sire JY,

600   Zylberberg L, Ricqlès Ad: **Microstructure and Mineralization of Vertebrate**

601   **Skeletal Tissues**. In: *Skeletal Biomineralization: Patterns, Processes and*

602   *Evolutionary Trends*. Edited by Carter JG. New York: Van Nostrand Reinhold;

603   1989: 471-530.

604 28. Castanet J, Francillon-Vieillot H, Meunier FJ, de Ricqlès A: **Bone and**

605   **individual aging**. In: *Bone Vol 7: Bone Growth*. Edited by Hall BK. Boca

606   Raton: CRC Press; 1993: 245–283.

607 29. de Margerie E, J. Cubo, Castanet J: **Bone typology and growth rate: testing**

608   **and quantifying 'Amprino's rule' in the mallard (*Anas platyrhynchos*)**.

609   *Comptes Rendus Biologies* 2002, **325**:221-230.

610 30. Woodward HN, Padian K, Lee AH: **Skeletochronology**. In: *Bone Histology of*

611   *Fossil Tetrapods: Advancing Methods, Analysis, and Interpretation*. Edited by

Padian K, Lamm ET. Berkeley: University of California Press; 2013: 195 – 216.

31.   Andrade RCLP, Bantim RA, Lima FJ, dos Santos Campos L, Eleutério LHS,

Sayão J: **New data about the presence and absence of the external**

fundamental system in archosaurs. *Cadernos de Cultura e Ciência* 2015,

14:200-211.

32.   Calderon T, DeMiguel D, Arnold W, Stalder G, Köhler M: **Calibration of life**

history traits with epiphyseal closure, dental eruption and bone histology in

captive and wild red deer. *J Anat* 2019, **235**(2):205-216.

33.   Jordana X, Marín-Moratalla N, Moncunill-Solè B, Nacarino-Meneses C, Köhler

M: **Ontogenetic changes in the histological features of zonal bone tissue of**

ruminants: A quantitative approach. *Comptes Rendus Palevol* 2016, **15**(1-

2):255-266.

34.   Woodward HN, Horner JR, Farlow JO: **Quantification of intraskeletal**

histovariability in *Alligator mississippiensis* **and implications for vertebrate**

osteohistology. *PeerJ* 2014, **2**:e422.

35.   Orlandi-Oliveras G, Nacarino-Meneses C, Koufos GD, Kohler M: **Bone**

histology provides insights into the life history mechanisms underlying

dwarfing in hipparionins. *Sci Rep* 2018, **8**(1):17203.

36.   Castanet J, Rogers KC, Cubo J, Boisard J-J: **Periosteal bone growth rates in**

extant ratites (ostriche and emu). Implications for assessing growth in

dinosaurs. *CR Acad Sci Paris, Sciences de la vie / Life Sciences* 2000, **323**:543-

550.

37.   Lee AH, Huttenlocker AK, Padian K, Woodward HN: **Analysis of growth rates**.

In: *Bone Histology of Fossil Tetrapods: Advancing Methods, Analysis, and*

Interpretation. Edited by Padian K, Lamm ET. Berkeley: University of

California Press; 2013: 217-251.

38.   Kubo MO, Fujita M, Matsu'ura S, Kondo M, Suwa G: **Mortality profiles of**

late Pleistocene deer remains of Okinawa Island: evidence from the

Hananda-Gama cave and Yamashita-cho cave I sites. *Anthropological*

Science 2011, **119**(2):183-201.

39.   Kubo MO, Yamada E, Fujita M, Oshiro I: **Paleoecological reconstruction of**

Late Pleistocene deer from the Ryukyu Islands, Japan: combined evidence

of mesowear and stable isotope analyses. *Palaeogeogr Palaeoclimatol*

Palaeoecol 2015, **435**:159-166.

40.   Pianka ER: **Evolutionary Ecology**. New York: Harper & Row; 1978.

41.   Lister AM: **Rapid dwarfing of red deer on Jersey in the Last Interglacial**.

Nature 1989, **342**:539-542.

42.   Shiroma T, Ohta H: **Geographical variation in cranial morphology in sika**

deer with a special reference on Kerama deer In: *A report of conservation*

and management of sika deer in Kerama Islands. Edited by Okinawa Prefectural

Board of Education. Okinawa: Okinawa Prefectural Board of Education; 1996:

13-55 (in Japanese).

43.   Tsujino R, Noma N, Yumoto T: **Growth in sika deer (*Cervus nippon***

***yakushimae*) population in the western lowland forest on Yakushima Island,**

Japan. *Mammal Study* 2004, **29**:105-111.

44.   Terada C, Saitoh T: **Phenotypic and genetic divergence among island**

populations of sika deer (*Cervus nippon*) in southern Japan: a test of the

local adaptation hypothesis. *Population Ecology* 2018, **60**(3):211-221.

45.   Terada C, Tatsuzawa S, Saitoh T: **Ecological correlates and determinants in**

661   **the geographical variation of deer morphology**. *Oecologia* 2012, **169**(4):981-

662   994.

663   46.   Kubo MO, Fujita M: **Diets of Pleistocene insular dwarf deer revealed by**

664   **dental microwear texture analysis**. *Palaeogeogr Palaeoclimatol Palaeoecol*

665   2021, **562**.

666   47.   Okinawa Prefectural Museum and Art Museum (ed.): **Excavation of the**

667   **Hananda-Gama Cave site, Okinawa**. Naha: Kokusai-insatsu Co.; 2009.

668   48.   Fujita M, Yamasaki S, Katagiri C, Oshiro I, Sano K, Kurozumi T, Sugawara H,

669   Kunikita D, Matsuzaki H, Kano A *et al*: **Advanced maritime adaptation in the**

670   **western Pacific coastal region extends back to 35,000-30,000 years before**

671   **present**. *Proc Natl Acad Sci U S A* 2016, **113**(40):11184-11189.

672   49.   Fujita M, Yamasaki S, Sugawara H, Eda M: **Body size reduction in wild boar**

673   (*Sus scrofa*) **from the late Pleistocene Maehira Fissure Site in Okinawa-jima**

674   **Island, Japan, with relevance to human arrival**. *Quaternary International*

675   2014, **339**:289-299.

676   50.   Sondaar PY: **Early human exploration and exploitation of islands**. *Tropics*

677   2000, **10**:203-230.

678   51.   Alcover JA, Sans A, Palmer M: **The Extent of Extinctions of Mammals on**

679   **Islands**. *J Biogeogr* 1998, **25**(5):913-918.

680   52.   Louys J, Braje TJ, Chang CH, Cosgrove R, Fitzpatrick SM, Fujita M, Hawkins S,

681   Ingicco T, Kawamura A, MacPhee RDE *et al*: **No evidence for widespread**

682   **island extinctions after Pleistocene hominin arrival**. *Proc Natl Acad Sci U S*

683   *A* 2021, **118**(20).

53. Raia P, Barbera C, Conte M: **The fast life of a dwarfed giant**. *Evol Ecol* 2003,
17:293-312.

54. Köhler M, Palombo MR, Pretus JL, Jordana X, Moncunill-Solé B, Madurell-
Malapeira J, Marín-Moratalla N, Bromage TG: **Bone histology of the dwarf**
elephant *Palaeoloxodon falconeri* from Sicily. In: *The Second International*
Symposium on Paleohistology ISPH 2013: 2013; Bozeman, Montana, USA. 42.

55. Jordana X, Marin-Moratalla N, DeMiguel D, Kaiser TM, Köhler M: **Evidence**
of correlated evolution of hypsodonty and exceptional longevity in endemic
insular mammals. *Proc Biol Sci* 2012, **279**(1741):3339-3346.

56. Kubo MO, Takatsuki S: **Geographical body size clines in sika deer: path**
analysis to discern amongst environmental influences. *Evol Biol* 2015,
42(1):115-127.

57. Okumura K, Ishida S, Taruno H, Kawamura Y: **Yabe's giant deer and elk**
remains from the Late Pleistocene of Kumaishi-do Cave, Gifu Prefecture
(Part 1): Antlers, a skull, mandibles, and teeth. *Bulletin of the Osaka Museum*
of Natural History 2016, **70**:1-82 (in Japanese with English abstract).

58. Chinsamy A, Raath MA: **Preparation of fossil bone for histological**
examination. *Palaeontologia Africana* 1992, **29**:39-44.

59. Sander PM: **Longbone histology of the Tendaguru sauropods: implications**
for growth and biology. *Paleobiology* 2000, **26**(3):466-488.

60. Castanet J, Croci S, Aujard F, Perret M, Cubo J, de Margerie E: **Lines of**
arrested growth in bone and age estimation in a small primate: *Microcebus*
murinus. *J Zool* 2004, **263**(1):31-39.

61. Lehman TM, Woodward HN: **Modeling growth rates for sauropod dinosaurs**.

Paleobiology 2008, **34**(2):264-281.

62.   Scott KM: **Postcranial dimensions of ungulates as predictors of body mass**.

In: *Body size in mammalian paleontology: estimation and biological*

implications. Edited by Damuth J, MacFadden BJ. Cambridge: Cambridge

University Press; 1990: 301-335.

63.   Caughley G: **Analysis of Vertebrate Populations**. New York: John Wiley &

Sons Inc.; 1977.

64.   Ozaki M, Kaji K, Matsuda N, Ochiai K, Asada M, Ohba T, Hosoi E, Tado H,

Koizumi T, Suwa G *et al*: **The relationship between food habits, molar wear**

and life expectancy in wild sika deer populations. *J Zool* 2010, **280**(2):202-

212.

65.   Hokkaido Institute of Environmental Sciences: **Results of a survey related to**

sika deer and brown bear sighting on Hokkaido. In. Sapporo: Hokkaido

Institute of Environmental Sciences (in Japanese); 1997.

66.   Ozaki M: **Estimation of age at death from tooth wear of lower molars of**

excavated deer and comparison of age structures with modern deer

populations. In: *Excavation of the Hananda-Gama Cave site, Okinawa.* Edited

by Okinawa Prefectural Museum and Art Museum. Naha: Kokusai-insatsu Co.;

2009: 24-34 (in Japanese).

67.   Chapman NG, Brown WAB, Rothery P: **Assessing the age of Reeves' muntjac**

(***Muntiacus reevesi***) **by scoring wear of the mandibular molars**. *J Zool* 2005,

267(03).

68.   Palombo MR: **Endemic elephants of the Mediterranean Islands: knowledge,**

problems and perspectives. In: *The World of Elephants - International*

Congress, Rome 2001; Rome. 2001: 486-491.

69.   Kawamura Y: **Fossil record of sika deer in Japan**. In: *Sika Deer: Biology and*

Management of Native and Introduced Populations. Edited by McCullough DR,

Takatsuki S, Kaji K. Tokyo: Springer; 2009: 11-25.

Figures

Fig. 1. Locations of the sampled cervids in Japan. [†]indicates fossils excavated from
Late Pleistocene deposits. The localities of extant taxa are shown in closed stars and
those of extinct taxa are in open stars. The approximate body height (BH) is indicated
for each species/population. Photography of fossil cervids courtesy of Gunma Museum
of Natural History (*Sinomegaceros yabei*) and Okinawa Prefectural Museum & Art
Museum (*Cervus astylodon* and Muntiacini gen. et sp. indet.).

Fig. 2. Growth (A) and survivorship curves (B-C) for Japanese cervids. Insular deer have a significantly slower growth rate and a delayed growth plateau (somatic maturity) compared with mainland deer (A), and there is an ecological shift in mortality pattern between the insular and mainland deer (B, C). As the age estimation of *C. astylodon* was based on the lower third molar height, we confined the survivorship curves to juvenile and onwards for *C. astylodon* and the sika deer (B).

Fig. 3. Bone histological features of the Japanese cervids. (A–C) Midshaft cross section (A), cortex in normal light (B), and cortex in polarized light (C) of a femur of fossil *Cervus astylodon* (OPM-HAN07-1603). (D–F) Midshaft cross section (D), cortex in normal light (E), and cortex in polarized light (F) of a femur of extant sika deer (*Cervus nippon*) from Yakushima Island (TPM-M-313). (G–I) Midshaft cross section (G), cortex in normal light (H), and cortex in polarized light (I) of a femur of extant sika deer from Honshu mainland (CBM-ZZ-412). (J–L) Midshaft cross section (J), cortex in normal light (K), and cortex in polarized light (L) of a femur of fossil *Sinomegaceros yabei* (OMNH-QV-4062). Red lines in A, D, G and J and yellow arrows in B, C, E, F, H, I, and K indicate lines of arrested growth. The outer cortex is in the upper right in B, C, E and F and on the right in H, I and K. BM, body mass; EFS, external fundamental system; FBL, fibro-lamellar bone tissue; LAG, line of arrested growth; LV, longitudinal vascular canal; MC, medullary cavity; PFB, parallel-fibered bone tissue; PV, plexiform to laminar vascular canal. † Fossil cervids. Note: only cervids from the Okinawa Islands have a distinctly longer-term geographical isolation history and exhibit bone tissue indicating a slower growth rate [29].

Fig. 4. Summary of the relationship between insularity, body size and life history found in extant and fossil large mammals. The cervid species/populations of Japanese Archipelago were investigated in the present study and information of animals in other islands were obtained from references: *Odocoileus hemionus* [8], *Hippopotamus minor* [2], *Candiacervus* spp. [2], *Elephas falconeri* [2, 68], *Myotragus balearicus* [6]. Photographs of fossil mammals of Mediterranean islands courtesy of Gunma Museum of Natural History (*Elephas falconeri*), Alexandra van der Geer (*Hippopotamus minor* and *Candiacervus* spp.), and Meike Köhler (*Myotragus balearicus*). † Fossil cervids. * Life history of *Elephas falconeri* is disputed [53, 54]. ** *Muntiacus reevesi* is an artificially introduced animal in 1980's, therefore the isolation period of > 0.43 Myr is not applicable to it [24].

Table 1. Summary of the general information of sampled Japanese insular cervids.

Species/Population	Locality (island size)	Timing of immigration/isolation [reference]	Habitat type[*]	EBM (kg)[‡]	N of histological samples (Femur, Tibia)	N of demographic analysis samples [reference of data]
Extant species/population						
Cervus nippon (Hokkaido mainland)	Hokkaido (83,424 km²)	Separation from Honshu mainland population: Late Pleistocene (ca. 0.02 Ma) [69]	Subarctic coniferous forests or mixed forests	105.2	F: 7, T: 7	1060 [65]
Cervus nippon (Honshu mainland)	Honshu (227,962 km²)	Middle Pleistocene (0.43 Ma) [69]	Temperate evergreen broadleaved forests	45.5	F: 8, T: 8	594 [38]
Muntiacus reevesi		Recent introduction (1960–1980) [24]		9.2	F: 5, T: 5	84 [67]
Cervus nippon (Yakushima Island)	Yakushima Island (505 km²)	Isolation from Kyushu mainland: Late Pleistocene (ca. 0.1–0.02 Ma; see SI Appendix for details and references there in)	Subtropical evergreen broadleaved forests	37.2	F: 4, T: 4	74 [this study]
Cervus nippon (Kerama Islands)	Kerama Islands (36 km²)	Historic introduction (ca. 400 years ago) [42]	Subtropical evergreen broadleaved forests	27.1	F: 7, T: 7	NA
Fossil species						
[†]*Sinomegaceros yabei*	Honshu (227,962 km²)	Middle Pleistocene (> 0.43 Ma) [18]	NA	564.5	F: 2, T: 2	NA
[†]*Cervus astylodon*				25.4	F: 6, T: 5	45 [38]
[†]Muntiacini gen. et sp. indet.	Okinawa Island (1,207 km²)	Early Pleistocene (ca. 1.5 Ma) [17]	NA	16.5	F: 4, T: 2	65 [66]

[*] Habitat type was obtained from a reference [56]. [†] Fossil cervids. [‡] The estimated body mass (EBM) was obtained from fitted growth curves (SI Appendix, Fig. S8, S9) by averaging asymptotes for each species/population (SI Appendix, Table S8).