Genomic diversity of SARS-CoV-2 can be accelerated by a mutation in the nsp14 gene

Kosuke Takada[1,2], Mahoko Takahashi Ueda[3], Tokiko Watanabe[1] and So Nakagawa[2]*.

[1]Department of Molecular Virology, Research Institute for Microbial Diseases, Osaka University,

Suita, Osaka, Japan.

[2]Department of Molecular Life Science, Tokai University School of Medicine, Isehara, Kanagawa,

Japan.

[3]Department of Genomic Function and Diversity, Medical Research Institute, Tokyo Medical and

Dental University, Bunkyo, Tokyo, Japan

*Corresponding author

E-mail: so@tokai.ac.jp (So Nakagawa)

Abstract

Nucleotide substitution rate of severe acute respiratory syndrome coronavirus 2

(SARS-CoV-2) is relatively low compared to the other RNA viruses because coronaviruses

including SARS-CoV-2 encode non-structural protein 14 (nsp14) that is an error-correcting

exonuclease protein. In this study, to understand genome evolution of SARS-CoV-2 in the current

pandemic, we examined mutations of SARS-CoV-2 nsp14 which could inhibit its error-correcting

function. First, to obtain functionally important sites of nsp14, we examined 62 representative

coronaviruses belonging to alpha, beta, gamma, delta, and unclassified coronaviruses. As a result, 99

out of 527 amino acid sites of nsp14 were evolutionarily conserved. We then examined nsp14

sequences obtained from 28,082 SARS-CoV-2 genomes and identified 6 amino acid changes in

nsp14 mutants that were not detected in the 62 representative coronaviruses. We examined genome

substitution rates of these mutants and found that an nsp14 mutant with a proline to leucine change at

position 203 (P203L) showed a higher substitution rate (35.9 substitutions/year) than SARS-CoV-2

possessing wild-type nsp14 (19.8 substitutions/year). We confirmed that the substitution rate of the

P203L is significantly higher than those of other variants containing mutations in structural proteins.

Although the number of SARS-CoV-2 variants containing P203L mutation of nsp14 is limited (26),

these mutants appeared at least 10 times independently in the current pandemic. These results

indicated that the molecular function of nsp14 is important for survival of various coronaviruses

including SARS-CoV-2 and that some mutations such as P203L of nsp14 inhibiting its

error-correcting function are removed rapidly due to their deleterious effects.

(250 words, Max 250 words)

Introduction

Coronavirus disease of 2019 (COVID-19) became a pandemic in 2020, infecting more than 60 million people worldwide as of the end of November 2020 with no end in sight. Severe acute respiratory syndrome (SARS) coronavirus-2 (SARS-CoV-2) is the causing agent of this symptom (Lu, et al. 2020; Wu, et al. 2020; Zhou, et al. 2020; Zhu, et al. 2020). SARS-CoV-2 is a positive-sense single-stranded RNA (+ssRNA) that belongs to the coronavirus family (*Coronaviridae*), which is a member of the order *Nidovirus* (Lu, et al. 2020; Wu, et al. 2020; Zhou, et al. 2020; Zhu, et al. 2020). According to the classification of the International Committee on Taxonomy of Viruses (ICTV), *Coronaviridae* are divided into two subfamilies, *Letovirinae* and *Orthocoronavirinae*. *Orthocoronavirinae* is composed of four genera (*Alpha-*, *Beta-*, *Gamma-*, and *Delta-*) of coronaviruses (CoVs). *Betacoronavirus* including SARS-CoV-2 as well as SARS-CoV are mainly composed of 5 subgenera: *Salbecovirus*, *Hibecovirus*, *Nobecovirus, Merbecovirus*, and *Embecovirus* (Woo, et al. 2010; Nakagawa and Miyazawa 2020). SARS-CoV-2 and SARS-CoV belongs to the subgenus *Salbecovirus* (Lu, et al. 2020; Wu, et al. 2020; Zhu, et al. 2020).

One of the unique genomic characteristics of CoVs is their lower mutation rates compared with other RNA viruses (Eckerle, et al. 2007; Eckerle, et al. 2010). RNA viruses usually have a high mutation rate, which is partially caused by the relatively low fidelity of RNA-dependent RNA polymerase (RdRp) and the lack of correction of nucleotide incorporation errors. In CoVs, however, nonstructural protein 14 (nsp14) in an open reading frame 1ab (ORF1ab) cleaved by internal proteases (present in nsp3 and nsp5) (Hegyi and Ziebuhr 2002; Thiel, et al. 2003) contains an exoribonuclease (ExoN) domain that performs proofreading during replication steps (Snijder, et al. 2003). Functional inactivation experiments of nsp14 of SARS-CoV demonstrated 10-20 times higher mutation rates (Eckerle, et al. 2010; Graham, et al. 2012; Smith, et al. 2013). It is known that CoVs have the largest RNA genome of all RNA viruses (Gorbalenya, et al. 2006), and, therefore, nsp14 could be important for maintaining their genomes.

62   To understand genome evolution of SARS-CoV-2 during the current pandemic, we

63 examined SARS-CoV-2 nsp14 mutants that may impair its error-correcting function. First, to

64 examine the evolution and diversity of nsp14 in CoVs, we generated a phylogenetic tree of the nsp14

65 gene obtained from a variety of 62 CoVs belonging to the four genera as well as an unclassified one

66 (Supplementary table 1). We then evaluated evolutionary conserved amino acid sites to infer

67 functionally important sites. We examined 28,082 SARS-CoV-2 genomes that are available in the

68 GISAID database, which includes not only sequencing data but also sampling date, country, and

69 institution information (Shu and McCauley 2017). We examined various nsp14 SARS-CoV-2

70 mutants that had spread in the viral population. Among them, highly frequent nsp14 mutants that

71 were not observed among the diversified 62 CoVs were identified. Based on those results, we found

72 that an amino acid replacement from proline (P) to leucine (L) at position 203 of nsp14 increased a

73 nucleotide substitution rate of SARS-CoV-2 genomes. We validated the higher mutation rate

74 compared with those of structural proteins. These results indicate that some of variations in nsp14

75 may increase diversity of SARS-CoV-2 genomes.

Results

Conserved region of nsp14 proteins in various coronaviruses

We first generated a phylogenetic tree of amino acid sequences of the nsp14 gene obtained from representative 62 coronaviruses (CoVs); the viruses used in the analysis was summarized in Supplementary table 1. The phylogenetic tree based on the 62 amino acid sequences of nsp14 clearly showed the relationship of the four genera of CoVs (Fig. 1), which is almost identical to the results obtained by using the partial amino acid sequences of ORF1ab described in a previous report (Nakagawa and Miyazawa 2020). Our results suggest that the nsp14 gene is under negative selection due to its molecular function, and that the conserved amino acid sites in particular could be important.

Subsequently, we examined the conserved amino acid sites of the nsp14 gene among the 62 CoVs (Fig 2A). The amino acid positions of nsp14 are based on SARS-CoV, three-dimensional structure of which is available (PDB ID: 5C8S). SARS-CoV is a coronavirus belonging to the subgenus *Sarbecovirus* in the genus *Betacoronavirus*, which is the same as SARS-CoV-2. For nsp14, the number of identical amino acids was 482/527 (91.5%) in 8 representative CoVs belonging to the subgenus *Sarbecovirus*, and 359/527 (68.1%) if we added nsp14 of the Bat Hp-betacoronavirus/Zhejiang2013 (GenBank ID: NC_025217.1) belonging to the subgenus *Hibecovirus* which is known as the outgroup of *Sarbecovirus*. Moreover, 200/527 (38.0%) positions were preserved among 29 viruses of the genus *Betacoronavirus*, and 108/527 (20.5%) positions were preserved among the 61 CoVs (Supplementary table 1). If we include an unclassified CoV found in *Tropidophorus sinicus* (Chinese waterside skink) (Shi, et al. 2018), the conserved amino acid positions were 99/527 (18.8%). These results suggest that various amino acid sites of nsp14 gene are conserved among diverse CoVs because they may be functionally important. We further validated this hypothesis by calculating nonsynonymous (dN) and synonymous (dS) substitution rates on a per-site basis for the nsp14 sequences of 62 CoVs using SLAC program (see Materials and Methods) (Kosakovsky Pond and Frost 2005). Note that codon position 504 of nsp14 was calculated using 61

CoVs excluding the Chinese waterside skink CoV because it has a mixed nucleotide at that position.

As a result, no positive selection sites were found in nsp14, whereas 379 out of 527 sites were under

negative selection ($P < 0.01$, Supplementary fig 1). In total, 91 amino acids were unchanged as well

as under negative selection in nsp14 of the 62 representative CoVs.

We investigated whether the amino acids that are known to be important for the

error-correcting function of nsp14 are conserved. CoV nsp14 is an enzyme that possesses both ExoN

and N7-MTase activities. The ExoN domain of CoV nsp14 was originally identified based on

sequence similarity to distant cell homologs (Snijder, et al. 2003) and has been assigned to the

DEDD exonuclease superfamily, which includes the proofreading domains of many DNA

polymerases and other eukaryotic and prokaryotic exonucleases. The nsp14 proteins with four

conserved active site residues were distributed in three standard motifs of the primary structure:

residues 90D/92E (motif I), 191E (motif II), and 273D (motif III) (Ogando, et al. 2019; Romano, et

al. 2020). We found that the four active sites of nsp14 were conserved among the 62 CoVs examined

in this analysis (Fig 2B). Moreover, there were three zinc fingers that were essential for nsp14

function: the first zinc finger (ZF1): 207C, 210C, 226C, and 229H; the second zinc finger (ZF2):

257H, 261C, 264H, and 279C; and the third zinc finger (ZF3): 452C, 473C, 484C, and 487H.

S-adenosyl methionine (SAM)-binding motif I include residues 331D, 333G, 335P and 337A (or

337G), which were involved in the (N7 guanine)-methyl transferase reaction by nsp14. We found

that all of these residues were conserved among the nsp14 sequences of the 62 CoVs examined in

this analysis (Supplementary fig 2). These results suggest that the amino acids of the nsp14 that are

functionally important are conserved among diversified CoVs.

Diversity of nsp14 in SARS-CoV-2

Amino acid comparisons of nsp14 in the 62 representative CoVs revealed that several sites

are strongly conserved, and that some substitutions may have deleterious effects for the survive of

SARS-CoV-2. SARS-CoV-2 was first detected in humans in 2019 and has spread rapidly all over the

world in a short period of time. Therefore, some mutations of nsp14 that impair its error-correcting

function may remain in the SARS-CoV-2 population. Therefore, we attempted to identify such

deleterious mutations in the current SARS-CoV-2 population.

We downloaded the aligned nucleotide sequence of SARS-CoV2 (87,625 sequences) from

the GISAID database as of September 7, 2020. The SARS-CoV2 genomes isolated from humans

were extracted (87,304 sequences), and the sequences containing undetermined and/or mixed

nucleotides were removed; the remaining 28,082 sequences were used in subsequent analyses. The

coding region of nsp14 is translated and compared. As a result, 25,913 nsp14 amino acid sequences

(92.27%) were identical to that of the reference sequence, while 2169 sequences contain at least one

amino acid replacement in nsp14 (Supplementary table 2). Among them, the top 20 variants

possessing nsp14 with amino acids that were different from the major sequence are summarized in

Table 1. Note that frameshift mutants were excluded in Table 1 because they may affect downstream

genes as well. It was revealed that SARS-CoV-2, which acquired various mutations in nsp14, was

present, although the number of examined sequences is limited. Notably, among the 20 amino acid

nsp14 mutants, 6 amino acids (177F, 377L, 369F, 501I, 203L and 297S) were not found in the 62

nsp14 of the representative CoVs (Supplementary fig 2).

We further analyzed the evolutionary selection pressures of these 6 amino acid sites using

73 CoVs belonging to the genus *Betacoronaviruses* that includes SARS-CoV-2 (Supplementary table

3). We confirmed the 6 amino acid variants were not found in the 73 Beta-CoVs (Supplementary

table 3). The 6 conserved positions (177, 203, 297, 369, 377, and 501) were also found to be under

negative selection ($P < 0.01$, Supplementary table 4). Overall, the results indicated that the conserved

amino acid sites of nsp14 are evolutionarily selected in the genus *Betacoronaviruses* as well.

Correlation between amino acid sequences of nsp14 and nucleotide substitution rates in the

viral genome

To examine whether the 6 amino acid replacements observed in the nsp14 gene of

SARS-CoV-2 variants affect its proofreading function, we investigated a correlation between the

amino acid variants of nsp14 and the genetic diversities of SARS-CoV-2 genomes. The nucleotide

substitution rates of SARS-CoV-2 possessing mutated nsp14 variants were compared to that of

SARS-CoV-2 with wild-type nsp14. Note that sequences for which sampling date information (year,

month, and day) was not available were excluded from this analysis. Nucleotide substitutions per

year were calculated based on the regression coefficient using hCoV-19/Wuhan/Hu-1/2019 (GISAID

ID: EPI_ISL_402125) as a reference sequence (see Materials and Methods).

We found that SARS-CoV-2 containing the wild type nsp14 had 19.8 nucleotide

substitutions in whole genome per year (Table 2). Conversely, 7 SARS-CoV-2 variants containing

nsp14 with amino acid replacements showed 27.3 (L177F), 59.8 (L177F/T372I), 35.9 (P203L), 17.1

(P297S), 26.2 (S369F), 13.0 (F377L), or 21.9 (M501I) nucleotide substitutions per year (Fig 3).

Although the substitution rates of the L177F/T372I and F377L variants were found to be not reliable

because of limited time ranges, those of the other variants were clearly correlated (Fig 3). Therefore,

SARS-CoV-2 variants possessing nsp14 with the L177F, P203L, S369F, or M501I replacement may

have a particularly higher genomic mutation rate than do SARS-CoV-2 variants with wild-type

nsp14.

We further verified the elevated substitution rates found in various nsp14 mutants. We used

all sequences that did not contain any ambiguous nucleotides (shown in Supplementary table 5) to

calculate the mutation rates; however, this could cause an overestimation because of sampling biases.

Therefore, we extracted representative SARS-CoV-2 genomes by selecting the sequence showing the

oldest sampling date if the genome sequences were identical. Using the representative dataset, we

analyzed the nucleotide substitutions per year of the amino acid replacements for each site of nsp14

if amino acid replacements were present in 20 or more sequences. Note that amino acid replacements

caused by frameshifts were removed. As a result, 16 amino acid replacements in nsp14 were

evaluated, and the mean value of nucleotide substitution rate was 20.7 (Supplementary fig 3A).

Among them, we found that the P203L variant of nsp14 (nsp14-P203L) showed the highest mutation

rate (35.7 nucleotide mutations per year).

We also evaluated the substitution rates of the 7 nsp14 variants for each single cluster in

which the maximum number was observed. We assumed that variants were in the same clade if their

Pangolin lineages (Rambaut, et al. 2020) and GISAID clades were identical. We found that

nsp14-P203L variants in the Pangolin lineage B.1.1.33 showed the largest nucleotide substitution

rate (58.8 nucleotide substitutions/year) (Fig 4A, B and Table 3). Although the number of variants

was limited (6), a correlation was clearly observed (Fig 4B and Table 3). In addition, other 6

nsp-P203L variants in a different cluster (B.1) also showed a relatively high nucleotide substitution

rate (28.0 nucleotide substitutions/year) (Table3 and Supplementary fig 4). These results clearly

indicate that SARS-CoV-2 variants containing nsp14-P203L could accumulate mutations in the viral

population.

Mutations in structural genes of SARS-CoV-2 and validation of substitution rates

It was reported that nucleotide mutations in the M and E structural genes have been used to

calculate genome-wide mutation load of SARS-CoV-2, the mutations of which may not affect RNA

replication (Eskier, Karakulah, et al. 2020; Eskier, Suner, et al. 2020). Therefore, we also analyzed

nucleotide mutations of the structural genes that do not directly interact with viral RNAs: the S, M,

and E genes. As the number of nucleotide mutations in the E or M gene was only 0 or 1 in most

sequences, we summed up the number of mutations in those three genes. We found that viruses

containing nsp14-L177F or nsp14-P203L had higher rates of nucleotide substitutions in the S, M and

E genes per year than did viruses with the wild-type nsp14 (Table 2). These results suggest that

viruses carrying nsp14-L177F and nsp14-P203L viruses have an increased substitution rate.

To examine whether the substitution rates observed in the nsp14 mutants are high, we

calculated the nucleotide substitution rates of the mutants of each structural gene if they were present

in 20 or more sequences. 40 mutants containing 22, 3 and 5 amino acid replacements in sites of the S,

E and M genes, respectively, were remained. The mean rate of nucleotide substitutions per year was

18.4, and its distribution is shown in Supplementary fig 3B. We then investigated a possibility that

the increase in nucleotide substitution rates observed in the nsp14 gene occurred by chance. We

found that the nucleotide substitution rate of the nsp14-P203L mutant was over 5% in the

distribution, although the largest rate was found in the S-V6F mutant (Fig 5). However, the elevated

nucleotide substitution rate of V6F was found to be due to the effects of an outlier sample shown in

the black dot in Supplementary fig 5. In fact, if we removed the outlier sample, the nucleotide

substitution rate was reduced to 26.2. Conversely, the amino acid replacement rate of nsp14 did not

show any extreme bias in the sample. These results suggest that P203L of nsp14 has a higher rate of

genomic mutation compared with other amino acid replacements.

nsp14-P203L variant in the SARS-CoV-2 population

The nsp14-P203L variants have been isolated 26 patients mainly in Europe and North

America, with the first being reported on March 5, 2020 in Netherlands (GISAID ID:

EPI_ISL_454756, Supplementary table 5). Considering their Pangolin lineages (Rambaut, et al.

2020) as well as GISAID clades (shown in Supplementary table 5), the nsp14-P203L variant

appeared at least 10 times. In fact, the nsp14-P203L variants were observed in both of the Spike

D614 and G614 variants, which changes the efficiency of infection and have spread worldwide

(Korber, et al. 2020). However, we also found that the nsp14-P203L variants did not increase

steadily over time. As we examined, the largest number of the nsp14-P203L variants found in the

B.1.1.33 and B.1 lineages was only 6 for each (Supplementary table 5).

nsp14-P203L mutation effect on protein structure.

To estimate an effect of the P203L mutation of nps14, we first investigated the location of

the residue 203 using a tertiary structure model of SARS-CoV nsp14 (PDB ID: 5C8S) (Ma, et al.

2015). As the protein structure of SARS-CoV-2 nsp14 is currently not available in the PDB database,

we used the protein structure of SARS-CoV nsp14. Note that the amino acid residues around

position 203 of nsp14 SARS-CoV and SARS-CoV-2 are well conserved (Supplementary fig 6A).

Residue 203 is exposed on the surface of the protein and is located in the loop stretching from

residues 200 to 214 where is in the interstices of zinc and magnesium binding sites (positions 207

and 191, respectively, Fig 6). We found that the loop is close to the amino acids 1-58, which

constitutes an interaction surface for binding a co-factor, SARS-CoV nsp10 (Supplementary fig 6B)

(Ogando, et al. 2020). It was reported that the interaction with nsp10 greatly increases nucleolytic

activity of SARS-CoV nsp14 (Bouvet, et al. 2012). Therefore, we investigated the possibility that

residues at or around position 203 play an important role in nsp10-nsp14 interaction. We searched

the interaction interface using SARS-CoV nsp14-nsp10 complex structure (PDB ID: 5C8S) (Ma, et

al. 2015) by focusing the residues around position 203. Residues 199-203 (VKIGP) of the nsp14

were identified as a part of interaction interface. However, the detailed analysis of the protein

complex showed that the residues 200 and 201 alone bound directly to the residues 9 and 21 of the

nsp10 which are also well conserved between SARS-CoV and SARS-CoV-2 (Supplementary fig 6A).

These results suggest that the residue 203 may support binding to nsp10 indirectly by constituting the

interaction interface around residues 200 and 201. We further predicted the impact of the P203L

mutation on protein-protein binding affinity using BeAtMuSiC program (Dehouck, et al. 2013), and

found that the mutation decreases the binding affinity of the protein (0.38 kcal/mol). The mutation of

P to L increases hydrophobicity at the position 203 (Supplementary fig 6C and 6D), which might

alter the local structure afforded by the residues around position 203, and lead to a change in the

affinity between nsp14 and nsp10.

Discussion

In this study, we found that nsp14 was well conserved in highly diversified CoVs (Fig 2),

and that 379 out of the 527 amino acid sites of nsp14 of the 62 representative CoVs were under

negative selection (Supplementary fig 1). For SARS-CoV-2, we found that 92.27% of the viruses

had an nsp14 amino acid sequence that was identical to the reference sequence in all of the genome

sequences examined in this study (Table 1). We found that the nsp14-P203L variants showing a

higher nucleotide substitution rate did not increase steadily over time (0.09% in the current

population, Table 1), although they appeared multiple times. In fact, the nsp14-nsp10 complex of

SARS-CoV suggests that the amino acid residue at position 203 of nsp14 could be important to

interact with nsp10, around which amino acids are well conserved between SARS-CoV-2 and

SARS-CoV (Supplementary fig 6A). The nsp14-P203L mutants were suggested to weakly interact

with nsp10 (Supplementary fig 6C and 6D), possibly resulting in reduction of the proofreading

activity of nsp14. Overall, these results suggest the biological importance of nsp14 in various CoVs

including SARS-CoV-2, and that nsp14 mutations may have deleterious effects on the survival of

these viruses. In RNA viruses, rapid mutation and substitution rates are sometimes beneficial

because certain mutations allow them to survive in various situations (Mansky and Cunningham

2000; Vignuzzi, et al. 2006; Holmes 2009; Duffy 2018). However, in CoVs, high mutation rates may

not be preferred, which may be beneficial for the development of vaccines and antiviral drugs.

The frequency of nucleotide substitutions of CoVs may be affected not only by the nsp14

proofreading mechanism but also by the characteristics of other proteins, such as fidelity of

RNA-dependent RNA polymerase (nonstructural protein 12; nsp12) (Graepel, et al. 2017). In

addition, the proofreading activity of nsp14 is also affected by the amino acid substitution of nsp10,

which contributes to its interaction with nsp14 (Bouvet, et al. 2012). It is possible that amino acid

replacements in other proteins involved in genome replication, rather than nsp14-203L, actually

affected the nucleotide substitution rates of SARS-CoV-2. Therefore, amino acid replacements were

examined among nsp14-203L variants with the following genes being involved in gene replication:

nsp7, nsp8, nsp9, nsp10, nsp12, nsp13, nsp15, and nsp16. As result, proline-to-lysine replacement at position 323 (P323L) of nsp12 was frequently found (20/25) in sequences of SARS-CoV-2 possessing nsp14-203L (Supplemental table 6). However, the P323L mutation in nsp12 (nsp12-P323L) is known to accompany an aspartate-to-glycine replacement at position 614 (D614G) of S protein (Hou, et al. 2020; Korber, et al. 2020). The S-D614G mutant was first reported in the end of January 2020 in China and Germany, and after March 2020, the mutant is major variant in various regions in all over the world (Korber, et al. 2020). Therefore, P323L mutation nsp12 was found in nsp14-203L variants as well. Indeed, the genomic mutation rate of SARS-CoV-2 with the P323L amino acid replacement in nsp12, was found to be 17.7 nucleotide mutations per year (Supplemental fig 7). Therefore, it is unlikely that the P323L in nsp12 directly affects the genomic mutation rate of SARS-CoV-2 with the P203L amino acid replacement in nsp14.

According to previous reports, SARS-CoV-2 base substitutions are mostly C to U nucleotide substitutions (Simmonds 2020). Our analysis also revealed that SARS-CoV-2 has many C to U nucleotide substitutions (Supplemental table 7). Furthermore, it was clarified that there are many nucleotide substitutions from C to U in the nucleotide substitution in the population possessing nsp14-P203L (Supplemental table 7). These results suggest that the proofreading activities of nsp14-203P and nsp14-203L may not have nucleotide preferences.

It is known that alanine substitution of an amino acid that is important for the proofreading activity of nsp14 of SARS-CoV increases viral genomic mutations in cultured cell (Eckerle, et al. 2010; Smith, et al. 2013) and *in vivo* (Graham, et al. 2012). In addition, Graham et al. reported that the SARS-CoV virus is attenuated by alanine replacement of an amino acid that is important for the proofreading activity of nsp14 (Graham, et al. 2012). The effects of amino acid replacements in nsp14 on coronavirus infections, which have been found in nature, have not been investigated to date. In our study, the candidates that might affect nsp14 function were searched by focusing on the amino acid replacements of nsp14 in epidemic viruses and comparing genomic mutation rates; subsequently, we proposed candidates for amino acid replacements that may affect viral properties. However, the

molecular mechanisms underlying the amino acid replacements in SARS-CoV-2 suggested by our

study are unclear. Further studies are required to clarify the effects of the candidate amino acids on

the properties of CoVs. In conclusion, we propose that rare amino acid replacements (particularly

leucine at amino acid site 203) in coronavirus nsp14 may increase the rate of genomic mutations.

References

Abascal F, Zardoya R, Telford MJ. 2010. TranslatorX: multiple alignment of nucleotide sequences
guided by amino acid translations. Nucleic Acids Res 38:W7-13.

Bouvet M, Imbert I, Subissi L, Gluais L, Canard B, Decroly E. 2012. RNA 3'-end mismatch excision
by the severe acute respiratory syndrome coronavirus nonstructural protein nsp10/nsp14
exoribonuclease complex. Proc Natl Acad Sci U S A 109:9372-9377.

Crooks GE, Hon G, Chandonia JM, Brenner SE. 2004. WebLogo: a sequence logo generator.
Genome Res 14:1188-1190.

Darriba D, Posada D, Kozlov AM, Stamatakis A, Morel B, Flouri T. 2020. ModelTest-NG: A New
and Scalable Tool for the Selection of DNA and Protein Evolutionary Models. Mol Biol Evol
37:291-294.

Darriba D, Taboada GL, Doallo R, Posada D. 2011. ProtTest 3: fast selection of best-fit models of
protein evolution. Bioinformatics 27:1164-1165.

Dehouck Y, Kwasigroch JM, Rooman M, Gilis D. 2013. BeAtMuSiC: Prediction of changes in
protein-protein binding affinity on mutations. Nucleic Acids Res 41:W333-339.

Duffy S. 2018. Why are RNA virus mutation rates so damn high? PLoS Biol 16:e3000003.

Eckerle LD, Becker MM, Halpin RA, Li K, Venter E, Lu X, Scherbakova S, Graham RL, Baric RS,
Stockwell TB, et al. 2010. Infidelity of SARS-CoV Nsp14-exonuclease mutant virus replication is
revealed by complete genome sequencing. PLoS Pathog 6:e1000896.

Eckerle LD, Lu X, Sperry SM, Choi L, Denison MR. 2007. High fidelity of murine hepatitis virus
replication is decreased in nsp14 exoribonuclease mutants. J Virol 81:12135-12144.

Eskier D, Karakulah G, Suner A, Oktay Y. 2020. RdRp mutations are associated with SARS-CoV-2
genome evolution. PeerJ 8:e9587.

Eskier D, Suner A, Oktay Y, Karakulah G. 2020. Mutations of SARS-CoV-2 nsp14 exhibit strong
association with increased genome-wide mutation load. PeerJ 8:e10181.

Gorbalenya AE, Enjuanes L, Ziebuhr J, Snijder EJ. 2006. Nidovirales: evolving the largest RNA
virus genome. Virus Res 117:17-37.

Graepel KW, Lu X, Case JB, Sexton NR, Smith EC, Denison MR. 2017. Proofreading-Deficient
Coronaviruses Adapt for Increased Fitness over Long-Term Passage without Reversion of
Exoribonuclease-Inactivating Mutations. mBio 8.

Graham RL, Becker MM, Eckerle LD, Bolles M, Denison MR, Baric RS. 2012. A live,
impaired-fidelity coronavirus vaccine protects in an aged, immunocompromised mouse model of
lethal disease. Nat Med 18:1820-1826.

Hegyi A, Ziebuhr J. 2002. Conservation of substrate specificities among coronavirus main proteases.
Journal of General Virology 83:595-599.

Holmes EC. 2009. The Evolutionary Genetics of Emerging Viruses. Annual Review of Ecology,
Evolution, and Systematics 40:353-372.

Hou YJ, Chiba S, Halfmann P, Ehre C, Kuroda M, Dinnon KH, Leist SR, Schafer A, Nakajima N, Takahashi K, et al. 2020. SARS-CoV-2 D614G Variant Exhibits Enhanced Replication ex vivo and Earlier Transmission in vivo. bioRxiv.

Katoh K, Standley DM. 2013. MAFFT multiple sequence alignment software version 7: improvements in performance and usability. Mol Biol Evol 30:772-780.

Korber B, Fischer WM, Gnanakaran S, Yoon H, Theiler J, Abfalterer W, Hengartner N, Giorgi EE, Bhattacharya T, Foley B, et al. 2020. Tracking Changes in SARS-CoV-2 Spike: Evidence that D614G Increases Infectivity of the COVID-19 Virus. Cell 182:812-827 e819.

Kosakovsky Pond SL, Frost SD. 2005. Not so different after all: a comparison of methods for detecting amino acid sites under selection. Mol Biol Evol 22:1208-1222.

Kozlov AM, Darriba D, Flouri T, Morel B, Stamatakis A, Wren J. 2019. RAxML-NG: a fast, scalable and user-friendly tool for maximum likelihood phylogenetic inference. Bioinformatics 35:4453-4455.

Kumar S, Stecher G, Li M, Knyaz C, Tamura K. 2018. MEGA X: Molecular Evolutionary Genetics Analysis across Computing Platforms. Mol Biol Evol 35:1547-1549.

Lanave C, Preparata G, Saccone C, Serio G. 1984. A new method for calculating evolutionary substitution rates. J Mol Evol 20:86-93.

Li W, Godzik A. 2006. Cd-hit: a fast program for clustering and comparing large sets of protein or nucleotide sequences. Bioinformatics 22:1658-1659.

Lu R, Zhao X, Li J, Niu P, Yang B, Wu H, Wang W, Song H, Huang B, Zhu N, et al. 2020. Genomic characterisation and epidemiology of 2019 novel coronavirus: implications for virus origins and receptor binding. The Lancet 395:565-574.

Ma Y, Wu L, Shaw N, Gao Y, Wang J, Sun Y, Lou Z, Yan L, Zhang R, Rao Z. 2015. Structural basis and functional analysis of the SARS coronavirus nsp14-nsp10 complex. Proc Natl Acad Sci U S A 112:9436-9441.

Mansky LM, Cunningham KS. 2000. Virus mutators and antimutators. Trends in Genetics 16:512-517.

Nakagawa S, Miyazawa T. 2020. Genome evolution of SARS-CoV-2 and its virological characteristics. Inflamm Regen 40:17.

Ogando NS, Ferron F, Decroly E, Canard B, Posthuma CC, Snijder EJ. 2019. The Curious Case of the Nidovirus Exoribonuclease: Its Role in RNA Synthesis and Replication Fidelity. Front Microbiol 10:1813.

Ogando NS, Zevenhoven-Dobbe JC, van der Meer Y, Bredenbeek PJ, Posthuma CC, Snijder EJ. 2020. The Enzymatic Activity of the nsp14 Exoribonuclease Is Critical for Replication of MERS-CoV and SARS-CoV-2. J Virol 94.

Rambaut A, Holmes EC, O'Toole A, Hill V, McCrone JT, Ruis C, du Plessis L, Pybus OG. 2020. A dynamic nomenclature proposal for SARS-CoV-2 lineages to assist genomic epidemiology. Nat Microbiol 5:1403-1407.

Romano M, Ruggiero A, Squeglia F, Maga G, Berisio R. 2020. A Structural View of SARS-CoV-2
RNA Replication Machinery: RNA Synthesis, Proofreading and Final Capping. Cells 9.
Shi M, Lin XD, Chen X, Tian JH, Chen LJ, Li K, Wang W, Eden JS, Shen JJ, Liu L, et al. 2018. The
evolutionary history of vertebrate RNA viruses. Nature 556:197-202.
Shu Y, McCauley J. 2017. GISAID: Global initiative on sharing all influenza data - from vision to
reality. Euro Surveill 22.
Simmonds P. 2020. Rampant C-->U Hypermutation in the Genomes of SARS-CoV-2 and Other
Coronaviruses: Causes and Consequences for Their Short- and Long-Term Evolutionary Trajectories.
mSphere 5.
Smith EC, Blanc H, Surdel MC, Vignuzzi M, Denison MR. 2013. Coronaviruses lacking
exoribonuclease activity are susceptible to lethal mutagenesis: evidence for proofreading and
potential therapeutics. PLoS Pathog 9:e1003565.
Snijder EJ, Bredenbeek PJ, Dobbe JC, Thiel V, Ziebuhr J, Poon LLM, Guan Y, Rozanov M, Spaan
WJM, Gorbalenya AE. 2003. Unique and Conserved Features of Genome and Proteome of
SARS-coronavirus, an Early Split-off From the Coronavirus Group 2 Lineage. Journal of Molecular
Biology 331:991-1004.
Thiel V, Ivanov KA, Putics A, Hertzig T, Schelle B, Bayer S, Weissbrich B, Snijder EJ, Rabenau H,
Doerr HW, et al. 2003. Mechanisms and enzymes involved in SARS coronavirus genome expression.
J Gen Virol 84:2305-2315.
Vignuzzi M, Stone JK, Arnold JJ, Cameron CE, Andino R. 2006. Quasispecies diversity determines
pathogenesis through cooperative interactions in a viral population. Nature 439:344-348.
Weaver S, Shank SD, Spielman SJ, Li M, Muse SV, Kosakovsky Pond SL. 2018. Datamonkey 2.0:
A Modern Web Application for Characterizing Selective and Other Evolutionary Processes. Mol
Biol Evol 35:773-777.
Woo PC, Huang Y, Lau SK, Yuen KY. 2010. Coronavirus genomics and bioinformatics analysis.
Viruses 2:1804-1820.
Wu F, Zhao S, Yu B, Chen YM, Wang W, Song ZG, Hu Y, Tao ZW, Tian JH, Pei YY, et al. 2020. A
new coronavirus associated with human respiratory disease in China. Nature 579:265-269.
Zhou P, Yang XL, Wang XG, Hu B, Zhang L, Zhang W, Si HR, Zhu Y, Li B, Huang CL, et al. 2020.
A pneumonia outbreak associated with a new coronavirus of probable bat origin. Nature
579:270-273.
Zhu N, Zhang D, Wang W, Li X, Yang B, Song J, Zhao X, Huang B, Shi W, Lu R, et al. 2020. A
Novel Coronavirus from Patients with Pneumonia in China, 2019. N Engl J Med 382:727-733.
420
421

Methods

Coronavirus sequence data

Amino acid sequences of open reading frame 1ab (ORF1ab) of the 62 representative CoVs (summarized in Supplementary Table 1) were obtained from NCBI (https://www.ncbi.nlm.nih.gov/) and the GISAID EpiCoV database (https://www.gisaid.org) as reported previously (Nakagawa and Miyazawa 2020). Aligned nucleotide sequences of SARS-CoV-2 and their metadata were obtained from the GISAID EpiCoV database as of September 7, 2020 (87,625 sequences). The CoV genome sequences of which host were annotated as "Human" were extracted, and sequences containing undetermined and/or mixed nucleotides were removed. The 28,082 remaining genomes were used for the analyses. The open reading frame (ORF) of each sequence was determined from the alignment sequence based on the ORF information of hCoV-19/Wuhan/Hu-1/2019 (GISAID ID: EPI_ISL_402125, GenBank ID: NC_045512.2), which was used as the reference sequence. We then removed the gaps from the alignment and translated the nucleotide sequences. To detect amino acid replacements for each site, the amino acid sequence after the stop codon was removed and amino acid sequences were aligned using the L-INS-i program in MAFFT version v7.453 (Katoh and Standley 2013). From the sequence alignment of nsp14 of the representative CoVs, the amino acids aligned at the amino acid positions of SARS-CoV are shown using WebLogo 3 (Crooks, et al. 2004). Note that, in the alignment sequence, gap sites were removed based on the SARS-CoV nsp14 sequence.

Molecular evolutionary phylogenetic analysis

To compute phylogenetic tree of nsp14 of the 62 representative CoVs, their nsp14 amino acid sequences were aligned using the L-INS-i program in MAFFT version 7.453 (Katoh and Standley 2013). To estimate the appropriate amino acid replacement model, we used ProtTest3 version 3.4.2 (Darriba, et al. 2011) based on the Bayesian information criterion (BIC) values. We

then computed a phylogenetic tree using RAxML-NG version 1.0.0 (Kozlov, et al. 2019) with rapid

1000 bootstrap replicates (Lanave, et al. 1984).

We also generated a phylogenetic tree of SARS-CoV-2 genomes belonging to the Pangolin

lineages B.1.1.33 including nsp14-P203L variants. Using CD-HIT-EST version 4.8.1 (Li and Godzik

2006), the identical genomic sequences were removed: from 200 to 142 sequences. The 142

nucleotide sequences were aligned by the L-INS-i program in MAFFT version 7.453 (Katoh and

Standley 2013). To estimate the appropriate nucleotide substitution model, we used ModelTest-NG

(Darriba, et al. 2020) based on the Akaike information criterion (AIC) values, and GTR+I+G4 was

selected. We then computed a phylogenetic tree using RAxML-NG version 1.0.0 (Kozlov, et al.

2019) with rapid 1000 bootstrap replicates (Lanave, et al. 1984).

Mutation rate analysis

A multiple sequence alignment of SARS-CoV-2 genomes obtained from the GISAID

database was used to analyze the genomic mutation rate. The number of nucleotide substitutions per

site from between sequences was counted. Analyses were conducted using the Kimura 2-parameter

model. Codon positions included were 1st+2nd+3rd+Noncoding. All gaps were removed for each

sequence pair with a pairwise deletion option. Counting of the number of nucleotide substitution was

conducted in MEGA X (Kumar, et al. 2018). The hCoV-19/Wuhan/Hu-1/2019 (GISAID ID:

EPI_ISL_402125, GenBank ID: NC_045512.2) was selected as the reference sequence. Sequences

with the information on the collection date were used for the analysis. Note that we omitted

sequences that were collected before December 26, 2019, the date on which the reference sequences

were collected. The approximate straight line was calculated using the least-squares method using

in-house python scripts.

Measurement of natural selection pressure

TranslatorX (Abascal, et al. 2010) was used to align the nucleotide sequence based on the

aligned amino acid sequence of nsp14 of the representative CoVs. To determine evolutionary

selection pressures operating the nsp14 gene in CoVs, we computed the ratio of nonsynonymous

substitution rates (dN) and synonymous substitution rates (dS) per site (dN/dS) using the

Datamonkey software (Kosakovsky Pond and Frost 2005; Weaver, et al. 2018) through MEGA X

(Kumar, et al. 2018).

Structural analysis

A PDB file (5C8S) of the SARS-CoV nsp14-nsp10 complex was obtained from the PDB database

(https://www.rcsb.org). Using chains C and D of the structure, further analyses were performed. The

residues at the interface of the SARS-CoV nsp10-nsp14 docked complex were determined using the

"InterfaceResidues" PyMol script (http://www.pymolwiki.org/index.php/InterfaceResidues/). We

used the SARS-CoV nsp10-nsp14 docked complex for the prediction of the effect of the P203L

mutation on protein-protein interactions using the BeAtMuSiC program version 1.0

(http://babylone.3bio.ulb.ac.be/beatmusic/query.php) (Dehouck, et al. 2013). P203L mutant

generation, hydrophobicity calculation, and visualization were performed using the Discovery Studio

Visualizer software.

Acknowledgments

We thank the GISAID database (https://www.gisaid.org), which shares sequence data of

SARS-CoV-2 and related viral species. This study was supported by KAKENHI Grant-in-Aid for

Scientific Research on Innovative Areas 16H06429 (to T.W. and S.N.), 16K21723 (to T.W. and

S.N.), 16H06434 (to T.W), and 19H04843 (to S.N.); AMED Research Program on Emerging and

Re-emerging Infectious Diseases JP19fk018113 (to T.W.) and 19fk0108171 (to S.N.); 2020 Tokai

University School of Medicine Research Aid (to S.N.) The super-computing resource was provided

by Human Genome Center (the Univ. of Tokyo) and the NIG supercomputer at ROIS National

Institute of Genetics.

Author information

Affiliations

Department of Molecular Virology, Research Institute for Microbial Diseases, Osaka University,

Osaka, Japan.

Kosuke Takada and Tokiko Watanabe.

Department of Genomic Function and Diversity, Medical Research Institute, Tokyo Medical and

Dental University, Bunkyo, Tokyo, Japan

Mahoko Takahashi Ueda.

Department of Molecular Life Science, Tokai University School of Medicine, Isehara, Kanagawa,

Japan.

Kosuke Takada and So Nakagawa.

Contributions

K.T., T.W. and S.N. conceived the concept of this study. K.T., M.T.U and S.N analyzed data. K.T, M.T.U and S.N wrote the manuscript. All authors read and approved the final manuscript.

Competing interests

K.T., M.T.U., T.W. and S.N. have no competing interests.

Data Availability Statements

The data underlying this article will be shared on reasonable request to the corresponding author.

Corresponding author

Correspondence to So Nakagawa <so@tokai.ac.jp>

Figure Legends

Figure 1. Phylogenetic tree of nsp14

Maximum likelihood (ML)-based phylogenetic tree of the 62 representative CoVs. Amino acid

sequences of nsp14 in ORF1ab were used for the analysis. A red or blue circle in an internal node

correspond to bootstrap values ≥ 95% or ≥ 80 %, respectively. The virus located at the top is

SARS-CoV-2 (hCoV-19/Wuhan/Hu-1/2019).

Figure 2. Amino acid sequence alignment of selected nsp14 derived from representative CoVs

(A) Conserved amino acid residues among each group. The amino acid sequence of nsp14 of the 62

representative CoVs is shown according to the amino acid position of SARS-CoV-2, where the

gapped positions in the SARS-CoV-2 sequence were not shown. The conserved amino acid positions

are indicated in gray. (B) Four conserved active site residues involved in proofreading. The amino

acid fractions for each site were calculated based on the sequence alignment of nsp14 of the 62

representative CoVs visualized by using WebLogo (Crooks, et al. 2004).

Figure 3. Scatter plot of nucleotide substitutions in SARS-CoV-2 genomes

Genome diversity of SARS-CoV-2 containing wild-type (WT) nsp14 and 7 nsp14 mutants is shown.

The X-axis indicates the sampling date and the Y-axis indicates the number of nucleotide

substitutions in the genomes. The dotted line corresponds to the regression line.

Figure 4. Phylogeny and nucleotide substitution rate of nsp14-P203L variants in the Pangolin

lineages B.1.1.33

(A) ML tree of SARS-CoV-2 genomes in B.1.1.33-lineage. The nsp14-P203L variant is highlighted

in orange. A gray circle corresponds to bootstrap values ≥ 70%. (B) Nucleotide substitution rates of

nsp14-P203L variants belonging to the B.1.1.33-lineages. All dots of samples are displayed. The

dotted line corresponds to the regression line.

Figure 5. Frequency of nucleotide substitution rate of SARS-CoV-2 genome sequences in nsp14

or structural gene variants

The nucleotide substitution rates of SARS-CoV-2 genome sequences were calculated for each

variant containing amino acid mutations in the nsp14 or in the S, M, and E genes. We used genomes

with 20 or more nucleotide substitutions. Frequency of the substitution rate of genome sequences

with amino acid replacements for each site in nsp14 or in S, M, and E are shown in dark gray or light

gray, respectively. The thin dotted line indicates the median value for each.

Figure 6. The three-dimensional position of amino acid position 203 in nsp14

(A) Domain structure of nsp14 with residue numbers. The interaction surface with nsp10 is indicated

by the hatched box. The ExoN and N7-Mtase domains are shown in light blue and gray, respectively.

The loop from positions 200 to 214 is marked by a dark gray bar. The amino acid at position 203 in

nsp14 is shown in magenta. The amino acid positions of magnesium binding (191), nsp10 binding

(200 and 201), and zinc binding (207 and 210) in nsp14 are shown in purple, light pink and orange,

respectively. (B) Schematic representation of nsp14. The coloring is the same as in (A).

Table 1. Frequent amino acid sequence of nsp14 in SARS-CoV-2 epidemic strains.

Ranking of appearance frequency	Substitution in nsp14	Number of sequences	Frequency of appearance (%)
1	–	25913	92.27
2	A320V	192	0.68
3	F233L	144	0.51
4	S374A	123	0.44
5	L177F	116	0.41
6	T250I	61	0.22
7	F377L	51	0.18
8	A138V	39	0.14
9	V120A	35	0.12
10	T16I	34	0.12
11	S369F	33	0.12
12	G481S	31	0.11
13	M501I	28	0.10
13	D345G	28	0.10
13	N129D	28	0.10
16	P203L	26	0.09
16	L177F, T372I	26	0.09
18	P297S	25	0.09
18	T31I	25	0.09
20	D324Y	24	0.09

576 28,082 sequences that were annotated as Human and free of undetermined or mixed nucleotides, were used in

577 the analysis. Amino acid replacements were defined with hCoV-19/Wuhan/Hu-1/2019 as the reference

578 sequence. —, no mutation was found compared with the reference sequence.

579   Table 2. Nucleotide substitution rates of SARS-CoV-2 with different nsp14 variants.

Ranking of nsp14	Substitution in nsp14	Number of comparison sequences	Substitution rate per year in	
			whole genome	S+M+E gene
1	–	25471	19.8	4.31
5	L177F	114	27.3	7.45
7	F377L	51	13.0	2.81
11	S369F	33	26.2	2.48
13	M501I	28	21.9	3.07
16	P203L	24	35.9	5.29
17	L177F, T372I	26	59.8	9.82
18	P297S	25	17.1	2.48

The number of mismatched nucleotides in the alignment sequences of the reference sequence and the
comparison sequence of each group was counted, and the average value was calculated. The

hCoV-19/Wuhan/Hu-1/2019 was selected as the reference sequence. —, no mutation was found compared

with the sequences from the hCoV-19/Wuhan/Hu-1/2019.

Table 3. Nucleotide substitution rates of the 7 nsp14 variants for each single cluster in which the maximum
number was observed

Ranking of nsp14	Substitution in nsp14	Lineage (Pangolin ID)	Number of comparison sequences	Substitution rate per year in whole genome
5	L177F	B.1.1.25	54	7.8
7	F377L	B	50	8.5
11	S369F	B.1.1	17	13.4
13	M501I	B.1	11	4.4
16	P203L	B.1.1.33	6	58.8
		B.1	6	28.0
17	L177F, T372I	B.1.113	20	50.1
18	P297S	B.4	11	16.1

586 The number of mismatched nucleotides in the alignment sequences of the reference sequence and the
587 comparison sequence of each group was counted, and the average value was calculated. The
588 hCoV-19/Wuhan/Hu-1/2019 was selected as the reference sequence. A scatter plot of these nsp14 is shown in
589 Fig 4B or Supplementary figure 4.

Figure 1

(A)

Orthocoronaviruses

Alp/Bet/Gam/Del

Betacoronaviruses

Sarbecovirus

SARS-CoV-1, -2

Conserved amino acid
Non-conserved amino acid

Amino acid position

(B)

90 92   193   273

Figure 2

Figure 3

(A)

6.0E-5

Figure 4 continue

(B)

Figure 4

Figure 5

(A)

1 58   203   287   527

(B)

Figure 6