Genomic signature of shifts in selection and alkaline adaptation in highland fish

Chao Tong[1,2], Miao Li[3], Yongtao Tang[1,4], Kai Zhao[1,*]

[1] Key Laboratory of Adaptation and Evolution of Plateau Biota, Northwest Institute of Plateau
Biology, Chinese Academy of Sciences, Xining, 810001, China;

[2] Department of Biology, University of Pennsylvania, Philadelphia, PA, 19104, USA;

[3] Center for Advanced Retinal and Ocular Therapeutics, Scheie Eye Institute, University of
Pennsylvania Perelman School of Medicine, Philadelphia, PA, 19104, USA;

[4] College of Fisheries, Henan Normal University, Xinxiang, 453007, China

* Corresponding author:

Chao Tong, tongchao1990@gmail.com

Kai Zhao, zhaokai@nwipb.cas.cn

Abstract

Understanding how organisms adapt to aquatic life at high altitude is fundamental in
evolutionary biology. This objective has been addressed primarily related to hypoxia adaptation
by recent comparative studies, whereas highland fish has also long suffered extreme alkaline
environment, insight into the genomic basis of alkaline adaptation has rarely been provided.
Here, we compared the genomes or transcriptomes of 15 fish species, including two alkaline
tolerant highland fish species and their six alkaline intolerant relatives, three alkaline tolerant
lowland fish species and four alkaline intolerant species. We found putatively consistent
patterns of molecular evolution in alkaline tolerant species in a large number of shared orthologs
within highland and lowland fish taxa. Remarkably, we identified consistent signatures of
accelerated evolution and positive selection in a set of shared genes associated with ion
transport, apoptosis, immune response and energy metabolisms in alkaline tolerant species
within both highland and lowland fish taxa. This is one of the first comparative studies that
began to elucidate the consistent genomic signature of alkaline adaptation shared by highland
and lowland fish. This finding also highlights the adaptive molecular evolution changes that
support fish adapting to extreme environments at high altitude.

Keywords

Comparative genomics; Molecular evolution; Alkaline adaptation; Schizothoracine fish

Significance Statement

Little is known about how wild fish responds to extreme alkaline stress besides hypoxia at high

altitude. Comparative genomics has begun to elucidate the genomic basis of alkaline adaptation

in lowland fish, such as killifish, but insight from highland fish has lagged behind. The common

role of adaptive molecular evolution during alkaline adaptation in highland and lowland fish has

rarely been discussed. We address this question by comparing 15 fish omics data. We find

numbers of shared orthologs exhibited consistent patterns of molecular evolution in alkaline

tolerant species relative to intolerant species. We further identify remarkably consistent

signatures of rapidly evolving and positive selection in a substantial shared core of genes in

both highland and lowland alkaline tolerant species.

Introduction

Environments shape the genetic landscape of the populations that inhabit them (Witt & Huerta-Sánchez 2019). The Tibetan Plateau had experienced continuous uplift during the India-Asia collision since approximately 45 million years ago, that triggered numerous environmental changes (Li & Fang 1999; Favre et al. 2015). As elevation above sea level increases, a decrease in barometric pressure results in fewer oxygen molecules in the air, which causes hypoxia. Besides, other harsh environments highland wildlife have encountered include the long-term low temperature, and intensified ultraviolet radiation (An et al. 2001). Large numbers of endemic Tibetan animals had developed unique morphological, physiological or genetic features to tolerate such harsh conditions (Wen 2014; Tong, Tian, et al. 2017; Tong, Fei, et al. 2017). Basically, understanding how organisms adapt to extreme environment is fundamental to address many evolutionary questions, but it remains a formidable task to fully uncover the mechanism of adaptive process (Scheinfeldt & Tishkoff 2010; Tong, Fei, et al. 2017; Tong, Tian, et al. 2017). Adaptation at molecular level can occur by adaptive mutation in key genes over prolonged evolutionary time scales (Orr 2005). Recent studies employing genome-wide approaches have identified candidate genes associated with hypoxia and long-term cold response in Tibetan terrestrial wildlife adaptation to high altitude (Qu et al. 2013; Wu et al. 2020). Nevertheless, the draft genomes of very few Tibetan aquatic wildlife are sequenced (Xiao et al. 2020; Liu et al. 2019), the genomic basis of highland adaptation in aquatic animals (e.g. fish) remains largely unknown.

The schizothoracine fishes (Teleostei: Cyprinidae), the predominant fish fauna in the Tibetan Plateau, had evolved specific phenotypic characteristics to adapt to extreme aquatic environments, including hypoxia and long-term low temperature (Wu 1992; Cao et al. 1981). Recent comparative studies have identified key genes showing signals of positive selection during adaptation to such harsh environments (Yang et al. 2014; Wang et al. 2015; Kang et al. 2017), such as Hypoxia-inducible factor (HIF) (Guan et al. 2014) and Erythropoietin (EPO) (Xu et al. 2016) associated with hypoxia response, ATPase Family AAA Domain Containing 2 (ATAD2) (Tong, Fei, et al. 2017) and cAMP-dependent protein kinase catalytic subunit alpha (PRKACA) that involved into energy metabolism (Tong, Fei, et al. 2017). The main focus of previous studies in schizothoracine fishes are still on hypoxia and cold response. Notably, an increasing number of lakes in the Tibetan Plateau have been existing or towards alkaline due to

the global climate changes and human activities (Zheng 1997). Thus, the increasing alkalization
of fresh water has been the potential challenge to schizothoracine fishes. Among the
schizothoracine fishes, *Gymnocypris przewalskii przewalskii* and *Gymnocypris przewalskii*
kelukehuensis are the only two species inhabited extremely alkaline environment (Wu 1992).
Unlike other broadly distributed schizothoracine fishes, such as *Gymnocypris eckloni*,
Schizopygopsis pylzovi and *Platypharodon extremus* that inhabit in the Yellow river basin (Cao
et al. 1981; Wu 1992; Qi et al. 2012), *G. p przewalskii* only inhabits in saline and alkaline lake.
As the largest salt lake in China, Lake Qinghai (FIG. 1) is highly saline (up to 13‰) and alkaline
(up to pH 9.4) water environment, a typical salt lake with unusually high sodium, potassium and
magnesium concentration (Zheng 1997; Zhu & Wu 1975). In addition, *G. p. kelukehuensis* only
inhabits in a soda lake located at the Tsaidam Basin in the northeastern Tibetan Plateau. Lake
Keluke (FIG. 1) is also a soda lake with low salinity of 0.79‰ and high pH value up to 9.3
(Zheng 1997). Both schizothoracine fish species had developed unique physiological or genetic
features to tolerate such harsh living conditions (Tong, Fei, et al. 2017). Therefore, this provides
an exceptional model to investigate the genetic mechanisms underlying alkaline adaptation, and
may provide novel insights to fully understand the mechanism of highland adaptation in fish as
complement.

Unlike highland alkaline tolerant fish, a huge amount of studies had explored the mechanisms of
high saline and high alkaline tolerance in lowland fish species, such as killifish (e.g. *Fundulus*
heteroclitus) (Wood et al. 2010;Brennan et al. 2018; Burnett et al. 2007), tilapia (e.g.
Oreochromis niloticus) (Zhao et al. 2020; Wood et al. 1994), and salmonids (e.g. *Salmo salar*)
(Levings 2016; Lien et al. 2016). These studies had provided insights in physiology of acid-base
balance in fishes response to high salinity or high pH environments, and suggested key genes,
such as ion transport associated genes under selection during the adaptation (Lien et al. 2016).

In this study, we generated and assembled the transcriptomes of two alkaline tolerant
schizothoracine fish species, *G. p. przewalskii* and *G. p. kelukehuensis* inhabited in high pH
environment in the northeastern Tibetan Plateau (FIG. 1). We performed a comparative
genomics study together with recently sequenced schizothoracine fish transcriptomes and other
lowland fish genomes (FIG. 2A & 2B), and sought to identify consistent genomic signature
associated with alkaline adaptation in highland and lowland fishes. Specifically, we focused our

comparisons on testing whether alkaline adaptation in highland and lowland alkaline tolerant

fishes is associated with the following signatures of molecular evolution: (1) consistent patterns

of molecular evolution in protein-coding genes across the phylogeny; (2) consistent shifts in

evolutionary rates for specific genes; and (3) consistent signals of positive selection in particular

genes.

Materials and Methods

Sample collection

We collected eight adult *G. p. przewalskii* (FIG. 1) individuals (four males and four females, 172

± 0.7 g) from Lake Qinghai and eight adult *G. p. kelukehuensis* (FIG. 1) individuals (four males

and four females, 139 ± 0.3 g) from Lake Keluke using gill nets. All the fish samples were

dissected after anesthesia with MS-222 (Solarbio, Beijing, China). All individuals were classified

based on the gender and dissected after anesthesia with MS-222 (Solarbio, Beijing, China).

Tissues from gill, kidney, brain, heart and liver from each individual were collected and

immediately stored in liquid nitrogen at -80 °C. All the animal experiments were approved by the

Animal Care and Use Committees of the Northwest Institute of Plateau Biology, Chinese

Academy of Sciences (NWIPB-CY-010).

Transcriptomics

Total RNA of each tissue sample was extracted using TRIzol reagent (Invitrogen, CA, USA) in

accordance with manufacturer's instructions, and detected for quality and quantity of RNAs with

Nanodrop 1000 (NanoDrop Technologies, DE, USA) and Agilent Bioanalyzer 2100 (Agilent

Technologies, CA, USA). Equal amount of RNA from eight individual of five tissue was pooled to

construct transcriptome library as previously described (Tong, Tian, et al. 2017; Tong, Fei, et al.

2017), and sequenced with an Illumina NovaSeq 6000 yielding 150-bp paired-end (PE150)

reads.

Sequencing reads were checked for quality using FastQC (

https://www.bioinformatics.babraham.ac.uk/projects/fastqc/). Sequencing adapters and reads

with a quality score < 20 were trimmed with Trimmomatic (Bolger et al. 2014). We built a *de*

novo transcriptome assembly based on clean reads using Trinity v2.6.5 (Grabherr et al. 2011)

with default parameters. Next, we removed the redundant transcripts using CD-HIT (Fu et al.

2012) with the threshold of 0.90 and extracted the longest transcript as putative genes. We

predicted the open reading frame of each putative genes using TransDecoder

(https://github.com/TransDecoder/TransDecoder).

Additional data retrieval

We downloaded six alkaline intolerant schizothoracine fish transcriptomes (Zhou et al. 2020)

including *Oxygymnocypris stewartii*, *Schizopygopsis younghusbandi*, *Gymnocypris namensis*,

Platypharodon extremus, *Schizopygopsis pylzovi* and *Gymnocypris eckloni* from NCBI SRA

database (https://www.ncbi.nlm.nih.gov/sra) (FIG. 2B, supplementary table S1), and performed

assembly following above pipeline. In addition, we downloaded the genomes of four alkaline

intolerant fish species of *Danio rerio*, *Ctenopharyngodon idellus*, *Cyprinus carpio*, and *Carassius*

auratus, and three alkaline tolerant fish species of *Fundulus heteroclitus*, *Oreochromis niloticus*

and *Salmo salar* (FIG. 2A, supplementary table S1).

Species phylogeny and gene orthology

We obtained the phylogenetic tree of 15 fish species by pruning the Fish Tree of Life

(https://fishtreeoflife.org/) using R package, phangorn (Schliep 2011). To obtain phylogeny-

based orthology relationships between different fish taxa, we included all the predicted

proteomes of seven lowland fish genomes, and translated nucleotide sequences of protein-

coding genes from eight schizothoracine fish transcriptome assemblies into amino acid

sequences, and pooled these datasets as input for an orthology inference tool, OMA (Altenhoff

et al. 2018). In this way, we identified one-to-one, one-to-many, and many-to-many orthologs

among these 15 fish species. For further comparison, we restricted our analysis to 1:1

orthologs, that is the geneset for which only one gene from each species representing the

orthology. In addition, we extracted the shared orthologs among all fish taxa. At last, gene

ontology (GO) terms were assigned to each ortholog using Trinotate (https://trinotate.github.io/).

Pattern of molecular evolution in shared orthologs

To determine whether highland fish and lowland fish showing consistent patterns of molecular

evolution in the set of alkaline tolerant species branches across the phylogeny, we

characterized the rates of non-synonymous to synonymous rate (dN/dS) in each shared

ortholog. For this, we performed the protein sequence alignment using MUSCLE v3.8.31

(https://www.ebi.ac.uk/Tools/msa/muscle). We prepared the codon alignments of shared

orthologs, that derived from protein alignments and the corresponding DNA sequences using

PAL2NAL v.14 (Suyama et al. 2006). Then, we executed the filtration for shared ortholog

alignments with length of at least 50 codons.

We took advantage of HyPHY pipeline (Kosakovsky Pond, Poon, et al. 2020) to test the

hypotheses by comparing selective pressures (dN/dS) between *a prori* defined alkaline tolerant

fish species branches (focal foreground branch) and alkaline intolerant species fish branches

(background branch) in the specified fish phylogeny at ortholog-wide scale. Before that, a

common approach to test the hypothesis is to perform separate analyses on subsets of

sequences, and compare the parameter estimated in a *post hoc* fashion, that is one branch to

the rest of branches in a phylogenetic tree like which we previously described (Tong, Tian, et al.

2017; Tong, Fei, et al. 2017; Tong et al. 2020). While, such approach is statistically suboptimal,

and not always applicable (Kosakovsky Pond, Wisotsky, et al. 2020). In this way, we estimated

two discrete categories of dN/dS for alkaline tolerant and alkaline intolerant species under the

MG94 rate matrices combined with HKY85 model using HyPHY (Kosakovsky Pond, Poon, et al.

2020), and compared nested models with constrained relationships among them. We detected

the shift of dN/dS ratios between alkaline tolerant and alkaline intolerant (alternative model, H1),

relative to the null model that assuming all lowland fish taxa have the same dN/dS ratio (H0).

We constructed the log-likelihood ratio score for each ortholog (ΔlnL) as: $\Delta lnL = 2(lnLH_1 - lnLH_0)$

and employed the likelihood ratio test. In this way, we repeated this analysis with shared

ortholog datasets of highland taxa.

Analysis of accelerated evolution

To determine if consistent shift in evolutionary rates (e.g. acceleration) in particular genes within

alkaline tolerant species across the phylogeny, we defined rapidly evolving genes with

significantly higher dN/dS ratio for alkaline tolerant species than alkaline intolerant species ($P <$

0.05, likelihood ratio test). In this way, we identified the rapidly evolving genes in alkaline

tolerant species within highland and lowland based on earlier estimated dataset including two

discrete dN/dS ratios for each shared ortholog, respectively. In addition, these sets of rapidly

evolving genes were examined for GO enrichment relative to the full set of all shared orthologs

using R package, topGO (https://bioconductor.org/packages/release/bioc/html/topGO.html). We

finally visualized all significantly enriched GO terms remaining after the REVIGO

(http://revigo.irb.hr/) similarity filter.

Analysis of positive selection

To further determine whether consistent signals of positive selection in a set of branches

representing alkaline tolerant species across the phylogeny in specific genes, we used a

branch-site model, BUSTED (Murrell et al. 2015) to test for positive selection signal in a gene at

any site on focal branches (i.e. at least one site on at least one focal foreground branch). In

addition, we use another branch-site model, aBSREL (Smith et al. 2015) to test if positive

selection has occurred on a proportion of branches (i.e. number of focal foreground branches).

In this way, we took advantage of both models to identify positively selected genes (i.e. gene

with sites under positive selection) in alkaline tolerant species within highland fish taxa and

lowland fish taxa, respectively. We finally performed GO enrichment analysis using topGO and

REVIGO as earlier described.

Results

224   Following *de novo* assembly and annotation, each of the eight schizothoracine fish

225   transcriptome assemblies had an average of 39,921 transcripts representing the complete or

226   partial protein-coding regions of genes (supplementary table S2). We further identified a total of

227   7,309 one-to-one orthologs shared by 15 fish species (i.e. range from 2 to 15 fish species), and

228   6,241 shared 1:1 orthologs including all 15 species (supplementary table S3).

Consistent pattern of molecular evolution

We estimated two categories of dN/dS ratios for 5,748 shared 1:1 orthologs (after restricting to

1:1 orthologs with at least 50 codons) in lowland fish taxa and highland fish taxa, separately. We

found consistent patterns of molecular evolution showing significant ($P < 0.05$, likelihood ratio

test) acceleration (increased dN/dS) or deceleration (decreased dN/dS) in a set of terminal

branches of alkaline tolerant species relative to the set of terminal branches of alkaline

intolerant species in large numbers of shared orthologs within lowland fish taxa (n = 1,630) and

highland fish taxa (n = 406) (FIG. 2C, supplementary table S4).

Consistent signature of accelerated evolution

Building on earlier dataset of two discrete categories of dN/dS ratios separately representing

branches of alkaline tolerant and alkaline intolerant species across the phylogeny, we focused

on genes with significantly higher dN/dS ($P < 0.05$, likelihood ratio test) in alkaline tolerant

species, that is rapidly evolving gene (REGs) repertoire in alkaline tolerant species within

highland and lowland fish taxa. We identified 240 REGs in highland fishes and 781 REGs in

lowland fishes (FIG. 3A). Out of 41 overlapping REGs, we found a set of ion transport and

transmembrane functions associated genes, such as sodium-dependent phosphate transport

protein 2A (SLC34A1), transient receptor potential cation channel subfamily V member 1

(TRPV1), solute carrier family 7, member 3 (SLC7A3), and sodium-dependent phosphate

transporter 1-B (SLC20A1b) (FIG, 3B, supplementary table S5). Besides, we identified

overlapping REGs related to energy metabolism process, such ADP-dependent glucokinase

(ADPGK) and mitochondrial pyruvate carrier 2 (MPC2) (FIG. 3B). Given that different REGs

identified in either highland or lowland alkaline tolerant fish species, we found a number of ion

transport and ATP synthesis related genes in REGs repertoire of highland fish (FIG. 3B,

supplementary table S5), such as solute carrier family 35 member F4 (SLC35F4), calcium-

activated potassium channel subunit beta-2 (KCNMB2), ATP-sensitive inward rectifier

potassium channel 15 (KCNJ15), sodium-dependent serotonin transporter (SLC6A4), NAD-

dependent protein deacetylase sirtuin-2 (SIRT2) and NADP-dependent malic enzyme (ME1).

Similarly, in lowland fish, we also identified genes like solute carrier family 45 member 3

(SLC45A3), potassium inwardly-rectifying channel subfamily J member 2a (KCNJ2a), and ATP

synthase F0 complex subunit B1 (ATP5PB) showing rapidly evolving in alkaline tolerant species

(FIG. 3B, supplementary table S5).

Further, we did GO enrichment analysis for both REGs datasets, showing 134 significantly

enriched GO terms (biological process) in highland fish ($P < 0.05$, fisher's exact test) and 137

significantly enriched GO terms in lowland fish. After the filtration by semantic similarity of GO

terms, we found a set of enriched dominant GO terms in highland fish were related to ion

transport and transmembrane functions (FIG. 3C, supplementary table S6), such as ion

transport (GO:0006811), regulation of calcium ion transport (GO:0051924), anion transport

(GO:0006820). In addition, in lowland fish, we observed a set of enriched dominant GO terms

associated with ion transport function (FIG. 3D, supplementary table S6), such as phosphate ion

transport (GO:0006817) and response to salt stress (GO:0009651). Finally, we found 5

overlapping GO terms related to ion transport and metabolism, such as phosphate ion transport

(GO:0006817), regulation of glycogen metabolic process (GO:0070873), and

phosphatidylserine metabolic process (GO:0006658) (supplementary table S6). Collectively, this

finding suggested the consistent signature of accelerated evolution in alkaline tolerant species

within highland taxa and lowland taxa.

Consistent signature of positive selection

Out of 5,748 shared orthologs, we identified 227 orthologs with at least one site under positive

selection on all branches of alkaline tolerant species ($P < 0.05$, likelihood ratio test), namely

positively selected genes (PSGs) in highland fish (FIG. 4A, supplementary table S7), and 274

PSGs in lowland alkaline tolerant fish species (FIG. 4A, supplementary table S7). Out of 16

overlapping PSGs, we found these genes were mainly related to apoptosis (cell death), ion

transport and immune response, such as vitamin K-dependent protein C precursor (PROC), E3

ubiquitin-protein ligase SH3RF1 (SH3RF1), 14-3-3 protein beta/alpha-A (YWHABA) and

cadherin-related family member 2 (Cdhr2). Besides, we found PSGs in highland alkaline

tolerant species were also mainly involved in similar functional categories as overlapping PSGs

(FIG. 4B, supplementary table S7), such as transmembrane protein 268 (TMEM268),

transmembrane protein 266 (TMEM266), solute carrier family 35 member F4 (SLC35F4), solute

carrier family 7, member 3 (SLC35F4) and solute carrier family 7, member 3 (SLC7A3) involved

in ion transport or transmembrane functions, probable phospholipid-transporting ATPase VD

(ATP10D) involved in energy metabolism, apoptosis-inducing factor 1 (AIFM1) involved in

apoptosis, toll-like receptor 8 (TLR8) and interleukin-2 receptor subunit beta (IL2RB) related to

immune response. Similarly, in lowland alkaline tolerant fish species, these PSGs were involved

in four main categories (FIG. 4B, supplementary table S7), such as solute carrier family 2

member 15b (SLC2A15b), transmembrane protein 69 (TMEM69) and potassium voltage-gated

channel subfamily E member 4 (KCNE4) associated with ion transport function,

phosphoinositide 3-kinase regulatory subunit 4 (PIK3R4) and lysosomal-associated membrane

protein 1 (LAMP1) associated with apoptosis, CD44 antigen (CD44) and immunoglobulin-like

domain containing receptor 1b precursor (ILDR1) involved in immune response, and plasma

membrane calcium-transporting ATPase 3 (ATP2B3) associated with energy metabolism.

Further, GO enrichment results showed that a number of significantly enriched GO terms ($P <$

0.05, fisher's exact test) are related to immune response, apoptosis, protein metabolisms for

PGSs in highland alkaline tolerant species, such as inflammatory response (GO:0006954),

natural killer cell activation (GO:0030101), cell killing (GO:0001906), muscle cell apoptotic

process (GO:0010656), phagocytosis (GO:0006909), regulation of protein metabolism

(GO:0051246) and negative regulation of macromolecule biosynthesis (GO:0010558) (FIG. 4C,

supplementary table S8). Similarly, in lowland alkaline tolerant species, the significantly

enriched GO terms were mainly related to four categories as well, such as T cell mediated

cytotoxicity (GO:0001913) related to immune response, regulation of cation channel activity

(GO:2001257) and regulation of transporter activity (GO:0032409) related to transport function,

macroautophagy (GO:0016236) and regulation of apoptotic signaling pathway (GO:1902253)

related to apoptosis (cell death), glucose metabolism (GO:0006006) and sulfur amino acid

metabolism (GO:0000096) (FIG. 4D, supplementary table S8). At last, we six overlapping

signature metabolism and cell death related GO terms, such as negative regulation of metabolic

process (GO:0009892) and regulation of chromosome organization (GO:0033044)

(supplementary table S8). Collectively, this finding suggested the consistent signature of

positive selection in alkaline tolerant species within highland taxa and lowland taxa.

Discussion

Our results support above three hypotheses that alkaline tolerant species shared the consistent

patterns of molecular evolution in protein-coding genes (dN/dS) and consistent signatures of

accelerated evolution (rapidly evolving gene) and positive selection (positively selected gene) in

highland and lowland fish. Specifically, these signatures include: genes experienced consistent

acceleration in evolutionary rates (increased dN/dS) in alkaline tolerant species, which are

mainly involved in ion transport, transmembrane and energy metabolism functions; genes

showing consistent signals of positive selection in alkaline tolerant species within highland and

lowland fish taxa, these are mainly associated with ion transport/transmembrane, apoptosis (cell

death), immune response and energy metabolism processes. Altogether, this study provides

insights in understanding the common role of adaptive molecular evolution in fish adaptation to

alkaline environment, as well as adaptation to extreme environment at high-altitude.

Acid-base balance and osmoregulation

In freshwater fish, Na^+ and Cl^- are actively taken up across the gill epithelium to counter the

passive loss of osmolytes to the more dilute environment. After transition to saline or alkaline

water, fish must increase its rate to balance the osmotic loss of water to the more solute-

concentrated environment and actively excrete Na^+ and Cl^- from the gill to maintain ionic and

osmoregulation (Marshall 2005). Thus, alkaline tolerant species requires enhanced

physiological abilities including acid-base balance and osmoregulation to respond the elevation

in alkalinity or salinity of freshwater (Evans et al. 2005; Marshall 2005). Extremely alkaline

environment may accelerate the evolution of genes associated with osmoregulation in these

species survived in such harsh environment (Xu et al. 2017; Tong, Fei, et al. 2017). In this

study, we identified a set of genes associated with ion transport and transmembrane that tended

to evolve rapidly in alkaline tolerant species than their alkaline intolerant relatives (FIG. 3). This

result echoed our previous finding in *G. przewalskii* compared with other teleost fishes (alkaline

intolerance) (Tong, Fei, et al. 2017), also was in line with the rapidly evolving gene repertoire of

a wild fish, Amur ide (*Leuciscus waleckii*) that survived in extremely alkaline environment (Xu et

al. 2017). Interestingly, osmoregulation related genes including three solute carrier (SLC) genes

and one transient receptor potential cation channel (TRPV) gene exhibited consistent signature

of accelerated evolution in both highland and lowland alkaline tolerant fishes. *SLC* genes

encode transmembrane transporters for inorganic ions, amino acids, neurotransmitters, sugars,

purines and fatty acids, and other solute substrates (Dorwart et al. 2008). Recent evidences

indicted that adaptive evolution of *SLC* genes contributed to fish adaptation to high salinity and

high pH environment (Wang & Guo 2019; Tong, Fei, et al. 2017; Xu et al. 2017). Besides, we

also identified numbers of ion transport and transmembrane genes under positive selection in

alkaline tolerant fish species, such as potassium voltage-gated channel (*KCN*) genes. In killifish,

an excellent model to study extreme environment adaptation, recent genome-wide studies also

found *KCN* genes have been implicated in freshwater adaptation (transition from marine to

freshwater environment) (Brennan et al. 2018).) Besides a set of same rapidly evolving or

positively selected genes identified in both highland and lowland alkaline tolerant fish species,

we found a large number of different genes under selection but involved in similar functions,

such as anion transport (GO:0006820) and phosphate ion transport (GO:0006817), indicating

the common role of adaptive molecular evolution during alkaline adaptation in fish.

Apoptosis and cell death

Extremely alkaline stress may cause extensive damage to fish, such as inducing cell apoptosis

(Monteiro et al. 2009; Zhao et al. 2016, 2020). However, several fish species can survive in this

harsh environment, such as schizothoracine (*Gymnocypris przewalskii*) (Tong, Fei, et al. 2017;

Tong & Li 2020), Magadi tilapia (*Alcolapia grahami*) (Wilkie & Wood 1996), and Amur ide

(*Leuciscus waleckii)* (Xu et al. 2017). This may also pose a barrier for alkaline tolerant fish as

compared with alkaline intolerant fish. Significantly, positively selected *AIFM1*, *SH3RF1*,

YWHABA, *PIK3R4* and *LAMP1* of alkaline tolerant species relative to alkaline intolerant species

have enrichment in a set of apoptosis pathways, such regulation of apoptotic signaling pathway

(GO:1902253), macroautophagy (GO:0016236) and phagocytosis (GO:0006909) (FIG. 4).

Apoptosis is a form of programmed cell death that occurs in multicellular organisms, it plays a

significant role in the biochemical events lead to characteristic cell changes (morphology) and

death (Green 2011). Few direct evidences suggested the roles of these candidate genes in

response to extreme alkaline stress in fish, but numerous studies had defined their functions in

tolerance to harsh environments. *AIFM1* is a ubiquitous mitochondrial oxidoreductase involved

in apoptosis, involved in sea bream (*Sparus aurata*) response to acute environmental stress

(Bermejo-Nogales et al. 2014). *YWHABA* (14-3-3 protein beta/alpha-A) is an important gene

showing the ability to bind a multitude of functionally diverse signaling proteins, such as

transmembrane receptors (Fu et al. 2000), that involved in spotted sea bass (*Lateolabrax*

maculatus) tolerance to saline stress (Zhang et al. 2019). In addition, *LAMP1* plays an important

role in lysosome biogenesis and autophagy (Eskelinen 2006), which involved in common carp

(*Cyprinus carpio*) response to hydrogen peroxide environment. Collectively, the presence of

apoptosis related genes under positive selection may contribute to the alkaline adaptation of

fish, how these adaptive molecular changes affect the ability programmed cell death remains

unknown.

Immune response

Extreme environments (including high pH) impact on the physiology of animals in a wide variety

of ways. Recent advances in the understanding of environmental impacts were identified in

relation to specific areas of immune function, such as increase in pH resulted in a general

increase in immune function (Bowden 2008; Sridhar et al. 2020). Intriguingly, we identified

different immune genes under positive selectin in alkaline tolerant species within highland

(IL2RB, TLR8, IF144) and lowland (ILDR1 and CD44), but they all involved similar immune

functions, such as inflammatory response (GO:0006954), natural killer cell activation

(GO:0030101), and T cell mediated cytotoxicity (GO:0001913). In another word, we found

different genes but conserved pathways may underlie fish adaptation to alkaline environment.

TLR8, IF144, IL2RB and ILDR1 are important components of toll-like receptor (TLR) signaling

pathway that play key roles in the innate immune system (Rebl et al. 2010). For instance, our

previous comparative studies in alkaline tolerant fish, *G. przewalskii* identified key genes

involved in TLR pathway under selection (Tong, Fei, et al. 2017; Tong et al. 2015). In Nile tilapia

(*Oreochromis niloticus*), another alkaline tolerant species and stands as ideal model to study

extreme environment adaptation, a most recent comparative study identified numbers of

immune genes involved in natural killer cell mediated cytotoxicity and NF-kappa B signaling

pathway in response to alkalinity stress (up to pH = 8.9) (Zhao et al. 2020), and echoes our

present results. Thus, it is possible that adaptive evolution changes (e.g. positive selection)

acting on innate immune genes in alkaline tolerant species contribute to their adaptation to

extremely alkaline environment.

Energy metabolism

It is not surprising that we identified a set of genes under either accelerated evolution or positive

selection, that enriched in diverse metabolisms, such as glucose metabolism, phosphate

metabolism, sulfur amino acid metabolism and protein metabolism in alkaline tolerant species

compared with their alkaline intolerant relatives. In general, metabolism processes were

involved in fish response to varies environmental stresses (including alkaline stress) by a huge

amount of researches (Wood 1991). A most recent study highlights the genes involved in

conserved mitochondrial pathways under selection in adaptation to extreme environment

(Greenway et al. 2020). In addition to our previous studies in highland alkaline tolerant fish, we

found a number of genes associated with energy metabolism processes under selection as well,

such as mitochondrial function and protein metabolism (Tong, Fei, et al. 2017). Moreover,

increasing studies on alkaline tolerant fish species (e.g. killifish) pointed out the roles of varied

metabolisms in extreme environment (e.g. high pH) adaptation (Scott et al. 2019; Zhao et al.
2020; Wang & Guo 2019; Xu et al. 2017). Altogether, our analysis infer that the adaptive
molecular evolution of metabolism associated genes may be indispensable and common
features to alkaline adaptation as well as extreme environment adaptation in fish.

Conclusion

In summary, this comparative genomics study of 15 fish species suggests the common role of
alkaline adaptation in fish, regardless of highland or lowland background environments. Our
results also highlight that the adaptive evolution of protein-coding genes are likely to play a
crucial role in fish response to extreme environment, such as extremely high PH. Notably, this
study provides putative genomic signatures of shift in selection and alkaline adaptation in
several alkaline tolerant fish species, further study should include large scale of omics data of
more alkaline tolerant fish species and their intolerant relatives as multiple pairs to demonstrate
the genetic basis of alkaline adaptation in fish at genome-wide scale.

Author Contributions

442 C.T. and K.Z. conceived this project. C.T. designed the project. C.T., Y.T. and K.Z. collected the
443 samples. C.T. and M.L. performed the comparative genomics analyses. C.T. wrote the paper.
444 All authors read and approved the final manuscript.

Acknowledgments

447 We would like to thank four anonymous reviewers for helpful comments on this manuscript.

Funding

450 This work was supported by grants from the National Natural Science Foundation of China
451 (31870365), Joint Grant from Chinese Academy of Sciences -People's Government of Qinghai
452 Province on Sanjiangyuan National Park (LHZX-2020-01), China Biodiversity Observation and
453 Research Network, Sino BON-Inland Water Fish Diversity Observation Network.

Conflicts of Interest

The authors declare no conflict of interest.

Data Availability Statement

All the scripts required to perform the analyses are available at github.com/jiyideanjiao/Alkaline_Adaptation/. The Illumina sequencing reads have been deposited at NCBI Sequence Read Archive under the NCBI BioProject accession PRJNA684806.

Literature Cited

Altenhoff AM et al. 2018. The OMA orthology database in 2018: retrieving evolutionary relationships among all domains of life through richer web and programmatic interfaces. Nucleic Acids Res. 46:D477–D485.

An Z, John K, Warren P, Stephen P. 2001. Evolution of Asian monsoons and phased uplift of the Himalaya–Tibetan plateau since Late Miocene times. Nature. 411:62–66.

Bermejo-Nogales A et al. 2014. Metabolic and transcriptional responses of gilthead sea bream (Sparus aurata L.) to environmental stress: new insights in fish mitochondrial phenotyping. Gen. Comp. Endocrinol. 205:305–315.

Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. Bioinformatics. 30:2114–2120.

Bowden TJ. 2008. Modulation of the immune system of fish by their environment. Fish Shellfish Immunol. 25:373–383.

Brennan RS., et al. 2018. Integrative population and physiological genomics reveals mechanisms of adaptation in killifish. Mol. Biol. Evol. 35.11: 2639-2653.

Burnett KG et al. 2007. Fundulus as the premier teleost model in environmental biology: opportunities for new insights using genomics Comp. Biochem. Physiol. Part D Genomics

Proteomics. 2.4: 257-286.

Cao W, Chen Y, Wu Y, Zhu S, Others. 1981. Origin and evolution of schizothoracine fishes in
relation to the upheaval of the Qinghai-Tibetan Plateau. Science Press: Beijng.

Dorwart MR, Shcheynikov N, Yang D, Muallem S. 2008. The solute carrier 26 family of proteins
in epithelial ion transport. Physiology . 23:104–114.

Eskelinen E-L. 2006. Roles of LAMP-1 and LAMP-2 in lysosome biogenesis and autophagy.
Mol. Aspects Med. 27:495–502.

Evans DH, Piermarini PM, Choe KP. 2005. The multifunctional fish gill: dominant site of gas
exchange, osmoregulation, acid-base regulation, and excretion of nitrogenous waste. Physiol.
Rev. 85:97–177.

Favre A et al. 2015. The role of the uplift of the Qinghai-Tibetan Plateau for the evolution of
Tibetan biotas. Biol. Rev. Camb. Philos. Soc. 90:236–253.

Fu H, Subramanian RR, Masters SC. 2000. 14-3-3 proteins: structure, function, and regulation.
Annu. Rev. Pharmacol. Toxicol. 40:617–647.

Fu L, Niu B, Zhu Z, Wu S, Li W. 2012. CD-HIT: accelerated for clustering the next-generation
sequencing data. Bioinformatics. 28:3150–3152.

Grabherr MG et al. 2011. Full-length transcriptome assembly from RNA-Seq data without a
reference genome. Nat. Biotechnol. 29:644–652.

Green DR. 2011. Means to an end: apoptosis and other cell death mechanisms. Cold Spring
Harbor Laboratory Press.

Greenway R et al. 2020. Convergent evolution of conserved mitochondrial pathways underlies
repeated adaptation to extreme environments. Proc. Natl. Acad. Sci. U. S. A. 117:21822–21822.

Guan L, Chi W, Xiao W, Chen L, He S. 2014. Analysis of hypoxia-inducible factor alpha
polyploidization reveals adaptation to Tibetan Plateau in the evolution of schizothoracine fish.
BMC Evol. Biol. 14:192.

Kang J, Ma X, He S. 2017. Evidence of high-altitude adaptation in the glyptosternoid fish,
Creteuchiloglanis macropterus from the Nujiang River obtained through transcriptome analysis.

BMC Evol. Biol. 17:229.

Kosakovsky Pond SL, Poon AFY, et al. 2020. HyPhy 2.5—A Customizable Platform for
Evolutionary Hypothesis Testing Using Phylogenies. Mol. Biol. Evol. 37:295–299.

Kosakovsky Pond SL, Wisotsky SR, Escalante A, Magalis BR, Weaver S. 2020. Contrast-FEL--
a test for differences in selective pressures at individual sites among clades and sets of
branches. Mol. Biol. Evol. 38.3: 1184-1198. .

Levings CD. 2016. Ecology of Salmonids in Estuaries around the World: Adaptations, Habitats,
and Conservation. UBC Press.

Lien S et al. 2016. The Atlantic salmon genome provides insights into rediploidization. Nature.
533:200–205.

Li J, Fang X. 1999. Uplift of the Tibetan Plateau and environmental changes. Chinese science
bulletin. 44:2117–2124.

Liu H-P et al. 2019. The sequence and de novo assembly of Oxygymnocypris stewartii genome.
Scientific Data. 6:190009.

Marshall WS. 2005. Ion transport, osmoregulation, and acid-base balance. The physiology of
fishes. CRC Press.

Monteiro SM, dos Santos NMS, Calejo M, Fontainhas-Fernandes A, Sousa M. 2009. Copper
toxicity in gills of the teleost fish, Oreochromis niloticus: effects in apoptosis induction and cell
proliferation. Aquat. Toxicol. 94:219–228.

Murrell B et al. 2015. Gene-wide identification of episodic selection. Mol. Biol. Evol. 32:1365–
1371.

Orr HA. 2005. The genetic theory of adaptation: a brief history. Nat. Rev. Genet. 6:119–127.

Qi D et al. 2012. Convergent, parallel and correlated evolution of trophic morphologies in the
subfamily schizothoracinae from the Qinghai-Tibetan plateau. PLoS One. 7:e34070.

Qu Y et al. 2013. Ground tit genome reveals avian adaptation to living at high altitudes in the
Tibetan plateau. Nat. Commun. 4:2071.

Rebl A, Goldammer T, Seyfert H-M. 2010. Toll-like receptor signaling in bony fish. Vet.
Immunol. Immunopathol. 134:139–150.

Scheinfeldt LB, Tishkoff SA. 2010. Living the high life: high-altitude adaptation. Genome Biol.
11:133.

Schliep KP. 2011. phangorn: phylogenetic analysis in R. Bioinformatics. 27:592–593.Scott WC
et al. 2019. Influence of salinity and pH on bioconcentration of ionizable pharmaceuticals by the
gulf killifish, Fundulus grandis. Chemosphere. 229:434–442.

Smith MD et al, 2015. Less is more: an adaptive branch-site random effects model for efficient
detection of episodic diversifying selection. Molecular biology and evolution 32.5: 1342-1353.

Sridhar A et al. 2020. Activity profile of innate immune-related enzymes and bactericidal of
freshwater fish epidermal mucus extract at different pH. Environ. Sci. Pollut. Res. Int. doi:
10.1007/s11356-020-11173-5.

Suyama M, Torrents D, Bork P. 2006. PAL2NAL: robust conversion of protein sequence
alignments into the corresponding codon alignments. Nucleic Acids Res. 34:W609–12.

Tong C et al. 2015. Transcriptome-wide identification, molecular evolution and expression
analysis of Toll-like receptor family in a Tibet fish, Gymnocypris przewalskii. Fish Shellfish
Immunol. 46:334–345.

Tong C, Fei T, Zhang C, Zhao K. 2017. Comprehensive transcriptomic analysis of Tibetan
Schizothoracinae fish Gymnocypris przewalskii reveals how it adapts to a high altitude aquatic
life. BMC evolutionary biology. 17:74.

Tong C, Li M. 2020. Genomic signature of accelerated evolution in a saline-alkaline lake-
dwelling Schizothoracine fish. Int. J. Biol. Macromol. 149:341–347.

Tong C, Najm GM, Pinter-Wollman N, Pruitt JN, Linksvayer TA. 2020. Comparative Genomics
Identifies Putative Signatures of Sociality in Spiders. Genome Biol. Evol. 12:122–133.

Tong C, Tian F, Zhao K. 2017. Genomic signature of highland adaptation in fish: a case study in
Tibetan Schizothoracinae species. BMC Genomics. 18:948.

Wang Y et al. 2015. Evidence for Adaptation to the Tibetan Plateau Inferred from Tibetan Loach
Transcriptomes. Genome Biol. Evol. 7:2970–2982.

Wang Y, Guo B. 2019. Adaption to extreme environments: a perspective from fish genomics.
Rev. Fish Biol. Fish. 29:735–747.

Wen LY. 2014. Uplift of the tibetan plateau influenced the morphological evolution of animals. J.
Agric. Sci. 6:244.

Wilkie MP, Wood CM. 1996. The adaptations of fish to extremely alkaline environments. Comp.
Biochem. Physiol. B Biochem. Mol. Biol. 113:665–673.

Witt KE, Huerta-Sánchez E. 2019. Convergent evolution in human and domesticate adaptation
to high-altitude environments. Philos. Trans. R. Soc. Lond. B Biol. Sci. 374:20180235.

Wood C et al. 1994. Urea production, acid-base regulation and their interactions in the Lake
Magadi tilapia, a unique teleost adapted to a highly alkaline environment. Journal of
Experimental Biology. 189:13–36.

Wood CM. 1991. Acid-Base and Ion Balance, Metabolism, and their Interactions, after
Exhaustive Exercise in Fish. J. Exp. Biol. 160:285–308.

Wood CM, Bucking C, Grosell M. 2010. Acid--base responses to feeding and intestinal Cl--
uptake in freshwater-and seawater-acclimated killifish, Fundulus heteroclitus, an agastric
euryhaline teleost. J. Exp. Biol. 213:2681–2692.

Wu DD, Yang CP, Wang MS, Dong KZ. 2020. Convergent genomic signatures of high-altitude
adaptation among domestic mammals. National Science Review. 7:952–963.

Wu Y. 1992. The Fishes of the Qinghai-Tibetan Plateau. Sichuan Publishing House of Science
& Technology.

Xiao S et al. 2020. Genome of Tetraploid Fish Schizothorax o'connori Provides Insights into
Early Re-diploidization and High-Altitude Adaptation. iScience. 23:101497.

Xu J et al. 2017. Genomic Basis of Adaptive Evolution: The Survival of Amur Ide (Leuciscus
waleckii) in an Extremely Alkaline Environment. Mol. Biol. Evol. 34:145–159.

Xu Q et al. 2016. Analysis of the erythropoietin of a Tibetan Plateau schizothoracine fish
(Gymnocypris dobula) reveals enhanced cytoprotection function in hypoxic environments. BMC
Evol. Biol. 16:11.

Yang L, Wang Y, Zhang Z, He S. 2014. Comprehensive transcriptome analysis reveals
accelerated genic evolution in a Tibet fish, Gymnodiptychus pachycheilus. Genome Biol. Evol.
7:251–261.

Zhang K-Q et al. 2019. 14-3-3 gene family in spotted sea bass (Lateolabrax maculatus):
Genome-wide identification, phylogenetic analysis and expression profiles after salinity stress.
Comp. Biochem. Physiol. A Mol. Integr. Physiol. 235:1–11.

Zhao Y et al. 2020. Transcriptome changes for Nile tilapia (Oreochromis niloticus) in response
to alkalinity stress. Comp. Biochem. Physiol. Part D Genomics Proteomics. 33:100651.

Zhao Y, Wu J-W, Wang Y, Zhao J-L. 2016. Role of miR-21 in alkalinity stress tolerance in
tilapia. Biochem. Biophys. Res. Commun. 471:26–33.

Zheng M. 1997. An Introduction to Saline Lakes on the Qinghai—Tibet Plateau. Springer
Science & Business Media.

Zhou C et al. 2020. Comprehensive transcriptome data for endemic Schizothoracinae fish in the
Tibetan Plateau. Sci Data. 7:28.

Zhu S, Wu Y. 1975. Study of fish fauna in Qinghai Lake. Science Press: Beijing.

Figure legend

FIG. 1. Overview of geographic distributions of two alkaline tolerant schizothoracine fish species

and their habitats in the northeastern Tibetan Plateau. Map depicting the geographic

distributions of *G. p. przewalskii* in Lake Qinghai, and *G. p. kelukehuensis* in Lake Keluke.

Photos showing the representative specimens of *G. p. przewalskii* and *G. p. kelukehuensis*.

Photo credit: Chao Tong and Kai Zhao.

FIG. 2. Consistent patterns of molecular evolution in alkaline tolerant fish species.

(A-B) Species trees for lowland and highland fish taxa were pruned from the Fish Tree of Life

(https://fishtreeoflife.org/). Alkaline tolerant taxa and alkaline intolerant taxa are depicted in

orange and sky blue, respectively. Two schematic dragram depicting the comparisons made

between alkaline tolerant and alkaline intolerant fish species.

(C) Number of tested orthologs (N = 5,748), and number of orthologs under consistent shifts in

rates of molecular evolution in alkaline tolerant relative to alkaline intolerant species

withinhighland fish taxa and lowland fish taxa ($P < 0.05$, likelihood ratio test).

FIG. 3. Consistent signature of accelerated evolution in alkaline tolerant species within highland

and lowland fish taxa.

(A) Venn diagram depicting numbers of rapidly evolving genes (REGs) in highland alkaline

tolerant species, lowland alkaline tolerant species and both.

(B) Highland fish specific REGs, lowland fish specific REGs and overlapping REGs mainly

related to ion transport, transmembrane and energy metabolism functions. The table showing

the representative REGs under each category.

(C) REVIGO plot depicting the dominant enriched GO terms for REGs in highland alkaline

tolerant fish. The scale of dark dot indicating the number of included enriched GO terms under

a dominant GO term, the color scale representing the P value transformed by log10.

(D) REVIGO plot depicting the dominant enriched GO terms for REGs in lowland alkaline

tolerant fish.

FIG. 4. Consistent signature of positive selection in alkaline tolerant species within highland and

lowland fish taxa.

(A) Venn diagram depicting numbers of positively selected genes (PSGs) in highland alkaline

tolerant species, lowland alkaline tolerant species and both.

(B) Highland fish specific PSGs, lowland fish specific PSGs and overlapping PSGs mainly

related to four categories, including ion transport/transmembrane, apoptosis/cell death, immune

response and energy metabolism. The table showing the representative PSGs under each

category.

(C) REVIGO plot depicting the dominant enriched GO terms for PSGs in highland alkaline

tolerant fish. The dark dot scale indicating the number of included GO terms under a dominant

GO term, the color scale representing the P value transformed by log10.

(D) REVIGO plot depicting the dominant enriched GO terms for PSGs in lowland alkaline

tolerant fish.

Supplementary table capture

supplementary table S1: Information for additional omics data.

supplementary table S2: Statistics of transcriptome assembly and protein-coding genes.

supplementary table S3: Statistics of orthologs in 15 fish species.

supplementary table S4: Patterns of molecular evolution in alkaline tolerant and alkaline intolerant species within lowland fish taxa and highland fish taxa.

supplementary table S5: Rapidly evolving gene repertoires in highland alkaline tolerant and lowland alkaline tolerant species.

supplementary table S6: Significantly enriched GO terms for rapidly evolving gene of highland alkaline tolerant and lowland alkaline tolerant species.

supplementary table S7: Positively selected gene repertoires in highland alkaline tolerant and lowland alkaline tolerant species.

supplementary table S8: Significantly enriched GO terms for positively selected gene of highland alkaline tolerant and lowland alkaline tolerant species.

Lake Keluke

Lake Qinghai

Gymnocypris przewalskii kelukehuensis

Gymnocypris przewalskii przewalskii

Lake Keluke

Lake Qinghai

Tibetan
Plateau

80   100   120

N

A

Lowland fish taxa

Alkaline tolerant
Alkaline intolerant

○ *Carassius auratus* (Caur)
○ *Cyprinus carpio* (Ccar)
○ *Ctenopharyngodon idellus* (Cide)
○ *Danio rerio* (Drer)
○ *Fundulus heteroclitus* (Fhet)
○ *Oreochromis niloticus* (Onil)
○ *Salmo salar* (Ssal)

B

Highland fish taxa

○ *Oxygymnocypris stewartii* (Oste)
○ *Schizopygopsis younghusbandi* (Syou)
○ *Gymnocypris namensis* (Gnam)
○ *Platypharodon extremus* (Pext)
○ *Schizopygopsis pylzovi* (Spyl)
○ *Gymnocypris eckloni* (Geck)
○ *Gymnocypris przewalskii przewalskii* (Gprz)
○ *Gymnocypris przewalskii kelukehuensis* (Gkel)

C

	Number of tested orthologs	Number of orthologs with significance
Lowland fish taxa	5,748	1,630
Highland fish taxa	5,748	406

A

B

ion transport/transmenbrane		apoptosis cell death	immune response	energy metabolism
TMEM268 TMEM266 TMEM216 CLDN15 TRPV1	SLC6A4 SLC35F4 SLC7A3 SCN2B SLC13A5	AIFM1	IL2RB TLR8 IF144	ATP10D
PROC Cdhr2		SH3RF1 YWHABA		TPI1b
SLC2A15b SLC12A9 TMEM69 KCNE4		PIK3R4 LAMP1	ILDR1 CD44	ATP2B3

C   D