Genomic signature of shifts in selection and alkaline adaptation in highland fish

Chao Tong[1,2], Miao Li[3], Yongtao Tang[1,4], Kai Zhao[1,*]

[1] Key Laboratory of Adaptation and Evolution of Plateau Biota, Northwest Institute of Plateau
Biology, Chinese Academy of Sciences, Xining, 810001, China;

[2] Department of Biology, University of Pennsylvania, Philadelphia, PA, 19104, USA;

[3] Center for Advanced Retinal and Ocular Therapeutics, Scheie Eye Institute, University of
Pennsylvania Perelman School of Medicine, Philadelphia, PA, 19104, USA;

[4] College of Fisheries, Henan Normal University, Xinxiang, 453007, China

* Corresponding author:

Chao Tong, tongchao1990@gmail.com, https://orcid.org/0000-0001-5202-5507

Kai Zhao, zhaokai@nwipb.cas.cn

Abstract

Understanding how organisms adapt to aquatic life at high altitude is fundamental in

evolutionary biology. This objective has been addressed primarily related to hypoxia adaptation

by recent comparative studies, whereas highland fish has also long suffered extreme alkaline

environment, insight into the genomic basis of alkaline adaptation has rarely been provided.

Here, we compared the genomes or transcriptomes of 15 fish species, including two alkaline

tolerant highland fish species and their six alkaline intolerant relatives, three alkaline tolerant

lowland fish species and four alkaline intolerant species. We found putatively consistent

patterns of molecular evolution in alkaline tolerant species in a large number of shared orthologs

within highland and lowland fish taxa. Remarkably, we identified consistent signatures of

accelerated evolution and positive selection in a set of shared genes associated with ion

transport, apoptosis, immune response and energy metabolisms in alkaline tolerant species

within both highland and lowland fish taxa. This is one of the first comparative studies that

began to elucidate the consistent genomic signature of alkaline adaptation shared by highland

and lowland fish. This finding also highlights the adaptive molecular evolution changes that

support fish adapting to extreme environments at high altitude.

Keywords

Comparative genomics; Molecular evolution; Alkaline adaptation; Schizothoracine fish

Significance Statement

Little is known about how wild fish responds to extreme alkaline stress besides hypoxia at high altitude. Comparative genomics has begun to elucidate the genomic basis of alkaline adaptation in lowland fish, such as killifish, but insight from highland fish has lagged behind. The common role of adaptive molecular evolution during alkaline adaptation in highland and lowland fish has rarely been discussed. We address this question by comparing 15 fish omics data. We find numbers of shared orthologs exhibited consistent patterns of molecular evolution in alkaline tolerant species relative to intolerant species. We further identify remarkably consistent signatures of rapidly evolving and positive selection in a substantial shared core of genes in both highland and lowland alkaline tolerant species.

Introduction

Environments shape the genetic landscape of the populations that inhabit them (Witt & Huerta-Sánchez 2019). The Tibetan Plateau had experienced continuous uplift during the India-Asia collision since approximately 45 million years ago, that triggered numerous environmental changes (Li & Fang 1999; Favre et al. 2015). As elevation above sea level increases, a decrease in barometric pressure results in fewer oxygen molecules in the air, which causes hypoxia. Besides, other harsh environments highland wildlife have encountered include the long-term low temperature, and intensified ultraviolet radiation (An et al. 2001). Large numbers of endemic Tibetan animals had developed unique morphological, physiological or genetic features to tolerate such harsh conditions (Wen 2014; Tong, Tian, et al. 2017; Tong, Fei, et al. 2017). Basically, understanding how organisms adapt to extreme environment is fundamental to address many evolutionary questions, but it remains a formidable task to fully uncover the mechanism of adaptive process (Scheinfeldt & Tishkoff 2010; Tong, Fei, et al. 2017; Tong, Tian, et al. 2017). Adaptation at molecular level can occur by adaptive mutation in key genes over prolonged evolutionary time scales (Orr 2005). Recent studies employing genome-wide approaches have identified candidate genes associated with hypoxia and long-term cold response in Tibetan terrestrial wildlife adaptation to high altitude (Qu et al. 2013; Wu et al. 2020). Nevertheless, the draft genomes of very few Tibetan aquatic wildlife are sequenced (Xiao et al. 2020; Liu et al. 2019), the genomic basis of highland adaptation in aquatic animals (e.g. fish) remains largely unknown.

The schizothoracine fishes (Teleostei: Cyprinidae), the predominant fish fauna in the Tibetan Plateau, had evolved specific phenotypic characteristics to adapt to extreme aquatic environments, including hypoxia and long-term low temperature (Wu 1992; Cao et al. 1981). Recent comparative studies have identified key genes showing signals of positive selection during adaptation to such harsh environments (Yang et al. 2014; Wang et al. 2015; Kang et al. 2017), such as Hypoxia-inducible factor (HIF) (Guan et al. 2014) and Erythropoietin (EPO) (Xu et al. 2016) associated with hypoxia response, ATPase Family AAA Domain Containing 2 (ATAD2) (Tong, Fei, et al. 2017) and cAMP-dependent protein kinase catalytic subunit alpha (PRKACA) that involved into energy metabolism (Tong, Fei, et al. 2017). The main focus of previous studies in schizothoracine fishes are still on hypoxia and cold response. Notably, an increasing number of lakes in the Tibetan Plateau have been existing or towards alkaline due to

the global climate changes and human activities (Zheng 1997). Thus, the increasing alkalization

of fresh water has been the potential challenge to schizothoracine fishes. Among the

schizothoracine fishes, *Gymnocypris przewalskii przewalskii* and *Gymnocypris przewalskii*

kelukehuensis are the only two species inhabited extremely alkaline environment (Wu 1992).

Unlike other broadly distributed schizothoracine fishes, such as *Gymnocypris eckloni*,

Schizopygopsis pylzovi and *Platypharodon extremus* that inhabit in the Yellow river basin (Cao

et al. 1981; Wu 1992; Qi et al. 2012), *G. p przewalskii* only inhabits in saline and alkaline lake.

As the largest salt lake in China, Lake Qinghai (FIG. 1) is highly saline (up to 13‰) and alkaline

(up to pH 9.4) water environment, a typical salt lake with unusually high sodium, potassium and

magnesium concentration (Zheng 1997; Zhu & Wu 1975). In addition, *G. p. kelukehuensis* only

inhabits in a soda lake located at the Tsaidam Basin in the northeastern Tibetan Plateau. Lake

Keluke (FIG. 1) is also a soda lake with low salinity of 0.79‰ and high pH value up to 9.3

(Zheng 1997). Both schizothoracine fish species had developed unique physiological or genetic

features to tolerate such harsh living conditions (Tong, Fei, et al. 2017). Therefore, this provides

an exceptional model to investigate the genetic mechanisms underlying alkaline adaptation, and

may provide novel insights to fully understand the mechanism of highland adaptation in fish as

complement.

Unlike highland alkaline tolerant fish, a huge amount of studies had explored the mechanisms of

high saline and high alkaline tolerance in lowland fish species, such as killifish (e.g. *Fundulus*

heteroclitus) (Wood et al. 2010;Brennan et al. 2018; Burnett et al. 2007), tilapia (e.g.

Oreochromis niloticus) (Zhao et al. 2020; Wood et al. 1994), and salmonids (e.g. *Salmo salar*)

(Levings 2016; Lien et al. 2016). These studies had provided insights in physiology of acid-base

balance in fishes response to high salinity or high pH environments, and suggested key genes,

such as ion transport associated genes under selection during the adaptation (Lien et al. 2016).

In this study, we generated and assembled the transcriptomes of two alkaline tolerant

schizothoracine fish species, *G. p. przewalskii* and *G. p. kelukehuensis* inhabited in high pH

environment in the northeastern Tibetan Plateau (FIG. 1). We performed a comparative

genomics study together with recently sequenced schizothoracine fish transcriptomes and other

lowland fish genomes (FIG. 2A & 2B), and sought to identify consistent genomic signature

associated with alkaline adaptation in highland and lowland fishes. Specifically, we focused our

comparisons on testing whether alkaline adaptation in highland and lowland alkaline tolerant

fishes is associated with the following signatures of molecular evolution: (1) consistent patterns

of molecular evolution in protein-coding genes across the phylogeny; (2) consistent shifts in

evolutionary rates for specific genes; and (3) consistent signals of positive selection in particular

genes.

Materials and Methods

Materials and Methods

Sample collection

We collected eight adult *G. p. przewalskii* (FIG. 1) individuals (four males and four females, 172

± 0.7 g) from Lake Qinghai and eight adult *G. p. kelukehuensis* (FIG. 1) individuals (four males

and four females, 139 ± 0.3 g) from Lake Keluke using gill nets. All the fish samples were

dissected after anesthesia with MS-222 (Solarbio, Beijing, China). All individuals were classified

based on the gender and dissected after anesthesia with MS-222 (Solarbio, Beijing, China).

Tissues from gill, kidney, brain, heart and liver from each individual were collected and

immediately stored in liquid nitrogen at -80 °C. All the animal experiments were approved by the

Animal Care and Use Committees of the Northwest Institute of Plateau Biology, Chinese

Academy of Sciences (NWIPB-CY-010).

Transcriptomics

Total RNA of each tissue sample was extracted using TRIzol reagent (Invitrogen, CA, USA) in

accordance with manufacturer's instructions, and detected for quality and quantity of RNAs with

Nanodrop 1000 (NanoDrop Technologies, DE, USA) and Agilent Bioanalyzer 2100 (Agilent

Technologies, CA, USA). Equal amount of RNA from eight individual of five tissue was pooled to

construct transcriptome library as previously described (Tong, Tian, et al. 2017; Tong, Fei, et al.

2017), and sequenced with an Illumina NovaSeq 6000 yielding 150-bp paired-end reads (FIG.2).

Sequencing reads were checked for quality using FastQC

(https://www.bioinformatics.babraham.ac.uk/projects/fastqc/). Sequencing adapters and reads

with a quality score < 20 were trimmed with Trimmomatic (Bolger et al. 2014). We built a *de*

novo transcriptome assembly based on clean reads using Trinity v2.6.5 (Grabherr et al. 2011)

with default parameters. Next, we removed the redundant transcripts using CD-HIT (Fu et al.

2012) with the threshold of 0.90 and extracted the longest transcript as putative genes. We

predicted the open reading frame of each putative genes using TransDecoder

(https://github.com/TransDecoder/TransDecoder) (FIG.2).

Additional data retrieval

We downloaded six alkaline intolerant schizothoracine fish transcriptomes (Zhou et al. 2020)

including *Oxygymnocypris stewartii*, *Schizopygopsis younghusbandi*, *Gymnocypris namensis*,

Platypharodon extremus, *Schizopygopsis pylzovi* and *Gymnocypris eckloni* from NCBI SRA

database (https://www.ncbi.nlm.nih.gov/sra) (FIG. 3B, supplementary table S1), and performed

assembly following above pipeline. In addition, we downloaded the genomes of four alkaline

intolerant fish species of *Danio rerio*, *Ctenopharyngodon idellus*, *Cyprinus carpio*, and *Carassius*

auratus, and three alkaline tolerant fish species of *Fundulus heteroclitus*, *Oreochromis niloticus*

and *Salmo salar* (FIG. 3A, supplementary table S1).

Species phylogeny and gene orthology

We obtained the phylogenetic tree of 15 fish species by pruning the Fish Tree of Life

(https://fishtreeoflife.org/) using R package, phangorn (Schliep 2011). To obtain phylogeny-

based orthology relationships between different fish taxa, we included all the predicted

proteomes of seven lowland fish genomes, and translated nucleotide sequences of protein-

coding genes from eight schizothoracine fish transcriptome assemblies into amino acid

sequences, and pooled these datasets as input for an orthology inference tool, OMA (Altenhoff

et al. 2018). In this way, we identified one-to-one, one-to-many, and many-to-many orthologs

among these 15 fish species. For further comparison, we restricted our analysis to 1:1 orthologs,

that is the geneset for which only one gene from each species representing the orthology. In

addition, we extracted the shared orthologs among all fish taxa. At last, gene ontology (GO)

terms were assigned to each ortholog using Trinotate (https://trinotate.github.io/) (FIG.2).

Pattern of molecular evolution in shared orthologs

To determine whether highland fish and lowland fish showing consistent patterns of molecular

evolution in the set of alkaline tolerant species branches across the phylogeny, we

characterized the rates of non-synonymous to synonymous rate (dN/dS) in each shared

ortholog. For this, we performed the protein sequence alignment using MUSCLE v3.8.31

(https://www.ebi.ac.uk/Tools/msa/muscle). We prepared the codon alignments of shared

orthologs, that derived from protein alignments and the corresponding DNA sequences using

PAL2NAL v.14 (Suyama et al. 2006). Then, we executed the filtration for shared ortholog

alignments with length of at least 50 codons.

We took advantage of HyPHY pipeline (Kosakovsky Pond, Poon, et al. 2020) to test the

hypotheses by comparing selective pressures (dN/dS) between *a prori* defined alkaline tolerant

fish species branches (focal foreground branch) and alkaline intolerant species fish branches

(background branch) in the specified fish phylogeny at ortholog-wide scale. Before that, a

common approach to test the hypothesis is to perform separate analyses on subsets of

sequences, and compare the parameter estimated in a *post hoc* fashion, that is one branch to

the rest of branches in a phylogenetic tree like which we previously described (Tong, Tian, et al.

2017; Tong, Fei, et al. 2017; Tong et al. 2020). While, such approach is statistically suboptimal,

and not always applicable (Kosakovsky Pond, Wisotsky, et al. 2020). In this way, we estimated

two discrete categories of dN/dS for alkaline tolerant and alkaline intolerant species under the

MG94 rate matrices combined with HKY85 model using HyPHY (Kosakovsky Pond, Poon, et al.

2020), and compared nested models with constrained relationships among them. We detected

the shift of dN/dS ratios between alkaline tolerant and alkaline intolerant (alternative model, H1),

relative to the null model that assuming all lowland fish taxa have the same dN/dS ratio (H0).

We constructed the log-likelihood ratio score for each ortholog (ΔlnL) as: ΔlnL = 2(lnLH$_1$-lnLH$_0$)

and employed the likelihood ratio test. Additionally, we applied a correction for multiple testing to

a false discovery rate (FDR) < 0.20 using R package, q-value (http://github.com/jdstorey/qvalue).

In this way, we repeated this analysis with shared ortholog datasets of highland taxa.

Analysis of accelerated evolution

To determine if consistent shift in evolutionary rates (e.g. acceleration) in particular genes within

alkaline tolerant species across the phylogeny, we defined rapidly evolving genes with

significantly higher dN/dS ratio for alkaline tolerant species than alkaline intolerant species (*P* <

0.05, likelihood ratio test, FDR < 0.2). In this way, we identified the rapidly evolving genes in

alkaline tolerant species within highland and lowland based on earlier estimated dataset

including two discrete dN/dS ratios for each shared ortholog, respectively. In addition, these

sets of rapidly evolving genes were examined for GO enrichment relative to the full set of all

shared orthologs using R package, topGO

(https://bioconductor.org/packages/release/bioc/html/topGO.html). We finally visualized all

significantly enriched GO terms remaining after the REVIGO (http://revigo.irb.hr/) similarity filter.

Analysis of positive selection

To further determine whether consistent signals of positive selection in a set of branches

representing alkaline tolerant species across the phylogeny in specific genes, we used three

complementary branch-site models to identify the positively selected genes (FIG.2). Specifically,

we first used the Branch-Site Unrestricted Statistical Test for Episodic Diversification (BUSTED)

model (Murrell et al. 2015) to test for positive selection signal in a gene at any site on focal

branches. In this model, it defines a positively selected gene (LRT P < 0.05, FDR < 0.2) with at

least one site under positive selection on at least one focal foreground branch, while does not

specify the exact branch with positively selected site, may include false positive. Then, we used

an adaptive Branch-Site Random Effects Likelihood (aBSREL) modelaBSREL (Smith et al.

2015) to test if positive selection has occurred on a proportion of branches (i.e. number of focal

foreground branches) at specific genes (Holm-Bonferroni corrected P < 0.05). This allowed us

to filter the false positive cases without positive selection signal on specific focal foreground

branches (i.e. alkaline tolerant fish) out of the gene sets determined by BUSTED (FIG.2).We

defined the positively selected genes that required the shift across all focal foreground branches.

Since the branch-site models can cause false positive in case of multinucleotide mutations

(MNMs, Venkat et al. 2018), we performed more conservative branch-site (BS) model test

covering MNM situation (BS + MNM) (github.com/JoeThorntonLab/MNM_SelectionTests). In

this model, an additional parameter δ represents the relative instantaneous rate of double

mutations compared to that of single mutations. We ran null models and alternative models in

BS + MNM and conducted LRTs to evaluate significance (LRT P < 0.05). In this way, we further

filter false positive cases out of the previously positively selected gene sets that determined by

BUSTED and aBSREL (FIG.2). In this way, we took advantage of these three models to finalize

the positively selected genes (i.e. gene with sites under positive selection) in alkaline tolerant

species within highland fish taxa and lowland fish taxa, respectively (FIG.2). We finally

performed GO enrichment analysis using topGO and REVIGO as earlier described.

Intersection of rapidly evolving and positively selected gene repertoire

To determine if there is both accelerated evolution and positive selection for specific genes, we

did the overlapping between rapidly evolving genes (REGs) and positively selected genes

(REGs) for highland and lowland alkaline tolerant fish species. In addition, we performed GO

enrichment analysis with topGO and REVIGO for the overlapping genes.

Results

Following *de novo* assembly and annotation, each of the eight schizothoracine fish

transcriptome assemblies had an average of 39,921 transcripts representing the complete or

partial protein-coding regions of genes (supplementary table S2). We further identified a total of

7,309 one-to-one orthologs shared by 15 fish species (i.e. range from 2 to 15 fish species), and

6,241 shared 1:1 orthologs including all 15 species (supplementary table S3).

Consistent pattern of molecular evolution

We estimated two categories of dN/dS ratios for 5,748 shared 1:1 orthologs (after restricting to

1:1 orthologs with at least 50 codons) in lowland fish taxa and highland fish taxa, separately. We

found consistent patterns of molecular evolution showing significant (LRT, $P < 0.05$, FDR < 0.2)

acceleration (increased dN/dS) or deceleration (decreased dN/dS) in a set of terminal branches

of alkaline tolerant species relative to the set of terminal branches of alkaline intolerant species

in large numbers of shared orthologs within lowland fish taxa (n = 952) and highland fish taxa (n

= 162) (FIG. 3C, supplementary table S4).

Consistent signature of accelerated evolution

Building on earlier dataset of two discrete categories of dN/dS ratios separately representing

branches of alkaline tolerant and alkaline intolerant species across the phylogeny, we focused

on genes with significantly higher dN/dS (LRT, $P < 0.05$, FDR < 0.2) in alkaline tolerant species,

that is rapidly evolving gene (REGs) repertoire in alkaline tolerant species within highland and

lowland fish taxa. We identified 110 REGs in highland fishes and 470 REGs in lowland fishes

(FIG. 4A). Out of 11 overlapping REGs, we found a set of ion transport and transmembrane

functions associated genes, such as sodium-dependent phosphate transport protein 2A

(SLC34A1) and sodium-dependent phosphate transporter 1-B (SLC20A1b) (FIG, 4B,

supplementary table S5). Besides, we identified overlapping REGs related to energy

metabolism process, such Ectonucleotide pyrophosphatase/phosphodiesterase family member

1 (ENPP1) (FIG. 4B). Given that different REGs identified in either highland or lowland alkaline

tolerant fish species, we found a number of ion transport and ATP synthesis related genes in

REGs repertoire of highland fish (FIG. 4B, supplementary table S5), such as solute carrier

family 35 member F4 (SLC35F4), ATP-sensitive inward rectifier potassium channel 15

(KCNJ15), sodium-dependent serotonin transporter (SLC6A4), and NADP-dependent malic

enzyme (ME1). Similarly, in lowland fish, we also identified genes like solute carrier family 45

member 3 (SLC45A3), Potassium Inwardly Rectifying Channel Subfamily J Member 5 (KCNJ5) ,

and ATP synthase F0 complex subunit B1 (ATP5PB) showing rapidly evolving in alkaline

tolerant species (FIG. 4B, supplementary table S5).

Further, we did GO enrichment analysis for both REGs datasets, showing 112 significantly

enriched GO terms (biological process) in highland fish ($P < 0.05$, fisher's exact test) and 189

significantly enriched GO terms in lowland fish. After the filtration by semantic similarity of GO

terms, we found a set of enriched dominant GO terms in highland fish were related to ion

transport and transmembrane functions (FIG. 4C, supplementary table S6), such as ion

transport (GO:0006811), regulation of calcium ion transport (GO:0051924), anion transport

(GO:0006820). In addition, in lowland fish, we observed a set of enriched dominant GO terms

associated with ion transport function (FIG. 4D, supplementary table S6), such as phosphate ion

transport (GO:0006817) and response to salt stress (GO:0009651). Finally, we found 5

overlapping GO terms related to ion transport and metabolism, including glycogen biosynthetic

process (GO:0005978), phosphate ion transport (GO:0006817), inorganic anion transport

(GO:0015698), negative regulation of ion transport (GO:0043271), regulation of glycogen

metabolic process (GO:0070873) and anion transport (GO:0006820) (supplementary table S6).

Collectively, this finding suggested the consistent signature of accelerated evolution in alkaline

tolerant species within highland taxa and lowland taxa.

Consistent signature of positive selection

After two rounds of filtration with the use of BUSTED, aBSREL and BS+MNM models, we

identified 162 positively selected genes in highland alkaline tolerant fish (FIG. 5A,

supplementary table S7), and 156 PSGs in lowland alkaline tolerant fish species (FIG. 5A,

supplementary table S7). Out of 7 overlapping PSGs, we found these genes were mainly

related to apoptosis (cell death), ion transport and immune response, such as vitamin K-

dependent protein C precursor (PROC), E3 ubiquitin-protein ligase SH3RF1 (SH3RF1), 14-3-3

protein beta/alpha-A (YWHABA) and cadherin-related family member 2 (Cdhr2). Besides, we

found PSGs in highland alkaline tolerant species were also mainly involved in similar functional

categories as overlapping PSGs (FIG. 5B, supplementary table S7), such as transmembrane

protein 268 (TMEM268), transmembrane protein 266 (TMEM266), solute carrier family 35

member F4 (SLC35F4), solute carrier family 7, member 3 (SLC35F4) and solute carrier family 7,

member 3 (SLC7A3) involved in ion transport or transmembrane functions, probable

phospholipid-transporting ATPase VD (ATP10D) involved in energy metabolism, apoptosis-

inducing factor 1 (AIFM1) involved in apoptosis, interleukin-2 receptor subunit beta (IL2RB)

related to immune response. Similarly, in lowland alkaline tolerant fish species, these PSGs

were involved in four main categories (FIG. 5B, supplementary table S7), such as solute carrier

family 2 member 15b (SLC2A15b and potassium voltage-gated channel subfamily E member 4

(KCNE4) associated with ion transport function, phosphoinositide 3-kinase regulatory subunit 4

(PIK3R4) and lysosomal-associated membrane protein 1 (LAMP1) associated with apoptosis,

CD22 antigen (CD22) and immunoglobulin-like domain containing receptor 1b precursor (ILDR1)

involved in immune response, and plasma membrane calcium-transporting ATPase 3 (ATP2B3)

associated with energy metabolism.

Further, GO enrichment results showed that a number of significantly enriched GO terms ($P <$

0.05, fisher's exact test) are related to immune response, apoptosis, protein metabolisms for

PGSs in highland alkaline tolerant species, such as inflammatory response (GO:0006954),

natural killer cell activation (GO:0030101), cell killing (GO:0001906), muscle cell apoptotic

process (GO:0010656), regulation of innate immune response (GO:0045088), immune

response (GO:0006955), ion transport (GO:0006811), sphingolipid metabolic process

(GO:0006665) and thyroid hormone metabolic process (GO:0042403) (FIG. 5C, supplementary

table S8). Similarly, in lowland alkaline tolerant species, the significantly enriched GO terms

were mainly related to four categories as well, such as T cell mediated cytotoxicity

(GO:0001913) related to immune response, negative regulation of transport (GO:0051051) and
negative regulation of ion transport (GO:0043271) related to transport function, macroautophagy
(GO:0016236) related to apoptosis (cell death), regulation of phospholipid metabolic process
(GO:1903725) and sulfur amino acid metabolism (GO:0000096) (FIG. 5D, supplementary table
S8). Collectively, this finding suggested the consistent signature of positive selection in alkaline
tolerant species within highland taxa and lowland taxa.

Genes with evidence for both accelerated evolution and positive selection

We sought to find the intersection of REGs and PSGs repertoires, we found 29 overlapping
genes in highland alkaline tolerant fish (FIG.6A), mainly related to ion transport, apoptosis (cell
death), and energy metabolism, such as SLC35F4, SLC7A3, SLC6A4, TMEM266, CLDN15 and
ALDH16A1 and ATP6V1C1 (supplementary table S9). GO enrichment also suggested that
overlapping genes were mainly enriched in three main functional categories, such ion transport
(GO:0006811), anion transport (GO:0006820), regulation of calcium ion import (GO:0090279),
alditol phosphate metabolic process (GO:0052646), oxoacid metabolic process (GO:0043436)
and regulation of muscle cell apoptotic process (GO:0010660) (FIG.6B, supplementary table
S10). Similarly, we observed 26 shared REG/PSG genes in lowland alkaline tolerant fish (FIG.
6C, supplementary table S9). The additional GO enrichment result showed that the overlapping
genes mainly enriched into transport and metabolic process, such as negative regulation of ion
transport (GO:0043271), negative regulation of potassium ion transmembrane transporter
activity (GO:1901017) and GTP metabolic process (GO:0046039) (FIG. 6D, supplementary
table S10). Thus, this finding emphasized the consistent signatures of accelerated evolution and
positive selection in alkaline tolerant species.

Discussion

Our results support above three hypotheses that alkaline tolerant species shared the consistent
patterns of molecular evolution in protein-coding genes (dN/dS) and consistent signatures of
accelerated evolution (rapidly evolving gene) and positive selection (positively selected gene) in
highland and lowland fish. Specifically, these signatures include: genes experienced consistent
acceleration in evolutionary rates (increased dN/dS) in alkaline tolerant species, which are
mainly involved in ion transport, transmembrane and energy metabolism functions; genes

showing consistent signals of positive selection in alkaline tolerant species within highland and

lowland fish taxa, these are mainly associated with ion transport/transmembrane, apoptosis (cell

death), immune response and energy metabolism processes. Altogether, this study provides

insights in understanding the common role of adaptive molecular evolution in fish adaptation to

alkaline environment, as well as adaptation to extreme environment at high-altitude.

Acid-base balance and osmoregulation

In freshwater fish, Na^+ and Cl^- are actively taken up across the gill epithelium to counter the

passive loss of osmolytes to the more dilute environment. After transition to saline or alkaline

water, fish must increase its rate to balance the osmotic loss of water to the more solute-

concentrated environment and actively excrete Na^+ and Cl^- from the gill to maintain ionic and

osmoregulation (Marshall 2005). Thus, alkaline tolerant species requires enhanced

physiological abilities including acid-base balance and osmoregulation to respond the elevation

in alkalinity or salinity of freshwater (Evans et al. 2005; Marshall 2005). Extremely alkaline

environment may accelerate the evolution of genes associated with osmoregulation in these

species survived in such harsh environment (Xu et al. 2017; Tong, Fei, et al. 2017). In this study,

we identified a set of genes associated with ion transport and transmembrane that tended to

evolve rapidly in alkaline tolerant species than their alkaline intolerant relatives (FIG. 4). This

result echoed our previous finding in *G. przewalskii* compared with other teleost fishes (alkaline

intolerance) (Tong, Fei, et al. 2017), also was in line with the rapidly evolving gene repertoire of

a wild fish, Amur ide (*Leuciscus waleckii*) that survived in extremely alkaline environment (Xu et

al. 2017). Interestingly, osmoregulation related genes including three solute carrier (SLC) genes

and one transient receptor potential cation channel (TRPV) gene exhibited consistent signature

of accelerated evolution in both highland and lowland alkaline tolerant fishes. *SLC* genes

encode transmembrane transporters for inorganic ions, amino acids, neurotransmitters, sugars,

purines and fatty acids, and other solute substrates (Dorwart et al. 2008). Recent evidences

indicted that adaptive evolution of *SLC* genes contributed to fish adaptation to high salinity and

high pH environment (Wang & Guo 2019; Tong, Fei, et al. 2017; Xu et al. 2017). Besides, we

also identified numbers of ion transport and transmembrane genes under positive selection in

alkaline tolerant fish species, such as potassium voltage-gated channel (*KCN*) genes. In killifish,

an excellent model to study extreme environment adaptation, recent genome-wide studies also

found *KCN* genes have been implicated in freshwater adaptation (transition from marine to

freshwater environment) (Brennan et al. 2018).) Besides a set of same rapidly evolving and

positively selected genes identified in both highland and lowland alkaline tolerant fish species,

we found a large number of different genes under selection but involved in similar functions,

such as anion transport (GO:0006820) and phosphate ion transport (GO:0006817), indicating

the common role of adaptive molecular evolution during alkaline adaptation in fish.

Apoptosis and cell death

Extremely alkaline stress may cause extensive damage to fish, such as inducing cell apoptosis

(Monteiro et al. 2009; Zhao et al. 2016, 2020). However, several fish species can survive in this

harsh environment, such as schizothoracine (*Gymnocypris przewalskii*) (Tong, Fei, et al. 2017;

Tong & Li 2020), Magadi tilapia (*Alcolapia grahami*) (Wilkie & Wood 1996), and Amur ide

(*Leuciscus waleckii)* (Xu et al. 2017). This may also pose a barrier for alkaline tolerant fish as

compared with alkaline intolerant fish. Significantly, positively selected *AIFM1*, *SH3RF1*,

YWHABA, *PIK3R4* and *LAMP1* of alkaline tolerant species relative to alkaline intolerant species

have enrichment in a set of apoptosis pathways, such regulation of apoptotic signaling pathway

(GO:1902253) and macroautophagy (GO:0016236)(FIG. 5). Apoptosis is a form of programmed

cell death that occurs in multicellular organisms, it plays a significant role in the biochemical

events lead to characteristic cell changes (morphology) and death (Green 2011). Few direct

evidence suggested the roles of these candidate genes in response to extreme alkaline stress

in fish, but numerous studies had defined their functions in tolerance to harsh environments.

AIFM1 is a ubiquitous mitochondrial oxidoreductase involved in apoptosis, involved in sea

bream (*Sparus aurata*) response to acute environmental stress (Bermejo-Nogales et al. 2014).

YWHABA (14-3-3 protein beta/alpha-A) is an important gene showing the ability to bind a

multitude of functionally diverse signaling proteins, such as transmembrane receptors (Fu et al.

2000), that involved in spotted sea bass (*Lateolabrax maculatus*) tolerance to saline stress

(Zhang et al. 2019). In addition, *LAMP1* plays an important role in lysosome biogenesis and

autophagy (Eskelinen 2006), which involved in common carp (*Cyprinus carpio*) response to

hydrogen peroxide environment. Collectively, the presence of apoptosis related genes under

positive selection may contribute to the alkaline adaptation of fish, how these adaptive

molecular changes affect the ability programmed cell death remains unknown.

Immune response

Extreme environments (including high pH) impact on the physiology of animals in a wide variety

of ways. Recent advances in the understanding of environmental impacts were identified in

relation to specific areas of immune function, such as increase in pH resulted in a general

increase in immune function (Bowden 2008; Sridhar et al. 2020). Intriguingly, we identified

different immune genes under positive selectin in alkaline tolerant species within highland

(IL2RB, TLR8, IF144) and lowland (ILDR1 and CD44), but they all involved similar immune

functions, such as inflammatory response (GO:0006954), natural killer cell activation

(GO:0030101), and T cell mediated cytotoxicity (GO:0001913). In another word, we found

different genes but conserved pathways may underlie fish adaptation to alkaline environment.

IF144, IL2RB and ILDR1 are important components of toll-like receptor (TLR) signaling pathway

that play key roles in the innate immune system (Rebl et al. 2010). For instance, our previous

comparative studies in alkaline tolerant fish, *G. przewalskii* identified key genes involved in TLR

pathway under selection (Tong, Fei, et al. 2017; Tong et al. 2015). In Nile tilapia (*Oreochromis*

niloticus), another alkaline tolerant species and stands as ideal model to study extreme

environment adaptation, a most recent comparative study identified numbers of immune genes

involved in natural killer cell mediated cytotoxicity and NF-kappa B signaling pathway in

response to alkalinity stress (up to pH = 8.9) (Zhao et al. 2020), and echoes our present results.

Thus, it is possible that adaptive evolution changes (e.g. positive selection) acting on innate

immune genes in alkaline tolerant species contribute to their adaptation to extremely alkaline

environment.

Energy metabolism

It is not surprising that we identified a set of genes under either accelerated evolution or positive

selection, that enriched in diverse metabolisms, such as glucose metabolism, phosphate

metabolism, sulfur amino acid metabolism and protein metabolism in alkaline tolerant species

compared with their alkaline intolerant relatives. In general, metabolism processes were

involved in fish response to varies environmental stresses (including alkaline stress) by a huge

amount of research (Wood 1991). A most recent study highlights the genes involved in

conserved mitochondrial pathways under selection in adaptation to extreme environment

(Greenway et al. 2020). In addition to our previous studies in highland alkaline tolerant fish, we

found a number of genes associated with energy metabolism processes under selection as well,

such as mitochondrial function and protein metabolism (Tong, Fei, et al. 2017). Moreover,

increasing studies on alkaline tolerant fish species (e.g. killifish) pointed out the roles of varied

metabolisms in extreme environment (e.g. high pH) adaptation (Scott et al. 2019; Zhao et al.
2020; Wang & Guo 2019; Xu et al. 2017). Altogether, our analysis infer that the adaptive
molecular evolution of metabolism associated genes may be indispensable and common
features to alkaline adaptation as well as extreme environment adaptation in fish.

Conclusion

In summary, this comparative genomics study of 15 fish species suggests the common role of
alkaline adaptation in fish, regardless of highland or lowland background environments. Our
results also highlight that the adaptive evolution of protein-coding genes are likely to play a
crucial role in fish response to extreme environment, such as extremely high PH. Notably, this
study provides putative genomic signatures of shift in selection and alkaline adaptation in
several alkaline tolerant fish species, further study should include large scale of omics data of
more alkaline tolerant fish species and their intolerant relatives as multiple pairs to demonstrate
the genetic basis of alkaline adaptation in fish at genome-wide scale.

Author Contributions

473 C.T. and K.Z. conceived this project. C.T. designed the project. C.T., Y.T. and K.Z. collected the
474 samples. C.T. and M.L. performed the comparative genomics analyses. C.T. wrote the paper.
475 All authors read and approved the final manuscript.

Acknowledgments

478 We would like to thank four anonymous reviewers for helpful comments on this manuscript.

Funding

481 This work was supported by grants from the National Natural Science Foundation of China
482 (31870365), Joint Grant from Chinese Academy of Sciences -People's Government of Qinghai
483 Province on Sanjiangyuan National Park (LHZX-2020-01), China Biodiversity Observation and
484 Research Network, Sino BON-Inland Water Fish Diversity Observation Network.

Conflicts of Interest

The authors declare no conflict of interest.

Data Availability Statement

The Illumina sequencing reads have been deposited at NCBI Sequence Read Archive under the NCBI BioProject accession PRJNA684806.

Literature Cited

Altenhoff AM et al. 2018. The OMA orthology database in 2018: retrieving evolutionary relationships among all domains of life through richer web and programmatic interfaces. Nucleic Acids Res. 46:D477–D485.

An Z, John K, Warren P, Stephen P. 2001. Evolution of Asian monsoons and phased uplift of the Himalaya–Tibetan plateau since Late Miocene times. Nature. 411:62–66.

Bermejo-Nogales A et al. 2014. Metabolic and transcriptional responses of gilthead sea bream (Sparus aurata L.) to environmental stress: new insights in fish mitochondrial phenotyping. Gen. Comp. Endocrinol. 205:305–315.

Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. Bioinformatics. 30:2114–2120.

Bowden TJ. 2008. Modulation of the immune system of fish by their environment. Fish Shellfish Immunol. 25:373–383.

Brennan RS., et al. 2018. Integrative population and physiological genomics reveals mechanisms of adaptation in killifish. Mol. Biol. Evol. 35.11: 2639-2653.

Burnett KG et al. 2007. Fundulus as the premier teleost model in environmental biology: opportunities for new insights using genomics Comp. Biochem. Physiol. Part D Genomics Proteomics. 2.4: 257-286.

Cao W, Chen Y, Wu Y, Zhu S, Others. 1981. Origin and evolution of schizothoracine fishes in
relation to the upheaval of the Qinghai-Tibetan Plateau. Science Press: Beijng.

Dorwart MR, Shcheynikov N, Yang D, Muallem S. 2008. The solute carrier 26 family of proteins
in epithelial ion transport. Physiology . 23:104–114.

Eskelinen E-L. 2006. Roles of LAMP-1 and LAMP-2 in lysosome biogenesis and autophagy.
Mol. Aspects Med. 27:495–502.

Evans DH, Piermarini PM, Choe KP. 2005. The multifunctional fish gill: dominant site of gas
exchange, osmoregulation, acid-base regulation, and excretion of nitrogenous waste. Physiol.
Rev. 85:97–177.

Favre A et al. 2015. The role of the uplift of the Qinghai-Tibetan Plateau for the evolution of
Tibetan biotas. Biol. Rev. Camb. Philos. Soc. 90:236–253.

Fu H, Subramanian RR, Masters SC. 2000. 14-3-3 proteins: structure, function, and regulation.
Annu. Rev. Pharmacol. Toxicol. 40:617–647.

Fu L, Niu B, Zhu Z, Wu S, Li W. 2012. CD-HIT: accelerated for clustering the next-generation
sequencing data. Bioinformatics. 28:3150–3152.

Grabherr MG et al. 2011. Full-length transcriptome assembly from RNA-Seq data without a
reference genome. Nat. Biotechnol. 29:644–652.

Green DR. 2011. Means to an end: apoptosis and other cell death mechanisms. Cold Spring
Harbor Laboratory Press.

Greenway R et al. 2020. Convergent evolution of conserved mitochondrial pathways underlies
repeated adaptation to extreme environments. Proc. Natl. Acad. Sci. U. S. A. 117:21822–21822.

Guan L, Chi W, Xiao W, Chen L, He S. 2014. Analysis of hypoxia-inducible factor alpha
polyploidization reveals adaptation to Tibetan Plateau in the evolution of schizothoracine fish.
BMC Evol. Biol. 14:192.

Kang J, Ma X, He S. 2017. Evidence of high-altitude adaptation in the glyptosternoid fish,
Creteuchiloglanis macropterus from the Nujiang River obtained through transcriptome analysis.
BMC Evol. Biol. 17:229.

538 Kosakovsky Pond SL, Poon AFY, et al. 2020. HyPhy 2.5—A Customizable Platform for
539 Evolutionary Hypothesis Testing Using Phylogenies. Mol. Biol. Evol. 37:295–299.

540 Kosakovsky Pond SL, Wisotsky SR, Escalante A, Magalis BR, Weaver S. 2020. Contrast-FEL--
541 a test for differences in selective pressures at individual sites among clades and sets of
542 branches. Mol. Biol. Evol. 38.3: 1184-1198. .

543 Levings CD. 2016. Ecology of Salmonids in Estuaries around the World: Adaptations, Habitats,
544 and Conservation. UBC Press.

545 Lien S et al. 2016. The Atlantic salmon genome provides insights into rediploidization. Nature.
546 533:200–205.

547 Li J, Fang X. 1999. Uplift of the Tibetan Plateau and environmental changes. Chinese science
548 bulletin. 44:2117–2124.

549 Liu H-P et al. 2019. The sequence and de novo assembly of Oxygymnocypris stewartii genome.
550 Scientific Data. 6:190009.

551 Marshall WS. 2005. Ion transport, osmoregulation, and acid-base balance. The physiology of
552 fishes. CRC Press.

553 Monteiro SM, dos Santos NMS, Calejo M, Fontainhas-Fernandes A, Sousa M. 2009. Copper
554 toxicity in gills of the teleost fish, Oreochromis niloticus: effects in apoptosis induction and cell
555 proliferation. Aquat. Toxicol. 94:219–228.

556 Murrell B et al. 2015. Gene-wide identification of episodic selection. Mol. Biol. Evol. 32:1365–
557 1371.

558 Orr HA. 2005. The genetic theory of adaptation: a brief history. Nat. Rev. Genet. 6:119–127.

559 Qi D et al. 2012. Convergent, parallel and correlated evolution of trophic morphologies in the
560 subfamily schizothoracinae from the Qinghai-Tibetan plateau. PLoS One. 7:e34070.

561 Qu Y et al. 2013. Ground tit genome reveals avian adaptation to living at high altitudes in the
562 Tibetan plateau. Nat. Commun. 4:2071.

563 Rebl A, Goldammer T, Seyfert H-M. 2010. Toll-like receptor signaling in bony fish. Vet.
564 Immunol. Immunopathol. 134:139–150.

Scheinfeldt LB, Tishkoff SA. 2010. Living the high life: high-altitude adaptation. Genome Biol.
11:133.

Schliep KP. 2011. phangorn: phylogenetic analysis in R. Bioinformatics. 27:592–593.Scott WC
et al. 2019. Influence of salinity and pH on bioconcentration of ionizable pharmaceuticals by the
gulf killifish, Fundulus grandis. Chemosphere. 229:434–442.

Smith MD et al, 2015. Less is more: an adaptive branch-site random effects model for efficient
detection of episodic diversifying selection. Molecular biology and evolution 32.5: 1342-1353.

Sridhar A et al. 2020. Activity profile of innate immune-related enzymes and bactericidal of
freshwater fish epidermal mucus extract at different pH. Environ. Sci. Pollut. Res. Int. doi:
10.1007/s11356-020-11173-5.

Suyama M, Torrents D, Bork P. 2006. PAL2NAL: robust conversion of protein sequence
alignments into the corresponding codon alignments. Nucleic Acids Res. 34:W609–12.

Tong C et al. 2015. Transcriptome-wide identification, molecular evolution and expression
analysis of Toll-like receptor family in a Tibet fish, Gymnocypris przewalskii. Fish Shellfish
Immunol. 46:334–345.

Tong C, Fei T, Zhang C, Zhao K. 2017. Comprehensive transcriptomic analysis of Tibetan
Schizothoracinae fish Gymnocypris przewalskii reveals how it adapts to a high altitude aquatic
life. BMC evolutionary biology. 17:74.

Tong C, Li M. 2020. Genomic signature of accelerated evolution in a saline-alkaline lake-
dwelling Schizothoracine fish. Int. J. Biol. Macromol. 149:341–347.

Tong C, Najm GM, Pinter-Wollman N, Pruitt JN, Linksvayer TA. 2020. Comparative Genomics
Identifies Putative Signatures of Sociality in Spiders. Genome Biol. Evol. 12:122–133.

Tong C, Tian F, Zhao K. 2017. Genomic signature of highland adaptation in fish: a case study in
Tibetan Schizothoracinae species. BMC Genomics. 18:948.

Venkat A, Matthew H, Joseph T. 2018. Multinucleotide mutations cause false inferences of
lineage-specific positive selection. Nature ecology & evolution 2.8: 1280-1288.

Wang Y et al. 2015. Evidence for Adaptation to the Tibetan Plateau Inferred from Tibetan Loach
Transcriptomes. Genome Biol. Evol. 7:2970–2982.

Wang Y, Guo B. 2019. Adaption to extreme environments: a perspective from fish genomics.
Rev. Fish Biol. Fish. 29:735–747.

Wen LY. 2014. Uplift of the tibetan plateau influenced the morphological evolution of animals. J.
Agric. Sci. 6:244.

Wilkie MP, Wood CM. 1996. The adaptations of fish to extremely alkaline environments. Comp.
Biochem. Physiol. B Biochem. Mol. Biol. 113:665–673.

Witt KE, Huerta-Sánchez E. 2019. Convergent evolution in human and domesticate adaptation
to high-altitude environments. Philos. Trans. R. Soc. Lond. B Biol. Sci. 374:20180235.

Wood C et al. 1994. Urea production, acid-base regulation and their interactions in the Lake
Magadi tilapia, a unique teleost adapted to a highly alkaline environment. Journal of
Experimental Biology. 189:13–36.

Wood CM. 1991. Acid-Base and Ion Balance, Metabolism, and their Interactions, after
Exhaustive Exercise in Fish. J. Exp. Biol. 160:285–308.

Wood CM, Bucking C, Grosell M. 2010. Acid--base responses to feeding and intestinal Cl--
uptake in freshwater-and seawater-acclimated killifish, Fundulus heteroclitus, an agastric
euryhaline teleost. J. Exp. Biol. 213:2681–2692.

Wu DD, Yang CP, Wang MS, Dong KZ. 2020. Convergent genomic signatures of high-altitude
adaptation among domestic mammals. National Science Review. 7:952–963.

Wu Y. 1992. The Fishes of the Qinghai-Tibetan Plateau. Sichuan Publishing House of Science
& Technology.

Xiao S et al. 2020. Genome of Tetraploid Fish Schizothorax o'connori Provides Insights into
Early Re-diploidization and High-Altitude Adaptation. iScience. 23:101497.

Xu J et al. 2017. Genomic Basis of Adaptive Evolution: The Survival of Amur Ide (Leuciscus
waleckii) in an Extremely Alkaline Environment. Mol. Biol. Evol. 34:145–159.

Xu Q et al. 2016. Analysis of the erythropoietin of a Tibetan Plateau schizothoracine fish
(Gymnocypris dobula) reveals enhanced cytoprotection function in hypoxic environments. BMC
Evol. Biol. 16:11.

Yang L, Wang Y, Zhang Z, He S. 2014. Comprehensive transcriptome analysis reveals
accelerated genic evolution in a Tibet fish, Gymnodiptychus pachycheilus. Genome Biol. Evol.
7:251–261.

Zhang K-Q et al. 2019. 14-3-3 gene family in spotted sea bass (Lateolabrax maculatus):
Genome-wide identification, phylogenetic analysis and expression profiles after salinity stress.
Comp. Biochem. Physiol. A Mol. Integr. Physiol. 235:1–11.

Zhao Y et al. 2020. Transcriptome changes for Nile tilapia (Oreochromis niloticus) in response
to alkalinity stress. Comp. Biochem. Physiol. Part D Genomics Proteomics. 33:100651.

Zhao Y, Wu J-W, Wang Y, Zhao J-L. 2016. Role of miR-21 in alkalinity stress tolerance in
tilapia. Biochem. Biophys. Res. Commun. 471:26–33.

Zheng M. 1997. An Introduction to Saline Lakes on the Qinghai—Tibet Plateau. Springer
Science & Business Media.

Zhou C et al. 2020. Comprehensive transcriptome data for endemic Schizothoracinae fish in the
Tibetan Plateau. Sci Data. 7:28.

Zhu S, Wu Y. 1975. Study of fish fauna in Qinghai Lake. Science Press: Beijing.

Figure legend

FIG. 1. Overview of geographic distributions of two alkaline tolerant schizothoracine fish species and their habitats in the northeastern Tibetan Plateau. Map depicting the geographic distributions of *G. p. przewalskii* in Lake Qinghai, and *G. p. kelukehuensis* in Lake Keluke. Photos showing the representative specimens of *G. p. przewalskii* and *G. p. kelukehuensis*. Photo credit: Chao Tong and Kai Zhao.

FIG.2. The flowchart represents the analysis pipeline: (1) sample collection and transcriptomics; (2) data assembling and annotation; (3) gene ortholog; (4) phenotype data and (5) molecular evolution analysis.

FIG. 3. Consistent patterns of molecular evolution in alkaline tolerant fish species. (A-B) Species trees for lowland and highland fish taxa were pruned from the Fish Tree of Life (https://fishtreeoflife.org/). Alkaline tolerant taxa and alkaline intolerant taxa are depicted in orange and sky blue, respectively. Two schematic diagrams depicting the comparisons made between alkaline tolerant and alkaline intolerant fish species, $\omega1$ representing the rate of molecular evolution of alkaline tolerant species and $\omega2$ for the alkaline intolerant species (C) Number of tested orthologs (N = 5,748), and number of orthologs under consistent shifts in rates of molecular evolution in alkaline tolerant relative to alkaline intolerant species within highland fish taxa and lowland fish taxa ($P < 0.05$, likelihood ratio test; false discovery rate < 0.2).

FIG. 4. Consistent signature of accelerated evolution in alkaline tolerant species within highland and lowland fish taxa.
(A) Venn diagram depicting numbers of rapidly evolving genes (REGs) in highland alkaline tolerant species, lowland alkaline tolerant species and both.
(B) Highland fish specific REGs, lowland fish specific REGs and overlapping REGs mainly related to ion transport, transmembrane and energy metabolism functions. The table showing the representative REGs under each category.
(C) REVIGO plot depicting the dominant enriched GO terms for REGs in highland alkaline tolerant fish. The scale of dark dot indicating the number of included enriched GO terms under a dominant GO term, the color scale representing the *P* value transformed by log10.
(D) REVIGO plot depicting the dominant enriched GO terms for REGs in lowland alkaline tolerant fish.

FIG. 5. Consistent signature of positive selection in alkaline tolerant species within highland and
lowland fish taxa.
(A) Venn diagram depicting numbers of positively selected genes (PSGs) in highland alkaline
tolerant species, lowland alkaline tolerant species and both.
(B) Highland fish specific PSGs, lowland fish specific PSGs and overlapping PSGs mainly
related to four categories, including ion transport/transmembrane, apoptosis/cell death, immune
response and energy metabolism. The table showing the representative PSGs under each
category.
(C) REVIGO plot depicting the dominant enriched GO terms for PSGs in highland alkaline
tolerant fish. The dark dot scale indicating the number of included GO terms under a dominant
GO term, the color scale representing the P value transformed by log10.
(D) REVIGO plot depicting the dominant enriched GO terms for PSGs in lowland alkaline
tolerant fish.
684

FIG.6. The intersection of rapidly evolving genes (REGs) and positively selected genes (PSGs)
in alkaline tolerant species within highland and lowland fish taxa.
(A) Venn diagram depicting numbers of REGs, PSGs and their overlapping genes in highland
alkaline tolerant species.
(B) REVIGO plot depicting the dominant enriched GO terms for overlapping genes in highland
alkaline tolerant species.
(C) Venn diagram depicting numbers of REGs, PSGs and their overlapping genes in lowland
alkaline tolerant species.
(D) REVIGO plot depicting the dominant enriched GO terms for overlapping genes in lowland
alkaline tolerant species.
695

Supplementary table capture

supplementary table S1: Information for additional omics data.

supplementary table S2: Statistics of transcriptome assembly and protein-coding genes.

supplementary table S3: Statistics of orthologs in 15 fish species.

supplementary table S4: Patterns of molecular evolution in alkaline tolerant and alkaline

intolerant species within lowland fish taxa and highland fish taxa.

supplementary table S5: Rapidly evolving gene repertoires in highland alkaline tolerant and

lowland alkaline tolerant species.

supplementary table S6: Significantly enriched GO terms for rapidly evolving gene of highland

alkaline tolerant and lowland alkaline tolerant species.

supplementary table S7: Positively selected gene repertoires in highland alkaline tolerant and

lowland alkaline tolerant species.

supplementary table S8: Significantly enriched GO terms for positively selected gene of

highland alkaline tolerant and lowland alkaline tolerant species.

supplementary table S9: Shared REG/PSG repertoires in highland alkaline tolerant and lowland

alkaline tolerant species.

supplementary table S10: significantly enriched GO terms for shared REG/PSG of highland

alkaline tolerant and lowland alkaline tolerant species.

Lake Keluke

Lake Qinghai

Gymnocypris przewalskii kelukehuensis

Gymnocypris przewalskii przewalskii

Lake Keluke

Lake Qinghai

Tibetan
Plateau

N

Sample collection and transcriptomics

tissue dissection

gill kidney brain heart liver

RNA extraction

quantity and quality control

library preparation

transcriptome sequencing

Illumina NovaSeq 6000

Date assembling and annotation

raw reads

quality control

fastqc, Trimmomatic

trim adaptors, contaminated and low-quality reads

assembly

Trinity, cd-hit

1. de novo assemble reads to transcripts
2. remove redundance and obtain unigenes

protein-coding genes

TransDecoder

obtain the open reading frame regions from transcripts

Annotation

Trinotate

assign gene ontology (GO) to each gene

Gene orthology

download additional genomes and transcriptoms from NCBI

data collection

pruning the Fish Tree of Life

phylogenetic tree

phangorn

identify orthologous groups (OG) in 15 fish species

ortholog identification

OMA

Accelerated evolution
[branch model]

shared orthologs

HyPHY

nucleotide substitution rate

dN/dS ω1: alkaline tolerant
ω2: alkaline intolerant

Likelihood-ratio test
False discovery rate

$P < 0.05$, FDR < 0.2

gene set 1

identification of rapidly evolving gene

ω1 > ω2

rapidly evolving genes

gene set enrichment analysis

topGO

Positive selection
[branch-site (BS) model]

shared orthologs

BUSTED

BS Unrestricted Statistical for Episodic Diversification

LRT $P < 0.05$, FDR < 0.2

gene set 1

1st filtration   aBSREL

adaptive BS Random Effects Likelihood

Holm-Bonferroni corrected $P < 0.05$

gene set 2

2nd filtration   BS + MNM

filter false positve cases with multinucleotide mutations (MNMs)

LRT $P < 0.05$

positively selected genes

gene set enrichment analysis

topGO

Phenotype data

lowland fish   highland fish

alkaline tolerant

Ssal Fhet   Gkel
Onil   Gprz

alkaline intolerant

Drer Cide   Oste Syou Gnam
Ccar Caur   Pext Spyl Geck

A Lowland fish taxa

- Carassius auratus (Caur)
- Cyprinus carpio (Ccar)
- Ctenopharyngodon idellus (Cide)
- Danio rerio (Drer)
- Fundulus heteroclitus (Fhet)
- Oreochromis niloticus (Onil)
- Salmo salar (Ssal)

● Alkaline tolerant
● Alkaline intolerant

B Highland fish taxa

- Oxygymnocypris stewartii (Oste)
- Schizopygopsis younghusbandi (Syou)
- Gymnocypris namensis (Gnam)
- Platypharodon extremus (Pext)
- Schizopygopsis pylzovi (Spyl)
- Gymnocypris eckloni (Geck)
- Gymnocypris przewalskii przewalskii (Gprz)
- Gymnocypris przewalskii kelukehuensis (Gkel)

C

	Number of tested orthologs	Number of orthologs with significance
Lowland fish taxa	5,748	952
Highland fish taxa	5,748	162

A

rapidly evolving genes (REGs)

highland fish

lowland fish

B

ion transport / transmembrane

SLC35F4	OSTM1
SLC6A4	TMEM192
SLC7A3	TMEM266
KCNJ15	TRPV1

energy metabolism

ME1
GET3
GPAM

SLC34A1	SLC20A1b

ENPP1

SLC45A3	SLC5A2	KCNE4
SLC35E1	SLC45A3	KCNG1
SLC51A	SLC4A4	KCNJ5
SLC22A16	TMEM168	KCNAB1
SLC5A2	TMEM26	KCNAB2

ATP5PB
ATP6V1C1
ADPGK
ADPRH
ABCB9

C Ontology for REGs in highland alkaline tolerant fish

D Ontology for REGs in lowland alkaline tolerant fish

A

rapidly evolving genes

Highland alkaline tolerant fish

positively selected genes

B

Ontology for shared REG/PSG in highland alkaline tolerant fish

amino acid import
basic amino acid transport
cardiac muscle cell apoptotic process
muscle cell apoptotic process
ion transport
carbohydrate transport
anion transport
anion transmembrane transport
aiditol phosphate metabolic process
mitochondrial fusion
response to heat
CDP-diacylglycerol metabolic process
oxoacid metabolic process
regulation of calcium ion import
purine nucleoside bisphosphate metabolic process
nucleoside bisphosphate metabolic process
branched-chain amino acid metabolic process
monovalent inorganic anion homeostasis
regulation of glucan biosynthetic process
pyrimidine-containing compound metabolic process
negative regulation of muscle cell apoptotic process
nucleobase-containing small molecule metabolic process
heme metabolic process
regulation of muscle cell apoptotic process

semantic space x

semantic space y

C

rapidly evolving genes

Lowland alkaline tolerant fish

positively selected genes

D

Ontology for shared REG/PSG in lowland alkaline tolerant fish

negative regulation of ion transport
regulation of membrane repolarization
nitric oxide biosynthetic process
guanosine-containing compound metabolic process
nuclear transport
mRNA metabolic process
miRNA metabolic process
nucleobase-containing compound transport
mRNA transport
reactive nitrogen species metabolic process

semantic space x

semantic space y

0
−0.5
−1.0
−1.5
−2.0
−2.5