Title: Red-beet betalain pigments inhibit amyloid-β aggregation and toxicity in

amyloid-β expressing *Caenorhabditis elegans*

Running title: Red-beet betalain pigments inhibit amyloid-β aggregation

Author names and affiliations:

Tomohiro Imamura [a*], Hironori Koga [b], Yasuki Higashimura [c], Noriyoshi Isozumi [d],

Kenji Matsumoto [c], Shinya Ohki [d], and Masashi Mori[a*]

[a]Research Institute for Bioresources and Biotechnology, Ishikawa Prefectural University,

308-1 Suematsu, Nonoichi, Ishikawa 921-8836, Japan

[b]Department of Bioproduction Science, Ishikawa Prefectural University, 308-1 Suematsu,

Nonoichi, Ishikawa 921-8836, Japan

[c]Department of Food Science, Ishikawa Prefectural University, 308-1 Suematsu,

Nonoichi, Ishikawa 921-8836, Japan

[d]Center for Nano Materials and Technology (CNMT), Japan Advanced Institute of

Science and Technology (JAIST), 1-1 Asahidai, Nomi, Ishikawa 923-1292, Japan

*Corresponding authors: Tomohiro Imamura and Masashi Mori

Address: Research Institute for Bioresources and Biotechnology, Ishikawa Prefectural

University, 308-1 Suematsu, Nonoichi, Ishikawa 921-8836, Japan (TI and MM)

Phone: +81-76-227-7527 (TI and MM)

Fax: +81-76-227-7557 (TI and MM)

E-mail: timamura@ishikawa-pu.ac.jp (TI), mori@ishikawa-pu.ac.jp (MM)

ORCID: 0000-0003-2277-9134 (TI), 0000-0002-9021-8104 (MM)

ABSTRACT

BACKGROUND: Betalain pigments are mainly produced by plants in the order Caryophyllales. Recent interest in the biological functions of betalain pigments has increased with antioxidant, anti-inflammatory, and anticancer activities reported.

RESULTS: We investigated the effects of betalain pigments derived from red-beet on amyloid-β (Aβ) aggregation, a cause of Alzheimer's disease. Inhibition of Aβ aggregation against Aβ40 and Aβ42 by betalain pigments *in vitro* was demonstrated by the Thioflavin T fluorescence assay, circular dichroism spectroscopy analysis and transmission electron microscopic observations. Moreover, we examined the ability of betalain pigments to interfere with Aβ toxicity by using the transgenic *Caenorhabditis elegans* strain CL2006, which expresses the human Aβ42 protein intracellularly within the body wall muscle and responds to Aβ-toxicity with paralysis. Treatment with 50 μM betalain pigments significantly delayed the paralysis of *Caenorhabditis elegans*.

CONCLUSION: These results suggest that betalain pigments reduce Aβ-induced toxicity by inhibiting Aβ aggregation and may lead to their use as inhibitors of Aβ aggregation.

Keywords: amyloid-beta aggregation, *Caenorhabditis elegans*, circular dichroism spectroscopy, red-beet betalain pigments, transmission electron microscopy, Thioflavin T fluorescence assay

INTRODUCTION

Betalains are tyrosine-derived pigments found exclusively in plants of the order

Caryophyllales,[1] some *Basidiomycota* fungi[2] and one bacterial species,

Gluconacetobacter diazotrophicus.[3] Betalain pigments are divided into betacyanins (red

and purple) and betaxanthins (yellow and orange). Unlike other plant pigments, e.g.,

flavonoids and carotenoids, betalains contain nitrogen. Previous studies have

demonstrated that betalains exhibit strong antioxidant activity[4] and are involved in plant

responses to environmental stimuli and stress.[5, 6] Betalains also have commercial

applications; they are often used as food additives because of their vivid colors. For

example, beetroot extract, designated by the label E162, is approved by the United States

Food and Drug Administration (FDA) and European Union regulatory agencies for use as

a natural food colorant.

Biological activities of betalain pigments have been reported. Betalain-rich extracts

possess anti-inflammatory, anticancer and anti-diabetic properties.[7-10] Furthermore,

betanin, the major red pigment of beetroot extract, has been demonstrated to inhibit

low-density lipoprotein oxidation.[11, 12] Our recent study found that the amaranthin

betalain pigment, which accumulates in amaranth and quinoa, inhibits HIV-1 protease activity.[13] Although various physiological functions of the betalain pigments have been reported, it is envisaged that many of the physiological activities of these pigments remain undiscovered.

Alzheimer's disease (AD) is an irreversible, progressive brain disorder that progressively impairs memory and the ability to think.[14] From a neuropathological viewpoint, AD is characterized by the aggregation of the amyloid-β (Aβ) protein in senile plaques and hyperphosphorylated tau protein in brain neurofibrillary tangles.[15] Thus, inhibiting the aggregation and accumulation of Aβ and phosphorylated tau in the brain is the key to treating AD. However, all current FDA-approved anti-AD drugs are only symptomatic treatments, which do not cure the disease. Recent studies have reported the efficacy of plant-derived polyphenolic compounds as candidate substances for AD treatment.[16, 17] Polyphenolic compounds such as resveratrol, curcumin, quercetin, genistein, morin, and epigallocatechin gallate have been shown to inhibit Aβ production and aggregation.[18, 19]

Caenorhabditis elegans (*C. elegans*) is currently used in many studies as a model

organism for AD.[20] *C. elegans* is a free-living, non-parasitic nematode. Approximately

38% of the *C. elegans* genome has human orthologs, including AD-related orthologs such

as the a*myloid-bata precursor protein* and *tau*.[21] Therefore, *C. elegans* has many

advantages as an *in vivo* model for AD research. Moreover, a transformant strain CL2006

expressing the human amyloid gene has been generated.[22] This transformed strain

exhibits a paralytic phenotype by accumulating human amyloid-β 1–42 (Aβ42) in

muscle.[22] This transformant phenotype is useful for evaluating substances that are

effective against AD. Many studies using this nematode to examine the bioactivity of

phytochemicals in Aβ aggregation have been reported;[23–25] however, the bioactivity of

betalain pigments have not been reported in AD studies using nematodes.

In this study, we evaluated the biological activity of betalain pigments against Aβ

aggregation using red-beet betalain pigments, mainly composed of betanin and

isobetanin. The *in vitro* ThT fluorescence assay and circular dichroic (CD) spectroscopy

analysis were performed to evaluate the effect of red-beet betalain pigments on Aβ

aggregation. Furthermore, sample of Aβ with and without treatment with red-beet

betalain pigments were examined by transmission electron microscopy (TEM). In all

experiments, red-beet betalain pigments were found to inhibit Aβ aggregation *in vitro*.

Additionally, we evaluated the biological activity of red-beet betalain pigments using an

Aβ expressing nematode strain CL2006. Feeding red-beet betalain pigments to Aβ

expressing *C. elegans* alleviated the severity of the phenotype associated with Aβ

aggregation.

MATERIAL AND METHODS

Materials

Lyophilized human amyloid-β 1–40 (Aβ40) and amyloid-β 1–42 (Aβ42) were obtained

from SCRUM Inc. (Tokyo, Japan). Red-beet betalains (red-beet betanin extract diluted

with dextrin) were obtained from Tokyo Chemical Industry (TCI, Tokyo, Japan). Morin

hydrate (TCI) was used as a control for the ThT fluorescence assay. Dextrin (TCI) was

used as a control in nematode experiments.

Sample preparation

Red-beet betalain pigments were mixed with dextrin for preservation. Betalain pigments

were purified from red-beet betalain pigments to remove any side effects of dextrin when

conducting *in vitro* experiments. First, an aqueous solution of red-beet betalain pigment

was prepared and added to DEAE Sephacel (Pharmacia, Uppsala, Sweden). The solution

was washed with water before eluting the pigments with 300 mM NaCl. Next, the eluate

was adsorbed on cellulose powder (Fuji Film, Osaka, Japan) and washed three times with

ethanol. Subsequently, the pigments were eluted with water and concentrated. The eluate

was added to reversed-phase Cosmosil 140C_{18}-OPN (Nacalai tesque, Kyoto, Japan),

washed with water and eluted with 10% acetonitrile. The eluate was evaporated to

dryness and residues were dissolved in water and stored at –20 °C until needed. These

purified red-beet betalain pigments were used for *in vitro* experiments.

Preparation of Aβ monomer

Aβ monomer was prepared according to previous studies.[26, 27] Aβ was dissolved in 100%

1,1,1,3,3,3-hexafluoro-2-propanol (TCI) at 1 mg/mL and incubated for 2 h at room

temperature and then sonicated for 30 min. Subsequently, the solution was centrifuged

15000 × *g* and 4 °C for 30 min to remove the existing Aβ aggregates. The supernatant

was collected and the solvent was removed by lyophilization overnight. The dried Aβ

peptide was stored at −20 °C until needed.

High-performance liquid chromatography (HPLC)

A Shimadzu LC-20AD system (Kyoto, Japan) was used for analytical HPLC

separations. Samples were separated on a Shim-pack GWS C18 column (5 µm; 200 ×

4.6 mm i.d.; Shimadzu GLC, Tokyo, Japan), and linear gradients were run from 0% B to

45% B over 45 min using 0.05% trifluoroacetic acid (TFA) in water (solvent A) and

0.05% TFA in acetonitrile (solvent B). Elution was monitored by measuring the

absorbance at 536 nm. The flow rate was 0.5 mL min^{-1} and the column temperature

25 °C.

UV−Vis spectroscopy

A UV-2450 (Shimadzu) spectrophotometer was used for UV-Vis spectroscopy. The

concentration of betalain pigments without dextrin was determined using the molar

extinction coefficient of $\varepsilon = 65{,}000$ M^{-1} cm^{-1} at 536 nm for purified red-beet betalain

pigments.[28, 29]

ThT fluorescence assay

The ThT fluorescence assay was performed using the SensoLyte Thioflavin T

β-Amyloid Aggregation Kit (ANASPEC, Fremont, CA, USA). Aβ and ThT solutions at

final concentrations of 50 μM and 200 μM, respectively, were used in the evaluation

system according to the manufacturers' protocol. Aβ40 dissolved in phosphate buffered

saline (PBS, pH7.4) or 10 mM NaOH and Aβ42 dissolved in 10 mM NaOH were used.

Aβ solutions with red-beet betalain extract at final concentrations of 6.5, 12.5, 25, and 50

μM were evaluated. Morin, a flavonoid with inhibitory acitivity against Aβ aggregation,[30]

was used as an inhibitor control at a final concentration of 50 μM. A water sample without

red-beet betalain extract or morin was used as the positive control, and a sample without

inhibitor and Aβ40 was used as the reference control. ThT fluorescence was monitored at

37 °C using a spectrofluorometer (VarioskanLUX; Thermo Fisher Scientific, Waltham,

MA, USA) at Ex/Em = 440/484 nm. Reading were taken every 5 min and each sample

was shaken for 15 s immediately prior to the reading. Fluorescence data were analyzed

using Skanlt software (Thermo Fisher Scientific). The reported values are the average of four wells.

Circular dichroic (CD) spectroscopy analysis

For CD spectra analysis, 40 µM Aβ40 incubated with or without 200 µM red-beet betalain pigments for 7 days and 20 µM Aβ42 incubated with or without 100 µM red-beet betalain pigments for 4 days. The 7 days incubation sample of Aβ42 could not be measured due to precipitation. These incubated samples were measured using a CD spectropolarimeter (J-820, JASCO, Tokyo, Japan) at room temperature. Far-UV (190–260 nm) was measured with a scan speed of 100 nm/min. Every sample was scanned three times, and spectra presented are shown as the average of these results.

TEM observation of Aβ aggregates

Aβ40 or Aβ42 was solubilized in DMSO or 10 mM NaOH to give a 1 mM solution, respectively. Aβ solutions were prepared with and without the addition of 50 µM red-beet betalain pigments and diluted to 20 µM with PBS. Aβ mixtures were incubated at 37 ℃

for 5 days before TEM observations.

Approximately 2 µL of sample solution was placed on a 150-mesh copper grid

covered with formvar. After 5 min the sample was soaked away, then the grid was

stained with 2.0% (w/v) uranyl acetate solution. TEM images of Aβ aggregates were

obtained using a Hitachi 7650 TEM (Hitachi Co., Ltd., Tokyo, Japan) with an

acceleration voltage of 80 kV.

Caenorhabditis elegans maintenance and synchronization

C. elegans were maintained on nematode growth medium (NGM) plates seeded with

Escherichia coli OP50 (OP50) as a food source at 15 °C unless otherwise indicated. The

transgenic strain CL2006 [dvIs (unc-54::human β-amyloid 1–42; pRF4)] was obtained

from the Caenorhabditis Genetics Center (CGC; MN, USA). In CL2006, the human

Aβ42 protein is expressed intracellularly within the body wall muscle. The expression

and subsequent aggregation of Aβ in the muscle result in progressive paralysis.[22]

For synchronization, the nematodes were cultured on fresh NGM plates for 2–3

generations without starvation. Young adult nematodes were collected and cleaned, then

disintegrated using a lysis solution (0.6% sodium hypochlorite and 200 mM NaOH).

After 12–14 h at 20 °C, the isolated eggs were hatched to obtain the synchronized L1

larvae.

Paralysis assay

The synchronized L1 larvae were fed OP50 and grown to young adults at 20 °C;

5-fluorodeoxyuridine (0.5 mg/ml) was added to prevent progeny production. The

resultant synchronized hermaphrodites were transferred to OP50-seeded NGM plates

containing various dextrin or red-beet betalain pigments concentrations (20 nematodes

per plate). Dextrin was used as a control because the red-beet betalain pigments used in

this study contained dextrin. The nematodes were tested for paralysis by tapping their

noses with a platinum wire. According to a previous report, nematodes that moved their

noses in response but fail to move their bodies were paralyzed.[31] Paralyzed worms were

calculated by the Kaplan-Meier method, and the difference in proportion of paralyzed

worms was tested for significance using the log-rank test. Statistical analyses were

performed using statistical software (GraphPad Prism ver. 8.0.1; GraphPad Software,

Inc., La Jolla, CA, USA). Differences that gave *$p < 0.05$ were defined as statistically

significance.

Animals and surgery

Twenty-five male C57BL/6J mice (six-week-old) were purchased from Charles River

Laboratories Japan (Yokohama, Kanagawa, Japan) and fed a standard rodent diet, CE-2

(CLEA Japan, Tokyo, Japan), for six weeks. At 12 weeks of age, mice were divided into

five groups of five based on their body weight. Red-beet betalain pigments were

suspended in distilled water at 83.3 mg/mL and its oral gavage carried out using a

feeding needle (2.7 μmol/kg body weight as red-beet betalain pigments). After 1, 2, 3

and 6 h, mice were euthanized by CO_2 inhalation and dissected. For the sham group,

oral gavage of distilled water was performed and mice were immediately euthanized by

CO_2 inhalation. After blood sampling from the postcaval vein, and brain tissues were

collected and frozen by using liquid nitrogen. Separated plasma and tissues were stored

at −80 °C until analysis. Experiments were conducted following the guidelines for the

appropriate conduct of animal experiments set by the Scientific Council of Japan (2006).

The Animal Experimentation Ethics Committee of Ishikawa Prefectural University

(Ishikawa, Japan) approved the study protocol (31–14–34).

Quantification of betalain in brain samples

The amount of red-beet betalain pigments within brain samples was evaluated at selected

time intervals (1, 2, 3, 6 h) after a single oral administration of the pigments. Brains,

plasma, and red blood cells (five per time point) were then isolated and homogenized in

PBS. Red-beet betalain pigments were extracted from samples with chloroform:methanol,

at a ratio of 2:1 by volume (1 g of tissue with three volumes of extraction mixture). The

methanol phase from all samples at each time point was dried, resuspended in 1% (v/v)

acetic acid in the water and analyzed by HPLC.

RESULTS

Pigment analysis of red-beet betalain pigments

HPLC was used to investigate the composition of the betalain pigments present in the

red-beet betalain extract. The red-beet betalain pigments, betanin and isobetanin were

found to be the major pigments present in the extract, and other betalain pigments were

not ascertained (Fig. 1). The proportions of betanin and isobetanin in this extract were

almost equal (Fig. 1).

Red-beet betalain extracts inhibit the aggregation of Aβ40

Initially, the effect of red-beet betalain pigments on Aβ40 aggregation was investigated

by using the ThT fluorescence assay. Thioflavin T used in this assay is a fluorescent dye

that binds to amyloid and increases in fluorescence intensity. Aβ40 was dissolved in PBS,

and red-beet betalain pigments at various concentrations (6.25, 12.5, 25 and 50 µM) were

added. All samples with red-beet betalain pigments present had lower ThT fluorescence

when compared with that of the Aβ40 alone sample (Fig. 2A), suggesting that red-beet

betalain pigments inhibit formation of Aβ40 amyloids. Furthermore, the inhibitory

effects were similar between 6.25 µM red-beet betalain pigments and 50 µM morin, an

amyloid inhibitor (control) isolated from mulberry fruit (Fig. 2A). The red-beet betalains

over 6.25 µM showed higher Aβ40 aggregation inhibitory activity than 50 µM morin.

Moreover, the ThT assay was performed using Aβ42 and Aβ40 dissolved in 10 mM

NaOH because Aβ42 hardly dissolved in PBS. The results showed that the fluorescence

of these Aβ samples containing red-beet betalains pigments were lower than that of

Aβ42 and Aβ40 alone, but higher than Aβ samples containing morin (Fig. 2B, C).

The effect of red-beet betalain pigments on the secondary structure of Aβ

CD spectroscopy was performed to detect secondary structure changes of Aβ in the

presence of red-beet betalain pigments (a 5:1 molar ratio of the pigment to Aβ, Fig. 3).

During the initial stage of Aβ aggregation, the CD spectra of Aβ40 and Aβ42 without

red-beet betalain pigments showed a negative minimum at 196–199 nm, demonstrating

that Aβ40 and Aβ42 adopted no secondary structure, i.e., random coil (Fig. 3A, C). After

incubating at room temperature, the CD spectrum of Aβ without red-beet betalain

pigments did not show a negative minimum at 196–199 nm but a new characteristic peak

was appeared at 217 nm (Fig. 3B and D). These CD spectra showed that Aβ40 and Aβ42

underwent a structural transition from random coil to the β-sheet secondary structure.[32]

After incubation at room temperature, the CD signals at 217 nm arising from Aβ40 and

Aβ42 in the presence of red-beet betalain pigments were significantly weaker when

compared with the spectra of Aβ alone (Fig. 3B and D). This indicates that the amount of

β-sheet structure in Aβ was clearly reduced by red-beet betalain pigments. These results

suggested that red-beet betalain pigments hamper the onset of the conformational

transition of Aβ40 and Aβ42 from random coil toβ-sheet.

Red-beet betalain extracts modulate the morphology of Aβ Aggregates

The inhibitory effect of red-beet betalain pigments on Aβ aggregation was observed using

TEM (Fig 4). Aβ40 and Aβ42 solutions (25 µM) with and without red-beet betalain

pigments (50 µM) were incubated at 37 °C for 5 days, after which Aβ aggregation was

monitored by TEM. Aβ fibrosis and amorphous aggregation were observed in Aβ40 and

Aβ42 samples without red-beet betalain pigments (Fig. 4A, C). In contrast, Aβ fibisis and

amorphous aggregation were significantly reduced in the sample with red-beet betalain

pigments (Fig. 4B, D).

Red-beet betalain pigments delayed Aβ-induced paralysis in a transgenic nematode

C. elegans CL2006 strain expressing human Aβ in muscle develops a paralysis

phenotype during grows. This transgenic nematode accumulates Aβ42, a major

component of the senile plaques observed in AD, and is considered a nematode model of

AD. We evaluated the inhibition of Aβ aggregation by red-beet betalain pigments *in vivo*

using CL2006 nematode, as indicated in the experimental schedule presented in Fig. 5A.

The nematodes were treated with dextrin or red-beet betalain pigments at concentrations

of 2, 10 and 50 µM. As shown in Kaplan-Meier plots, dextrin added as a stabilizing

agent for red-beet betalain pigments did not change the phenotype of the transgenic

nematodes (Fig. 5B). In contrast, the highest concentration of red-beet betalain pigments

(50 µM) delayed Aβ-induced paralysis in the transgenic nematodes significantly when

compared with that of control (no betalain pigments), whereas the 2 and 10 µM

treatments did not lead to a change in phenotype when compared with that of the control

(Fig. 5C). Representative microscopic images treated (i.e., 50 µM red-beet betalain

pigments) and control groups at day 15 are shown in Fig. 5D. By microscopic

observation, many individual nematodes in the control group adopted a linear paralyzed

body phenotype, whereas linear paralysis was not as distinct for nematodes in the group

treated with 50 µM red-beet betalain pigments.

DISCUSSION

In this study, results from the ThT fluorescence assay, CD spectroscopy and TEM showed

that red-beet betalain pigments composed of betanin and isobetanin inhibits Aβ

aggregation. Furthermore, in feeding experiments using Aβ expressing nematodes, the

ingestion of red-beet betalain pigments delayed the onset of paralytic symptoms caused

by Aβ aggregation. Together, these results strongly suggest that betanin and isobetanin,

which are the main components of red-beet betalain pigments, inhibit the aggregation of

Aβ.

Previous studies have reported that betalain pigments have various biological

effects, including anticancer,[10, 13] anti-inflammatory,[33] inhibition of low-density

lipoprotein oxidation[11, 12] and inhibition of HIV-1 protease activity.[13] Recently,

nematode feeding experiments have shown that specific ingestion of betalain pigments

produce lifespan-prolonging effects.[34, 35] This study discovered an inhibitory effect on Aβ

aggregation as a new physiological effect of betalain pigments.

PBS and 10 mM NaOH were used as solvents for Aβin the ThT assay. The results

showed that the inhibitory activity of betalain pigments against Aβ aggregation was

lower when the sample was dissolved in 10 mM NaOH versus dissolution in PBS. This observation indicated that pH affects the stability of betalain pigments. Betalains are reported to be stable between weakly acidic and neutral condition but are unstable outside this pH range.[36] Therefore, the use of 10 mM NaOH may have destabilized the betalain pigments, which reduced their inhibitory activity against Aβ aggregation during the ThT assay. Nonetheless, red-beet betalain pigments are anticipated to be active against Aβ aggregation because the *in vivo* pH in human is near neutral.

Red-beet betalain pigments were found to suppress Aβ induced toxicity in transgenic nematodes. These transformed nematodes express Aβ42 in muscle but as a model organism do not have a blood-brain barrier (BBB). In mammals, red-beet betalain pigments must cross the BBB to accumulate in the brain and inhibit aggregation of Aβ. The betalain pigment, indicaxanthin, has been reported to cross the BBB in rats.[37] As a preliminary experiment, we fed red-beet betalain pigments to mice and evaluated whether the pigments accumulated in the brain. However, betanin accumulation in the mouse brain was not confirmed (Suppl. Table 1). Nonetheless, a recent study reported that feeding betalain pigments to Alzheimer's model rats alleviates

AD symptoms.[38] Thus, further *in vivo* studies are required to clarify whether betalains

act directly to inhibit amyloid polymerization by passing across the BBB or function via

a different mechanism such as modulating oxidative stress in blood.

CONCLUSIONS

We found that beet-derived betalains inhibit the aggregation of Aβ. Additionally,

feeding tests using Aβ-expressing *C. elegans* suggested that betalains inhibit Aβ

aggregation *in vivo*. As a prospect, further analysis of betalain pigments may lead to

their use as inhibitors of Aβ aggregation and as supplements to prevent AD. In this study,

we conducted a bioactivity evaluation using *C. elegans*. The evaluation of

phytochemicals bioactivity using nematodes is an effective initial selection process.

There are many types of molecular species in betalain pigments.[39] Future research efforts

may lead to the discovery of betalain molecular species that function more effectively in

inhibiting Aβ aggregation than red-beet betalain pigments. Furthermore, by using

various model nematodes,we expect to find new bioactivities of betalain pigments.

ACKNOWLEDGMENTS

The authors thank Akiko Mizuno, Hiroko Hayashi, Mami Awatani and Hitomi Nishikawa for their excellent technical assistance. This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors. We thank Edanz Group (https://en-author-services.edanz.com/ac) for editing a draft of this manuscript.

CONFLICT OF INTERESTS

There is no conflict of interests to declare.

CONTRIBUTIONS

MM conceived the study. TI and MM designed the experiments. TI performed the ThT fluorescence assay. HK performed TEM observations. YH conducted the nematode experiment. KM and YH conducted the experiment using mice. NI and SO performed CD spectroscopy analysis. TI, YH, KM, and MM wrote the draft manuscript. All authors

have read and approved the final manuscript.

SUPPORTING INFORMATION

Supporting information may be found in the online version of this article.

REFERENCES

1.   Brockington SF, Walker RH, Glover BJ, Soltis PS and Soltis DE, Complex

     pigment evolution in the Caryophyllales. *New Phytol* **190**:854–864 (2011).

2.   Michelot D and Melendez-Howell LM, Amanita muscaria: chemistry, biology,

     toxicology, and ethnomycology. *Mycol Res* **107**:131–146 (2003).

3.   Contreras-Llano LE, Guerrero-Rubio MA, Lozada-Ramírez JD, García-Carmona

     F and Gandía-Herrero F, First betalain-producing bacteria break the exclusive

     presence of the pigments in the plant kingdom. *mBio* **10**:e00345–00319 (2019).

4.   Slimen IB, Najar T and Abderrabba M, Chemical and antioxidant properties of

     betalains. *J Agric Food Chem* **65**:675–689 (2017).

5.   Jain G, Schwinn KE and Gould KS, Betalain induction by l-DOPA application

confers photoprotection to saline-exposed leaves of Disphyma australe. *The New phytol* **207**:1075–1083 (2015).

6.   Polturak G, Grossman N, Vela-Corcia D, Dong Y, Nudel A, Pliner M, Levy M, Rogachev I and Aharoni A, Engineered gray mold resistance, antioxidant capacity, and pigmentation in betalain-producing crops and ornamentals. *Proc Natl Acad Sci USA* **114**:9062–9067 (2017).

7.   Lechner JF, Wang LS, Rocha CM, Larue B, Henry C, McIntyre CM, Riedl KM, Schwartz SJ and Stoner GD, Drinking water with red beetroot food color antagonizes esophageal carcinogenesis in N-nitrosomethylbenzylamine-treated rats. *J Med Food* **13**:733–739 (2010).

8.   Lugo-Radillo A, Delgado-Enciso I and Peña-Beltrán E, Betanidin significantly reduces blood glucose levels in BALB/c mice fed with an atherogenic diet. *Nat Prod Bioprospect* **2**:154–155 (2012).

9.   Martinez RM, Longhi-Balbinot DT, Zarpelon AC, Staurengo-Ferrari L, Baracat MM, Georgetti SR, Sassonia RC, Verri WA, Jr. and Casagrande R, Anti-inflammatory activity of betalain-rich dye of *Beta vulgaris:* effect on edema,

leukocyte recruitment, superoxide anion and cytokine production. *Arch Pharmacal Res* **38**:494–504 (2015).

10. Nowacki L, Vigneron P, Rotellini L, Cazzola H, Merlier F, Prost E, Ralanairina R, Gadonna JP, Rossi C and Vayssade M, Betanin-enriched red beetroot (*Beta vulgaris* L.) extract induces apoptosis and autophagic cell death in MCF-7 cells. *Phytother Res* **29**:1964–1973 (2015).

11. Tesoriere L, Butera D, D'Arpa D, Di Gaudio F, Allegra M, Gentile C and Livrea MA, Increased resistance to oxidation of betalain-enriched human low density lipoproteins. *Free Rradic Res* **37**:689–696 (2003).

12. Tesoriere L, Allegra M, Butera D and Livrea MA, Absorption, excretion, and distribution of dietary antioxidant betalains in LDLs: potential health effects of betalains in humans. *Am J Clin Nutr* **80**:941–945 (2004).

13. Imamura T, Isozumi N, Higashimura Y, Miyazato A, Mizukoshi H, Ohki S and Mori M, Isolation of amaranthin synthetase from *Chenopodium quinoa* and construction of an amaranthin production system using suspension-cultured tobacco BY-2 cells. *Plant Biotechnol J* **17**:969–981 (2019).

14. Shankar GM, Li S, Mehta TH, Garcia-Munoz A, Shepardson NE, Smith I, Brett FM, Farrell MA, Rowan MJ, Lemere CA, Regan CM, Walsh DM, Sabatini BL and Selkoe DJ, Amyloid-beta protein dimers isolated directly from Alzheimer's brains impair synaptic plasticity and memory. *Nature Med* **14**:837–842 (2008).

15. Cohen E, Bieschke J, Perciavalle RM, Kelly JW and Dillin A, Opposing activities protect against age-onset proteotoxicity. *Science* **313**:1604–1610 (2006).

16. Ayaz M, Sadiq A, Junaid M, Ullah F, Ovais M, Ullah I, Ahmed J and Shahid M, Flavonoids as prospective neuroprotectants and their therapeutic propensity in aging associated neurological disorders. *Front Aging Neurosci* **11**:155 (2019).

17. Williams RJ and Spencer JP, Flavonoids, cognition, and dementia: actions, mechanisms, and potential therapeutic utility for Alzheimer disease. *Free Radic Biol Med* **52**:35–45 (2012).

18. Kobayashi H, Murata M, Kawanishi S and Oikawa S, Polyphenols with anti-amyloid β aggregation show potential risk of toxicity via pro-oxidant properties. *Int J Mol Sci* **21**: 3561 (2020).

19. Vaiserman A, Koliada A and Lushchak O, Neuroinflammation in pathogenesis of

Alzheimer's disease: Phytochemicals as potential therapeutics. *Mech Ageing Dev* **189**:111259 (2020).

20. Alexander AG, Marfil V and Li C, Use of Caenorhabditis elegans as a model to study Alzheimer's disease and other neurodegenerative diseases. *Front Genet* **5**:279 (2014).

21. Shaye DD and Greenwald I, OrthoList: a compendium of *C. elegans* genes with human orthologs. *PLoS ONE* **6**:e20085 (2011).

22. Link CD, Expression of human beta-amyloid peptide in transgenic *Caenorhabditis elegans*. *Proc Natl Acad Sci USA* **92**:9368–9372 (1995).

23. Guo H, Cao M, Zou S, Ye B and Dong Y, Cranberry extract standardized for proanthocyanidins alleviates β-amyloid peptide toxicity by improving proteostasis through HSF-1 in *Caenorhabditis elegans* model of Alzheimer's disease. *J Gerontol A Biol Sci Med Sci* **71**:1564–1573 (2016).

24. Hao S, Li X, Han A, Yang Y, Luo X, Fang G, Wang H, Liu J and Wang S, Hydroxycinnamic acid from corncob and its structural analogues inhibit Aβ40 fibrillation and attenuate Aβ40-induced cytotoxicity. *J Agric Food Chem*

68:8788–8796 (2020).

25. Jagota S and Rajadas J, Effect of phenolic compounds against Aβ aggregation and

    Aβ-induced toxicity in transgenic *C. elegans*. *Neurochem Res* **37**:40–48 (2012).

26. Wang Q, Shah N, Zhao J, Wang C, Zhao C, Liu L, Li L, Zhou F and Zheng J,

    Structural, morphological, and kinetic studies of β-amyloid peptide aggregation

    on self-assembled monolayers. *Phys Chem Chem Phys* **13**:15200–15210 (2011).

27. Xiong N, Dong XY, Zheng J, Liu FF and Sun Y, Design of LVFFARK and

    LVFFARK-functionalized nanoparticles for inhibiting amyloid β-protein

    fibrillation and cytotoxicity. *ACS Appl Mater Interfaces* **7**:5650–5662 (2015).

28. Gandía-Herrero F, Escribano J and García-Carmona F, Structural implications on

    color, fluorescence, and antiradical activity in betalains. *Planta* **232**:449–460

    (2010).

29. Schwartz SJ and Von Elbe JH, Quantitative determination of individual

    betacyanin pigments by high-performance liquid chromatography. *J Agric Food

    Chem* **28**:540–543 (1980).

30. Hanaki M, Murakami K, Akagi K-i and Irie K, Structural insights into

mechanisms for inhibiting amyloid β42 aggregation by non-catechol-type

flavonoids. *Bioorg Med Chem* **24**:304–313 (2016).

31. Chen XQ and Mobley WC, Alzheimer disease pathogenesis: Insights from

molecular and cellular biology studies of oligomeric Aβ and tau species. *Front*

Neurosci **13**:659 (2019).

32. Pellarin R and Caflisch A, Interpreting the aggregation kinetics of amyloid

peptides. *J Mol Biol* **360**:882–892 (2006).

33. Allegra M, Ianaro A, Tersigni M, Panza E, Tesoriere L and Livrea MA,

Indicaxanthin from cactus pear fruit exerts anti-inflammatory effects in

carrageenin-induced rat pleurisy. *J Nutr* **144**:185–192 (2014).

34. Guerrero-Rubio MA, Hernández-García S, García-Carmona F and

Gandía-Herrero F, Extension of life-span using a RNAi model and in vivo

antioxidant effect of Opuntia fruit extracts and pure betalains in *Caenorhabditis*

elegans. *Food Chem* **274**:840–847 (2019).

35. Guerrero-Rubio MA, Hernández-García S, Escribano J, Jiménez-Atiénzar M,

Cabanes J, García-Carmona F and Gandía-Herrero F, Betalain health-promoting

effects after ingestion in *Caenorhabditis elegans* are mediated by DAF-16/FOXO
and SKN-1/Nrf2 transcription factors. *Food Chem* **330**:127228 (2020).

36. Herbach KM, Stintzing FC and Carle R, Betalain stability and degradation —
structural and chromatic aspects. *J Food Sci* **71**:R41–R50 (2006).

37. Allegra M, Carletti F, Gambino G, Tutone M, Attanzio A, Tesoriere L, Ferraro G,
Sardo P, Almerico AM and Livrea MA, Indicaxanthin from opuntia ficus-indica
crosses the blood-brain barrier and modulates neuronal bioelectric activity in rat
hippocampus at dietary-consistent amounts. *J Agric Food Chem* **63**:7353–7360
(2015).

38. Shunan D, Yu M, Guan H and Zhou Y, Neuroprotective effect of betalain against
AlCl$_3$-induced Alzheimer's disease in Sprague Dawley Rats via putative
modulation of oxidative stress and nuclear factor kappa B (NF-κB) signaling
pathway. *Biomed Pharmacother* **137**:111369 (2021).

39. Delgado-Vargas F, Jiménez AR and Paredes-López O, Natural pigments:
carotenoids, anthocyanins, and betalains--characteristics, biosynthesis, processing,
and stability. *Crit Rev Food Sci Nutr* **40**:173–289 (2000).

Figure legends

Figure 1. Composition of red-beet betalain pigments (A) Chemical structure of betanin and isobetanin (B) HPLC chromatogram of red-beet betalain pigments. HPLC elution sample peaks at 23.5 min and 24.6 min represent betanin and isobetanin, respectively.

Figure 2. ThT fluorescence during amyloid fibrillation of human amyloid-β (Aβ). (A) Thioflavin T (ThT) fluorescence changes for Aβ40 incubated with/without red-beet betalain pigments. Aβ40 was dissolved in PBS. (B) ThT fluorescence changes for Aβ42 incubated with/without red-beet betalain pigments. Aβ42 was dissolved in 10 mM NaOH. (C) ThT fluorescence changes for Aβ40 incubated with/without red-beet betalain pigments. Aβ40 was dissolved in 10 mM NaOH. Aβ was incubated with 6.25, 12.5, 25 and 50 μM red-beet betalain pigments at 37 ℃ for 200 min. Error bars represent the means ± SD ($n = 4$). ThT relative fluorescence was expressed as a percentage of fluorescence for the Aβ alone sample, which had a maximum value of 100%. "Beet" and "Morin" represent sample with red-beet betalain pigments and inhibitor control, respectively.

Figure 3. CD spectra of Aβ incubated with or without red-beet betalain pigment. (A) 40

μM Aβ40 incubated with or without 200 μM red-beet betalain pigment at 0 h. (B) 40

μM Aβ40 incubated with or without 200 μM red-beet betalain pigment at day 7. (C) 20

μM Aβ42 incubated with or without 100 μM red-beet betalain pigment at 0 h. (D) 20

μM Aβ42 incubated with or without 100 μM red-beet betalain pigment at day 4. Beet

indicates red-beet betalain pigments.

Figure 4. Transmission electron microscope images of human amyloid-β aggregates.

(A) Aβ40 alone; (B) Aβ40 with 50 μM red-beet betalain pigments; (C) Aβ42 alone; (D)

Aβ42 with 50 μM red-beet betalain pigments; Scale bars, 200 μm; Incubation time, 5

days.

Figure 5. Effect of red-beet betalain pigments on amyloid-β-induced paralysis of *C.*

elegans transgenic strain CL2006. (A) Experimental schedule of red-beet betalain

pigments treatments in *C. elegans.* Image of *C. elegans* transgenic strain CL2006

morphology taken with a light microscope on day 0 (left) and day 15 (right) of

adulthood. (B, C) The synchronized hermaphrodites were exposed to 2, 10, and 50 μM

of dextrin (B) and red-beet betalain pigments (C). The paralysis rate was scored three

times per week and is expressed as the percentage of not-paralyzed nematodes. ($n =$

109–119 nematodes/group). Differences in paralysis rates were tested for significance

using the log-rank test (*$p < 0.05$). Results are representative of two independent

experiments. (D) Morphological assessment of transgenic nematodes in the presence

and absence of red-beet betalain pigments using a light microscope. Beet indicate

red-beet betalain pigments.

A

betanin   isobetanin

B

betanin

isobetanin

A

Hatching

|——————————|————————————————————————————————→|
| Days of adulthood |

−4   0   5   10   15   20 (Days)

Not paralyzed

Paralyzed

Beet treatment

Hatching

|——————————|————————————————————————————————→|
| Days of adulthood |

−4   0   5   10   15   20 (Days)

B

% of not paralyzed

Days of adulthood

······ No dextrin
—— 2 μM dextrin
—— 10 μM dextrin
—— 50 μM dextrin

C

% of not paralyzed

Days of adulthood

······ No Beet
—— 2 μM Beet
—— 10 μM Beet
—— 50 μM Beet *

D

No betanin   50 μM Beet