CLN5 and CLN3 function as a complex to regulate endolysosome function

Running title: Endolysosome function of CLN3 and CLN5

Seda Yasa [1], Etienne Sauvageau [1], Graziana Modica [1], and Stephane Lefrancois [1, 2, #]

[1] Centre Armand-Frappier Santé Biotechnologie, Institut national de la recherche scientifique, Laval, Canada H7V 1B7
[2] Department of Anatomy and Cell Biology, McGill University, Montreal, Canada H3A 0C7 and Centre d'Excellence en Recherche sur les Maladies Orphelines - Fondation Courtois (CERMO-FC), Université du Québec à Montréal (UQAM), Montréal, Canada H2X 3Y7

To whom correspondence should be addressed: stephane.lefrancois@inrs.ca

Keywords: endosomes, CLN5, CLN3, retromer, sortilin, palmitoylation, Rab7A

Abstract

CLN5 is a soluble endolysosomal protein that regulates the itinerary of the lysosomal sorting receptor sortilin. Mutations in this protein cause neuronal ceroid lipofuscinosis, a rare neurodegenerative disorder, and have also been associated with Alzheimer's disease, suggesting functional defects in a common pathway. We previously found that depletion of CLN5 leads to dysfunctional retromer, resulting in the degradation of the lysosomal sorting receptor, sortilin. However, how a soluble lysosomal protein can modulate the function of a cytosolic protein is not known. In this work, we show that deletion of CLN5 not only results in retromer dysfunction, but also in impaired endolysosome fusion events. This results in delayed degradation of endocytic proteins and in defective autophagy. CLN5 modulates these various pathways by regulating downstream interactions between CLN3, an integral membrane protein, Rab7A and a subset of Rab7A effectors. Mutations in CLN3 are also a cause of neuronal ceroid lipofuscinosis. Our data supports a model where CLN3 and CLN5 function as an endolysosome complex regulating several endosomal functions.

Summary Statement

We have previously demonstrated that CLN3 is required for efficient endosome-to-trans Golgi Network (TGN) trafficking of sortilin by regulating retromer function. In this work, we show that CLN5, which interacts with CLN3, regulates retromer function by modulating key

interactions between CLN3, Rab7A, retromer, and sortilin. Therefore, CLN3 and CLN5 serve as endosomal switch regulating the itinerary of the lysosomal sorting receptors.

Introduction

The Neuronal Ceroid Lipofuscinoses (NCLs) are a group of inherited lysosomal diseases with over 430 mutations in 13 genetically distinct genes (*CLN1-8* and *CLN10-14*) (Mole and Cotman, 2015). Germline mutations in *CLN5,* resulting in either amino acid conversion (R112H, N192S, D279N) or truncations due to premature stop codons (W75X and Y392X), are causes of CLN5 disease. The truncation mutation, CLN5^{Y392X}, is the most common mutation in patients. This form of NCL has an early onset between the ages of 3 - 7, with a lifespan into the teenage years. Patients exhibit symptoms including retinopathy, motor disorders, epilepsy, and cognitive regression. The principal characteristics leading to the identification of NCL disorder is aberrant lysosomal function and excess accumulation of auto fluorescent ceroid lipopigments in neurons and peripheral tissues (Anderson et al., 2012).

CLN5 is a glycosylated endolysosomal protein that is translated as a type II integral membrane protein (residues 1 - 407), prior to being cleaved into a mature soluble form (93 - 407), which is then transported to endolysosomes (Jules et al., 2017). Recent work has suggested that CLN5 could function as a glycoside hydrolase, but endogenous targets have not been identified (Huber and Mathavarajah, 2018). Furthermore, CLN5 has been linked to the regulation of lysosomal pH (Best et al., 2017) and mitophagy (Doccini et al., 2020). To further our understanding of the this protein, using small interfering RNA (siRNA), we generated CLN5 knockdown (CLN5^{KD}) HeLA cells. In these cells, we found less membrane bound Rab7A (Mamo et al., 2012). Rab7A is a small GTPases that can bind to and recruit retromer, a protein complex that mediates endosome-to-trans Golgi Network (TGN) trafficking of the lysosomal sorting receptors cationic independent mannose 6-phosphate receptor (CI-MPR) and sortilin. Once recruited, retromer interacts with the cytosolic tails of the receptors to drive endosome-to-TGN retrieval (Canuel et al., 2008; Rojas et al., 2008; Seaman et al., 2009). In CLN5^{KD} HeLa cells, the lack of Rab7A recruitment resulted in the lysosomal degradation of CI-MPR and sortilin, and misrouting of the lysosomal enzyme cathepsin D, due to lack of retromer recruitment (Mamo et al., 2012). Beyond retromer recruitment, Rab7A also regulates the degradation of endocytic cargo such as the epidermal growth factor (EGF) receptor (EGFR) (Vanlandingham and Ceresa, 2009) by interacting with PLEKHM1 (McEwan et al., 2015), regulates organelle movement by in interacting with RILP (Pankiv et al., 2010; van der Kant et al., 2013), and regulates the late stages of autophagy by mediating autophagosome/lysosome fusion through HOPS and other tethering factors (Ao et al., 2014).

Recently, we have shown that CLN3 is also required in the endosome-to-TGN trafficking of the lysosomal sorting receptors (Yasa et al., 2020). Mutations in the gene encoding for CLN3 results in the juvenile variant of NCL (JNCL), commonly known as Batten disease. JNCL is the most common cause of childhood dementia, and has an age of onset between 5 to 10 years (Butz et al., 2019). CLN3 is a protein of 438 amino acids with six transmembrane domains, whose N- and C- terminal ends are located in the cytosol. This glycosylated protein localizes to endolysosomal membranes among other intracellular locations (Oetjen et al., 2016; Storch et al., 2004), can interact with other CLN proteins including CLN5, however the molecular function of these interactions are not known (Vesa et al., 2002). Unlike depletion of CLN5, deletion of CLN3 did not affect the localization or activation of Rab7A or its effector retromer. Rather, CLN3 is required for the efficient Rab7A interaction with retromer, and for the retromer/sortilin interaction, most likely acting as a scaffold protein (Yasa et al., 2020).

In this study, we generated CLN5 knockout (CLN5^{KO}) HeLa cells on the same parental cell line used to generate CLN3 knockout (CLN3^{KO}) and Rab7A knockout (Rab7A^{KO}) cells (Yasa et al., 2020). This knockout system has enabled us to study the effects of a CLN5 disease-causing mutation. We extended our studies to identify defects in endocytic degradation and autophagy. Importantly, we identified significant decreases in previously known interactions between Rab7A/retromer, sortilin/retromer, and sortilin/CLN3 in CLN5^{KO} cells (Yasa et al., 2020). Overall, our data suggest that CLN3 and CLN5 form a late endolysosome complex that modulates a subset of Rab7A functions. These finding provides insights into the pathogenic mechanisms observed in CLN3 and CLN5 disease.

Results

Retromer is not efficiently recruited to endosomal membranes in CLN5^{KO} HeLa cells

In our previous work conducted in CLN5^{KD} HeLa cells, we showed decreased membrane recruitment of Rab7A leading to a significant decrease in the membrane recruitment of its effector, retromer (Mamo et al., 2012). In order to study the the role of CLN5 disease-causing mutations, we generated CLN5^{KO} HeLa cells using CRISPR/Cas9. Our sequencing data and Western blot analysis confirms the deletion of CLN5 in this cell line (**Fig. S1A and B)**. To confirm our previous results obtained in CLN5^{KD} cells, we performed a membrane isolation experiment in our CLN5^{KO} HeLa cells as we have previously done (Mamo et al., 2012; Modica et al., 2017; Yasa et al., 2020), and compared the membrane distribution of Rab7A in wild-type and CLN5^{KO} HeLa cells (**Fig. 1A**). Our membrane separation was successful, as the integral membrane protein Lamp2 was found in the pellet fraction (P) containing membranes, and not in the supernatant fraction (S) (**Fig. 1A**). Quantification of 3 independent experiments showed that CLN5 deletion had no significant effect on Rab7A membrane distribution (**Fig. 1B**). We sought

to determine the effect of a CLN5 mutation known to cause human disease on Rab7A membrane recruitment. Thus, we engineered a CLN5 disease-causing mutation expression plasmid using site directed mutagenesis to generate CLN5^{Y392X}-HA, and expressed it in CLN5^{KO} HeLa cells (**Fig. 1A**). Rab7A was was not affected in CLN5^{KO} HeLa cells expressing CLN5-HA or expressing the disease causing mutation, CLN5^{Y392X}-HA (**Fig. 1B**). The small GTPase Rab7A regulates the spatiotemporal recruitment of retromer (Rojas et al., 2008; Seaman et al., 2009). Therefore, we repeated the membrane assay as above to test if CLN5 is required for membrane recruitment of retromer. This time we included Rab7A^{KO} HeLa cells as a control, as we previously demonstrated a reduction of retromer recruitment in these cells (Yasa et al., 2020). Lamp2 was used as a marker of the membrane fraction (P) (**Fig. 1A**). Compared to wild-type HeLa cells, we observed a significant decrease in retromer recruitment (as detected by the retromer subunit Vps26A) in CLN5^{KO} and Rab7A^{KO} HeLa cells (**Fig. 1A**). Quantification of 6 independent experiments found a reduction of 21.4% in the membrane recruitment of retromer in CLN5^{KO}, compared to a 31.4% decrease in recruitment in Rab7A^{KO} HeLa cells (**Fig. 1C**). These values are comparable to previous studies (Rojas et al., 2008; Seaman et al., 2009). This decrease in retromer membrane recruitment was rescued by expressing CLN5-HA in CLN5^{KO} cells, indicating the effect seen on Vps26A recruitment is specific to CLN5 deletion, and likely not an off-target effect (**Fig. 1A and C**). The expression of CLN5^{Y392X}-HA in CLN5^{KO} cells partially rescued retromer recruitment, but not to levels comparable as that of wild-type CLN5 (**Fig. 1C**).

The question remained as to how Rab7A was membrane bound, but not able to recruit retromer. In recent years, Rab7A has been shown to be phosphorylated (Francavilla et al., 2016; Heo et al., 2018; Lin et al., 2017), ubiquitinated (Song et al., 2016) and palmitoylated (Modica et al., 2017). These post-translational modifications have been demonstrated to regulate various functions of this small GTPase, often by modulating the interaction with its effectors. In particular, we have previously shown that Rab7A palmitoylation is required for the recruitment of retromer to endosomal membranes. While non-palmitoylatable Rab7A is still membrane bound, it does not efficiently interact with retromer, and is not capable of rescuing retromer membrane recruitment in Rab7A^{KO} HEK293 cells (Modica et al., 2017). Since Rab7A was still membrane bound in CLN5^{KO} HeLa cells, but retromer was not, we used Acyl-RAC to test if Rab7A palmitoylation is affected in these cells (**Fig. 1D**). We found that Rab7A palmitoylation is significantly decreased in CLN5^{KO} HeLa cells compared to wild-type cells (**Fig. 1E**).

CLN5 is required for efficient retromer interactions

Rab7A coordinates the spatiotemporal recruitment of retromer to endosomal membranes (Rojas et al., 2008; Seaman et al., 2009). Since we observed decreased retromer membrane recruitment along with decreased Rab7A palmitoylation, we expected to observe a weaker interaction between Rab7A and retromer in CLN5^{KO} HeLa cells. To test our hypothesis, we used bioluminescence resonance energy transfer (BRET), as we have previously done (Modica et al., 2017; Yasa et al., 2020). Compared to co-immunoprecipitation, BRET is performed in live cells, with proteins localized to their native environment. From BRET titration curves, the BRET$_{50}$ can be calculated, which is the value at which the concentration of the acceptor is required to obtain 50% of the maximal BRET signal (BRET$_{max}$). The BRET$_{50}$ is indicative of the propensity of a pair of protein to interact, as the smaller the BRET$_{50}$, the stronger the interaction. (Kobayashi et al., 2009; Mercier et al., 2002). The energy donor Renilla Luciferase II (RlucII) was fused at the N-terminus to wild-type Rab7A (RlucII-Rab7A), while the energy acceptor GFP10 was fused at the C-terminus of the retromer subunit Vps26. Neither of these tags appear to interfere with the function of the proteins, as we have previously shown that RlucII had little effect on the distribution or function of Rab7A (Modica et al., 2017), while we confirmed that Vps26A-GFP10 is efficiently integrated into the retromer trimer (Yasa et al., 2020). We have previously used BRET to study the Rab7A/Vps26A interaction (Modica et al., 2017). This interaction is specific as Rab7A did not interact with AP-1 subunits (a clathrin adaptor that has been localized to both the TGN and endosomes), while Vps26A did not interact with RlucII-Rab1a, which is localized to the Golgi apparatus (Modica et al., 2017). Wild-type and CLN5^{KO} HeLa cells were co-transfected with a constant amount of RlucII-Rab7A, and increasing amounts of Vps26A-GFP10 to generate BRET titration curves (**Fig. 2A, blue and red curves**). The BRET signal between RlucII-Rab7A and Vps26A-GFP10 rapidly increased with increasing amounts of expressed Vps26A-GFP10 until it reached saturation, suggesting a specific interaction. Compared to wild-type HeLa cells, we found a 10-fold increase in the BRET$_{50}$ value for Rab7A binding to Vps26A in CLN5^{KO} cells, suggesting a weaker interaction (**Fig. 2B**). To test the effect of the disease-causing mutation in CLN5 affected the Rab7A/retromer interaction, we expressed wild-type HA-CLN5 (**Figure 2A, green curve**) or HA-CLN5^{Y392X} (**Figure 2A, purple curve**) in CLN5^{KO} cells and generated BRET titration curves as above. While expressing HA-CLN5 rescued the Rab7/retromer interaction as the BRET$_{50}$ was similar to the BRET$_{50}$ in wild-type HeLa cells (**Fig. 2B**), expression of HA-CLN5^{Y392X} did not (**Fig. 2B**). Our previous work suggests that palmitoylation favours the association of Rab7A to a specific endosomal domain to optimize its interaction with retromer (Modica et al., 2017), without affecting the overall ability of the small GTPase to interact with this effector. As such, when the Rab7A/retromer interaction was analyzed with BRET, hence preserving cellular membranes, we observed a decreased interaction, while no

change in the interaction was observed in co-immunoprecipitation where cellular compartments are lost. To confirm if this was the case in CLN5[KO] HeLa cells, we performed co-immunoprecipitation between Rab7A and retromer in wild-type and CLN5[KO] HeLa cells (**Fig. S1C**). We found no change in the Rab7A/retromer interaction using this method, supporting our previous data.

CI-MPR and sortilin are known to interact with retromer, which is necessary for their endosome-to-TGN trafficking (Arighi et al., 2004; Canuel et al., 2008; Seaman, 2004). Using BRET, we have previously shown that nanoLuciferase-tagged Vps26A (Vps26A-nLuc) and YFP-tagged sortilin (sortilin-YFP) interact, and that this interaction is significantly weaker in CLN3[KO] cells compared to wild-type HeLa cells (Yasa et al., 2020). We tested the sortilin/retromer interaction in CLN5[KO] HeLa cells to determine if this protein plays a role in this interaction. Wild-type and CLN5[KO] HeLa cells were co-transfected with a constant amount of Vps26A-nLuc and increasing amounts of sortilin-YFP to generate BRET titration curves (**Fig. 2C, blue and red curves**). We found a significantly weakened interaction between retromer and sortilin in CLN5[KO] cells compared to wild-type HeLa cells, as we observed a 4 fold increase in the $BRET_{50}$ value (**Fig. 2D**). The retromer/sortilin interaction in CLN5[KO] HeLa cells could be rescued by expressing HA-CLN5 (**Fig. 2C, green curve**), as the $BRET_{50}$ value was similar to the one we calculated in wild-type HeLa cells. We next performed BRET experiments to determine the impact of the CLN5[Y392X] disease-causing mutation on the Vps26A/sortilin interaction. Expression of HA-CLN5[Y392X] in CLN5[KO] HeLa cells (**Fig. 2C, purple curve**) did not rescue the interaction as shown by the $BRET_{50}$ value (**Fig. 2D**). To confirm our BRET data, we immunoprecipitated endogenous Vps26A with Vps26A antibody, and blotted for endogenous sortilin (**Fig. S1C**). By co-immunoprecipitation, we observed no change in the Vps26A/sortilin interaction in CLN5[KO] cells, suggesting that the proteins can interact when the contribution of membrane distribution is not considered.

Sortilin is degraded in CLN5[KO] HeLa cells

In cells lacking functional retromer, sortilin does not efficiently recycle to the TGN, accumulates in late endosomes, and is subsequently degraded in lysosomes (Yasa et al., 2020). We have previously demonstrated the same phenotype in cells depleted of CLN5 (Mamo et al., 2012). Using our CLN5[KO] HeLa cells, we performed a cycloheximide chase experiment to determine receptor stability as we have previously done (Mamo et al., 2012; McCormick et al., 2008; Yasa et al., 2020). Wild-type, Rab7[KO] and CLN5[KO] HeLa cells were incubated with serum free medium containing 50 µg/ml cycloheximide and collected after 0, 3, and 6 hours of incubation. Western blot (Wb) analysis shows decreased levels of sortilin in CLN5[KO] HeLa cells at 3 and 6 hours compared to Rab7[KO] and wild-type HeLa cells (**Fig. 3A**). Transfecting CLN5-

HA in CLN5^{KO} cells rescued this phenotype (**Fig. 3A**). We next aimed to determine the impact of CLN5^{Y392X} on the stability of sortilin. Expression of CLN5^{Y392X}-HA in CLN5^{KO} cells did not rescue the degradation of sortilin (**Fig. 3A**). Actin staining was used as a loading control, while HA staining was used to demonstrate the expression level of HA tagged CLN5 constructs. Quantitative analysis of 3 independent cycloheximide chase experiments showed that compared to wild-type cells which had 77.8% and 75.7% of sortilin remaining at 3 and 6 hours respectively (**Fig. 3B**), sortilin was significantly degraded in CLN5^{KO} cells, as only 45.2% and 11% of sortilin remained after 3 and 6 hours (**Fig. 3B**). As expected, we observed no significant degradation in Rab7A^{KO} cells, which had 72.9% and 87.8% sortilin remaining at 3 and 6 hours respectively (**Fig. 3B**). Expressing CLN5-HA in CLN5^{KO} cells rescued the stability and hence recycling of sortilin, as it was no longer degraded and had protein levels remaining of 83.2% and 79% respectively at 3 and 6 hours, which is similar to wild-type HeLa cells (**Fig. 3B**). The degradation of sortilin remained significant in CLN5^{KO} cells expressing CLN5^{Y392X}-HA (20.4% and 20.5% remaining at 3 and 6 hours), suggesting that this disease-causing mutation did not rescue the knockout phenotype.

Since we observed the degradation of sortilin in CLN5^{KO} HeLa cells, we aimed to determine if this had an impact on lysosomal function. Cathepsin D is generated as a 53 kDa pro-cathepsin D (proCatD), which is processed to a 47 kDa intermediate form (iCatD), before being fully processed to a 31 kDa mature cathepsin D (mCatD) once it reaches the lysosome (Laurent-Matha et al., 2006). Depletion of Rab7A by siRNA affects this processing, resulting in increased amounts of the pro and intermediate forms (Rojas et al., 2008). Total cell lysate from wild-type, CLN5^{KO}, and Rab7A^{KO} HeLa cells was run on a SDS-PAGE and Western blotting was performed using anti-cathepsin D antibody (**Fig. 3C**). Compared to wild-type HeLa cells which had 24.6% pro-cathepsin (proCatD), 9.8% intermediate (iCatD) and 65.6% mature cathepsin D (mCatD), CLN5^{KO} HeLa had significantly increased iCatD (24.6%) and significantly reduced mCatD (48%), with similar levels of proCatD (27.4%). On the other hand, Rab7A^{KO} HeLa cells had increased levels of proCatD (37.6%) and iCatD (30.3%), and had decreased amounts of mCatD (32.1%) (**Fig. 3D**).

CLN5 is required for efficient CLN3 interactions

Recently, we showed that CLN3 functions as a scaffold protein to ensure the Rab7A/ retromer and retromer/sortilin interactions, which are sequentially required to regulate the endosome-to-TGN trafficking of sortilin (Yasa et al., 2020). In this current study, we demonstrated a role for CLN5 in regulating the stability of sortilin. However, the question remains how a soluble lysosomal protein, CLN5, can regulate the Rab7A/retromer and retromer/ sortilin interactions, which occur in the cytosol. In order to better understand the relationship of

CLN5 and CLN3 in this trafficking pathway, we wanted to determine the role of CLN5 in modulating CLN3 interactions. Previous studies have shown that CLN3 and CLN5 interact (Lyly et al., 2009; Vesa et al., 2002). We confirmed the CLN3/CLN5 interaction using co-immunoprecipitation, and tested the impact of a disease-causing mutation in CLN5 on this interaction (**Fig. S2A**). The disease-causing mutation we tested, $CLN5^{Y392X}$, had no impact on the CLN3/CLN5 interaction (**Fig. S2A**). The CLN3/Rab7A interaction has been previously shown using co-immunoprecipitation and BRET experiments (Uusi-Rauva et al., 2012; Yasa et al., 2020). To determine if CLN5 plays a role in this interaction, wild-type (**Fig. 4A, blue curve**) and $CLN5^{KO}$ HeLa cells (**Fig. 4A, red curve**) were co-transfected with a constant amount of RlucII-Rab7A and increasing amounts of GFP10-CLN3 to generate BRET titration curves. We extrapolated the $BRET_{50}$ for the interaction between CLN3 and Rab7A and found that the $BRET_{50}$ was 2.85 fold larger in $CLN5^{KO}$ HeLa cells compared to wild-type cells, indicating a weaker CLN3/Rab7A interaction in cells lacking CLN5 (**Fig. 4B**). Expressing HA-CLN5 in $CLN5^{KO}$ cells rescued the CLN3/Rab7A interaction (**Fig. 4A, green curve**) as the $BRET_{50}$ was similar to wild-type HeLa cells (**Fig. 4B**). Expressing $HA-CLN5^{Y392X}$ in $CLN5^{KO}$ cells had a partial rescue effect on the CLN3/Rab7A interaction (**Fig. 4A, purple curve**), but the $BRET_{50}$ was still more than 2 fold larger (**Fig. 4B**).

We previously demonstrated an interaction between CLN3 and retromer (Yasa et al., 2020). To test if CLN5 plays a role in modulating this interaction, we generated BRET titration curves using RlucII-CLN3 and Vps26A-GFP10 in wild-type (**Fig. 4C, blue curve**) and $CLN5^{KO}$ HeLa cells (**Fig. 4C, red curve**). We found a significantly weaker CLN3/retromer interaction in $CLN5^{KO}$ cells compared to wild-type HeLa cells as shown by the 4 fold increase of the $BRET_{50}$ in $CLN5^{KO}$ HeLa cells compared to wild-type cells (**Fig. 4D**). As expected, expressing HA-CLN5 in $CLN5^{KO}$ cells rescued the CLN3/retromer interaction (**Fig. 4B, green curve**) as the $BRET_{50}$ was similar to wild-type HeLA cells (**Fig. 4C**). Expressing $HA-CLN5^{Y392X}$ in $CLN5^{KO}$ HeLa cells had a partial rescue effect on the CLN3/retromer interaction (**Fig. 4C, purple curve**), but the $BRET_{50}$ was still more than 3 fold larger (**Fig. 4D**).

We have also shown an interaction between CLN3 and sortilin (Yasa et al., 2020) and between CLN5 and sortilin (Mamo et al., 2012). To test if a disease-causing mutation in CLN5 affected its ability to interact with sortilin, we performed a co-immunoprecipitation assay (**Fig. S2C**). Wild-type CLN5, as well as the disease-causing mutant tested, $CLN5^{Y392X}$, were able to interact with sortilin. We next generated BRET titration curves using RlucII-CLN3 and sortilin-YFP in wild-type (**Fig. 4E, blue curve**) and $CLN5^{KO}$ HeLa cells (**Fig. 4E, red curve**). We extrapolated the $BRET_{50}$ for the interaction between CLN3 and sortilin and found that the $BRET_{50}$ in $CLN5^{KO}$ HeLa cells was 6 fold larger than in wild-type cells, indicating a decreased interaction in $CLN5^{KO}$ cells (**Fig. 4F**). Expressing HA-CLN5 in $CLN5^{KO}$ HeLa cells rescued the

phenotype as the $BRET_{50}$ value was similar to wild-type cells (**Fig. 4F, green curve**), while expressing HA-CLN5[Y392X] (**Fig. 4F, purple curve**) did not. Although expression of CLN5[Y392X] had a partial rescue of the CLN3/sortilin interaction, the $BRET_{50}$ values were not restored to levels comparable to rescuing with wild-type CLN5 (**Fig. 4F**). To confirm our BRET results, we performed a co-immunoprecipitation experiment to determine the role of CLN5 in the modulating the CLN3/Rab7A, CLN3/retromer and sortilin/CLN3 interactions (**Fig. S2B**). We found that retromer (Vps26A), Rab7A and sortilin could co-immunoprecipitate with Flag-CLN3 in wild-type HeLa cells, but not in CLN5[KO] HeLa cells. The interaction was specific as expressing empty Flag vector did not co-immunoprecipitate retromer or Rab7A (**Fig. S2B**).

CLN5 is required for the efficient degradation of proteins following internalization

Another function of Rab7A is mediating key steps in the degradation of endocytic cargo (Ceresa and Bahr, 2006; Vanlandingham and Ceresa, 2009). Upon EGF stimulation, EGF receptor (EGFR) is internalized and can be either recycled to the cell surface, or degraded in lysosomes (Ceresa and Peterson, 2014). At least two Rab7A effectors have been implicated in EGFR degradation, RILP and PLEKHM1. Depletion of either of these proteins results in significant delays in the degradation kinetics of EGFR (Marwaha et al., 2017; McEwan et al., 2015; Progida et al., 2007). To determine whether CLN5 is implicated in this pathway, we investigated the degradation kinetics of EGFR in wild-type and CLN5[KO] HeLa cells. The cells were serum starved for 1 hour in the presence of cycloheximide and then stimulated with 100 ng/ml of EGF in the presence of cycloheximide for the indicated periods of time. The level of endogenous EGFR was determined by Western blot. Anti-actin staining was used as a loading control (**Fig. 5A**). In wild-type cells, EGFR degradation was observed after 15 minutes and quantification of 3 independent experiments found substantial degradation at 15 (34.3% remaining) and 120 minutes (9% remaining) (**Fig. 5B**). When we compared the degradation kinetics of EGFR in CLN5[KO] cells, we found significant delay at 15 (75.6% remaining) compared to wild-type cells, with no significant difference at 120 minutes (10.3% remaining) (**Fig. 5B**).

Next, we tested the degradation kinetics of Alexa-488 labeled EGF (EGF-488) using the same cell lines to confirm our EGFR degradation results. Following 2 hours of serum starvation, wild type (**Fig. 5C - E**) and CLN5[KO] (**Fig. 5F - H**) HeLa cells were incubated with 300 ng/ml EGF-488 for 30 minutes, washed and then chased for 0 (**Fig. 5C and F**), 15 (**Fig. 5D and G**) or 30 minutes (**Fig. 5E and H**). Images were acquired at random from the different conditions and the number of EGF-488 puncta from 35 cells per condition were counted using Image J as we have previously done (Yasa et al., 2020). At time 0 min, both wild-type and CLN5[KO] HeLa cells had comparable number of EGF-488 puncta (an average of 8.7 versus 9.7 respectively). After 15 minutes of chase, wild-type HeLa cells had an average of 5.4 puncta per cell, while CLN5[KO]

HeLa had on average 7.4 puncta per cell, an increase of 37% (**Fig. 5I**). After 30 minutes of chase, wild-type HeLa cells had an average of 3.4 puncta per cell, while CLN5[KO] HeLa had on average 3.8 puncta per cell (**Fig. 5I**).

In order to understand the mechanism behind the delayed EGF and EGFR degradation, we used BRET to determine if the Rab7A/PLEKHM1 or Rab7A/RILP interactions were affected (**Fig. 6A - D**). RILP and PLEKHM1 are implicated in the degradation of internalized cargo such as EGFR by either participating in the tethering of vesicles, as is the case for PLEKHM1 (McEwan et al., 2015), or by mediating the movement of vesicles, as is the case for RILP (Wijdeven et al., 2016). We generated BRET titration curves by transfecting a constant amount of RlucII-Rab7A and increasing amounts of PLEKHM1-GFP10 (**Fig. 6A**) or increasing amounts of RILP-GFP10 (**Fig. 6C**). We found no significant change in the interaction between PLEKHM1 and Rab7A in CLN5[KO] HeLa cells compared to wild-type HeLa cells as shown by the similar BRET$_{50}$ values (**Fig. 6B**). On the other hand, the interaction between RILP and Rab7A was significantly disrupted in CLN5[KO] cells, as the BRET$_{50}$ value was 3 fold higher for the Rab7A/ RILP interaction in CLN5[KO] cells compared to wild-type cells, suggesting a weaker interaction (**Fig. 6D**). Rab7A can interact with RILP to subsequently engage dynein motors for minus-end transport of lysosomes (Johansson et al., 2007; van der Kant et al., 2013). If this transport system is affected due to the decreased Rab7A/RILP interaction, we would expect dysfunctional minus-end movement of lysosomes following starvation and the induction of autophagy. We compared the positioning of lysosomes, by staining with the lysosomal membrane protein CD63, in wild-type and CLN5[KO] cells that were starved in EBSS for 3 hours or maintained in DMEM (**Fig. 6E - H**). In both wild-type (**Fig. 6E**) and CLN5[KO] (**Fig. 6F**) HeLa cells cultured in DMEM, CD63 positive lysosomes were distributed throughout the cells. In nutrient starved wild-type cells, lysosomes efficiently moved in a retrograde manner towards the perinuclear region of the cell (**Fig. 6G, white arrows**). However, while CLN5[KO] HeLa cells were able to move some lysosomes, they were not as efficient as wild-type HeLa cells (**Fig. 6H**). Quantification of 20 cells per condition showed that as a percentage of total CD63 fluorescence, CLN5[KO] cells had significantly less perinuclear lysosomes compared to wild-type HeLa cells (**Fig. 6I**).

A role for minus-end movement of lysosomes is to enable efficient fusion with autophagosomes, a step required for efficient autophagic degradation (Cabukusta and Neefjes, 2018). Previous studies have reported autophagic defects in cells depleted of CLN5 by RNAi (Adams et al., 2019), or CLN5 deficient cells (Doccini et al., 2020; Leinonen et al., 2017). Therefore, in CLN5[KO] HeLa cells, we would expect less autophagosome/lysosome fusions and defective autophagic flux. To test fusion ability, we expressed mCherry-LC3 and Lamp1-GFP in wild-type (**Fig. 7A - C**), CLN5[KO] (**Fig. 7D - F**) and Rab7A[KO] HeLa cells (**Fig. 7G - I**). LC3 is a marker of autophagosomes, while Lamp1 is a marker of lysosomes. Upon the initiation of

autophagy by starvation, LC3-positive autophagosomes fuse with Lamp1-positive lysosomes, forming autolysosomes in order to degrade material (Nakamura and Yoshimori, 2017). Cells were nutrient starved for 3 hours and the co-localization of mCherry-LC3 and Lamp1-GFP was determined. Quantification of the co-localization using Pearson's coefficient showed a high level of co-localization of mCherry-LC3 and Lamp1-GFP in wild-type HeLa cells, suggesting that autophagosomes were fusing with lysosomes (**Fig. 7J**). Rab7A participates not only in the movement of lysosomes, but also in the fusion process (Birgisdottir and Johansen, 2020). As such, we would expect significantly less co-localization in Rab7A[KO] HeLa cells. Compared to wild-type cells, the co-localization between LC3 and Lamp1 was significantly reduced in Rab7A[KO] HeLa cells (**Fig. 7J**). We also found significantly less co-localization between LC3 and Lamp1 in CLN5[KO] HeLa cells (**Fig. 7J**), suggesting that the deficient retrograde movement of lysosomes was preventing fusion between autophagosomes and lysosomes.

Next, we used a tandem LC3 probe (mTagRFP-mWasabi-LC3) to determine autophagic flux (Zhou et al., 2012). mWasabi is pH sensitive (more sensitive compared to EGFP), while mTagRFP is not. If cells can proceed with autophagy, the fusion of autophagosomes with lysosomes quenches the mWasabi so red is observed, while late blocks in autophagy (no autophagosome fusion with lysosomes) results in an increase in yellow puncta, with a corresponding decrease in red signal, since mWasabi is not quenched. We expressed mTagRFP-mWasabi-LC3 in wild-type (**Fig. 8K**), CLN5[KO] (**Fig. 8L**) and Rab7A[KO] HeLa cells (**Fig. 8M**), and cells were starved in EBSS for 3 hours. In wild-type cells, autophagosome/lysosome fusion occurs, resulting in a red signal as the mWasabi is quenched (**Fig. 8K and N**). In CLN5[KO] HeLa cells, fusion is deficient, and therefore mWasabi is not exposed to the acidic pH of the lysosomal lumen and no quenching occurs. Therefore, the red signal is significantly decreased (**Fig. 8L and N**). Since Rab7A is required for efficient fusion, a similar observation was made in Rab7A[KO] HeLa cells as expected (**Fig. 8M and N**).

Discussion

CLN5 is a soluble endolysosomal protein whose function remains poorly understood. Mutations in this protein cause a rare neurodegenerative disease that affects children, so understanding the function of CLN5 could lead to new therapeutic targets. Work using the model organism *Dictyostelium discoideum* showed that CLN5 has glycoside hydrolase activity (Huber and Mathavarajah, 2018), and previous reports found that CLN5 regulates lysosomal pH (Best et al., 2017) and autophagy (Adams et al., 2019; Leinonen et al., 2017). More recent work has implicated CLN5 in mitochondrial function (Doccini et al., 2020). Using siRNA (CLN5[KD]), we previously demonstrated that CLN5 regulates the stability of the lysosomal sorting receptor sortilin by enabling its endosome-to-TGN trafficking (Mamo et al., 2012). However, using this KD

system, we never tested the effects of disease-causing mutations on this trafficking pathway, or whether CLN5 modulated other Rab7A functions. Furthermore, it is not known how a soluble endolysosomal protein can regulate protein-protein interactions in the cytosol. In this current work, we generated a CLN5 knockout (CLN5[KO]) HeLa cell line that can be rescued with wild-type or mutant CLN5. Using this tool, we have confirmed our previous work and expanded our understanding of the function of this protein.

In our CLN5[KD] HeLa cells, we observed less membrane bound Rab7A and retromer (Mamo et al., 2012). We found less membrane bound retromer in our CLN5[KO] HeLa cells, which was rescued by expressing wild-type CLN5, but not the disease-causing mutant (CLN5[Y392X]). Although retromer recruitment was deficient in CLN5[KO] HeLa cells, Rab7A was still membrane bound. This raised two points. Why the discrepancy between the KD and KO models, and how could Rab7A be membrane bound but not recruit retromer? We hypothesize that chronic loss of CLN5 in the KO model results in the cell overcoming the effect, while acute depletion of CLN5 by siRNA does not allow time for the cell to compensate. We have previously shown that Rab7A palmitoylation is required for the efficient recruitment of retromer to endosomal membranes, but is not required for Rab7A membrane localization (Modica et al., 2017). We found that Rab7A palmitoylation is significantly decreased in CLN5[KO] HeLa cells, providing an explanation for the decrease in the Rab7A/retromer interaction as measured in live cells by BRET, and the subsequent lack of retromer recruitment. Palmitoylation of Rab7A is not required to modulate the Rab7A/retromer interaction *per se*, but serves to localize Rab7A to specific endosomal membrane domains to favour the interaction (Modica et al., 2017). When membranes are disrupted such as in co-immunoprecipitation, non-palmitoylatable Rab7A can interact with retromer (Modica et al., 2017). As such, it is not surprising that Rab7A and retromer interacted by co-immunoprecipitation in CLN5[KO] HeLa cells.

The recruitment of retromer to endosomal membranes enables the endosome-to-TGN retrieval of the lysosomal sorting receptor sortilin (Canuel et al., 2008). The sortilin/retromer interaction was perturbed in CLN5[KO] HeLa cells compared to wild-type HeLa cells. While the expression of wild-type CLN5 in CLN5[KO] HeLa cells rescued the sortilin/retromer interaction, the expression CLN5[Y392X] did not. This would suggest that the residues at the C-terminal end of CLN5 are required for the sortilin/retromer interaction, however how those residues modulate this interaction is yet to be determined. The decrease in the sortilin/retromer interaction observed in the CLN5[KO] HeLa cells should result in the lysosomal degradation of the lysosomal sorting receptor. As in the CLN5[KD] HeLa cells (Mamo et al., 2012), sortilin is degraded in CLN5[KO] HeLa cells. Expression of wild-type CLN5 rescued this degradation, while expressing the disease-causing mutant CLN5[Y392X] did not rescue. Taken together, these observations point to a crucial role of CLN5 in modulating Rab7A-dependent retromer recruitment and function.

Indeed by modulating the palmitoylation level of the small GTPase, CLN5 establishes the conditions for an optimal Rab7/retromer interaction, hence efficient retromer/sortilin binding and recycling to the TGN, which in turns translates in proper lysosomal function.

In our recently published paper, we demonstrated that CLN3 functions as a scaffold to ensure efficient protein-protein interactions. In that study, we demonstrated decreased Rab7A/ retromer and retromer/sortilin interactions in CLN3[KO] HeLa cells, which consequently lead to the degradation of sortilin (Yasa et al., 2020). We found similar results in our CLN5[KO] HeLa cells. Furthermore, we found that CLN5 is required for the CLN3/retromer and CLN3/sortilin interactions. Although wild-type CLN5 rescued the CLN3/sortilin interaction in CLN5[KO] HeLa cells, the expression of the disease-causing mutant did not. Based on this data, we propose a model where CLN3 and CLN5 function as an endosomal complex required for the efficient endosome-to-TGN trafficking of sortilin. This inefficient recycling results in defective lysosomal function, which is a hallmark of CLN5 disease. Our results suggest a potential pathogenic mechanism for CLN5 disease.

We next asked if other Rab7A mediated pathways are affected in CLN5[KO] HeLa cells. We found significant delays in the degradation of both EGFR and EGF. This delayed degradation could be due to less efficient lysosomal function, or due to decreased fusion of endocytic cargo with lysosomes, a mechanism also modulated by Rab7A. The Rab7A effector RILP is implicated in the degradation of internalized cargo such as EGFR by mediating the movement of vesicles (Wijdeven et al., 2016), while the Rab7A effector PLEKHM1 is a tether required for fusion in both endocytic degradation and autophagy (McEwan et al., 2015). In CLN5[KO] HeLa cells, we found no changes to the Rab7A/PLEKMH1 interaction, but we found a decreased Rab7A/RILP interaction. This resulted in deficient movement of CD63 positive lysosomes toward the perinuclear region, an important step in the autophagic process. Indeed, we observed less co-localization of LC3II containing autophagosomes with Lamp1 positive lysosomes upon starvation in CLN5[KO] HeLa cells, and less autophagic flux as demonstrated using a tandem LC3 probe (mTagRFP-mWasabi-LC3). While decreased lysosomal function plays a role in the pathogenic mechanism of CLN5 disease, defective endocytic degradation and autophagy are most likely due to defective lysosomal movement and thus decreased fusion events.

In conclusion, in this study we demonstrate a role for CLN5, together with CLN3, acting as a complex to regulate endosome-to-TGN trafficking. While CLN3 behaves like an interaction platform, CLN5 ensures those CLN3 interactions, which are vital for lysosomal functioning are maintained and regulated. These findings enlighten the molecular mechanisms behind NCL pathogenesis caused by defected CLN5 and CLN3 proteins.

Materials and Methods

Plasmids and Mutagenesis

RlucII-CLN3, GFP10-CLN3, Vps26A-nLuc, PLEKHM1-GFP10, RlucII-Rab7A, Vps26A-GFP10, RILP-GFP10 and sortilin-myc were previously described (Lefrancois et al., 2003; Modica et al., 2017; Yasa et al., 2020). The various CLN5 mutants were engineered using site-directed mutagenesis from the previously described HA-CLN5 and CLN5-HA constructs (Jules et al., 2017). Sortilin-YFP was a generous gift from Dr. Makoto Kanzaki, Tohoku University. mCherry-LC3 was a generous gift from Dr. Peter McCormick, Queen Mary University. Lamp1-GFP was a generous gift from Dr. Juan Bonifacino, NICHD, NIH. mTagRFP-mWasabi-LC3 was a generous gift from Jian Lin, Peking University.

Antibodies

The following mouse monoclonal antibodies were used: anti-actin (Wb: 1:3000, BD Biosciences 612657); anti-Lamp2 (Wb: 1:500, Abcam ab25631); anti-HA (Wb: 1:1000, Cedarlane Labs 901503); anti-myc (Wb: 1:1000, ThermoFisher Scientific LS132500), anti-Cathepsin D (Wb: 1:100, Sigma-Aldrich, IM03); anti-Flag (Wb: 1:1000, Sigma-Aldrich F1804). The following rabbit monoclonal antibodies were used: anti-Rab7A (Wb: 1:1000, Cell Signalling Technology D95F2); anti-EGFR (Wb: 1:1000, Abcam ab52894); anti-CLN5 (Wb: 1:1000 Abcam ab1700899). The following rabbit polyclonal antibodies were used: anti-Vps26A (Wb: 1:2000, Abcam ab23892); anti-FLAG (Wb: 1:1000, BioLegend, 902401); anti-sortilin (Wb: 1 µg/ml, Abcam ab16640).

Cell culture and transient transfections

HeLa cells were cultured in Dulbecco's modified Eagle's medium (DMEM) supplemented with 2mM L-Glutamine, 100 U/ml penicillin, 100g/ml streptomycin and 10% FBS (Wisent Bioproducts, St-Bruno, QC) at 37 °C in a humidified chamber at 95% air and 5% CO_2. Cells were seeded at a density of 2 X 10^5/well for 12 well plates and 5 X 10^5/well for 6 well plates 24 hours prior to transfection. Transfections were performed with polyethylenimine (PEI) (ThermoFisher Scientific, Ottawa, ON). Briefly, solution 1 was prepared by diluting plasmid into Opti-MEM (ThermoFisher Scientific). Solution 2 was prepared by diluting PEI (1 µg/µl) into Opti-MEM in a ration of 1:3 with the DNA to be transfected. After a 5 min incubation, the two solutions were mixed, vortexed for 3 s, incubated at room temperature (RT) for 15 min and subsequently added to the cells.

CRISPR/Cas9 Editing

HeLa cells were transfected with an all-in-one CRISPR/Cas9 plasmid for CLN5 (plasmid number HCP202087-CG01-1-B, Genecopoeia, Rockville, MD). 72 hours post-transfection, cells were treated with 1mg/ml Geneticin (ThermoFisher Scientific) for 1 week. Limited dilution was performed to isolate single cells, which were allowed to grow for 2 weeks. Western blotting was used to identify CLN5[KO] cells. Genomic DNA extracted form CLN5[KO] HeLa cells was sequenced to confirm CLN5[KO] cells. Rab7A[KO] HeLa cells were previously described (Yasa et al., 2020).

BRET titration experiments

HeLa cells were seeded in 12-well plates and transfected with the indicated plasmids. 48 hours post transfection, cells were washed in PBS, detached with 5mM EDTA in PBS and collected in 500 µl of PBS. Cells were transferred to opaque 96-well plates (VWR Canada, Mississauga, ON) in triplicates. Total fluorescence was first measured with the Tecan Infinite M1000 Pro plate reader (Tecan Group Ltd., Mannedorf, Switzerland) with the excitation and emission set at 400 nm and 510 nm respectively for BRET[2] and 500 nm and 530 nm for BRET[1]. The BRET[2] substrate coelenterazine 400a and BRET[1] substrate h-coelenterazine were then added to all wells (5 µM final concentration) and the BRET[2] or BRET[1] signals measured 2 min later. The BRET signals were calculated as a ratio of the light emitted at 525 ± 15 nm over the light emitted at 410 ± 40 nm. The BRET$_{net}$ signals were calculated as the difference between the BRET signal in cells expressing both fluorescence and luminescence constructs and the BRET signal from cells where only the luminescence fused construct was expressed.

Western blotting

Cell were detached using 5mM EDTA in PBS, washed in 1X PBS and collected by centrifugation. TNE buffer (150 mM NaCl, 50 mM Tris, pH 7.5, 2 mM EDTA, 0.5% Triton X-100 and protease inhibitor cocktail) was used to lyse cells by incubating them for 30 minutes on ice. Lysates were centrifuged at high speed for 10 minutes and the supernatants (whole cell lysate) were collected to be analyzed by Western blotting. Samples were mixed with sample buffer 3X to obtain a final concentration of 1X (62.5 mM Tris-HCl pH 6.5, 2.5% SDS, 10% glycerol, 0.01% bromophenol blue). Samples were incubated at 95°C for 5 minutes and resolved on SDS-PAGE followed by wet-transfer to nitrocellulose membranes. Detection was done by immunoblotting using the indicated antibodies.

Membrane separation assay

24hrs after transfection, cells were collected in 5mM EDTA in PBS. The cells were subsequently snap frozen in liquid nitrogen and allowed to thaw at RT for 5min. The cells were

resuspended in Buffer 1 (0.1 M Mes-NaOH pH 6.5, 1 mM MgAc, 0.5 mM EGTA, 200 M sodium orthovanadate, 0.2 M sucrose) and centrifuged at 10,000 g for 5 minutes at 4°C. The supernatants containing the cytosolic proteins (S, soluble fraction) were collected. The remaining pellet was resuspended in Buffer 2 (50 mM Tris, 150 mM NaCl, 1 mM EDTA, 0.1% SDS, 1% Triton X-100) and centrifuged at 10,000 g for 5 minutes at 4°C. Samples were loaded into SDS-PAGE gels in equal volumes. Fiji was used to quantify the intensity of the bands (Schindelin et al., 2012). The intensity of each fraction was calculated and divided by the total intensity to determine the distribution of proteins.

Cycloheximide chase

Wild-type, CLN5[KO], Rab7A[KO] HeLa cells were seeded in 6-well plates the day prior to transfection. 500 ng of HA fused wild-type and mutation harbouring CLN5 were transfected into CLN5[KO] HeLa cells. 48 hours after transfection, cells were treated with 50 µg/ml of cycloheximide in Opti-MEM. Lysates were collected at 0, 3 or 6 hour-time points and run on 10% SDS-PAGE gels. Fiji was used to quantify the intensity of the bands (Schindelin et al., 2012). All protein levels were standardized to the actin loading control. Amount of remaining protein is expressed as a percentage of the 0 time point in each group.

EGFR degradation assay

Wild-type, CLN5[KO] and Rab7A[KO] cells were seeded in 6-well plates the day before the assay. In order to prevent *de novo* synthesis EGFR during EGF stimulation, cells were treated with 50 µg/ml cycloheximide in Opti-MEM for 1 hour. Stimulation of EGF was performed with 100 ng/ml EGF in Opti-MEM containing 50 µg/ml of cycloheximide. Cell lysates were collected as indicated above and Western blotting was performed. Fiji was used to quantify the intensity of the bands (Schindelin et al., 2012). All protein levels were standardized to the actin loading control. Amount of remaining protein is expressed as a percentage of the 0 time point in each group.

EGF-488 pulse-chase

Wild-type and CLN5[KO] HeLa cells were seeded on coverslips the day before the experiment. Cells were serum-starved in Opti-Mem for 1 h followed by a 30-min pulse of 300 ng/ml of EGF-488. Cells were then washed with PBS, fixed in 4% paraformaldehyde at 0 and 30 min and mounted onto slide using Fluoromount-G (ThermoFisher Scientific). The coverslips were sealed using nail polish. Cells were imaged using fluorescence microscope using a 63X objective. The number of puncta per cell was counted using Fiji. Briefly, the images were split into their individual channels. Using the green channel, the threshold was adjusted to 0,93% in

order to get signal only from the puncta. The number of puncta counted from an image was divided by the number of nuclei within that image, which gave us the number of puncta per cell. 50 cells per condition for each time point were analyzed.

Acyl-RAC to isolate palmitoylated proteins

The isolation of palmitoylated protein was adapted from Modica et al., 2017. Briefly, cells were lysed in TNE (150 mM NaCl, 50 mM Tris, pH 7.5, 2 mM EDTA, 0.5% Triton X-100 and protease inhibitor cocktail) supplemented with 50mM N-Ethylmaleimide (NEM) and incubated 30 minutes in rotating wheel at 4°C. Samples were centrifuged 10 minutes at 10000g at 4°C and the collected supernatants were incubated 2 hours at RT on a rotating wheel. Samples were then precipitated over night with two volumes of cold acetone at -20°C to remove NEM. After washing with cold acetone, the pellets were resuspended in binding buffer (100 mM HEPES, 1 mM EDTA, 1% SDS) with 250 mM hydroxylamine (Sigma-Aldrich) (NH_2OH) pH7.5 to cleave palmitate residues off of proteins. Control samples were resuspended in binding buffer containing 250mM NaCl. When the pellets were completely resuspended, water-swollen thiopropyl sepharose 6B beads (GE Healthcare Life Sciences, Mississauga, ON) were added and samples were incubated 2 hrs at RT on rotating wheel. Beads were then washed 4 times with binding buffer and captured proteins were eluted with sample buffer containing 100mM DTT.

Autophagy assays

Cells were seeded on 6-well plates containing glass coverslips. The next day, the cells were transfected with the indicated plasmids. 48 hours post-transfection cells, the cells were starved in EBSS for 3 hours. Then, the coverslips were washed with PBS and fixed with 4% paraformaldehyde in PBS for 12 min at room temperature and washed twice with PBS. Fixed cells on the coverslips are mounted on cover slides with Fluoromount G (Thermo Fisher Scientific). Images were acquired using a fluorescence microscope using a 63X objective. Quantification of the fluorescence to analyze the colocalization of LC3 and Lamp1 was performed using Fiji software by splitting the red and green channels and manually outlining the cells to obtain the region of interest (ROI). Coloc2 was performed to determine the Pearson's coefficient. Red (LC3) in green (Lamp1) was chosen together with the bisection option for the threshold regression. Quantification of the fluorescence to analyze the quenching of the green signal was performed using the Fiji software. Red and green channels were split, and the cells manually outlined to get the ROI. Red and green fluorescence total intensities coming from each selected cells are analyzed. For autophagic flux comparison, intensity of the red signal is divided by the green signal for each cell.

Statistics

Statistical analysis was performed using GraphPad Prism Version 7. The statistical tests used are described in the corresponding figure legends.

Acknowledgements

We would like to thank Michel Bouvier (IRIC and University de Montreal), Regis Grailhe, (Pasteur Institute Korea), Paul Odgren (University of Massachusetts Medical School, Addgene #73592), Makoto Kanzaki (Tohoku University), Takeshi Nakamura (Tokyo University of Science) , Peter J. McCormick (Queen Mary University), Juan S. Bonifacino (NICHD, NIH), Jian Lin (Peking University) for providing plasmids. We would like to thank Guido Hermey (Institute for Molecular and Cellular Cognition, Center for Molecular Neurobiology Hamburg, University Medical Center Hamburg-Eppendorf) for providing the parental HeLa cells.

Competing interests

The authors have no financial or competing interests to declare.

Author Contributions

Conceptualization: S.Y., G.M., E.S., S.L.; Methodology: S.Y., G.M., E.S., S.L.; Validation: S.L.; Formal analysis: S.Y., G.M., E.S., S.L.; Investigation: S.Y., G.M., E.S.,; Writing - original draft: S.Y., S.L.; Writing - review & editing: S.Y., G.M., E.S., S.L.; Supervision: S.L.; Project administration: S.L.; Funding acquisition: S.L.

Funding

This work was supported by the Joint Programme in Neurodegenerative Diseases Grant (Neuronode), the Canadian Institutes for Health Research (ENG-155186), and the Canadian Foundation for Innovation (35258) to SL. SY and GM are supported by a scholarship from Fond de recherche du Quebec - Santé.

Figure Legends

Figure 1. Rab7A palmitoylation is reduced in CLN5^{KO} HeLa cells affecting retromer recruitment

(**A**) A membrane separation assay was performed on wild-type, Rab7A^{KO}, CLN5^{KO}, and CLN5^{KO} HeLa cells expressing CLN5-HA or CLN5^{Y392X}-HA. Western blotting (Wb) was performed with anti-Rab7A, anti-Vps26A, anti-Lamp2 (as a marker of the membrane fraction) and anti-HA (to identify cells expressing CLN5-HA) antibodies. S; supernatant fraction, P; pellet fraction. (**B**) Quantification of 3 separate membrane isolation assay experiments for Rab7A distribution. Data

is shown as mean±s.d.; ns, not significant; two-way ANOVA followed by Tukey's post hoc test. (**C**) Quantification of 3 separate membrane isolation assay experiments for Vps26A distribution. Data is shown as mean±s.d.; ns, not significant; *P≤0.01, **P≤0.01,****P≤0.0001, two-way ANOVA followed by Tukey's post hoc test (**D**) Whole cell lysate (WCL) from wild-type and CLN5[KO] HeLa cells were subjected to Acyl-RAC analysis to determine the palmitoylation status of Rab7A. NH_2OH: hydroxylamine, PD: pull-down (**E**) Quantification of 3 separate Acyl-RAC assay experiments. Data is shown as mean±s.d.; **P≤0.01, Student's *t*-test.

Figure 2. Weaker Rab7A/retromer and retromer/sortilin interactions in CLN5[KO] HeLa cells
(**A**) Wild-type, CLN5[KO] and CLN5[KO] HeLa rescued with wild-type HA-CLN5 or HA-CLN5[Y392X] were transfected with a constant amount of RlucII-Rab7A and increasing amounts of Vps26A-GFP10 to generate BRET titration curves. BRET signals are plotted as a function of the ratio between the GFP10 fluorescence over RlucII luminescence. (**B**) $BRET_{50}$ was extrapolated from 3 independent experiments. Data is shown as mean±s.d.; ns, not significant, ***P≤0.001, ****P≤0.0001, one-way ANOVA followed by Tukey's post hoc test (**C**) Wild-type, CLN5[KO] and CLN5[KO] HeLa rescued with wild-type HA-CLN5 or HA-CLN5[Y392X] were transfected with a constant amount of Vps26A-nLuc and increasing amounts of sortilin-YFP to generate BRET titration curves. BRET signals are plotted as a function of the ratio between the YFP fluorescence over nLuc luminescence. (**D**) Quantification of 3 independent experiments. Data is shown as mean±s.d.; ns, not significant; **P≤0.01, ***P≤0.001, one-way ANOVA followed by Tukey's post hoc test

Figure 3. The lysosomal sorting receptor sortilin is degraded in CLN5[KO] HeLa cells
(**A**) Wild-type, Rab7A[KO], CLN5[KO], and CLN5[KO] HeLa cells expressing CLN5-HA or CLN5[Y392X]-HA were treated with 50 µg/ml of cycloheximide in serum free media for the indicated times. Whole cell lysate was run on a SDS-PAGE and Western blotting (Wb) was performed with anti-sortilin, anti-actin and anti-HA antibodies. (**B**) Quantification of 3 independent experiments. Data is shown as mean±s.d.; ns, not significant; **P≤0.01,****P≤0.0001, two-way ANOVA followed by Tukey's post hoc test. (**C**) Whole cell lysate from wild-type, CLN5KO and Rab7AKO HeLa cells was resolved by SDS-PAGE and Western blotting (Wb) was performed using anti-cathepsin D antibody. (**D**) Quantification of pro-cathepsin D (proCatD, 53 kDa), intermediate cathepsin D (iCatD, 48 kDa) and mature cathepsin D (mCatD, 31 kDa). The amount of each is expressed as a percentage of total cathepsin D. Data are mean±s.d.; ns, not significant; **P≤0.01,***P≤0.001, two-way ANOVA followed by Tukey's post hoc test.

Figure 4. CLN5 modulates CLN3 interactions

(**A**) Wild-type, CLN5[KO] and CLN5[KO] HeLa rescued with wild-type HA-CLN5 or HA-CLN5[Y392X] were transfected with a constant amount of RlucII-Rab7A and increasing amounts of GFP10-CLN3 to generate BRET titration curves. BRET signals are plotted as a function of the ratio between the GFP10 fluorescence over RlucII luminescence. (**B**) $BRET_{50}$ was extrapolated from 3 independent experiments. Data is shown as mean±s.d.; ns, not significant, **$P \leq 0.01$, ****$P \leq 0.0001$, one-way ANOVA followed by Tukey's post hoc test. (**C**) Wild-type, CLN5[KO] and CLN5[KO] HeLa rescued with wild-type HA-CLN5 or HA-CLN5[Y392X] were transfected with a constant amount of RlucII-CLN3 and increasing amounts of Vps26A-GFP10 to generate BRET titration curves. BRET signals are plotted as a function of the ratio between the GFP10 fluorescence over RlucII luminescence. (**D**) $BRET_{50}$ was extrapolated from 3 independent experiments. Data is shown as mean±s.d.; ns, not significant, **$P \leq 0.01$, ***$P \leq 0.0001$, one-way ANOVA followed by Tukey's post hoc test. (**E**) Wild-type, CLN5[KO] and CLN5[KO] HeLa rescued with wild-type HA-CLN5 or HA-CLN5[Y392X] were transfected with a constant amount of RlucII-CLN3 and increasing amounts of sortilin-YFP to generate BRET titration curves. BRET signals are plotted as a function of the ratio between the YFP fluorescence over RlucII luminescence. (**F**) Quantification of 3 independent experiments. Data is shown as mean±s.d.; ns, not significant, **$P \leq 0.01$, ****$P \leq 0.0001$, one-way ANOVA followed by Tukey's post hoc test

Figure 5. EGF and EGFR degradation is delayed in CLN5[KO] cells

(**A**) Wild-type and CLN5[KO] HeLa cells were incubated with 50 µg/ml cycloheximide for 1 h and subsequently treated with 100 ng/ml EGF in Opti-MEM for 0, 15 and 120 min. Whole cell lysate was then resolved by SDS-PAGE and a Western blot (Wb) was performed using anti-EGFR antibody. Anti-actin staining was used as a loading control. (**B**) Quantification of the remaining EGFR as detected in A was performed in 3 independent experiments. Data are mean±s.d. ns, not significant; **$P \leq 0.01$; one-way ANOVA followed by Tukey's post hoc test. Scale bar: 10 µm. (**C - H**) Wild-type (**C - E**) and CLN5[KO] (**F - H**) HeLa cells were grown on coverslips and incubated with 300 ng/ml EGF-488 for 0, 15 or 30 min. The cells were fixed with 4% PFA for 12 min, followed by staining with DAPI to visualize the nucleus. Images were taken on a Zeiss Fluorescence microscope using a 63× objective. Stars indicate remaining EGF-488. (**I**) EGF-488 (green puncta) were counted using ImageJ in 50 cells per condition. The results shown are the average number of puncta per cell per condition. Data are mean±s.d. ns, not significant; *$P \leq 0.05$; one-way ANOVA followed by Tukey's post hoc test. Scale bar: 10 µm.

Figure 6. Retrograde transport of lysosomes is deficient in CLN5^KO HeLa cells

(**A**) Wild-type and CLN5^KO HeLa cells were transfected with a constant amount of RlucII-Rab7A and increasing amounts of PLEKHM1-GFP10 to generate BRET titration curves. BRET signals are plotted as a function of the ratio between the GFP10 fluorescence over RlucII luminescence. (**B**) BRET$_{50}$ was extrapolated from 3 independent experiments. Data is shown as mean±s.d.; ns, not significant. (**C**) Wild-type and CLN5^KO HeLa cells were transfected with a constant amount of RlucII-Rab7A and increasing amounts of RILP-GFP10 to generate BRET titration curves. BRET signals are plotted as a function of the ratio between the GFP10 fluorescence over RlucII luminescence. (**D**) BRET$_{50}$ was extrapolated from 3 independent experiments. Data is shown as mean±s.d.; **P≤0.01, Student's *t*-test. (**E - H**) Wild-type (**E and G**) and CLN5^KO (F and H) HeLa cells were cultured in standard DMEM (**E and F**) or starved (**G and H**) for 3 hours in EBSS. The cells were subsequently immunostained with anti-CD63 antibody. The cells were then fixed with 4% PFA for 12 min, followed by staining with DAPI to visualize the nucleus. Images were taken on a Zeiss Fluorescence microscope using a 63× objective. Scale bar: 10 µm. (**I**) Fluorescence intensity of CD63 in the perinuclear region and total cellular CD63 fluorescence was determined using ImageJ. Results shown are the ratio of perinuclear divided by total CD63 fluorescence. Data is shown as mean±s.d.; **P≤0.01, Student's *t*-test.

Figure 7. Lack of fusion of autophagosomes to lysosomes in CLN5^KO HeLa cells

(**A - I**) mCherry-LC3 and Lamp1-GFP were co-transfected into wild-type (**A - C**), CLN5^KO (**D - F**) and Rab7A^KO (**G - I**) HeLa cells and imaged on a Zeiss Fluorescence microscope using a 63× objective. Representative images are shown. Scale bar = 10 µm. (**J**) Quantification of the fluorescence to determine the Pearson's coefficient of mCherry-LC3 and Lamp1-GFP in wild-type, CLN5^KO and Rab7A^KO HeLa cells. **P≤0.01; ****P≤0.0001; one-way ANOVA followed by Tukey's post hoc test. (**K - M**) mTagRFP-mWasabi-LC3 was transfected into wild-type (K), CLN5^KO (L) and Rab7A^KO (M) HeLa cells. Representative images are shown. Scale bar: 10 µm (N) Quantification of mCherry divided by mWasabi was calculated. ****P≤0.0001; one-way ANOVA followed by Tukey's post hoc test.

Bibliography

Adams, J., M. Feuerborn, J.A. Molina, A.R. Wilden, B. Adhikari, T. Budden, and S.Y. Lee. 2019. Autophagy-lysosome pathway alterations and alpha-synuclein up-regulation in the subtype of neuronal ceroid lipofuscinosis, CLN5 disease. *Sci Rep*. 9:151.

Anderson, G.W., H.H. Goebel, and A. Simonati. 2012. Human pathology in NCL. *Biochim Biophys Acta*.

Ao, X., L. Zou, and Y. Wu. 2014. Regulation of autophagy by the Rab GTPase network. *Cell Death Differ*. 21:348-358.

Arighi, C.N., L.M. Hartnell, R.C. Aguilar, C.R. Haft, and J.S. Bonifacino. 2004. Role of the mammalian retromer in sorting of the cation-independent mannose 6-phosphate receptor. *J Cell Biol*. 165:123-133.

Best, H.L., N.J. Neverman, H.E. Wicky, N.L. Mitchell, B. Leitch, and S.M. Hughes. 2017. Characterisation of early changes in ovine CLN5 and CLN6 Batten disease neural cultures for the rapid screening of therapeutics. *Neurobiol Dis*. 100:62-74.

Birgisdottir, A.B., and T. Johansen. 2020. Autophagy and endocytosis - interconnections and interdependencies. *J Cell Sci*. 133.

Butz, E.S., U. Chandrachud, S.E. Mole, and S.L. Cotman. 2019. Moving towards a new era of genomics in the neuronal ceroid lipofuscinoses. *Biochim Biophys Acta Mol Basis Dis*:165571.

Cabukusta, B., and J. Neefjes. 2018. Mechanisms of lysosomal positioning and movement. *Traffic*. 19:761-769.

Canuel, M., S. Lefrancois, J. Zeng, and C.R. Morales. 2008. AP-1 and retromer play opposite roles in the trafficking of sortilin between the Golgi apparatus and the lysosomes. *Biochem Biophys Res Commun*. 366:724-730.

Ceresa, B.P., and S.J. Bahr. 2006. rab7 activity affects epidermal growth factor:epidermal growth factor receptor degradation by regulating endocytic trafficking from the late endosome. *J Biol Chem*. 281:1099-1106.

Ceresa, B.P., and J.L. Peterson. 2014. Cell and molecular biology of epidermal growth factor receptor. *Int Rev Cell Mol Biol*. 313:145-178.

Doccini, S., F. Morani, C. Nesti, F. Pezzini, G. Calza, R. Soliymani, G. Signore, S. Rocchiccioli, K.M. Kanninen, M.T. Huuskonen, M.H. Baumann, A. Simonati, M.M. Lalowski, and F.M. Santorelli. 2020. Proteomic and functional analyses in disease models reveal CLN5 protein involvement in mitochondrial dysfunction. *Cell Death Discov*. 6:18.

Francavilla, C., M. Papetti, K.T. Rigbolt, A.K. Pedersen, J.O. Sigurdsson, G. Cazzamali, G. Karemore, B. Blagoev, and J.V. Olsen. 2016. Multilayered proteomics reveals molecular switches dictating ligand-dependent EGFR trafficking. *Nat Struct Mol Biol*. 23:608-618.

Heo, J.M., A. Ordureau, S. Swarup, J.A. Paulo, K. Shen, D.M. Sabatini, and J.W. Harper. 2018. RAB7A phosphorylation by TBK1 promotes mitophagy via the PINK-PARKIN pathway. *Sci Adv*. 4:eaav0443.

Huber, R.J., and S. Mathavarajah. 2018. Cln5 is secreted and functions as a glycoside hydrolase in Dictyostelium. *Cell Signal*. 42:236-248.

Johansson, M., N. Rocha, W. Zwart, I. Jordens, L. Janssen, C. Kuijl, V.M. Olkkonen, and J. Neefjes. 2007. Activation of endosomal dynein motors by stepwise assembly of Rab7-RILP-p150Glued, ORP1L, and the receptor betalll spectrin. *J Cell Biol*. 176:459-471.

Jules, F., E. Sauvageau, K. Dumaresq-Doiron, J. Mazzaferri, M. Haug-Kroper, R. Fluhrer, S. Costantino, and S. Lefrancois. 2017. CLN5 is cleaved by members of the SPP/SPPL family to produce a mature soluble protein. *Exp Cell Res*. 357:40-50.

Kobayashi, H., K. Ogawa, R. Yao, O. Lichtarge, and M. Bouvier. 2009. Functional rescue of beta-adrenoceptor dimerization and trafficking by pharmacological chaperones. *Traffic*. 10:1019-1033.

Laurent-Matha, V., D. Derocq, C. Prebois, N. Katunuma, and E. Liaudet-Coopman. 2006. Processing of human cathepsin D is independent of its catalytic function and auto-activation: involvement of cathepsins L and B. *J Biochem*. 139:363-371.

Lefrancois, S., J. Zeng, A.J. Hassan, M. Canuel, and C.R. Morales. 2003. The lysosomal trafficking of sphingolipid activator proteins (SAPs) is mediated by sortilin. *Embo J*. 22:6430-6437.

Leinonen, H., V. Keksa-Goldsteine, S. Ragauskas, P. Kohlmann, Y. Singh, E. Savchenko, J. Puranen, T. Malm, G. Kalesnykas, J. Koistinaho, H. Tanila, and K.M. Kanninen. 2017. Retinal Degeneration In A Mouse Model Of CLN5 Disease Is Associated With Compromised Autophagy. *Sci Rep*. 7:1597.

Lin, X., J. Zhang, L. Chen, Y. Chen, X. Xu, W. Hong, and T. Wang. 2017. Tyrosine phosphorylation of Rab7 by Src kinase. *Cell Signal*. 35:84-94.

Lyly, A., C. von Schantz, C. Heine, M.L. Schmiedt, T. Sipila, A. Jalanko, and A. Kyttala. 2009. Novel interactions of CLN5 support molecular networking between Neuronal Ceroid Lipofuscinosis proteins. *BMC Cell Biol*. 10:83.

Mamo, A., F. Jules, K. Dumaresq-Doiron, S. Costantino, and S. Lefrancois. 2012. The role of ceroid lipofuscinosis neuronal protein 5 (CLN5) in endosomal sorting. *Mol Cell Biol*. 32:1855-1866.

Marwaha, R., S.B. Arya, D. Jagga, H. Kaur, A. Tuli, and M. Sharma. 2017. The Rab7 effector PLEKHM1 binds Arl8b to promote cargo traffic to lysosomes. *J Cell Biol*. 216:1051-1070.

McCormick, P.J., K. Dumaresq-Doiron, A.S. Pluviose, V. Pichette, G. Tosato, and S. Lefrancois. 2008. Palmitoylation controls recycling in lysosomal sorting and trafficking. *Traffic*. 9:1984-1997.

McEwan, D.G., D. Popovic, A. Gubas, S. Terawaki, H. Suzuki, D. Stadel, F.P. Coxon, D. Miranda de Stegmann, S. Bhogaraju, K. Maddi, A. Kirchof, E. Gatti, M.H. Helfrich, S. Wakatsuki, C. Behrends, P. Pierre, and I. Dikic. 2015. PLEKHM1 regulates autophagosome-lysosome fusion through HOPS complex and LC3/GABARAP proteins. *Mol Cell*. 57:39-54.

Mercier, J.F., A. Salahpour, S. Angers, A. Breit, and M. Bouvier. 2002. Quantitative assessment of beta 1- and beta 2-adrenergic receptor homo- and heterodimerization by bioluminescence resonance energy transfer. *J Biol Chem*. 277:44925-44931.

Modica, G., O. Skorobogata, E. Sauvageau, A. Vissa, C.M. Yip, P.K. Kim, H. Wurtele, and S. Lefrancois. 2017. Rab7 palmitoylation is required for efficient endosome-to-TGN trafficking. *J Cell Sci*. 130:2579-2590.

Mole, S.E., and S.L. Cotman. 2015. Genetics of the neuronal ceroid lipofuscinoses (Batten disease). *Biochim Biophys Acta*. 1852:2237-2241.

Nakamura, S., and T. Yoshimori. 2017. New insights into autophagosome-lysosome fusion. *J Cell Sci*. 130:1209-1216.

Oetjen, S., D. Kuhl, and G. Hermey. 2016. Revisiting the neuronal localization and trafficking of CLN3 in juvenile neuronal ceroid lipofuscinosis. *J Neurochem*. 139:456-470.

Pankiv, S., E.A. Alemu, A. Brech, J.A. Bruun, T. Lamark, A. Overvatn, G. Bjorkoy, and T. Johansen. 2010. FYCO1 is a Rab7 effector that binds to LC3 and PI3P to mediate microtubule plus end-directed vesicle transport. *J Cell Biol*. 188:253-269.

Progida, C., L. Malerod, S. Stuffers, A. Brech, C. Bucci, and H. Stenmark. 2007. RILP is required for the proper morphology and function of late endosomes. *J Cell Sci*. 120:3729-3737.

Rojas, R., T. van Vlijmen, G.A. Mardones, Y. Prabhu, A.L. Rojas, S. Mohammed, A.J. Heck, G. Raposo, P. van der Sluijs, and J.S. Bonifacino. 2008. Regulation of retromer recruitment to endosomes by sequential action of Rab5 and Rab7. *J Cell Biol*. 183:513-526.

Schindelin, J., I. Arganda-Carreras, E. Frise, V. Kaynig, M. Longair, T. Pietzsch, S. Preibisch, C. Rueden, S. Saalfeld, B. Schmid, J.Y. Tinevez, D.J. White, V. Hartenstein, K. Eliceiri, P. Tomancak, and A. Cardona. 2012. Fiji: an open-source platform for biological-image analysis. *Nat Methods*. 9:676-682.

Seaman, M.N. 2004. Cargo-selective endosomal sorting for retrieval to the Golgi requires retromer. *J Cell Biol*. 165:111-122.

Seaman, M.N., M.E. Harbour, D. Tattersall, E. Read, and N. Bright. 2009. Membrane recruitment of the cargo-selective retromer subcomplex is catalysed by the small GTPase Rab7 and inhibited by the Rab-GAP TBC1D5. *J Cell Sci*. 122:2371-2382.

Song, P., K. Trajkovic, T. Tsunemi, and D. Krainc. 2016. Parkin Modulates Endosomal Organization and Function of the Endo-Lysosomal Pathway. *J Neurosci*. 36:2425-2437.

Storch, S., S. Pohl, and T. Braulke. 2004. A dileucine motif and a cluster of acidic amino acids in the second cytoplasmic domain of the batten disease-related CLN3 protein are required for efficient lysosomal targeting. *J Biol Chem*. 279:53625-53634.

Uusi-Rauva, K., A. Kyttala, R. van der Kant, J. Vesa, K. Tanhuanpaa, J. Neefjes, V.M. Olkkonen, and A. Jalanko. 2012. Neuronal ceroid lipofuscinosis protein CLN3 interacts with motor proteins and modifies location of late endosomal compartments. *Cell Mol Life Sci*. 69:2075-2089.

van der Kant, R., A. Fish, L. Janssen, H. Janssen, S. Krom, N. Ho, T. Brummelkamp, J. Carette, N. Rocha, and J. Neefjes. 2013. Late endosomal transport and tethering are coupled processes controlled by RILP and the cholesterol sensor ORP1L. *J Cell Sci*. 126:3462-3474.

Vanlandingham, P.A., and B.P. Ceresa. 2009. Rab7 regulates late endocytic trafficking downstream of multivesicular body biogenesis and cargo sequestration. *J Biol Chem*. 284:12110-12124.

Vesa, J., M.H. Chin, K. Oelgeschlager, J. Isosomppi, E.C. DellAngelica, A. Jalanko, and L. Peltonen. 2002. Neuronal ceroid lipofuscinoses are connected at molecular level: interaction of CLN5 protein with CLN2 and CLN3. *Mol Biol Cell*. 13:2410-2420.

Wijdeven, R.H., H. Janssen, L. Nahidiazar, L. Janssen, K. Jalink, I. Berlin, and J. Neefjes. 2016. Cholesterol and ORP1L-mediated ER contact sites control autophagosome transport and fusion with the endocytic pathway. *Nat Commun*. 7:11808.

Yasa, S., G. Modica, E. Sauvageau, A. Kaleem, G. Hermey, and S. Lefrancois. 2020. CLN3 regulates endosomal function by modulating Rab7A-effector interactions. *J Cell Sci*. 133.

Zhou, C., W. Zhong, J. Zhou, F. Sheng, Z. Fang, Y. Wei, Y. Chen, X. Deng, B. Xia, and J. Lin. 2012. Monitoring autophagic flux by an improved tandem fluorescent-tagged LC3 (mTagRFP-mWasabi-LC3) reveals that high-dose rapamycin impairs autophagic flux in cancer cells. *Autophagy*. 8:1215-1226.

Yasa et al., 2020 Figure 1

Yasa et al., 2020 Figure 2

Yasa et al., 2020 Figure 3

Yasa et al., 2020 Figure 4

Yasa et al., 2020 Figure 5

A

HeLa   CLN5^{KO}
0   15 120   0   15 120 (min)

150 — Wb: EGFR

40 — Wb: actin

C 0' **D** 15' **E** 30' — HeLa

F 0' **G** 15' **H** 30' — CLN5^{KO}

B

0'
15'
120'

HeLa   CLN5^{KO}

**

I

# of EGF-448 Punta

0'
15'
30'

HeLa   CLN5^{KO}

*

Yasa et al., 2020 Figure 6

Yasa et al., 2020 Figure 7

mTagRFP- mWasabi-LC3