Cytoplasmic and nuclear Sw-5b NLR act both independently and synergistically to dictate full host defense against tospovirus infection

Hongyu Chen[1,¶], Xin Qian[1,¶], Xiaojiao Chen[1,¶], Tongqing Yang[1], Mingfeng Feng[1], Jing Chen[1],

Ruixiang Cheng[1], Hao Hong[1], Ying Zheng[1], Yuzhen Mei[4], Danyu Shen[1], Yi Xu[1], Min Zhu[1], Xin

Shun Ding[1] and Xiaorong Tao[1,*]

[1] Key Laboratory of Plant Immunity, Department of Plant Pathology, College of Plant Protection,

Nanjing Agricultural University, Nanjing 210095, P. R. China.

[2] Huaiyin Institute of Agricultural Sciences of Xuhuai Region in Jiangsu，Huaian 223001, Jiangsu,

P. R. China.

[3] College of Plant Protection, Yunnan Agricultural University, Kunming 650201, Yunnan, P. R.

China.

[4] State Key Laboratory of Rice Biology, Institute of Biotechnology, Zhejiang University,

Hanghzou 310029, P. R. China.

[¶] These authors contributed equally to this work.

Author for correspondence:

Xiaorong Tao

Tel: (+86)-25-84399027

Email: taoxiaorong@njau.edu.cn

RUNNING HEAD

Independent role of cytoplasmic and nuclear Sw-5b in immunity

Word Count: Main body of the text, 8136

Title 18, Summary 249, Keywords 8, Introduction 1286, Materials and Methods 1562,

Results 3030, Discussion 1884, Acknowledgments 35, References 3687, Figure

Legends 1103, Supporting Information Legends 786

9 Figures (Colors); Supporting information: 8 Supplemental Figures, 6 Supplemental

Tables

Summary

● Plant intracellular nucleotide binding-leucine-rich repeat (NLR) receptors play critical roles in mediating host immunity to pathogen attack. We use tomato Sw-5b::tospovirus as a model system to study the specific role of the compartmentalized plant NLR in dictating host defense against virus at different infection steps.

● We demonstrated here that tomato NLR Sw-5b translocates to cytoplasm and nucleus, respectively, to play different roles in inducing host resistances against *Tomato spotted wilt tospovirus* (TSWV) infection. The cytoplasmic Sw-5b functions to induce a strong cell death response to inhibit TSWV replication. This host response is, however, insufficient to block viral intercellular and long-distance movement. The nucleus-localized Sw-5b triggers a host defense that weakly inhibits viral replication but strongly impedes virus intercellular and systemic movement. Furthermore, the cytoplasmic and nuclear Sw-5b act synergistically to dictate full host defense to TSWV infection.

● We further demonstrated that the extended N-terminal *Solanaceae* domain (SD) of Sw-5b plays critical roles in cytoplasm/nucleus partitioning. Sw-5b nucleotide-binding leucine-rich repeat (NB-LRR) controls its cytoplasm localization. Strikingly, the SD but not coil-coil (CC) domain is crucial for Sw-5b receptor to translocate from cytoplasm to nucleus to trigger the immunity. The SD was found to interact with importins. Silencing both importin α and β expression disrupted Sw-5b nucleus translocation and host immunity

against TSWV systemic infection.

● Collectively, our findings suggest that Sw-5b bifurcates disease resistances

by cytoplasm/nucleus partitioning to block different infection steps of TSWV.

The findings also identified a new regulatory role of extra domain of a plant NLR

in mediating host innate immunity.

Keywords:

NLRs, Cytoplasm, Nuclear, Plant innate immunity, Tomato spotted wilt virus,

Replication, Cell-to-cell movement and Long distance movement

Introduction

Plant innate immunity plays critical roles in host defense against pathogen invasions,

and is triggered by cell-surface receptors or intracellular nucleotide-binding

leucine-rich repeat (NLR) receptors (Soosaar *et al.*, 2005; Dodds & Rathjen, 2010;

Cui *et al.*, 2015; Li *et al.*, 2015; Jones *et al.*, 2016; Kourelis & van der Hoorn, 2018;

Kapos *et al.*, 2019; van Wersch, 2020). Plant intracellular NLRs are the largest

classes of resistance proteins that function to detect pathogen effectors, and to activate

host immunity upon pathogen attack (Caplan, J *et al.*, 2008; Takken & Goverse, 2012;

Li *et al.*, 2015; Jones *et al.*, 2016; Kourelis & van der Hoorn, 2018; Kapos *et al.*,

2019). Plant NLRs typically contain an N-terminal domain, a central

nucleotide-binding domain (NB), a nucleotide-binding adaptor (ARC domain shared

by Apaf-1, certain resistance proteins, and CED-4), and a C-terminal leucine-rich

repeat (LRR) domain (Ea & Jones, 1998; Jones *et al.*, 2016; Ma *et al.*, 2018; Wang *et*

al., 2019a; Wang *et al.*, 2019b; Ma *et al.*, 2020). Based on the differences among the

N-terminal domains, plant NLRs can be further divided into two main categories,

known as the coiled-coil NLR (refers to as CNL) category and the Toll/interleukin-1

receptor NLR (TNL) category (Meyers *et al.*, 2003; Collier & Moffett, 2009; Qi &

Innes, 2013). The CC- or the TIR-domain-bearing NLRs have distinct genetic

requirements and can regulate different functions in the downstream of defense

signaling (Collier & Moffett, 2009; Qi & Innes, 2013; Horsefield *et al.*, 2019; Jubic *et*

al., 2019; van Wersch & Li, 2019; Wan *et al.*, 2019).

In addition to classical domains, non-canonical domains were frequently found to

integrate into certain NLRs. The additional domain, called BED, was first found in 32

poplar NLR proteins (Germain & Seguin, 2011). This extra BED domain was also

found in nine rice NLRs (Das *et al.*, 2014). The RATX1/HMA domain in the rice

NLRs RGA5 and Pik‑1 was found to act as integrated decoys to detect the cognate

pathogen effectors (Kanzaki *et al.*, 2012; Cesari *et al.*, 2013; Cesari *et al.*, 2014). The

WRKY domain on Arabidopsis thaliana NLR RRS1 was further found to function as

an integrated decoy that recognizes the effectors AvrRps4 and PopP2 (Le Roux *et al.*,

2015; Sarris *et al.*, 2015). Genome-wide analyses of plant NLR receptors revealed

that about 3.5 % of the NLRs carried specific non-canonical domains (Cesari *et al.*,

2014; Kroj *et al.*, 2016; Sarris *et al.*, 2016), and some of these non-canonical domains

were shown to be targeted by pathogen effectors during pathogen infections (Sarris *et*

al., 2016). However, molecular functions of most non-canonical domains in plant

NLRs remain largely unexplored.

Translocations of plant NLRs into proper subcellular compartments are critical for

the induction of innate immunity (Cui *et al.*, 2015; van Wersch, 2020). Multiple plant

NLRs and immune regulators, including tobacco N, Arabidopsis snc1, RRS1/RPS4,

barley MLA10, and Arabidopsis EDS1, NPR1, have been shown to accumulate in

both cytoplasm and nucleus, and for several nucleocytoplasmic NLRs accumulation

in nucleus is required for triggering host resistance to pathogen infections (Deslandes

et al., 2003; Burch-Smith *et al.*, 2007; Shen *et al.*, 2007; Wirthmueller *et al.*, 2007;

Tasset *et al.*, 2010; Bai *et al.*, 2012; Inoue *et al.*, 2013; Padmanabhan *et al.*, 2013).

Wheat Sr33, a homolog of barley MLA10, however, was reported to accumulate in

cytoplasm to induce host resistance against stem rust pathogen (Cesari *et al.*, 2016).

For potato Rx, a balanced cytoplasm and nucleus accumulation of Rx is needed to

induce the host immunity (Slootweg *et al.*, 2010; Tameling *et al.*, 2010). Other studies

have shown that *Arabidopsis* Rpm1 (Gao *et al.*, 2011), RPS2 (Axtell & Staskawicz,

2003), RPS5 (Qi *et al.*, 2012), rice Pit (Takemoto *et al.*, 2012), and tomato Tm-2^2

(Chen *et al.*, 2017; Wang *et al.*, 2020) need to associate with plasma membrane in

order to trigger cell death and host immunity. Latest studies have shown that the

activated *Arabidopsis* ZAR1 can bind to cellular membrane, leading to a membrane

leakage followed by cell death and host immunity (Wang *et al.*, 2019a; Wang *et al.*,

2019b). Flax L6 and M have been shown to accumulate in both Golgi apparatus and

tonoplast, and these compartmentalized localizations are necessary for the induction

of host resistance (Kawano *et al.*, 2014). The re-distribution of potato R3a from

cytosol to endosomal compartments is crucial for the induction of host resistance to

Phytophthora infestans infection (Engelhardt *et al.*, 2012). Different plant NLRs have

diverse subcellular localizations for their proper functions. However, how the

compartmentalized plant NLRs specifically dictate defense signaling remains largely

unknown.

Tomato spotted wilt tospovirus (TSWV) is one of most destructive plant NSVs,

infecting more than 1000 plant species, and causes crop losses more than one billion

US dollars annually worldwide (Kormelink *et al.*, 2011; Scholthof *et al.*, 2011; Oliver

& Whitfield, 2016). Tomato NLR Sw-5b confers strong resistance to TSWV infection

and has been widely used in tomato breeding projects to produce tospovirus resistant

tomato cultivars (Brommonschenkel *et al.*, 2000; Spassova M I, 2001; Turina *et al.*,

2016; Zhu *et al.*, 2019). Upon recognition of TSWV movement protein, NSm, Sw-5b

can trigger a hypersensitive response (HR), which typically associated with localized

cell death (Lopez *et al.*, 2011; Hallwass *et al.*, 2014; Peiro *et al.*, 2014; De Oliveira *et*

al., 2016; Zhao *et al.*, 2016; Leastro *et al.*, 2017). Tospoviruses are divided into

American and Euro-Asia type based on their geographic distribution and amino acid

sequence identity of viral nucleocapsid protein. We have previously shown that

Sw-5b can confer a broad-spectrum resistance to American type tospoviruses,

including TSWV, through recognition of a conserved 21 amino acid PAMP-like

region in the viral movement protein NSm (Zhu *et al.*, 2017). Sw-5b carries an

extended N-terminal *Solanaceae* domain (SD), a CC domain, a NB-ARC domain, and

a LRR domain (Brommonschenkel *et al.*, 2000; Spassova M I, 2001;

Lukasik-Shreepaathy *et al.*, 2012). Similar SDs have also been found in the Mi-1.2,

R8, Rpi-blb2, and Hero (Milligan *et al.*, 1998; Vos *et al.*, 1998; Ernst *et al.*, 2002; van

der Vossen *et al.*, 2005; Lukasik-Shreepaathy *et al.*, 2012; Vossen *et al.*, 2016). More

recently, Seong and others reported that the extended CNL has been evolved initially

in the ancestor of *Asterids* and *Amaranthaceae*, predated the *Solanaceae* family

(Seong *et al.*, 2020). In the presence of the extended N-terminal SD, Sw-5b is in an

autoinhibited state through multilayered interactions between SD, CC, NB-ARC, and

LRR domains (Chen *et al.*, 2016). For activation, the extra SD also recognizes NSm.

Sw-5b adopts a two-step NSm recognition strategy through SD and then LRR domain

(Li *et al.*, 2019). This two-step recognition mechanism significantly enhances the

sensitivity of the detection on TSWV NSm (Li *et al.*, 2019). Although Sw-5b is

known to localize in both cytoplasm and nucleus (De Oliveira *et al.*, 2016), the

biological roles of the cytoplasm- and the nucleus-accumulated Sw-5b in host

immunity signaling are unknown.

In this study, we investigated the subcellular distribution pattern of Sw-5b and the

functions of the compartmentalized Sw-5b in the induction of host immunity to

TSWV infection. We determined here that cytoplasm- and nucleus-accumulated

Sw-5b functions differently in inducing host defense response to inhibit multiple

tospovirus infection steps. The cytoplasmic Sw-5b can induce a strong cell death

response to suppress TSWV replication, whereas the nucleus-accumulated Sw-5b can

induce a strong defense against viral intercellular movement and systemic infection.

The combination of cytoplasmic and nuclear Sw-5b induces a synergistic and strong

plant immunity against tospovirus infection. We also found that the extended SD

functions as the key regulator for this critical intracellular translocation. The SD was

also found to interact with importins α and β to mediate Sw-5b nucleus translocation,

and to confer the full host immunity against tospovirus infection.

Materials and Methods

Plasmid construction

p2300S-YFP-Sw-5b was from a previously described source (Chen *et al.*, 2016).

Different domains of Sw-5b were PCR-amplified from p2300S-Sw-5b (Chen *et al.*,

2016) and cloned individually behind the *YFP* gene in the p2300S vector using a

two-step overlap PCR procedure as described (Li *et al.*, 2019). All the primers used in

this study are listed in Table S1.

To visualize the subcellular localization patterns of various fusion proteins, a SV40

T-Ag-derived nuclear localization signal (NLS, QPKKKRKVGG) (Lanford & Butel,

1984) or a PK1 nuclear export signal (NES, NELALKLAGLDINK) (Wen *et al.*, 1995)

was fused to the N-terminus of YFP-Sw-5b or the C-terminus of NSm-YFP, as

described (Kong *et al.*, 2017), to produce pNES-YFP-Sw-5b, pNLS-YFP-Sw-5b,

pNSm-YFP-NES, and pNSm-YFP-NLS, respectively. In addition, YFP-Sw-5b and

NSm-YFP were fused individually with a mutant NLS (nls, QPKKTRKVGG) or a

mutant NES (nes, NELALKAAGADANK) to produce pnes-YFP-Sw-5b,

pnls-YFP-Sw-5b, pNSm-YFP-nes, and pNSm-YFP-nls. The constructs were then

transformed individually into *Agrobacterium tumefaciens* strain GV3101 cells.

Transient gene expression, stable plant transformation, and virus inoculation

Nicotiana benthamiana were grown in soil in pots inside a greenhouse maintained at

25°C and a 16 h light/8 h dark photoperiod. Six-to-eight week-old *N. benthamiana*

plants were used for various assays. Transient gene expression assays were performed

in *N. benthamiana* leaves through agro-infiltration using Agrobacterium cultures

carrying specific expressing constructs as described previously (Feng *et al.*, 2016; Ma

et al., 2017). Transgenic *N. benthamiana* lines expressing YFP-Sw-5b or its

derivatives were made using constructs with a 35S promoter or a Sw-5b native

promoter via a standard leaf-disc transformation method (Chen *et al.*, 2016). The

resulting transgenic *N. benthamiana* lines were named as NES-YFP-Sw-5b,

NLS-YFP-Sw-5b, nes-YFP-Sw-5b, nls-YFP-Sw-5b, YFP-Sw-5b, and EV

(transformed with an empty vector), respectively. Inoculation of transgenic *N.*

benthamiana plants with TSWV was done by rubbing plant leaves with TSWV-YN

isolate-infected crude saps as described (Zhu *et al.*, 2017). TRV-mediated VIGS in *N.*

benthamiana plants was done as described (Ma *et al.*, 2015). The agro-infiltrated or

the virus-inoculated plants were growning inside a growth chamber maintained at

25/23 °C (day/night) with a 16/8 h light and dark photoperiod.

Particle bombardment

The particle bombardment is described (Feng et al., 2016). Briefly, 60 mg Tungsten

M-10 microcarrier (Bio-RAD) was placed into a 1.5 ml Eppendorf tube with 1 mL

70% ethanol. The tube was vortexes for 3 minutes, and then stood at room

temperature for 15 minutes. After centrifuge at low speed for 5 seconds, the

supernatant was removed and the pellet was rinsed with 70% ethanol for 3 times. One

mL 50% sterile glycerol solution was added and divided Tungsten M-10 microcarrier

into 50 µl. Five µg pRTL2-YFP, pRTL2-YFP-Sw-5b or pRTL2-YFP-Sw-5bD857V

plasmid DNA, 50 µl of 2.5 M CaCl2, and 20 µl of 0.1 M spermidine, respectively

were added and mixed with microcarrier. After centrifuge at low speed for 5 seconds

and the supernatant removed. The pellet was resuspended in 200 µl 70% ethanol and

centrifugation as described above. Use 48 µl of 100% ethanol to resuspend the

tungsten particle::plasmid DNA complexes, and load 15 48 µl mixture onto the center

of carrier (Bio-RAD), air dry, and use He/1000 particle transport system (BIO-RAD)

to bombard tomato leaves harvested from 3- or 4-week-old of Money Marker. The

bombarded leaves were incubated in Petri dishes for 24 hours at 25°C followed with

Confocal Microscope analysis.

Trypan blue staining

N. benthamiana leaves were harvested at 3 days post agro-infiltration (dpai) and

boiled for 5 min in a 1.15:1 (v/v) mixed ethanol and trypan blue staining solution (10

g phenol, 10 mL glycerol, 10 mL lactic acid, and 20 mg trypan blue in 10 mL distilled

water). The stained leaves were then de-stained in a chloral hydrate solution (2.5 g per

mL distilled water) as described (Bai *et al.*, 2012).

Electrolyte leakage assay

Electrolyte leakage assay was performed as previously described (Mittler *et al.*, 1999;

Zhu *et al.*, 2017) with slight modifications. Briefly, five leaf discs (9 mm in diameter

each) were taken from the agro-infiltrated leaves per treatment and at various dpai.

The harvested leaf discs from a specific treatment were floated on a 10 mL distilled

water for 3 h at room temperature (RT), and the conductivity of each bathing water

was measured (referred to as value A) using a Multiparameter Meter as instructed

(Mettler Toledo, Zurich, Switzerland). After the first measurement, the leaf discs were

returned to the bathing water and incubated at 95°C for 25 min. After cooling down to

RT, the conductivity of each bathing sample was measured again (referred to as value

B). The ion leakage was expressed as the ratio determined by value A/value B × 100.

The mean value and standard error of each treatment were calculated using the data

from three biological replicates per treatment at each sampling time point.

Confocal laser scanning microscopy

Tissue samples were collected from the leaves of transiently expressing YFP-Sw-5b

or one of the fusion proteins at 24–36 hours post agro-infiltration (hpai). The collected

tissue samples were mounted in water between a glass slide and a coverslip. Images

of individual samples were captured under a Carl Zeiss LSM 710 confocal laser

scanning microscope. YFP fusions were excited at 488 nm and the emission was

captured at 497–520 nm. The resulting images were further processed using the Zeiss

710 CLSM software followed by the Adobe Photoshop software (San Jose, CA,

USA).

Nucleus and cytoplasm fractionations

N. benthamiana leaf tissues (1 g per sample), representing a specific treatment, were

collected at 24 hpai, frozen in liquid nitrogen, ground to fine powders, and then

homogenized in 2 mL (per sample) extraction buffer 1 (20 mM Tris-HCl, pH 7.5, 20

mM KCl, 2.5 mM $MgCl_2$, 2 mM EDTA, 25% glycerol, 250 mM sucrose, 1 × Protease

Inhibitor Cocktail, and 5 mM DTT). The resulting lysate was filtered through 30 μm

filters to remove debris, and the filtrate was centrifuged at 2,000 $\times$ g for 5 minutes

to pellet nuclei. The supernatant from a sample was transferred into a new tube and

centrifuged at 10,000$\times g$ for 10 min. The resulting supernatant was used as the

cytoplasm fraction. The nuclei containing pellet was resuspended in 5 mL extraction

buffer 2 (20 mM Tris-HCl, pH 7.4, 25% glycerol, 2.5 mM MgCl2, and 0.2% Triton

X-100), centrifuged for 10 min at 2,000 $\times$ g followed by 4-6 cycles of resuspension

and centrifugation as described above. The resulting pellet was resuspended again in

500 μl extraction buffer 3 (20 mM Tris-HCl, pH 7.5, 0.25 M sucrose, 10 mM MgCl2,

0.5% Triton X-100, and 5 mM β-mercaptoethanol). The nuclei fraction was carefully

layered on the top of 500 mL extraction buffer 4 (20 mM Tris-HCl, pH 7.5, 1.7 M

sucrose, 10 mM MgCl2, 0.5% Triton X-100, 1$\times$Protease Inhibitor Cocktail, and 5

mM β-mercaptoethanol), and then centrifuged at 16,000 $\times$ g for 1 h. The resulting

pellet was resuspended in 500 μL extraction buffer 1 and stored at −80°C until use or

used immediately for SDS-PAGE assays. All the processes were performed on ice or

at 4°C. In this study, actin and histone H3 were used as the cytoplasmic and the

nuclear markers, respectively.

Western blot, co-immunoprecipitation and mass spectrometry analysis

Western blot and co-immunoprecipitation assays were performed as described (Zhu *et*

al., 2017). Briefly, agro-infiltrated leaf samples (1 g per sample) were harvested and

homogenized individually in pre-chilled mortars with pestles in 2 mL extraction

buffer [10% (v/v) glycerol, 25 mM Tris, pH 7.5, 1 mM EDTA, 150 mM NaCl, 10 mM

DTT, 2% (w/v) polyvinylpolypyrrolidone, and $1 \times$ protease inhibitor cocktail (Sigma,

Shanghai, China)]. Each crude slurry was transferred into a 2 mL Eppendorf tube, and

spun for 2 min at full speed in a refrigerated microcentrifuge. The supernatant was

transferred into a clean 1.5 mL Eppendorf tube and spun for 10 min at 4°C. For

Western blot assays, 50 μL supernatant from a sample was mixed with 150 μL

Laemmli buffer, boiled for 5 min, and analyzed in SDS-PAGE gels through

electrophoresis. For immunoprecipitation assays, 1 mL supernatant was mixed with

25 μL GFP-trap agarose beads (ChromoTek, Planegg-Martinsried, Germany),

incubated for 2 h at 4°C on an orbital shaker, and then pelleted through low speed

centrifugation. The blots were probed with a 1:2,500 (v/v) diluted anti-YFP antibody

or other specific antibodies followed a 1:10,000 (v/v) diluted horseradish peroxidase

(HRP)-conjugated goat anti-rabbit or a goat anti-mouse antibody (Sigma-Aldrich, St.

Louis, MO, USA). The detection signal was developed using the ECL substrate kit as

instructed (Thermo Scientific, Hudson, NH, USA).

For mass spectrometry analysis, the immunoprecipitation samples of YFP-SD

and SD without tag were processed by The Beijing Genomics Institute (BGI) for mass

spectrometry analysis. The immunoprecipitation samples of YFP-Sw-5b and Sw-5b

without tag were processed by Applied Protein Technology in Shanghai. Database

searches were performed using the Mascot search engine against *N. benthamiana*.

(https://solgenomics.net/organism/Nicotiana_benthamiana/genome).

RT-PCR detection of TSWV infection

Total RNA was extracted from TSWV-inoculated *N. benthamiana* plant leaves using

an RNA Purification Kit (Tiangen Biotech, Beijing, China), and then treated with

RNase-free DNase I (TaKaRa, Dalian, China). First-strand cDNA was synthesized

using a TSWV-specific primer (S3 Table). PCR reactions were as follows: initial

denaturation at 94°C for 2 min followed by 35 cycles of 94°C for 30 s, 52°C for 30 s,

and 72°C for 1 min. The final extension was 72°C for 10 min. The resulting PCR

products were visualized in 1.0% (w/v) agarose gels through electrophoresis.

Results

Determination of Sw-5b subcellular localization pattern

Expression of YFP-Sw-5b in *N. benthamiana* leaves resulted in a strong HR cell death

as well as Sw-5b (Chen *et al.*, 2016; Zhu *et al.*, 2017). To investigate the subcellular

localization pattern of Sw-5b, we transiently expressed YFP and YFP-Sw-5b in *N.*

benthamiana leaves, respectively, through agro-infiltration. Confocal Microscopy

results showed that the YFP-Sw-5b fusion accumulated in both cytoplasm and nucleus

in *N. benthamiana* leaf cells (Fig. 1b, middle image). This subcellular localization

pattern was similar to that of YFP (Fig. 1b, left image). When a D857V mutation,

which keeps Sw-5b in an autoactivated state (Chen *et al.*, 2016), was introduced into

Sw-5b to produce pYFP-Sw-5b^{D857V} and expressed in *N. benthamiana* leaves, the

mutant YFP-Sw-5b^{D857V} fusion also accumulated in both cell cytoplasm and nucleus

(Fig. 1b, right image). We also making a construct pNativePro::YFP-Sw-5b

expressing YFP-Sw-5b under native Sw-5b promoter. However, the expression of

YFP-Sw-5b by native Sw-5b promoter is too low to detect green fluorescence signal.

To investigate the subcellular localization pattern of Sw-5b in tomato leaf cells,

we transiently expressed YFP, YFP-Sw-5b, and YFP-Sw-5b^{D857V}, respectively,

through particle bombardment. Confocal Microscopy results showed that these three

proteins exhibited the same subcellular localization pattern as that in the *N.*

benthamiana leaf cells (Fig. 1c).

To further confirm the above results, we harvested *N. benthamiana* leaves

expressing YFP-Sw-5b or YFP-Sw-5b^{D857V} and analyzed by cytoplasm and nucleus

fractionation assay. Leaf samples agro-infiltrated with the empty vector (p2300S)

were also harvested and used as controls. Analyses of total protein, cytoplasm

fractions, and nucleus fractions from these harvested leaves using Western blot assays

showed that both YFP-Sw-5b and YFP-Sw-5b^{D857V} were accumulated in the

cytoplasm and nucleus (Fig. 1d).

Sw-5b recognizes TSWV NSm in the cytoplasm

TSWV NSm is known to reside in cytoplasm and plasmodesmata, but not in nucleus

(Kormelink *et al.*, 1994; Feng *et al.*, 2016). To determine where Sw-5b can recognize

TSWV NSm, we fused a NES, a nes, a NLS or a nls signal peptide to the C-terminus

of NSm-YFP to produce NSm-YFP-NES, NSm-YFP-nes, NSm-YFP-NLS, and

NSm-YFP-nls, respectively. Transient expressions of these fusions in *N. benthamiana*

leaves showed that NSm-YFP-NES accumulated exclusively in the cytoplasm, while

NSm-YFP-NLS accumulated in the nucleus (Fig. S1a). As expected, NSm-YFP-nes

and NSm-YFP-nls showed the same accumulation pattern as that of NSm-YFP (Fig.

S1a). When Sw-5b was co-expressed with one of the above four fusions in *N.*

benthamiana leaves through agro-infiltration, the leaf tissues co-expressing Sw-5b

and NSm-YFP-NES (Sw-5b + NSm-YFP-NES), Sw-5b and NSm-YFP-nes (Sw-5b +

NSm-YFP-nes), or Sw-5b and NSm-YFP-nls (Sw-5b + NSm-YFP-nls) developed a

strong HR cell death (Fig. S1b). In contrast, the leaf tissues co-expressing Sw-5b and

NSm-YFP-NLS (Sw-5b + NSm-YFP-NLS) did not. Western blot assays using a YFP

specific antibody confirmed that all the assayed proteins were expressed in the

infiltrated tissues (Fig. S1c), indicating that Sw-5b recognizes NSm in the cytoplasm.

Sw-5b activity in cell death induction is enhanced in the cytoplasm but

suppressed in the nucleus

To investigate the roles of the cytoplasmic and nuclear Sw-5b in the induction of cell

death and host immunity, we produced constructs to express YFP-Sw-5b,

NLS-YFP-Sw-5b, nls-YFP-Sw-5b, NES-YFP-Sw-5b, and nes-YFP-Sw-5b,

respectively, and then tested their abilities to elicit cell death and host immunity to

tospovirus infection. Transient expressions of these fusions in *N. benthamiana* leaves

showed that NES-YFP-Sw-5b accumulated exclusively in the cytoplasm, while

NLS-YFP-Sw-5b accumulated only in the nucleus (Fig. 2a). In addition,

nes-YFP-Sw-5b and nls-YFP-Sw-5b showed the same accumulation pattern as that

shown by YFP-Sw-5b. We then tested cell death induction through co-expressions of

NSm and YFP-Sw-5b (NSm + YFP-Sw-5b), NSm and NES-YFP-Sw-5b (NSm +

NES-YFP-Sw-5b), NSm and nes-YFP-Sw-5b (NSm + nes-YFP-Sw-5b), NSm and

NLS-YFP-Sw-5b (NSm + NLS-YFP-Sw-5b), or NSm and nls-YFP-Sw-5b (NSm +

nls-YFP-Sw-5b) in *N. benthamiana* leaves through agro-infiltration. Results of this

study showed that the NSm + NES-YFP-Sw-5b-induced cell death was stronger than

that induced by NSm + nes-YFP-Sw-5b or NSm + nls-YFP-Sw-5b co-expression (Fig.

2b). In addition, the cell death induced by NSm + NLS-YFP-Sw-5b co-expression

was suppressed (Fig. 2b). Western blot results showed that the stronger cell death

caused by NSm + NES-YFP-Sw-5b co-expression was not due to a greater

accumulation of NES-YFP-Sw-5b in the leaves (Fig. 2c). The ion leakage assay

results (Fig. 2d) agreed with the phenotype observation results, and indicated that

co-expression of NSm + NES-YFP-Sw-5b in leaves lead to a greater ion leakage

compared with that induced by the co-expression of NSm + nes-YFP-Sw-5b at 24 and

48 hours post agro-infiltration (hpai). The ion leakage caused by the co-expression of

NSm + NLS-YFP-Sw-5b was significantly weaker than that caused by the

co-expression of NSm + nls-YFP-Sw-5b (Fig. 2d).

Cytoplasmic Sw-5b induces a strong host defense against tospovirus replication

Virus infection in plant starts with virus replication in the initially infected cells

followed by spreading into adjacent cells for further infection. To monitor tospovirus

replication in plant cells, we recently developed a TSWV mini-replicon-based reverse

genetic system (Feng et al., 2020). In this study, co-expression of TSWV

mini-replicon SR$_{(+)eGFP}$, L$_{(+)opt}$ (with a codon usage optimized RdRp), VSRs and NSm

resulted in a cell-to-cell movement of SR$_{(+)eGFP}$. In contrast, co-expression of SR$_{(+)eGFP}$,

L$_{(+)opt}$, VSRs, and NSm$^{H93A\&H94A}$ mutant, a defective movement protein but can be

recognized by Sw-5b to cause a strong HR (Li *et al.*, 2009; Zhao *et al.*, 2016), in cells

resulted in the expression of SR$_{(+)eGFP}$ in only single cells (Fig. S2), thus dissecting the

viral replication from viral cell-to-cell movement. We then co-expressed SR$_{(+)eGFP}$,

L$_{(+)opt}$, VSRs, NSm$^{H93A\&H94A}$ mutant and one of the five proteins (i.e., Sw-5b,

NES-Sw-5b, nes-Sw-5b, NLS-Sw-5b, nls-Sw-5b) in *N. benthamiana* leaves. Leaves

co-expressing SR$_{(+)eGFP}$, L$_{(+)opt}$, VSRs, NSm$^{H93A\&H94A}$ mutant and p2300 (empty vector,

EV) were used as controls. The results showed that in the presence of Sw-5b or one of

its derivatives, the expression of SR$_{(+)eGFP}$ was strongly suppressed compared with

that expressed in the presence of EV (Fig. 3a). It is noteworthy that the expression of

SR$_{(+)eGFP}$ was less inhibited in the presence of NLS-Sw-5b (Fig. 3a). Western blot

result indicated that the GFP accumulation of SR$_{(+)eGFP}$ was strongly inhibited in the

presence of Sw-5b, NES-Sw-5b, nes-Sw-5b or nls-Sw-5b compared with that

expressed in the presence of NLS-Sw-5b or EV (Fig. 3b). This finding indicates that

the cytoplasmic Sw-5b can inhibit SR$_{(+)eGFP}$ expression, possibly through induction of

a host defense against TSWV replication.

Sw-5b induces a host defense against viral NSm intercellular movement

In our previous study, we used pmCherry-HDEL//NSm-GFP vector (Fig. 4a) to

investigate TSWV NSm cell-to-cell movement (Feng *et al.*, 2016). The expressed

mCherry-HDEL binds ER membrane in the initial cells but NSm-eGFP traffics between cells. To investigate whether the Sw-5b-induced host defense can affect TSWV NSm cell-to-cell movement, we co-expressed mCherry-HDEL, NSm-GFP, and Sw-5b or mCherry-HDEL, NSm-GFP, and EV in *N. benthamiana* leaves through agro-infiltration. Under the fluorescence microscope, both NSm-GFP and mCherry-HDEL were found in single cells in the presence of Sw-5b. In the presence of EV, however, NSm-GFP moved into multiple cells, while mCherry-HDEL accumulated in the initial cells (Fig. 4b, upper two panels). The result suggested that Sw-5b elicited a defense that strongly inhibited cell-to-cell movement of viral NSm.

To make sure this inhibition to viral NSm cell-to-cell movement is not caused by overexpression of Sw-5b, we also used the NSm^{T120N} mutant, from the resistance‑breaking (RB) TSWV isolates, which cannot be recognized by Sw-5b (Zhao et al., 2016). The assays showed that in the presence of either Sw-5b or EV, NSm^{T120N}-GFP moved into multiple cells, while mCherry-HDEL retained in the initial cells (Fig S3a).

Sw-5b in the nucleus but not in the cytoplasm triggers a defense against NSm cell-to-cell movement

To determine the effects of the cytoplasmic and nuclear Sw-5b on host defense against TSWV NSm intercellular movement, we co-expressed mCherry-HDEL and NSm-GFP with NES-Sw-5b, nes-Sw-5b, NLS-Sw-5b, or nls-Sw-5b in *N. benthamiana* leaves via agro-infiltration. The results showed that in the presence of NLS-Sw-5b, the cell-to-cell movement of NSm-GFP was inhibited (Fig. 4b). Similar

results were also obtained in the leaves co-expressing mCherry-HDEL and NSm-GFP

with nls-YFP-Sw-5b or nes-YFP-Sw-5b (Fig S3b). In the presence of

NES-YFP-Sw-5b, however, NSm-GFP did move into surrounding cells. (Fig. 4b).

This finding indicates that the Sw-5b in the nucleus but not in the cytoplasm induced

a host defense that inhibited TSWV NSm cell-to-cell movement.

Nuclear Sw-5b confers host immunity to TSWV systemic infection

To dissect the host immunity induced by the cytoplasmic and the nuclear Sw-5b, we

generated transgenic *N. benthamiana* lines expressing YFP-Sw-5b, NES-YFP-Sw-5b,

nes-YFP-Sw-5b, NLS-YFP-Sw-5b, and nls-YFP-Sw-5b, respectively (Tables S1 and

S2). After inoculation of these transgenic lines with TSWV-YN isolate, the EV

(control) transgenic plants developed typical viral symptoms including stunt, leaf curl

and mosaic at 7 to 15 days post inoculation (dpi). The NES-YFP-Sw-5b transgenic

plants developed a strong HR trailing in the systemic leaves by 7 to 15 days post

inoculation (dpi) (Fig. 5a and Fig. S4a), suggesting that NES-YFP-Sw-5b transgenic

plant did not block TSWV systemic infection and caused virus infection-related

systemic HR. In contrast, no systemic virus infection symptoms were observed in the

YFP-Sw-5b and the nes-YFP-Sw-5b transgenic plants. The RT-PCR agreed with the

symptom observation results and showed that TSWV-YN genomic RNA was

accumulated in the systemic leaves of the TSWV-YN-inoculated NES-YFP-Sw-5b or

the EV transgenic plants, but not in the systemic leaves of the TSWV-YN-inoculated

YFP-Sw-5b or nes-YFP-Sw-5b transgenic plants (Fig. 5c and Fig. S4b). Also in this

study, the TSWV-YN-inoculated NLS-YFP-Sw-5b or nls-YFP-Sw-5b transgenic

plants did not show virus like symptoms in their systemic leaves by 7-15 dpi (Fig. 5a,

and Fig. S4a). The RT-PCR result confirmed that TSWV-N genomic RNA had not

accumulated in the systemic leaves of the NLS-YFP-Sw-5b or the nls-YFP-Sw-5b

transgenic plants (Fig. 5c, and Fig. S4b), indicating that the nuclear Sw-5b is

responsible for the host immunity against TSWV systemic infection.

The cytoplasmic and the nuclear Sw-5b act synergistically to confer a strong

immunity to TSWV infection in *N. benthamiana*

To investigate whether cytoplasm-targeted and nucleus-targeted Sw-5b have joint

effects on the defense against TSWV infection, we constructed a $M_{(-)opt}$-$pSR_{(+)eGFP}$

vector by inserting a cassette expressing optimized TSWV M genomic sequence

(Feng *et al.*, 2020) into the $pSR_{(+)eGFP}$ mini-replicon to express NSm, N, and eGFP

simultaneously in the same cells (Fig. 6a). The construct $M_{(-)opt}$-$pSR_{(+)eGFP}$ couples the

functions for both viral replication and viral cell-to-cell movement, mimicking the

virus infection in plant leaves. After co-expressing this vector, the $L_{(+)opt}$ and the EV in

N. benthamiana leaves through agro-infiltration, the eGFP fluorescence was observed

in many cells, due to the presence of the NSm movement protein and the $RdRp_{opt}$ (Fig.

6b, upper left image). When $M_{(-)opt}$-$SR_{(+)eGFP}$, $L_{(+)opt}$ and Sw-5b were co-expressed in

N. benthamiana leaves, the eGFP fluorescence was hardly detected and some were

observed only in single leaf cells (Fig. 6b upper right image, Fig. S5a and b). When

$M_{(-)opt}$-$SR_{(+)eGFP}$, $L_{(+)opt}$ and NES-Sw-5b were co-expressed in *N. benthamiana* leaves,

the eGFP fluorescence was observed in clusters of a few cells (Fig. 6b, Fig. S5a),

indicating that limited cell-to-cell movement had occurred in these leaves (Fig. S5b).

When $M_{(-)opt}$-$SR_{(+)eGFP}$, $L_{(+)opt}$ and NLS-Sw-5b were co-expressed in leaves, a few of

eGFP fluorescence were detected but they were in single cells only. When leaves

co-expressing $M_{(-)opt}$-$SR_{(+)eGFP}$, $L_{(+)opt}$ and NES-Sw-5b + NLS-Sw-5b, the eGFP

fluorescence was also hardly detected and some were observed only in single leaf

cells. Western blot results showed that more eGFP had accumulated in the leaves

co-expressing $M_{(-)opt}$-$SR_{(+)eGFP}$, $L_{(+)opt}$, and EV, followed by the leaves co-expressing

$M_{(-)opt}$-$SR_{(+)eGFP}$, $L_{(+)opt}$, and NLS-Sw-5b, and then the leaves co-expressing

$M_{(-)opt}$-$SR_{(+)eGFP}$, $L_{(+)opt}$, and NES-Sw-5b. Much less eGFP had accumulated in the

leaves co-expressing $M_{(-)opt}$-$SR_{(+)eGFP}$, $L_{(+)opt}$, and NLS-Sw-5b + NES-Sw-5b, and in

the leaves co-expressing $M_{(-)opt}$-$SR_{(+)eGFP}$, $L_{(+)opt}$, and Sw-5b (Fig. 6c and d). The

accumulation of eGFP was lower in the leaves co-expressing NES-Sw-5b +

NLS-Sw-5b than that in the leaves co-expressing NES-Sw-5b or NLS-Sw-5b (Fig. 6c

and d), indicating that NES-Sw-5b and NLS-Sw-5b have additive role in mediating

host immunity against different TSWV infection steps.

The Sw-5b NB-ARC-LRR control its cytoplasm localization whereas the

extended N-terminal SD domain is crucial for targeting Sw-5b into nucleus, and

for inducing host systemic immunity

Sw-5b has an extended N-terminal SD domain, a CC domain, a NB-ARC domain, and

a C-terminal LRR domain (Chen *et al.*, 2016). To determine which domain(s) of

Sw-5b is/are responsible for nucleoplasm/nucleolus targeting and for plant immunity,

we tested these domains using various deletion mutants and YFP fusion proteins (Fig.

7a). We reported previously that the Sw-5b NB-ARC-LRR region was able to induce

HR cell death in plant in the presence of NSm (Chen *et al.*, 2016). In this study, we

fused YFP to the N-terminus of NB-ARC-LRR (Fig. 7a). Transient expression of

YFP-NB-ARC-LRR (112 kDa) in *N. benthamiana* leaf cells resulted in a localization

of the fusion protein in cytoplasm exclusively (Fig. 7b).

A previous study had shown that the CC domain of potato NLR receptor Rx was

required for targeting this protein to nucleus (Slootweg *et al.*, 2010). To determine the

function of Sw-5b CC domain in intracellular trafficking, we inserted the CC domain

between the YFP and NB-ARC-LRR to generate an YFP-CC-NB-ARC-LRR

construct or fused the CC domain to YFP to produce an YFP-CC construct. Transient

expression of these two fusion proteins individually in *N. benthamiana* leaves, and

examined the leaves under a confocal microscope, we determined that the YFP-CC

fusion protein accumulated in both cytoplasm and nucleus of the cells while the

YFP-CC-NB-ARC-LRR fusion protein was in the cytoplasm only (Fig. 7b). This

result indicated that addition of the CC domain to YFP-NB-ARC-LRR was not

sufficient to traffic the fusion protein into the nucleus.

An extended N-terminal SD domain is known to be present at the upstream of the

Sw-5b CC domain. In this study, we first generated an YFP-SD and an YFP-SD-CC

constructs, and transiently expressed them individually in *N. benthamiana* leaf cells.

Confocal Microscopy showed that both YFP-SD and YFP-SD-CC fusion proteins

accumulated in the cytoplasm and nucleus (Fig. 7b). We then inserted a SD between the YFP and CC-NB-ARC-LRR to produce an YFP-SD-CC-NB-ARC-LRR construction. Transient expression of this construct in *N. benthamiana* leaf cells showed that this fusion protein accumulated in the cytoplasm and nucleus (Fig. 7b).

We next generated stable transgenic *N. benthamiana* plants expressing YFP-NB-ARC-LRR, YFP-CC-NB-ARC-LRR and YFP-SD-CC-NB-ARC-LRR. Upon inoculation transgenic *N. benthamiana* plants expressing YFP-NB-ARC-LRR with TSWV, large HR foci were observed in the TSWV-inoculated leaves and later, HR trailing was seen in the systemic leaves of most assayed plants (Fig. 7c; Table S2 and S3). RT-PCR results confirmed the presence of TSWV genomic RNA in these systemic leaves (Fig. S6b). We also inoculate *N. benthamiana* plants expressing YFP-CC-NB-ARC-LRR with TSWV. By 7 dpi, no systemic resistance to TSWV infection was observed in these plants (Fig. 6a and B, Table S4). RT-PCR results showed that systemic infection of TSWV did occur in the TSWV-inoculated YFP-CC-NB-ARC-LRR transgenic plants (Fig. S6b). In contrast, transgenic plants expressing YFP-SD-CC-NB-ARC-LRR fusion exhibited a systemic immunity to TSWV infection (Fig. 7c, Fig. S6b).

These data indicated that Sw-5b NB-ARC-LRR control its cytoplasm localization, CC domain of Sw-5b alone was not sufficient to transport the NB-ARC-LRR into nucleus and the extended SD domain is required for targeting Sw-5b to the nucleus, and for inducing host immunity.

The extended SD domain interacted with *importin α1*, *α2* and *β*

To identify the cellular machinery needed for transporting Sw-5b into nucleus, we co-expressed YFP-SD and YFP-Sw-5b in *N. benthamiana* leaves followed by a co-immunoprecipitation (co-IP) and Mass Spectrometry. The results identified *N. benthamiana* importin α as one of candidate proteins interacted with YFP-SD and YFP-Sw-5b (Table S5 and S6). The co-IP and Mass Spectrometry also identified nuclear pore complex protein TPRb and nuclear pore complex protein Nup160a interacted with YFP-Sw-5b (Table S6). *Importins* play important roles in translocating proteins from cytoplasm into nucleus (Kanneganti *et al.*, 2007). We used BiFC analysis to confirm the interaction between YFP-SD with *N. benthamiana* importin homologs α1, α2 and β. The result showed that co-expression of cYFP-SD with nYFP-IMP α1, nYFP-IMP α2 or nYFP-IMP β produced a strong YFP fluorescence signal in nucleus. Co-expression of cYFP-Sw-5b with nYFP-IMP α1, nYFP-IMP α2 or nYFP-IMP β also detected a strong YFP fluorescence signal in nucleus (Fig. S7). In contrast, co-expression of controls cYFP-SD and nYFP, cYFP-Sw-5b and nYFP, cYFP and nYFP-IMP α1, cYFP and nYFP-IMP α2 or cYFP and nYFP-IMP β did not show fluorescence signal in *N. benthamiana* leaf cells (Fig. S7).

Silencing *importin α1*, *α2* and *β* expression abolished Sw-5b nucleus accumulation and host resistance to TSWV systemic infection

To determine the functions of importin α1, α2 and β in Sw-5b nucleus localization, we

silenced *importin* $\alpha 1$, $\alpha 2$, β, $\alpha 1$ and $\alpha 2$, and $\alpha 1$ and $\alpha 2$ and β expressions, respectively,

in *N. benthamiana* leaves using a tobacco rattle virus (TRV)-based virus-induced gene

silencing (VIGS) vector, and then transiently expressed YFP-Sw-5b in these plants.

Analyses of these plants through RT-PCR using gene specific primers showed that

silencing of these *importin* genes in *N. benthamiana* leaves were successful (Fig.

S8a). However, silencing individual *importin* gene or both *importin* $\alpha 1$ and $\alpha 2$ was

not enough to block the nucleus accumulation of YFP-Sw-5b (Fig. 8a). In contrast,

after *importin* $\alpha 1$, $\alpha 2$ and β were all silenced through VIGS, the nucleus accumulation

of YFP-Sw-5b was inhibited (Fig. 8a, the middle image in the bottom panel).

To investigate the effects of nuclear import defected Sw-5b on host immunity to

TSWV systemic infection, we silenced these *importin* genes in the Sw-5b transgenic

N. benthamiana plants as described above, and then inoculated them with TSWV. The

results showed that the plants silenced for *importin* $\alpha 1$, $\alpha 2$, and β gene, individually,

did not show TSWV systemic infection (Fig. 8b and Fig. S8b). In addition, the

transgenic plants silenced for both *importin* $\alpha 1$ and $\alpha 2$ genes also did not show

TSWV systemic infection (Fig. 8b and Fig. S8b). In contrast, after silencing *importin*

$\alpha 1$, $\alpha 2$ and β together, the plants developed clear TSWV symptoms in systemic

leaves followed by HR (Fig. 8b, white arrow and Fig. S8b), indicating that the nucleus

accumulation of Sw-5b is indispensable for the induction of host immunity against

TSWV systemic infection.

Discussion

In this report, we provide evidence to show that the cytoplasm-accumulated and the

nucleus-accumulated Sw-5b, a tomato immune receptor, play different roles in

inducing host defense against TSWV infection in plant. The cytoplasmic Sw-5b

functions to induce a strong cell death response to inhibit TSWV replication. This

host response is, however, insufficient to block virus intercellular and long-distance

movement. The nuclear-localized Sw-5b triggers a host defense that weakly inhibit

viral replication but strongly inhibit tospovirus intercellular and systemic movement.

These findings suggest that tomato Sw-5b NLR induces different types of defense

responses by cytoplasm and nucleus partitioning to combat virus at different infection

steps. Furthermore, the cytoplasmic and the nuclear Sw-5b act synergistically to

confer a strong host immunity to TSWV infection in plant. We also demonstrated that

the extra SD domain functioned as a critical intracellular translocation modulator,

allowing Sw-5b receptor to translocate from cytoplasm to nucleus to trigger the

immunity. The Sw-5b NB-LRR controls its cytoplasm localization. Unlike Rx CC

domain, Sw-5b CC domain is not sufficient to translocate NB-LRR into nucleus.

Strikingly, the SD is crucial for Sw-5b to translocate from cytoplasm for nucleus. This

SD-mediated receptor translocation is dependent on importins α and β.

Successful virus infection in plant requires several steps including viral replication

in the initially infected cells followed by cell-to-cell and long-distance movement

(Heinlein, 2015; Wang, 2015). After entering into plant cells, virus first encode

multiple proteins needed for its replication. Once the initial replication is established,

virus will encode specific protein(s), known as movement proteins (MPs), to traffic

viral genome or virions into adjacent cells through plasmodesmata in cell walls, and

then long-distantly into other parts of the plant to cause a systemic infection (Rao,

2002; Lucas, 2006; Taliansky *et al.*, 2008). To date, multiple plant NLRs, conferring

host resistance against plant viruses have been identified (Soosaar *et al.*, 2005; Meier

et al., 2019), but how these plant NLRs induce host resistance against virus infection

remain largely unknown.

In this study, we have determined that the forced cytoplasm accumulation of

Sw-5b can induce a stronger cell death than that caused by the accumulation of Sw-5b

in both cytoplasm and nucleus. While, the cell death induced by the forced nucleus

accumulation of Sw-5b was significantly weakened. We then analyzed

Sw-5b-mediated immunity against TSWV replication using a TSWV mini-replicon

system and a movement defective NSm mutant. Our results showed that the forced

cytoplasm accumulation of Sw-5b can induce a strong host defense against virus

replication in cells. This finding implies that cytoplasm is one of the main source of

defense signaling against TSWV replication. The defense signaling generated in

nucleus can only induce a weak defense against TSWV replication. Therefore, the

nuclear localized Sw-5b is only partially responsible for the induction of host defense

against TSWV replication. It is also possible that this nuclear localized

Sw-5b-induced weak host response is caused by a trace of NLS-YFP-Sw-5b

maintained in the cytoplasm that maybe below the detection limit of Confocal

Microscope. It has been shown to accumulate in both cytoplasm and nucleus, and the

forced cytoplasm accumulation of Barley MLA10 enhance cell death signaling (Bai *et*

al., 2012). We speculate that, for both MLA10 and Sw-5b, the cytoplasm

accumulation is crucial for the initiation and/or amplification of the cell death

signaling. The CC and the TIR domain of several plant NLRs have been shown to

trigger cell death (Swiderski *et al.*, 2009; Krasileva *et al.*, 2010; Bernoux *et al.*, 2011;

Collier *et al.*, 2011; Maekawa *et al.*, 2011; Bai *et al.*, 2012; Chen *et al.*, 2017; Wang

et al., 2020). Analyses of the three dimensional structures of Arabidopsis ZAR1

resistosome have also shown that its CC domain can form pentamer structures that

was able to target into host cell membranes, leading to ion leakage and cell death

(Wang *et al.*, 2019a; Wang *et al.*, 2019b). We speculate that cell death likely cause the

toxicity on viral replicase or other proteins associated with virus replication in cells.

Plant virus encodes specific movement protein(s) to traffic viral genome between

cells and then leaves to cause systemic infection (Rao, 2002; Lucas, 2006; Taliansky

et al., 2008). We reported previously that TSWV NSm alone can move between plant

cells (Feng *et al.*, 2016). In this study, we investigated the effect of the

Sw-5b-mediated host defense on TSWV intercellular movement. Through this study,

we have determined that after the recognition of NSm, Sw-5b receptor induced a

strong reaction to block NSm intercellular trafficking. Previous reports have some

indications on the role of plant NLRs in viral movement. Nevertheless it has no direct

evidence showing that plant NLRs induce resistance against viral movement. Deom

and colleagues had shown that the 9.4-kDa fluorescein isothiocyanate-labeled dextran

was unable to move between cells in the transgenic tobacco *N* leaves expressing

tobacco mosaic virus (TMV) movement protein at 24°C, an HR-permissive

temperature (Deom *et al.*, 1991). However, that study did not involve a TMV Avr

protein. In a different report, TMV-GFP showed a limited cell-to-cell movement in

leaves of tobacco cv. Sumsan NN at 33°C, an HR-nonpermissive temperature (Canto

& Palukaitis, 2002). Li and colleagues found that after treatment of SMV-inoculated

Jidou 7 resistant plants with a callose synthase inhibitor, the plants showed enlarged

HR lesions (Li *et al.*, 2012). The soybean *Rsv3* induced extreme resistance. However,

after this extremely resistant soybean line was treated with a callose synthase inhibitor,

the plants developed HR lesions upon SMV-G5H inoculation (Seo *et al.*, 2014). These

reports indicate that plant NLRs likely involves the defense against viral movement.

Here we provide the direct evidence that Sw-5b NLR can induce a strong defense

response to impede NSm intercellular trafficking. More importantly, we have

determined that the induction of host immunity to TSWV intercellular movement

requires the accumulation of Sw-5b in nucleus. Although the cytoplasmic Sw-5b can

induce a strong cell death response, it cannot prevent TSWV NSm cell-to-cell

movement. Consequently, we propose that nucleus is a key compartment to generate

defense signaling to block TSWV cell-to-cell movement.

In this study, although the NES-YFP-Sw-5b transgenic *N. benthamiana* plants

showed an HR, they were unable to stop TSWV systemic infection. We also showed

that Sw-5b YFP-NB-ARC-LRR (112 kDa) accumulates in cytoplasm exclusively (Fig.

7b), however, transgenic *N. benthamiana* plants expressing YFP-NB-ARC-LRR show

strong systemic HR trailing caused by TSWV infection. Based on these findings, we

conclude that HR cell death alone is not sufficient to block TSWV long-distance

movement. In our study, the NLS-YFP-Sw-5b transgenic plants were resistant to

TSWV systemic infection. After silencing the expressions of *importin α1*, *α2* and *β*

simultaneously to inhibit the nucleus accumulation of Sw-5b, however, the resistance

to TSWV systemic infection was abolished. These findings indicate that the

Sw-5b-mediated resistance signaling against viral systemic infection is generated in

nucleus. Some plant NLRs are known to interact with specific transcription factors in

nucleus upon recognition of pathogen effectors (Cui *et al.*, 2015; Kapos *et al.*, 2019).

The immune regulator EDS1 has also been shown to accumulate in nucleus to

reprogram RNA transcription (Garcia *et al.*, 2010; Heidrich *et al.*, 2011; Cui *et al.*,

2015; Lapin *et al.*, 2020). How Sw-5b regulates host immunity in nucleus requires

further investigations.

Several plant immune receptors and immune regulators, including, e.g. potato Rx

(Slootweg *et al.*, 2010; Tameling *et al.*, 2010), tobacco N (Burch-Smith *et al.*, 2007;

Caplan, JL *et al.*, 2008), barley MLA10 (Shen *et al.*, 2007), Arabidopsis

RRS1-R/RPS4, and snc1 (Deslandes *et al.*, 2003; Wirthmueller *et al.*, 2007; Cheng *et*

al., 2009), as well as Arabidopsis NPR1 (Katagiri & Tsuda, 2010), and EDS1 (Lapin

et al., 2020) have been found to be nucleocytoplasmic. For some of them, nuclear

accumulation of NLRs are required for the induction of plant immunity to pathogen

attacks. Moreover, the MLA10-YFP-NES fusion was found to induce a strong cell

death response, but not a strong host resistance to powdery mildew fungus infection.

In contrast, the MLA10-YFP-NLS fusion inhibited its activity to induce a cell death

response, but caused a host immunity to this pathogen (Bai *et al.*, 2012). In many

plant-pathogen interactions, cell death responses can be uncoupled from disease

resistance (Bendahmane *et al.*, 1999; Gassmann, 2005; Coll *et al.*, 2010; Heidrich *et*

al., 2011). This separation raises questions about how host resistance prevents

pathogen invasion and what are the roles of cell death during pathogen infection. It is

unclear whether the MLA10-YFP-NES-induced cell death has some inhibitory effects

on powdery mildew fungus infection. In this study, we determined that cytoplasm-

and nuclear-accumulation of Sw-5b have different functions. The

cytoplasm-accumulated Sw-5b induces a strong defense against virus replication,

whereas the nuclear-accumulated Sw-5b induced an inhibition of virus cell-to-cell and

long distance movement. Both cytoplasmic and nuclear Sw-5b are needed to confer a

synergistic and full defense against tospovirus infection.

We have also determined that Sw-5b NB-ARC-LRR and SD domains are

important to regulate the proper subcellular localization of Sw-5b and the proper

nucleoplasmic distribution of Sw-5b is needed to elicit full immune responses to

inhibit different TSWV infection steps. Sw-5b NB-ARC-LRR controls its cytoplasm

localization. The CC domain of the Sw-5b is not sufficient to target the receptor into

nuclear. Importantly, the extended SD of Sw-5b is absolutely required for the nucleus

translocation. Because non-canonical domains are frequently found in other NLRs

and are quite diversified, our findings have broad implications to investigate the

potential new functions of non-canonical domains that integrated in the plant NLRs to

regulate the plant immunity against pathogen invasions.

Through co-IP, Mass Spectrometry and BiFC analysis, we found that the extended

SD domain and Sw-5b interacted with host importin machineries to translocate the Sw-5b receptor from cytoplasm into nucleus to mediate local and systemic resistance to tospovirus. Recent studies have shown that nuclearporin MOS3, MOS6 and nuclear pore complex component MOS7/Nup88 proteins played important roles in regulating Arabidopsis innate immunity (Palma *et al.*, 2005; Zhang & Li, 2005; Cheng *et al.*, 2009). We found that when *importin α1*, *importin α2* and *importin β* gene were all silenced through VIGS, the nucleus targeting of Sw-5b were completely blocked, and consequently, the Sw-5b-mediated systemic immunity to tospovirus infection was compromised. Importin α and β are known to form a nucleus import complex. Binding with importin β could activate importin α to form a binding surface for NLS proteins (Stewart, 2007). Hence, disruption of either importin α or importin β would block the nucleus import of NLS proteins. Because silencing *importin α* or *importin β* gene expression through VIGS did not disrupt the nuclear targeting of Sw-5b, we speculate that a non-canonical nuclear import pathway may take part in importing Sw-5b into nucleus.

Based on the above results, we have created a working model for the Sw-5b NLR-induced host resistance against TSWV replication, and intercellular and long-distance movement in plant (Fig. 9). Upon recognition of NSm in cytoplasm, Sw-5b switched from an autoinhibited state to an activated state. The activated Sw-5b accumulated in cytoplasm and also translocate into nucleus via importins α and β. The cytoplasm-accumulated and the nucleus-accumulated Sw-5b play different roles in inducing host immunity against TSWV infection. The cytoplasmic Sw-5b functions

to induce a cell death response to inhibit TSWV replication, while the nuclear Sw-5b

functions to induce a weak host defense against TSWV replication, but a strong

defense against TSWV cell-to-cell and long-distance movement. The concerted

defense signaling generated in the cytoplasm and nucleus resulted in a strong host

resistance to tospovirus infection.

Acknowledgments

This work was supported by the National Natural Science Foundation of China

(31630062, 31925032 and 31870143), the Fundamental Research Funds for the

Central Universities (JCQY201904 and KYXK202012), Youth Science and

Technology Innovation Program to XT.

Author contributions

HC XQ, XC and XT, designed the research; HC, XQ and XC, TY, MF, JC, RC, HH,

YZ, YM, DS, YX, MZ performed the experiments; HC, XSD and XT interpreted the

result and wrote the paper.

ORCID

Hongyu Chen, 0000-0001-8142-0653

Yi Xu, 0000-0002-1913-4530

Min Zhu, 0000-0002-9354-4300

Xiaorong Tao, 0000-0003-1259-366X

Competing interests

The authors declare that nocompeting interests exist.

Data availability

All data produced in this study are presented in this manuscript or as the supporting

files

References

Axtell MJ, Staskawicz BJ. 2003. Initiation of RPS2-specified disease resistance in Arabidopsis is coupled to the AvrRpt2-directed elimination of RIN4. *Cell* **112**(3): 369-377.

Bai S, Liu J, Chang C, Zhang L, Maekawa T, Wang Q, Xiao W, Liu Y, Chai J, Takken FL, Schulze-Lefert P, Shen QH. 2012. Structure-function analysis of barley NLR immune receptor MLA10 reveals its cell compartment specific activity in cell death and disease resistance. *PLoS Pathog* **8**(6): e1002752.

Bendahmane A, Kanyuka K, Baulcombe DC. 1999. The Rx gene from potato controls separate virus resistance and cell death responses. *Plant Cell* **11**(5): 781-792.

Bernoux M, Ve T, Williams S, Warren C, Hatters D, Valkov E, Zhang X, Ellis JG, Kobe B, Dodds PN. 2011. Structural and functional analysis of a plant resistance protein TIR domain reveals interfaces for self-association, signaling, and autoregulation. *Cell Host Microbe* **9**(3): 200-211.

Brommonschenkel SH, Frary A, Frary A, Tanksley SD. 2000. The broad-spectrum tospovirus resistance gene Sw-5 of tomato is a homolog of the root-knot nematode resistance gene Mi. *Mol Plant Microbe Interact* **13**(10): 1130-1138.

Burch-Smith TM, Schiff M, Caplan JL, Tsao J, Czymmek K, Dinesh-Kumar SP. 2007. A novel role for the TIR domain in association with pathogen-derived elicitors. *PLoS Biol* **5**(3): e68.

Canto T, Palukaitis P. 2002. Novel N gene-associated, temperature-independent resistance to the movement of tobacco mosaic virus vectors neutralized by a cucumber mosaic virus RNA1 transgene. *J Virol* **76**(24): 12908-12916.

Caplan J, Padmanabhan M, Dinesh-Kumar SP. 2008. Plant NB-LRR Immune Receptors: From Recognition to Transcriptional Reprogramming. *Cell Host Microbe* **3**(3): 126-135.

Caplan JL, Mamillapalli P, Burch-Smith TM, Czymmek K, Dinesh-Kumar SP. 2008.
Chloroplastic protein NRIP1 mediates innate immune receptor recognition of a viral effector.
Cell **132**(3): 449-462.

Cesari S, Bernoux M, Moncuquet P, Kroj T, Dodds PN. 2014. A novel conserved mechanism for
plant NLR protein pairs: the "integrated decoy" hypothesis. *Front Plant Sci* **5**: 606.

Cesari S, Moore J, Chen C, Webb D, Periyannan S, Mago R, Bernoux M, Lagudah ES, Dodds
PN. 2016. Cytosolic activation of cell death and stem rust resistance by cereal MLA-family
CC-NLR proteins. *Proc Natl Acad Sci U S A* **113**(36): 10204-10209.

Cesari S, Thilliez G, Ribot C, Chalvon V, Michel C, Jauneau A, Rivas S, Alaux L, Kanzaki H,
Okuyama Y, Morel JB, Fournier E, Tharreau D, Terauchi R, Kroj T. 2013. The rice
resistance protein pair RGA4/RGA5 recognizes the Magnaporthe oryzae effectors AVR-Pia
and AVR1-CO39 by direct binding. *Plant Cell* **25**(4): 1463-1481.

Chen T, Liu D, Niu X, Wang J, Qian L, Han L, Liu N, Zhao J, Hong Y, Liu Y. 2017. Antiviral
Resistance Protein Tm-2(2) Functions on the Plasma Membrane. *Plant Physiol* **173**(4):
2399-2410.

Chen X, Zhu M, Jiang L, Zhao W, Li J, Wu J, Li C, Bai B, Lu G, Chen H, Moffett P, Tao X. 2016.
A multilayered regulatory mechanism for the autoinhibition and activation of a plant
CC-NB-LRR resistance protein with an extra N-terminal domain. *New Phytol* **212**(1):
161-175.

Cheng YT, Germain H, Wiermer M, Bi D, Xu F, Garcia AV, Wirthmueller L, Despres C, Parker
JE, Zhang Y, Li X. 2009. Nuclear pore complex component MOS7/Nup88 is required for
innate immunity and nuclear accumulation of defense regulators in Arabidopsis. *Plant Cell*
21(8): 2503-2516.

Coll NS, Vercammen D, Smidler A, Clover C, Van Breusegem F, Dangl JL, Epple P. 2010.
Arabidopsis type I metacaspases control cell death. *Science* **330**(6009): 1393-1397.

Collier SM, Hamel LP, Moffett P. 2011. Cell death mediated by the N-terminal domains of a unique
and highly conserved class of NB-LRR protein. *Mol Plant Microbe Interact* **24**(8): 918-931.

Collier SM, Moffett P. 2009. NB-LRRs work a "bait and switch" on pathogens. *Trends Plant Sci*
14(10): 521-529.

Cui H, Tsuda K, Parker JE. 2015. Effector-triggered immunity: from pathogen perception to robust
defense. *Annu Rev Plant Biol* **66**: 487-511.

Das B, Sengupta S, Prasad M, Ghose TK. 2014. Genetic diversity of the conserved motifs of six
bacterial leaf blight resistance genes in a set of rice landraces. *BMC Genet* **15**: 82.

De Oliveira AS, Koolhaas I, Boiteux LS, Caldararu OF, Petrescu AJ, Oliveira Resende R,
Kormelink R. 2016. Cell death triggering and effector recognition by Sw-5 SD-CNL proteins
from resistant and susceptible tomato isolines to Tomato spotted wilt virus. *Mol Plant Pathol*
17(9): 1442-1454.

Deom CM, Wolf S, Holt CA, Lucas WJ, Beachy RN. 1991. Altered function of the tobacco mosaic
virus movement protein in a hypersensitive host. *Virology* **180**(1): 251-256.

Deslandes L, Olivier J, Peeters N, Feng DX, Khounlotham M, Boucher C, Somssich I, Genin S,
Marco Y. 2003. Physical interaction between RRS1-R, a protein conferring resistance to
bacterial wilt, and PopP2, a type III effector targeted to the plant nucleus. *Proc Natl Acad Sci*
U S A **100**(13): 8024-8029.

Dodds PN, Rathjen JP. 2010. Plant immunity: towards an integrated view of plant-pathogen

interactions. *Nat Rev Genet* **11**(8): 539-548.

Ea VDB, Jones JD. 1998. Plant disease-resistance proteins and the gene-for-gene concept. *Trends in*
Biochemical Sciences **23**(12): 454-456(453).

Engelhardt S, Boevink PC, Armstrong MR, Ramos MB, Hein I, Birch PR. 2012. Relocalization of
late blight resistance protein R3a to endosomal compartments is associated with effector
recognition and required for the immune response. *Plant Cell* **24**(12): 5142-5158.

Ernst K, Kumar A, Kriseleit D, Kloos DU, Phillips MS, Ganal MW. 2002. The broad-spectrum
potato cyst nematode resistance gene (Hero) from tomato is the only member of a large gene
family of NBS-LRR genes with an unusual amino acid repeat in the LRR region. *Plant J* **31**(2):
127-136.

Feng M, Cheng R, Chen M, Guo R, Li L, Feng Z, Wu J, Xie L, Hong J, Zhang Z, Kormelink R,
Tao X. 2020. Rescue of tomato spotted wilt virus entirely from complementary DNA clones.
Proc Natl Acad Sci U S A **117**(2): 1181-1190.

Feng Z, Xue F, Xu M, Chen X, Zhao W, Garcia-Murria MJ, Mingarro I, Liu Y, Huang Y, Jiang
L, Zhu M, Tao X. 2016. The ER-Membrane Transport System Is Critical for Intercellular
Trafficking of the NSm Movement Protein and Tomato Spotted Wilt Tospovirus. *PLoS*
Pathog **12**(2): e1005443.

Gao Z, Chung EH, Eitas TK, Dangl JL. 2011. Plant intracellular innate immune receptor Resistance
to Pseudomonas syringae pv. maculicola 1 (RPM1) is activated at, and functions on, the
plasma membrane. *Proc Natl Acad Sci U S A* **108**(18): 7619-7624.

Garcia AV, Blanvillain-Baufume S, Huibers RP, Wiermer M, Li G, Gobbato E, Rietz S, Parker
JE. 2010. Balanced nuclear and cytoplasmic activities of EDS1 are required for a complete
plant innate immune response. *PLoS Pathog* **6**: e1000970.

Gassmann W. 2005. Natural variation in the Arabidopsis response to the avirulence gene hopPsyA
uncouples the hypersensitive response from disease resistance. *Mol Plant Microbe Interact*
18(10): 1054-1060.

Germain H, Seguin A. 2011. Innate immunity: has poplar made its BED? *New Phytol* **189**(3):
678-687.

Hallwass M, de Oliveira AS, de Campos Dianese E, Lohuis D, Boiteux LS, Inoue-Nagata AK,
Resende RO, Kormelink R. 2014. The Tomato spotted wilt virus cell-to-cell movement
protein (NSM) triggers a hypersensitive response in Sw-5-containing resistant tomato lines
and in Nicotiana benthamiana transformed with the functional Sw-5b resistance gene copy.
Mol Plant Pathol **15**(9): 871-880.

Heidrich K, Wirthmueller L, Tasset C, Pouzet C, Deslandes L, Parker JE. 2011. Arabidopsis
EDS1 connects pathogen effector recognition to cell compartment-specific immune responses.
Science **334**(6061): 1401-1404.

Heinlein M. 2015. Plant virus replication and movement. *Virology* **479-480**: 657-671.

Horsefield S, Burdett H, Zhang X, Manik MK, Shi Y, Chen J, Qi T, Gilley J, Lai JS, Rank MX,
Casey LW, Gu W, Ericsson DJ, Foley G, Hughes RO, Bosanac T, von Itzstein M,
Rathjen JP, Nanson JD, Boden M, Dry IB, Williams SJ, Staskawicz BJ, Coleman MP,
Ve T, Dodds PN, Kobe B. 2019. NAD(+) cleavage activity by animal and plant TIR domains
in cell death pathways. *Science* **365**(6455): 793-799.

Inoue H, Hayashi N, Matsushita A, Xinqiong L, Nakayama A, Sugano S, Jiang CJ, Takatsuji H.
2013. Blast resistance of CC-NB-LRR protein Pb1 is mediated by WRKY45 through

protein-protein interaction. *Proc Natl Acad Sci U S A* **110**(23): 9577-9582.

Jones JD, Vance RE, Dangl JL. 2016. Intracellular innate immune surveillance devices in plants and
animals. *Science* **354**(6316).

Jubic LM, Saile S, Furzer OJ, El Kasmi F, Dangl JL. 2019. Help wanted: helper NLRs and plant
immune responses. *Curr Opin Plant Biol* **50**: 82-94.

Kanneganti TD, Bai X, Tsai CW, Win J, Meulia T, Goodin M, Kamoun S, Hogenhout SA. 2007.
A functional genetic assay for nuclear trafficking in plants. *Plant J* **50**(1): 149-158.

Kanzaki H, Yoshida K, Saitoh H, Fujisaki K, Hirabuchi A, Alaux L, Fournier E, Tharreau D,
Terauchi R. 2012. Arms race co-evolution of Magnaporthe oryzae AVR-Pik and rice Pik
genes driven by their physical interactions. *Plant J* **72**(6): 894-907.

Kapos P, Devendrakumar KT, Li X. 2019. Plant NLRs: From discovery to application. *Plant Sci* **279**:
3-18.

Katagiri F, Tsuda K. 2010. Understanding the plant immune system. *Mol Plant Microbe Interact*
23(12): 1531-1536.

Kawano Y, Fujiwara T, Yao A, Housen Y, Hayashi K, Shimamoto K. 2014.
Palmitoylation-dependent membrane localization of the rice resistance protein pit is critical
for the activation of the small GTPase OsRac1. *J Biol Chem* **289**(27): 19079-19088.

Kong L, Qiu X, Kang J, Wang Y, Chen H, Huang J, Qiu M, Zhao Y, Kong G, Ma Z, Wang Y, Ye
W, Dong S, Ma W, Wang Y. 2017. A Phytophthora Effector Manipulates Host Histone
Acetylation and Reprograms Defense Gene Expression to Promote Infection. *Curr Biol* **27**(7):
981-991.

Kormelink R, Garcia ML, Goodin M, Sasaya T, Haenni AL. 2011. Negative-strand RNA viruses:
the plant-infecting counterparts. *Virus Res* **162**(1-2): 184-202.

Kormelink R, Storms M, Van Lent J, Peters D, Goldbach R. 1994. Expression and subcellular
location of the NSM protein of tomato spotted wilt virus (TSWV), a putative viral movement
protein. *Virology* **200**(1): 56-65.

Kourelis J, van der Hoorn RAL. 2018. Defended to the Nines: 25 Years of Resistance Gene Cloning
Identifies Nine Mechanisms for R Protein Function. *Plant Cell* **30**(2): 285-299.

Krasileva KV, Dahlbeck D, Staskawicz BJ. 2010. Activation of an Arabidopsis resistance protein is
specified by the in planta association of its leucine-rich repeat domain with the cognate
oomycete effector. *Plant Cell* **22**(7): 2444-2458.

Kroj T, Chanclud E, Michel-Romiti C, Grand X, Morel JB. 2016. Integration of decoy domains
derived from protein targets of pathogen effectors into plant immune receptors is widespread.
New Phytol **210**(2): 618-626.

Lanford RE, Butel JS. 1984. Construction and characterization of an SV40 mutant defective in
nuclear transport of T antigen. *Cell* **37**(3): 801-813.

Lapin D, Bhandari DD, Parker JE. 2020. Origins and Immunity Networking Functions of EDS1
Family Proteins. *Annu Rev Phytopathol* **58**: 253-276.

Le Roux C, Huet G, Jauneau A, Camborde L, Tremousaygue D, Kraut A, Zhou B, Levaillant M,
Adachi H, Yoshioka H, Raffaele S, Berthome R, Coute Y, Parker JE, Deslandes L. 2015.
A receptor pair with an integrated decoy converts pathogen disabling of transcription factors
to immunity. *Cell* **161**(5): 1074-1088.

Leastro MO, Pallas V, Resende RO, Sanchez-Navarro JA. 2017. The functional analysis of distinct
tospovirus movement proteins (NSM) reveals different capabilities in tubule formation,

cell-to-cell and systemic virus movement among the tospovirus species. *Virus Res* **227**: 57-68.

Li J, Huang H, Zhu M, Huang S, Zhang W, Dinesh-Kumar SP, Tao X. 2019. A Plant Immune
Receptor Adopts a Two-Step Recognition Mechanism to Enhance Viral Effector Perception.
Mol Plant **12**(2): 248-262.

Li W, Lewandowski DJ, Hilf ME, Adkins S. 2009. Identification of domains of the Tomato spotted
wilt virus NSm protein involved in tubule formation, movement and symptomatology.
Virology **390**(1): 110-121.

Li W, Zhao Y, Liu C, Yao G, Wu S, Hou C, Zhang M, Wang D. 2012. Callose deposition at
plasmodesmata is a critical factor in restricting the cell-to-cell movement of Soybean mosaic
virus. *Plant Cell Rep* **31**(5): 905-916.

Li X, Kapos P, Zhang Y. 2015. NLRs in plants. *Curr Opin Immunol* **32**: 114-121.

Lopez C, Aramburu J, Galipienso L, Soler S, Nuez F, Rubio L. 2011. Evolutionary analysis of
tomato Sw-5 resistance-breaking isolates of Tomato spotted wilt virus. *J Gen Virol* **92**(Pt 1):
210-215.

Lucas WJ. 2006. Plant viral movement proteins: agents for cell-to-cell trafficking of viral genomes.
Virology **344**(1): 169-184.

Lukasik-Shreepaathy E, Slootweg E, Richter H, Goverse A, Cornelissen BJ, Takken FL. 2012.
Dual regulatory roles of the extended N terminus for activation of the tomato MI-1.2
resistance protein. *Mol Plant Microbe Interact* **25**(8): 1045-1057.

**Ma S, Lapin D, Liu L, Sun Y, Song W, Zhang X, Logemann E, Yu D, Wang J, Jirschitzka J, Han
Z, Schulze-Lefert P, Parker JE, Chai J. 2020.** Direct pathogen-induced assembly of an
NLR immune receptor complex to form a holoenzyme. *Science* **370**(6521).

**Ma Y, Guo H, Hu L, Martinez PP, Moschou PN, Cevik V, Ding P, Duxbury Z, Sarris PF, Jones
JDG. 2018.** Distinct modes of derepression of an Arabidopsis immune receptor complex by
two different bacterial effectors. *Proc Natl Acad Sci U S A* **115**(41): 10218-10227.

**Ma Z, Song T, Zhu L, Ye W, Wang Y, Shao Y, Dong S, Zhang Z, Dou D, Zheng X, Tyler BM,
Wang Y. 2015.** A Phytophthora sojae Glycoside Hydrolase 12 Protein Is a Major Virulence
Factor during Soybean Infection and Is Recognized as a PAMP. *Plant Cell* **27**(7): 2057-2072.

**Ma Z, Zhu L, Song T, Wang Y, Zhang Q, Xia Y, Qiu M, Lin Y, Li H, Kong L, Fang Y, Ye W,
Wang Y, Dong S, Zheng X, Tyler BM, Wang Y. 2017.** A paralogous decoy protects
Phytophthora sojae apoplastic effector PsXEG1 from a host inhibitor. *Science* **355**(6326):
710-714.

**Maekawa T, Cheng W, Spiridon LN, Toller A, Lukasik E, Saijo Y, Liu P, Shen QH, Micluta MA,
Somssich IE, Takken FLW, Petrescu AJ, Chai J, Schulze-Lefert P. 2011.** Coiled-coil
domain-dependent homodimerization of intracellular barley immune receptors defines a
minimal functional module for triggering cell death. *Cell Host Microbe* **9**(3): 187-199.

Meier N, Hatch C, Nagalakshmi U, Dinesh-Kumar SP. 2019. Perspectives on intracellular
perception of plant viruses. *Mol Plant Pathol* **20**(9): 1185-1190.

Meyers BC, Kozik A, Griego A, Kuang H, Michelmore RW. 2003. Genome-wide analysis of
NBS-LRR-encoding genes in Arabidopsis. *Plant Cell* **15**(4): 809-834.

Milligan SB, Bodeau J, Yaghoobi J, Kaloshian I, Zabel P, Williamson VM. 1998. The root knot
nematode resistance gene Mi from tomato is a member of the leucine zipper, nucleotide
binding, leucine-rich repeat family of plant genes. *Plant Cell* **10**(8): 1307-1319.

Mittler R, Herr EH, Orvar BL, van Camp W, Willekens H, Inze D, Ellis BE. 1999. Transgenic

tobacco plants with reduced capability to detoxify reactive oxygen intermediates are hyperresponsive to pathogen infection. *Proc Natl Acad Sci U S A* **96**(24): 14165-14170.

Oliver JE, Whitfield AE. 2016. The Genus Tospovirus: Emerging Bunyaviruses that Threaten Food Security. *Annu Rev Virol* **3**(1): 101-124.

Padmanabhan MS, Ma S, Burch-Smith TM, Czymmek K, Huijser P, Dinesh-Kumar SP. 2013. Novel positive regulatory role for the SPL6 transcription factor in the N TIR-NB-LRR receptor-mediated plant innate immunity. *PLoS Pathog* **9**(3): e1003235.

Palma K, Zhang Y, Li X. 2005. An importin alpha homolog, MOS6, plays an important role in plant innate immunity. *Curr Biol* **15**(12): 1129-1135.

Peiro A, Canizares MC, Rubio L, Lopez C, Moriones E, Aramburu J, Sanchez-Navarro J. 2014. The movement protein (NSm) of Tomato spotted wilt virus is the avirulence determinant in the tomato Sw-5 gene-based resistance. *Mol Plant Pathol* **15**(8): 802-813.

Qi D, DeYoung BJ, Innes RW. 2012. Structure-function analysis of the coiled-coil and leucine-rich repeat domains of the RPS5 disease resistance protein. *Plant Physiol* **158**(4): 1819-1832.

Qi D, Innes RW. 2013. Recent Advances in Plant NLR Structure, Function, Localization, and Signaling. *Front Immunol* **4**: 348.

Rao ALNC, Y. G，Khan, J. A，Dijkstra, J. 2002. Molecular biology of plant virus movement. *Plant Viruses As Molecular Pathogens*.

Sarris PF, Cevik V, Dagdas G, Jones JD, Krasileva KV. 2016. Comparative analysis of plant immune receptor architectures uncovers host proteins likely targeted by pathogens. *BMC Biol* **14**: 8.

Sarris PF, Duxbury Z, Huh SU, Ma Y, Segonzac C, Sklenar J, Derbyshire P, Cevik V, Rallapalli G, Saucet SB, Wirthmueller L, Menke FL, Sohn KH, Jones JD. 2015. A Plant Immune Receptor Detects Pathogen Effectors that Target WRKY Transcription Factors. *Cell* **161**(5): 1089-1100.

Scholthof KB, Adkins S, Czosnek H, Palukaitis P, Jacquot E, Hohn T, Hohn B, Saunders K, Candresse T, Ahlquist P, Hemenway C, Foster GD. 2011. Top 10 plant viruses in molecular plant pathology. *Mol Plant Pathol* **12**(9): 938-954.

Seo JK, Kwon SJ, Cho WK, Choi HS, Kim KH. 2014. Type 2C protein phosphatase is a key regulator of antiviral extreme resistance limiting virus spread. *Sci Rep* **4**: 5905.

Seong K, Seo E, Witek K, Li M, Staskawicz B. 2020. Evolution of NLR resistance genes with noncanonical N-terminal domains in wild tomato species. *New Phytol* **227**(5): 1530-1543.

Shen QH, Saijo Y, Mauch S, Biskup C, Bieri S, Keller B, Seki H, Ulker B, Somssich IE, Schulze-Lefert P. 2007. Nuclear activity of MLA immune receptors links isolate-specific and basal disease-resistance responses. *Science* **315**(5815): 1098-1103.

Slootweg E, Roosien J, Spiridon LN, Petrescu AJ, Tameling W, Joosten M, Pomp R, van Schaik C, Dees R, Borst JW, Smant G, Schots A, Bakker J, Goverse A. 2010. Nucleocytoplasmic distribution is required for activation of resistance by the potato NB-LRR receptor Rx1 and is balanced by its functional domains. *Plant Cell* **22**(12): 4195-4215.

Soosaar JL, Burch-Smith TM, Dinesh-Kumar SP. 2005. Mechanisms of plant resistance to viruses. *Nat Rev Microbiol* **3**(10): 789-798.

Spassova M I PTW, Folkertsma R T, et al. 2001. The tomato gene Sw-5 is a member of the coiled coil, nucleotide binding, leucine-rich repeat class of plant resistance genes and confers resistance to TSWV in tobacco. *Molecular Breeding*. **7(2)**: 151.

Stewart M. 2007. Molecular mechanism of the nuclear protein import cycle. *Nat Rev Mol Cell Biol* **8**(3): 195-208.

Swiderski MR, Birker D, Jones JD. 2009. The TIR domain of TIR-NB-LRR resistance proteins is a signaling domain involved in cell death induction. *Mol Plant Microbe Interact* **22**(2): 157-165.

Takemoto D, Rafiqi M, Hurley U, Lawrence GJ, Bernoux M, Hardham AR, Ellis JG, Dodds PN, Jones DA. 2012. N-terminal motifs in some plant disease resistance proteins function in membrane attachment and contribute to disease resistance. *Mol Plant Microbe Interact* **25**(3): 379-392.

Takken FLW, Goverse A. 2012. How to build a pathogen detector: structural basis of NB-LRR function. *Current Opinion in Plant Biology* **15**(4): 375-384.

Taliansky M, Torrance L, Kalinina NO. 2008. Role of plant virus movement proteins. *Methods Mol Biol* **451**: 33-54.

Tameling WI, Nooijen C, Ludwig N, Boter M, Slootweg E, Goverse A, Shirasu K, Joosten MH. 2010. RanGAP2 mediates nucleocytoplasmic partitioning of the NB-LRR immune receptor Rx in the Solanaceae, thereby dictating Rx function. *Plant Cell* **22**(12): 4176-4194.

Tasset C, Bernoux M, Jauneau A, Pouzet C, Briere C, Kieffer-Jacquinod S, Rivas S, Marco Y, Deslandes L. 2010. Autoacetylation of the Ralstonia solanacearum effector PopP2 targets a lysine residue essential for RRS1-R-mediated immunity in Arabidopsis. *PLoS Pathog* **6**(11): e1001202.

Turina M, Kormelink R, Resende RO. 2016. Resistance to Tospoviruses in Vegetable Crops: Epidemiological and Molecular Aspects. *Annu Rev Phytopathol* **54**: 347-371.

van der Vossen EA, Gros J, Sikkema A, Muskens M, Wouters D, Wolters P, Pereira A, Allefs S. 2005. The Rpi-blb2 gene from Solanum bulbocastanum is an Mi-1 gene homolog conferring broad-spectrum late blight resistance in potato. *Plant J* **44**(2): 208-222.

van Wersch S, Li X. 2019. Stronger When Together: Clustering of Plant NLR Disease resistance Genes. *Trends Plant Sci* **24**(8): 688-699.

van Wersch S, Tian, L., Hoy, R., Li, X. . 2020. Plant NLRs: The whistleblowers of plant immunity. *Plant Commun*: doi: https://doi.org/10.1016/j.xplc.2019.100016.

Vos P, Simons G, Jesse T, Wijbrandi J, Heinen L, Hogers R, Frijters A, Groenendijk J, Diergaarde P, Reijans M, Fierens-Onstenk J, de Both M, Peleman J, Liharska T, Hontelez J, Zabeau M. 1998. The tomato Mi-1 gene confers resistance to both root-knot nematodes and potato aphids. *Nat Biotechnol* **16**(13): 1365-1369.

Vossen JH, van Arkel G, Bergervoet M, Jo KR, Jacobsen E, Visser RG. 2016. The Solanum demissum R8 late blight resistance gene is an Sw-5 homologue that has been deployed worldwide in late blight resistant varieties. *Theor Appl Genet* **129**(9): 1785-1796.

Wan L, Essuman K, Anderson RG, Sasaki Y, Monteiro F, Chung EH, Osborne Nishimura E, DiAntonio A, Milbrandt J, Dangl JL, Nishimura MT. 2019. TIR domains of plant immune receptors are NAD(+)-cleaving enzymes that promote cell death. *Science* **365**(6455): 799-803.

Wang A. 2015. Dissecting the molecular network of virus-plant interactions: the complex roles of host factors. *Annu Rev Phytopathol* **53**: 45-66.

Wang J, Chen T, Han M, Qian L, Li J, Wu M, Han T, Cao J, Nagalakshmi U, Rathjen JP, Hong Y, Liu Y. 2020. Plant NLR immune receptor Tm-2² activation requires NB-ARC domain-mediated self-association of CC domain. *PLoS Pathog* **16**(4): e1008475.

Wang J, Hu M, Wang J, Qi J, Han Z, Wang G, Qi Y, Wang HW, Zhou JM, Chai J. 2019a.
Reconstitution and structure of a plant NLR resistosome conferring immunity. *Science*
364(6435).

Wang J, Wang J, Hu M, Wu S, Qi J, Wang G, Han Z, Qi Y, Gao N, Wang HW, Zhou JM, Chai J.
2019b. Ligand-triggered allosteric ADP release primes a plant NLR complex. *Science*
364(6435).

Wen W, Meinkoth JL, Tsien RY, Taylor SS. 1995. Identification of a signal for rapid export of
proteins from the nucleus. *Cell* **82**(3): 463-473.

Wirthmueller L, Zhang Y, Jones JD, Parker JE. 2007. Nuclear accumulation of the Arabidopsis
immune receptor RPS4 is necessary for triggering EDS1-dependent defense. *Curr Biol* **17**(23):
2023-2029.

Zhang Y, Li X. 2005. A putative nucleoporin 96 Is required for both basal defense and constitutive
resistance responses mediated by suppressor of npr1-1,constitutive 1. *Plant Cell* **17**(4):
1306-1316.

Zhao W, Jiang L, Feng Z, Chen X, Huang Y, Xue F, Huang C, Liu Y, Li F, Liu Y, Tao X. 2016.
Plasmodesmata targeting and intercellular trafficking of Tomato spotted wilt tospovirus
movement protein NSm is independent of its function in HR induction. *J Gen Virol* **97**(8):
1990-1997.

Zhu M, Jiang L, Bai B, Zhao W, Chen X, Li J, Liu Y, Chen Z, Wang B, Wang C, Wu Q, Shen Q,
Dinesh-Kumar SP, Tao X. 2017. The Intracellular Immune Receptor Sw-5b Confers
Broad-Spectrum Resistance to Tospoviruses through Recognition of a Conserved 21-Amino
Acid Viral Effector Epitope. *Plant Cell* **29**(9): 2214-2232.

Zhu M, van Grinsven IL, Kormelink R, Tao X. 2019. Paving the Way to Tospovirus Infection:
Multilined Interplays with Plant Innate Immunity. *Annu Rev Phytopathol* **57**: 41-62.

1115

1116

1117

1118

 FIGURE LEGENDS

 Fig. 1. Subcellular localization of Sw-5b in *Nicotiana benthamiana* and tomato leaf

cells. (a) Schematic diagrams of Sw-5b. (b) Subcellular localizations of free YFP

(left), YFP-Sw-5b (middle) and autoactive YFP-Sw-5b^{D857V} mutant (right) in *N.*

benthamiana leaf cells at 24 hours post agro-infiltration (hpi). (c) Subcellular

localization of free YFP (left), YFP-Sw-5b (middle) and autoactive YFP-Sw-5b^{D857V}

mutant (right) in tomato leaf cells at 24 hpi. N nucleus, and C cytoplasm inside the

cell are indicated. Bar = 10 μm. (d) Nucleocytoplasmic partitioning analysis of

YFP-Sw-5b and autoactive YFP-Sw-5b^{D857V}. Total lysate (T) from p2300S empty

vector (EV), YFP-Sw-5b or YFP-Sw-5b^{D857V} expressing leaves were fractionated into

cytoplasm and nucleus, and analyzed by immunoblots using antibodies against YFP.

The actin and histone were used as a cytoplasm marker and nucleus marker,

respectively, in the fractionation analysis. Ponceau S staining was also used as

cytoplasm marker.

1133

Fig. 2. Effect of Sw-5b subcellular localization pattern on HR induction. (a) Confocal

images of *N. benthamiana* leaf cells transiently expressing NES-YFP-Sw-5b,

nes-YFP-Sw-5b, NLS-YFP-Sw-5b or nls-YFP-Sw-5b fusion. The images were taken

at 24–36 hpi. N nucleus and C cytoplasm (c). Bar = 10 μm. (b) Induction of HR in *N.*

benthamiana leaf tissues co-expressing NSm and one of the five Sw-5b fusion

proteins. The infiltrated *N. benthamiana* leaf was photographed at 3 dpi (left image).

Induction of HR in the infiltrated tissues were visualized using a trypan blue staining

method (right image). (c) Immunoblot analysis of NES-YFP-Sw-5b, nes-YFP-Sw-5b,

NLS-YFP-Sw-5b, and nls-YFP-Sw-5b expressions in the infiltrated *N. benthamiana*

leaf tissues. These fusion proteins were enriched using the GFP-Trap beads prior to

SDS-PAGE, and the blot was probed using an YFP specific antibody. Ponceau-S

staining was used to estimate sample loadings. (d) Time course analysis of ion

leakage in Nicotiana benthamiana leaves co-expressing NSm with one of the five

Sw-5b fusion proteins. Measurements were performed at 4 h intervals starting from

24 to 48 hpi. Error bars (SEs) were calculated using the results from three biological

replicates per treatment collected at each time point.

1150

Fig. 3. The effect of cytoplasm- and nucleus-targeted Sw-5b on viral replication. (a)

Schematic representation of binary constructs to express TSWV $SR_{(-)eGFP}$

mini-genome replicon, TSWV L RNA segment containing an optimized RdRp and

$NSm^{H93A\&H94A}$ mutant that defected in viral movement. Minus sign (-) and 5′ to 3′

designation represent the negative (genomic)-strand of tospovirus RNA. 35S: a

double 35S promoter; HH: hammerhead ribozyme; RZ: hepatitis delta virus (HDV)

ribozyme; NOS: nopaline synthase terminator; 35S Ter: a 35S transcription terminator.

(b) Accumulation of eGFP fluorescence in *N. benthamiana* leaves co-expressing

p2300S empty vector (EV), Sw-5b, NES-Sw-5b, nes-Sw-5b, NLS-Sw-5b, or

nls-Sw-5b with TSWV $SR_{(-)eGFP}$, L, and $NSm^{H93A\&H94A}$ at 4 days post infiltration (dpi)

viewed with a fluorescence microscope. Bar represents 400 μm. (c) Immunoblot

analysis of expression of eGFP proteins in leaves shown in panel (b) using specific

antibodies against YFP. Ponceau S staining of rubisco large subunit is shown for

protein loading control.

1165

Fig. 4. Effect of subcellular localization of Sw-5b on cell-to-cell movement of NSm

in leaf epidermis of *N. benthamiana.* (a) Schematic diagram of the binary construct to

co-express mCherry-HDEL and NSm-GFP. (b) Cell-to-cell movement analysis of

NSm-GFP in *N. benthamiana* leaves co-expressing p2300S empty vector (EV), Sw-5b,

NES-Sw-5b, NLS-Sw-5b, or nls-Sw-5b with the construct harboring both

mCherry-HDEL and NSm-GFP. *Agrobacterium* containing the construct to co-express

mCherry-HDEL and NSm-GFP was diluted 500 times for expression in a single

epidermal cell. All other *Agrobacterium* were infiltrated at the concentration of OD_{600}

= 0.2. Bar, 50 μm.

1175

Fig. 5. Analysis of cytoplasm- and nucleus-targeted Sw-5b-mediated host immunity

to TSWV systemic infection. (a) TSWV systemic infection in transgenic *N.*

benthamiana plants expressing NES-YFP-Sw-5b, nes-YFP-Sw-5b, NLS-YFP-Sw-5b,

nls-YFP-Sw-5b, YFP-Sw-5b or p2300S empty vector (EV) driven by 35S promoter.

TSWV-inoculated plants were photographed at 15 dpi. White arrow indicates the

systemic leaves showing HR trailing. White arrowhead indicates the systemic leaves

showing mosaic. (b) Immunoblot analysis of NES-YFP-Sw-5b, nes-YFP-Sw-5b,

NLS-YFP-Sw-5b, nls-YFP-Sw-5b and YFP-Sw-5b expressions in different transgenic

N. benthamiana plants. EV plants transformed with an empty vector and were used as

a negative control. (c) RT-PCR analysis of TSWV accumulation in the systemic

leaves of different transgenic *N. benthamiana* plants at 15 dpi.

1187

Fig. 6. Joint effects of cytoplasm- and nucleus-targeted Sw-5b on defenses against

tospovirus infection in *Nicotiana benthamiana* leaves. (a) Schematic representation of

binary constructs to express TSWV $SR_{(+)eGFP}$-$M_{(-)op}$ and TSWV L RNA segment

containing an optimized RdRp. 35S: a double 35S promoter; HH: hammerhead

ribozyme; RZ: hepatitis delta virus (HDV) ribozyme; NOS: nopaline synthase

terminator. (b) Accumulation of eGFP fluorescence in *N. benthamiana* leaves

co-expressing p2300S empty vector (EV), Sw-5b, NES-Sw-5b, NLS-Sw-5b, or

NES-Sw-5b+NLS-Sw-5b with TSWV $SR_{(+)eGFP}$-$M_{(-)op}$ at 4 days post infiltration (dpi)

viewed with a fluorescence microscope. Bar represents 400 μm. (c) Immunoblot

analysis of expression of eGFP proteins in leaves shown in panel (b) using specific

antibodies against YFP. Ponceau S staining of rubisco large subunit is shown for

protein loading control. (d) Quantification of eGFP proteins in leaves shown in panel

(c).

1201

Fig. 7. Functional analysis and subcellular localization patterns of individual or

combined Sw-5b domains. (a) Schematic diagrams showing a full length Sw-5b or

Sw-5b domains fused with YFP. (b) Confocal images of *N. benthamiana* leaf

epidermal cells expressing various YFP fusions. Images of the cells were taken at 24

hpi. N nucleus, Nu nucleolus, and C cytoplasm. Bar = 10 μm. (c) TSWV-inoculated

transgenic *N. benthamiana* plants expressing these various YFP fusions and

photographed at 15 dpi.

1209

Fig. 8. Roles of *importins* α and β in YFP-Sw-5b nucleus targeting and

Sw-5b-mediated immunity to TSWV systemic infection. (a) Transient expression of

YFP-Sw-5b in *N. benthamiana* leaf epidermal cells silenced for *importin*

$\alpha 1$ (IMP α1 KD)*, importin* $\alpha 2$ (IMP α2 KD), *importin* $\alpha 1$ and $\alpha 2$ (IMP α1 & α2 KD),

importin β (IMP β KD) or *importin* $\alpha 1$ and $\alpha 2$ and β (IMP α1 & α2 & β KD) through

VIGS. Images of the cells were captured using a confocal microscope at 26 hpi. N

nucleus, C cytoplasm. Bar = 10 μm. (b) YFP-Sw-5b transgenic *N. benthamiana* plants

were silenced for *importin* $\alpha 1$*, importin* $\alpha 2$*, importin* $\alpha 1$ and $\alpha 2$, *importin* β or

importin $\alpha 1$ and $\alpha 2$ and β expression through VIGS followed by inoculation with

TSWV. TSWV-inoculated YFP-Sw-5b transgenic *N. benthamiana* plants were

photographed at 15 dpi. White arrowhead indicates HR trailing in systemic leaves.

1221

Fig. 9. A working model for Sw-5b. Sw-5b furcates disease resistances by proper

nucleocytoplasmic partition to block different infection steps of tomato spotted wilt

tospovirus. Sw-5b switched from the autoinhibited state to an activated state upon

recognition of NSm in the cytoplasm. Cytoplasm portion of Sw-5b induce cell death

and defense that inhibit viral replication. The activated Sw-5b also translocated into

nucleus via *importins* α and β. Nucleus-localized Sw-5b induces a defense that block

viral cell-to-cell and long-distance movement. Cytoplasm- and nucleus-localized

Sw-5b have additively effects on defense to inhibit viral replication, intercellular and

long-distance movement during tospovirus infection.

1231

Supporting Information

Short legends

Fig. S1 Sw-5b recognizes TSWV NSm in cytoplasm.

Fig. S2 Analysis of virus replication monitoring system using a TSWV-based mini-genome replicon and a movement defective NSm mutant.

Fig. S3 Effects of nes-Sw-5b and nls-Sw-5b on NSm-GFP cell-to-cell movement and effects of Sw-5b and EV on NSm^{T120N}-GFP cell-to-cell movement.

Fig. S4 Effects of cytoplasmic and nuclear Sw-5b on host immunity to TSWV systemic infection.

Fig. S5 Cytoplasmic and nuclear Sw-5b activity on TSWV-GFP cell-to-cell movement in *N. benthamiana* leaves.

Fig. S6 An immunoblot showing the accumulations of various YFP-tagged proteins expressed in different transgenic *N. benthamiana* plants and RT-PCR analysis of TSWV accumulation in the systemic leaves.

Fig. S7 Bimolecular fluorescence complementation (BiFC) assay of cYFP-SD, cYFP-Sw-5b and nYFP-Importin α1 (nYFP-IMP α1), nYFP-Importin α2 (nYFP-IMP α2), nYFP-Importin β (nYFP-IMP β) interaction in *N. benthamiana* leaf epidermal cells.

Fig. S8 RT-PCR analyses of *importin a1*, *a2* and *b* expressions in the assayed plants and their effects on TSWV systemic infection.

Table S1. List of primers used in this study.

Table S2. Response of six different types of transgenic *Nicotiana benthamina* plants

driven by 35S promoter to TSWV infection.

Table S3. Response of six different types of transgenic *Nicotiana benthamina* plants

driven by Sw-5b native promoter to TSWV infection.

Table S4. Response of six different types of transgenic *Nicotiana benthamina* plants

to TSWV infection.

Table S5. Mass spectrum data of YFP-SD

Table S6. Mass spectrum data of YFP-Sw-5b

1261

Cytoplasmic and nuclear Sw-5b NLR act both independently and synergistically to dictate full host defense against tospovirus infection

Hongyu Chen[1,¶], Xin Qian[1,¶], Xiaojiao Chen[1,¶], Tongqing Yang[1], Mingfeng Feng[1], Jing Chen[1],

Ruixiang Cheng[1], Hao Hong[1], Ying Zheng[1], Yuzhen Mei[4], Danyu Shen[1], Yi Xu[1], Min Zhu[1], Xin

Shun Ding[1] and Xiaorong Tao[1,*]

[1] Key Laboratory of Plant Immunity, Department of Plant Pathology, College of Plant Protection,

Nanjing Agricultural University, Nanjing 210095, P. R. China.

[2] Huaiyin Institute of Agricultural Sciences of Xuhuai Region in Jiangsu，Huaian 223001, Jiangsu,

P. R. China.

[3] College of Plant Protection, Yunnan Agricultural University, Kunming 650201, Yunnan, P. R.

China.

[4] State Key Laboratory of Rice Biology, Institute of Biotechnology, Zhejiang University,

Hanghzou 310029, P. R. China.

[¶] These authors contributed equally to this work.

Author for correspondence:

Xiaorong Tao

Tel: (+86)-25-84399027

Email: taoxiaorong@njau.edu.cn

RUNNING HEAD

Independent role of cytoplasmic and nuclear Sw-5b in immunity

Word Count: Main body of the text, 8136

Title 18, Summary 249, Keywords 8, Introduction 1286, Materials and Methods 1562, Results 3030, Discussion 1884, Acknowledgments 35, References 3687, Figure Legends 1103, Supporting Information Legends 786

9 Figures (Colors); Supporting information: 8 Supplemental Figures, 6 Supplemental Tables

Summary

● Plant intracellular nucleotide binding-leucine-rich repeat (NLR) receptors play critical roles in mediating host immunity to pathogen attack. We use tomato Sw-5b::tospovirus as a model system to study the specific role of the compartmentalized plant NLR in dictating host defense against virus at different infection steps.

● We demonstrated here that tomato NLR Sw-5b translocates to cytoplasm and nucleus, respectively, to play different roles in inducing host resistances against *Tomato spotted wilt tospovirus* (TSWV) infection. The cytoplasmic Sw-5b functions to induce a strong cell death response to inhibit TSWV replication. This host response is, however, insufficient to block viral intercellular and long-distance movement. The nucleus-localized Sw-5b triggers a host defense that weakly inhibits viral replication but strongly impedes virus intercellular and systemic movement. Furthermore, the cytoplasmic and nuclear Sw-5b act synergistically to dictate full host defense to TSWV infection.

● We further demonstrated that the extended N-terminal *Solanaceae* domain (SD) of Sw-5b plays critical roles in cytoplasm/nucleus partitioning. Sw-5b nucleotide-binding leucine-rich repeat (NB-LRR) controls its cytoplasm localization. Strikingly, the SD but not coil-coil (CC) domain is crucial for Sw-5b receptor to translocate from cytoplasm to nucleus to trigger the immunity. The SD was found to interact with importins. Silencing both importin α and β expression disrupted Sw-5b nucleus translocation and host immunity

against TSWV systemic infection.

● Collectively, our findings suggest that Sw-5b bifurcates disease resistances

by cytoplasm/nucleus partitioning to block different infection steps of TSWV.

The findings also identified a new regulatory role of extra domain of a plant NLR

in mediating host innate immunity.

Keywords:

NLRs, Cytoplasm, Nuclear, Plant innate immunity, Tomato spotted wilt virus,

Replication, Cell-to-cell movement and Long distance movement

Introduction

Plant innate immunity plays critical roles in host defense against pathogen invasions, and is triggered by cell-surface receptors or intracellular nucleotide-binding leucine-rich repeat (NLR) receptors (Soosaar *et al.*, 2005; Dodds & Rathjen, 2010; Cui *et al.*, 2015; Li *et al.*, 2015; Jones *et al.*, 2016; Kourelis & van der Hoorn, 2018; Kapos *et al.*, 2019; van Wersch, 2020). Plant intracellular NLRs are the largest classes of resistance proteins that function to detect pathogen effectors, and to activate host immunity upon pathogen attack (Caplan, J *et al.*, 2008; Takken & Goverse, 2012; Li *et al.*, 2015; Jones *et al.*, 2016; Kourelis & van der Hoorn, 2018; Kapos *et al.*, 2019). Plant NLRs typically contain an N-terminal domain, a central nucleotide-binding domain (NB), a nucleotide-binding adaptor (ARC domain shared by Apaf-1, certain resistance proteins, and CED-4), and a C-terminal leucine-rich repeat (LRR) domain (Ea & Jones, 1998; Jones *et al.*, 2016; Ma *et al.*, 2018; Wang *et al.*, 2019a; Wang *et al.*, 2019b; Ma *et al.*, 2020). Based on the differences among the N-terminal domains, plant NLRs can be further divided into two main categories, known as the coiled-coil NLR (refers to as CNL) category and the Toll/interleukin-1 receptor NLR (TNL) category (Meyers *et al.*, 2003; Collier & Moffett, 2009; Qi & Innes, 2013). The CC- or the TIR-domain-bearing NLRs have distinct genetic requirements and can regulate different functions in the downstream of defense signaling (Collier & Moffett, 2009; Qi & Innes, 2013; Horsefield *et al.*, 2019; Jubic *et*

al., 2019; van Wersch & Li, 2019; Wan *et al.*, 2019).

In addition to classical domains, non-canonical domains were frequently found to

integrate into certain NLRs. The additional domain, called BED, was first found in 32

poplar NLR proteins (Germain & Seguin, 2011). This extra BED domain was also

found in nine rice NLRs (Das *et al.*, 2014). The RATX1/HMA domain in the rice

NLRs RGA5 and Pik‑1 was found to act as integrated decoys to detect the cognate

pathogen effectors (Kanzaki *et al.*, 2012; Cesari *et al.*, 2013; Cesari *et al.*, 2014). The

WRKY domain on Arabidopsis thaliana NLR RRS1 was further found to function as

an integrated decoy that recognizes the effectors AvrRps4 and PopP2 (Le Roux *et al.*,

2015; Sarris *et al.*, 2015). Genome-wide analyses of plant NLR receptors revealed

that about 3.5 % of the NLRs carried specific non-canonical domains (Cesari *et al.*,

2014; Kroj *et al.*, 2016; Sarris *et al.*, 2016), and some of these non-canonical domains

were shown to be targeted by pathogen effectors during pathogen infections (Sarris *et*

al., 2016). However, molecular functions of most non-canonical domains in plant

NLRs remain largely unexplored.

Translocations of plant NLRs into proper subcellular compartments are critical for

the induction of innate immunity (Cui *et al.*, 2015; van Wersch, 2020). Multiple plant

NLRs and immune regulators, including tobacco N, Arabidopsis snc1, RRS1/RPS4,

barley MLA10, and Arabidopsis EDS1, NPR1, have been shown to accumulate in

both cytoplasm and nucleus, and for several nucleocytoplasmic NLRs accumulation

in nucleus is required for triggering host resistance to pathogen infections (Deslandes

et al., 2003; Burch-Smith *et al.*, 2007; Shen *et al.*, 2007; Wirthmueller *et al.*, 2007;

Tasset *et al.*, 2010; Bai *et al.*, 2012; Inoue *et al.*, 2013; Padmanabhan *et al.*, 2013).

Wheat Sr33, a homolog of barley MLA10, however, was reported to accumulate in

cytoplasm to induce host resistance against stem rust pathogen (Cesari *et al.*, 2016).

For potato Rx, a balanced cytoplasm and nucleus accumulation of Rx is needed to

induce the host immunity (Slootweg *et al.*, 2010; Tameling *et al.*, 2010). Other studies

have shown that *Arabidopsis* Rpm1 (Gao *et al.*, 2011), RPS2 (Axtell & Staskawicz,

2003), RPS5 (Qi *et al.*, 2012), rice Pit (Takemoto *et al.*, 2012), and tomato Tm-2^2

(Chen *et al.*, 2017; Wang *et al.*, 2020) need to associate with plasma membrane in

order to trigger cell death and host immunity. Latest studies have shown that the

activated *Arabidopsis* ZAR1 can bind to cellular membrane, leading to a membrane

leakage followed by cell death and host immunity (Wang *et al.*, 2019a; Wang *et al.*,

2019b). Flax L6 and M have been shown to accumulate in both Golgi apparatus and

tonoplast, and these compartmentalized localizations are necessary for the induction

of host resistance (Kawano *et al.*, 2014). The re-distribution of potato R3a from

cytosol to endosomal compartments is crucial for the induction of host resistance to

Phytophthora infestans infection (Engelhardt *et al.*, 2012). Different plant NLRs have

diverse subcellular localizations for their proper functions. However, how the

compartmentalized plant NLRs specifically dictate defense signaling remains largely

unknown.

Tomato spotted wilt tospovirus (TSWV) is one of most destructive plant NSVs,

infecting more than 1000 plant species, and causes crop losses more than one billion

US dollars annually worldwide (Kormelink *et al.*, 2011; Scholthof *et al.*, 2011; Oliver

& Whitfield, 2016). Tomato NLR Sw-5b confers strong resistance to TSWV infection

and has been widely used in tomato breeding projects to produce tospovirus resistant

tomato cultivars (Brommonschenkel *et al.*, 2000; Spassova M I, 2001; Turina *et al.*,

2016; Zhu *et al.*, 2019). Upon recognition of TSWV movement protein, NSm, Sw-5b

can trigger a hypersensitive response (HR), which typically associated with localized

cell death (Lopez *et al.*, 2011; Hallwass *et al.*, 2014; Peiro *et al.*, 2014; De Oliveira *et*

al., 2016; Zhao *et al.*, 2016; Leastro *et al.*, 2017). Tospoviruses are divided into

American and Euro-Asia type based on their geographic distribution and amino acid

sequence identity of viral nucleocapsid protein. We have previously shown that

Sw-5b can confer a broad-spectrum resistance to American type tospoviruses,

including TSWV, through recognition of a conserved 21 amino acid PAMP-like

region in the viral movement protein NSm (Zhu *et al.*, 2017). Sw-5b carries an

extended N-terminal *Solanaceae* domain (SD), a CC domain, a NB-ARC domain, and

a LRR domain (Brommonschenkel *et al.*, 2000; Spassova M I, 2001;

Lukasik-Shreepaathy *et al.*, 2012). Similar SDs have also been found in the Mi-1.2,

R8, Rpi-blb2, and Hero (Milligan *et al.*, 1998; Vos *et al.*, 1998; Ernst *et al.*, 2002; van

der Vossen *et al.*, 2005; Lukasik-Shreepaathy *et al.*, 2012; Vossen *et al.*, 2016). More

recently, Seong and others reported that the extended CNL has been evolved initially

in the ancestor of *Asterids* and *Amaranthaceae*, predated the *Solanaceae* family

(Seong *et al.*, 2020). In the presence of the extended N-terminal SD, Sw-5b is in an

autoinhibited state through multilayered interactions between SD, CC, NB-ARC, and

LRR domains (Chen *et al.*, 2016). For activation, the extra SD also recognizes NSm.

Sw-5b adopts a two-step NSm recognition strategy through SD and then LRR domain

(Li *et al.*, 2019). This two-step recognition mechanism significantly enhances the

sensitivity of the detection on TSWV NSm (Li *et al.*, 2019). Although Sw-5b is

known to localize in both cytoplasm and nucleus (De Oliveira *et al.*, 2016), the

biological roles of the cytoplasm- and the nucleus-accumulated Sw-5b in host

immunity signaling are unknown.

In this study, we investigated the subcellular distribution pattern of Sw-5b and the

functions of the compartmentalized Sw-5b in the induction of host immunity to

TSWV infection. We determined here that cytoplasm- and nucleus-accumulated

Sw-5b functions differently in inducing host defense response to inhibit multiple

tospovirus infection steps. The cytoplasmic Sw-5b can induce a strong cell death

response to suppress TSWV replication, whereas the nucleus-accumulated Sw-5b can

induce a strong defense against viral intercellular movement and systemic infection.

The combination of cytoplasmic and nuclear Sw-5b induces a synergistic and strong

plant immunity against tospovirus infection. We also found that the extended SD

functions as the key regulator for this critical intracellular translocation. The SD was

also found to interact with importins α and β to mediate Sw-5b nucleus translocation,

and to confer the full host immunity against tospovirus infection.

Materials and Methods

Plasmid construction

p2300S-YFP-Sw-5b was from a previously described source (Chen *et al.*, 2016).

Different domains of Sw-5b were PCR-amplified from p2300S-Sw-5b (Chen *et al.*,

2016) and cloned individually behind the *YFP* gene in the p2300S vector using a

two-step overlap PCR procedure as described (Li *et al.*, 2019). All the primers used in

this study are listed in Table S1.

To visualize the subcellular localization patterns of various fusion proteins, a SV40

T-Ag-derived nuclear localization signal (NLS, QPKKKRKVGG) (Lanford & Butel,

1984) or a PK1 nuclear export signal (NES, NELALKLAGLDINK) (Wen *et al.*, 1995)

was fused to the N-terminus of YFP-Sw-5b or the C-terminus of NSm-YFP, as

described (Kong *et al.*, 2017), to produce pNES-YFP-Sw-5b, pNLS-YFP-Sw-5b,

pNSm-YFP-NES, and pNSm-YFP-NLS, respectively. In addition, YFP-Sw-5b and

NSm-YFP were fused individually with a mutant NLS (nls, QPKKTRKVGG) or a

mutant NES (nes, NELALKAAGADANK) to produce pnes-YFP-Sw-5b,

pnls-YFP-Sw-5b, pNSm-YFP-nes, and pNSm-YFP-nls. The constructs were then

transformed individually into *Agrobacterium tumefaciens* strain GV3101 cells.

Transient gene expression, stable plant transformation, and virus inoculation

Nicotiana benthamiana were grown in soil in pots inside a greenhouse maintained at

25°C and a 16 h light/8 h dark photoperiod. Six-to-eight week-old *N. benthamiana*

plants were used for various assays. Transient gene expression assays were performed

in *N. benthamiana* leaves through agro-infiltration using Agrobacterium cultures

carrying specific expressing constructs as described previously (Feng *et al.*, 2016; Ma

et al., 2017). Transgenic *N. benthamiana* lines expressing YFP-Sw-5b or its

derivatives were made using constructs with a 35S promoter or a Sw-5b native

promoter via a standard leaf-disc transformation method (Chen *et al.*, 2016). The

resulting transgenic *N. benthamiana* lines were named as NES-YFP-Sw-5b,

NLS-YFP-Sw-5b, nes-YFP-Sw-5b, nls-YFP-Sw-5b, YFP-Sw-5b, and EV

(transformed with an empty vector), respectively. Inoculation of transgenic *N.*

benthamiana plants with TSWV was done by rubbing plant leaves with TSWV-YN

isolate-infected crude saps as described (Zhu *et al.*, 2017). TRV-mediated VIGS in *N.*

benthamiana plants was done as described (Ma *et al.*, 2015). The agro-infiltrated or

the virus-inoculated plants were growning inside a growth chamber maintained at

25/23 °C (day/night) with a 16/8 h light and dark photoperiod.

Particle bombardment

The particle bombardment is described (Feng et al., 2016). Briefly, 60 mg Tungsten

M-10 microcarrier (Bio-RAD) was placed into a 1.5 ml Eppendorf tube with 1 mL

70% ethanol. The tube was vortexes for 3 minutes, and then stood at room

temperature for 15 minutes. After centrifuge at low speed for 5 seconds, the

supernatant was removed and the pellet was rinsed with 70% ethanol for 3 times. One

mL 50% sterile glycerol solution was added and divided Tungsten M-10 microcarrier

into 50 µl. Five µg pRTL2-YFP, pRTL2-YFP-Sw-5b or pRTL2-YFP-Sw-5bD857V

plasmid DNA, 50 µl of 2.5 M CaCl2, and 20 µl of 0.1 M spermidine, respectively

were added and mixed with microcarrier. After centrifuge at low speed for 5 seconds

and the supernatant removed. The pellet was resuspended in 200 µl 70% ethanol and

centrifugation as described above. Use 48 µl of 100% ethanol to resuspend the

tungsten particle::plasmid DNA complexes, and load 15 48 µl mixture onto the center

of carrier (Bio-RAD), air dry, and use He/1000 particle transport system (BIO-RAD)

to bombard tomato leaves harvested from 3- or 4-week-old of Money Marker. The

bombarded leaves were incubated in Petri dishes for 24 hours at 25°C followed with

Confocal Microscope analysis.

Trypan blue staining

N. benthamiana leaves were harvested at 3 days post agro-infiltration (dpai) and

boiled for 5 min in a 1.15:1 (v/v) mixed ethanol and trypan blue staining solution (10

g phenol, 10 mL glycerol, 10 mL lactic acid, and 20 mg trypan blue in 10 mL distilled

water). The stained leaves were then de-stained in a chloral hydrate solution (2.5 g per

mL distilled water) as described (Bai *et al.*, 2012).

Electrolyte leakage assay

Electrolyte leakage assay was performed as previously described (Mittler *et al.*, 1999;

Zhu *et al.*, 2017) with slight modifications. Briefly, five leaf discs (9 mm in diameter

each) were taken from the agro-infiltrated leaves per treatment and at various dpai.

The harvested leaf discs from a specific treatment were floated on a 10 mL distilled

water for 3 h at room temperature (RT), and the conductivity of each bathing water

was measured (referred to as value A) using a Multiparameter Meter as instructed

(Mettler Toledo, Zurich, Switzerland). After the first measurement, the leaf discs were

returned to the bathing water and incubated at 95°C for 25 min. After cooling down to

RT, the conductivity of each bathing sample was measured again (referred to as value

B). The ion leakage was expressed as the ratio determined by value A/value B × 100.

The mean value and standard error of each treatment were calculated using the data

from three biological replicates per treatment at each sampling time point.

Confocal laser scanning microscopy

Tissue samples were collected from the leaves of transiently expressing YFP-Sw-5b

or one of the fusion proteins at 24–36 hours post agro-infiltration (hpai). The collected

tissue samples were mounted in water between a glass slide and a coverslip. Images

of individual samples were captured under a Carl Zeiss LSM 710 confocal laser

scanning microscope. YFP fusions were excited at 488 nm and the emission was

captured at 497–520 nm. The resulting images were further processed using the Zeiss

710 CLSM software followed by the Adobe Photoshop software (San Jose, CA,

USA).

Nucleus and cytoplasm fractionations

N. benthamiana leaf tissues (1 g per sample), representing a specific treatment, were

collected at 24 hpai, frozen in liquid nitrogen, ground to fine powders, and then

homogenized in 2 mL (per sample) extraction buffer 1 (20 mM Tris-HCl, pH 7.5, 20

mM KCl, 2.5 mM MgCl$_2$, 2 mM EDTA, 25% glycerol, 250 mM sucrose, 1×Protease

Inhibitor Cocktail, and 5 mM DTT). The resulting lysate was filtered through 30 μm

filters to remove debris, and the filtrate was centrifuged at 2,000 $\times$ g for 5 minutes

to pellet nuclei. The supernatant from a sample was transferred into a new tube and

centrifuged at 10,000$\times g$ for 10 min. The resulting supernatant was used as the

cytoplasm fraction. The nuclei containing pellet was resuspended in 5 mL extraction

buffer 2 (20 mM Tris-HCl, pH 7.4, 25% glycerol, 2.5 mM MgCl2, and 0.2% Triton

X-100), centrifuged for 10 min at 2,000 $\times$ g followed by 4-6 cycles of resuspension

and centrifugation as described above. The resulting pellet was resuspended again in

500 μl extraction buffer 3 (20 mM Tris-HCl, pH 7.5, 0.25 M sucrose, 10 mM MgCl2,

0.5% Triton X-100, and 5 mM β-mercaptoethanol). The nuclei fraction was carefully

layered on the top of 500 mL extraction buffer 4 (20 mM Tris-HCl, pH 7.5, 1.7 M

sucrose, 10 mM MgCl2, 0.5% Triton X-100, 1$\times$Protease Inhibitor Cocktail, and 5

mM β-mercaptoethanol), and then centrifuged at 16,000 $\times$ g for 1 h. The resulting

pellet was resuspended in 500 μL extraction buffer 1 and stored at −80°C until use or

used immediately for SDS-PAGE assays. All the processes were performed on ice or

at 4°C. In this study, actin and histone H3 were used as the cytoplasmic and the

nuclear markers, respectively.

Western blot, co-immunoprecipitation and mass spectrometry analysis

Western blot and co-immunoprecipitation assays were performed as described (Zhu *et*

al., 2017). Briefly, agro-infiltrated leaf samples (1 g per sample) were harvested and

homogenized individually in pre-chilled mortars with pestles in 2 mL extraction

buffer [10% (v/v) glycerol, 25 mM Tris, pH 7.5, 1 mM EDTA, 150 mM NaCl, 10 mM

DTT, 2% (w/v) polyvinylpolypyrrolidone, and 1 × protease inhibitor cocktail (Sigma,

Shanghai, China)]. Each crude slurry was transferred into a 2 mL Eppendorf tube, and

spun for 2 min at full speed in a refrigerated microcentrifuge. The supernatant was

transferred into a clean 1.5 mL Eppendorf tube and spun for 10 min at 4°C. For

Western blot assays, 50 μL supernatant from a sample was mixed with 150 μL

Laemmli buffer, boiled for 5 min, and analyzed in SDS-PAGE gels through

electrophoresis. For immunoprecipitation assays, 1 mL supernatant was mixed with

25 μL GFP-trap agarose beads (ChromoTek, Planegg-Martinsried, Germany),

incubated for 2 h at 4°C on an orbital shaker, and then pelleted through low speed

centrifugation. The blots were probed with a 1:2,500 (v/v) diluted anti-YFP antibody

or other specific antibodies followed a 1:10,000 (v/v) diluted horseradish peroxidase

(HRP)-conjugated goat anti-rabbit or a goat anti-mouse antibody (Sigma-Aldrich, St.

Louis, MO, USA). The detection signal was developed using the ECL substrate kit as

instructed (Thermo Scientific, Hudson, NH, USA).

For mass spectrometry analysis, the immunoprecipitation samples of YFP-SD

and SD without tag were processed by The Beijing Genomics Institute (BGI) for mass

spectrometry analysis. The immunoprecipitation samples of YFP-Sw-5b and Sw-5b

without tag were processed by Applied Protein Technology in Shanghai. Database

searches were performed using the Mascot search engine against *N. benthamiana*.

(https://solgenomics.net/organism/Nicotiana_benthamiana/genome).

RT-PCR detection of TSWV infection

Total RNA was extracted from TSWV-inoculated *N. benthamiana* plant leaves using

an RNA Purification Kit (Tiangen Biotech, Beijing, China), and then treated with

RNase-free DNase I (TaKaRa, Dalian, China). First-strand cDNA was synthesized

using a TSWV-specific primer (S3 Table). PCR reactions were as follows: initial

denaturation at 94°C for 2 min followed by 35 cycles of 94°C for 30 s, 52°C for 30 s,

and 72°C for 1 min. The final extension was 72°C for 10 min. The resulting PCR

products were visualized in 1.0% (w/v) agarose gels through electrophoresis.

342   Results

Determination of Sw-5b subcellular localization pattern

Expression of YFP-Sw-5b in *N. benthamiana* leaves resulted in a strong HR cell death

as well as Sw-5b (Chen *et al.*, 2016; Zhu *et al.*, 2017). To investigate the subcellular

localization pattern of Sw-5b, we transiently expressed YFP and YFP-Sw-5b in *N.*

benthamiana leaves, respectively, through agro-infiltration. Confocal Microscopy

results showed that the YFP-Sw-5b fusion accumulated in both cytoplasm and nucleus

in *N. benthamiana* leaf cells (Fig. 1b, middle image). This subcellular localization

pattern was similar to that of YFP (Fig. 1b, left image). When a D857V mutation,

which keeps Sw-5b in an autoactivated state (Chen *et al.*, 2016), was introduced into

Sw-5b to produce pYFP-Sw-5b^{D857V} and expressed in *N. benthamiana* leaves, the

mutant YFP-Sw-5b^{D857V} fusion also accumulated in both cell cytoplasm and nucleus

(Fig. 1b, right image). We also making a construct pNativePro::YFP-Sw-5b

expressing YFP-Sw-5b under native Sw-5b promoter. However, the expression of

YFP-Sw-5b by native Sw-5b promoter is too low to detect green fluorescence signal.

To investigate the subcellular localization pattern of Sw-5b in tomato leaf cells,

we transiently expressed YFP, YFP-Sw-5b, and YFP-Sw-5b^{D857V}, respectively,

through particle bombardment. Confocal Microscopy results showed that these three

proteins exhibited the same subcellular localization pattern as that in the *N.*

benthamiana leaf cells (Fig. 1c).

To further confirm the above results, we harvested *N. benthamiana* leaves

expressing YFP-Sw-5b or YFP-Sw-5b^{D857V} and analyzed by cytoplasm and nucleus

fractionation assay. Leaf samples agro-infiltrated with the empty vector (p2300S)

were also harvested and used as controls. Analyses of total protein, cytoplasm

fractions, and nucleus fractions from these harvested leaves using Western blot assays

showed that both YFP-Sw-5b and YFP-Sw-5b^{D857V} were accumulated in the

cytoplasm and nucleus (Fig. 1d).

Sw-5b recognizes TSWV NSm in the cytoplasm

TSWV NSm is known to reside in cytoplasm and plasmodesmata, but not in nucleus

(Kormelink *et al.*, 1994; Feng *et al.*, 2016). To determine where Sw-5b can recognize

TSWV NSm, we fused a NES, a nes, a NLS or a nls signal peptide to the C-terminus

of NSm-YFP to produce NSm-YFP-NES, NSm-YFP-nes, NSm-YFP-NLS, and

NSm-YFP-nls, respectively. Transient expressions of these fusions in *N. benthamiana*

leaves showed that NSm-YFP-NES accumulated exclusively in the cytoplasm, while

NSm-YFP-NLS accumulated in the nucleus (Fig. S1a). As expected, NSm-YFP-nes

and NSm-YFP-nls showed the same accumulation pattern as that of NSm-YFP (Fig.

S1a). When Sw-5b was co-expressed with one of the above four fusions in *N.*

benthamiana leaves through agro-infiltration, the leaf tissues co-expressing Sw-5b

and NSm-YFP-NES (Sw-5b + NSm-YFP-NES), Sw-5b and NSm-YFP-nes (Sw-5b +

NSm-YFP-nes), or Sw-5b and NSm-YFP-nls (Sw-5b + NSm-YFP-nls) developed a

strong HR cell death (Fig. S1b). In contrast, the leaf tissues co-expressing Sw-5b and

NSm-YFP-NLS (Sw-5b + NSm-YFP-NLS) did not. Western blot assays using a YFP

specific antibody confirmed that all the assayed proteins were expressed in the

infiltrated tissues (Fig. S1c), indicating that Sw-5b recognizes NSm in the cytoplasm.

Sw-5b activity in cell death induction is enhanced in the cytoplasm but

suppressed in the nucleus

To investigate the roles of the cytoplasmic and nuclear Sw-5b in the induction of cell

death and host immunity, we produced constructs to express YFP-Sw-5b,

NLS-YFP-Sw-5b, nls-YFP-Sw-5b, NES-YFP-Sw-5b, and nes-YFP-Sw-5b,

respectively, and then tested their abilities to elicit cell death and host immunity to

tospovirus infection. Transient expressions of these fusions in *N. benthamiana* leaves

showed that NES-YFP-Sw-5b accumulated exclusively in the cytoplasm, while

NLS-YFP-Sw-5b accumulated only in the nucleus (Fig. 2a). In addition,

nes-YFP-Sw-5b and nls-YFP-Sw-5b showed the same accumulation pattern as that

shown by YFP-Sw-5b. We then tested cell death induction through co-expressions of

NSm and YFP-Sw-5b (NSm + YFP-Sw-5b), NSm and NES-YFP-Sw-5b (NSm +

NES-YFP-Sw-5b), NSm and nes-YFP-Sw-5b (NSm + nes-YFP-Sw-5b), NSm and

NLS-YFP-Sw-5b (NSm + NLS-YFP-Sw-5b), or NSm and nls-YFP-Sw-5b (NSm +

nls-YFP-Sw-5b) in *N. benthamiana* leaves through agro-infiltration. Results of this

study showed that the NSm + NES-YFP-Sw-5b-induced cell death was stronger than

that induced by NSm + nes-YFP-Sw-5b or NSm + nls-YFP-Sw-5b co-expression (Fig.

2b). In addition, the cell death induced by NSm + NLS-YFP-Sw-5b co-expression

was suppressed (Fig. 2b). Western blot results showed that the stronger cell death

caused by NSm + NES-YFP-Sw-5b co-expression was not due to a greater

accumulation of NES-YFP-Sw-5b in the leaves (Fig. 2c). The ion leakage assay

results (Fig. 2d) agreed with the phenotype observation results, and indicated that

co-expression of NSm + NES-YFP-Sw-5b in leaves lead to a greater ion leakage

compared with that induced by the co-expression of NSm + nes-YFP-Sw-5b at 24 and

48 hours post agro-infiltration (hpai). The ion leakage caused by the co-expression of

NSm + NLS-YFP-Sw-5b was significantly weaker than that caused by the

co-expression of NSm + nls-YFP-Sw-5b (Fig. 2d).

Cytoplasmic Sw-5b induces a strong host defense against tospovirus replication

Virus infection in plant starts with virus replication in the initially infected cells

followed by spreading into adjacent cells for further infection. To monitor tospovirus

replication in plant cells, we recently developed a TSWV mini-replicon-based reverse

genetic system (Feng et al., 2020). In this study, co-expression of TSWV

mini-replicon $SR_{(+)eGFP}$, $L_{(+)opt}$ (with a codon usage optimized RdRp), VSRs and NSm

resulted in a cell-to-cell movement of $SR_{(+)eGFP}$. In contrast, co-expression of $SR_{(+)eGFP}$,

$L_{(+)opt}$, VSRs, and $NSm^{H93A\&H94A}$ mutant, a defective movement protein but can be

recognized by Sw-5b to cause a strong HR (Li *et al.*, 2009; Zhao *et al.*, 2016), in cells

resulted in the expression of $SR_{(+)eGFP}$ in only single cells (Fig. S2), thus dissecting the

viral replication from viral cell-to-cell movement. We then co-expressed $SR_{(+)eGFP}$,

$L_{(+)opt}$, VSRs, $NSm^{H93A\&H94A}$ mutant and one of the five proteins (i.e., Sw-5b,

NES-Sw-5b, nes-Sw-5b, NLS-Sw-5b, nls-Sw-5b) in *N. benthamiana* leaves. Leaves

co-expressing $SR_{(+)eGFP}$, $L_{(+)opt}$, VSRs, $NSm^{H93A\&H94A}$ mutant and p2300 (empty vector,

EV) were used as controls. The results showed that in the presence of Sw-5b or one of

its derivatives, the expression of $SR_{(+)eGFP}$ was strongly suppressed compared with

that expressed in the presence of EV (Fig. 3a). It is noteworthy that the expression of

$SR_{(+)eGFP}$ was less inhibited in the presence of NLS-Sw-5b (Fig. 3a). Western blot

result indicated that the GFP accumulation of $SR_{(+)eGFP}$ was strongly inhibited in the

presence of Sw-5b, NES-Sw-5b, nes-Sw-5b or nls-Sw-5b compared with that

expressed in the presence of NLS-Sw-5b or EV (Fig. 3b). This finding indicates that

the cytoplasmic Sw-5b can inhibit $SR_{(+)eGFP}$ expression, possibly through induction of

a host defense against TSWV replication.

Sw-5b induces a host defense against viral NSm intercellular movement

In our previous study, we used pmCherry-HDEL//NSm-GFP vector (Fig. 4a) to

investigate TSWV NSm cell-to-cell movement (Feng *et al.*, 2016). The expressed

mCherry-HDEL binds ER membrane in the initial cells but NSm-eGFP traffics

between cells. To investigate whether the Sw-5b-induced host defense can affect

TSWV NSm cell-to-cell movement, we co-expressed mCherry-HDEL, NSm-GFP,

and Sw-5b or mCherry-HDEL, NSm-GFP, and EV in *N. benthamiana* leaves through

agro-infiltration. Under the fluorescence microscope, both NSm-GFP and

mCherry-HDEL were found in single cells in the presence of Sw-5b. In the presence

of EV, however, NSm-GFP moved into multiple cells, while mCherry-HDEL

accumulated in the initial cells (Fig. 4b, upper two panels). The result suggested that

Sw-5b elicited a defense that strongly inhibited cell-to-cell movement of viral NSm.

To make sure this inhibition to viral NSm cell-to-cell movement is not caused by

overexpression of Sw-5b, we also used the NSm^{T120N} mutant, from the resistance‐

breaking (RB) TSWV isolates, which cannot be recognized by Sw-5b (Zhao et al.,

2016). The assays showed that in the presence of either Sw-5b or EV, NSm^{T120N}-GFP

moved into multiple cells, while mCherry-HDEL retained in the initial cells (Fig S3a).

Sw-5b in the nucleus but not in the cytoplasm triggers a defense against NSm

cell-to-cell movement

To determine the effects of the cytoplasmic and nuclear Sw-5b on host defense against

TSWV NSm intercellular movement, we co-expressed mCherry-HDEL and

NSm-GFP with NES-Sw-5b, nes-Sw-5b, NLS-Sw-5b, or nls-Sw-5b in *N.*

benthamiana leaves via agro-infiltration. The results showed that in the presence of

NLS-Sw-5b, the cell-to-cell movement of NSm-GFP was inhibited (Fig. 4b). Similar

results were also obtained in the leaves co-expressing mCherry-HDEL and NSm-GFP

with nls-YFP-Sw-5b or nes-YFP-Sw-5b (Fig S3b). In the presence of

NES-YFP-Sw-5b, however, NSm-GFP did move into surrounding cells. (Fig. 4b).

This finding indicates that the Sw-5b in the nucleus but not in the cytoplasm induced

a host defense that inhibited TSWV NSm cell-to-cell movement.

Nuclear Sw-5b confers host immunity to TSWV systemic infection

To dissect the host immunity induced by the cytoplasmic and the nuclear Sw-5b, we

generated transgenic *N. benthamiana* lines expressing YFP-Sw-5b, NES-YFP-Sw-5b,

nes-YFP-Sw-5b, NLS-YFP-Sw-5b, and nls-YFP-Sw-5b, respectively (Tables S1 and

S2). After inoculation of these transgenic lines with TSWV-YN isolate, the EV

(control) transgenic plants developed typical viral symptoms including stunt, leaf curl

and mosaic at 7 to 15 days post inoculation (dpi). The NES-YFP-Sw-5b transgenic

plants developed a strong HR trailing in the systemic leaves by 7 to 15 days post

inoculation (dpi) (Fig. 5a and Fig. S4a), suggesting that NES-YFP-Sw-5b transgenic

plant did not block TSWV systemic infection and caused virus infection-related

systemic HR. In contrast, no systemic virus infection symptoms were observed in the

YFP-Sw-5b and the nes-YFP-Sw-5b transgenic plants. The RT-PCR agreed with the

symptom observation results and showed that TSWV-YN genomic RNA was

accumulated in the systemic leaves of the TSWV-YN-inoculated NES-YFP-Sw-5b or

the EV transgenic plants, but not in the systemic leaves of the TSWV-YN-inoculated

YFP-Sw-5b or nes-YFP-Sw-5b transgenic plants (Fig. 5c and Fig. S4b). Also in this

study, the TSWV-YN-inoculated NLS-YFP-Sw-5b or nls-YFP-Sw-5b transgenic

plants did not show virus like symptoms in their systemic leaves by 7-15 dpi (Fig. 5a,

and Fig. S4a). The RT-PCR result confirmed that TSWV-N genomic RNA had not

accumulated in the systemic leaves of the NLS-YFP-Sw-5b or the nls-YFP-Sw-5b

transgenic plants (Fig. 5c, and Fig. S4b), indicating that the nuclear Sw-5b is

responsible for the host immunity against TSWV systemic infection.

The cytoplasmic and the nuclear Sw-5b act synergistically to confer a strong

immunity to TSWV infection in *N. benthamiana*

To investigate whether cytoplasm-targeted and nucleus-targeted Sw-5b have joint

effects on the defense against TSWV infection, we constructed a $M_{(-)opt}$-$pSR_{(+)eGFP}$

vector by inserting a cassette expressing optimized TSWV M genomic sequence

(Feng *et al.*, 2020) into the $pSR_{(+)eGFP}$ mini-replicon to express NSm, N, and eGFP

simultaneously in the same cells (Fig. 6a). The construct $M_{(-)opt}$-$pSR_{(+)eGFP}$ couples the

functions for both viral replication and viral cell-to-cell movement, mimicking the

virus infection in plant leaves. After co-expressing this vector, the $L_{(+)opt}$ and the EV in

N. benthamiana leaves through agro-infiltration, the eGFP fluorescence was observed

in many cells, due to the presence of the NSm movement protein and the $RdRp_{opt}$ (Fig.

6b, upper left image). When $M_{(-)opt}$-$SR_{(+)eGFP}$, $L_{(+)opt}$ and Sw-5b were co-expressed in

N. benthamiana leaves, the eGFP fluorescence was hardly detected and some were

observed only in single leaf cells (Fig. 6b upper right image, Fig. S5a and b). When

$M_{(-)opt}$-$SR_{(+)eGFP}$, $L_{(+)opt}$ and NES-Sw-5b were co-expressed in *N. benthamiana* leaves,

the eGFP fluorescence was observed in clusters of a few cells (Fig. 6b, Fig. S5a),

indicating that limited cell-to-cell movement had occurred in these leaves (Fig. S5b).

When $M_{(-)opt}$-$SR_{(+)eGFP}$, $L_{(+)opt}$ and NLS-Sw-5b were co-expressed in leaves, a few of

eGFP fluorescence were detected but they were in single cells only. When leaves

co-expressing $M_{(-)opt}$-$SR_{(+)eGFP}$, $L_{(+)opt}$ and NES-Sw-5b + NLS-Sw-5b, the eGFP

fluorescence was also hardly detected and some were observed only in single leaf

cells. Western blot results showed that more eGFP had accumulated in the leaves

co-expressing $M_{(-)opt}$-$SR_{(+)eGFP}$, $L_{(+)opt}$, and EV, followed by the leaves co-expressing

$M_{(-)opt}$-$SR_{(+)eGFP}$, $L_{(+)opt}$, and NLS-Sw-5b, and then the leaves co-expressing

$M_{(-)opt}$-$SR_{(+)eGFP}$, $L_{(+)opt}$, and NES-Sw-5b. Much less eGFP had accumulated in the

leaves co-expressing $M_{(-)opt}$-$SR_{(+)eGFP}$, $L_{(+)opt}$, and NLS-Sw-5b + NES-Sw-5b, and in

the leaves co-expressing $M_{(-)opt}$-$SR_{(+)eGFP}$, $L_{(+)opt}$, and Sw-5b (Fig. 6c and d). The

accumulation of eGFP was lower in the leaves co-expressing NES-Sw-5b +

NLS-Sw-5b than that in the leaves co-expressing NES-Sw-5b or NLS-Sw-5b (Fig. 6c

and d), indicating that NES-Sw-5b and NLS-Sw-5b have additive role in mediating

host immunity against different TSWV infection steps.

The Sw-5b NB-ARC-LRR control its cytoplasm localization whereas the

extended N-terminal SD domain is crucial for targeting Sw-5b into nucleus, and

for inducing host systemic immunity

Sw-5b has an extended N-terminal SD domain, a CC domain, a NB-ARC domain, and

a C-terminal LRR domain (Chen *et al.*, 2016). To determine which domain(s) of

Sw-5b is/are responsible for nucleoplasm/nucleolus targeting and for plant immunity,

we tested these domains using various deletion mutants and YFP fusion proteins (Fig.

7a). We reported previously that the Sw-5b NB-ARC-LRR region was able to induce

HR cell death in plant in the presence of NSm (Chen *et al.*, 2016). In this study, we

fused YFP to the N-terminus of NB-ARC-LRR (Fig. 7a). Transient expression of

YFP-NB-ARC-LRR (112 kDa) in *N. benthamiana* leaf cells resulted in a localization

of the fusion protein in cytoplasm exclusively (Fig. 7b).

A previous study had shown that the CC domain of potato NLR receptor Rx was

required for targeting this protein to nucleus (Slootweg *et al.*, 2010). To determine the

function of Sw-5b CC domain in intracellular trafficking, we inserted the CC domain

between the YFP and NB-ARC-LRR to generate an YFP-CC-NB-ARC-LRR

construct or fused the CC domain to YFP to produce an YFP-CC construct. Transient

expression of these two fusion proteins individually in *N. benthamiana* leaves, and

examined the leaves under a confocal microscope, we determined that the YFP-CC

fusion protein accumulated in both cytoplasm and nucleus of the cells while the

YFP-CC-NB-ARC-LRR fusion protein was in the cytoplasm only (Fig. 7b). This

result indicated that addition of the CC domain to YFP-NB-ARC-LRR was not

sufficient to traffic the fusion protein into the nucleus.

An extended N-terminal SD domain is known to be present at the upstream of the

Sw-5b CC domain. In this study, we first generated an YFP-SD and an YFP-SD-CC

constructs, and transiently expressed them individually in *N. benthamiana* leaf cells.

Confocal Microscopy showed that both YFP-SD and YFP-SD-CC fusion proteins

accumulated in the cytoplasm and nucleus (Fig. 7b). We then inserted a SD between the YFP and CC-NB-ARC-LRR to produce an YFP-SD-CC-NB-ARC-LRR construction. Transient expression of this construct in *N. benthamiana* leaf cells showed that this fusion protein accumulated in the cytoplasm and nucleus (Fig. 7b).

We next generated stable transgenic *N. benthamiana* plants expressing YFP-NB-ARC-LRR, YFP-CC-NB-ARC-LRR and YFP-SD-CC-NB-ARC-LRR. Upon inoculation transgenic *N. benthamiana* plants expressing YFP-NB-ARC-LRR with TSWV, large HR foci were observed in the TSWV-inoculated leaves and later, HR trailing was seen in the systemic leaves of most assayed plants (Fig. 7c; Table S2 and S3). RT-PCR results confirmed the presence of TSWV genomic RNA in these systemic leaves (Fig. S6b). We also inoculate *N. benthamiana* plants expressing YFP-CC-NB-ARC-LRR with TSWV. By 7 dpi, no systemic resistance to TSWV infection was observed in these plants (Fig. 6a and B, Table S4). RT-PCR results showed that systemic infection of TSWV did occur in the TSWV-inoculated YFP-CC-NB-ARC-LRR transgenic plants (Fig. S6b). In contrast, transgenic plants expressing YFP-SD-CC-NB-ARC-LRR fusion exhibited a systemic immunity to TSWV infection (Fig. 7c, Fig. S6b).

These data indicated that Sw-5b NB-ARC-LRR control its cytoplasm localization, CC domain of Sw-5b alone was not sufficient to transport the NB-ARC-LRR into nucleus and the extended SD domain is required for targeting Sw-5b to the nucleus, and for inducing host immunity.

The extended SD domain interacted with *importin α1, α2 and β*

To identify the cellular machinery needed for transporting Sw-5b into nucleus, we co-expressed YFP-SD and YFP-Sw-5b in *N. benthamiana* leaves followed by a co-immunoprecipitation (co-IP) and Mass Spectrometry. The results identified *N. benthamiana* importin α as one of candidate proteins interacted with YFP-SD and YFP-Sw-5b (Table S5 and S6). The co-IP and Mass Spectrometry also identified nuclear pore complex protein TPRb and nuclear pore complex protein Nup160a interacted with YFP-Sw-5b (Table S6). *Importins* play important roles in translocating proteins from cytoplasm into nucleus (Kanneganti *et al.*, 2007). We used BiFC analysis to confirm the interaction between YFP-SD with *N. benthamiana* importin homologs α1, α2 and β. The result showed that co-expression of cYFP-SD with nYFP-IMP α1, nYFP-IMP α2 or nYFP-IMP β produced a strong YFP fluorescence signal in nucleus. Co-expression of cYFP-Sw-5b with nYFP-IMP α1, nYFP-IMP α2 or nYFP-IMP β also detected a strong YFP fluorescence signal in nucleus (Fig. S7). In contrast, co-expression of controls cYFP-SD and nYFP, cYFP-Sw-5b and nYFP, cYFP and nYFP-IMP α1, cYFP and nYFP-IMP α2 or cYFP and nYFP-IMP β did not show fluorescence signal in *N. benthamiana* leaf cells (Fig. S7).

Silencing *importin α1, α2* and *β* expression abolished Sw-5b nucleus accumulation and host resistance to TSWV systemic infection

To determine the functions of importin α1, α2 and β in Sw-5b nucleus localization, we

silenced *importin α1, α2, β, α1* and *α2,* and *α1* and *α2* and *β* expressions, respectively,

in *N. benthamiana* leaves using a tobacco rattle virus (TRV)-based virus-induced gene

silencing (VIGS) vector, and then transiently expressed YFP-Sw-5b in these plants.

Analyses of these plants through RT-PCR using gene specific primers showed that

silencing of these *importin* genes in *N. benthamiana* leaves were successful (Fig.

S8a). However, silencing individual *importin* gene or both *importin α1* and *α2* was

not enough to block the nucleus accumulation of YFP-Sw-5b (Fig. 8a). In contrast,

after *importin α1, α2* and *β* were all silenced through VIGS, the nucleus accumulation

of YFP-Sw-5b was inhibited (Fig. 8a, the middle image in the bottom panel).

To investigate the effects of nuclear import defected Sw-5b on host immunity to

TSWV systemic infection, we silenced these *importin* genes in the Sw-5b transgenic

N. benthamiana plants as described above, and then inoculated them with TSWV. The

results showed that the plants silenced for *importin α1, α2,* and *β* gene, individually,

did not show TSWV systemic infection (Fig. 8b and Fig. S8b). In addition, the

transgenic plants silenced for both *importin α1* and *α2* genes also did not show

TSWV systemic infection (Fig. 8b and Fig. S8b). In contrast, after silencing *importin*

α1, α2 and *β* together, the plants developed clear TSWV symptoms in systemic

leaves followed by HR (Fig. 8b, white arrow and Fig. S8b), indicating that the nucleus

accumulation of Sw-5b is indispensable for the induction of host immunity against

TSWV systemic infection.

Discussion

In this report, we provide evidence to show that the cytoplasm-accumulated and the

nucleus-accumulated Sw-5b, a tomato immune receptor, play different roles in

inducing host defense against TSWV infection in plant. The cytoplasmic Sw-5b

functions to induce a strong cell death response to inhibit TSWV replication. This

host response is, however, insufficient to block virus intercellular and long-distance

movement. The nuclear-localized Sw-5b triggers a host defense that weakly inhibit

viral replication but strongly inhibit tospovirus intercellular and systemic movement.

These findings suggest that tomato Sw-5b NLR induces different types of defense

responses by cytoplasm and nucleus partitioning to combat virus at different infection

steps. Furthermore, the cytoplasmic and the nuclear Sw-5b act synergistically to

confer a strong host immunity to TSWV infection in plant. We also demonstrated that

the extra SD domain functioned as a critical intracellular translocation modulator,

allowing Sw-5b receptor to translocate from cytoplasm to nucleus to trigger the

immunity. The Sw-5b NB-LRR controls its cytoplasm localization. Unlike Rx CC

domain, Sw-5b CC domain is not sufficient to translocate NB-LRR into nucleus.

Strikingly, the SD is crucial for Sw-5b to translocate from cytoplasm for nucleus. This

SD-mediated receptor translocation is dependent on importins α and β.

Successful virus infection in plant requires several steps including viral replication

in the initially infected cells followed by cell-to-cell and long-distance movement

(Heinlein, 2015; Wang, 2015). After entering into plant cells, virus first encode

multiple proteins needed for its replication. Once the initial replication is established,

virus will encode specific protein(s), known as movement proteins (MPs), to traffic

viral genome or virions into adjacent cells through plasmodesmata in cell walls, and

then long-distantly into other parts of the plant to cause a systemic infection (Rao,

2002; Lucas, 2006; Taliansky *et al.*, 2008). To date, multiple plant NLRs, conferring

host resistance against plant viruses have been identified (Soosaar *et al.*, 2005; Meier

et al., 2019), but how these plant NLRs induce host resistance against virus infection

remain largely unknown.

In this study, we have determined that the forced cytoplasm accumulation of

Sw-5b can induce a stronger cell death than that caused by the accumulation of Sw-5b

in both cytoplasm and nucleus. While, the cell death induced by the forced nucleus

accumulation of Sw-5b was significantly weakened. We then analyzed

Sw-5b-mediated immunity against TSWV replication using a TSWV mini-replicon

system and a movement defective NSm mutant. Our results showed that the forced

cytoplasm accumulation of Sw-5b can induce a strong host defense against virus

replication in cells. This finding implies that cytoplasm is one of the main source of

defense signaling against TSWV replication. The defense signaling generated in

nucleus can only induce a weak defense against TSWV replication. Therefore, the

nuclear localized Sw-5b is only partially responsible for the induction of host defense

against TSWV replication. It is also possible that this nuclear localized

Sw-5b-induced weak host response is caused by a trace of NLS-YFP-Sw-5b

maintained in the cytoplasm that maybe below the detection limit of Confocal

Microscope. It has been shown to accumulate in both cytoplasm and nucleus, and the

forced cytoplasm accumulation of Barley MLA10 enhance cell death signaling (Bai *et*

al., 2012). We speculate that, for both MLA10 and Sw-5b, the cytoplasm

accumulation is crucial for the initiation and/or amplification of the cell death

signaling. The CC and the TIR domain of several plant NLRs have been shown to

trigger cell death (Swiderski *et al.*, 2009; Krasileva *et al.*, 2010; Bernoux *et al.*, 2011;

Collier *et al.*, 2011; Maekawa *et al.*, 2011; Bai *et al.*, 2012; Chen *et al.*, 2017; Wang

et al., 2020). Analyses of the three dimensional structures of Arabidopsis ZAR1

resistosome have also shown that its CC domain can form pentamer structures that

was able to target into host cell membranes, leading to ion leakage and cell death

(Wang *et al.*, 2019a; Wang *et al.*, 2019b). We speculate that cell death likely cause the

toxicity on viral replicase or other proteins associated with virus replication in cells.

Plant virus encodes specific movement protein(s) to traffic viral genome between

cells and then leaves to cause systemic infection (Rao, 2002; Lucas, 2006; Taliansky

et al., 2008). We reported previously that TSWV NSm alone can move between plant

cells (Feng *et al.*, 2016). In this study, we investigated the effect of the

Sw-5b-mediated host defense on TSWV intercellular movement. Through this study,

we have determined that after the recognition of NSm, Sw-5b receptor induced a

strong reaction to block NSm intercellular trafficking. Previous reports have some

indications on the role of plant NLRs in viral movement. Nevertheless it has no direct

evidence showing that plant NLRs induce resistance against viral movement. Deom

and colleagues had shown that the 9.4-kDa fluorescein isothiocyanate-labeled dextran

was unable to move between cells in the transgenic tobacco *N* leaves expressing

tobacco mosaic virus (TMV) movement protein at 24°C, an HR-permissive

temperature (Deom *et al.*, 1991). However, that study did not involve a TMV Avr

protein. In a different report, TMV-GFP showed a limited cell-to-cell movement in

leaves of tobacco cv. Sumsan NN at 33°C, an HR-nonpermissive temperature (Canto

& Palukaitis, 2002). Li and colleagues found that after treatment of SMV-inoculated

Jidou 7 resistant plants with a callose synthase inhibitor, the plants showed enlarged

HR lesions (Li *et al.*, 2012). The soybean *Rsv3* induced extreme resistance. However,

after this extremely resistant soybean line was treated with a callose synthase inhibitor,

the plants developed HR lesions upon SMV-G5H inoculation (Seo *et al.*, 2014). These

reports indicate that plant NLRs likely involves the defense against viral movement.

Here we provide the direct evidence that Sw-5b NLR can induce a strong defense

response to impede NSm intercellular trafficking. More importantly, we have

determined that the induction of host immunity to TSWV intercellular movement

requires the accumulation of Sw-5b in nucleus. Although the cytoplasmic Sw-5b can

induce a strong cell death response, it cannot prevent TSWV NSm cell-to-cell

movement. Consequently, we propose that nucleus is a key compartment to generate

defense signaling to block TSWV cell-to-cell movement.

In this study, although the NES-YFP-Sw-5b transgenic *N. benthamiana* plants

showed an HR, they were unable to stop TSWV systemic infection. We also showed

that Sw-5b YFP-NB-ARC-LRR (112 kDa) accumulates in cytoplasm exclusively (Fig.

7b), however, transgenic *N. benthamiana* plants expressing YFP-NB-ARC-LRR show

strong systemic HR trailing caused by TSWV infection. Based on these findings, we

conclude that HR cell death alone is not sufficient to block TSWV long-distance

movement. In our study, the NLS-YFP-Sw-5b transgenic plants were resistant to

TSWV systemic infection. After silencing the expressions of *importin α1*, *α2* and *β*

simultaneously to inhibit the nucleus accumulation of Sw-5b, however, the resistance

to TSWV systemic infection was abolished. These findings indicate that the

Sw-5b-mediated resistance signaling against viral systemic infection is generated in

nucleus. Some plant NLRs are known to interact with specific transcription factors in

nucleus upon recognition of pathogen effectors (Cui *et al.*, 2015; Kapos *et al.*, 2019).

The immune regulator EDS1 has also been shown to accumulate in nucleus to

reprogram RNA transcription (Garcia *et al.*, 2010; Heidrich *et al.*, 2011; Cui *et al.*,

2015; Lapin *et al.*, 2020). How Sw-5b regulates host immunity in nucleus requires

further investigations.

Several plant immune receptors and immune regulators, including, e.g. potato Rx

(Slootweg *et al.*, 2010; Tameling *et al.*, 2010), tobacco N (Burch-Smith *et al.*, 2007;

Caplan, JL *et al.*, 2008), barley MLA10 (Shen *et al.*, 2007), Arabidopsis

RRS1-R/RPS4, and snc1 (Deslandes *et al.*, 2003; Wirthmueller *et al.*, 2007; Cheng *et*

al., 2009), as well as Arabidopsis NPR1 (Katagiri & Tsuda, 2010), and EDS1 (Lapin

et al., 2020) have been found to be nucleocytoplasmic. For some of them, nuclear

accumulation of NLRs are required for the induction of plant immunity to pathogen

attacks. Moreover, the MLA10-YFP-NES fusion was found to induce a strong cell

death response, but not a strong host resistance to powdery mildew fungus infection.

In contrast, the MLA10-YFP-NLS fusion inhibited its activity to induce a cell death

response, but caused a host immunity to this pathogen (Bai *et al.*, 2012). In many

plant-pathogen interactions, cell death responses can be uncoupled from disease

resistance (Bendahmane *et al.*, 1999; Gassmann, 2005; Coll *et al.*, 2010; Heidrich *et*

al., 2011). This separation raises questions about how host resistance prevents

pathogen invasion and what are the roles of cell death during pathogen infection. It is

unclear whether the MLA10-YFP-NES-induced cell death has some inhibitory effects

on powdery mildew fungus infection. In this study, we determined that cytoplasm-

and nuclear-accumulation of Sw-5b have different functions. The

cytoplasm-accumulated Sw-5b induces a strong defense against virus replication,

whereas the nuclear-accumulated Sw-5b induced an inhibition of virus cell-to-cell and

long distance movement. Both cytoplasmic and nuclear Sw-5b are needed to confer a

synergistic and full defense against tospovirus infection.

We have also determined that Sw-5b NB-ARC-LRR and SD domains are

important to regulate the proper subcellular localization of Sw-5b and the proper

nucleoplasmic distribution of Sw-5b is needed to elicit full immune responses to

inhibit different TSWV infection steps. Sw-5b NB-ARC-LRR controls its cytoplasm

localization. The CC domain of the Sw-5b is not sufficient to target the receptor into

nuclear. Importantly, the extended SD of Sw-5b is absolutely required for the nucleus

translocation. Because non-canonical domains are frequently found in other NLRs

and are quite diversified, our findings have broad implications to investigate the

potential new functions of non-canonical domains that integrated in the plant NLRs to

regulate the plant immunity against pathogen invasions.

Through co-IP, Mass Spectrometry and BiFC analysis, we found that the extended

SD domain and Sw-5b interacted with host importin machineries to translocate the

Sw-5b receptor from cytoplasm into nucleus to mediate local and systemic resistance

to tospovirus. Recent studies have shown that nuclearporin MOS3, MOS6 and nuclear

pore complex component MOS7/Nup88 proteins played important roles in regulating

Arabidopsis innate immunity (Palma *et al.*, 2005; Zhang & Li, 2005; Cheng *et al.*,

2009). We found that when *importin α1, importin α2* and *importin β* gene were all

silenced through VIGS, the nucleus targeting of Sw-5b were completely blocked, and

consequently, the Sw-5b-mediated systemic immunity to tospovirus infection was

compromised. Importin α and β are known to form a nucleus import complex.

Binding with importin β could activate importin α to form a binding surface for NLS

proteins (Stewart, 2007). Hence, disruption of either importin α or importin β would

block the nucleus import of NLS proteins. Because silencing *importin α* or

importin β gene expression through VIGS did not disrupt the nuclear targeting of

Sw-5b*,* we speculate that a non-canonical nuclear import pathway may take part in

importing Sw-5b into nucleus.

Based on the above results, we have created a working model for the Sw-5b

NLR-induced host resistance against TSWV replication, and intercellular and

long-distance movement in plant (Fig. 9). Upon recognition of NSm in cytoplasm,

Sw-5b switched from an autoinhibited state to an activated state. The activated Sw-5b

accumulated in cytoplasm and also translocate into nucleus via importins α and β .

The cytoplasm-accumulated and the nucleus-accumulated Sw-5b play different roles

in inducing host immunity against TSWV infection. The cytoplasmic Sw-5b functions

to induce a cell death response to inhibit TSWV replication, while the nuclear Sw-5b

functions to induce a weak host defense against TSWV replication, but a strong

defense against TSWV cell-to-cell and long-distance movement. The concerted

defense signaling generated in the cytoplasm and nucleus resulted in a strong host

resistance to tospovirus infection.

Acknowledgments

This work was supported by the National Natural Science Foundation of China

(31630062, 31925032 and 31870143), the Fundamental Research Funds for the

Central Universities (JCQY201904 and KYXK202012), Youth Science and

Technology Innovation Program to XT.

Author contributions

HC XQ, XC and XT, designed the research; HC, XQ and XC, TY, MF, JC, RC, HH,

YZ, YM, DS, YX, MZ performed the experiments; HC, XSD and XT interpreted the

result and wrote the paper.

ORCID

Hongyu Chen, 0000-0001-8142-0653

Yi Xu, 0000-0002-1913-4530

Min Zhu, 0000-0002-9354-4300

Xiaorong Tao, 0000-0003-1259-366X

Competing interests

798   The authors declare that nocompeting interests exist.

Data availability

801   All data produced in this study are presented in this manuscript or as the supporting

802   files

References

Axtell MJ, Staskawicz BJ. 2003. Initiation of RPS2-specified disease resistance in Arabidopsis is
coupled to the AvrRpt2-directed elimination of RIN4. *Cell* **112**(3): 369-377.
Bai S, Liu J, Chang C, Zhang L, Maekawa T, Wang Q, Xiao W, Liu Y, Chai J, Takken FL,
Schulze-Lefert P, Shen QH. 2012. Structure-function analysis of barley NLR immune
receptor MLA10 reveals its cell compartment specific activity in cell death and disease
resistance. *PLoS Pathog* **8**(6): e1002752.
Bendahmane A, Kanyuka K, Baulcombe DC. 1999. The Rx gene from potato controls separate virus
resistance and cell death responses. *Plant Cell* **11**(5): 781-792.
Bernoux M, Ve T, Williams S, Warren C, Hatters D, Valkov E, Zhang X, Ellis JG, Kobe B,
Dodds PN. 2011. Structural and functional analysis of a plant resistance protein TIR domain
reveals interfaces for self-association, signaling, and autoregulation. *Cell Host Microbe* **9**(3):
200-211.
Brommonschenkel SH, Frary A, Frary A, Tanksley SD. 2000. The broad-spectrum tospovirus
resistance gene Sw-5 of tomato is a homolog of the root-knot nematode resistance gene Mi.
Mol Plant Microbe Interact **13**(10): 1130-1138.
Burch-Smith TM, Schiff M, Caplan JL, Tsao J, Czymmek K, Dinesh-Kumar SP. 2007. A novel
role for the TIR domain in association with pathogen-derived elicitors. *PLoS Biol* **5**(3): e68.
Canto T, Palukaitis P. 2002. Novel N gene-associated, temperature-independent resistance to the
movement of tobacco mosaic virus vectors neutralized by a cucumber mosaic virus RNA1
transgene. *J Virol* **76**(24): 12908-12916.
Caplan J, Padmanabhan M, Dinesh-Kumar SP. 2008. Plant NB-LRR Immune Receptors: From
Recognition to Transcriptional Reprogramming. *Cell Host Microbe* **3**(3): 126-135.

Caplan JL, Mamillapalli P, Burch-Smith TM, Czymmek K, Dinesh-Kumar SP. 2008.
Chloroplastic protein NRIP1 mediates innate immune receptor recognition of a viral effector.
Cell **132**(3): 449-462.
Cesari S, Bernoux M, Moncuquet P, Kroj T, Dodds PN. 2014. A novel conserved mechanism for
plant NLR protein pairs: the "integrated decoy" hypothesis. *Front Plant Sci* **5**: 606.
Cesari S, Moore J, Chen C, Webb D, Periyannan S, Mago R, Bernoux M, Lagudah ES, Dodds
PN. 2016. Cytosolic activation of cell death and stem rust resistance by cereal MLA-family
CC-NLR proteins. *Proc Natl Acad Sci U S A* **113**(36): 10204-10209.
Cesari S, Thilliez G, Ribot C, Chalvon V, Michel C, Jauneau A, Rivas S, Alaux L, Kanzaki H,
Okuyama Y, Morel JB, Fournier E, Tharreau D, Terauchi R, Kroj T. 2013. The rice
resistance protein pair RGA4/RGA5 recognizes the Magnaporthe oryzae effectors AVR-Pia
and AVR1-CO39 by direct binding. *Plant Cell* **25**(4): 1463-1481.
Chen T, Liu D, Niu X, Wang J, Qian L, Han L, Liu N, Zhao J, Hong Y, Liu Y. 2017. Antiviral
Resistance Protein Tm-2(2) Functions on the Plasma Membrane. *Plant Physiol* **173**(4):
2399-2410.
Chen X, Zhu M, Jiang L, Zhao W, Li J, Wu J, Li C, Bai B, Lu G, Chen H, Moffett P, Tao X. 2016.
A multilayered regulatory mechanism for the autoinhibition and activation of a plant
CC-NB-LRR resistance protein with an extra N-terminal domain. *New Phytol* **212**(1):
161-175.
Cheng YT, Germain H, Wiermer M, Bi D, Xu F, Garcia AV, Wirthmueller L, Despres C, Parker
JE, Zhang Y, Li X. 2009. Nuclear pore complex component MOS7/Nup88 is required for
innate immunity and nuclear accumulation of defense regulators in Arabidopsis. *Plant Cell*
21(8): 2503-2516.
Coll NS, Vercammen D, Smidler A, Clover C, Van Breusegem F, Dangl JL, Epple P. 2010.
Arabidopsis type I metacaspases control cell death. *Science* **330**(6009): 1393-1397.
Collier SM, Hamel LP, Moffett P. 2011. Cell death mediated by the N-terminal domains of a unique
and highly conserved class of NB-LRR protein. *Mol Plant Microbe Interact* **24**(8): 918-931.
Collier SM, Moffett P. 2009. NB-LRRs work a "bait and switch" on pathogens. *Trends Plant Sci*
14(10): 521-529.
Cui H, Tsuda K, Parker JE. 2015. Effector-triggered immunity: from pathogen perception to robust
defense. *Annu Rev Plant Biol* **66**: 487-511.
Das B, Sengupta S, Prasad M, Ghose TK. 2014. Genetic diversity of the conserved motifs of six
bacterial leaf blight resistance genes in a set of rice landraces. *BMC Genet* **15**: 82.
De Oliveira AS, Koolhaas I, Boiteux LS, Caldararu OF, Petrescu AJ, Oliveira Resende R,
Kormelink R. 2016. Cell death triggering and effector recognition by Sw-5 SD-CNL proteins
from resistant and susceptible tomato isolines to Tomato spotted wilt virus. *Mol Plant Pathol*
17(9): 1442-1454.
Deom CM, Wolf S, Holt CA, Lucas WJ, Beachy RN. 1991. Altered function of the tobacco mosaic
virus movement protein in a hypersensitive host. *Virology* **180**(1): 251-256.
Deslandes L, Olivier J, Peeters N, Feng DX, Khounlotham M, Boucher C, Somssich I, Genin S,
Marco Y. 2003. Physical interaction between RRS1-R, a protein conferring resistance to
bacterial wilt, and PopP2, a type III effector targeted to the plant nucleus. *Proc Natl Acad Sci*
U S A **100**(13): 8024-8029.
Dodds PN, Rathjen JP. 2010. Plant immunity: towards an integrated view of plant-pathogen

871    interactions. *Nat Rev Genet* **11**(8): 539-548.

872 **Ea VDB, Jones JD. 1998.** Plant disease-resistance proteins and the gene-for-gene concept. *Trends in*
873    *Biochemical Sciences* **23**(12): 454-456(453).

874 **Engelhardt S, Boevink PC, Armstrong MR, Ramos MB, Hein I, Birch PR. 2012.** Relocalization of
875    late blight resistance protein R3a to endosomal compartments is associated with effector
876    recognition and required for the immune response. *Plant Cell* **24**(12): 5142-5158.

877 **Ernst K, Kumar A, Kriseleit D, Kloos DU, Phillips MS, Ganal MW. 2002.** The broad-spectrum
878    potato cyst nematode resistance gene (Hero) from tomato is the only member of a large gene
879    family of NBS-LRR genes with an unusual amino acid repeat in the LRR region. *Plant J* **31**(2):
880    127-136.

881 **Feng M, Cheng R, Chen M, Guo R, Li L, Feng Z, Wu J, Xie L, Hong J, Zhang Z, Kormelink R,**
882    **Tao X. 2020.** Rescue of tomato spotted wilt virus entirely from complementary DNA clones.
883    *Proc Natl Acad Sci U S A* **117**(2): 1181-1190.

884 **Feng Z, Xue F, Xu M, Chen X, Zhao W, Garcia-Murria MJ, Mingarro I, Liu Y, Huang Y, Jiang**
885    **L, Zhu M, Tao X. 2016.** The ER-Membrane Transport System Is Critical for Intercellular
886    Trafficking of the NSm Movement Protein and Tomato Spotted Wilt Tospovirus. *PLoS*
887    *Pathog* **12**(2): e1005443.

888 **Gao Z, Chung EH, Eitas TK, Dangl JL. 2011.** Plant intracellular innate immune receptor Resistance
889    to Pseudomonas syringae pv. maculicola 1 (RPM1) is activated at, and functions on, the
890    plasma membrane. *Proc Natl Acad Sci U S A* **108**(18): 7619-7624.

891 **Garcia AV, Blanvillain-Baufume S, Huibers RP, Wiermer M, Li G, Gobbato E, Rietz S, Parker**
892    **JE. 2010.** Balanced nuclear and cytoplasmic activities of EDS1 are required for a complete
893    plant innate immune response. *PLoS Pathog* **6**: e1000970.

894 **Gassmann W. 2005.** Natural variation in the Arabidopsis response to the avirulence gene hopPsyA
895    uncouples the hypersensitive response from disease resistance. *Mol Plant Microbe Interact*
896    **18**(10): 1054-1060.

897 **Germain H, Seguin A. 2011.** Innate immunity: has poplar made its BED? *New Phytol* **189**(3):
898    678-687.

899 **Hallwass M, de Oliveira AS, de Campos Dianese E, Lohuis D, Boiteux LS, Inoue-Nagata AK,**
900    **Resende RO, Kormelink R. 2014.** The Tomato spotted wilt virus cell-to-cell movement
901    protein (NSM) triggers a hypersensitive response in Sw-5-containing resistant tomato lines
902    and in Nicotiana benthamiana transformed with the functional Sw-5b resistance gene copy.
903    *Mol Plant Pathol* **15**(9): 871-880.

904 **Heidrich K, Wirthmueller L, Tasset C, Pouzet C, Deslandes L, Parker JE. 2011.** Arabidopsis
905    EDS1 connects pathogen effector recognition to cell compartment-specific immune responses.
906    *Science* **334**(6061): 1401-1404.

907 **Heinlein M. 2015.** Plant virus replication and movement. *Virology* **479-480**: 657-671.

908 **Horsefield S, Burdett H, Zhang X, Manik MK, Shi Y, Chen J, Qi T, Gilley J, Lai JS, Rank MX,**
909    **Casey LW, Gu W, Ericsson DJ, Foley G, Hughes RO, Bosanac T, von Itzstein M,**
910    **Rathjen JP, Nanson JD, Boden M, Dry IB, Williams SJ, Staskawicz BJ, Coleman MP,**
911    **Ve T, Dodds PN, Kobe B. 2019.** NAD(+) cleavage activity by animal and plant TIR domains
912    in cell death pathways. *Science* **365**(6455): 793-799.

913 **Inoue H, Hayashi N, Matsushita A, Xinqiong L, Nakayama A, Sugano S, Jiang CJ, Takatsuji H.**
914    **2013.** Blast resistance of CC-NB-LRR protein Pb1 is mediated by WRKY45 through

protein-protein interaction. *Proc Natl Acad Sci U S A* **110**(23): 9577-9582.

Jones JD, Vance RE, Dangl JL. 2016. Intracellular innate immune surveillance devices in plants and

animals. *Science* **354**(6316).

Jubic LM, Saile S, Furzer OJ, El Kasmi F, Dangl JL. 2019. Help wanted: helper NLRs and plant

immune responses. *Curr Opin Plant Biol* **50**: 82-94.

Kanneganti TD, Bai X, Tsai CW, Win J, Meulia T, Goodin M, Kamoun S, Hogenhout SA. 2007.

A functional genetic assay for nuclear trafficking in plants. *Plant J* **50**(1): 149-158.

Kanzaki H, Yoshida K, Saitoh H, Fujisaki K, Hirabuchi A, Alaux L, Fournier E, Tharreau D,

Terauchi R. 2012. Arms race co-evolution of Magnaporthe oryzae AVR-Pik and rice Pik

genes driven by their physical interactions. *Plant J* **72**(6): 894-907.

Kapos P, Devendrakumar KT, Li X. 2019. Plant NLRs: From discovery to application. *Plant Sci* **279**:

3-18.

Katagiri F, Tsuda K. 2010. Understanding the plant immune system. *Mol Plant Microbe Interact*

23(12): 1531-1536.

Kawano Y, Fujiwara T, Yao A, Housen Y, Hayashi K, Shimamoto K. 2014.

Palmitoylation-dependent membrane localization of the rice resistance protein pit is critical

for the activation of the small GTPase OsRac1. *J Biol Chem* **289**(27): 19079-19088.

Kong L, Qiu X, Kang J, Wang Y, Chen H, Huang J, Qiu M, Zhao Y, Kong G, Ma Z, Wang Y, Ye

W, Dong S, Ma W, Wang Y. 2017. A Phytophthora Effector Manipulates Host Histone

Acetylation and Reprograms Defense Gene Expression to Promote Infection. *Curr Biol* **27**(7):

981-991.

Kormelink R, Garcia ML, Goodin M, Sasaya T, Haenni AL. 2011. Negative-strand RNA viruses:

the plant-infecting counterparts. *Virus Res* **162**(1-2): 184-202.

Kormelink R, Storms M, Van Lent J, Peters D, Goldbach R. 1994. Expression and subcellular

location of the NSM protein of tomato spotted wilt virus (TSWV), a putative viral movement

protein. *Virology* **200**(1): 56-65.

Kourelis J, van der Hoorn RAL. 2018. Defended to the Nines: 25 Years of Resistance Gene Cloning

Identifies Nine Mechanisms for R Protein Function. *Plant Cell* **30**(2): 285-299.

Krasileva KV, Dahlbeck D, Staskawicz BJ. 2010. Activation of an Arabidopsis resistance protein is

specified by the in planta association of its leucine-rich repeat domain with the cognate

oomycete effector. *Plant Cell* **22**(7): 2444-2458.

Kroj T, Chanclud E, Michel-Romiti C, Grand X, Morel JB. 2016. Integration of decoy domains

derived from protein targets of pathogen effectors into plant immune receptors is widespread.

New Phytol **210**(2): 618-626.

Lanford RE, Butel JS. 1984. Construction and characterization of an SV40 mutant defective in

nuclear transport of T antigen. *Cell* **37**(3): 801-813.

Lapin D, Bhandari DD, Parker JE. 2020. Origins and Immunity Networking Functions of EDS1

Family Proteins. *Annu Rev Phytopathol* **58**: 253-276.

Le Roux C, Huet G, Jauneau A, Camborde L, Tremousaygue D, Kraut A, Zhou B, Levaillant M,

Adachi H, Yoshioka H, Raffaele S, Berthome R, Coute Y, Parker JE, Deslandes L. 2015.

A receptor pair with an integrated decoy converts pathogen disabling of transcription factors

to immunity. *Cell* **161**(5): 1074-1088.

Leastro MO, Pallas V, Resende RO, Sanchez-Navarro JA. 2017. The functional analysis of distinct

tospovirus movement proteins (NSM) reveals different capabilities in tubule formation,

cell-to-cell and systemic virus movement among the tospovirus species. *Virus Res* **227**: 57-68.

Li J, Huang H, Zhu M, Huang S, Zhang W, Dinesh-Kumar SP, Tao X. 2019. A Plant Immune
Receptor Adopts a Two-Step Recognition Mechanism to Enhance Viral Effector Perception.
Mol Plant **12**(2): 248-262.

Li W, Lewandowski DJ, Hilf ME, Adkins S. 2009. Identification of domains of the Tomato spotted
wilt virus NSm protein involved in tubule formation, movement and symptomatology.
Virology **390**(1): 110-121.

Li W, Zhao Y, Liu C, Yao G, Wu S, Hou C, Zhang M, Wang D. 2012. Callose deposition at
plasmodesmata is a critical factor in restricting the cell-to-cell movement of Soybean mosaic
virus. *Plant Cell Rep* **31**(5): 905-916.

Li X, Kapos P, Zhang Y. 2015. NLRs in plants. *Curr Opin Immunol* **32**: 114-121.

Lopez C, Aramburu J, Galipienso L, Soler S, Nuez F, Rubio L. 2011. Evolutionary analysis of
tomato Sw-5 resistance-breaking isolates of Tomato spotted wilt virus. *J Gen Virol* **92**(Pt 1):
210-215.

Lucas WJ. 2006. Plant viral movement proteins: agents for cell-to-cell trafficking of viral genomes.
Virology **344**(1): 169-184.

Lukasik-Shreepaathy E, Slootweg E, Richter H, Goverse A, Cornelissen BJ, Takken FL. 2012.
Dual regulatory roles of the extended N terminus for activation of the tomato MI-1.2
resistance protein. *Mol Plant Microbe Interact* **25**(8): 1045-1057.

**Ma S, Lapin D, Liu L, Sun Y, Song W, Zhang X, Logemann E, Yu D, Wang J, Jirschitzka J, Han
Z, Schulze-Lefert P, Parker JE, Chai J. 2020.** Direct pathogen-induced assembly of an
NLR immune receptor complex to form a holoenzyme. *Science* **370**(6521).

**Ma Y, Guo H, Hu L, Martinez PP, Moschou PN, Cevik V, Ding P, Duxbury Z, Sarris PF, Jones
JDG. 2018.** Distinct modes of derepression of an Arabidopsis immune receptor complex by
two different bacterial effectors. *Proc Natl Acad Sci U S A* **115**(41): 10218-10227.

**Ma Z, Song T, Zhu L, Ye W, Wang Y, Shao Y, Dong S, Zhang Z, Dou D, Zheng X, Tyler BM,
Wang Y. 2015.** A Phytophthora sojae Glycoside Hydrolase 12 Protein Is a Major Virulence
Factor during Soybean Infection and Is Recognized as a PAMP. *Plant Cell* **27**(7): 2057-2072.

**Ma Z, Zhu L, Song T, Wang Y, Zhang Q, Xia Y, Qiu M, Lin Y, Li H, Kong L, Fang Y, Ye W,
Wang Y, Dong S, Zheng X, Tyler BM, Wang Y. 2017.** A paralogous decoy protects
Phytophthora sojae apoplastic effector PsXEG1 from a host inhibitor. *Science* **355**(6326):
710-714.

**Maekawa T, Cheng W, Spiridon LN, Toller A, Lukasik E, Saijo Y, Liu P, Shen QH, Micluta MA,
Somssich IE, Takken FLW, Petrescu AJ, Chai J, Schulze-Lefert P. 2011.** Coiled-coil
domain-dependent homodimerization of intracellular barley immune receptors defines a
minimal functional module for triggering cell death. *Cell Host Microbe* **9**(3): 187-199.

Meier N, Hatch C, Nagalakshmi U, Dinesh-Kumar SP. 2019. Perspectives on intracellular
perception of plant viruses. *Mol Plant Pathol* **20**(9): 1185-1190.

Meyers BC, Kozik A, Griego A, Kuang H, Michelmore RW. 2003. Genome-wide analysis of
NBS-LRR-encoding genes in Arabidopsis. *Plant Cell* **15**(4): 809-834.

Milligan SB, Bodeau J, Yaghoobi J, Kaloshian I, Zabel P, Williamson VM. 1998. The root knot
nematode resistance gene Mi from tomato is a member of the leucine zipper, nucleotide
binding, leucine-rich repeat family of plant genes. *Plant Cell* **10**(8): 1307-1319.

Mittler R, Herr EH, Orvar BL, van Camp W, Willekens H, Inze D, Ellis BE. 1999. Transgenic

tobacco plants with reduced capability to detoxify reactive oxygen intermediates are hyperresponsive to pathogen infection. *Proc Natl Acad Sci U S A* **96**(24): 14165-14170.

Oliver JE, Whitfield AE. 2016. The Genus Tospovirus: Emerging Bunyaviruses that Threaten Food Security. *Annu Rev Virol* **3**(1): 101-124.

Padmanabhan MS, Ma S, Burch-Smith TM, Czymmek K, Huijser P, Dinesh-Kumar SP. 2013. Novel positive regulatory role for the SPL6 transcription factor in the N TIR-NB-LRR receptor-mediated plant innate immunity. *PLoS Pathog* **9**(3): e1003235.

Palma K, Zhang Y, Li X. 2005. An importin alpha homolog, MOS6, plays an important role in plant innate immunity. *Curr Biol* **15**(12): 1129-1135.

Peiro A, Canizares MC, Rubio L, Lopez C, Moriones E, Aramburu J, Sanchez-Navarro J. 2014. The movement protein (NSm) of Tomato spotted wilt virus is the avirulence determinant in the tomato Sw-5 gene-based resistance. *Mol Plant Pathol* **15**(8): 802-813.

Qi D, DeYoung BJ, Innes RW. 2012. Structure-function analysis of the coiled-coil and leucine-rich repeat domains of the RPS5 disease resistance protein. *Plant Physiol* **158**(4): 1819-1832.

Qi D, Innes RW. 2013. Recent Advances in Plant NLR Structure, Function, Localization, and Signaling. *Front Immunol* **4**: 348.

Rao ALNC, Y. G，Khan, J. A，Dijkstra, J. 2002. Molecular biology of plant virus movement. *Plant Viruses As Molecular Pathogens*.

Sarris PF, Cevik V, Dagdas G, Jones JD, Krasileva KV. 2016. Comparative analysis of plant immune receptor architectures uncovers host proteins likely targeted by pathogens. *BMC Biol* **14**: 8.

Sarris PF, Duxbury Z, Huh SU, Ma Y, Segonzac C, Sklenar J, Derbyshire P, Cevik V, Rallapalli G, Saucet SB, Wirthmueller L, Menke FL, Sohn KH, Jones JD. 2015. A Plant Immune Receptor Detects Pathogen Effectors that Target WRKY Transcription Factors. *Cell* **161**(5): 1089-1100.

Scholthof KB, Adkins S, Czosnek H, Palukaitis P, Jacquot E, Hohn T, Hohn B, Saunders K, Candresse T, Ahlquist P, Hemenway C, Foster GD. 2011. Top 10 plant viruses in molecular plant pathology. *Mol Plant Pathol* **12**(9): 938-954.

Seo JK, Kwon SJ, Cho WK, Choi HS, Kim KH. 2014. Type 2C protein phosphatase is a key regulator of antiviral extreme resistance limiting virus spread. *Sci Rep* **4**: 5905.

Seong K, Seo E, Witek K, Li M, Staskawicz B. 2020. Evolution of NLR resistance genes with noncanonical N-terminal domains in wild tomato species. *New Phytol* **227**(5): 1530-1543.

Shen QH, Saijo Y, Mauch S, Biskup C, Bieri S, Keller B, Seki H, Ulker B, Somssich IE, Schulze-Lefert P. 2007. Nuclear activity of MLA immune receptors links isolate-specific and basal disease-resistance responses. *Science* **315**(5815): 1098-1103.

Slootweg E, Roosien J, Spiridon LN, Petrescu AJ, Tameling W, Joosten M, Pomp R, van Schaik C, Dees R, Borst JW, Smant G, Schots A, Bakker J, Goverse A. 2010. Nucleocytoplasmic distribution is required for activation of resistance by the potato NB-LRR receptor Rx1 and is balanced by its functional domains. *Plant Cell* **22**(12): 4195-4215.

Soosaar JL, Burch-Smith TM, Dinesh-Kumar SP. 2005. Mechanisms of plant resistance to viruses. *Nat Rev Microbiol* **3**(10): 789-798.

Spassova M I PTW, Folkertsma R T, et al. 2001. The tomato gene Sw-5 is a member of the coiled coil, nucleotide binding, leucine-rich repeat class of plant resistance genes and confers resistance to TSWV in tobacco. *Molecular Breeding.* **7(2)**: 151.

Stewart M. 2007. Molecular mechanism of the nuclear protein import cycle. *Nat Rev Mol Cell Biol*
8(3): 195-208.

Swiderski MR, Birker D, Jones JD. 2009. The TIR domain of TIR-NB-LRR resistance proteins is a
signaling domain involved in cell death induction. *Mol Plant Microbe Interact* **22**(2):
157-165.

Takemoto D, Rafiqi M, Hurley U, Lawrence GJ, Bernoux M, Hardham AR, Ellis JG, Dodds PN,
Jones DA. 2012. N-terminal motifs in some plant disease resistance proteins function in
membrane attachment and contribute to disease resistance. *Mol Plant Microbe Interact* **25**(3):
379-392.

Takken FLW, Goverse A. 2012. How to build a pathogen detector: structural basis of NB-LRR
function. *Current Opinion in Plant Biology* **15**(4): 375-384.

Taliansky M, Torrance L, Kalinina NO. 2008. Role of plant virus movement proteins. *Methods Mol*
Biol **451**: 33-54.

Tameling WI, Nooijen C, Ludwig N, Boter M, Slootweg E, Goverse A, Shirasu K, Joosten MH.
2010. RanGAP2 mediates nucleocytoplasmic partitioning of the NB-LRR immune receptor
Rx in the Solanaceae, thereby dictating Rx function. *Plant Cell* **22**(12): 4176-4194.

Tasset C, Bernoux M, Jauneau A, Pouzet C, Briere C, Kieffer-Jacquinod S, Rivas S, Marco Y,
Deslandes L. 2010. Autoacetylation of the Ralstonia solanacearum effector PopP2 targets a
lysine residue essential for RRS1-R-mediated immunity in Arabidopsis. *PLoS Pathog* **6**(11):
e1001202.

Turina M, Kormelink R, Resende RO. 2016. Resistance to Tospoviruses in Vegetable Crops:
Epidemiological and Molecular Aspects. *Annu Rev Phytopathol* **54**: 347-371.

van der Vossen EA, Gros J, Sikkema A, Muskens M, Wouters D, Wolters P, Pereira A, Allefs S.
2005. The Rpi-blb2 gene from Solanum bulbocastanum is an Mi-1 gene homolog conferring
broad-spectrum late blight resistance in potato. *Plant J* **44**(2): 208-222.

van Wersch S, Li X. 2019. Stronger When Together: Clustering of Plant NLR Disease resistance
Genes. *Trends Plant Sci* **24**(8): 688-699.

van Wersch S, Tian, L., Hoy, R., Li, X. . 2020. Plant NLRs: The whistleblowers of
plant immunity. *Plant Commun*: doi: https://doi.org/10.1016/j.xplc.2019.100016.

Vos P, Simons G, Jesse T, Wijbrandi J, Heinen L, Hogers R, Frijters A, Groenendijk J,
Diergaarde P, Reijans M, Fierens-Onstenk J, de Both M, Peleman J, Liharska T,
Hontelez J, Zabeau M. 1998. The tomato Mi-1 gene confers resistance to both root-knot
nematodes and potato aphids. *Nat Biotechnol* **16**(13): 1365-1369.

Vossen JH, van Arkel G, Bergervoet M, Jo KR, Jacobsen E, Visser RG. 2016. The Solanum
demissum R8 late blight resistance gene is an Sw-5 homologue that has been deployed
worldwide in late blight resistant varieties. *Theor Appl Genet* **129**(9): 1785-1796.

Wan L, Essuman K, Anderson RG, Sasaki Y, Monteiro F, Chung EH, Osborne Nishimura E,
DiAntonio A, Milbrandt J, Dangl JL, Nishimura MT. 2019. TIR domains of plant immune
receptors are NAD(+)-cleaving enzymes that promote cell death. *Science* **365**(6455): 799-803.

Wang A. 2015. Dissecting the molecular network of virus-plant interactions: the complex roles of host
factors. *Annu Rev Phytopathol* **53**: 45-66.

Wang J, Chen T, Han M, Qian L, Li J, Wu M, Han T, Cao J, Nagalakshmi U, Rathjen JP, Hong
Y, Liu Y. 2020. Plant NLR immune receptor Tm-2² activation requires NB-ARC
domain-mediated self-association of CC domain. *PLoS Pathog* **16**(4): e1008475.

Wang J, Hu M, Wang J, Qi J, Han Z, Wang G, Qi Y, Wang HW, Zhou JM, Chai J. 2019a.
Reconstitution and structure of a plant NLR resistosome conferring immunity. *Science*
364(6435).

Wang J, Wang J, Hu M, Wu S, Qi J, Wang G, Han Z, Qi Y, Gao N, Wang HW, Zhou JM, Chai J.
2019b. Ligand-triggered allosteric ADP release primes a plant NLR complex. *Science*
364(6435).

Wen W, Meinkoth JL, Tsien RY, Taylor SS. 1995. Identification of a signal for rapid export of
proteins from the nucleus. *Cell* **82**(3): 463-473.

Wirthmueller L, Zhang Y, Jones JD, Parker JE. 2007. Nuclear accumulation of the Arabidopsis
immune receptor RPS4 is necessary for triggering EDS1-dependent defense. *Curr Biol* **17**(23):
2023-2029.

Zhang Y, Li X. 2005. A putative nucleoporin 96 Is required for both basal defense and constitutive
resistance responses mediated by suppressor of npr1-1,constitutive 1. *Plant Cell* **17**(4):
1306-1316.

Zhao W, Jiang L, Feng Z, Chen X, Huang Y, Xue F, Huang C, Liu Y, Li F, Liu Y, Tao X. 2016.
Plasmodesmata targeting and intercellular trafficking of Tomato spotted wilt tospovirus
movement protein NSm is independent of its function in HR induction. *J Gen Virol* **97**(8):
1990-1997.

Zhu M, Jiang L, Bai B, Zhao W, Chen X, Li J, Liu Y, Chen Z, Wang B, Wang C, Wu Q, Shen Q,
Dinesh-Kumar SP, Tao X. 2017. The Intracellular Immune Receptor Sw-5b Confers
Broad-Spectrum Resistance to Tospoviruses through Recognition of a Conserved 21-Amino
Acid Viral Effector Epitope. *Plant Cell* **29**(9): 2214-2232.

Zhu M, van Grinsven IL, Kormelink R, Tao X. 2019. Paving the Way to Tospovirus Infection:
Multilined Interplays with Plant Innate Immunity. *Annu Rev Phytopathol* **57**: 41-62.

1115

1116

1117

1118

 FIGURE LEGENDS

Fig. 1. Subcellular localization of Sw-5b in *Nicotiana benthamiana* and tomato leaf

cells. (a) Schematic diagrams of Sw-5b. (b) Subcellular localizations of free YFP

(left), YFP-Sw-5b (middle) and autoactive YFP-Sw-5b^{D857V} mutant (right) in *N.*

benthamiana leaf cells at 24 hours post agro-infiltration (hpi). (c) Subcellular

localization of free YFP (left), YFP-Sw-5b (middle) and autoactive YFP-Sw-5b^{D857V}

mutant (right) in tomato leaf cells at 24 hpi. N nucleus, and C cytoplasm inside the

cell are indicated. Bar = 10 μm. (d) Nucleocytoplasmic partitioning analysis of

YFP-Sw-5b and autoactive YFP-Sw-5b^{D857V}. Total lysate (T) from p2300S empty

vector (EV), YFP-Sw-5b or YFP-Sw-5b^{D857V} expressing leaves were fractionated into

cytoplasm and nucleus, and analyzed by immunoblots using antibodies against YFP.

The actin and histone were used as a cytoplasm marker and nucleus marker,

respectively, in the fractionation analysis. Ponceau S staining was also used as

cytoplasm marker.

1133

Fig. 2. Effect of Sw-5b subcellular localization pattern on HR induction. (a) Confocal

images of *N. benthamiana* leaf cells transiently expressing NES-YFP-Sw-5b,

nes-YFP-Sw-5b, NLS-YFP-Sw-5b or nls-YFP-Sw-5b fusion. The images were taken

at 24–36 hpi. N nucleus and C cytoplasm (c). Bar = 10 μm. (b) Induction of HR in *N.*

benthamiana leaf tissues co-expressing NSm and one of the five Sw-5b fusion

proteins. The infiltrated *N. benthamiana* leaf was photographed at 3 dpi (left image).

Induction of HR in the infiltrated tissues were visualized using a trypan blue staining

method (right image). (c) Immunoblot analysis of NES-YFP-Sw-5b, nes-YFP-Sw-5b,

NLS-YFP-Sw-5b, and nls-YFP-Sw-5b expressions in the infiltrated *N. benthamiana*

leaf tissues. These fusion proteins were enriched using the GFP-Trap beads prior to

SDS-PAGE, and the blot was probed using an YFP specific antibody. Ponceau-S

staining was used to estimate sample loadings. (d) Time course analysis of ion

leakage in Nicotiana benthamiana leaves co-expressing NSm with one of the five

Sw-5b fusion proteins. Measurements were performed at 4 h intervals starting from

24 to 48 hpi. Error bars (SEs) were calculated using the results from three biological

replicates per treatment collected at each time point.

1150

Fig. 3. The effect of cytoplasm- and nucleus-targeted Sw-5b on viral replication. (a)

Schematic representation of binary constructs to express TSWV $SR_{(-)eGFP}$

mini-genome replicon, TSWV L RNA segment containing an optimized RdRp and

$NSm^{H93A\&H94A}$ mutant that defected in viral movement. Minus sign (-) and 5′ to 3′

designation represent the negative (genomic)-strand of tospovirus RNA. 35S: a

double 35S promoter; HH: hammerhead ribozyme; RZ: hepatitis delta virus (HDV)

ribozyme; NOS: nopaline synthase terminator; 35S Ter: a 35S transcription terminator.

(b) Accumulation of eGFP fluorescence in *N. benthamiana* leaves co-expressing

p2300S empty vector (EV), Sw-5b, NES-Sw-5b, nes-Sw-5b, NLS-Sw-5b, or

nls-Sw-5b with TSWV $SR_{(-)eGFP}$, L, and $NSm^{H93A\&H94A}$ at 4 days post infiltration (dpi)

viewed with a fluorescence microscope. Bar represents 400 μm. (c) Immunoblot

analysis of expression of eGFP proteins in leaves shown in panel (b) using specific

antibodies against YFP. Ponceau S staining of rubisco large subunit is shown for

protein loading control.

1165

Fig. 4. Effect of subcellular localization of Sw-5b on cell-to-cell movement of NSm

in leaf epidermis of *N. benthamiana.* (a) Schematic diagram of the binary construct to

co-express mCherry-HDEL and NSm-GFP. (b) Cell-to-cell movement analysis of

NSm-GFP in *N. benthamiana* leaves co-expressing p2300S empty vector (EV), Sw-5b,

NES-Sw-5b, NLS-Sw-5b, or nls-Sw-5b with the construct harboring both

mCherry-HDEL and NSm-GFP. *Agrobacterium* containing the construct to co-express

mCherry-HDEL and NSm-GFP was diluted 500 times for expression in a single

epidermal cell. All other *Agrobacterium* were infiltrated at the concentration of OD_{600}

= 0.2. Bar, 50 μm.

1175

Fig. 5. Analysis of cytoplasm- and nucleus-targeted Sw-5b-mediated host immunity

to TSWV systemic infection. (a) TSWV systemic infection in transgenic *N.*

benthamiana plants expressing NES-YFP-Sw-5b, nes-YFP-Sw-5b, NLS-YFP-Sw-5b,

nls-YFP-Sw-5b, YFP-Sw-5b or p2300S empty vector (EV) driven by 35S promoter.

TSWV-inoculated plants were photographed at 15 dpi. White arrow indicates the

systemic leaves showing HR trailing. White arrowhead indicates the systemic leaves

showing mosaic. (b) Immunoblot analysis of NES-YFP-Sw-5b, nes-YFP-Sw-5b,

NLS-YFP-Sw-5b, nls-YFP-Sw-5b and YFP-Sw-5b expressions in different transgenic

N. benthamiana plants. EV plants transformed with an empty vector and were used as

a negative control. (c) RT-PCR analysis of TSWV accumulation in the systemic

leaves of different transgenic *N. benthamiana* plants at 15 dpi.

1187

Fig. 6. Joint effects of cytoplasm- and nucleus-targeted Sw-5b on defenses against

tospovirus infection in *Nicotiana benthamiana* leaves. (a) Schematic representation of

binary constructs to express TSWV SR$_{(+)eGFP}$-M$_{(-)op}$ and TSWV L RNA segment

containing an optimized RdRp. 35S: a double 35S promoter; HH: hammerhead

ribozyme; RZ: hepatitis delta virus (HDV) ribozyme; NOS: nopaline synthase

terminator. (b) Accumulation of eGFP fluorescence in *N. benthamiana* leaves

co-expressing p2300S empty vector (EV), Sw-5b, NES-Sw-5b, NLS-Sw-5b, or

NES-Sw-5b+NLS-Sw-5b with TSWV SR$_{(+)eGFP}$-M$_{(-)op}$ at 4 days post infiltration (dpi)

viewed with a fluorescence microscope. Bar represents 400 μm. (c) Immunoblot

analysis of expression of eGFP proteins in leaves shown in panel (b) using specific

antibodies against YFP. Ponceau S staining of rubisco large subunit is shown for

protein loading control. (d) Quantification of eGFP proteins in leaves shown in panel

(c).

1201

Fig. 7. Functional analysis and subcellular localization patterns of individual or

combined Sw-5b domains. (a) Schematic diagrams showing a full length Sw-5b or

Sw-5b domains fused with YFP. (b) Confocal images of *N. benthamiana* leaf

epidermal cells expressing various YFP fusions. Images of the cells were taken at 24

hpi. N nucleus, Nu nucleolus, and C cytoplasm. Bar = 10 μm. (c) TSWV-inoculated

transgenic *N. benthamiana* plants expressing these various YFP fusions and

photographed at 15 dpi.

1209

Fig. 8. Roles of *importins* α and β in YFP-Sw-5b nucleus targeting and

Sw-5b-mediated immunity to TSWV systemic infection. (a) Transient expression of

YFP-Sw-5b in *N. benthamiana* leaf epidermal cells silenced for *importin*

$\alpha1$ (IMP α1 KD), *importin* $\alpha2$ (IMP α2 KD), *importin* $\alpha1$ and $\alpha2$ (IMP α1 & α2 KD),

importin β (IMP β KD) or *importin* $\alpha1$ and $\alpha2$ and β (IMP α1 & α2 & β KD) through

VIGS. Images of the cells were captured using a confocal microscope at 26 hpi. N

nucleus, C cytoplasm. Bar = 10 μm. (b) YFP-Sw-5b transgenic *N. benthamiana* plants

were silenced for *importin* $\alpha1$, *importin* $\alpha2$, *importin* $\alpha1$ and $\alpha2$, *importin* β or

importin $\alpha1$ and $\alpha2$ and β expression through VIGS followed by inoculation with

TSWV. TSWV-inoculated YFP-Sw-5b transgenic *N. benthamiana* plants were

photographed at 15 dpi. White arrowhead indicates HR trailing in systemic leaves.

1221

Fig. 9. A working model for Sw-5b. Sw-5b furcates disease resistances by proper

nucleocytoplasmic partition to block different infection steps of tomato spotted wilt

tospovirus. Sw-5b switched from the autoinhibited state to an activated state upon

recognition of NSm in the cytoplasm. Cytoplasm portion of Sw-5b induce cell death

and defense that inhibit viral replication. The activated Sw-5b also translocated into

nucleus via *importins* α and β. Nucleus-localized Sw-5b induces a defense that block

viral cell-to-cell and long-distance movement. Cytoplasm- and nucleus-localized

Sw-5b have additively effects on defense to inhibit viral replication, intercellular and

long-distance movement during tospovirus infection.

1231

Table S2. Response of six different types of transgenic *Nicotiana benthamina* plants

driven by 35S promoter to TSWV infection.

Table S3. Response of six different types of transgenic *Nicotiana benthamina* plants

driven by Sw-5b native promoter to TSWV infection.

Table S4. Response of six different types of transgenic *Nicotiana benthamina* plants

to TSWV infection.

Table S5. Mass spectrum data of YFP-SD

Table S6. Mass spectrum data of YFP-Sw-5b

1261

Fig. 1. Subcellular localization of Sw-5b in *Nicotiana benthamiana* and tomato leaf cells. (a) Schematic diagram of Sw-5b. (b) Subcellular localizations of free YFP (left), YFP-Sw-5b (middle) and autoactive YFP-Sw-5b^{D857V} mutant (right) in *N. benthamiana* leaf cells at 24 hours post agro-infiltration (hpi). (c) Subcellular localization of free YFP (left), YFP-Sw-5b (middle) and autoactive YFP-Sw-5b^{D857V} mutant (right) in tomato leaf cells at 24 hpi. N nucleus, and C cytoplasm inside the cell are indicated. Bar = 10 μm. (d) Nucleocytoplasmic partitioning analysis of YFP-Sw-5b and autoactive YFP-Sw-5b^{D857V}. Total lysate (T) from p2300S empty vector (EV), YFP-Sw-5b or YFP-Sw-5b^{D857V} expressing leaves were fractionated into cytoplasm and nucleus, and analyzed by immunoblots using antibodies against YFP. The actin and histone were used as a cytoplasm marker and nucleus marker, respectively, in the fractionation analysis. Ponceau S staining was also used as cytoplasm marker.

Fig. 2. Effect of Sw-5b subcellular localization pattern on HR induction. (a) Confocal images of *N. benthamiana* leaf cells transiently expressing NES-YFP-Sw-5b, nes-YFP-Sw-5b, NLS-YFP-Sw-5b or nls-YFP-Sw-5b fusion. The images were taken at 24–36 hpi. N nucleus and C cytoplasm (c). Bar = 10 μm. (b) Induction of HR in *N. benthamiana* leaf tissues co-expressing NSm and one of the five Sw-5b fusion proteins. The infiltrated *N. benthamiana* leaf was photographed at 3 dpi (left image). Induction of HR in the infiltrated tissues were visualized using a trypan blue staining method (right image). (c) Immunoblot analysis of NES-YFP-Sw-5b, nes-YFP-Sw-5b, NLS-YFP-Sw-5b, and nls-YFP-Sw-5b expressions in the infiltrated *N. benthamiana* leaf tissues. These fusion proteins were enriched using the GFP-Trap beads prior to SDS-PAGE, and the blot was probed using an YFP specific antibody. Ponceau-S staining was used to estimate sample loadings. (d) Time course analysis of ion leakage in Nicotiana benthamiana leaves co-expressing NSm with one of the five Sw-5b fusion proteins. Measurements were performed at 4 h intervals starting from 24 to 48 hpi. Error bars (SEs) were calculated using the results from three biological replicates per treatment collected at each time point.

(a)

(b)

SR$_{(+)eGFP}$ + L$_{(+)opt}$ + NSm$^{H93A\&H94A}$ + VSRs

EV	NES-Sw-5b	nes-Sw-5b
Sw-5b	NLS-Sw-5b	nls-Sw-5b

(c)

SR$_{(+)eGFP}$ + L$_{(+)opt}$ + NSm$^{H93A\&H94A}$ + VSRs

Fig. 3. The effect of cytoplasm- and nucleus-targeted Sw-5b on viral replication. (a) Schematic representation of binary constructs to express TSWV $SR_{(-)eGFP}$ mini-genome replicon, TSWV L RNA segment containing an optimized RdRp and $NSm^{H93A\&H94A}$ mutant that defected in viral movement. Minus sign (-) and 5′ to 3′ designation represent the negative (genomic)-strand of tospovirus RNA. 35S: a double 35S promoter; HH: hammerhead ribozyme; RZ: hepatitis delta virus (HDV) ribozyme; NOS: nopaline synthase terminator; 35S Ter: a 35S transcription terminator. (b) Accumulation of eGFP fluorescence in *N. benthamiana* leaves co-expressing p2300S empty vector (EV), Sw-5b, NES-Sw-5b, nes-Sw-5b, NLS-Sw-5b, or nls-Sw-5b with TSWV $SR_{(-)eGFP}$, L, and $NSm^{H93A\&H94A}$ at 4 days post infiltration (dpi) viewed with a fluorescence microscope. Bar represents 400 μm. (c) Immunoblot analysis of expression of eGFP proteins in leaves shown in panel (b) using specific antibodies against YFP. Ponceau S staining of rubisco large subunit is shown for protein loading control.

(a)

mCherry-HDEL//NSm-GFP

35S | mCherry-HDEL | 35St | 35S | NSm | GFP | 35St

(b)

Fig. 4. Effect of subcellular localization of Sw-5b on cell-to-cell movement of NSm in leaf epidermis of *N. benthamiana.* (a) Schematic diagram of the binary construct to co-express mCherry-HDEL and NSm-GFP. (b) Cell-to-cell movement analysis of NSm-GFP in *N. benthamiana* leaves co-expressing p2300S empty vector (EV), Sw-5b, NES-Sw-5b, NLS-Sw-5b, or nls-Sw-5b with the construct harboring both mCherry-HDEL and NSm-GFP. *Agrobacterium* containing the construct to co-express mCherry-HDEL and NSm-GFP was diluted 500 times for expression in a single epidermal cell. All other *Agrobacterium* were infiltrated at the concentration of OD_{600} = 0.2. Bar = 50 μm.

Fig. 5. Analysis of cytoplasm- and nucleus-targeted Sw-5b-mediated host immunity to TSWV systemic infection. (a) TSWV systemic infection in transgenic *N. benthamiana* plants expressing NES-YFP-Sw-5b, nes-YFP-Sw-5b, NLS-YFP-Sw-5b, nls-YFP-Sw-5b, YFP-Sw-5b or p2300S empty vector (EV) driven by 35S promoter. TSWV-inoculated plants were photographed at 15 dpi. White arrow indicates the systemic leaves showing HR trailing. White arrowhead indicates the systemic leaves showing mosaic. (b) Immunoblot analysis of NES-YFP-Sw-5b, nes-YFP-Sw-5b, NLS-YFP-Sw-5b, nls-YFP-Sw-5b and YFP-Sw-5b expressions in different transgenic *N. benthamiana* plants. EV plants transformed with an empty vector and were used as a negative control. (c) RT-PCR analysis of TSWV accumulation in the systemic leaves of different transgenic *N. benthamiana* plants at 15 dpi.

A

B

L$_{(+)opt}$+M$_{(-)opt}$-SR$_{(+)eGFP}$

C

L$_{(+)opt}$+M$_{(-)opt}$-SR$_{(+)eGFP}$

kDa

1.00   0.55   0.17   0.05   0.02   0.00

α-YFP

Ponceau S

D

L$_{(+)opt}$+M$_{(-)opt}$-SR$_{(+)eGFP}$

Fig. 6. Joint effects of cytoplasm- and nucleus-targeted Sw-5b on defenses against tospovirus infection in *Nicotiana benthamiana* leaves. (a) Schematic representation of binary constructs to express TSWV $SR_{(+)eGFP}$-$M_{(-)op}$ and TSWV L RNA segment containing an optimized RdRp. 35S: a double 35S promoter; HH: hammerhead ribozyme; RZ: hepatitis delta virus (HDV) ribozyme; NOS: nopaline synthase terminator. (b) Accumulation of eGFP fluorescence in *N. benthamiana* leaves co-expressing p2300S empty vector (EV), Sw-5b, NES-Sw-5b, NLS-Sw-5b, or NES-Sw-5b+NLS-Sw-5b with TSWV $SR_{(+)eGFP}$-$M_{(-)op}$ at 4 days post infiltration (dpi) viewed with a fluorescence microscope. Bar represents 400 µm. (c) Immunoblot analysis of expression of eGFP proteins in leaves shown in panel (b) using specific antibodies against YFP. Ponceau S staining of rubisco large subunit is shown for protein loading control. (d) Quantification of eGFP proteins in leaves shown in panel (c).

Fig. 7. Functional analysis and subcellular localization patterns of individual or combined Sw-5b domains. (a) Schematic diagram showing a full length Sw-5b or Sw-5b domains fused with YFP. (b) Confocal images of *N. benthamiana* leaf epidermal cells expressing various YFP fusions. Images of the cells were taken at 24 hpi. N nucleus, and C cytoplasm. Bar = 10 μm. (c) TSWV-inoculated transgenic *N. benthamiana* plants expressing these various YFP fusions and photographed at 15 dpi.

Fig. 8. Roles of *importins α* and *β* in YFP-Sw-5b nucleus targeting and Sw-5b-mediated immunity to TSWV systemic infection. (a) Transient expression of YFP-Sw-5b in *N. benthamiana* leaf epidermal cells silenced for *importin α1* (IMP α1 KD)*, importinα2* (IMP α2 KD), *importin α1* and *α2* (IMP α1 & α2 KD), *importin β* (IMP β KD) or *importin α1 and β2* and *β* (IMP α1 & α2 & β KD) through VIGS. Images of the cells were captured using a confocal microscope at 26 hpi. N nucleus, C cytoplasm. Bar = 10 mm. (b) YFP-Sw-5b transgenic *N. benthamiana* plants were silenced for *importin a1, importin α2, importin α1* and *α2, importin β* or *importin α1* and *α2* and *β* expression through VIGS followed by inoculation with TSWV. TSWV-inoculated YFP-Sw-5b transgenic *N. benthamiana* plants were photographed at 15 dpi. White arrowhead indicates HR trailing in systemic leaves.

Fig. 9. A working model for Sw-5b. Sw-5b furcates disease resistances by proper nucleocytoplasmic partition to block different infection steps of tomato spotted wilt tospovirus. Sw-5b switched from the autoinhibited state to an activated state upon recognition of NSm in the cytoplasm. Cytoplasm portion of Sw-5b induce cell death and defense that inhibit viral replication. The activated Sw-5b also translocated into nucleus via *importins a* and *b* . Nucleus-localized Sw-5b induces a defense that block viral cell-to-cell and long-distance movement. Cytoplasm- and nucleus-localized Sw-5b have additively effects on defense to inhibit viral replication, intercellular and long-distance movement during tospovirus infection.

Fig. S1 Sw-5b recognizes TSWV NSm in cytoplasm. (a) Transient expressions of NSm-YFP-NES, NSm-YFP-nes, NSm-YFP-NLS, and NSm-YFP-nls, respectively, in *N. benthamiana* leaves through agro-infiltration. Epidermal cells expressing various fusion proteins were imaged under a confocal microscope at 24 hpai. The numbers in each image indicate the number of cells showing this subcellular localization pattern and the total number of cells examined per treatment. N, nucleus; C, cytoplasm. Bar = 10 μm. (b) Various fusion proteins described in (a) were, individually, co-expressed with Sw-5b in *N. benthamiana* leaves. A representative leaf was photographed at 5 dpai. (c) Western blot analysis of various NSm fusion protein expressions in the assayed *N. benthamiana* leaves using a YFP specific antibody. Leaf areas co-expressing Sw-5b and EV were used as negative controls. The Ponceau S stained Rubisco large subunit gel was used to show sample loadings.

Fig. S2 Analysis of virus replication monitoring system using a TSWV-based mini-genome replicon and a movement defective NSm mutant. (a) SR$_{(+)eGFP}$, L$_{(+)opt}$, VSRs and NSm or NSm$^{H93A\&H94A}$ mutant were transiently co-expressed in *N. benthamiana* leaves through agro-infiltration. The infiltrated leaves were examined and imaged under a confocal microscope at 4 dpai. The numbers in each image indicate the number of cells showing similar expression pattern and the total number of cells examined per treatment. Bar = 400 μm. (b) Western blot analysis of eGFP accumulation in assayed leaves using a YFP specific antibody. Leaves co-expressing SR$_{(+)eGFP}$, L$_{(+)opt}$, VSRs and EV were used as negative controls. The Ponceau S stained Rubisco large subunit gel was used to show sample loadings.

(a)

mCherry-HDEL//NSm^T120N^-GFP

(b)

mCherry-HDEL//NSm-GFP

Fig. S3 Effects of nes-Sw-5b and nls-Sw-5b on NSm-GFP cell-to-cell movement and effects of Sw-5b and EV on NSm^{T120N}-GFP cell-to-cell movement. (a) Sw-5b and EV were, respectively, co-expressed with mCherry-HDEL//NSm^{T120N}-GFP in *N. benthamiana* leaves through agro-infiltration. (b) nes-Sw-5b and nls-Sw-5b were, respectively, co-expressed with mCherry-HDEL//NSm-GFP in *N. benthamiana* leaves through agro-infiltration. The Agrobacterium culture carrying pmCherry-HDEL//NSm-GFP or pmCherry-HDEL//NSm^{T120N}-GFP was first adjusted to OD$_{600}$ = 0.2 and then further diluted 500 times prior to use. All other Agrobacterium cultures were adjusted to OD$_{600}$ = 0.2 prior to use. The numbers in each image indicate the number of cells showing similar expression pattern and the total number of cells examined per treatment. Bar = 50 μm.

Fig. S4 Effects of cytoplasmic and nuclear Sw-5b on host immunity to TSWV systemic infection. (a) Transgenic *N. benthamiana* lines expressing NES-YFP-Sw-5b, nes-YFP-Sw-5b, NLS-YFP-Sw-5b, nls-YFP-Sw-5b or YFP-Sw-5b, driven by the Sw-5b promoter, were used in this study. The EV transgenic plants were used as controls. The transgenic plants were inoculated with TSWV and photographed at 15 dpi. White arrow indicate the systemic leaves showing HR trailing. White arrowhead indicates the systemic leaves showing mosaic symptoms. (b) RT-PCR detection of TSWV infection in the systemic leaves of the assayed *N. benthamiana* plants at 15 dpi.

Fig. S5 Cytoplasmic and nuclear Sw-5b activity on TSWV-GFP cell-to-cell movement in *N. benthamiana* leaves. (a) $pL_{(+)opt}$ and $pSR_{(+)eGFP}$-$M_{(-)opt}$ were co-inoculated with TSWV-GFP into *Nicotiana benthamiana* leaves through agro-infiltration. The inoculated leaves were examined and imaged under a confocal microscope at 4 dpai. Bar = 400 μm. (b) Statistic analysis of TSWV-GFP cell-to-cell movement in the assayed *N. benthamiana* leaves from Figure 7B. A total of 9 assayed leaves were used for each treatment.

Fig. S6 An immunoblot showing the accumulations of various YFP-tagged proteins expressed in different transgenic *N. benthamiana* plants and RT-PCR analysis of TSWV accumulation in the systemic leaves. (a) The fusion proteins were detected using an YFP specific antibody. Arrows indicate the positions of the expressed fusion proteins. Ponceau S stained Rubisco large subunits were used to estimate sample loadings. (b) RT-PCR analysis of TSWV accumulation in the systemic leaves of different transgenic *N. benthamiana* plants at 15 dpi.

Fig. S7 Bimolecular fluorescence complementation (BiFC) assay of cYFP-SD, cYFP-Sw-5b and nYFP-Importin α1 (nYFP-IMP α1), nYFP-Importin α2 (nYFP-IMP α2), nYFP-Importin β (nYFP-IMP β) interaction in N. benthamiana leaf epidermal cells. Confocal images were taken at 26 hour post agro-infiltration. Leaf epidermal cells co-expressing cYFP-SD and nYFP, cYFP-Sw-5b and nYFP or cYFP and nYFP-IMP α1, nYFP-IMP α2, nYFP-IMP β were used as controls. Bar = 50 μm.

(a)

(b)

Fig. S8 RT-PCR analyses of *importin* α*1*, α*2* and *β* expressions in the assayed plants and their effects on TSWV systemic infection. (a) Expressions of *importin* α*1,* α*2,* and *β* in Sw-5b transgenic *N. benthamiana* plants were silenced individually or together using a TRV-based VIGS vector. The gene silencing results were determined through semi-quantitative RT-PCR using gene specific primers. PCR products obtained after 25 cycles of PCR reaction were visualized in 1% agrose gel through electrophoresis. (b) RT-PCR detection of TSWV systemic infection in the assayed plants. The resulting PCR products were visualized in 1% agrose gel through electrophoresis.