**Host-generalist fungal pathogens of seedlings may maintain forest diversity via host-**

**specific impacts and differential susceptibility among tree species**

Erin R. Spear[1][*] and Kirk Broders[1]

1. Smithsonian Tropical Research Institute, Panama City, Panama, Republic of

Panama.

[*]Corresponding author: erin.r.spear@gmail.com

**Summary**

• Host-specialized pathogens are credited with the maintenance of tropical forest

diversity under the Janzen-Connell hypothesis (JCH). Yet, in diverse forests,

selection should favor pathogens with broad host ranges given their passive

dispersal and the relative rarity of tree species.

• We surveyed the host associations of potential pathogens isolated from

symptomatic seedlings in the forests in Panama; and used inoculations to assess

(i) the pathogenicity and host ranges of 27 fungal isolates, and (ii) differences

among tree species in susceptibility.

• Thirty-one of the 33 non-singleton OTUs isolated from seedlings are multi-host.

All 31 multi-host OTUs exhibit low to moderate specialization and

phylogenetically dispersed host use, with no phylogenetic signal. The

pathogenicity of 10 isolates was experimentally confirmed; nine caused disease in

seedlings in multiple families. However, the outcome of infection differs among

tree species susceptible to a given multi-host pathogen. Furthermore, some tree

species were seemingly resistant to all fungi tested, while others were susceptible

to multiple fungi. Tree species adapted to environments with lower disease

pressure were most likely to exhibit disease symptoms.

• Our results suggest that generalist pathogens contribute to the maintenance of

local forest diversity via host-specific impacts rather than the host specificity

originally envisioned under JCH.

**Key words:** *Clonostachys*, host generalists, host-specific impacts, multi-host pathogens,

*Mycoleptodiscus*, Nectriaceae, seedlings, tropical forests

## Introduction

The maintenance of local diversity is commonly attributed to host-specific natural enemies believed to prevent any single host species from becoming competitively dominant. Under the Janzen-Connell hypothesis (JCH), conspecific seeds and seedlings at high densities near conspecific adults suffer disproportionately higher mortality relative to heterospecific seeds and seedlings due to an accumulation of host-specialized enemies (Gillett, 1962; Janzen, 1970; Connell, 1971; Comita *et al*., 2014). However, support for the JCH has often been based on spatial and temporal patterns of conspecific negative density-dependent mortality (e.g., Harms *et al*., 2000; Comita *et al*., 2010), which cannot identify the mechanism(s) driving the observed patterns.

Experimental studies suggest that pathogens may contribute to density- and distance-dependent mortality of conspecific seedlings (Augspurger & Kelly, 1984; Bell *et al*., 2006; Mangan *et al*., 2010; Bagchi *et al*., 2014). However, the pathogens are rarely identified, and most studies fail to experimentally address a crucial assumption of the JCH, that the pathogens exhibit high host specificity with regard to locally available hosts (but see Packer & Clay, 2000; Y. Liu *et al*., 2012). Furthermore, the relative rarity of tree species in diverse tropical forests and the passive dispersal of fungi may select for pathogens with intermediate to broad host ranges (May, 1991; Coley & Barone, 1996). Indeed, mounting evidence from tropical and temperate forests suggests that multi-host pathogens are prevalent (Augspurger & Wilkinson, 2007; Gallery *et al*., 2007; Gilbert & Webb, 2007; Kluger *et al*., 2008; Hersh *et al*., 2012; Schweizer *et al*., 2013; Sarmiento *et al*., 2017; Chen *et al*., 2019).

Pathogens are classified as specialists or generalists based on the number of host species and the phylogenetic relatedness among hosts (Barrett *et al*., 2009). Host-specialized multi-host pathogens are predicted to exhibit a phylogenetic signal to their host range (Barrett *et al*., 2009). A variety of plant traits influence the outcome of plant-pathogen interactions, specifically susceptibility to infection, and resistance and tolerance post-infection (Coley & Barone, 1996; Bradley *et al*., 2003; Barrett *et al*., 2009). Plant defense traits are often, but not always, phylogenetically conserved (Fine *et al*., 2006; Agrawal, 2007; Kursar *et al.*, 2009). Under the assumption that closely related plants are more likely to possess similar defense traits, the likelihood of tree species sharing a

pathogen should decrease with decreasing phylogenetic relatedness (Barrett *et al.*., 2009).

Pathogens specialized at the clade-level could still maintain plant community diversity,

but at higher taxonomic levels (Parker *et al.*, 2015).

While specialist pathogens are invoked under JCH, generalist pathogens may

contribute to the maintenance of host community diversity via effective specialization

(Benítez *et al.*, 2013). One form is host-specific impacts (Sarmiento *et al.*, 2017);

specifically, disease severity differs among the hosts of a given pathogen. Host-specific

impacts could serve as an equalizing mechanism that maintains plant community

diversity if the growth and survival of infected seedlings varies among host species

(Barrett *et al.*, 2009); particularly, if differential impacts reflect ecological trade-offs

between allocation of resources to defense, competitive ability, and/or tolerance of

abiotic stressors. For example, for tree species adapted to environments with relatively

high and/or consistent disease pressure, the fitness benefits of physical and chemical

resistance and/or tolerance traits may outweigh the associated allocation costs of those

traits, while the costs may outweigh the benefits for tree species adapted to environments

with relatively low and/or inconsistent disease pressure (Haak *et al.*, 2012). The risk of

pathogen attack, referred to here as pathogen pressure, varies with environmental

conditions that impact pathogen and plant fitness (Barrett *et al.*, 2009). High pathogen

pressure has been observed in areas with low light, high soil moisture, and persistent fog

and dew (Augspurger & Kelly, 1984; Bradley *et al.*, 2003; Mordecai, 2012; Spear *et al.*,

2015). Plants adapted to environments with high and/or consistent pathogen pressure may

be under selection for increased defenses, whereas plants adapted to low and/or

inconsistent pathogen pressure environments may be poorly defended (Coley & Barone,

1996, Talley *et al.*, 2002).

Here we describe (i) pathogens infecting tree seedlings in the forests of Panama,

(ii) their host ranges, (iii) interspecific variability among tree species in pathogen

susceptibility, and (iv) the relationship between disease susceptibility and plant habitat

associations. We had four central hypotheses. First, in these diverse tropical forests,

seedling pathogens tend to be multi-host. Second, there is a phylogenetic signal to the

host ranges of these multi-host pathogens (i.e., pathogen sharing is positively correlated

with host relatedness). Third, tree species differ in their susceptibility to pathogens

because the cost-to-benefit ratio of defense traits differs among species (Strauss &

Agrawal 1999; Endara & Coley, 2010). Fourth, plants adapted to environments with

relatively low and/or inconsistent disease pressure are more susceptible to disease than

those adapted to relatively high and/or consistent disease pressure.

**Materials and Methods**

*Isolation and identification of putative pathogens*

Potentially pathogenic fungi and oomycetes were isolated from 124 seedlings,

representing 26 tree species in 13 families, with disease on the leaf, stem, and/or root

(Supporting Information Fig. **S1**; Table **S1**). Symptomatic seedlings were collected from

seven forest sites in 2007, 2010-2012, and 2019 by: (1) opportunistic collection of

naturally occurring seedlings ($n = 59$); (2) collection of seedlings germinated in a

shadehouse and then transplanted to forest sites ($n = 22$), and (3) collection of seedlings

that germinated from surface-sterilized seeds planted directly in the forests ($n = 43$)

(Table **S2**). Most putative pathogens were isolated from leaves (52.13%), followed by

stems (37.44%) and roots (10.43%). Molecular identification was completed by

bidirectional Sanger sequencing of either the nuclear ribosomal Internal Transcribed

Spacer (ITS) region (149 isolates) or the ITS plus an adjacent portion of the large subunit

(ITS+LSU) (62 isolates). See Table **S2** and Spear (2017) for details.

All 211 unique isolates were identified and assigned to operational taxonomic

units (OTUs) based on 99% sequence similarity (minimum overlap of 40 bp) in

Sequencher v. 5.4.6. We used a stringent similarity threshold that splits rather than lumps,

while accounting for ≤1% sequencing error, because we make statements about host

specificity and to avoid masking strain-specific effects (Higgins *et al*., 2011). We

estimated the taxonomic placement of the fungal isolates by querying all 209 edited

sequences against the well-curated Full UNITE+INSD dataset for Fungi (v. 8.2, released

2020-02-04; Abarenkov *et al*., 2020) (detailed in Methods **S1**). For the two oomycetes,

we queried their sequences against GenBank (accessed 8-Oct-2020; Benson *et al*., 2012)

and the curated database Phytopthora-ID (v. 2.0, accessed 8-Oct-2020; Grünwald *et al*.,

2011). All assigned nomenclature should be interpreted as an estimate as: (i) the ITS

region is an imprecise barcode for species discrimination for certain taxa; (ii) multi-locus

phylogenies are required for accuracy; and (iii) precision is limited by incomplete

taxonomic and geographic coverage of databases (Kang *et al*., 2010; Hofstetter *et al*.,

2019; Lücking *et al*., 2020). Additionally, sequences in the UNITE database are clustered

into hypothesized species-level groups (Species Hypotheses, SHs) based on sequence

similarity thresholds ranging from 97–100% (Kõljalg *et al*., 2013; Robbertse *et al*.,

2017), while we used 99% sequence similarity to designate OTUs because 97-98.5% is

too relaxed for species delimitation for some taxa (Lücking *et al*., 2020). Because our

grouping strategy often differed from that of the UNITE database, multiple, distinct

OTUs in our study sometimes share the same nomenclature based on the SH of the best-

matching UNITE reference sequence (e.g., OTUs Z and AAG are both annotated as

*Colletotrichum brevisporum*). Finally, for certain groups of fungi, UNITE SHs may

erroneously include distinct species (Robbertse *et al*., 2017).

Sequences for the 90 previously published fungal isolates (Spear, 2017) were

submitted to the NCBI GenBank database (KY413686–KY413775). Sequence data for

the 121 isolates collected in 2019 will be submitted to GenBank (XXXXXXX–

XXXXXXX).

To characterize sampling efficacy and the relative abundance of OTUs, we

calculated the richness of observed OTUs and plotted the taxon accumulation curve

(vegan package; Oksanen *et al*., 2019) and rank-abundance distribution (BiodiversityR

package; Kindt & Coe, 2005) ($n$ = 211 isolates from 124 seedlings of 26 tree species).

We defined OTUs observed ≥10 times as abundant.

*Survey-based assessment of host associations and ranges of putative pathogens*

Our survey-based assessment of the host associations and ranges of putative

pathogens utilize our full dataset ($n$ = 211 isolates, representing 66 OTUs, from 26 tree

species) and three subsets to ameliorate the influence of undersampled taxa. Subset A

excludes singleton OTUs ($n$ = 178 isolates, representing 33 OTUs, from 24 tree species),

B excludes singleton and doubleton OTUs and tree species ($n$ = 144 isolates, representing

22 OTUs, from 13 tree species), and C excludes singleton OTUs and the two non-

singleton OTUs only isolated from one host species ($n$ = 174 isolates, representing 31

OTUs, from 24 tree species). All four datasets include isolates collected in 2007, 2010-

2012, and 2019.

We used four approaches to characterize host associations. We (1) summed and

plotted the number of observed host tree species (unweighted links) (full dataset: $n = 211$

isolates), and (2) calculated Blüthgen's weighted specialization index d' for each of the

33 non-singleton OTUs (subset A: $n = 178$ isolates; bipartite package; Dormann *et al*.,

2008; Blüthgen *et al*., 2006). The d′ index is useful for identifying discriminatory host

use because it accounts for interaction strength (i.e., strong, frequent vs. weak, occasional

interactions) and sampling intensity, while the number of interacting partners does neither

(Dormann, 2011; Blüthgen *et al*., 2006). The index estimates specialization by

calculating the degree to which observed host use differs from expected host use given

partner availability (Blüthgen *et al*., 2006; Dormann, 2011). We could not adjust for the

relative abundances of tree species when calculating d′ as the seven forest sites span a

marked gradient in plant diversity, and tree species are not uniformly distributed spatially

(Pyke *et al*., 2001). Thus, we relied on the interaction matrix's marginal sums (column-

and row-wise totals; total interactions per species) as a measure of partner availability,

meaning that, for our data, a generalized microbe infects tree species proportional to their

sampling intensities, while a specialized microbe infects tree species independent of their

sampling intensities (Blüthgen *et al*., 2006). We considered d' values ranging from 0–

0.33, 0.34–0.67, and 0.68–1.00 to represent low, moderate, and high specialization,

respectively (Blüthgen *et al*., 2006; Spear & Mordecai, 2018). We (3) calculated the

observed number of shared OTUs (rich package; Rossi, 2011) and, because that metric

generally results in an underestimation of similarity due to undetected species, we (4)

estimated similarities in the communities of putative pathogens shared between pairs of

tree species via the inverse of the Chao dissimilarity index (vegan package; Oksanen *et*

*al*., 2019) (full dataset: $n = 211$ isolates; subset B: $n = 144$ isolates). The Chao index

accounts for the relative abundances of observed OTUs and unseen shared OTUs, making

it relatively robust to unequal sampling and diverse guilds with many rare taxa (Chao *et*

*al*., 2005).

To determine if there is a phylogenetic signal to the host ranges of the OTUs, (i)

we used the mean pairwise phylogenetic distance (MPD) to determine if a given multi-

host OTU infects phylogenetically clustered or dispersed hosts, and (ii) we used two

binomial generalized linear models (GLMs) (logit link) to determine whether putative

pathogen community similarity is positively correlated with host relatedness.

Specifically, for the 31 OTUs isolated from ≥2 tree species (subset C: $n = 174$ isolates),

we compared the observed MPD between host tree species infected by a given OTU to

the MPD generated in a null model with random host-OTU associations (999

permutations generated via randomization of the host phylogeny tips; picante and ape

packages; Kembel *et al*., 2010; Paradis & Schliep, 2018). For the GLMs, we used our full

dataset ($n = 211$ isolates) and subset B ($n = 144$ isolates), and we regressed and plotted

the pairwise estimated community similarities (1 - Chao index) against the pairwise

phylogenetic distances. The phylogeny of tree species was generated by Phylomatic

(phylodiversity.net/phylomatic/), which called the supertree published by Zanne *et al*.

(2014).

We tested for correlations between (i) the frequency with which a given fungal

OTU was isolated and the number of tree species from which it was isolated, and (ii)

sampling intensity (number of seedlings collected) and the number of OTUs isolated

from a given tree species using one-tailed, non-parametric Spearman's rank tests (full

dataset: $n = 211$ isolates; stats package; R Core Team, 2020). One-tailed tests were used

because we had directional, literature-based *a priori* hypotheses (Quinn & Keough, 2002;

Ruxton & Neuhäuser, 2010) that: (i) the most abundant fungal taxa would be multi-host,

and (ii) observed fungal richness associated with a given plant species would increase

with increased sampling (Ferrer & Gilbert, 2003; Spear & Mordecai, 2018).

*Experimental assessments of pathogenicity and host range*

In 2011 and 2012, inoculation experiments were used to assess the pathogenicity

and host specificities of 27 fungal isolates collected in 2010 and 2011 (Fig. **S2a**). We

tested the 27 isolates, representing 18 OTUs, against seedlings of 35 tree species,

representing 20 families (Table **S1**). The tree species-by-fungal isolate combinations

were not fully reciprocal (285 of 945 possible combinations; 3,154 seedlings total) due to

space and time constraints and because seeds were unavailable or available in insufficient

quantities. A given fungal isolate was tested against three to 28 tree species and 48% of

isolates evaluated were tested against seedlings of ≥10 tree species. Because an objective

of the inoculation experiments was to describe pathogen host range, once isolates

displayed pathogenicity, we prioritized testing those isolates against additional tree

species (12-28 tree species, median 15) over expanding the testing of isolates failing to
generate significant disease. A given tree species was tested against one to 26 fungal
isolates and 29% of tree species were tested against ≥10 isolates.

Our inoculation technique simulated infested plant material in the soil to passively
inoculate the seedlings with the fungal isolates selected for screening (modified from
Augspurger & Wilkinson, 2007). To generate seedlings for the inoculation experiments,
seeds were collected from forests bordering the Panama Canal (May-Nov. of 2011, May-
June 2012), surface-sterilized (sequential washes of 95% EtOH for 10 s, 10% commercial
bleach for 2 min, 70% EtOH for 2 min), and germinated in flats of autoclave-sterilized (1
hour at 121º C, twice) commercial soil (Fig. **S2b**). Seedlings were transplanted to
individual pots containing autoclaved commercial soil and either rice visibly colonized
by one of the fungal isolates or inoculum-free, autoclave-sterilized rice (control/sham
treatment) (Fig. **S2c-f**). Depending on seed availability, five to 12 seedlings per species
were included in each treatment.

Inoculation experiments were conducted in Smithsonian Tropical Research
Institute shadehouses in Gamboa, Panama during the wet seasons of 2011 and 2012 (Fig.
**S2a**). The two screened-in shadehouses were covered on all sides with shadecloth and
their roofs were covered with plastic to minimize external contamination and to mimic
the forest understory. Shadehouse air temperatures (26.1 and 25.6ºC) did not significantly
differ from those recorded simultaneously in a nearby forest understory (25.4 and 25.3ºC;
Paton, 2019a). Relative humidity was significantly lower in the shadehouses (85.1 and
87.6%) than in the forest understory (95.6 and 96.7%; Paton, 2019b). Photosynthetically
active radiation (PAR) reaching the shadehouse seedlings was slightly higher (1.4 and
1.7% of full PAR) than observed in a nearby forest understory (1.3%; Brenes-Arguedas
*et al*., 2011). See Table **S3** for details. Light was somewhat variable within a given
shadehouse. To counteract this potentially confounding factor, seedlings were rotated
among the tables weekly.

Disease symptoms were documented every three days, for up to five weeks, and
were categorized as: (i) seedling mortality; (ii) stem damage (Fig. **S2g-o**); (iii) wilted
tissue (Fig. **S2p,q**); and (iv) stunted seedling growth (Fig. **S2r,s**). A seedling was
classified as morbid if it had any of the four disease symptoms. Wilted tissue was

considered a symptom of disease because the soil was not allowed to dry out (seedlings
were hand-watered every three days) and pathogens attacking roots can interfere with
water uptake (Yadeta & Thomma, 2013). Similarly, stunted growth can be indicative of
pathogen-caused damage to roots and/or the vascular system (Yadeta & Thomma, 2013).
Additionally, stunting may occur because energy and resources are allocated to defenses
and/or tissue replacement rather than growth (Huot *et al*., 2014). Root and foliar damage
could not be reliably observed and were not included in any analyses. Regarding the
latter, we observed a synchronous occurrence of morphologically similar foliar damage
across distinct treatments and the controls, which suggested that causative biotic agent
(*Xylaria* sp.; E. R. Spear, unpublished data) may have blown into the shadehouses.
To compare the proportion of symptomatic seedlings in the inoculated versus
inoculum-free sham (control) treatments, we used generalized linear models (GLMs)
with binomial error distributions, probit link functions, and mean bias-reducing score
adjustments (brglm package; Firth, 1993; Kosmidis & Firth, 2009; Kosmidis, 2019). We
used bias-reduced GLMs because we observed complete and quasi-complete separation
of data for some fungal isolate-by-tree species combinations (i.e., treatment perfectly, or
near perfectly, predicted the binary outcome; Albert & Anderson, 1984). The response
variables were presence/absence of (i) general morbidity and (ii) mortality. Each tree
species-by-fungal isolate combination was considered individually for each response
variable. We considered an isolate pathogenic if, for at least one tree species, a
significantly greater proportion of inoculated seedlings were symptomatic than paired
control seedlings (using $\alpha = 0.10$ because small sample sizes limited our power to detect
treatment effects; Quinn & Keough, 2002; Thiese *et al*., 2016). We tested for a
correlation between the frequency with which a fungal OTU was isolated and the number
of tree species in which it caused statistically significant disease using a non-parametric
Spearman's rank test. We used a one-tailed test because we had the *a priori* hypothesis
that the most abundant fungal taxa would be multi-host.
*Plant habitats, seed sizes, and disease susceptibility*
We used logit-link beta-binomial generalized linear models to explore disease
susceptibility as a function of habitat associations and seed size (aod package; Lesnoff &
Lancelot, 2012). We obtained data for seed size, shade tolerance, and spatial distribution

relative to annual rainfall from published sources (Table **S1**). In all models, the response

variable was presence/absence of morbidity in inoculated seedlings, controlling for the

morbidity observed in paired control seedlings and considering all 27 isolates tested

(pathogens and non-pathogens alike). Because negative relative proportions are

biologically equivalent to zero (no disease caused), and for statistical purposes, we

converted all negative values to zero. We used two global models that contained (i) the

additive main effects of seed size (continuous) and shade tolerance (categorical), and

their interaction, or (ii) the additive main effects of seed size and distribution relative to

rainfall (categorical), and their interaction. We excluded seeds with average dry masses

greater than 3,000 mg to improve model stability. For both categorical predictors, we

limited our comparisons to binary extremes (i.e., only shade-intolerant and -tolerant tree

species, and only wet- and dry-site tree species). We analyzed disease as a function of

shade tolerance (26 tree species, 179 observations: 42 for intolerant spp., 137 for tolerant

spp.) and spatial distribution (15 tree species, 154 observations: 108 for dry-site spp., 46

for wet-site spp.) separately due to differences in the availability of trait data. For the two

sets of models, automated model selection and averaging was done based on the global

models, the Akaike information criterion with a correction for small sample sizes (AICc),

and $\Delta$AICc $\leq 2$ (MuMIn package, Bartoń, 2020).

All data were processed in R v. 4.0.0 (R Core Team, 2020). In addition to the

packages used for analyses, we used these for data formatting and figures: fossil (Vavrek,

2011), ggplot2 (Wickham, 2016), otuSummary (Yang, 2020), plot.matrix (Klinke, 2019),

plyr (Wickham, 2011), reshape2 (Wickham, 2007).

**Results**

*Relative abundances and taxonomic assignments of putative pathogens*

The 211 isolates represent 66 operational taxonomic units (OTUs); however, the

taxon accumulation curve is non-asymptotic, indicating incomplete sampling (Fig. **S3b**).

Most of the OTUs are relatively rare, 33 are singletons (Figures **1**, **S3a**; Table **S4**). Four

OTUs are common (observed $\geq$10) and comprised 35% of all isolates: A–*Colletotrichum*

*xanthorrhoeae*, B–Nectriaceae, C–*Mycoleptodiscus suttonii*, and F–*Diaporthe tulliensis*

(Figures **1**, **S3a**; Table **S4**). The 209 fungal and two oomycete isolates represent five

classes (Sordariomycetes, Dothideomycetes, Agaricomycetes, Eurotiomycetes, and

Oomycetes) and 11 orders. Most (81.5%) of the isolates could be assigned to 26 genera.

*Colletotrichum*, *Diaporthe,* and *Mycoleptodiscus* were the three genera most frequently

isolated (Fig. **1**; Table **S4**). We observed overlap in fungal OTUs among the three tissues

sampled: leaves, stems, and roots (Fig. **S4d**). While sampling occurred over five years,

there was overlap in the tree species and forest sites sampled and the isolation and

molecular techniques used (Table **S2**). Accordingly, we observed overlap in fungal OTUs

among the years, seedling collection methods, and isolation media (Fig. **S4a-c**).

*Fungi associated with symptomatic seedlings are host generalists*

The fungi isolated from symptomatic seedlings are predominately multi-host

generalists with wide host ranges. Of the 33 non-singleton OTUs, 31 were isolated from

2-9 tree species (median = 3) (Fig. **1**; Table **S4**). The number of tree species from which a

fungal OTU was isolated is positively correlated with isolation frequency (one-tailed

Spearman's rank correlation: rho = 0.963, $P$ < 0.001; blue points Fig. **S5**). A–*C.*

*xanthorrhoeae* was isolated with the greatest frequency (35 times) and from the greatest

number of tree species (9 spp. in 7 families) (Fig. **1**; Table **S4**). Only two of the non-

singleton OTUs (T–*Macrophomina* and X–*Ceratobasidium*) were not multi-host, but

each was isolated only twice, and from the best-sampled tree species, suggesting their

observed host ranges may be a sampling artifact (Fig. **1**; Tables **S1, S4**). Based on the

specialization index d', 57.6% of 33 the non-singleton OTUs, including A–*C.*

*xanthorrhoeae* and B–Nectriaceae, exhibit low host-specialization (d' ≤ 0.33) and 42.4%,

including C–*M. suttonii* and F–*D. tulliensis*, exhibit moderate host-specialization (d'

0.33–0.67) at the species level (Table **S4**). Twelve of the 31 multi-host OTUs infect hosts

that are more closely related than expected by chance, but none significantly so (Table

**S4**). For two of those OTUs (Q–*Diaporthe cf. columnaris* and R–*Diaporthe fraxini-*

*angustifoliae*), phylogenetic clustering of host use was marginally significant ($P$ = 0.09;

Fig. **1**; Table **S4**), which is tentative evidence of a phylogenetic signal to host range. Both

OTUs were solely isolated from *Anacardium excelsum* (Anacardiaceae, Sapindales) and

*Protium panamense* (Burseraceae, Sapindales) seedlings.

*Tree species are infected by various microbes, and their microbial communities overlap*

Multiple OTUs were isolated for all tree species with >2 seedlings collected (Fig.

**1**; Table **S1**). Predictably, the number of OTUs isolated from a given tree species is

strongly positively correlated with sampling intensity (one-tailed Spearman's rank

correlation: rho = 0.951, *P* < 0.001; Fig. **1**). The greatest richness of fungi (24 OTUs) was

recovered from the best-sampled tree species (41 isolates), *A. excelsum* (Table **S1**).

We observed overlap among the microbial communities of distantly related tree

species (Fig. **1**). Based on subset B, excluding singleton and doubleton observations of

tree species and OTUs, the estimated average similarity (1 - Chao index) between the

microbial communities of pairs of tree species is 0.21 (median = 0.15; Table **S5**). The

GLMs suggest that the similarity of microbial communities is unrelated to phylogenetic

distance between tree species (Fig. **2**; full dataset: Est. ± SE = 0.002 ± 0.003, *z*-value =

0.60, *P* = 0.55; subset B: Est. ± SE = 0.002 ± 0.004, *z*-value = 0.36, *P* = 0.72).

*Identities and impacts of fungi with confirmed pathogenicity*

We experimentally confirmed the pathogenicity of 10 of the 27 fungal isolates

tested (Fig. **3**). The isolates with confirmed pathogenicity are assigned to six OTUs, all in

the phylum Ascomycota, and were originally isolated from stems, roots, and leaves of

seedlings of seven tree species in seven families (Fig. **3**; Table **1**). Based on morbidity

(i.e., any of the four specific disease symptoms: wilt, stunting, stem damage, or

mortality), significant pathogenicity was observed for 20 of the 285 tree species-by-

fungal isolate combinations (Fig. **3**; Table **1**). For 13 of these combinations, infection led

to seedling mortality (Table **1**). Multiple disease symptoms were often observed for a

single tree species-by-pathogen combination (e.g., *Annona glabra* seedlings suffered

stem damage, wilt, and mortality when inoculated with isolate ERS029).

*Fungi with confirmed pathogenicity are multi-host*

We show that fungal pathogens can attack tree species belonging to

phylogenetically dispersed families. Five of the 10 pathogenic isolates successfully

attacked multiple tree species (Table **1**). The 10 isolates with confirmed pathogenicity

generated significant disease symptoms in 6-31% of the tree species against which they

were tested (median = 10%). At the high end, ERS029 (C–*Mycoleptodiscus suttonii*)

caused significant disease in 31% of the 13 tree species it was tested against and the four

susceptible tree species belong to three orders (Fig. **3**; Tables **1**, **S1**).

Three of the OTUs were represented by multiple isolates: B–Nectriaceae, G–

*Clonostachys rosea*, and C–*M. suttonii*. The two isolates of B–Nectriaceae were

recovered from *Trichilia tuberculata* and *Calophyllum longifolium*, and combined were pathogenic on six of the 28 tree species tested (Fig. **3**; Table **1**). The three isolates of G–*C. rosea* were isolated from *Virola surinamensis* and *Protium tenuifolium*, and combined caused significant disease on four tree species (Fig. **3**; Table **1**). The two isolates of C–*M. suttonii* were isolated from *V. surinamensis* and *Randia armata*, and combined caused significant disease on five species (Fig. **3**; Table **1**). While the tree species-by-fungal isolate combinations were not fully reciprocal, limiting conclusions, we observed possible strain-specific effects (e.g., B–Nectriaceae: ERS034 caused significant disease on seedlings of *Guapira standleyana* while ERS036 did not; Fig. **3**). Finally, there is a marginally significant positive association between the frequency with which a fungal OTU was isolated and the number of tree species in which it caused statistically significant ($\alpha = 0.10$) disease (one-tailed Spearman's rank correlation rho = 0.721, $P = 0.053$; green points Fig. **S5**).

*Tree species are differentially impacted by pathogens*

We documented interspecific differences in disease susceptibility and impact among tree species. Most tree species tested were seemingly resistant to, or tolerant of, all fungal isolates tested, suffering minimal to no fitness impacts despite soil inoculation (Fig. **3**). For 13 of the 35 tree species tested, at least one of the 27 fungal isolates tested caused significant disease (Table **1**). Five tree species experienced significant disease from two to three of the isolates tested (Table **1**). Those five tree species experienced significant disease from 10-50% (median = 15%) of the fungal isolates against which they were tested (Fig. **3**).

The 13 susceptible tree species differed in their tolerance of attack. For most of the susceptible tree species, pathogen attack did not consistently lead to death, the impact from infection differed depending on the pathogenic isolate with which it was inoculated. For example, while *Castilla elastica* seedlings suffered significant disease when inoculated with two fungal isolates (ERS036 and ERS040), they only suffered significant seedling mortality when inoculated with ERS040 (AAF–Nectriaceae) (Table **1**). Furthermore, for those tree species that were susceptible to the same multi-host pathogen, the impact from attack differed among species. For example, isolate ERS036 (B–Nectriaceae) caused significant morbidity on seedlings of *C. elastica, Cochlospermum*

*vitifolium,* and *Dalbergia retusa*, but only caused significant mortality on *C. vitifolium*

seedlings (Table **1**).

*Plant habitats, seed sizes, and disease susceptibility*

When considering all 27 fungal isolates tested (pathogens and non-pathogens), the

proportion of seedlings with morbidity was influenced by a tree species' seed size, spatial

distribution relative to annual rainfall, and shade tolerance (Fig. **4a-d**; Tables **S6**, **S7**).

Shade-intolerant tree species were more likely to exhibit disease than shade-tolerant

species ($P = 0.003$; Fig. **4a,b**; Table **S6**). For both groups, the probability of disease

decreases with increasing seed size, but to a greater extent for shade-intolerant species ($P$

$= 0.014$; Fig. **4a,b**; Table **S6**). Tree species restricted to, or more abundant in, drier sites

(dry-site species) were more likely to exhibit disease than those restricted to, or more

abundant in, wetter sites (wet-site species) ($P = 0.042$; Fig. **4c,d**; Table **S7**). While we

found no correlation between seed size and disease risk for wet-site tree species, disease

risk decreases with increasing seed size for dry-site tree species (Fig. **4c,d**; Table **S7**).

**Discussion**

Pathogens are believed to shape the distributions and abundances of their hosts,

effects often ascribed to host-specialized pathogens (Comita *et al*., 2014). However, there

is increasing evidence that multi-host pathogens are prevalent, in natural plant

communities (Augspurger & Wilkinson, 2007; Gallery *et al*., 2007; Gilbert & Webb,

2007; Kluger *et al*., 2008; Hersh *et al*., 2012; Schweizer *et al*., 2013; Sarmiento *et al*.,

2017; Spear & Mordecai, 2018). These multi-host pathogens may contribute to the

maintenance of these communities via host-specific impacts (disease severity differs

among a pathogen's hosts), a form of effective specialization (Augspurger & Wilkinson

2007; Benítez *et al*., 2013; Sarmiento *et al*., 2017). Using a multifaceted approach, we

identified fungal pathogens attacking seedlings in the forests of Panama, and show that

they can attack phylogenetically dispersed hosts. Furthermore, we demonstrate that tree

species differ in their susceptibility to pathogens, with trees adapted to environments with

relatively low and/or inconsistent disease pressure being more susceptible to disease.

Collectively, these results suggest that generalist pathogens contribute to the maintenance

of local and regional plant diversity.

*Pathogens and their impacts*

We observed 66 OTUs, but only four were observed more than 10 times (Figures **1, S3a**; Table **S4**). The relative rarity of the OTUs, coupled with a non-asymptomatic taxon accumulation curve, indicate incomplete sampling and a diverse community of seedling-associated fungi (Fig. **S3**). The most frequently isolated fungal genera (Fig. **1**) and those with experimentally confirmed pathogenicity (Table **1**) include known plant pathogens (Nguyen *et al*., 2016).

Seven of the 10 isolates with confirmed pathogenicity caused significant seedling mortality for at least one tree species during the inoculation experiments (Table **1**), reinforcing the importance of pathogens as a selective pressure and source of nonrandom seedling mortality that can directly shape plant community composition (Green *et al*., 2014). We also observed pathogen-caused stem damage, stunted growth, and wilting. These nonlethal infections can also be a structuring force by compromising a seedling's ability to survive future stressors (Ditommaso & Watson, 1995).

Seventeen of the 27 fungal isolates tested appear to be nonpathogenic. However, some of the isolates categorized as nonpathogenic may be pathogenic under optimal conditions. These isolates may not have caused disease in our experiments because we did not replicate the environmental conditions necessary for disease development (Álvarez-Loayza *et al*., 2011) or due to the absence of a susceptible host (each isolate was only tested against three to 28 tree species, a fraction of the locally available hosts). While a number of the fungi tested can infect multiple tissue types (Table **1**; Fig. **S4d**), some of the isolates deemed nonpathogenic based on our soil inoculation technique may only be able to colonize leaves or stems.

*Pathogens can infect phylogenetically dispersed hosts*

The host associations of the fungi isolated from symptomatic seedlings, in combination with the results of the inoculation experiments, provide strong support for our hypothesis that seedling pathogens in tropical forests tend to be multi-host (Figures **1, 3**; Table **1**). Multi-host pathogens of seedlings, leaves, and seeds have been observed in previous Panama-based studies (Augspurger & Wilkinson, 2007; Gilbert & Webb, 2007; Schweizer *et al*., 2013; Sarmiento *et al*., 2017). We speculate that our study

underestimates the true breadth of the fungi's host ranges, as we sampled and tested a

small fraction of the >824 tree species present in the local forests (Pyke *et al*., 2001).

In general, a given fungal OTU was isolated from heterofamilial tree species, host use

is phylogenetically dispersed, and pairwise similarities in community composition of

seedling-associated fungi are unrelated to pairwise phylogenetic distances between tree

species (Figures **1, 2**; Tables **S4, S5**). Together these results suggest that, contrary to our

second hypothesis, there is no phylogenetic signal to host range for the pathogens

attacking seedlings in these tropical forests. The absence of a significant phylogenetic

signal is consistent with one Panama-based study of foliar pathogens (Schweizer *et al*.,

2013), but conflicts with other studies in temperate and tropical plant communities

(Gilbert & Webb, 2007; Gilbert *et al*., 2012; Parker *et al*., 2015; Chen *et al*., 2019). Our

study did not include as many congeneric or confamilial host pairs as Gilbert and Webb

(2007) or Gilbert *et al*. (2012), which may have hindered our ability to detect a signal.

Alternatively, pathogen host use may track phylogenetically labile plant defenses (Kursar

*et al*., 2009) rather than phylogenetically conserved defenses (e.g., Fine *et al*., 2006).

Parallel evolution of similar defenses in unrelated lineages would explain why the

pathogens in this and other studies can infect many distantly related tree species.

The positive relationships between OTU isolation frequency and the number of hosts

observed during our survey and inoculation experiments (Fig. **S5**) suggest that host

generalists are abundant members of the pathogen community and that, if our sampling

effort were expanded, a given fungal OTU would be detected in additional tree species.

Many seedling pathogens, such as the soil-borne fungi and oomycetes responsible for

seedling damping-off, are passively dispersed (e.g., by splashing rain), with limited

dispersal distances (Madden, 1997). Thus, the ability to infect a wide range of hosts,

particularly spatially aggregated hosts, would increase a pathogen's likelihood of

persistence. For pathogens with limited dispersal, host use may be more tightly linked

with plant habitat associations than with phylogenetic relatedness, making them more

likely to infect tree species that commonly co-occur spatially and temporally (Roy, 2001).

Furthermore, just as selection should favor multi-host pathogens in species-rich

assemblages, selection should favor true generalists, whose host ranges are decoupled

from plant relatedness, in plant communities that are phylogenetically overdispersed on

the small spatial scale relevant to dispersal-limited pathogens (Swenson *et al*., 2007;

Swenson *et al*., 2012; Pearse *et al*., 2013).

*Tree species share pathogens, but are differentially impacted*

A given plant species is often infected by multiple microorganisms and these

microorganisms are frequently shared among members of a plant community (Hersh *et*

*al*., 2012; Sarmiento *et al*., 2017; Spear & Mordecai, 2018). We observed overlap among

the fungal communities of distantly related tree species (Fig. **1**; Table **S5**), a pattern likely

to intensify with increased sampling. The presence of a particular fungus in the seedlings

of multiple host tree species does not mean that those hosts are equally likely to be

infected by that fungus. Pathogens may disproportionally infect certain hosts based on

availability, defense traits, and/or habitat (Ferrer & Gilbert, 2003).

Disease progression among susceptible hosts is mediated by multiple factors,

including defense traits. Based on our inoculation experiments, tree species were

differentially impacted by shared pathogens (Fig. **3**; Table **1**); in support of our third

hypothesis and consistent with the host-specific impacts of multi-host pathogens

observed in previous studies (Davidson *et al*., 2000; Augspurger & Wilkinson, 2007;

Hersh *et al*., 2012; Sarmiento *et al*., 2017). Most of the tree species we tested were

seemingly resistant to the 10 pathogenic isolates (Fig. **3**), which does not preclude their

susceptibility to numerous untested fungi. Moreover, these tree species may actually be

susceptible to infection by the pathogenic isolates in our study, but their tolerance

mechanisms prevent extensive pathogen colonization and symptom development. In

which case, these tolerant tree species may act as reservoir hosts that facilitate the

persistence of pathogens within the community (Haas *et al*., 2011).

*Plant habitats, seed sizes, and disease susceptibility*

Disease pressure varies spatially, particularly across environmental gradients

(Augspurger, 1984a; Brenes-Arguedas *et al*., 2009; Defossez *et al*., 2011; Haak *et al*.,

2012; Hersh *et al*., 2012; Spear *et al*., 2015). For plants adapted to areas with relatively

low and/or inconsistent disease pressure, such as areas with higher light, lower humidity,

and inconsistent precipitation (Augspurger, 1984a; Spear *et al*., 2015), the allocation

costs of physical and chemical defenses may outweigh the benefits and vice versa for

plants adapted to areas with relatively high and/or consistent disease pressure (Coley &

Barone, 1996; Strauss & Agrawal, 1999; Endara & Coley, 2010). Consistent with variable selective pressures, and previous field and greenhouse studies (Augspurger, 1984b; Augspurger & Kelly, 1984; McCarthy-Neumann & Kobe, 2008), seedlings of shade-intolerant tree species and tree species restricted to, or more abundant in, drier forests (dry-site species) were more likely to suffer disease than seedlings of shade-tolerant and wet-site tree species in our inoculation experiments (Fig. **4**; Tables **S6**, **S7**). Engelbrecht *et al*. (2007) found no correlation between a species' light requirements and its occurrence in dry- versus wet-sites, but evergreen trees are more common in wet-sites, while deciduous trees dominate dry-sites (Santiago *et al*., 2004). Shade-tolerant and evergreen tree species, both with long-lived leaves, tend to invest more in pest defenses than shade-intolerant and deciduous tree species (Coley & Barone, 1996; Santiago *et al*., 2004).

For shade-tolerant, shade-intolerant, and dry-site tree species, the probability of disease decreases with increasing seed size (Fig. **4**). This is perhaps unsurprising as large-seeded species can invest more in defenses and better compensate for lost tissue (Armstrong & Westoby, 1993; Green & Juniper, 2004). Our results are inconsistent with two Panama-based studies, which found no correlation between seed size and pathogen susceptibility, possibly because they only included wind-dispersed species while our analyses included wind- and vertebrate-dispersed species (average seed dry masses: 1.9 - 686 versus 3 - 2334 mg) (Augspurger, 1984a; Augspurger & Kelly, 1984).

*Consequences for plant communities*

Neither disturbance nor microhabitat specialization are believed to be sufficient to maintain the hyperdiversity of tropical forests (Leigh *et al*., 2004). While host-specialized pests are commonly touted as the true drivers of diversity maintenance (Leigh *et al*., 2004; Comita *et al*., 2014), we show that generalist pathogens are frequently responsible for seedling death and disease. The host-specific impacts of multi-host pathogens and differences among tree species in pathogen susceptibility observed by this and other studies (Augspurger & Wilkinson, 2007; Hersh *et al*., 2012; Spear *et al*., 2015) suggest that generalized pathogens have the potential to influence plant community diversity by unevenly affecting seedling recruitment. First, variability in host competence may maintain plant community diversity by creating a dilution effect (Ostfeld & Keesing,

2000). Overall disease risk should be lowered if the most competent host(s) is/are
relatively rare within the community because pathogen transmission would be slowed
between individuals of susceptible species. Accordingly, the presence of less competent
hosts should protect poorly defended host species from pathogen-caused local extinction.
Second, generalist pathogens with host-specific impacts may alter direct and indirect
interspecific competitive interactions among co-occurring species (Alexander & Holt,
1998; Barrett *et al.*, 2009; Mordecai, 2011). Host-generalized pathogens could promote
coexistence if a competition-defense tradeoff exists among plant species (i.e., if superior
competitors are more susceptible; Mordecai, 2011). Finally, interspecific variation in host
susceptibility to generalist pathogens may maintain diversity because disease-sensitive
tree species are excluded from disease-prone habitats. By reinforcing resource
partitioning and limiting the geographic distributions of species, generalist pathogens
contribute to the spatial turnover of plant species and the maintenance of local and
regional forest diversity.

**Acknowledgments**
This research was funded by Simons Foundation Award 429440, Center for Tropical
Forest Science, Smithsonian Tropical Research Institute, Garden Club of America,
University of Utah's Global Change and Sustainability Center, and A. Herbert and
Marian W. Gold Scholarship. Additional logistical support and facilities were provided
by the Smithsonian Tropical Research Institute, S. Mangan, and the Arnold Lab at
University of Arizona. We thank J. Sertich, D. Martinez, B. Wolfe, J. Ibarra, H. Espinosa,
C. Swatek, M. Schmoker, D. Sandberg, J. Shaffer, and C. Gonzalez for field and lab
assistance; R. Perez for help with tree species identification; J. Wright, G. Gilbert, S.
Mangan, A. E. Arnold, S. Higginbotham, T. Kursar, and P. Coley for advice; T. Brenes-
Arguedas for sharing her culture collection, and N. Anaya and G. LaRosa, who
contributed to that collection; and Parque Natural Metropolitano, D. Roubik, Barro
Colorado Nature Monument, and the Ministerio de Ambiente for authorizing this
research.

**Author Contributions:** ERS designed and conducted the research, analyzed the data,
and wrote the first draft of the manuscript, ERS and KDB contributed to manuscript
revisions and approved the final version.

**References**

**Abarenkov K, Zirk A, Piirmann T, Pöhönen R, Ivanov F, Nilsson RH, Kõljalg U. 2020.** Full UNITE+INSD dataset for Fungi. UNITE Community. 10.15156/BIO/786372

**Agrawal AA. 2007.** Macroevolution of plant defense strategies. *Trends in Ecology & Evolution* **22**: 103-109.

**Albert A, Anderson JA. 1984.** On the existence of maximum likelihood estimates in logistic regression models. *Biometrika* **71**: 1-10.

**Álvarez-Loayza P, White JF Jr, Torres MS, Balslev H, Kristiansen T, Svenning JC, Gil N. 2011.** Light converts endosymbiotic fungus to pathogen, influencing seedling survival and niche-space filling of a common tropical tree, *Iriartea deltoidea*. *PloS ONE* **6**: e16386.

**Armstrong DP, Westoby M. 1993.** Seedlings from large seeds tolerated defoliation better: a test using phylogeneticaly independent contrasts. *Ecology* **74**: 1092-1100.

**Augspurger CK. 1984a.** Seedling survival of tropical tree species: interactions of dispersal distance, light-gaps, and pathogens. *Ecology* **65**:1705-1712.

**Augspurger CK. 1984b.** Light requirements of neotropical tree seedlings: a comparative study of growth and survival. *Journal of Ecology* **72**: 777-795.

**Augspurger CK, Kelly CK. 1984.** Pathogen mortality of tropical tree seedlings: experimental studies of the effects of dispersal distance, seedling density, and light conditions. *Oecologia* **61**: 211-217.

**Augspurger CK, Wilkinson HT. 2007.** Host specificity of pathogenic *Pythium* species: Implications for tree species diversity. *Biotropica* **39**: 702–708.

**Bagchi R, Gallery RE, Gripenberg S, Gurr SJ, Narayan L, Addis CE, Freckleton RP, Lewis OT. 2014.** Pathogens and insect herbivores drive rainforest plant diversity and composition. *Nature* **506**: 85-88.

**Barrett LG, Kniskern JM, Bodenhausen N, Zhang W, Bergelson J. 2009.** Continua of specificity and virulence in plant host–pathogen interactions: causes and consequences. *New Phytologist* **183**: 513-529.

**Bartoń K. 2020.** MuMIn: Multi-Model Inference. R package version 1.43.17. [WWW
document] URL https://CRAN.R-project.org/package=MuMIn [accessed 2 June
2020].

**Bell T, Freckleton RP, Lewis OT. 2006.** Plant pathogens drive density-dependent
seedling mortality in a tropical tree. *Ecology Letters* **9**: 569–574.

**Benítez MS, Hersh MH, Vilgalys R, Clark JS. 2013.** Pathogen regulation of plant
diversity via effective specialization. *Trends in Ecology and Evolution* **28**: 705–
711.

**Benson DA, Cavanaugh M, Clark K, Karsch-Mizrachi I, Lipman DJ, Ostell J,**
**Sayers EW. 2012.** GenBank. *Nucleic Acids Research* **41**: D36-D42.

**Blüthgen N, Menzel F, Blüthgen N. 2006.** Measuring specialization in species
interaction networks. *BMC Ecology* **6**: 9.

**Bradley DJ, Gilbert GS, Parker IM. 2003.** Susceptibility of clover species to fungal
infection: the interaction of leaf surface traits and environment. American *Journal*
*of Botany* **90**: 857-864.

**Brenes-Arguedas T, Coley PD, Kursar TA. 2009.** Pests versus drought as determinants
of plant distribution along a tropical rainfall gradient. *Ecology* **90**: 1751-1761.

**Brenes-Arguedas T, Roddy AB, Coley PD, Kursar TA. 2011.** Do differences in
understory light contribute to species distributions along a tropical rainfall
gradient? *Oecologia* **166**: 443-456.

**Chao A, Chazdon RL, Colwell RK, Shen T. 2005.** A new statistical approach for
assessing similarity of species composition with incidence and abundance data.
*Ecology Letters* **8**: 148-159.

**Chen L, Swenson NG, Ji N, Mi X, Ren H, Guo L, Ma K. 2019.** Differential soil fungus
accumulation and density dependence of trees in a subtropical forest. *Science* **366**:
124-128.

**Coley PD, Barone JA. 1996.** Herbivory and plant defenses in tropical forests. *Annual*
*Review of Ecology and Systematics* **27**: 305-335.

**Connell JH. 1971.** On the role of natural enemies in preventing competitive exclusion in
some marine animals and in rain forest trees. In Boer PJD, Gradwell GR, eds.

*Dynamics of numbers in populations.* Wageningen, Netherlands: Centre for
Agricultural Publication and Documentation, 298–312.

**Comita LS, Muller-Landau HC, Aguilar S, Hubbell SP. 2010.** Asymmetric density
dependence shapes species abundances in a tropical tree community. *Science* **329**:
330-332.

**Comita LS, Queenborough SA, Murphy SJ, Eck JL, Xu K, Krishnadas M, Beckman
N, Zhu Y. 2014**. Testing predictions of the Janzen–Connell hypothesis: a meta-
analysis of experimental evidence for distance-and density-dependent seed and
seedling survival. *Journal of Ecology* **102**: 845-856.

**Davidson JM, Rehner SA, Santana M, Lasso E, Urena de Chapet O, Herre EA.
2000.** First report of *Phytophthora heveae* and *Pythium* spp. on tropical tree
seedlings in Panama. *Plant Disease* **84**: 706-706.

**Defossez E, Courbaud B, Marcais B, Thuiller W, Granda E, Kunstler G. 2011.** Do
interactions between plant and soil biota change with elevation? A study on *Fagus
sylvatica. Biology Letters* **7**: 699-701.

**Ditommaso A, Watson AK. 1995.** Impact of a fungal pathogen, *Colletotrichum
coccodes* on growth and competitive ability of *Abutilon theophrasti. New
Phytologist* **131**: 51-60.

**Dormann CF. 2011.** How to be a specialist? Quantifying specialisation in pollination
networks. *Network Biology* **1**: 1-20.

**Dormann CF, Gruber B, Fruend J. 2008.** Introducing the bipartite Package: analysing
ecological networks. *R news* **8**: 8-11.

**Endara MJ, Coley PD. 2011.** The resource availability hypothesis revisited: a meta-
analysis. *Functional Ecology* **25**: 389-398.

**Engelbrecht BM, Comita LS, Condit R, Kursar TA, Tyree MT, Turner BL, Hubbell
SP. 2007.** Drought sensitivity shapes species distribution patterns in tropical
forests. *Nature* **447**: 80–82.

**Fine PVA, Miller ZJ, Mesones I, Irazuzta S, Appel HM, Stevens MHH, Sääksjärvi I,
Schultz JC, Coley PD. 2006.** The growth-defense trade-off and habitat
specialization by plants in Amazonian forests. *Ecology* **87**: S150–S162.

**Firth D. 1993.** Bias reduction of maximum likelihood estimates. *Biometrika* **80**: 27–38.

**Ferrer A, Gilbert GS. 2003.** Effect of tree host species on fungal community

composition in a tropical rain forest in Panama. *Diversity and Distributions* **9**:

455-468.

**Gallery RE, Dalling JW, Arnold AE. 2007.** Diversity, host affinity, and distribution of

seed-infecting fungi: A case study with *Cecropia*. *Ecology* **88**: 582–588.

**Gilbert GS, Magarey R, Suiter K, Webb CO. 2012.** Evolutionary tools for

phytosanitary risk analysis: phylogenetic signal as a predictor of host range of

plant pests and pathogens. *Evolutionary Applications* **5**: 869-878.

**Gilbert GS, Webb CO. 2007.** Phylogenetic signal in plant pathogen-host range.

*Proceedings of the National Academy of Sciences, USA* **104**: 4979-4983.

**Gillett JB. 1962.** Pest pressure, and underestimated factor in evolution. *Systematics*

*Association Publication* **4**: 37–46.

**Green PT, Harms KE, Connell JH. 2014.** Nonrandom, diversifying processes are

disproportionately strong in the smallest size classes of a tropical forest.

*Proceedings of the National Academy of Sciences, USA* **111**: 18649-18654.

**Green PT, Juniper PA. 2004.** Seed mass, seedling herbivory and the reserve effect in

tropical rainforest seedlings. *Functional Ecology* **18**: 539-547.

**Grünwald NJ, Martin FN, Larsen MM, Sullivan CM, Coffey MD, Hansen EM,**

**Parke JL. 2011.** Phytophthora-ID.org: A sequence-based Phytophthora

identification tool. *Plant Disease* **95**: 337-342.

**Haak DC, McGinnis LA, Levey DJ, Tewksbury JJ. 2012.** Why are not all chilies hot?

A trade-off limits pungency. *Proceedings of the Royal Society B: Biological*

*Sciences* **279**: 2012-2017.

**Haas SE, Hooten MB, Rizzo DM, Meentemeyer RK. 2011.** Forest species diversity

reduces disease risk in a generalist plant pathogen invasion. *Ecology Letters* **14**:

1108-1116.

**Harms KE, Wright SJ, Calderon O, Hernandez A, Herre EA. 2000.** Pervasive

density-dependent recruitment enhances seedling diversity in a tropical forest.

*Nature* **404**: 493–495.

**Hersh MH, Vilgalys R, Clark JS. 2012**. Evaluating the impacts of multiple generalist

fungal pathogens on temperate tree seedling survival. *Ecology* **93**: 511-520.

**Higgins KL, Coley PD, Kursar TA, Arnold AE. 2011.** Culturing and direct PCR

suggest prevalent host generalism among diverse fungal endophytes of tropical

forest grasses. *Mycologia* **103**: 247-260.

**Hofstetter V, Buyck B, Eyssartier G, Schnee S, Gindro K. 2019.** The unbearable

lightness of sequenced-based identification. *Fungal Diversity* **96**: 243-284.

**Huot B, Yao J, BMontgomery BL, He SY. 2014.** Growth–defense tradeoffs in plants: a

balancing act to optimize fitness. *Molecular Plant* **7**: 1267-1287.

**Janzen DH. 1970.** Herbivores and the number of tree species in tropical forests.

*American Naturalist* **104**: 501–528.

**Kang S, Mansfield MA, Park B, Geiser DM, Ivors KL, Coffey MD, Grünwald NJ,**

**Martin FN, Lévesque CA, Blair JE. 2010.** The promise and pitfalls of sequence-

based identification of plant-pathogenic fungi and oomycetes. *Phytopathology*

**100**: 732-737.

**Kembel SW, Cowan PD, Helmus MR, Cornwell WK, Morlon H, Ackerly DD,**

**Blomberg SP, Webb CO. 2010.** Picante: R tools for integrating phylogenies and

ecology. *Bioinformatics* **26**: 1463-1464.

**Kindt R, Coe R. 2005.** *Tree diversity analysis. A manual and software for common*

*statistical methods for ecological and biodiversity studies*. Nairobi, Kenya: World

Agroforestry Centre (ICRAF).

**Klinke S. 2019.** plot.matrix: Visualizes a Matrix as Heatmap. R package version 1.4.

[WWW document] URL https://CRAN.R-project.org/package=plot.matrix

[accessed 2 June 2020].

**Kluger CG, Dalling JW, Gallery RE, Sanchez E, Weeks-Galindo C, Arnold AE.**

**2008.** Host generalists dominate fungal communities associated with seeds of four

neotropical pioneer species. *Journal of Tropical Ecology* **24**: 351–354.

**Kõljalg U, Nilsson RH, Abarenkov K, Tedersoo L, Taylor AFS, Bahram M, Bates**

**ST, Bruns TD, Bengtsson-Palme J, Callaghan TM *et al.* 2013.** Towards a

unified paradigm for sequence-based identification of fungi. *Molecular Ecology*

**22**: 5271–5277.

**Kosmidis I. 2019.** brglm: Bias Reduction in Binary-Response Generalized Linear
Models. R package version 0.6.2. [WWW document] URL https://cran.r-
project.org/package=brglm [accessed 2 June 2020].

**Kosmidis I, Firth D. 2009.** Bias reduction in exponential family nonlinear models.
*Biometrika* **96**: 793–804.

**Kursar TA, Dexter KG, Lokvam J, Pennington RT, Richardson JE, Weber MG,**
**Murakami ET, Drake C, McGregor R, Coley PD. 2009.** The evolution of
antiherbivore defenses and their contribution to species coexistence in the tropical
tree genus *Inga. Proceedings of the National Academy of Sciences, USA* **106**:
18073-18078.

**Leigh EG, Davdar P, Dick CW, Puyravaud J-P, Terborgh J, ter Steege H, Wright**
**SJ. 2004.** Why do some tropical forests have so many species of trees? *Biotropica*
**36**: 447-473.

**Lesnoff M, Lancelot R. 2012.** aod: Analysis of Overdispersed Data. R package version
1.3.1. [WWW document] URL http://cran.r-project.org/package=aod [accessed 2
June 2020].

**Liu Y, Yu S, Xie Z, Staehelin C. 2012.** Analysis of a negative plant–soil feedback in a
subtropical monsoon forest. *Journal of Ecology* **100**: 1019–1028.

**Lücking R, Aime MC, Robbertse B, Miller AN, Ariyawansa HA, Aoki T, Cardinali**
**G, Crous PW, Druzhinina IS, Geiser DM. 2020.** Unambiguous identification of
fungi: where do we stand and how accurate and precise is fungal DNA barcoding?
*IMA Fungus* **11**: 1-32.

**Madden LV. 1997.** Effects of rain on splash dispersal of fungal pathogens. *Canadian*
*Journal of Plant Pathology* **19**: 225-230.

**Mangan SA, Schnitzer SA, Herre EA, Mack KM, Valencia MC, Sanchez EI, Bever**
**J. 2010.** Negative plant-soil feedback predicts tree-species relative abundance in a
tropical forest. *Nature* **466**: 752-755.

**May RM. 1991.** A fondness for fungi. *Nature* **352**: 475-476.

**McCarthy-Neumann S, Kobe RK. (2008).** Tolerance of soil pathogens co-varies with
shade tolerance across species of tropical tree seedlings. *Ecology* **89**: 1883-1892.

**Mordecai EA. 2011.** Pathogen impacts on plant communities: Unifying theory, concepts,
and empirical work. *Ecological Monograph* **81**: 429-441.

**Mordecai EA. 2012.** Soil moisture and fungi affect seed survival in California grassland
annual plants. *PloS ONE* **7:** e39083.

**Nguyen NH, Song Z, Bates ST, Branco, S, Tedersoo L, Menke J, Schilling JS,**
**Kennedy PG. 2016.** FUNGuild: an open annotation tool for parsing fungal
community datasets by ecological guild. *Fungal Ecology* **20**: 241-248.

**Oksanen J, Blanchet FG, Friendly M, Kindt R, Legendre P, McGlinn D, Minchin**
**PR, O'Hara RB, Simpson GL, Solymos P *et al*. 2019.** vegan: Community
Ecology Package. R package version 2.5-6. [WWW document] URL
https://CRAN.R-project.org/package=vegan [accessed 2 June 2020].

**Ostfeld RS, Keesing F. 2000.** Biodiversity and disease risk: the case of Lyme disease.
*Conservation Biology* **14**: 722–728.

**Packer A, Clay K. 2000.** Soil pathogens and spatial patterns of seedling mortality in a
temperate tree. *Nature* **404**: 278-281.

**Paradis E, Schliep K. 2018.** ape 5.0: an environment for modern phylogenetics and
evolutionary analyses in R. *Bioinformatics* **35**: 526-528.

**Parker IM, Saunders M, Bontrager M, Weitz AP, Hendricks R, Magarey R, Suiter**
**K, Gilbert GS. 2015.** Phylogenetic structure and host abundance drive disease
pressure in communities. *Nature* **520**: 542-544.

**Paton S. 2019a.** Barro Colorado Island, Lutz tower 1m_Air Temperature. The
Smithsonian Institution. Dataset. https://doi.org/10.25573/data.10042394.v7

**Paton S. 2019b.** Barro Colorado Island, Lutz tower 1m_Relative Humidity. The
Smithsonian Institution. Dataset. https://doi.org/10.25573/data.10042400.v7

**Pearse WD, Jones FA, Purvis A. 2013.** Barro Colorado Island's phylogenetic
assemblage structure across fine spatial scales and among clades of different ages.
*Ecology* **94**: 2861-2872.

**Pyke CR, Condit R, Aguilar S, Lao S. 2001.** Floristic composition across a climatic
gradient in a neotropical lowland forest. *Journal of Vegetation Science* **12**: 553–
566.

**Quinn GP, Keough MJ. 2002.** *Experimental design and data analysis for biologists*.

Cambridge University Press.

**R Core Team. 2020.** R: a language and environment for statistical computing. R

Foundation for Statistical Computing, Vienna, Austria.

**Robbertse B, Strope PK, Chaverri P, Gazis R, Ciufo S, Domrachev M, Schoch CL.**

**2017.** Improving taxonomic accuracy for fungi in public sequence databases:

applying 'one name one species' in well-defined genera with

*Trichoderma*/*Hypocrea* as a test case. *Database* **2017**

**Roy BA. 2001.** Patterns of association between crucifers and their flower-mimic

pathogens: host jumps are more common than coevolution or cospeciation.

*Evolution* **55**: 41-53.

**Rossi J-P. 2011.** rich: an R package to analyse species richness. *Diversity* **3**: 112-120.

**Ruxton GD, Neuhäuser M. 2010.** When should we use one-tailed hypothesis testing?

*Methods in Ecology and Evolution* **1**: 114-117.

**Sarmiento C, Zalamea PC, Dalling JW, Davis AS, Stump SM, U'Ren JM, Arnold**

**AE. 2017.** Soilborne fungi have host affinity and host-specific effects on seed

germination and survival in a lowland tropical forest. *Proceedings of the National*

*Academy of Sciences, USA* **114**: 11458-11463.

**Santiago LS, Kitajima K, Wright SJ, Mulkey SS. 2004.** Coordinated changes in

photosynthesis, water relations and leaf nutritional traits of canopy trees along a

precipitation gradient in lowland tropical forest. *Oecologia* **139**: 495-502.

**Schweizer D, Gilbert GS, Holl KD. 2013.** Phylogenetic ecology applied to enrichment

planting of tropical native tree species. *Forest Ecology and Management* **297**: 57-

66.

**Spear ER. 2017.** Phylogenetic relationships and spatial distributions of putative fungal

pathogens of seedlings across a rainfall gradient in Panama. *Fungal Ecology* **26**:

65-73.

**Spear ER, Coley PD, Kursar TA. 2015.** Do pathogens limit the distributions of tropical

trees across a rainfall gradient? *Journal of Ecology* **103**: 165-174.

**Spear ER, Mordecai EA. 2018.** Foliar pathogens of California grasses are multi-host
and spatially widespread: implications for grassland diversity. *Ecology* **99**: 2250-
2259.

**Strauss SY, Agrawal AA. 1999.** The ecology and evolution of plant tolerance to
herbivory. *Trends in Ecology and Evolution* **14**: 179–185.

**Swenson NG, Enquist BJ, Thompson J, Zimmerman JK. 2007.** The influence of
spatial and size scale on phylogenetic relatedness in tropical forest communities.
*Ecology* **88**: 1770-1780.

**Swenson NG, Erickson DL, Mi X, Bourg NA, Forero-Montaña J, Ge X, Howe R,**
**Lake JK, Liu X, Ma K** *et al*. **2012.** Phylogenetic and functional alpha and beta
diversity in temperate and tropical tree communities. *Ecology* **93**: S112-S125.

**Talley SM, Coley PD, Kursar TA. 2002.** Anti-fungal leaf surface metabolites correlate
with fungal abundance in sagebrush populations. *Journal of Chemical Ecology*
**28**: 2141–2168.

**Vavrek MJ. 2011.** fossil: palaeoecological and palaeogeographical analysis tools.
*Palaeontologia Electronica* **14**: 1T.

**Wickham H. 2007.** Reshaping data with the reshape package. *Journal of Statistical*
Software **21**: 1-20.

**Wickham H. 2011.** The split-apply-combine strategy for data analysis. *Journal of*
*Statistical Software* **40**: 1-29.

**Wickham H. 2016.** *ggplot2: elegant graphics for data analysis*. New York, NY, USA:
Springer-Verlag.

**Yadeta KA, Thomma, BPHJ. 2013.** The xylem as battleground for plant hosts and
vascular wilt pathogens. *Frontiers in Plant Science* **4**: 97.

**Yang S. 2020.** otuSummary: summarizing OTU table regarding the composition,
abundance and beta diversity of abundant and rare biospheres. R package version
0.1.1. [WWW document] URL https://CRAN.R-
project.org/package=otuSummary [accessed 2 June 2020].

**Zanne AE, Tank DC, Cornwell WY, Eastman JM, Smith SA, FitzJohn RG,**
**McGlinn DJ, O'Meara BC, Moles AT, Reich PB** *et al*. **2014.** Three keys to the
radiation of angiosperms into freezing environments. *Nature* **506**: 89-92.

**Supporting Information**

**Figure S1** Examples of disease in the forests of Panama.

**Figure S2** Details of shadehouse-based inoculation experiments.

**Figure S3** Rank abundance plot and OTU accumulation curve.

**Figure S4** Overlap in fungal OTUs among sampling years, methods used to obtain

symptomatic seedlings, isolation media, and tissue sampled.

**Figure S5** Correlation between OTU host range and isolation frequency.

**Table S1** Taxonomic assignments, traits, sampling effort, and observed OTUs for tree

species evaluated in our survey and experiments.

**Table S2** Methodological details pertaining to the multi-year collection of symptomatic

seedlings, and microbial isolation and sequencing.

**Table S3** Average light levels, air temperatures, and relative humidities of the

shadehouses used for the inoculation experiments versus ambient conditions.

**Table S4** Estimated taxonomic placement, isolation frequency, number of observed

hosts, estimated host specialization, and phylogenetic pattern of host use of the OTUs.

**Table S5** Overlap in seedling-associated OTUs among tree species.

**Table S6** Results of the beta-binomial generalized linear regression with the proportion

of diseased seedlings as a function of seed size and shade tolerance.

**Table S7** Average estimates based on the best-ranked beta-binomial generalized linear

regressions with the proportion of diseased seedlings as a function of seed size and spatial

distribution relative to annual rainfall.

**Methods S1** Methods used to estimate the taxonomic placements of the 66 OTUs and

assign nomenclature.

**Figure 1.** Host associations of the fungi and oomycetes isolated from symptomatic

seedlings (full dataset: *n* = 211 isolates). The 26 host tree species (left) are arranged

according to their phylogenetic relationships. The estimated taxonomic placement of each

of the 66 microbial OTUs (right) is preceded by a unique letter identifier. Because our

OTU grouping strategy (99% sequence similarity) often differed from that of the UNITE

database, distinct OTUs in our study sometimes share the same nomenclature based on

the species hypothesis of the best-matching reference sequence in UNITE (e.g., Z and

AAG). For the non-singleton OTUs, nomenclature is followed by an L or M, indicating

low or moderate estimated specialization (d'). The width of each line represents the

number of times a given OTU was isolated from seedlings of a given tree species. For

both tree species and OTUs, font size is scaled to the number of interactions. For two

OTUs (Q and R, starred), phylogenetic clustering of host use was marginally significant

($P = 0.09$). The six fungal OTUs with members that exhibited statistically significant ($\alpha =$

0.10) pathogenicity during the inoculation experiments are listed in red.

**Figure 2.** Estimated pairwise similarities of OTU communities are unrelated to pairwise
phylogenetic distances (time of independent evolution in Myr). The blue circles and solid
line (GLM: Est. ± SE = 0.002 ± 0.003, $z$-value = 0.60, $P$ = 0.55) are based on the full
dataset ($n$ = 211 isolates, representing 66 OTUs, from 26 tree species), while the green
triangles and dotted line (GLM: Est. ± SE = 0.002 ± 0.004, $z$-value = 0.36, $P$ = 0.72) are
based on subset B, excluding singleton and doubleton observations of tree species and
OTUs ($n$ = 144 isolates, representing 22 OTUs, from 13 tree species).

**Figure 3.** Pathogenicity and host ranges of the 27 fungal isolates tested, and

susceptibility of the 35 tree species tested, based on seedling morbidity. Fungal isolates

are grouped according to their OTU (horizontal axis). Tree species (Table **S1**) are

arranged based on their phylogenetic relationships (vertical axis). Tree species-by-fungal

isolate combinations for which evidence of pathogenicity was observed (proportion of

inoculated seedlings with symptoms > proportion in paired control group) are depicted in

shades of yellow (pale for minimal) to red (dark for substantial). Combinations for which

there was no evidence of pathogenicity (proportion of inoculated seedlings with

symptoms ≤ proportion in paired control group) are depicted in black and shades of

green. Gray cells represent an untested combination. For 20 of the 285 combinations

tested, we observed significant pathogenicity (***$P \leq 0.001$, **$P \leq 0.01$, *$P \leq 0.05$, †$P \leq$

0.10; Table **1**).

**Figure 4.** Relationships between disease susceptibility and seed size and shade tolerance

(a,b), and seed size and spatial distribution relative to annual rainfall (c,d). Shade-

intolerant tree species (red) are more likely to experience disease than shade-tolerant

species (blue) ($P = 0.003$), and for both groups the probability of disease decreases with

increasing seed size (a,b) (Table **S6**). Dry-site tree species (orange) are more likely to

experience disease than wet-site species (green) ($P = 0.042$; Table **S7**). For dry-site

species (orange) the probability of disease decreases with increasing seed size (c,d)

(Table **S7**).

**Table 1.** The 20 fungal isolate-by-tree species combinations for which significant pathogenicity was observed based on the presence/absence of (i) general morbidity and (ii) mortality in the inoculated versus control seedlings, analyzed with bias-reduced generalized linear models assuming binomial error distributions and probit link functions and using α = 0.10 because small sample sizes limited our power to detect treatment effects.

| OTU – Estimated taxonomic placement (isolate, origin tree sp. & tissue, no. of tree spp. tested) | Susceptible tree species | $n$ Cntrl | Inoc | Disease metric | Prop. rel. to control | Est (SE) | z | P |
|---|---|---|---|---|---|---|---|---|
| | D. retusa[1] | 5 | 5 | morbidity | 0.8 | 2.169 (1.044) | 2.076 | 0.038 |
| | | | | mortality | 1 | 2.932 (1.195) | 2.454 | 0.014 |
| | G. standleyana | 10 | 10 | morbidity | 0.6 | 1.609 (0.651) | 2.471 | 0.014 |
| B - Nectriaceae (ERS034, TT root, 28) | J. copaia[1] | 10 | 10 | morbidity | 0.6 | 1.996 (0.828) | 2.411 | 0.016 |
| | | | | mortality | 0.6 | 1.996 (0.828) | 2.411 | 0.016 |
| | L. seemannii[1] | 5 | 5 | morbidity | 1 | 2.932 (1.195) | 2.454 | 0.014 |
| | | | | mortality | 1 | 2.932 (1.195) | 2.454 | 0.014 |
| | C. elastica | 10 | 10 | morbidity[8] | 0.8 | 2.252 (0.711) | 3.168 | 0.002 |
| B - Nectriaceae (ERS036, CL stem[2], 13) | C. vitifolium | 10 | 10 | morbidity[9] | 0.4 | 1.126 (0.640) | 1.759 | 0.079 |
| | | | | mortality | 0.4 | 1.126 (0.640) | 1.759 | 0.079 |
| | D. retusa | 10 | 10 | morbidity[10] | 0.5 | 1.36 (0.643) | 2.117 | 0.034 |

| | | | | | | | | |
|---|---|---|---|---|---|---|---|---|
| G – *Clonostachys rosea*, Bionectriaceae (ERS011, VS stem[3], 17) | *C. longifolium* | 5 | 5 | morbidity | 0.6 | 1.684 (1.017) | 1.657 | 0.098 |
| G – *C. rosea* (ERS012, PT leaf[4], 15) | *C. aromatica* | 5 | 5 | morbidity[11,12] | 0.6 | 1.684 (1.017) | 1.657 | 0.098 |
| G – *C. rosea* (ERS019, VS root, 12) | *L. seemannii* | 10 | 10 | morbidity | 0.4 | 1.126 (0.640) | 1.759 | 0.079 |
| | | | | mortality | 0.4 | 1.126 (0.640) | 1.759 | 0.079 |
| | *T. tuberculata* | 8 | 9 | morbidity | 0.56 | 1.796 (0.867) | 2.072 | 0.038 |
| | | | | mortality | 0.44 | 1.540 (0.867) | 1.776 | 0.076 |
| C - *Mycoleptodiscus suttoni*, Muyocopronaceae (ERS029, VS stem[5], 13) | *A. glabra* | 12 | 12 | morbidity | 0.75 | 2.462 (0.800) | 3.078 | 0.002 |
| | | | | mortality | 0.58 | 2.033 (0.788) | 2.579 | 0.010 |
| | *D. retusa* | 10 | 10 | morbidity | 0.7 | 1.891 (0.669) | 2.826 | 0.005 |
| | | | | mortality | 0.8 | 2.526 (0.848) | 2.977 | 0.003 |
| | *G. americana* | 10 | 10 | morbidity | 0.5 | 1.360 (0.643) | 2.117 | 0.034 |
| | *P. limonensis* | 10 | 10 | morbidity | 0.4 | 1.527 (0.828) | 1.844 | 0.065 |
| | | | | mortality | 0.4 | 1.527 (0.828) | 1.844 | 0.065 |
| C - *M. suttoni* (ERS033, RA stem[6], 16) | *L. seemannii* | 10 | 10 | morbidity[13] | 0.9 | 2.887 (0.882) | 3.274 | 0.001 |
| | | | | mortality | 0.9 | 2.887 (0.882) | 3.274 | 0.001 |
| AAF - Nectriaceae (ERS040, ANE stem[7], 12) | *C. elastica* | 10 | 9 | morbidity[14] | 0.9 | 2.843 (0.895) | 3.178 | 0.002 |
| | | | | mortality | 0.89 | 2.827 (0.890) | 3.177 | 0.002 |
| AE – *Pestalotiopsis*, Sporocadaceae (ERS031, TT root, 18) | *B. utile* | 10 | 10 | morbidity | 0.4 | 1.527 (0.828) | 1.844 | 0.065 |
| | | | | mortality | 0.4 | 1.527 (0.828) | 1.844 | 0.065 |
| | *G. americana* | 10 | 10 | morbidity | 0.4 | 1.126 (0.640) | 1.759 | 0.079 |
| | | | | mortality | 0.4 | 1.126 (0.640) | 1.759 | 0.079 |

| I - Sordariomycetes (ERS032, HC stem, 14) | C. aromatica | 5 | 5 | morbidity | 0.6 | 1.684 (1.017) | 1.657 | 0.098 |

See Table **S1** for the full names of the tree species. Prop. rel. to control is the proportion of inoculated seedlings with the disease

metric minus the proportion in the paired control group. We considered an isolate pathogenic if, for at least one tree species, a

significantly greater proportion of inoculated seedlings were symptomatic than paired control seedlings.

[1] Experiment end date = date all inoculated seedlings had died (i.e., used truncated observations of paired control seedlings).

[2,3,4,5,6,7] See **Fig. S1j,t,f,q,r,v**, respectively, for photos.

[8,9,10,11,12,13,14] See **Fig. S2r,n,k,g,h,m,s**, respectively, for photos.