The mechanism of SARS-CoV-2 nucleocapsid protein
recognition by the human 14-3-3 proteins

Kristina V. Tugaeva [1,a] , Dorothy E. D. P. Hawkins [2,a] , Jake L. R. Smith [2], Oliver W.
Bayfield [2], De-Sheng Ker [2], Andrey A. Sysoev [1], Oleg I. Klychnikov [3], Alfred A. Antson
[2*], Nikolai N. Sluchanko [1*]

[1] A.N. Bach Institute of Biochemistry, Federal Research Center of Biotechnology of the
Russian Academy of Sciences, 119071, Moscow, Russia

[2] York Structural Biology Laboratory, Department of Chemistry, University of York, York
YO10 5DD, United Kingdom

[3] Department of Biochemistry, School of Biology, M.V. Lomonosov Moscow State
University, 119991, Moscow, Russia

[a] Co-first; *Corresponding authors: nikolai.sluchanko@mail.ru , fred.antson@york.ac.uk

Highlights

SARS-CoV-2 nucleocapsid protein (N) binds to all seven human 14-3-3 isoforms.

This association with 14-3-3 strictly depends on phosphorylation of N.

The two proteins interact in 2:2 stoichiometry and with the Kd in a µM range.

Affinity of interaction depends on the specific 14-3-3 isoform.

Conserved Ser197-phosphopeptide of N is critical for the interaction.

Keywords

Host-pathogen interactions; protein-protein complex; phosphorylation;
nucleocytoplasmic shuttling; stoichiometry

Abstract

The coronavirus nucleocapsid protein (N) controls viral genome packaging and contains numerous phosphorylation sites located within unstructured regions. Binding of phosphorylated SARS-CoV N to the host 14-3-3 protein in the cytoplasm was reported to regulate nucleocytoplasmic N shuttling. All seven isoforms of the human 14-3-3 are abundantly present in tissues vulnerable to SARS-CoV-2, where N can constitute up to ~1% of expressed proteins during infection. Although the association between 14-3-3 and SARS-CoV-2 N proteins can represent one of the key host-pathogen interactions, its molecular mechanism and the specific critical phosphosites are unknown. Here, we show that phosphorylated SARS-CoV-2 N protein (pN) dimers, reconstituted via bacterial co-expression with protein kinase A, directly associate, in a phosphorylation-dependent manner, with the dimeric 14-3-3 protein, but not with its monomeric mutant. We demonstrate that pN is recognized by all seven human 14-3-3 isoforms with various efficiencies and deduce the apparent K_D to selected isoforms, showing that these are in a low micromolar range. Serial truncations pinpointed a critical phosphorylation site to Ser197, which is conserved among related zoonotic coronaviruses and located within the functionally important, SR-rich region of N. The relatively tight 14-3-3/pN association can regulate nucleocytoplasmic shuttling and other functions of N via occlusion of the SR-rich region, while hijacking cellular pathways by 14-3-3 sequestration. As such, the assembly may represent a valuable target for therapeutic intervention.

Introduction

The new coronavirus-induced disease, COVID19, has caused a worldwide health crisis with more than 90 million confirmed cases and 1.9 million deaths as of January 2021 [1]. The pathogen responsible, Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2), is highly similar to the causative agent of the SARS outbreak in 2002-2003 (SARS-CoV) and, to a lesser extent, to the Middle East Respiratory Syndrome Coronavirus (MERS-CoV) [2, 3]. Each is vastly more pathogenic and deadly than human coronaviruses HCoV-OC43, HCoV-NL63, HCoV-229E, and HCoV-HKU1 which cause seasonal respiratory diseases [4]. Like SARS-CoV and HCoV-NL63, SARS-CoV-2 uses angiotensin-converting enzyme 2 (ACE2) as entry receptor [5]. The ACE2 expression roughly correlates with the evidenced SARS-CoV-2 presence in different tissue types, which explains the multi-organ character of the disease [6] (Fig. 1).

In contrast to multiple promising COVID19 vaccine clinical trials [7-9], treatment of the disease is currently limited by the absence of approved efficient drugs [10]. The failure of several leading drug candidates in 2020 warrants the search for novel therapeutic targets including not only viral enzymes, but also heterocomplexes involving viral and host cell proteins. Unravelling mechanisms of interaction between the host and pathogen proteins may provide the platform for such progress.

The positive-sense single-stranded RNA genome of SARS-CoV-2 coronavirus encodes approximately 30 proteins which enable cell penetration, replication, viral gene transcription and genome assembly amongst other functions [11]. The 46-kDa SARS-CoV-2 Nucleocapsid (N) protein is 89.1% identical to SARS-CoV N. Genomic analysis

of human coronaviruses indicated that N might be the major factor conferring the

enhanced pathogenicity to SARS-CoV-2 [4]. N represents the most abundant viral

protein in the infected cell [12-14], with each assembled virion containing approximately

one thousand molecules of N [15]. Given that each infected cell can contain up to 10^5

virions (infectious, defective and incomplete overall) [14], the number of N molecules in

an infected cell can reach 10^8, accounting for ~1% of a total number of cellular proteins

(~10^{10} [16]).

The N protein interacts with viral genomic RNA, the membrane (M) protein and

self-associates to provide for the efficient virion assembly [17-19]. It consists of two

structured domains and three unstructured regions, including a functionally important

central Ser/Arg-rich region (Fig. 2A) [20-22]. Such organization allows for a vast

conformational change, which in combination with positively charged surfaces [23],

facilitates nucleic acid binding [24]. Indeed, the crystal structure of the N-terminal

domain (NTD) reveals an RNA binding groove [25-27], while crystal structures of the C-

terminal domain (CTD) show a highly interlaced dimer with additional nucleic acid

binding capacity [28, 29]. The N protein shows unusual properties in the presence of

RNA, displaying concentration-dependent liquid-liquid phase separation [22, 23, 30, 31]

that is pertinent to the viral genome packaging mechanism [32, 33]. In human cells, the

assembly of condensates is down-regulated by phosphorylation of the SR-rich region

[30, 34]. SARS-CoV-2 N protein is a major target of phosphorylation by host cell protein

kinases, with 22 phosphosites identified in vivo throughout the protein (Supplementary

table 1 and [13, 35]). Functions of N and viral replication can be regulated by a complex,

hierarchical phosphorylation of the SR-rich region of SARS-CoV-2 N by a cascade of

protein kinases [36]. Nevertheless, the potential functional role of N phosphorylation at

each specific site is not understood.

Using immunofluorescence, immunoprecipitation, siRNA silencing and kinase

inhibition, it has been shown that SARS-CoV N protein shuttles between the nucleus

and the cytoplasm in COS-1 cells [37]. This process is regulated by N protein

phosphorylation by several protein kinases including glycogen synthase kinase-3,

protein kinase A, casein kinase II and cyclin-dependent kinase [37-39]. Consequently,

phosphorylated N associates with 14-3-3 proteins in the cytoplasm [37]. Notably,

treatment with a kinase inhibitor cocktail eliminated the N/14-3-3 interaction, whereas

inhibition of 14-3-3θ expression by siRNA led to accumulation of N protein in the

nucleus [37]. These data suggest that 14-3-3 proteins directly shuttle SARS-CoV N

protein in a phosphorylation-dependent manner: a role which may be universal for N

proteins of all coronaviruses, including SARS-CoV-2. However, the molecular

mechanism of the 14-3-3/N interaction remains ill-defined.

14-3-3 proteins are amongst the top 1% of highest-expressed human proteins in

many tissues, with particular abundance in tissues vulnerable to SARS-CoV-2 infection

including the lungs, gastrointestinal system and brain [40, 41] (Fig. 1). 14-3-3 proteins

recognize hundreds of phosphorylated partner proteins involved in a magnitude of

cellular processes ranging from apoptosis to cytoskeleton rearrangements [42, 43].

Human 14-3-3 proteins are present in most of the tissues as seven conserved

"isoforms" (β, γ, ε, ζ, η, σ, τ/θ) (Fig. 1), with all-helical topology, forming dimers

possessing two identical antiparallel phosphopeptide-binding grooves, located at ~35 Å

distance from each other [44]. By recognizing phosphorylated Ser/Thr residues within

the structurally flexible (R/K)X$_{2-3}$(pS/pT)X(P/G) consensus motif [44, 45], 14-3-3 binding

is known to regulate the stability of partner proteins, their intracellular localization and

interaction with other factors [46]. In addition to their high abundance in many tissues

susceptible to SARS-CoV-2 infection (Fig. 1), 14-3-3 proteins were reported as one of

nine key host proteins during SARS-CoV-2 infection [47], indicating their potential

association with viral proteins.

In this work, we dissected the molecular mechanism of the interaction between

SARS-CoV-2 N and human 14-3-3 proteins. SARS-CoV-2 N protein containing several

phosphosites reported to occur during infection, was produced using the efficient

Escherichia coli system that proved successful for the study of polyphosphorylated

proteins [48]. We have observed the direct phosphorylation-dependent association

between polyphosphorylated SARS-CoV-2 N and all seven human 14-3-3 isoforms and

determined the affinity and stoichiometry of the interaction. Series of truncated mutants

of N localized the key 14-3-3-binding site to a single phosphopeptide residing in the

functionally important SR-rich region of N. These findings suggest a topology model for

the heterotetrameric 14-3-3/pN assembly occluding the SR-region, which presents a

feasible target for further characterization and therapeutic intervention.

Results

1.   Characterization of the polyphosphorylated N protein obtained by co-

expression with PKA in *E.coli*

Host-expressed SARS-CoV-2 N protein represents a phosphoprotein, harboring

multiple phosphorylation sites scattered throughout its sequence. The most densely

phosphorylated locus is the SR-rich region (Fig. 2A, Supplementary table 1,

Supplementary data file 1 and [13, 35]). Remarkably, this region is conserved in N

proteins of several coronaviruses [39, 49], including SARS-CoV (Fig. 2A). Although a

number of protein kinases have been implicated in SARS-CoV N phosphorylation [36,

37, 39, 50], the precise enzymes responsible for identified phosphosites and the

functional outcomes are largely unknown. Of note, many of the reported phosphosites

within the SARS-CoV-2 N protein are predicted to be phosphorylated by protein kinase

A (PKA), (Supplementary table 1). Hence, PKA was used for production of

phosphorylated SARS-CoV-2 N in *E.coli* [48], using the same approach that was

successfully applied for production of several phosphorylated eukaryotic proteins [48,

51-54] including the polyphosphorylated human tau competent for specific 14-3-3

binding [48].

Indeed, co-expression with PKA yielded a heavily phosphorylated SARS-CoV-2

N (Fig. 2B) containing more than 20 phosphosites according to LC-MS analysis

(Supplementary table 1 and Supplementary data file 1). Especially dense

phosphorylation occurred within the SR-rich region, involving recently reported in vivo

sites Ser23, Thr24, Ser180, Ser194, Ser197, Thr198, Ser201, Ser202, Thr205 and

Thr391 [13, 35] (Fig. 1A, C and Supplementary data file 1), implicating the success of

the PKA co-expression at emulating native phosphorylation. Interestingly, due to the

frequent occurrence of Arg residues, it was possible to characterize the

polyphosphorylation of the SR-rich region only with the use of an alternative protease

such as chymotrypsin, in addition to datasets obtained separately with trypsin (4

independent experiments overall, see Supplementary data file 1). Due to high

conservation between SARS-CoV and SARS-CoV-2 N proteins, many phosphosites

identified in the PKA-co-expressed N are likely shared by SARS-CoV N (Fig. 2A).

Importantly, many identified phosphosites lie within the regions predicted to be

disordered, and contribute to predicted 14-3-3-binding motifs, albeit deviating from the

optimal 14-3-3-binding sequence RXX(pS/pT)X(P/G) [44] (Fig. 2A and Supplementary

table 1).

Of note, the bacterially expressed SARS-CoV-2 N protein avidly binds random

E.coli nucleic acid, which results in a high 260/280 nm absorbance ratio in the eluate

from the nickel-affinity chromatography column. This is unchanged by

polyphosphorylation (Fig. 2D) and nucleic acid remained bound even after further

purification using heparin chromatography (data not shown). To quantitatively remove

nucleic acid from N preparations, we used continuous on-column washing of the His-

tagged protein with 3 M salt. This yielded clean protein preparations with the 260/280

nm absorbance ratio of 0.6 (Fig. 2D) of the unphosphorylated and polyphosphorylated N

(pN) with high electrophoretic homogeneity (Fig. 2E), enabling thorough investigation of

the 14-3-3 binding mechanism.

SEC-MALS suggested that the nucleic acid-free protein was a ~95 kDa dimer

(Fig. 2E), based on the calculated Mw of the His-tagged N monomer of 48 kDa,

regardless of its phosphorylation status. A significant overestimation of the apparent Mw

of pN from column calibration, ~160 kDa for the 95 kDa dimeric species, indicates

presence of elements with expanded loose conformation, in agreement with the

presence of unstructured regions. This necessitates the use of SEC-MALS for absolute

Mw determination that is independent of the assumptions on density and shape.

The skewed Mw distribution across the SEC peak indicated the propensity of the

SARS-CoV-2 N to oligomerization (Fig. 2E), which was instigated by the addition of

nucleic acid (Supplementary Fig. 1A and B). Using tRNA isolated from *E.coli* DH5□

cells, SEC and agarose gel electrophoresis, we showed that our nucleic acid-free,

polyphosphorylated N preparation is capable of avid binding of noncognate nucleic acid

(Supplementary Fig. 1A-C).

2.   Polyphosphorylated SARS-CoV-2 N and human 14-3-3γ form a tight

complex with defined stoichiometry

Next, we compared the ability of native full-length SARS-CoV-2 N, both

unphosphorylated and polyphosphorylated (N.1-419 and pN.1-419, respectively), to be

recognized by a human 14-3-3 protein. For the initial analysis we chose 14-3-3γ as one

of the strongest phosphopeptide binders among the 14-3-3 family [55].

SEC-MALS (Fig. 3A) shows that 14-3-3γ elutes as a dimer with Mw of ~55.2 kDa

(calculated monomer Mw 28.3 kDa, see also Fig. 3B). The position and amplitude of

this peak did not change in the presence of the N.1-419 dimer with Mw of 94.5 kDa

(calculated monomer Mw 48 kDa) where the SEC profile shows two distinct peaks. This

is corroborated by SDS-PAGE analysis, suggesting no interaction between

unphosphorylated N (pI >10) and 14-3-3 (pI ~4.5) despite the large difference in their pI

values (Fig. 3C,D). Thus, the presence of 200 mM NaCl was sufficient to prevent

nonspecific interactions between the two proteins.

In sharp contrast, co-expression of SARS-CoV-2 N with PKA, and subsequent

polyphosphorylation (Fig. 2), allows for tight complex formation between pN.1-419 and

human 14-3-3γ. This is evident from the peak shift and corresponding increased Mw

from ~95 to 150.7 kDa (Fig. 3E), perfectly matching the addition of the 14-3-3 dimer

mass (calculated dimer Mw 56.6 kDa) to the pN.1-419 dimer (calculated dimer Mw 96

kDa). The presence of both proteins in the complex was confirmed by SDS-PAGE (Fig.

3F). Collectively these data pointed toward the equimolar binding upon saturation.

Given the dimeric state of both proteins in their individual states (Fig. 2D and Fig. 3) and

the Mw of the 14-3-3γ/pN.1-419 complex, the most likely stoichiometry is 2:2. The ratio

of the proteins does not change across the peak of the complex (Fig. 3F), implying that

they form a relatively stable complex with the well-defined stoichiometry.

3. SARS-CoV-2 N interacts with all human 14-3-3 isoforms

We then questioned whether the interaction with pN is preserved for other human

14-3-3 isoforms. Analytical SEC clearly showed that the phosphorylated SARS-CoV-2 N

can be recognized by all seven human 14-3-3 isoforms, regardless of the presence of a

His-tag or disordered C-terminal tails on the corresponding 14-3-3 constructs (Fig. 4A).

However, the efficiency of complex formation differed for each isoform. Judging by the

repartition of 14-3-3 between free and the pN-bound peaks, the apparent efficiency of

pN binding was higher for 14-3-3γ, 14-3-3η, 14-3-3ζ and 14-3-3ε, and much lower for

14-3-3β, 14-3-3τ and 14-3-3σ, in a roughly descending order (Fig. 4A). The interaction

also appeared dependent on the oligomeric state of 14-3-3, since the monomeric

mutant form of 14-3-3ζ, 14-3-3ζm-S58E [56] (apparent Mw 29 kDa) showed virtually no

interaction relative to the wild-type dimeric 14-3-3ζ counterpart (apparent Mw 58 kDa),

(Fig. 4B).

This apparent variation in binding efficiency (Fig. 4A) justified more detailed

assessment of the binding parameters between pN.1-419 and selected 14-3-3 isoforms.

4. Affinity of the phosphorylated SARS-CoV-2 N towards selected human 14-3-3 isoforms

In light of the relative positions of the two proteins, separately and complexed, on

SEC profiles (Fig. 3 and 4), we used analytical SEC to track titration of a fixed

concentration of pN.1-419 (around 10 µM) against increasing quantities (0-100 µM) of

either of two selected full-length human 14-3-3 isoforms, γ and ε (Fig. 5A and B). A

saturation binding curve showed the maximal concentration of bound 14-3-3γ to

asymptotically approach 10 µM (Fig. 5C), supporting 2:2 stoichiometry. The apparent K_D

of the 14-3-3γ/pN.1-419 complex was estimated as 1.5 ± 0.3 µM. A similar binding

mechanism could be observed for 14-3-3ε, however in this case we could achieve pN.1-

419 saturation only at much higher 14-3-3ε concentrations (Fig. 5B and C), and the

resulting apparent K_D was ~7 times higher than for 14-3-3γ (Fig. 5C). Nevertheless,

once again the stoichiometry was close to 2:2. These findings strongly disfavor the

earlier hypothesis that 14-3-3 binding affects dimerization status of N [57].

We further asked what are the specific regions of SARS-CoV-2 N that are

responsible for interaction with human 14-3-3.

5. The N-terminal part of SARS-CoV-2 N is responsible for 14-3-3 binding

Among multiple phosphosites identified in our pN.1-419 preparations

(Supplementary table 1), the two most interesting regions are located in the intrinsically

disordered or loop segments, normally favored by 14-3-3 proteins [45]. The first

represents the C-terminally located RTA[pT^{265}]KAY site, which is predicted by the 14-3-

3-Pred webserver [58] as the 14-3-3-binding site. The second, SR-rich region features

multiple experimentally confirmed phosphosites including several suboptimal predicted

14-3-3-binding sites (Fig. 2A). To narrow down the 14-3-3-binding locus we used

several constructs representing its N- and C-terminal parts (N.1-211 and N.212-419,

respectively). The individual CTD included the C-terminal phosphosite around Thr265

(N.247-364), and the longer N-terminal construct extended toward the C-terminus to

include the predicted NES sequence (N.1-238) (Fig. 6A). This contains Asp225 and a

cluster of Leu residues which together resemble the unphosphorylated 14-3-3-binding

segments from ExoS/T [59] and therefore could be important for 14-3-3 binding.

As for the wild-type protein, the truncated SARS-CoV-2 N constructs were cloned

and expressed in the absence or presence of PKA to produce unphosphorylated or

polyphosphorylated proteins. PKA could be easily detected in the eluate from the Ni-

affinity column and typically led to an upward shift on SDS-PAGE and a shift on the

SEC profile (Supplementary Fig. 2). Each indicates efficient phosphorylation. Perhaps

with the exception of N.212-419, phosphorylation did not affect the oligomeric state of

the N constructs, and the observed shifts could be accounted for by the affected flexible

regions (Fig. 4 and Supplementary Fig. 2).

SEC-MALS (Supplementary Fig. 2) indicates the N.247-364 construct (calculated

monomer Mw 15.3 kDa), previously crystallized and kindly provided by Prof.

Baumeister's laboratory, exists as a ~30 kDa dimer. This is in perfect agreement with

earlier published data [28]. A dimeric state (Mw of 47.3 kDa) was also confirmed for the

longer C-terminal construct, N.212-419 (Supplementary Fig. 3, calculated monomer Mw

23.3 kDa), however, in this case a skewed Mw distribution indicative of the

polydispersity and tendency for oligomerization was observed (Supplementary Fig. 3).

In contrast, N-terminal constructs including the RNA-binding domain, such as

N.1-211, are monomeric (Supplementary Fig. 3). Thus, our data align with the low-

resolution structural model of SARS-CoV and SARS-CoV-2 N proteins, in which the

CTD (residues 247-364), largely responsible for dimerization, and NTD, involved in

RNA-binding, are only loosely associated [24, 28, 29].

Of the constructs analyzed, the N-terminal constructs N.1-211 and N.1-238 both

interacted with 14-3-3γ by forming distinct complexes on SEC profiles, and this

interaction was strictly phosphorylation-dependent (Fig. 6). Given the similarity of the

elution profiles for pN.1-211 and pN.1-238, it may be concluded that the presence of

NES in the latter is dispensable for the 14-3-3 binding.

Under similar conditions, only a very weak interaction between the dimeric

pN.212-419 construct and 14-3-3γ could be observed, whereas the phosphorylated

dimeric CTD (pN.247-364) displayed virtually no binding (Fig. 6). Neither

unphosphorylated construct interacted with 14-3-3. Thus, the Thr265 phosphosite can

be broadly excluded as the critical binding site. Separate phosphosites outside the CTD,

for instance, in the last ~30 C-terminal residues (Supplementary table 1) likely account

for residual binding of the pN.212-419 construct. Only its SEC profile showed a

significant positional peak shift with phosphorylation (Fig. 6B and C), indicating a

potential change in the oligomeric state. It is tempting to speculate that such

phosphorylation outside the CTD could affect higher order oligomerization associated

with the so-called N3 C-terminal segment (Fig. 2A) [20, 21, 49, 60].

According to SEC-MALS, the 14-3-3γ dimer (Mw 57 kDa) interacts with the pN.1-

238 monomer (Mw of 31 kDa) by forming a ~82 kDa complex (Fig. 7A) with an apparent

2:1 stoichiometry. It is remarkable that despite a moderate molar excess of pN.1-238 a

2:2 complex (one 14-3-3 dimer with two pN monomers) is not observed. The well-

defined 2:2 stoichiometry of the 14-3-3γ complex with the full-length pN (Fig. 3)

suggests that the dimeric pN is anchored using two equivalent, key 14-3-3-binding sites,

each located in a separate subunit of N. It is tempting to speculate that, in the absence

of the second subunit, the interaction involves the key phosphosite and an additional

phosphosite which is separated by a sufficiently long linker (≥ 15 residues [61]), to

secure occupation of both phosphopeptide-binding grooves of 14-3-3 (Fig. 7B). Such

bidentate binding would prevent the recruitment of a second pN.1-238 monomer and is

in line with the observed data. Similar 2:1 binding was observed qualitatively for the

interaction of 14-3-3γ with pN.1-211 (Fig. 6C and data not shown).

Our finding that the minimal N-terminal construct pN.1-211 exhibits firm binding

to 14-3-3γ indicates that the key 14-3-3-binding phosphosite(s) is/are located

exclusively within this region. Given the presence of numerous candidate 14-3-3-binding

sites within its most C-terminal part, i.e., the SR-rich region, we further focused on the

1-211 sequence in search of the 14-3-3-binding phosphosite(s).

6. Localization of the main 14-3-3-binding site within the SR-rich region of

SARS-CoV-2 N

Further N mutants were designed to disrupt the most probable 14-3-3-binding

phosphosites. These are located in the intrinsically flexible phosphorylatable SR-rich

segment centered at positions 197 and 205 (Fig. 8A). Both represent suboptimal 14-3-

3-binding motifs **S**RN[pS^{197}]T**P** and **S**RG[pT^{205}]S**P** in lacking an Arg/Lys residue in

position -3 (bold font) relative to the phosphorylation site (squared brackets). However,

each also features a Pro residue in position +2 (bold underlined font), which is highly

favorable for 14-3-3 binding [44] and absent from the other potential 14-3-3-binding

phosphosites in the SR-rich region (Fig. 2A). These conflicting factors complicate

predictions for the true 14-3-3 binding site. Moreover, even beyond the SR-rich region

the NTD is predicted to host further possible 14-3-3-binding phosphosites, including the

RRA[pT^{91}]RR site, which is the highest-scoring in 14-3-3-Pred [58] prediction

(Supplementary table 1).

We conceived stepwise truncations to remove the most probable 14-3-3-binding

phosphosites, aiming to identify the iteration at which binding (observed for the pN.1-

211 construct) ceased. 14-3-3 can bind incomplete consensus motifs at the extreme C-

terminus of some proteins [62], so truncations were designed to remove the critical

phosphorylated residue. However, upstream residues of each candidate 14-3-3-binding

site were preserved, in light of the sheer number of overlapping potential binding motifs

in the SR-rich region (Fig. 8A).

The new truncated constructs of N, namely N.1-204, N.1-196 and N.1-179, were

obtained in unphosphorylated and phosphorylated states, as before, and again washed

with high salt to remove potentially disruptive nucleic acid. SEC-MALS confirmed the

monomeric state of the N-terminal N mutants (the exemplary data are presented for

N.1-196, Supplementary Fig. 3), consistent with the proposed architecture of the N

protein [21, 24].

None of the truncated mutants interacted with 14-3-3γ in the unphosphorylated state

(Fig. 8B). No binding to 14-3-3γ was detected with phosphorylated N.1-179 (Fig. 8C)

and only very limited binding could be observed with phosphorylated N.1-196 (Fig. 8C).

This strongly indicated that all phosphosites of the 1-196 segment (including at least

three phosphosites within the SR-rich region, i.e., Ser180, Ser188 and Ser194, see Fig.

8A) are dispensable for 14-3-3 binding and at most could contribute only as auxiliary

sites (as suggested by the scheme in Fig. 7B). This narrowed the 14-3-3-binding region

within SARS-CoV-2 N down to 15 residues from 196 to 211, leaving only two possible

sites centered at Ser197 and Thr205 (Fig. 8A).

By contrast, pN.1-204 showed a similar interaction with 14-3-3γ to that of pN.1-211

(Fig. 8C). Although we cannot fully exclude that Thr205 phosphosite contributes to 14-3-

3 binding in the context of the full-length pN, it is clearly not critical for 14-3-3

recruitment, unlike Ser197. Moreover, in contrast to Thr205, Ser197 is preserved in

most related coronavirus N proteins (see Fig. 2A, Fig. 9).

354 Discussion

In this work, we investigated the molecular association between the SARS-CoV-2

N protein and human phosphopeptide-binding proteins of the 14-3-3 family. The former

is the most abundant viral protein [12, 13], the latter is a major protein-protein

interaction hub involved in multiple cellular signaling cascades, expressed at high levels

in many human tissues including those susceptible to SARS-CoV-2 infection (Fig. 1)

[40]. SARS-CoV-2 N is heavily phosphorylated in infected cells [13, 35, 36], which

poses a significant challenge for proteomic approaches: the densely phosphorylated

SR-rich region alone, functionally implicated in numerous viral processes [34, 49, 63],

hosts seven closely spaced Arg residues (Fig. 2A). These arginines restrict the length of

tryptic phosphopeptides and decrease the probability of their unambiguous identification

and phosphosite assignment [64]. The multiplicity of implicated protein kinases

(including GSK-3, protein kinase C, casein kinase II and mitogen-activated protein

kinase [36, 37, 50]) further hinders study of specific phosphorylations. Assuming that

the mechanistic implication of a specific phosphorylation is independent of the acting

kinase, we produced polyphosphorylated N protein (pN) via bacterial co-expression with

a catalytic subunit of PKA. A combination of orthogonal cleavage enzymes and LC-MS

phosphoproteomics mapped > 20 phosphosites (Supplementary table 1) including

Ser23, Thr24, Ser180, Ser194, Ser197, Thr198, Ser201, Ser202, Thr205 and Thr391

reported recently at SARS-CoV-2 infection [13, 35]. At least six of the identified

phosphosites are located in the unstructured regions and represent potential 14-3-3-

binding sites (Fig. 2A).

Biochemical analysis confirmed that polyphosphorylated N is competent for

binding to all seven human 14-3-3 isoforms (Fig. 3 and 4), but revealed remarkable

variation in binding efficiency between them (Fig. 4). This was supported by the

quantified affinities to two selected isoforms, 14-3-3γ (K_D of 1.5 µM) and 14-3-3ε (K_D of

10.7 µM) (Fig. 5). Our observations are in line with the recent finding that 14-3-3γ and

14-3-3η systematically bind phosphopeptides with higher affinities than 14-3-3ε and 14-

3-3σ [55]. The low micromolar-range K_D values compare well to those reported for other

physiologically relevant partners of 14-3-3 [65-67], indicating a stable and specific

interaction. Meanwhile, the well-defined 2:2 stoichiometry of the ~150-kDa 14-3-3/pN

complex, supported by titration experiments and SEC-MALS analysis (Fig. 3 and 5),

excludes the possibility that 14-3-3 binding disrupts pN dimerization [57]. It is

reasonable to assume that the principally bivalent 14-3-3 dimer [44, 46] recognizes just

one phosphosite in each pN subunit because a bidentate 14-3-3 binding to different

phosphosites within a single pN subunit would inevitably alter the observed 2:2

stoichiometry. Identification of the single phosphosite responsible for 14-3-3 recruitment

proved challenging, as none of the potential sites were a perfect match to the currently

known optimal 14-3-3-binding motifs [65].

To restrict the search, we analyzed the interaction of various N constructs with

14-3-3γ. This eliminated the high-scoring potential 14-3-3-binding phosphosite

RTApT^{265}KAY, present in the C-terminal N fragments, as the true site of interaction

despite its conservation in many related coronavirus N proteins (Supplementary table 2

and 3). The residual binding of pN.212-419 to 14-3-3γ suggested the existence of

auxiliary 14-3-3-binding sites located outside the folded CTD (residues 247-364). Both

pN.1-211 and pN.1-238 bound 14-3-3 with equivalent efficiency suggesting the binding

lies between amino acid 1 and 211. Interestingly, the N-terminal constructs existed as

monomers (Supplementary Fig. 3), which could potentially lower the binding affinity to

14-3-3 dimers in light of the 2:2 stoichiometry. Nonetheless, sufficient binding was

clearly observed between the dimeric 14-3-3γ and the monomeric N-terminal constructs

(Fig. 6 and 7).

Truncation of the SR-rich region streamlined the search for the 14-3-3-binding

site to the 15-residue stretch of amino acids 196-211. This sequence hosts two

principally similar potential 14-3-3-binding sites, RNpS^{197}TP and RGpT^{205}SP (Fig. 2A

and C). Importantly, the proximity of these sites rules out bidentate binding to the 14-3-3

dimer: 14-3-3 binds in an antiparallel manner requiring a minimum of 13-15 residues

between phosphosites on a single peptide [44, 68]. Thus, binding to Ser197 and Thr205

sites must be mutually exclusive.

The markedly different binding between pN.1-204 and pN.1-196 to 14-3-3 (Fig. 8)

prompted us to propose Ser197 as the critical phosphosite. This finding aided the

design of a topology model for the complex (Fig. 9A), in which the 14-3-3 dimer is

anchored by two identical Ser197 phosphosites from the SR-rich region in the two

equivalent pN chains. Noteworthily, the RN(pS/pT)[197]TP site is conserved in not only

SARS-CoV and SARS-CoV-2 but also in N proteins from several bat and pangolin

coronaviruses (Fig. 9B). Meanwhile, plausible phosphorylation of residue 205 is

possible in a smaller subset of coronaviruses (Fig. 9B).

The model shown in Fig. 9A notably does not exclude the possibility of 14-3-3

binding to alternative phosphosites under significantly different phosphorylation

conditions. Moreover, we speculate that hierarchical phosphorylation within the SR-rich

region [36] would alter 14-3-3 binding with phosphorylation at adjacent Ser/Thr

residues, likely to inhibit the interaction [69]. The SR-rich region is phosphorylated by

both Pro-directed and non-Pro-directed protein kinases [36, 37, 39, 50]. Since 14-3-3

proteins typically reject peptides with a proline adjacent to the phosphorylated residue,

the interplay between these aforementioned kinases could be regulatory. In theory, this

could create a phosphorylation code and conditional binding of 14-3-3, as has been

discussed recently for alternative polyphosphorylated 14-3-3 partners such as LRRK2,

CFTR and tau protein [48, 67, 70, 71].

The recruitment of the 14-3-3 dimer is expected to occlude the SR-rich region of

N by masking 10-20 residues surrounding the Ser197 phosphosite within the complex.

Apart from the likely effects on the properties of N and its ability to phase separate and

bind RNA, the 14-3-3 binding at the SR-rich region, triggered by its phosphorylation, can

potentially interfere with N binding to the M protein, an event clearly relevant to the

virion assembly [17]. In support of this hypothesis, the 14-3-3-occluded area reported

here overlaps with the N region (residues 168-208) proposed to mediate its association

with the M protein in SARS-CoV [19].

The presence of SR-rich regions in many viral N proteins suggests a more broad

interaction of 14-3-3 with N proteins. Indeed, using 14-3-3-Pred prediction [58] a

number of 14-3-3-binding phosphosites in N proteins from human (Supplementary table

2) and bat coronaviruses (Supplementary table 3) could be identified. Some display

strong conservation of the 14-3-3-binding binding site around Ser197 (Fig. 9B), whereas

others contain separate high-scoring potential 14-3-3-binding sites beyond the SR-rich

region. Given the reasonable threat that other zoonotic coronaviruses may ultimately

enter the human population [72, 73], the relevant N proteins are highly likely to undergo

phosphorylation and 14-3-3 binding, as seen for SARS-CoV-2 N. This is particularly

likely, given the high concentration of both proteins in the infected cells (see above).

Our findings underline the essential role of the SR-rich region in the biology of N

proteins and host-virus interactions [49]. Unrelated proteins with similar domains also

tend to show RNA-binding capability (e.g., the splicing factors) [74, 75] and are subject

to multisite phosphorylation [76]. Such proteins are often associated with phase

separation as a means to regulate membraneless compartmentalization within the

cytoplasm. Likewise, SARS-CoV-2 N protein has been shown to undergo phase

separation in vitro upon RNA addition: a phenomenon dependent on the concentration

of salt, presence of divalent ions, phosphorylation state of N and on RNA sequence [30,

34, 77, 78]. Furthermore, the N protein has been shown to recruit the RNA-dependent

RNA-polymerase complex, and granule-associated heterogeneous nuclear

ribonucleoproteins forming phase separated granules which can aid SARS-CoV-2

replication [30, 34]. Thus 14-3-3 potentially influences phase separation by binding the

SR-rich region and may also affect the accessibility of the NES sequence located

nearby (Fig. 2A). This in turn may impact nucleocytoplasmic shuttling of N, as seen for

SARS-CoV N [37].

Conclusively, 14-3-3 binding to the SR-rich region of N holds potential to regulate

multiple cell host processes affected by N. 14-3-3 binding to pN may present a cell

immune-like response to the viral infection aimed at arresting or neutralizing N activities

[57]. On the other hand, in light of the abundance of N protein in the infected cell [12,

13], pN may instead arrest 14-3-3 proteins in the cytoplasm and indirectly disrupt

cellular processes involving 14-3-3. For example, 14-3-3ε and 14-3-3η each play a role

in the innate immune response via RIG-1 and MDA5 signaling respectively [79, 80]. The

N protein:14-3-3 interaction would modulate these and other signaling pathways

involving 14-3-3 proteins. As such, understanding the molecular mechanism of pN

association with 14-3-3 proteins may inform the development of novel therapeutic

approaches.

Materials and methods

Protein expression and purification

The tagless SARS-CoV-2 N gene coding for Uniprot ID P0DTC9 protein

sequence was commercially synthesized (GeneWiz) and cloned into pET-SUMO

bacterial expression vectors using *Nde*I and *Hind*III restriction endonuclease sites. To

obtain constructs of N carrying the N-terminal His$_6$-tag cleavable by HRV-3C protease

(N.1-419, N.1-211, N.1-238, N.212-419), the N gene was PCR amplified using primers

listed in Supplementary table 4 and cloned into the pET-YBSL-Lic-3C plasmid vector

using an established ligation-independent cloning procedure [81]. After 3C cleavage

each construct contained extra GPA residues at the N terminus. The N.247-364

construct carrying an N-terminal uncleavable His$_6$-tag and corresponding to the

previously crystallized folded CTD dimer [28], was kindly provided by Prof. W.

Baumeister's laboratory.

Truncated mutants of N.1-211, namely N.1-179, N.1-196 and N.1-204 were

derived from the wild-type N plasmid using the standard T7 forward primer with varying

reverse primers listed in Supplementary table 4. Each carried a *BamH*I site to enable

consequent cloning into a pET28 vector using *Nco*I and *BamH*I sites. This preserved

the N-terminal His$_6$-tag cleavable by 3C protease. All constructs were validated by DNA

sequencing.

Untagged full-length human 14-3-3γ (Uniprot ID P61981) and 14-3-3ζ (Uniprot ID

P63104) cloned in a pET21 vector, and full-length human 14-3-3ε (Uniprot ID P62258)

carrying an N-terminal His$_6$-tag cleavable by Tobacco Etch Virus (TEV) protease, and

cloned into a pRSET-A vector have been previously described [82-84]. Truncated

versions of human 14-3-3η (Uniprot ID Q04917), 14-3-3β (Uniprot ID P31946), 14-3-3ε

(Uniprot ID P62258), 14-3-3τ (Uniprot ID P27348), 14-3-3γ (Uniprot ID P61981) and 14-

3-3σ (Uniprot ID P31947) devoid of the short disordered segment at the C terminus,

cloned into pProExHtb vector and carrying TEV-cleavable His$_6$-tags on their N-terminal

ends were obtained as described previously [66, 85]. The monomeric mutant form of

untagged full-length human 14-3-3ζ carrying monomerizing amino acid substitutions

and pseudophosphorylation in the subunit interface (14-3-3ζm-S58E) was obtained as

before [56].

14-3-3γ, 14-3-3ζ and 14-3-3ζm-S58E were expressed and purified using

ammonium sulfate fractionation and column chromatography as previously described

[82]. His$_6$-tagged N.247-364 was also expressed and purified as before [28]. All other

proteins carrying cleavable His$_6$-tags were expressed in *E.coli* BL21(DE3) cells and

purified using subtractive immobilized metal-affinity chromatography (IMAC) and gel-

filtration. For the N protein and its various constructs the eluate from the first IMAC

column, performed at 1 M NaCl, harbored a significant quantity of random bound

nucleic acid (the 260/280 nm absorbance ratio of 1.7-2.0). This necessitated an

additional long on-column washing step with 3 M NaCl (50 column volumes) to ensure

the 260/280 nm absorbance ratio of ~0.6 and removal of all nucleic acid. Tagless N

protein was purified after treatment of cell lysate with RNAse, using heparin and

subsequent ion-exchange chromatography on a sulfopropyl Sepharose (elution

gradients 100-1000 mM NaCl) followed by gel-filtration. Here, It proved challenging to

entirely eliminate nucleic acid and the typical 260/280 nm absorbance ratio was 0.7-0.9.

Protein concentration was determined by spectrophotometry at 280 nm on a N80

Nanophotometer (Implen, Munich, Germany) using sequence-specific extinction

coefficients calculated using the ProtParam tool in ExPASy (see Supplementary table

5).

Isolation of E.coli tRNA

The DH5◻ cells were incubated in 30 ml of liquid medium (LB without antibiotics)

for 16 h at 37 °C with maximum aeration and then harvested by centrifugation at 7000 g

for 10 min. The pellet was gently resuspended in RNAse-free Tris-acetate buffer,

followed by alkali-SDS lysis and neutralization by cold ammonium acetate. The

suspension was incubated on ice for 5 min and centrifuged at 21000 g at 4 °C for 10

min. The resulting supernatant (8 ml) was incubated with 12.5 ml isopropanol for 15 min

at 25 °C and centrifuged at 12100 g for 5 min. The pellet was resuspended in 800 µl of

2 M ammonium acetate, incubated for 5 min at 25 °C and centrifuged (12100 g, 10 min).

RNA was precipitated from supernatant by 800 µl of isopropanol, incubated for 5 min at

25 °C and centrifuged (12100 g, 5 min). After supernatant removal the pellet was

washed with 70% ice-cold ethanol, dried and dissolved in 100-200 µl milliQ-water.

SARS-CoV-2 N protein co-expression with PKA

For phosphorylation in cells, SARS-CoV-2 N was bacterially co-expressed with a

catalytic subunit of mouse protein kinase A (PKA), as described previously [48]. PKA

was cloned into a low-copy pACYC vector [48] which ensured that the target protein

was expressed in excess of kinase.

PKA and SARS-CoV-2 N were co-transformed into *E.coli* BL21(DE3) cells

against Kanamycin and Chloramphenicol resistance. Cells were grown in LB to an

OD_{600} reading of 0.6 before inducing with 500 µM of IPTG. After induction, cultivation

was continued for 1.5 h, 3 h, 4 h and overnight at 37 ◻C, and 4 h was found to be

sufficient to provide for the saturating interaction with 14-3-3. For all truncated

constructs of N, overnight co-expression was used. For unphosphorylated controls we

expressed proteins in the absence of PKA.

The cells with overexpressed proteins were harvested by centrifugation and

resuspended in 20 mM Tris-HCl buffer pH 8.0 containing 1 M NaCl, 10 mM imidazole,

0.01 mM phenylmethylsulfonyl fluoride, as well as trace amounts of lysozyme to

facilitate lysis. Phosphorylated proteins were purified using subtractive IMAC and gel-

filtration, whereby the His$_6$-tagged PKA was efficiently removed. Phosphorylated N and

its constructs typically showed significant shifts on SDS-PAGE and PhosTag gels

indicating phosphorylation.

Identification of phosphosites within N

Sample treatment for proteomics analysis. For phosphopeptide mapping, the

SARS-CoV-2 N protein co-expressed with PKA for 4 h at 30 □C was purified as above.

An aliquot (35 µg) was subjected to enzymatic hydrolysis "in solution" either with trypsin

(Sequencing Grade Modified Trypsin, Promega) or with chymotrypsin (Analytical Grade,

Sigma-Aldrich). Briefly, the sample preparation was as follows. The sample was

reduced with 2 mM Tris(2-carboxyethyl)phosphine (TCEP) and then alkylated with 4

mM S-Methyl methanethiosulfonate (MMTS); a protein:enzyme ratio was kept at 50:1

(w/w); digestion was performed overnight at 37 □C and pH 7.8 (for chymotrypsin, 10

mM Ca^{2+} was added to the reaction solution). The reaction was stopped by adjusting it

to pH 2 with formic acid. The resulting peptides were purified on a custom micro-tip SPE

column with Oasis HLB (Waters) as a stationary phase, using elution with an

acetonitrile:water:formic acid mix (50:49.9:0.1 v/v/v%). The eluate of resulting peptides

was dried out and stored at -30 □C prior to the LC-MS experiment.

LC-MS/MS experiment. Peptides were separated on a nano-flow

chromatographic system with a flow rate of 440 nl/min (Ultimate 3000 Nano RSLC,

Thermo Fisher Scientific) and ESI coupled to a mass-analyzer (Q Exactive Plus,

Thermo Fisher Scientific). Briefly, the protocol was as follows. Dried peptide samples

were rehydrated with 0.1% formic acid. An aliquot of rehydrated peptides (5 µl) was

injected onto a precolumn (100 µm x 2 cm, Reprosil-Pur C18-AQ 1.9 µm, Dr. Maisch

GmbH) and eluted on a nano-column (100 µm x 30 cm, Reprosil-Pur C18-AQ 1.9 µm,

Dr. Maisch GmbH) with a linear gradient of mobile phase A (water:formic acid 99.9:0.1

v/v%) with mobile phase B (acetonitrile:water:formic acid 80:19.9:0.1 v/v/v%) from 2%

till 80% B within 80 min. MS data were acquired by a data-dependent acquisition

approach after a MS-scan of a 350–1500 m/z range. Top 12 peaks were subjected to

high collision energy dissociation (HCD) or electron transfer dissociation (ETD). After a

round of MS/MS, masses were dynamically excluded from further analysis for 35 s.

MS data analysis. For Mascot (MatrixScience) database search, raw LC-MS

files were converted to general mgf format using MSConvert with its default settings.

For peptide database search, a concatenated database including SARS-CoV-2 and

general contaminants was constructed. The parameters of the search were: Enzyme –

Trypsin with one miscleavage allowed, or in case of Chymotrypsin - no enzyme; MS

tolerance – 5 ppm; MSMS – 0.2 Da; Fixed modifications – Methylthio(C); Variable

modifications – Phospho(ST), Phospho(Y), Oxidation(M). For PEAKS (PEAKS Studio

Xpro, Bioinformatics Solutions Inc.) analysis the parameters of a search were MS

tolerance - 10 ppm, MSMS – 0.05 Da; Fixed modifications – Methylthio(C); Variable

modifications – Phospho(STY), Oxidation(M), Ammonia-loss (N), Deamidation (NQ).

For estimation of FDR, a target-decoy approach was used.

Analytical size-exclusion chromatography (SEC)

The oligomeric state of proteins as well as protein-protein and protein-tRNA

interactions were analyzed by loading 50 µl samples on a Superdex 200 Increase 5/150

column (GE Healthcare) operated at a 0.45 ml/min flow rate using a Varian ProStar 335

system (Varian Inc., Melbourne, Australia). When specified, a Superdex 200 Increase

10/300 column (GE Healthcare) operated at a 0.8 ml/min flow rate was used (100 µl

loading). The columns were equilibrated by a 20 mM Tris-HCl buffer, pH 7.6, containing

200 mM NaCl and 3 mM NaN_3 (SEC buffer) and calibrated by the following protein

markers: BSA trimer (198 kDa), BSA dimer (132 kDa), BSA monomer (66 kDa),

ovalbumin (43 kDa), α-lactalbumin (15 kDa). The profiles were followed by 280 nm and

(optionally) at 260 nm absorbance. Diode array detector data were used to retrieve full-

absorbance spectral information about the eluted samples including protein and nucleic

acid. All SEC experiments were performed at least three times and the most typical

results are presented.

To assess binding parameters for the 14-3-3/pN.1-419 interaction, we used serial

loading of the samples containing a fixed pN concentration and increasing

concentrations of 14-3-3 in a constant volume of 50 µl and a Superdex 200 Increase

5/150 column operated at 0.45 ml/min. To validate linear augmentation of the peak

amplitude with increasing protein concentration, serial loading of 14-3-3 alone at

different concentrations was used. Such linear dependence allowed conversion of the

amplitude of the 14-3-3 peak into the molar concentration of its unbound form at each

point of titration. Concentration of pN-bound 14-3-3 was determined as the difference

between the total and free concentration. Binding curves representing the dependence

of bound 14-3-3 on its total concentration were approximated using the quadratic

equation to determine apparent K_D values. Graphing and fitting were performed in

Origin 9.0 (OriginLab Corporation, Northampton, MA, USA).

Size-exclusion chromatography coupled to multi-angle light scattering (SEC-MALS)

To determine the absolute masses of various N constructs and their complexes

with 14-3-3, we coupled a SEC column to a ProStar 335 UV/Vis detector (Varian Inc.,

Melbourne, Australia) and a multi-angle laser light scattering detector miniDAWN (Wyatt

Technologies). Either Superdex 200 Increase 10/300 (~24 ml, flow rate 0.8 ml/min) or

Superdex 200 Increase 5/150 (~3 ml, flow rate 0.45 ml/min) columns (GE Healthcare)

were used. The miniDAWN detector was calibrated relative to the scattering from

toluene and, together with concentration estimates obtained from UV detector at 280

nm, was exploited for determining the Mw distribution of the eluted protein species. All

processing was performed in ASTRA 8.0 software (Wyatt Technologies) taking dn/dc

equal to 0.185 and using extinction coefficients listed in Supplementary table 5. Protein

content in the eluted peaks was additionally analyzed by SDS-PAGE.

Data availability

The source LC-MS data on SARS-CoV-2 N phosphoproteomics are available

along with the paper as a Supplementary data file 1.

Acknowledgements

N.N.S. is thankful to Prof. W. Baumeister and Dr. L. Zinzula (Martinsried,

Germany) for providing the N.247-364 plasmid. MALDI and MALS were carried out at the Shared-Access Equipment Centre "Industrial Biotechnology" of the Federal Research Center "Fundamentals of Biotechnology" of the Russian Academy of Sciences. This work was supported by the Russian Science Foundation (grant 19-74-10031 to N.N.S.) and the Wellcome Trust (206377 award to A.A.A.). Protein-protein interactions were studied in the framework of the Program of the Russian Ministry of Science and Higher Education (K.V.T. and N.N.S.). O.I.K. acknowledges support by the Interdisciplinary Scientific and Educational School of Moscow University «Molecular Technologies of the Living Systems and Synthetic Biology». The authors declare no competing interests.

References

[1] World-Health-Organization. WHO Coronavirus Disease (COVID-19) Dashboard, https://covid19.who.int. 2020.

[2] Guarner J. Three Emerging Coronaviruses in Two Decades. Am J Clin Pathol. 2020;153:420-1.

[3] Masood N, Malik SS, Raja MN, Mubarik S, Yu C. Unraveling the Epidemiology, Geographical Distribution, and Genomic Evolution of Potentially Lethal Coronaviruses (SARS, MERS, and SARS CoV-2). Frontiers in cellular and infection microbiology. 2020;10:499.

[4] Gussow AB, Auslander N, Faure G, Wolf YI, Zhang F, Koonin EV. Genomic determinants of pathogenicity in SARS-CoV-2 and other human coronaviruses. Proceedings of the National Academy of Sciences. 2020;117:15193-9.

[5] Zamorano Cuervo N, Grandvaux N. ACE2: Evidence of role as entry receptor for SARS-CoV-2 and implications in comorbidities. eLife. 2020;9.

[6] Trypsteen W, Van Cleemput J, Snippenberg WV, Gerlo S, Vandekerckhove L. On the

whereabouts of SARS-CoV-2 in the human body: A systematic review. PLoS Pathog.

2020;16:e1009037.

[7] van Doremalen N, Lambe T, Spencer A, Belij-Rammerstorfer S, Purushotham JN, Port JR,

et al. ChAdOx1 nCoV-19 vaccination prevents SARS-CoV-2 pneumonia in rhesus macaques.

bioRxiv. 2020.

[8] Corbett KS, Flynn B, Foulds KE, Francica JR, Boyoglu-Barnum S, Werner AP, et al.

Evaluation of the mRNA-1273 Vaccine against SARS-CoV-2 in Nonhuman Primates. N Engl J

Med. 2020;383:1544-55.

[9] Mulligan MJ, Lyke KE, Kitchin N, Absalon J, Gurtman A, Lockhart S, et al. Phase I/II study of

COVID-19 RNA vaccine BNT162b1 in adults. Nature. 2020;586:589-93.

[10] Twomey JD, Luo S, Dean AQ, Bozza WP, Nalli A, Zhang B. COVID-19 update: The race to

therapeutic development. Drug resistance updates : reviews and commentaries in antimicrobial

and anticancer chemotherapy. 2020;53:100733.

[11] Marra MA, Jones SJ, Astell CR, Holt RA, Brooks-Wilson A, Butterfield YS, et al. The

Genome sequence of the SARS-associated coronavirus. Science. 2003;300:1399-404.

[12] Bojkova D, Klann K, Koch B, Widera M, Krause D, Ciesek S, et al. Proteomics of SARS-

CoV-2-infected host cells reveals therapy targets. Nature. 2020;583:469-72.

[13] Bouhaddou M, Memon D, Meyer B, White KM, Rezelj VV, Correa Marrero M, et al. The

Global Phosphorylation Landscape of SARS-CoV-2 Infection. Cell. 2020;182:685-712 e19.

[14] Sender R, Bar-On YM, Flamholz A, Gleizer S, Bernsthein B, Phillips R, et al. The total

number and mass of SARS-CoV-2 virions in an infected person. medRxiv.

2020:2020.11.16.20232009.

[15] Bar-On YM, Flamholz A, Phillips R, Milo R. SARS-CoV-2 (COVID-19) by the numbers.

eLife. 2020;9.

[16] Milo R. What is the total number of protein molecules per cell volume? A call to rethink

some published values. Bioessays. 2013;35:1050-5.

[17] McBride R, van Zyl M, Fielding BC. The coronavirus nucleocapsid is a multifunctional

protein. Viruses. 2014;6:2991-3018.

[18] Yao H, Song Y, Chen Y, Wu N, Xu J, Sun C, et al. Molecular Architecture of the SARS-

CoV-2 Virus. Cell. 2020;183:730-8 e13.

[19] He R, Leeson A, Ballantine M, Andonov A, Baker L, Dobie F, et al. Characterization of

protein-protein interactions between the nucleocapsid protein and membrane protein of the

SARS coronavirus. Virus research. 2004;105:121-5.

[20] Chang CK, Chen CM, Chiang MH, Hsu YL, Huang TH. Transient oligomerization of the

SARS-CoV N protein--implication for virus ribonucleoprotein packaging. PLoS One.

2013;8:e65045.

[21] Ye Q, West AMV, Silletti S, Corbett KD. Architecture and self-assembly of the SARS-CoV-2

nucleocapsid protein. Protein Sci. 2020;29:1890-901.

[22] Cubuk J, Alston JJ, Incicco JJ, Singh S, Stuchell-Brereton MD, Ward MD, et al. The SARS-

CoV-2 nucleocapsid protein is dynamic, disordered, and phase separates with RNA. bioRxiv.

2020.

[23] Chen H, Cui Y, Han X, Hu W, Sun M, Zhang Y, et al. Liquid-liquid phase separation by

SARS-CoV-2 nucleocapsid protein and RNA. Cell Res. 2020.

[24] Zeng W, Liu G, Ma H, Zhao D, Yang Y, Liu M, et al. Biochemical characterization of SARS-

CoV-2 nucleocapsid protein. Biochem Biophys Res Commun. 2020;527:618-23.

[25] Kang S, Yang M, Hong Z, Zhang L, Huang Z, Chen X, et al. Crystal structure of SARS-

CoV-2 nucleocapsid protein RNA binding domain reveals potential unique drug targeting sites.

Acta pharmaceutica Sinica B. 2020;10:1228-38.

[26] Khan A, Tahir Khan M, Saleem S, Junaid M, Ali A, Shujait Ali S, et al. Structural insights

into the mechanism of RNA recognition by the N-terminal RNA-binding domain of the SARS-

CoV-2 nucleocapsid phosphoprotein. Computational and structural biotechnology journal.

2020;18:2174-84.

[27] Chen CY, Chang CK, Chang YW, Sue SC, Bai HI, Riang L, et al. Structure of the SARS

coronavirus nucleocapsid protein RNA-binding dimerization domain suggests a mechanism for

helical packaging of viral RNA. J Mol Biol. 2007;368:1075-86.

[28] Zinzula L, Basquin J, Bohn S, Beck F, Klumpe S, Pfeifer G, et al. High-resolution structure

and biophysical characterization of the nucleocapsid phosphoprotein dimerization domain from

the Covid-19 severe acute respiratory syndrome coronavirus 2. Biochem Biophys Res

Commun. 2020.

[29] Zhou R, Zeng R, von Brunn A, Lei J. Structural characterization of the C-terminal domain of

SARS-CoV-2 nucleocapsid protein. Molecular Biomedicine. 2020;1:2.

[30] Perdikari TM, Murthy AC, Ryan VH, Watters S, Naik MT, Fawzi NL. SARS-CoV-2

nucleocapsid protein phase-separates with RNA and with human hnRNPs. EMBO J.

2020:e106478.

[31] Cascarina SM, Ross ED. A proposed role for the SARS-CoV-2 nucleocapsid protein in the

formation and regulation of biomolecular condensates. FASEB J. 2020;34:9832-42.

[32] Laude H, Masters PS. The Coronavirus Nucleocapsid Protein. In: Siddell SG, editor. The

Coronaviridae. Boston, MA: Springer US; 1995. p. 141-63.

[33] Masters PS. Coronavirus genomic RNA packaging. Virology. 2019;537:198-207.

[34] Savastano A, Ibanez de Opakua A, Rankovic M, Zweckstetter M. Nucleocapsid protein of

SARS-CoV-2 phase separates into RNA-rich polymerase-containing condensates. Nat

Commun. 2020;11:6041.

[35] Davidson AD, Williamson MK, Lewis S, Shoemark D, Carroll MW, Heesom KJ, et al.

Characterisation of the transcriptome and proteome of SARS-CoV-2 reveals a cell passage

induced in-frame deletion of the furin-like cleavage site from the spike glycoprotein. Genome

medicine. 2020;12:68.

[36] Yaron TM, Heaton BE, Levy TM, Johnson JL, Jordan TX, Cohen BM, et al. The FDA-

approved drug Alectinib compromises SARS-CoV-2 nucleocapsid phosphorylation and inhibits

viral infection in vitro. bioRxiv. 2020:2020.08.14.251207.

[37] Surjit M, Kumar R, Mishra RN, Reddy MK, Chow VT, Lal SK. The severe acute respiratory

syndrome coronavirus nucleocapsid protein is phosphorylated and localizes in the cytoplasm by

14-3-3-mediated translocation. J Virol. 2005;79:11476-86.

[38] Rana AK, Rahmatkar SN, Kumar A, Singh D. Glycogen synthase kinase-3: A putative

target to combat severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) pandemic.

Cytokine & growth factor reviews. 2020.

[39] Peng TY, Lee KR, Tarn WY. Phosphorylation of the arginine/serine dipeptide-rich motif of

the severe acute respiratory syndrome coronavirus nucleocapsid protein modulates its

multimerization, translation inhibitory activity and cellular localization. FEBS J. 2008;275:4152-

63.

[40] Wang M, Weiss M, Simonovic M, Haertinger G, Schrimpf SP, Hengartner MO, et al. PaxDb,

a database of protein abundance averages across all three domains of life. Mol Cell Proteomics.

2012;11:492-500.

[41] Boston PF, Jackson P, Thompson RJ. Human 14-3-3 protein: radioimmunoassay, tissue

distribution, and cerebrospinal fluid levels in patients with neurological disorders. J Neurochem.

1982;38:1475-82.

[42] van Hemert MJ, Steensma HY, van Heusden GP. 14-3-3 proteins: key regulators of cell

division, signalling and apoptosis. Bioessays. 2001;23:936-46.

[43] Aitken A. 14-3-3 proteins: a historic overview. Semin Canc Biol. 2006;16:162-72.

[44] Yaffe MB, Rittinger K, Volinia S, Caron PR, Aitken A, Leffers H, et al. The structural basis

for 14-3-3:phosphopeptide binding specificity. Cell. 1997;91:961-71.

[45] Bustos DM, Iglesias AA. Intrinsic disorder is a key characteristic in partners that bind 14-3-3

proteins. Proteins. 2006;63:35-42.

[46] Obsil T, Obsilova V. Structural basis of 14-3-3 protein functions. Semin Cell Dev Biol.

2011;22:663-72.

[47] Messina F, Giombini E, Montaldo C, Sharma AA, Piacentini M, Zoccoli A, et al. Looking for

pathways related to COVID-19 phenotypes: Confirmation of pathogenic mechanisms by SARS-

CoV-2 - Host interactome. bioRxiv. 2020:2020.11.03.366666.

[48] Tugaeva KV, Tsvetkov PO, Sluchanko NN. Bacterial co-expression of human Tau protein

with protein kinase A and 14-3-3 for studies of 14-3-3/phospho-Tau interaction. PLoS One.

2017;12:e0178933.

[49] Nikolakaki E, Giannakouros T. SR/RS Motifs as Critical Determinants of Coronavirus Life

Cycle. Frontiers in molecular biosciences. 2020;7:219.

[50] Wu CH, Yeh SH, Tsay YG, Shieh YH, Kao CL, Chen YS, et al. Glycogen synthase kinase-3

regulates the phosphorylation of severe acute respiratory syndrome coronavirus nucleocapsid

protein and viral replication. J Biol Chem. 2009;284:5229-39.

[51] Sluchanko NN, Tugaeva KV, Greive SJ, Antson AA. Chimeric 14-3-3 proteins for unraveling

interactions with intrinsically disordered partners. Sci Rep. 2017;7:12014.

[52] Tugaeva KV, Kalacheva DI, Cooley RB, Strelkov SV, Sluchanko NN. Concatenation of 14-

3-3 with partner phosphoproteins as a tool to study their interaction. Sci Rep. 2019;9:15007.

[53] Tugaeva KV, Remeeva A, Gushchin I, Cooley RB, Sluchanko NN. Design, expression,

purification and crystallization of human 14-3-3zeta protein chimera with phosphopeptide from

proapoptotic protein BAD. Protein Expr Purif. 2020:105707.

[54] Tugaeva KV, Titterington J, Sotnikov DV, Maksimov EG, Antson AA, Sluchanko NN.

Molecular basis for the recognition of steroidogenic acute regulatory protein by the 14-3-3

protein family. FEBS J. 2020;287:3944-66.

[55] Gogl G, Tugaeva KV, Eberling P, Kostmann C, Trave G, Sluchanko NN. Recognition of

high-risk HPV E6 oncoproteins by 14-3-3 proteins studied by interactomics and crystallography.

bioRxiv. 2020:2020.07.24.220376.

[56] Sluchanko NN, Uversky VN. Hidden disorder propensity of the N-terminal segment of

universal adapter protein 14-3-3 is manifested in its monomeric form: Novel insights into protein

dimerization and multifunctionality. Biochim Biophys Acta. 2015;1854:492-504.

[57] Tung HYL, Limtung P. Mutations in the phosphorylation sites of SARS-CoV-2 encoded

nucleocapsid protein and structure model of sequestration by protein 14-3-3. Biochem Biophys

Res Commun. 2020;532:134-8.

[58] Madeira F, Tinti M, Murugesan G, Berrett E, Stafford M, Toth R, et al. 14-3-3-Pred:

improved methods to predict 14-3-3-binding phosphopeptides. Bioinformatics. 2015;31:2276-83.

[59] Karlberg T, Hornyak P, Pinto AF, Milanova S, Ebrahimi M, Lindberg M, et al. 14-3-3

proteins activate Pseudomonas exotoxins-S and -T by chaperoning a hydrophobic surface. Nat

Commun. 2018;9:3785.

[60] Lo YS, Lin SY, Wang SM, Wang CT, Chiu YL, Huang TH, et al. Oligomerization of the

carboxyl terminal domain of the human coronavirus 229E nucleocapsid protein. FEBS Lett.

2013;587:120-7.

[61] Johnson C, Crowther S, Stafford MJ, Campbell DG, Toth R, MacKintosh C. Bioinformatic

and experimental survey of 14-3-3-binding sites. Biochem J. 2010;427:69-78.

[62] Coblitz B, Wu M, Shikano S, Li M. C-terminal binding: an expanded repertoire and function

of 14-3-3 proteins. FEBS Lett. 2006;580:1531-5.

[63] Tylor S, Andonov A, Cutts T, Cao J, Grudesky E, Van Domselaar G, et al. The SR-rich

motif in SARS-CoV nucleocapsid protein is important for virus replication. Canadian journal of

microbiology. 2009;55:254-60.

[64] Lin L, Shao J, Sun M, Liu J, Xu G, Zhang X, et al. Identification of phosphorylation sites in

the nucleocapsid protein (N protein) of SARS-coronavirus. International journal of mass

spectrometry. 2007;268:296-303.

[65] Sluchanko NN. Association of Multiple Phosphorylated Proteins with the 14-3-3 Regulatory

Hubs: Problems and Perspectives. J Mol Biol. 2018;430:20-6.

[66] Sluchanko NN, Beelen S, Kulikova AA, Weeks SD, Antson AA, Gusev NB, et al. Structural

Basis for the Interaction of a Human Small Heat Shock Protein with the 14-3-3 Universal

Signaling Regulator. Structure. 2017;25:305-16.

[67] Stevers LM, Lam CV, Leysen SF, Meijer FA, van Scheppingen DS, de Vries RM, et al.

Characterization and small-molecule stabilization of the multisite tandem binding between 14-3-

3 and the R domain of CFTR. Proc Natl Acad Sci U S A. 2016;113:E1152-61.

[68] Kostelecky B, Saurin A, Purkiss A, Parker P, McDonald N. Recognition of an intra-chain

tandem 14-3-3 binding site within PKCepsilon. EMBO reports. 2009;10:983-9.

[69] Manschwetus JT, Wallbott M, Fachinger A, Obergruber C, Pautz S, Bertinetti D, et al.

Binding of the Human 14-3-3 Isoforms to Distinct Sites in the Leucine-Rich Repeat Kinase 2.

Front Neurosci. 2020;14:302.

[70] Sluchanko NN. Reading the phosphorylation code: binding of the 14-3-3 protein to

multivalent client phosphoproteins. Biochem J. 2020;477:1219-25.

[71] Stevers LM, de Vries RM, Doveston RG, Milroy LG, Brunsveld L, Ottmann C. Structural

interface between LRRK2 and 14-3-3 protein. Biochem J. 2017;474:1273-87.

[72] Vilcek S. SARS-CoV-2: Zoonotic origin of pandemic coronavirus. Acta virologica.

2020;64:281-7.

[73] Ge XY, Li JL, Yang XL, Chmura AA, Zhu G, Epstein JH, et al. Isolation and characterization

of a bat SARS-like coronavirus that uses the ACE2 receptor. Nature. 2013;503:535-8.

[74] Boucher L, Ouzounis CA, Enright AJ, Blencowe BJ. A genome-wide survey of RS domain

proteins. RNA. 2001;7:1693-701.

[75] Mitrea DM, Kriwacki RW. Phase separation in biology; functional organization of a higher

order. Cell communication and signaling : CCS. 2016;14:1.

[76] Aubol BE, Plocinik RM, Keshwani MM, McGlone ML, Hagopian JC, Ghosh G, et al. N-

terminus of the protein kinase CLK1 induces SR protein hyperphosphorylation. Biochem J.

2014;462:143-52.

[77] Iserman C, Roden CA, Boerneke MA, Sealfon RSG, McLaughlin GA, Jungreis I, et al.

Genomic RNA elements drive phase separation of the SARS-CoV-2 nucleocapsid. Molecular

Cell. 2020.

[78] Carlson CR, Asfaha JB, Ghent CM, Howard CJ, Hartooni N, Safari M, et al.

Phosphoregulation of Phase Separation by the SARS-CoV-2 N Protein Suggests a Biophysical

Basis for its Dual Functions. Molecular Cell. 2020.

[79] Lin JP, Fan YK, Liu HM. The 14-3-3eta chaperone protein promotes antiviral innate

immunity via facilitating MDA5 oligomerization and intracellular redistribution. PLoS Pathog.

2019;15:e1007582.

[80] Liu HM, Loo YM, Horner SM, Zornetzer GA, Katze MG, Gale M, Jr. The mitochondrial

targeting chaperone 14-3-3epsilon regulates a RIG-I translocon that mediates membrane

association and innate antiviral immunity. Cell host & microbe. 2012;11:528-37.

[81] Fogg MJ, Wilkinson AJ. Higher-throughput approaches to crystallization and crystal

structure determination. Biochem Soc Trans. 2008;36:771-5.

[82] Chernik I, Seit-Nebi A, Marston S, Gusev N. Small heat shock protein Hsp20 (HspB6) as a

partner of 14-3-3gamma. Mol Cell Biochem. 2007;295:9-17.

[83] Ramteke MP, Shelke P, Ramamoorthy V, Somavarapu AK, Gautam AK, Nanaware PP, et

al. Identification of a novel ATPase activity in 14-3-3 proteins--evidence from enzyme kinetics,

structure guided modeling and mutagenesis studies. FEBS Lett. 2014;588:71-8.

[84] Sluchanko NN, Chebotareva NA, Gusev NB. Modulation of 14-3-3/phosphotarget

interaction by physiological concentrations of phosphate and glycerophosphates. PLoS One.

2013;8:e72597.

[85] Molzan M, Weyand M, Rose R, Ottmann C. Structural insights of the MLF1/14-3-3

interaction. FEBS J. 2012;279:563-71.

Figures

	ACE2 expression *	SARS-CoV-2 presence**	14-3-3 total	14-3-3ζ (zeta)	14-3-3ε (epsilon)	14-3-3β (beta)	14-3-3γ (gamma)	14-3-3τ (tau)	14-3-3σ (sigma)	14-3-3η (eta)
Oral cavity			9450	1383	2334	776	625	577	3234	521
Gastrointestinal system			9044	2254	1681	903	656	884	2042	624
Lungs			17790	6976	4373	2746	971	1384	156	1184
Urinary system and kidney			5519	1409	1531	1005	588	582	94	310
Male reproductive system			6638	1611	1992	794	648	910	225	458
Nervous system/brain			23494	6644	6003	3024	4122	1807	468	1426
Cardiovascular system/heart			5333	1182	860	894	1094	587	203	513
Pancreas			6989	1593	2301	921	704	751	171	548
Skin and adipose tissue			5976	1854	1520	659	340	637	694	272
Gall bladder			4920	969	1525	705	614	581	129	397
Liver			4352	1115	1086	855	620	365	147	164
Female reproductive system			8283	2083	2246	1152	742	1371	211	478

very low	low	high

scale

2423	2288	1203	977	870	648	575

average abundance of a 14-3-3 isoform in 12 selected tissues, ppm

Fig. 1. 14-3-3 proteins are highly abundant in human tissues with SARS-CoV-2 presence. Correlation of ACE2 expression levels (*) and SARS-CoV-2 reported presence (**) in various tissues of COVID19 patients based on the data from [6], shown with abundances (indicated in ppm, part per million, i.e., one molecule of a given protein per 1 million of all proteins from a given tissue) of the seven human 14-3-3 isoforms, extracted from the PAXdb database [40]. Different tissues are shown in the order corresponding to the SARS-CoV-2 presence, starting, at the top, from the highest virus presence [6]. The shown relative scale of ACE2 expression is also taken from [6]. The total abundance of the seven human 14-3-3 isoforms in a given tissue and the average abundance of an isoform in 12 selected tissues are also indicated. The latter values were used for ordering the data for 14-3-3 isoforms, left to right, from the highest average abundance (14-3-3ζ; 2423 ppm, or ~0.24%) to the lowest average abundance (14-3-3η; 575 ppm, or ~0.06%). Note that the average abundance of all seven 14-3-3 proteins in three tissues with the highest SARS-CoV-2 presence (oral cavity, gastrointestinal tract, lungs) reaches 1.21% of all proteins.

Fig. 2. Characterization of the SARS-CoV-2 N phosphoprotein. A. Schematic representation of the N sequence and its main features. The subdomains are named above, predicted Nuclear Export Signal (NES, yellow), Nuclear Localization Signal (NLS, violet), N- and C-terminal domains (NTD, CTD) and two main phosphorylation loci

(cyan rectangles) are marked. Sequences of the SR-rich region and short
phosphorylatable section of the CTD are shown aligned between SARS-CoV and
SARS-CoV-2 N proteins, highlighting multiple conserved phosphorylation sites (bold
font indicates phosphosites present in both N proteins and experimentally confirmed for
SARS-CoV-2 N). Some phosphosites (red spheres and labeled) are predicted as
suboptimal 14-3-3 binding sites (green arrows). B. A Phos-tag gel showing that bacterial
co-expression of the full-length N with PKA yields polyphosphorylated protein. C. A
fragmentation spectrum of the representative phosphopeptide carrying phosphorylation
at Ser197 and Thr205. Z- (red) and c-series (blue) of ETD fragmentation are shown.
Error did not exceed 5 ppm. D. Absorbance spectra show that both recombinant
unphosphorylated and PKA-phosphorylated N proteins elute from the Ni-affinity column
bound to random *E.coli* nucleic acid. On-column washing with 3 M NaCl (50 column
volumes) eliminates bound nucleic acid. E. Analysis of the oligomeric state of pN using
size-exclusion chromatography on a Superdex 200 Increase 10/300 column at 200 mM
NaCl, with multi-angle light scattering detection (SEC-MALS) and SDS-PAGE of the
eluted fractions. Flow rate was 0.8 ml/min. Apparent Mw determined from column
calibration based on protein standards (arrows above) is compared to the absolute
mass determined from SEC-MALS. The experiments were carried out three times, and
the most typical results are shown.

Fig. 3. Interaction of SARS-CoV-2 N with human 14-3-3γ. Interaction between SARS-CoV-2 N protein (50 µM) and human 14-3-3γ (160 µM) was analyzed by SEC-MALS and SDS-PAGE of the eluted fractions. SEC-MALS/SDS-PAGE of 14-3-3γ alone (A,B), with unphosphorylated N (C,D) and with phosphorylated N (E,F) are presented. Mw distributions obtained using MALS are shown on each peak. "L" shows contents of the samples loaded on a column, elution volumes corresponding to the maxima of the peaks are indicated by arrows below the gels. Mw markers are shown to the left of each gel. A black bar on panel F indicates the fractions of the complex between 14-3-3γ and pN. A Superdex 200 Increase 10/300 column equilibrated with 20 mM Tris-HCl buffer pH 7.6 containing 200 mM NaCl and 3 mM NaN$_3$ was operated at a flow rate of 0.8 ml/min. The experiment was performed three times and the most typical results are shown.

Fig. 4. SARS-CoV-2 N interacts with all seven dimeric human 14-3-3 isoforms, but
not with the monomeric mutant of 14-3-3ζ. A. Interaction between phosphorylated N
protein (46.6 μM) and human isoforms 14-3-3γ (69.4 μM), his-tagged 14-3-3γΔC (73.9
μM), his-tagged 14-3-3ηΔC (73.8 μM), 14-3-3ζ (77.6 μM), his-tagged 14-3-3ε (76.2 μM),
14-3-3εΔC (60.3 μM), 14-3-3βΔC (69.4 μM), 14-3-3τΔC (71.7 μM), 14-3-3σΔC (75.4
μM) studied by analytical SEC on a Superdex 200 Increase 5/150 column at 200 mM
NaCl. B. Phosphorylated N protein (46.6 μM) interacts with the wild-type dimeric 14-3-3ζ
(77.6 μM) but not with its monomeric mutant 14-3-3ζm-S58E (74.9 μM). Apparent Mw
indicated in kDa were determined from column calibration.

Fig. 5. Binding affinity. pN at a fixed concentration (~10 µM) was titrated against either of the two human 14-3-3 isoforms and monitored by analytical SEC with serial sample loading. A. Titration of pN with 14-3-3γ. B. Titration of pN with 14-3-3ε. Changes of the elution profiles associated with the increasing 14-3-3 concentration are shown by arrows. C. Binding curves used for apparent K_D determination. Note that, regardless of the observed differences in the affinities, the maximum 14-3-3 concentration complexed with pN equals the concentration of pN (~10 µM) for both 14-3-3 isoforms. Typical results from two independent experiments are shown.

Fig. 6. **Interaction of human 14-3-3γ with individual SARS-CoV-2 N protein domains.** A. Schematic representation of the N protein sequence with the main domains/regions highlighted. Yellow indicates NES, small red circles designate phosphosites within the two corresponding phosphorylatable loci. B. SEC profiles of individual 14-3-3γ (black traces), individual N (red traces), and 14-3-3 with N (blue traces) obtained using a Superdex 200 Increase 5/150 column at 200 mM NaCl in running buffer and operated at a 0.45 ml/min flow rate. The upper row corresponds to unphosphorylated N fragments, the bottom row corresponds to their phosphorylated counterparts, as indicated on each panel. In all cases, 50 μM of N was mixed with 50 μM of 14-3-3 and loaded in an equal volume of 50 μl. In the case of N.247-364 and pN.247-364, 90 μM of N was used. Note that the well-defined complexes with 14-3-3 were observed only for pN.1-211 and pN.1-238 fragments. The experiment was repeated twice and the most typical results are presented.

Fig. 7. Interaction of human 14-3-3γ with the phosphorylated monomeric SARS-CoV-2 N construct 1-238. A. SEC-MALS analysis of individual pN, 14-3-3γ and their complex formed at the indicated molar excess of pN. Protein concentration, MALS-derived Mw distributions across the peaks and the respective average Mw values are indicated. Note that the difference shown by a two-headed arrow roughly corresponds to a pN.1-238 monomer mass. B. Schematic representation of the possible bidentate binding mode where the 14-3-3 dimer interacts with the tentative key 14-3-3-binding phosphosite and another sterically allowed phosphosite separated by a sufficiently long linker.

Fig. 8. Localization of the 14-3-3-binding sites within the SR-rich region of the SARS-CoV-2 N protein. A. Sequence of the SR-rich region showing potential 14-3-3-binding sites (green and blue font) and the truncations designed to exclude one or two main 14-3-3-binding sites (blue font). # denotes the designed C-terminus. B, C. Analysis by SEC of the interaction between human 14-3-3γ and the truncated mutants of SARS-CoV-2 N in their unphosphorylated (B) or phosphorylated forms (C). Each graph contains SEC profiles for the individual 14-3-3 (black line), individual N or pN construct (red line), and the 14-3-3γ N/pN mixture (blue line) where 50 μM of each protein was used throughout. Inserts on panel C show the upward shift on SDS-PAGE as the result of phosphorylation (U, unphosphorylated; P, phosphorylated protein, arrowheads indicate the shift). Note that only the phosphorylated mutants interacted with 14-3-3γ and that only pN.1-211 and pN.1-204 formed a defined complex with 14-3-3γ. Column: Superdex 200 Increase 5/150, flow rate: 0.45 ml/min. The experiment was repeated twice and the most typical results are presented.

Fig. 9. Association of SARS-CoV-2 N with human 14-3-3. A. A topology model for the complex of SARS-CoV-2 N protein dimer with the dimer of human 14-3-3 illustrating the occlusion of the SR-rich region. B. Local alignment of SR-rich regions of the most similar coronavirus N proteins in order of descending sequence identity (s.id.) determined using entire N protein sequences. Alignment was performed using Clustal omega and visualized using Mview (https://www.ebi.ac.uk/Tools/msa/mview/). Residues identical to those in the SARS-CoV-2 sequence are shadowed by grey, phosphorylatable residues in positions 197 and 205 are in green color, residues blocking phosphorylation in position 205 are in red.