Road salt compromises functional morphology of larval gills in populations of an

amphibian

Running title: Salt alters amphibian gill structure

Keywords: osmoregulation; osmotic stress; pollution; road effects; road salt; secondary

salinization

Manuscript type: Primary research

The manuscript contains a table, a single panel figure, four multi-panel figures, and a

supplemental table.

Richard V. Szeligowski[1]

Jules A. Scanley[2]

Christine C. Broadbridge[2]

Steven P. Brady[1]

[1]Biology Department, Southern Connecticut State University. New Haven, CT 06515.

[2]Center for Nanotechnology, Connecticut State Colleges and Universities. New Haven, CT

06515.

Corresponding Author:

Richard Szeligowski

Richard.szeligowski@yale.edu

Summary Statement:

Chronic road salt exposure alters the functional morphology of gills in larval amphibians, potentially compromising osmoregulation, feeding, and respiration.

Abstract

Throughout much of the world, winter deicing practices have led to secondary salinization of freshwater habitats, where numerous taxa are vulnerable to elevated salinity. Many amphibians are of particular concern because of their permeable skin and reliance on small ponds and pools, where salinity levels can be high. The early life-history stages of amphibians that develop in these habitats are especially sensitive to salt exposure. Larvae developing in salt-polluted environments must osmoregulate through ion exchange in gills. While salt-induced changes to the physiology of ion exchange in amphibian gills is generally understood, functionally relevant changes in gill morphology remain poorly described. Yet the structure of gills should be an important component affecting their ionoregulatory capacity, for instance in terms available surface area. Larval amphibian gills also play critical roles in gas exchange and foraging. Thus, changes in gill morphology due to salt pollution potentially affect not only osmoregulation, but also respiration and feeding. Here, we used a chronic exposure experiment to quantify the effect of salinity on larval gill morphology in populations of the wood frog (*Rana sylvatica*). We measured a suite of morphological traits on gill tufts, where ionoregulation and gas exchange occur, and on gill filters, which are used in feeding. Larvae raised in high salinity conditions had gill tufts with lower surface area to volume ratio, while epithelial cells on these tufts were less circular but occurred at higher densities. Gill filters showed increased spacing, which can potentially reduce their efficiency in filtering food particles. Together, these changes seem likely to diminish the ionoregulatory and respiratory capacity of gill tufts, and compromise feeding functionality of gill filters. Thus, a singular change in the aquatic environment from a widespread pollutant has the potential to generate a suite of consequences via changes in gill morphology. Critically, this suite of negative effects is likely most detrimental in salinized environments, where ionoregulatory demands are higher, which in turn should increase respiratory demands along with energy acquisition demands through foraging.

Introduction

Salt pollution: a widespread problem for freshwater organisms

Freshwater habitats are becoming salinized by pollution from human activities including

industrial processes, mining effluent, agricultural runoff, and most substantially, road salt used

for de-icing (Lauer et al. 2016; Kaushal et al., 2005; Dugan et al., 2017; Maloney et al., 2017).

Road salt applied to roadways, driveways, walkways, and parking lots runs off into adjacent

habitats and/or stormwater infrastructure, and much of this salt is carried into surface waters.

For instance, one estimate suggests that about 70% of lakes in the North American Lakes

Region are polluted by road salt, and that if current salting practices continue, chloride pollution

levels in many lakes will exceed the EPA's water quality criterion 2050 (Dugan et al., 2017).

Elevated salinity in freshwater habitats spurs a suite of negative effects across biological levels,

from genetic to ecosystem (Hintz et al., 2017; Gibbons et al., 2018; Merrick and Searle, 2019;

Venâncio et al., 2019). For many freshwater taxa, salt pollution causes numerous sublethal

effects, including changes in foraging and anti-predator behavior (Hall et al., 2017); changes in

life history traits (Faulkner et al., 2019); and increased malformations (Brady, 2013; Hopkins et

al., 2013; Alam et al., 2020). Salt can also be directly toxic (Sanzo and Hecnar, 2006; Karraker

et al., 2008; Mahrosh et al., 2014). Together, these impacts can lead to loss of genetic diversity,

population declines, and changes in community composition (Van Meter and Swan, 2014;

Castillo et al., 2018), which collectively can modify ecosystem function (Schuler et al., 2017).

Salt pollution also affects biogeochemical cycles, and these changes can scale up through food

webs (Judd et al., 2005). For instance, salt pollution increases nitrogen and carbon export in

streams (Reviewed in Hintz and Relyea, 2019) while plant and zooplankton richness have both

been shown to decline in freshwater habitats polluted by salt (Richburg et al., 2001; Gutierrez et

al., 2018).

While many freshwater taxa are affected by salt pollution, amphibians are of particular concern

in part because of ongoing global population declines (Green et al., 2020). Although a subset of

amphibians is tolerant of saline conditions (Hopkins and Brodie, 2015), many are considered

vulnerable to increased salinization because they have narrow osmotic ranges (Chinathamby et

al., 2006; Sanzo and Hecnar, 2006; Brady, 2013), possess highly permeable skin (Pough,

2007), and rely on freshwater habitats for reproduction, development, and dwelling (Vences and

Köhler, 2008). While even modest increases in salt can be detrimental (Venâncio et al., 2019),

concentrations can become especially high in shallow and/or temporary habitats (e.g., vernal

pools), which many amphibians require to complete their lifecycle (Semlitsch and Skelly, 2007).

Consequences of exposure to secondary salinization in amphibians include osmotic stress

(Gomez-Mestre et al., 2004) and increased energy allocated to osmoregulation (Peña-Villalobos

et al., 2016); increased risk of predation (Denoël et al., 2010); behavioral changes including

reduced swim performance (Squires et al., 2008) and reduced foraging activity (Hall et al.,

2017); decreased abundance (Albecker and McCoy, 2017); lower larval survival rates and

ultimately, population size (Karraker et al., 2008).

Osmoregulation in amphibians

Many amhibians have complex life cycles (Semlitsch and Skelly, 2007), and the negative effects

of salt exposure tend to be most severe during early stages of development as embryos and

larvae (Gomez-Mestre et al., 2004; Alexander et al., 2012; C.S. Wu et al., 2014; Park and Do,

2020). In anuran larvae, gills are the primary site of ionoregulation—a key mode of

osmoregulation in which ions are exchanged between internal and external environments to

maintain internal ion concentrations, especially Na^+ and Cl^- (Alvarado and Moody, 1970; Dietz

and Alvarado, 1974). For amphibians in typical (unpolluted) freshwater habitats, the tendency is

for ions to be lost to the external environment. Osmoregulation for amphibians in these

environments requires actively pumping ions in through gill epithelium into systemic plasma

(Boonkoom and Alvarado, 1971). The primary mechanism responsible for ion exchange is the

Na^+/K^+ ATPase ion pump (NKA). NKA is a transmembrane pump found in all animals that

maintains body fluid homeostasis by exporting 3 Na^+ ions while importing 2 K^+ ions per ATP

molecule against their respective concentration gradients (Skou, 1957; Pivovarov et al., 2019;

Fig. 1). NKA-mediated osmoregulation operates via passive influx of Na^+ from the external

environment into an apical epithelial cell. From there, NKA actively transports Na^+ ions into the

Na^+-rich plasma while exchanging K^+ from the plasma into the cytosol, where K^+ concentrations

are higher. K^+ ions exit the cytosol back into the plasma via passive membrane channels, which

creates an electric potential between the external environment and the plasma, leading to

passive influx of Cl^- ions and creating a hyperosmotic internal environment (Campbell et al.,

2012).

Anuran gills are thought to function much like those of teleosts (Ultsch et al., 1999) in that NKA

is highly expressed in mitochondrion-rich cells (MRCs) in the gill epithelium, where MRCs

function as the primary mechanism of ionoregulation (Hwang and Lee, 2007). When freshwater

amphibian species experience osmoregulatory challenges through salinization, they must

prevent water loss by increasing ion absorption, thereby maintaining a hyperosmotic internal

environment (Gomez-Mestre et al., 2004). Bernabò et al. (2013) found that larvae of the

common toad (*Bufo bufo*) and Balearic green toad (*Bufo balearicus*) increased NKA abundance

in response to increasing salinity, presumably to maintain a hyperosmotic internal environment

and prevent water loss. Euryhaline (i.e., salinity tolerant) amphibians such as the Indian rice frog

(*Fejervarya limnocharis*) seem to utilize the same strategy for osmoregulation, although the

specific mechanisms that allow increased salinity tolerance are not fully understood. Wu *et al.*

(2014) found that crab-eating frog (*Rana cancrivora*) tadpoles acclimated to salinity levels

corresponding to 15,700 µS/cm had a six-fold increase in NKA abundance when compared to

those raised in 4,000 µS/cm water. Although crab-eating frog tadpoles can survive in full

strength seawater after acclimation (Gordon and Tucker, 1965), Hsu *et al.* (2012) found that

metamorphosis occurred in only 40% of tadpoles in salinity levels corresponding to 60%

seawater (30,000 µS/cm) after salinity acclimation.

Toward improved understanding of gill morphology in osmoregulation

In contrast to our knowledge of *physiological* mechanisms underpinning osmoregulation in larval

amphibians, we have limited understanding of the functional role of gill morphology, and how it

might be affected by salt exposure. Yet changes in gill morphology are important adaptations in

fish that transition between freshwater and seawater (Uchida et al., 1996; Hiroi et al., 1999), and

thus might be important in freshwater amphibians exposed to salt. Limited, primarily qualitative

insights from an acute exposure experiment show that high salinity can modify gill morphology

in larvae of the green toad (*Bufo balearicus*) and common toad (*Bufo bufo*) (Bernabò et al.,

2013). Specifically, gill tufts became "dehydrated" and "invaginated" in larvae exposed to 12,700

µS/cm. For larvae exposed to 15,000 µS/cm, hyperplasia was observed in pavement cells,

which play a limited role in ion exchange (Wilson, 2011). While these qualitative insights are

important, they were based on short-term exposure to salt concentrations far above those found

in most secondarily-salinized freshwater habitats. Thus, a quantitative assessment of

morphology, based on chronic exposure to ecologically relevant salinity levels for pollution is

critically needed.

Changes in larval amphibian gill filter morphology could also affect gas exchange and therefore

respiration. And because gills are also important structures for larval feeding, salt-induced

morphological changes may hamper foraging efficiency. For instance, if gill morphology

becomes less well suited to trapping food particles, larvae should experience relative increases

in foraging demand. Conceivably, these various effects could be multiplicative: increased

demands for ionoregulation in salt-polluted environments could change gill morphology in ways

that reduce efficiency for ionoregulation and gas exchange, driving energetic demands higher.

At the same time, reduced foraging efficiency could further increase energetic demands. Thus,

any changes in gill morphology have the potential to generate a cascade of effects on

amphibian physiology and fitness.

Here, we begin exploring these potential effects by quantifying changes in gill morphology in

populations of the wood frog (*Rana sylvatica*). We raised wild-collected embryos in low and high

salinity conditions, from early embryonic stages until nearing metamorphosis. We used

scanning electron microscopy to image and quantify changes in gill morphology in response to

salinity. We hypothesized that increased salinity would alter gill morphology, particularly at the

gill tufts and filters. These sites, respectively, are where most osmoregulation occurs via NKA

activity (Brunelli et al., 2004), and where food particles are filtered for feeding (Seale and

Wassersug, 1979). We predicted that salt would modify gill morphology in ways that

compromise ionoregulatory capacity, gas exchange, and feeding.

Methods

Natural History and larval gill function

Wood frogs are found throughout the northern ranges of North America, from the southern

Appalachian Mountains through Canadian and Alaskan forests in habitats including tundra,

deciduous and coniferous forests, marshes, and grasslands (Green et al., 2014). Wood frog

breeding is seasonal. In our study region in the northeastern U.S., adults migrate from upland

terrestrial habitat to nearby small, shallow, and often temporary ponds, typically during the first

warm rain of early spring following hibernation. Breeding is generally explosive, occurring over a

1-2 week period in each pond. Each female lays one clutch as a single egg masses containing

about 1,000 eggs, which are fertilized externally during amplexus. Eggs develop for about 2-3

weeks before hatching, after which time larvae (i.e., tadpoles) develop for about two months

before metamorphosing into terrestrial juveniles.

Among amphibians, the wood frog stands out as having a particularly puzzling response to salt

exposure. Not surprisingly, this species shows a variety of negative responses to salt pollution

and roadside dwelling including many of the known negative effects associated with salt

exposure, such as direct mortality and increased malformation in embryos and larvae (Karraker

et al., 2008; Brady, 2013) as well as reduced growth rates and increased developmental

variance (Brady, 2013; Dananay et al., 2015). However, reciprocal transplants and common

garden experiments reveal that aquatic-stage wood frogs from salt-polluted roadside ponds are

more sensitive to road salt and roadside conditions than nearby populations from unpolluted
ponds (Brady, 2013; 2017). These findings suggest that roadside populations might have
recently evolved lower tolerance to salt than individuals from unpolluted populations, consistent
with 'local maladaptation' processes (Brady et al., 2019). Alternatively, this pattern might reflect
a fitness tradeoff between aquatic- and terrestrial-stage frogs, for instance if lower salt tolerance
is correlated with adult lifetime reproductive success. In either case, we speculate that the
survival disadvantage in roadside populations might arise through effects on gill morphology.

During larval development, the primary site of respiration changes. Prior to feeding stage -
Gosner (Gosner, 1960) stage 25 - respiration occurs primarily via cutaneous diffusion and the
limited activity of external gills before transitioning to gas exchange in a bilateral pair of internal
gills (Feder and Burggren, 1992; Brunelli, 2004). Each internal gill contains three prominent
branchial arches that originate in the rostral end of the gill, extending caudally. A reduced fourth
branchial arch lateral to the first three arches is present in some species but has decreased tuft
density and filter plate size (McIndoe and Smith, 1984; Brunelli et al., 2004). Gill tufts are the
primary site of gas exchange (McIndoe and Smith, 1984; Brunelli et al., 2004), protruding
ventrally from the branchial arch. Each tuft has numerous smaller structures that split from the
main body (hereafter 'tuft branches'). A cartilaginous filter plate protrudes dorsally from the
branchial arch (McIndoe and Smith, 1984), and rows of gill filters extend laterally from the filter
plate, with main folds running ventrodorsally in a zigzag pattern (McIndoe and Smith, 1984).
Secondary folds branch from the main folds at corners, often continuing the zigzag pattern of
main folds, tertiary folds branch away from secondary folds at their corners, often forming a
rectangular filter niche with the main and secondary folds (McIndoe and Smith, 1984). We
termed points where folds changed directions, branched into other folds, or terminated 'nodes'.

Gill filters function as a screen for food particulates as water travels over them towards the tufts
(Feder et al., 1984; McIndoe and Smith, 1984; Brunelli et al., 2004). Both gill tufts and filters
play roles in ionoregulation. However, tufts seem to play a greater part than filters in this
capacity. For instance, Bernabò et al. (2013) showed NKA is more strongly expressed in gill
tufts than filters in both common toad and Balearic green toad gills, although NKA expression
increased in tufts and filters for both species when exposed to salt levels between 5%-25% sea
water.

Embryo collection and rearing

We collected egg masses from six natural ponds in Norwich, VT during the breeding season in
April 2019. To represent a range of populations, we sampled egg masses from three salt-
polluted 'roadside' ponds located within 10 m of a paved road and three unpolluted 'woodland'
ponds, which were located at least 500 m from any road. From each pond, a cluster of
approximately 20 embryos were removed from each of five egg masses. Selection of egg
masses and embryo groups was haphazard. Embryo clusters were placed in plastic 700 ml
(16.5 x 7.5 x 12.5 cm) containers filled with natal pond water and were immediately transported
to the lab in New Haven, CT where they were stored at 4˚C for 24 hours until the start of the
chronic salt-exposure experiment.

Four salinity treatments were prepared using aged, conditioned tap water and NaCl road salt
obtained from the Connecticut Department of Transportation. Embryos were exposed to one of
four treatments: 275 µS/cm (no salt added), 1,000 µS/cm, 4,000 µS/cm, and 7,000 µS/cm.
These values were chosen, respectively, to represent control conditions, average roadside
conditions, maximum roadside conditions, and values exceeding roadside conditions to
consider a broad range of exposure scenarios. Embryo clusters were assigned haphazardly to
one of four salinity treatments in a 5.6 L container (30.5 x 10 x 20 cm) containing 5 L of
assigned treatment solution such that each pond had representative embryos raised in each
treatment solution. Containers were arranged randomly across four levels of a shelving unit and
were rotated weekly within and between shelves to distribute potential block effects. When
tadpoles reached feeding stage, they were fed 5-day food rations containing a 3:1 ratio of rabbit
chow to fish food concurrently with water changes. Tadpoles were raised until they reached
approximately Gosner stage 35. After recording tadpole developmental stage scores, individuals
were euthanized via immersion in buffered Tricaine (2g/L MS222 ethyl 3-aminobenzoate
methanesulfonate, Sigma-Aldrich; pH 7.0) and gills were immediately excised and prepped for
scanning electron microscopy (SEM). All methods complied with institutional animal care
guidelines

Scanning electron microscopy sample preparation

We selected tadpoles haphazardly for SEM, imaging 19 individual tadpoles representing
different salt treatments and populations (see Supp Table 1). This sample size was chosen to
balance SEM constraints (e.g., costs, sample prep time, personnel time) with study goals.
Selected individuals ranged in Gosner developmental stage from 34-37. For each tadpole, the
left gill was excised, measured for length using the reticle of a dissecting scope, and
immediately transferred to 3% glutaraldehyde in phosphate buffer at 4˚C for 1.5 hours, then

transferred into 1% osmium tetroxide in phosphate buffer for 2 hours at 4˚C. Gills were then

dehydrated in a series of graded ethanol steps (30%, 50%, 70%, 80%, 90% and 100%),

modified from Brunelli et. al (2004). Gills were dried in a Tousimis Samdri-780 critical point drier

and fixed to SEM stubs. Gills were imaged with a Zeiss Sigma VP field emission scanning

electron microscope.

Gill Morphology

Images were analyzed using ImageJ software (v. 1.51, Schneider et al. 2012). To describe

potential effects of salt on gill morphology, we targeted measuring 12 different traits per gill

(Table 1). We measured gill tuft traits (N= 7) corresponding to available surface area (e.g. tuft

branch length and width), gill filter traits (N=5) corresponding with surface area available for ion

exchange (e.g. internode length), and traits related to filtering efficacy (e.g. internode length and

angle). Tuft branch cell attributes were also measured to assess how cell morphology may

contribute to ultrastructure (observable through electron microscopy) changes. Each gill was

assessed for multiple traits; however, SEM imaging could not guarantee comprehensive

coverage of all traits on each gill (see Supplemental Table 1).

For gill tufts, we measured epithelial cell attributes including cell area (C_A) and cell perimeter

(C_P) by tracing the perimeter of cells where boundaries were clearly definable, targeting 25 cells

per gill (illustrated in Fig. 2a). Cell aspect ratio and circularity were calculated from these values

to characterize cell shape. Tuft cell density (T_D) was calculated by tracing clusters of cells within

tuft branches and counting the number of cells in each defined area, then computing the value

of cell count over area. Tuft branch measurements were obtained by averaging three linear

measurements in tuft branch length (T_L) and tuft branch width (T_W), respectively (illustrated in

Fig. 3a). We averaged three measurements per trait per branch to account for nonuniform width

along branches and imprecise branch origins.

For gill filters, we measured internode distance for both main folds (D_M) and secondary folds

(D_S) (Fig. 4a). We also measured angles between main folds (φ_{MM}) and secondary folds (φ_{MS})

(Fig. 4a). Measurements for φ_{MS} angles were measured starting from the ventral end of the

main fold (proximal to branchial arch) and tracing the angle of the secondary fold, φ_{MM} angles

were measured ventrodorsally. The distance between main folds (D_{MM}) on filter plates (Fig. 4a)

was measured by taking the average of three measurements of the distance from a point on the

main fold apparatus (corresponding to nodes and points midway between nodes) to the nearest

point on an adjacent main fold apparatus.

Analysis

To maximize power given our sample size, we composited data into two groups instead of the

original four treatment groups – 'high salinity' comprised the three different salt concentrations

while 'low salinity' comprised the control group. We used the 'lme4' package (Haubo Bojesen et

al., 2017) in RStudio (v. 3.6.2, RStudio Team, 2020) to compose linear mixed effects models

analyzing each gill trait in response to salinity. We included gill length and developmental stage

as fixed effect covariates because of their potential to influence morphology. In each model,

pond and gill identity were included as random effects to account for correlation within ponds

and among repeated measurements on each gill. For each trait, we used AIC selection to

choose the preferred random effects structure with a delta AIC threshold value of 2. Initial

models were fit for AIC selection using restricted maximum likelihood, and selected models for

inference were refit with maximum likelihood. *P*-values for fixed effects were based on degrees

of freedom calculated with Satterthwaite's approach implemented in the R package

'lmerTest' (Haubo Bojesen et al., 2017). We estimated surface area and volume for gill tuft

branches by treating them as idealized cylinders and used model estimates for T_L. Because T_W

did not vary significantly, the average of T_W estimates was used for surface area and volume

estimates. We estimated the filter niche area (i.e., size of spaces within filters) by treating them

as rectangles with sides D_M and D_S and used model estimates for D_S. As D_M did not vary

significantly, we used the average of D_M model estimates for low and high salinity treatments.

Results

Tuft branches were longer in high versus low salinity treatments ($F_{1, 7.4}$ = 8.3, *P* = 0.022; fig. 3b).

Specifically, model estimates for tuft branch length were 31.6 μm for gills in the low salinity

treatments compared to 47.2 μm in the high salinity treatments. Thus, gill tuft branch length

increased by an estimated 49.5% in tadpoles raised in high salinity conditions. Gill tuft branch

width ($F_{1, 9.4}$ = 0.5, *P* = 0.484) did not vary between salinity treatments. Calculations of branch

surface area and volume using these model estimates gave a 41.86% increase in surface area

from low salinity to high salinity treatments, but a 49.46% increase in tuft branch volume,

reducing the surface area to volume ratio by 5.4% in high salinity treatments.

Traits associated with tuft cells also varied in response to salinity: cell aspect ratio ($F_{1, 294}$ = 15.6,

P = 0.001; fig. 2b), cell circularity ($F_{1, 5.0}$ = 13.1, P = 0.0152; fig. 2c) and cell density ($F_{1, 9.9}$ = 7.0,

P = 0.025; fig. 2d) each varied in response to salinity. Model estimates for aspect ratio was 1.45

in low salinity treatments versus 1.63 in high salinity treatments, representing an 11.9%

increase. Circularity in low salinity was estimated to be 0.611 compared to 0.562 in high salinity,

revealing a 7.96% decrease in circularity for high salinity treatments. Cell density was 47.4%

higher in high salinity tadpoles, with 0.025 cells per μm^2 compared to 0.017 cells per μm^2 in low

salinity treatments. Tuft cell perimeter ($F_{1, 8.0}$ = 1.5, P = 0.260) and tuft cell area ($F_{1, 8.0}$ = 1.0, P =

0.354). No significant variation in any tuft trait occurred in response to Gosner developmental

stage or gill length (all P > 0.166).

The distance between gill filter secondary nodes (D_S) varied between low and high salinity

treatments ($F_{1, 4.8}$ = 10.3, P = 0.026; fig. 4b). In low salinity treatments, D_S was estimated to be

13.52 μm compared to 22.96 μm for high salinity treatments, representing a 68.52% increase in

D_S for tadpoles raised in high salinity. There was marginal support for an effect of salinity on

main fold internode distance (D_M) ($F_{1, 4.8}$ = 6.0, P = 0.060; fig. 4c) and distance between main

folds (D_{MM}) ($F_{1, 4.0}$ = 5.4, P = 0.081; fig. 4d). Specifically, D_M model estimates were 14.43 μm in

low salinity treatments and 21.34 μm in high salinity, representing a 32.36% increase in D_M in

high salinity. D_{MM} model estimates were 45.76 μm in low salinity treatments and 79.26 μm in

high salinity treatments, representing a 57.74% increase in high salinity. Main fold-main fold

angles (φ_{MM}) ($F_{1, 4.3}$ = 4.0, P = 0.111), and main fold-secondary fold angles (φ_{MS}) ($F_{1, 3.9}$ = 0.8, P

= 0.431) did not vary significantly between animals raised in low salinity and high salinity

conditions. Calculating changes in filter niche area by using model estimates gave a 69.23%

increase for filter niche area in high salinity. While neither Gosner developmental stage nor gill

length covariates were correlated with filter trait variation in most of these models (D_M, D_S, φ_{MM},

φ_{MS}; P > 0.411), D_{MM} was correlated with gill length ($F_{1, 4.0}$ = 5.4, P > 0.060), increasing 25.9 μm

with each 1 mm of gill length.

Discussion

High salinity had numerous effects on tadpole gill morphology. Tuft branch length increased by

nearly half while its width was unchanged. Cells on these tufts were more tightly packed

together and were less circular, with relatively longer major than minor axes, but cell perimeter

and area were unchanged. For gill filters, the distance between nodes was longer, which should

result in larger filter size. Together, these results demonstrate that chronic exposure to road salt

modifies gill morphology in ways that reduce available surface area to volume ratio and increase

the spacing of gill filters. These changes appear likely to compromise the functional capacity of

gills for osmoregulation, gas exchange, and foraging.

Gill tufts are the primary site of ionoregulation and gas exchange. Increased tuft length (T_L) in

the absence of increased width (T_W), strongly suggests that changes in morphology due to salt

reduce the available surface area to volume ratio, thus reducing the efficiency and capacity of

ion and gas exchange. Indeed, we estimated a 5.4% reduction in surface area to volume ratio

for larvae from high salinity treatments. In addition to presumably lowering the efficiency of ion

exchange, this change should also reduce the efficiency of gas exchange, which in turn likely

exacerbates the already-increased metabolic demands of elevated salinity (Peña-Villalobos et

al. 2016). The increase in cell density on tufts was attributable to changes in cell shape, allowing

tighter packing of cells on the gill tufts. An increase in cell density could be a response to

increase the rate of ion exchange, as no difference in cell area was found in high salinity

treatments. While this could offset the reduction in the surface area to volume ratio found in high

salinity treatments, a higher cell density is likely to incur a higher energetic cost. However, more

information about cell-specific metabolic rates is needed to determine the extent of this tradeoff.

Gill filters screen food and particulates out of the water prior to gas exchange at the tufts, and

thus play an important role in foraging and feeding (Feder et al. 1984; McIndoe and Smith 1984;

Brunelli et al. 2004). Here, filter size appeared to increase in response to salinity, as evidenced

by increased length of secondary node distance and (cautiously) the marginal evidence for

increased distance between main folds and main fold internode distance. Increased filter size

presumably reduces the efficacy of filtering food particles, as more food particles are likely to be

flushed out through the operculum instead of being retained for consumption (Seale and

Wassersug, 1979). In turn, foraging efficiency should decrease due to high salinity, and this

particular change in morphology could bear an energetic cost as larvae might be required to

increase foraging efforts. However, this change in filter structure likely increases the available

area for ion exchange along the matrix (so-called 'ramifications') of the gill filters. Thus, there

might be a tradeoff between foraging efficacy and osmoregulatory capacity.

A reduction in the ability of gill filters to both capture food and remove particulates from

incoming water before it reaches the tufts may have compounding effects, leading to the widely

reported lethal and sublethal consequences of salinity exposure on amphibians. Allowing larger

particles to contact and potenitally entangle within the tufts (dowstream of the filters) could

reduce the available area for gas exchange. This effect would be magnified by increases in NKA

expression in response to salt (Bernabò et al., 2013; Wu et al., 2014) and the associated

increase in the metabolic demands of osmoregulation (Peña-Villalobos et al., 2016). When

these physiological effects are considered along with observed reductions in foraging activity

(Hall et al., 2017), a decrease in the filter's ability to capture food could have substantial impacts

on survival. Diminshed feeding capacity coupled with larger, less efficient gill tufts likely

constrains larval metabolic resources that are necessary to cope with the osmoregulatory

burden of salt-polluted water.

It is interesting to consider that even though gill tuft surface area to volume ratio declined in

response to high salinity, the length of gill tuft branches and filter secondary folds increased.

This pattern of increased size contrasts with other morphological trait responses for amphibians

exposed to high salinity. For instance, body length and mass have been shown to decrease in

response to salt in numerous studies (e.g., Gomez-Mestre and Tejedo, 2003; Wu and Kam,

2009). Thus, the effect of high salinity on amphibian morphology appears to be tissue-specific.

Osmoregulatory mechanisms in tadpole gill tissues might play a role in these differences via

changes in voltage gradients across membranes. Non-excitable cell depolarization is associated

with tissue growth and development (Shi and Borgens, 1995; Reid and Zhao, 2014), and the

ionic gradient that tadpoles maintain across gill epithelium for osmoregulatory function creates a

voltage gradient that Alvarado and Moody (1970) measured at -5 to -10 mV in pond water,

which depolarized as salinity increased to 20mM/L (1,950 µS/cm). While further work is needed

to characterize the mechanism behind increased gill tuft and filter size in response to salinity,

we are intrigued by the possibility of salinity-induced depolarization contributing to the increases

in gill structure size.

Acknowledgements

We acknowledge the use of facilities and support (personnel and supplies) from the CT State

Colleges and Universities Center for Nanotechnology, Werth Industry Academic Fellowship, and

New Haven Innovation Collaborative Undergraduate Research Fellowship Programs. We thank

E. Brunelli for providing expertise in amphibian gill anatomy. We are grateful for field and

laboratory work by M. Forgione, L. Frymus, J. Priester, F. Senturk, and A. Tucker.

Competing Interests

No competing interests declared

Data Availability

All data will be made available in a public repository upon acceptance.

References

Alam, M. R., Sharmin, S., Islam, S. M., Alam, M. A., Ehiguese, F. O., Pattadar, S. N. and
Shahjahan, M. (2020). Salinity intrusion affects early development of freshwater aquaculture
species pabda, *Ompok pabda*. *Aquac. Reports.* **18,** 100476.

Albecker, M.A. and McCoy, M.W. (2017). Adaptive responses to salinity stress across multiple
life stages in anuran amphibians. *Front. Zool*. **14**, 1–16.

Alexander, L.G., Lailvaux, S.P., Pechmann, J.H.K. and DeVries, P.J. (2012). Effects of
Salinity on Early Life Stages of the Gulf Coast Toad, Incilius nebulifer (Anura: Bufonidae).
Copeia. **2012,** 106–114.

Alvarado, R.H. and Moody, A. (1970). Sodium and chloride transport in tadpoles of the bullfrog
Rana catesbeiana. *Am. J. Physiol*. **218**, 1510–1516.

Bernabò, I., Bonacci, A., Coscarelli, F., Tripepi, M. and Brunelli, E. (2013). Effects of salinity
stress on Bufo balearicus and Bufo bufo tadpoles : Tolerance , morphological gill alterations and
Na + / K + -ATPase localization. *Aquat. Toxicol*. **132–133**, 119–133.

Boonkoom, V. and Alvarado, R. H. (1971). Adenosinetriphosphatase activity in gills of larval
Rana catesbeiana. *Am. J. Physiol.* **220(6)**, 1820–1824.

Brady, S.P. (2012). Road to evolution? Local adaptation to road adjacency in an amphibian
(*Ambystoma maculatum*). *Sci. Rep.* **2,** 1–5.

Brady, S.P. (2013). Microgeographic maladaptive performance and deme depression in
response to roads and runoff. *PeerJ*. **2013,** e163.

Brady, S. P. (2017) Environmental exposure does not explain putative maladaptation in road-

adjacent populations. *Oecologia*. **184(4)**, 931–942.

Brady, S.P., Bolnick, D.I., Barrett, R.D.H., Chapman, L., Crispo, E., Derry, A.M., Eckert,
C.G., Fraser, D.J., Fussmann, G.F., Gonzalez, A., et al. (2019). Understanding maladaptation
by uniting ecological and evolutionary perspectives. *Am. Nat.* **194**, 495–515.

Bridgeman, T.B., Wallace, C.D., Carter, G.S., Carvajal, R., Schiesari, L.C., Aslam, S.,
Cloyd, E., Elder, D., Field, A., Schulz, K.L., et al. (2000). A limnological survey of third sister
lake, michigan with historical comparisons. *Lake Reserv. Manag.* **16**, 253–267.

Brunelli E., Perrotta E. and Tripepi, S. (2004). Ultrastructure and development of the gills in
Rana dalmatina (Amphibia, Anura). *Zoomorphology*. **123**, 203–211.

Campbell, C.R., Voyles, J., Cook, D.I. and Dinudom, A. (2012). Frog skin epithelium:
Electrolyte transport and chytridiomycosis. *Int. J. Biochem. Cell Biol.* **44,** 431–434.

Castillo, A.M., Sharpe, D.M.T., Ghalambor, C.K. and De León, L.F. (2018). Exploring the
effects of salinization on trophic diversity in freshwater ecosystems: a quantitative review.
Hydrobiologia **807**, 1–17.

Chinathamby, K., Reina, R. D., Bailey, P. C. E. and Lees, B. K. (2006). Effects of salinity on
the survival, growth and development of tadpoles of the brown tree frog, *Litoria ewingii*. *Aust. J.*
Zool. **54**, 97.

Dananay, K. L., Krynak, K. L., Krynak, T. J. and Benard, M. F. (2015). Legacy of road salt:
Apparent positive larval effects counteracted by negative postmetamorphic effects in wood
frogs. Environ. *Toxicol. Chem.* **34**, 2417–2424.

Denoël, M., Bichot, M., Ficetola, G.F., Delcourt, J., Ylieff, M., Kestemont, P. and Poncin, P. (2010).
Cumulative effects of road de-icing salt on amphibian behavior. *Aquatic Toxicology*, **99(2)**, 275-280.

Dietz, T. H. and Alvarado, R. H. (1974). Na and Cl transport across gill chamber epithelium of
Rana catesbeiana tadpoles. *American Journal of Physiology*, **226(4)**. 764–770.

Dugan, H. A., Bartlett, S. L., Burke, S. M., Doubek, J. P., Krivak-Tetley, F. E., Skaff, N. K.,
Summers, J. C., Farrell, K. J., McCullough, I. M., Morales-Williams, A. M., et al. (2017).
Salting our freshwater lakes. *Proc. Natl. Acad. Sci. U. S. A.* **114**, 4453–4458.

Faulkner, P. C., Hala, D., Rahman, M. S. and Petersen, L. H. (2019). Short-term exposure to

12‰ brackish water has significant effects on the endocrine physiology of juvenile American
alligator (*Alligator mississippiensis*). *Comp. Biochem. Physiol. -Part A Mol. Integr. Physiol.* **236**,
110531.

Feder, M. E., Seale, D. B., Boraas, M. E., Wassersug, R. J. and Gibbs, A. G. (1984).
Functional conflicts between feeding and gas exchange in suspension-feeding tadpoles,
Xenopus laevis. J. Exp. Biol. **110**, 91-98.

Findlay, S. E. G. and Kelly, V. R. (2011). Emerging indirect and long-term road salt effects on
ecosystems. Ann. *N.Y. Acad. Sci*. **1223**, 58-68.

Gibbons, T. C., McBryan, T. L. and Schulte, P. M. (2018). Interactive effects of salinity and
temperature acclimation on gill morphology and gene expression in threespine stickleback.
Comparative Biochemistry and Physiology -Part A : Molecular and Integrative Physiology. **221**,
55–62.

Gomez-Mestre, I. and Tejedo, M. (2003). Local adaptation of an anuran amphibian to
osmotically stressful environments. *Evolution*. **57**, 1889–1899.

Gomez-Mestre, I., Tejedo, M., Ramayo, E., Estepa, J. and Luisa, M. (2004). Developmental
Alterations and Osmoregulatory Physiology of a Larval Anuran under Osmotic Stress. *Physiol.*
Biochem. Zoo. **77(2)**, 267-274

Gordon, M. S. and Tucker, V. A. (1965). Osmotic regulation in the tadpoles of the crab eating
frog (*Rana cancrivora*). *J. Exp. Bio.* **42**, 437-445.

Gosner, K. L. (1960) A simplified table for staging anuran embryos and larvae with notes
onIdentification. *Herpetologica.* **16(3)**, 183–190.

Gutierrez, M. F., Tavşanoğlu, Ü. N., Vidal, N., Yu, J., Teixeira-de Mello, F., Çakiroglu, A. I.,
He, H., Liu, Z. and Jeppesen, E. (2018). Salinity shapes zooplankton communities and
functional diversity and has complex effects on size structure in lakes. *Hydrobiologia.* **813(1)**,
237–255.

Green D. M., Weir L. A., Casper G. S. and Lannoo M. J. (2014). *North American Amphibians:*
Distribution and Diversity. University of California Press, Berkeley, California.

Green, D. M., Lannoo, M.J., Lesbarrères, D. and Muths, E. (2020). Amphibian population
declines: 30 years of progress in confronting a complex problem. *Herpetologica*. **76(2)**, 97–100.

Hall, E. M., Brady, S. P., Mattheus, N. M., Earley, R. L., Diamond, M. and Crespi, E. J.

(2017). Physiological consequences of exposure to salinized roadside ponds on wood frog
larvae and adults. *Bio. Conserv.* **208**, 98-106.

Haubo Bojesen, R., Kuznetsova, A., Brockhoff, P. B. and B. Christensen, R. H. (2017).
General rights lmerTest Package: Tests in Linear Mixed Effects Models lmerTest Package:
Tests in Linear Mixed Effects Models. *orbit.dtu.dk* 82. doi:10.18637/jss.v082.i13.

Hintz, W. D., Mattes, B. M., Schuler, M. S., Jones, D. K., Stoler, A. B., Lind, L. and Relyea,
R. A. (2017). Salinization triggers a trophic cascade in experimental freshwater communities
with varying food-chain length. *Ecological Applications*. **27(3)**, 833–844.

Hintz, W.D. and Relyea, R.A. (2019). A review of the species, community, and ecosystem
impacts of road salt salinisation in fresh waters. *Freshw. Biol.* **64**, 1081–1097.

Hiroi, J., Kaneko, T. and Tanaka, M. (1999). Sequential changes in chloride cell morphology.
J. Exp. Bio. **202**, 3485-3495.

Hopkins, G. R. and Brodie, E. D. (2015). Occurrence of Amphibians in Saline Habitats: A
Review and Evolutionary Perspective. *Herpetol. Monogr.* **29**,1–27.

Hopkins, G. R., French, S. S. and Brodie, E. D. (2013). Increased frequency and severity of
developmental deformities in rough-skinned newt (*Taricha granulosa*) embryos exposed to road
deicing salts (NaCl & MgCl2). *Environ. Pollut.* **173**, 264–269.

Hsu, W. T., Wu, C. S., Lai, J. C., Chia, Y. K., Hsu, C. H. and Kam, Y. C. (2012) Salinity
acclimation affects survival and metamorphosis of crab-eating frog tadpoles. *Herpetologica*.
68(1), 14–21.

Hwang, P. P. and Lee, T. H. (2007). New insights into fish ion regulation and mitochondrion-
rich cells. *Comparative Biochemistry and Physiology - A Molecular and Integrative Physiology*.
148(3), 479–497.

Judd, K. E., Adams, H. E., Bosch, N. S., Kostrzewski, J. M., Scott, C. E., Schultz, B. M.,
Wang, D. H. and Kling, G. W. (2005). A case history: Effects of mixing regime on nutrient
dynamics and community structure in third sister lake, Michigan during late winter and early
spring 2003. *Lake Reserv. Manag.* **21**, 316–329.

Karraker, N. E., Gibbs, J. P. and Vonesh, J. R. (2008). Impacts of road deicing salt on the
demography of vernal pool-breeding amphibians. *Ecol. Appl.* **18**, 724–734.

Kaushal, S. S., Groffman, P. M., Likens, G. E., Belt, K. T., Stack, W. P., Kelly, V. R., Band,
L. E. and Fisher, G. T. (2005). Increased salinization of fresh water in the northeastern United
States. *Proc. Natl. Acad. Sci. U.S.A.* **102**, 13517-13520.

Lauer, N. E., Harkness, J. S. and Vengosh, A. (2016) Brine Spills Associated with
Unconventional Oil Development in North Dakota. *Env. Sci. Technol.* **50**, 5389-5397.

Mahrosh, U., Kleiven, M., Meland, S., Rosseland, B. O., Salbu, B. and Teien, H. C. (2014).
Toxicity of road deicing salt (NaCl) and copper (Cu) to fertilization and early developmental
stages of Atlantic salmon (*Salmo salar*). *J. Hazard. Mater.* **280**, 331–339.

Maloney K. O., Baruch-Mordo S., Patterson L. A., Nicot J. P., Entrekin S. A., Fargione J.
E., Kiesecker J. M., Konschnik K. E., Ryan J. N., Trainor A. M., et al. (2017). Unconventional
oil and gas spills: Materials, volumes, and risks to surface waters in four states of the U.S. *Sci.*
Total Environ. **581–582**, 369–377.

McIndoe, R. and Smith, D. G. (1984). Functional anatomy of the internal gills of the tadpole of
Litoria ewingii (Anura, Hylidae). *Zoomorphology.* **104**, 280–291.

Merrick, A. M. and Searle, C. L. (2019). Combined effects of salinity and infectious disease on
Daphnia dentifera at multiple scales. *Freshw. Biol.* **64**, 601–607.

Van Meter, R. J. and Swan, C.M. (2014). Road Salts as Environmental Constraints in Urban
Pond Food Webs. *PLoS One* **9**, e90168.

Park, J.-K. and Do, Y. (2020). Physiological Response of *Pelophylax nigromaculatus* Adults to
Salinity Exposure. *Animals*. **10(9)**, 1698.

Peña-Villalobos, I., Narváez, C. and Sabat, P. (2016). Metabolic cost of osmoregulation in a
hypertonic environment in the invasive African clawed frog *Xenopus laevis. Biol. Open* **5**, 955–
961.

Pivovarov, A. S., Calahorro, F. and Walker, R. J.. (2019). Na+/K+-pump and neurotransmitter
membrane receptors. *Invertebr. Neurosci.* **19**, 1-16.

Pough, F. H. (2007). Amphibian Biology and Husbandry. *ILAR J.* **48**, 203–213.

Reid, B. and Zhao, M. (2014). The electrical response to injury : molecular mechanisms and
wound healing. *Adv. Wound Care.* **3**, 184–201.

Richburg, J. A., Patterson, W. A. and Lowenstein, F. (2001). Effects of road salt and
phragmites Australis invasion on the vegetation of a western Massachusetts calcareous lake-
basin fen. *Wetlands.* **21**, 247–255.

RStudio Team. (2020). RStudio: Integrated Development for R. RStudio, PBC, Boston,
MA. *http://www.rstudio.com/*.

Sanzo, D. and Hecnar, S. J. (2006). Effects of road de-icing salt (NaCl) on larval wood frogs
(*Rana sylvatica*). *Environ. Pollut.* **140**, 247–256.

Schuler, M. S., Hintz, W. D., Jones, D. K., Lind, L. A., Mattes, B. M., Stoler, A. B., Sudol, K.
A. and Relyea, R. A. (2017). How common road salts and organic additives alter freshwater
food webs: in search of safer alternatives. *J. Appl. Ecol.* **54**, 1353–1361.

Seale, D.B., and Wassersug, R. J. (1979). Suspension feeding dynamics of anuran larvae
related to their functional morphology. *Oecologia.* **39**, 259–272.

Semlitsch, R. D., and Skelly, D. K. (2007). Ecology and Conservation of Pool-Breeding
Amphibians. In *Science and Conservation of Vernal Pools in Northeastern North America*. (ed.
A. Calhoun and P. deMaynadier). pp. 127-147. Boca Raton: CRC Press.

Shi, R. and Borgens, R. B. (1995). Three-Dimensional Gradients of Voltage During
Development of the Nervous System as Invisible Coordinates for the Establishment of
Embryonic Pattern. *Dev. Dyn.* **202**, 110-114.

Skou, J. C. (1957). The influence of some cations on an adenosine triphosphatase from
peripheral nerves. *BBA - Biochem. Biophys. Acta* **23**, 394–401.

Squires, Z. E., Bailey, P. C. E., Reina, R. D. and Wong, B. B. M. (2008). Environmental
deterioration increases tadpole vulnerability to predation. *Biol. Lett.* **4**, 392–394.

Uchida, K., Kaneko, T., Yamauchi, K. and Hirano, T. (1996). Morphometrical analysis
chloride cell activity in the gill filaments and lamellae and changes in Na+, K+-ATPase activity
during seawater adaptation in chum salmon fry. *Journal of Experimental Zoology*, **276(3)**, 193-
200.

Uchiyama, M. and Yoshizawa, H. (1992). Salinity tolerance and structure of external and
internal gills in tadpoles of the crab-eating frog, *Rana cancrivora*. *Cell Tissue Res.* **267**, 35–44.

Ultsch, G. R., Bradford, D. F. and Freda, J. (1999). *Tadpoles: the biology of anuran larvae*.

Chicago: University of Chicago Press.

Venâncio, C., Castro, B. B., Ribeiro, R., Antunes, S. C. and Lopes, I. (2019). Sensitivity to
salinization and acclimation potential of amphibian (*Pelophylax perezi*) and fish (*Lepomis*
gibbosus) models. *Ecotoxical. Env. Saf.* **172**, 348-355.

Vences, M. and Köhler, J. (2008). Global diversity of amphibians (Amphibia) in freshwater.
Hydrobiologia **595**, 569–580.

Wilson, J. M. (2011). Role of the gills | Morphology of Branchial Ionocytes. In *Encyclopedia of*
Fish Physiology. Vol. 2. (ed. A. P. Farrell). pp. 1381–1388. Oxford, UK: Elsevier Inc.

Wu, C. S. and Kam, Y. C. (2009). Effects of salinity on the survival, growth, development, and
metamorphosis of *Fejervarya limnocharis* tadpoles living in brackish water. *Zoolog. Sci.* **26**,
476–482.

Wu, C. S., Yang, W K ., Lee, T. H., Gomez-Mestre, I. and Kam, Y.C. (2014). Salinity
acclimation enhances salinity tolerance in tadpoles living in brackish water through increased
Na+, K+-ATPase expression. *J. Exp. Zool. Part A Ecol. Genet. Physiol.* **321**, 57–64.

Figures

Figure 1. Ionic and Osmotic Control of Osmoregulation Across Gill Epithelium.

Mitochondria rich cells (MRC) create a gradient of increasing positive charge through the action

of Na^+/K^+ ATPase (NKA) pumps in the basal cell membrane. 3 Na^+ ions are actively pumped out

of the cell and 2 K^+ ions are pumped in for every molecule of ATP. This creates a chemical

gradient for Na^+ to be drawn into the cytosol through passive Na^+ channels. K^+ is present in

higher concentrations inside the cell and so passively exits through K^+ channels into the plasma

or the external environment. This high concentration of positive ions in the plasma generates a

voltage gradient that allows Cl^- from the environment to pass through the cytosol into the

plasma. The osmolality of the plasma is thus increased and aquaporins present in principle cells

allow the passive diffusion of water.

Figure 2. Variation in Tuft Branch Cell Shape and Density. Gill tuft branch at 7000x magnification (A) with a tuft cell circled in white. Scale bar, 2 μm. Cell shape was measured by tracing cells with well-defined borders and measuring area, perimeter, aspect ratio (B), and circularity (C). Tuft cell density (D) was obtained by tracing cell branches and counting cells within measured area. Points represent individual measurements and were jittered to reduce overlap; violin plots show density distributions, overlayed with a boxplot. Means with SE bars are shown with white diamond. Aspect ratio increased with salinity and circularity decreased. Altered cell shape in high salt treatments coincided with higher cell density.

Figure 3. Gill Tuft Branch Variation in response to salt. Gill tuft (A) with representative tuft branch length (T_L) and tuft branch width (T_W) depicted in white. Scale bar, 10 µm. Each datapoint for tuft dimensions is the average of three measurements taken per trait per tuft branch. Tuft branches are nonuniform in width, measurements were taken at towards the base, midsection, and apex accordingly. Tuft branch length (B) was averaged due to the boundary of the tuft branch at its base being imprecise. Points represent individual measurements and were jittered to reduce overlap; violin plots show density distributions, overlayed with a boxplot. Means with SE bars are shown with white diamond. T_L increased in response to salt, was less densely distributed around the mean and had a wider range of values. T_W was not found to vary in response to salt.

Figure 4. Gill Filter Fold Variation in Response to Salt. Gill plate with filters of epithelial
tissue exposed (A). Scale bar, 100 μm. Main folds (arrows) run ventrodorsally along the gill
plate (right to left in image above). Representative main internode distances (D_M) are depicted
in red, and main fold angle (φ_{MM}) is represented by the red arc between lines. A representative
secondary internode distance (D_S) is depicted in green, and the main fold-secondary fold angle
(φ_{MS}) is depicted by the blue arc. A representative measurement of the distance between main
fold rows (D_{MM}) is shown in white. D_S (B), D_M (C), and D_{MM} (D) distances. Points represent
individual measurements and were jittered to reduce overlap; violin plots show density
distributions, overlayed with a boxplot. Means with SE bars are shown with white diamond. D_S
increased in response to salinity.

Figure 5. Comparison of Gill Tufts from Control and Salt Treatments. (A) Tufts from animals in control treatments were contracted relative to the fuller appearance of (B) tufts from salt treatments. Scale bars,10 μm.

Gill Trait	Description
Gill Tufts	
Branch Length (T_L)	Length of tuft branch from base of branch to tip. Each datapoint for this trait in our study is the average of three such measurements per branch to account for imprecise branch origins.
Branch Width (T_W)	Width of tuft branch. Each datapoint for this trait in our study is the average of three such measurements per branch to account for tapering of branches from broader bases to narrower termini.
Cell Area	Area of gill tuft cells as defined by cell borders at apical epithelium.
Cell Perimeter	Perimeter of gill tuft cells as defined by cell borders at apical epithelium.
Cell Aspect Ratio	Ratio of the major axis, or longest diameter through the cell, to the minor axis, or shortest diameter through the cell.
Cell Circularity	A measure of how closely the perimeter of the cell describes a circle. Given by the formula $$4\pi \frac{Area\ of\ ellipse}{perimeter\ of\ ellipse^2}$$ A circularity value of 1 denotes a perfect circle.
Cell Density	The number of cells present within tuft cell branches.
Gill Filters	
Main Fold Spacing	The distance between rows of main folds. Multiple measures were taken between folds to account for differences arising from the zigzagging morphology of gill filter folds.
Main Fold Internode Distance	The distance between points in main folds where either the main fold changed direction, a secondary fold branched from the main fold, or the main fold terminated.
Main Fold to Main Fold Angle	The angle described by two adjacent sections of main fold, joined by a node, that do not lie on the same line
Secondary Fold Internode Distance	The distance between points on secondary folds where either the secondary node joined the main fold, changed direction, a tertiary fold branched from the secondary fold, or the secondary fold terminated.
Main Fold to Secondary Fold Angle	The angle described by two adjacent sections of main and secondary folds of the gill filter. Angles were always measured from the ventral side of the main fold onto the secondary fold.

674   **Table 1**. Description of gill traits measured in response to salinity.