<u>**CKS1-dependent proteostatic regulation has dual roles combating acute**</u>

<u>**myeloid leukemia whilst protecting normal hematopoiesis**</u>

<u>**Running Title: Dual roles for CKS1 in AML**</u>

Grey, W.[1,*], Rio-Machin, A[2]. Casado-Izquierdo, P.[3], Miettinen, J.J.[4], Copley, F.[2],

Parsons, A.[4], Heckman, C.A.[4], Cutillas, P.[3], Gribben, J.[5], Fitzgibbon, J.[2], Bonnet,

D.[1,*]

1. Haematopoietic Stem Cell Laboratory, The Francis Crick Institute, London, U.K.

2. Centre for Genomics and Computational Biology, Barts Cancer Institute, London,

U.K.

3. Cell signalling and proteomics group, Centre for Genomics and Computational

Biology, Barts Cancer Institute, London, U.K.

4. Institute for Molecular Medicine Finland – FINN, HiLIFE – Helsinki Institute of Life

Science, iCAN Digital Precision Cancer Medicine Flagship, University of Helsinki,

Helsinki, Finland.

5. Centre for Haemato-Oncology, Barts Cancer Institute, London, U.K.

*   Corresponding authors: dominique.bonnet@crick.ac.uk, william.grey@crick.ac.uk

Key Words: CKS1, Proteostasis, AML, Hematopoiesis, Chemotherapy

Word count: 3,764

Figures: 5 main figures, 7 supplementary figures

Tables: 3 data tables, 1 resources table

Abstract

Acute myeloid leukemia (AML) is an aggressive hematological disorder comprising a

hierarchy of quiescent leukemic stem cells (LSCs) and proliferating blasts with

limited self-renewal ability. AML has a dismal prognosis, with extremely low two-year

survival rates in the poorest cytogenetic risk patients, primarily due to the failure of

intensive chemotherapy protocols unable to deplete LSCs, which reconstitute the

disease *in vivo*, and the significant toxicity towards healthy hematopoietic cells.

Whilst much work has been done to identify genetic and epigenetic vulnerabilities in

AML LSCs, little is known about protein dynamics and the role of protein degradation

in drug resistance and relapse. Here, using a highly specific inhibitor of the $SCF^{SKP2-CKS1}$

complex, we report a dual role for CKS1-dependent protein degradation in

reducing AML blasts *in vivo*, and importantly depleting LSCs. Whilst many AML LSC

targeted therapies show significant toxicity to healthy hematopoiesis, inhibition of

CKS1-dependent protein degradation has the opposite effect, protecting normal

hematopoietic cells from chemotherapeutic toxicity. Together these findings

demonstrate CKS1-dependent proteostasis is key for normal and malignant

hematopoiesis.

Significance

CKS1-dependent protein degradation is a specific vulnerability in AML LSCs.

Specific inhibition of $SCF^{SKP2-CKS1}$ is lethal to *CKS1B^{high}* AML blasts and all AML

LSCs. Normal hematopoiesis is protected from chemotherapeutic toxicity by

inhibition of CKS1-dependent protein degradation, substantiating a dual role for

CKS1-dependent protein degradation in clinical treatment of AML.

Introduction

Acute myeloid leukaemia (AML) is a heterogeneous, aggressive disease of the

hematopoietic system, arising from hematopoietic stem/progenitor cells. In recent

years, several reports have demonstrated that the current approach (induction

chemotherapy) and new protocols (epigenetic targeting) still have severe limitations

due to significant plasticity in the AML epigenome, metabolic adaptions and the
presence of drug resistance leukemic stem cells (LSCs). With the average two-year
survival rate as low as 5-15% in poor risk, older patients (>65yr), there is an unmet
critical need for new therapeutic approaches(1). Recent developments, targeting the
anti-apoptotic protein BCL2, has demonstrated that therapies affecting protein
networks holds great promise for the poorest prognosis AMLs(2,3), yet resistance
still emerges for a subset of patients through mitochondrial adaptions in residual
leukemic cells(4,5).

New approaches such as BCL2 targeting in combination with classical induction
chemotherapy still hold excellent promise, but a critical failure is still the severe off-
target toxicity produced by induction chemotherapy protocols and new targeted
therapies alike(6). Indeed, reducing blast count with cytarabine/doxorubicin
treatment severely affects normal hematopoietic progenitor cells, stressing the
hematopoietic system(7). Whilst bone marrow transplantation remains the gold
standard consolidation therapy in AML(8), boosting normal hematopoiesis to
outcompete residual AML, combined with a reduction in severe cytopenia
immediately after chemotherapy, would be beneficial to overall survival.

A better understanding of the biological differences between normal and malignant
hematopoietic cells is needed to achieve selective AML targeting, without toxicity to
normal cells. We previously reported a proteostatic axis between the cyclin-
dependent kinase subunits Cks1 and Cks2, and the mixed lineage leukaemia 1
protein (Mll1), a key protein hijacked during neoplastic transformation of the
hematopoietic system(9). Cks1 and Cks2 have many overlapping and independent
roles in balancing protein homeostasis (proteostasis) throughout the cell cycle,
ensuring correct G0/G1 transition(10), chromatin separation(11–13) and DNA
repair(10,14,15). Whilst it was originally thought Cks1 and Cks2 function solely
through CDK-dependent activities(16–18), CDK-independent functions were later
reported, in concert with the SCF^{SKP2} and APC^{CDC20} E3 ubiquitin ligase complexes,
important for selective protein degradation(10,11,19).

The many functions of Cks1 and Cks2 place this axis at the centre of normal cell
growth and development, and potentially central to AML development. Here, we
investigated the role of CKS1-dependent protein degradation in a poor risk AML
cohort with few treatment options. We explored the therapeutic potential in *CKS1B*
expressing poor risk AML and demonstrated high efficacy in reducing total AML

burden in $CKS1B^{high}$ expressing samples. Critically, in both $CKS1B^{high\ and\ low}$ AML we

demonstrate a significant reduction of the chemotherapy refractory LSCs. In

contrast, CKS1 inhibition has the opposite effect on normal hematopoiesis,

improving stem cell functionality and conferring protection from chemotherapeutic

toxicity.

Results

<u>*CKS1B* expression dictates the susceptibility of AML to specific chemotherapy</u>

The expression of *CKS1B* varies in normal hematopoiesis, with moderate expression

in hematopoietic stem cells (HSCs), the highest expression in myeloid progenitors,

and the lowest expression in terminally differentiated cells (Supp. Fig. 1A). We

previously reported *CKS1B* upregulation in *MLL1*-rearranged AML, acute

lymphoblastic leukemia (ALL) and chronic myeloid leukemia (CML) compared to

peripheral blood mononuclear cells (PBMCs)(9). This trend of *CKS1B* expression is

conserved amongst most AML cytogenetic subtypes (Supp. Fig. 1A), despite high

variability of *CKS1B* expression within AML subtypes, with large standard deviations

compared to one of its key upstream proteostatic regulation partners *SKP2* (Supp.

Fig. 1B). Whilst *CKS1B* expression can be prognostic in a variety of cancers(20–22),

it is not prognostic in AML at the RNA level (Supp. Fig. 1C).

Considering that the key function of CKS1 is substrate recognition, and adapting the

binding of phosphoproteins to their kinase or ubiquitin ligase regulators(18,19), we

hypothesized that high *CKS1B* AMLs may show a selective susceptibility to inhibition

of CKS1-dependent protein degradation (SCF$^{SKP2-CKS1}$ E3 ligase inhibitor), and

inhibition of associated signalling pathways. To address this key question, we

screened a cohort of cytogenetically poor risk AMLs, spanning a variety of FAB and

molecular subtypes, with a broad range of clinically approved and early development

phase compounds and correlated drug sensitivity with *CKS1B* expression (Figure

1A-B, Supp. Table 1).

CKS1B expression significantly correlated with response to 52 individual compounds

including classical DNA damaging chemotherapy (e.g. Cisplatin, R=0.43 p=0.04),

kinase inhibitors (e.g. Ensartinib, R=0.70, p=0.001), BCL2 family inhibitors (e.g.

Sabutoclax, R=0.58, p=0.008) and the SCF$^{SKP2-CKS1}$ E3 ligase inhibitor (hereafter

referred to as CKS1i; R=0.61, p=0.008; Figure 1B-C, Supp. Fig. 1D). Susceptibility to

CKS1i significantly correlated with *CKS1B* expression overall (Supp. Fig. 1D), and
especially well in complex karyotype patients (Supp. Fig. 1E). Separating patients at
the 50th percentile by *CKS1B* expression revealed significantly increased drug
sensitivity in *CKS1B*^{high} AML versus *CKS1B*^{low} AML patients (Figure 1D-E), indicating
that RNA expression of *CKS1B* can be a selection criterion for targeting SCF$^{SKP2-CKS1}$
dependent protein degradation in AML.

The susceptibility of patient AML samples to CKS1i based on *CKS1B* expression is
well conserved in cell lines used to study molecular characteristics of AML. Dose
dependent response (IC$_{50}$) was lower in *CKS1B*^{high} AML cell lines THP-1 (*MLL-AF9*)
and KOPN-8 (*MLL-ENL*) than in *CKS1B*^{low} cell lines ML-1 (*MLL-AF6*) and HL60
(*cMyc*, Figure 1F) and IC$_{50}$ values correlate well with *CKS1B* expression (Figure 1G).
At higher doses, CKS1i induces cell death in both high and low *CKS1B* cell lines
(5μM; Figure 1H), a dose permissive to healthy cells *in vitro*(9). Susceptibility of AML
cell lines to CKS1i is maintained *in vivo* accordingly with *CKS1B* expression.
CKS1B^{high} THP-1 and *CKS1B*^{low} HL60 cells engrafted in immunodeficient mice
(NSG) showed similarly diverging sensitivity to a single course of CKS1i treatment
(10mg/kg, 5 days I.P.). THP-1 leukemic burden in NSG mice was significantly
reduced by CKS1i treatment (Figure 1I), whereas in HL60, leukemic burden
remained similar between control and CKS1i treated mice (Figure 1J). Interestingly,
despite the difference in overall tumor burden after chemotherapy, both xenograft
models showed significant overall improved survival when treated with CKS1i
compared to controls (Figure 1K).

In order to investigate the effect of CKS1i on primary patient AML *in vivo*, we
selected five primary patient samples with a range of *CKS1B* expression (Figure 2A,
Supp. Table 1), for which we have previously reported robust engraftment in NSG
mice(23,24) (Figure 2B). A single course of CKS1i (10mg/kg, 5 days I.P.) was
administered and significantly reduced the AML burden in patients with the highest
CKS1B expression (AML12 and AML21). A trend towards reduced AML burden was
seen at an intermediate level of *CKS1B* expression (AML26), but had no significant
effect was seen on patients with the lowest *CKS1B* expression (AML27 and AML32).
As such, *CKS1B* expression levels correlated well with outcome (R=-0.446; Figure
2C, Supp. Fig 2A). In agreement with our observations of AML cell line *in vivo*
responses, all CKS1i treated AML xenografts showed significantly improved overall

survival compared to untreated controls (Figure 2D), indicating that overall survival
conferred by CKS1i was not only due to changes in bulk leukemic burden.
Whilst reducing leukemic blast count is the current backbone of clinical
chemotherapeutic protocols, typically these approaches do not clear the most
quiescent leukemic stem cells, the subset of cells at the origin of relapse *in*
vivo(25,26). The observed effect on overall survival upon CKS1i treatment in
CKS1B[low] AMLs, without significant reduction of leukemic burden could indicate a
direct effect of CKS1i on LSCs. We sought to assess this by using a leukemic-long-
term culture initiating cell assay (L-LTC-IC)(27).
We selected seven patients with a range of *CKS1B* expression (Figure 2E) for *ex*
vivo study (Figure 2F). After treatment, all surviving cells were re-plated at limiting
dilution to assay the frequency of L-LTC-IC between control and CKS1i-treated cells.
All patient samples showed significant reduction in L-LTC-IC frequency, regardless
of bulk *CKS1B* expression at the start of the experiment (Figure 2G-H, Supp. Fig.
2B-C).
These data indicate that CKS1i can target *CKS1B*[high] AML blasts, but more
importantly is efficient at targeting the leukemic stem cell compartment. Considering
these results, we hypothesized that the overall survival advantage *in vivo* of both
CKS1B[high] and *CKS1B*[low] AMLs is likely due to depletion of LSCs, and therefore a
smaller residual pool of these cells able to repopulate the AML overtime.
182
<u>Normal and malignant hematopoietic cells have divergent responses to CKS1i</u>
To understand the key mechanisms by which CKS1i kills AML, and what effects
CKS1i has on normal hematopoietic cells, we performed mass spectrometry on
CKS1B[high] THP-1 AML cells and umbilical cord blood derived healthy CD34[+] HSPCs
(hereafter referred to as CD34[+]) pre- and post-treatment with CKS1i *in vitro* (1μM;
Figure 3A).
Upon CKS1i treatment, the majority of differentially expressed proteins in THP-1
cells were phosphoproteins (387), which can be alternatively spliced at the RNA
level (375) and also may be acetylated (257), demonstrating integration with post-
translational modification (Figure 3B). CKS1i treatment induced divergent proteomic
profiles in THP-1 and CD34[+] cells, with differentially abundant proteins
predominantly independent between healthy and malignant cells (Figure 3C-E).

Deeper pathway analysis of CKS1i-induced proteomic changes in THP-1 cells
identified the upregulation of the small GTP-binding molecular switch protein RAC1,
and its associated upstream and downstream regulators (Figure 3F-G), that are part
of a pathway known to be involved in regulating cell growth and survival in response
to a variety of external factors(28,29), and important for HSC homing and HSC/LSC
niche interactions(30). Although multiple guanine nucleotide exchange factors
(GEFs) are downregulated in CKS1i treated THP-1 cells, upregulation of the GTP
loaders CRK and Paxilin indicate potential hyperactivity of RAC1, and in keeping
with this, downstream MAP2K3 is also upregulated (Figure 3G).

The relative lack of RAC1 signalling pathway member changes in CD34$^+$ proteomes
are in keeping with RAC1-GTPase activity being upregulated in neoplastic
transformation of hematopoiesis(30) and may identify a specific molecular switch in
AML vulnerable to CKS1 inhibition.

<u>CKS1i drives RAC1-dependent ROS accumulation in AML</u>

The active GTP-bound form of RAC1 is known to bind to p67^{Phox} and catalyzes
NADP/NADPH production. This in turn can lead to upregulation of intracellular
reactive oxygen species (ROS), a process that has been shown to drive apoptosis
and eliminate quiescent LSCs (Figure 3H)(31–33). In AML cell lines, CKS1i
increased total NADP/NADPH levels in a dose dependent manner (Figure 3I, Supp.
Fig. 3A), by increasing the total amount of NADPH compared to control (Figure 3J,
Supp. Fig. 3B). The accumulation of NADPH in AML cell lines can partially be
rescued by inhibiting RAC1 activity (Figure 3I-J, Supp. Fig. 3A-B), validating
increased RAC1-GTP/p67^{Phox} activity upon CKS1 inhibition.

CKS1i-induced NADPH accumulation lead to significantly increased intracellular
ROS in all AML cell lines at sublethal doses of CKS1i, regardless of *CKS1B*
expression (Figure 3K-L). Reversal of this phenotype, by RAC1 inhibition, was most
significant in *CKS1B*^{high} cell lines (Figure 3K-L). Finally, the reduction in AML cell line
viability could be rescued by RAC1 inhibition, with *CKS1B*^{high} cell lines showing the
strongest sensitivity (Supp. Fig. 3C-F). These data indicate that AML requires
SCF$^{SKP2-CKS1}$ functions to regulate RAC1 activity and maintain the fine balance of
intracellular ROS, which are critical for LSC viability.

<u>CKS1 inhibition protects normal hematopoiesis from stress</u>

Since CKS1i is able to selectively kill *CKS1B^high* bulk AML, reduce the LSC
compartment, and prolong AML xenograft survival regardless of leukemic reduction
(Figure 2D), we hypothesized that normal hematopoiesis is spared by CKS1i
treatment.

Proteomic analyses of CD34$^+$ cells treated with CKS1i revealed a clear separation
from THP-1 AML cells (Figure 3C-E). Key proteins differentially abundant in CD34$^+$
cells and not THP-1 cells were integrated in three fundamental pathways in normal
hematopoiesis: Wnt signalling, cell cycle and NFkB signalling (Supp. Fig. 4A-D). The
changes in cell cycle are also consistent with previous reports of *Cks1^-/-* and CKS1i
cellular phenotypes(10,34), and our previously reported suppression of Wnt
signalling in *Cks1^-/-* mouse embryonic fibroblasts(9). The substantial change of
proteins in these key pathways are hallmarks of growth suppression and
differentiation, rather than an induction of cell death by CKS1i, and indeed, treatment
of CD34$^+$ cells *in vitro* with CKS1i resulted in reduced overall cell growth, with a
blockage in cell cycle resulting in increased quiescence of CD34$^+$ cells (Figure 4A-
B).

Considering that suppression of the cell cycle in CD34$^+$ by CKS1i would reduce the
potential for integration of nucleotide analogues and the requirement for
topoisomerase activity, we hypothesized that CKS1i treatment may be "chemo-
protective" for normal hematopoietic cells. To test this, we engrafted normal CD34$^+$
cells in NSG mice and treated mice with the clinical chemotherapy protocol of
cytarabine plus doxorubicin (as has been previously published; 5+3(35)), with or
without CKS1i (Figure 4C). Human CD45$^+$ bone marrow engraftment increased in
untreated control mice between weeks 4 and 6 (Figure 4D). Treatment at week 4,
with doxorubicin/cytarabine (DA) significantly reduced bone marrow engraftment by
week 6, but addition of CKS1i (DAC) was able to rescue this effect, returning
engraftment to comparable levels to control (Figure 4E-G). Better engraftment at
week 6 was complemented by a reduction in apoptotic human CD45 cells in the
bone marrow of recipient mice (Figure 4H-I), indicating that CKS1i treatment
prevents DA-induced cell death in normal hematopoietic cells.

Mechanistically, mass cytometric analysis of CKS1i-treated CD34$^+$ cells revealed a
conserved reduction in key active signalling and transcriptional components that we
recently reported to be important for proliferation and differentiation of HSPCs during
in vitro amplification, including NFkB, PU.1, CREB and mTOR (Supp. Fig. 5)(36),

further indicating an overall block in normal hematopoietic stem cell function. Indeed,
it is important to note that whilst classical cell cycle phosphorylation marks (e.g.
$CDK1^{pT14/Y15}$) are reduced, the protein levels of differentiation regulators are also
reduced (e.g. CREB, PU.1), indicating a potential block in differentiation as well as
growth (Supp. Fig. 5). Additionally, fewer cells have active β-catenin, indicating that
the Wnt pathway – a fundamental pathway that requires a tight balance for normal
hematopoiesis to proceed(37) – is also suppressed (Figure 4J-K). Further
suppression of markers of metabolically active cells (e.g. $mTOR^{pS2448}$), inflammatory
responses (e.g. $NFkB^{pS529}$), and suppression of translation machinery in our mass
spectrometry analyses, lead to a reduction in protein production in CKS1i treated
$CD34^+$ cells (Figure 4L-M). Together, these signalling pathways are fundamentally
linked to the control of ROS accumulation in HSCs, with high activity in these
pathways leading to high intracellular ROS, reduced cell viability and reduced stem
cell potential(38). In agreement, CKS1i treatment reduced intracellular ROS in
cultured $CD34^+$ cells (Figure 4N-O) and *ex vivo* CKS1i treatment increased LTC-IC
frequency by more than two-fold (1 in 1232 vs 1 in 2650, Figure 4P).
Together, these data indicate that the temporary suppression of HSPC activity
conferred by CKS1i leads to suppression of cell growth, protection from general
metabolic stress, improved stem cell functionality and overall increased healthy
hematopoietic capacity during induction chemotherapy.
283
<u>Combining CKS1 inhibition with induction chemotherapy protects normal</u>
<u>hematopoiesis, reduces leukemic stem cells and improves overall survival</u>
The current frontline chemotherapeutic protocols for AML include the use of DA as
induction therapy agents. To test the potential for combining classical DA
chemotherapy with CKS1i *in vivo*, we transplanted NSG mice with primary AML
samples with varying *CKS1B* expression (Figure 5A-B), and after stratifying for
engraftment at week 4, we treated the mice with either DA or DAC. One-week post
chemotherapy, xenografts showed strong reduction in leukemic burden in both DA
and DAC treatment for all AMLs, regardless of *CKS1B* expression, indicating that
CKS1i does not interfere with normal chemotherapeutic killing (Figure 5C). At the
same time point, resident murine CD45 cells co-extracted from aspirated tibias had
significantly higher colony forming potential upon the addition of CKS1i compared to

untreated mice and DA treated mice (Figure 5D), indicating that CKS1i treatment can both selectively reduce AML, whilst protecting normal hematopoiesis from chemotherapeutic toxicity. Overall, all xenografts showed improved overall survival with DA treatment, and in three of the four patients this was significantly improved by the addition of CKS1i, in line with *CKS1B* expression of these patients (Figure 5E-H). Examination of the normal hematopoietic compartment of xenografted mice at the end point of survival, revealed a severe reduction in total number of long-term HSCs (SLAM/LT-HSC) in the DA treated group, whereas addition of CKS1i to DA abolished this effect (Figure 5I). In addition, serial colony forming ability of normal murine hematopoietic cells was improved in DAC conditions, indicating that rescued LT-HSCs are still functional (Figure 5J).

We and others have well documented the refractory nature of LSCs to induction chemotherapy(24,39), and here we set out to investigate the potential conflict or beneficial contribution between DA enrichment of LSCs and CKS1i depletion of LSCs. In *ex vivo* conditions, *CKS1B* high and low AMLs (Figure 5K) showed significant reduction in total cell number one week after DA or DAC treatment (Figure 5L), yet whilst DA treatment enriched for L-LTC-IC frequency in three of the six patients, CKS1i markedly reduced L-LTC-IC frequency in all patients (DAC; Figure 5M & Supp. Fig 6A-B).

Finally, to test the reduction in LSC frequency conferred by CKS1i *in vivo*, we engrafted AML cells obtained from AML26 and AML32 (which had the smallest improvement in overall survival after chemotherapy) in secondary recipients in a limiting dilution manner. Whilst control AMLs retained strong LSC frequency and show robust engraftment after 6 weeks, this was increased by DA treatment in AML26 and was notably reduced in AML32 (Figure 5N-O, Supp. Fig. 7A-B). The addition of CKS1i counteracted the effect of DA by decreasing the LSC frequency in AML26 and further reducing LSC frequency in AML32 compared to control mice, demonstrating strong reduction in LSCs after CKS1i treatment (Figure 5N-O, Supp. Fig. 7A-B). Overall survival was tested for AML32, with DA-AML mice surviving significantly longer than controls, and DAC-AML mice demonstrating a further improvement, with no overt signs of sickness at 120 days in all but one case at the highest cell dose (Figure 5P).

328   Together, these data indicate that inhibition of the SCF$^{SKP2-CKS1}$ complex has dual
329   roles, reducing the viability of AML, and importantly of the LSC fraction, whilst
330   protecting normal HSPCs from chemotherapeutic stress.
331
332   *Discussion*
333
In this study, we demonstrate that proteostatic targeting of the SCF$^{SKP2-CKS1}$ E3
ubiquitin ligase complex selectively eliminates *CKS1B*^{high} AML blasts and reduced
the LSCs compartment of *CKS1B*$^{high \& low}$ AMLs, while sparing normal hematopoietic
cells from chemotherapeutic toxicity.
Poor risk AML is a heterogeneous group that includes patients with different
cytogenetic abnormalities, very limited treatment options and extremely low overall
survival rates(1,40), even accounting for newer therapies (Venetoclax plus
Azacitidine)(2,3). The selective reduction in viability of *CKS1B*^{high} AML blasts by
CKS1 inhibition (Figure 1D) indicate that, whilst *CKS1B* is not predictive of overall
survival at the RNA level, proteostatic vulnerabilities exist in AML and can be
identified through better understanding of leukemic proteomes. While gene
expression profiles, particularly those with single cell resolution, are improving our
understanding of AML heterogeneity, the origins of leukemic relapse and revealing
new clinical targets(25,26), the role of proteostasis has been comparatively
understudied(41,42).
In vivo, AML blasts and LSCs are relatively quiescent compared to AML cell lines
and patient samples screened *in vitro*, so the consistently high sensitivity of
CKS1B^{high} AMLs observed *in vivo* (Figure 2C) and the reduction of LSC frequency
regardless of the bulk *CKS1B* expression (Figure 2E-H), demonstrates a role for the
SCF$^{SKP2-CKS1}$ complex beyond targeting highly proliferative cells. Therefore, the role
of CKS1 likely reaches further than cell cycle regulation, as has been previously
reported(10,43). Here we provide evidence for CKS1 regulating RAC1/NADPH/ROS
signalling (Figure 3H), a fundamental pathway involved in amplifying extrinsic and
intrinsic signals in normal hematopoiesis and AML(4,44). The balance of intracellular
ROS in normal and malignant hematopoietic cells has been of great interest in
recent years(33,38), and changes in mitochondrial functions due to *RAS* mutations
and nicotinamide-NAD metabolism underline the critical role for this pathway in
primary patient resistance to Venetoclax(4,5). The induction of ROS in AML cell lines

upon CKS1 inhibition, regardless of *CKS1B* expression, demonstrates that the
balance of CKS1-dependent protein degradation is key to maintaining stress
responses in AML. This, together with LSCs requiring low ROS to maintain their
stem cell potential, would explain the strong reduction in LSC frequency conferred by
CKS1i in primary patient AML (Figure 2G-H & 5M-O).

The effect of CKS1i on normal hematopoiesis is clearly separate to AML (Figure 3C).
Indeed, cell cycle blockage was suggested by the relatively few protein changes
induced by CKS1i in CD34$^+$ but not AML cells. This is highly beneficial, as patients
treated with induction chemotherapy, which targets cycling cells, suffer from severe
toxicity and cytopenia upon treatment. Classical induction chemotherapy is known to
reduce the pool of hematopoietic progenitors, whilst quiescent HSCs are refractory
to treatment, but ultimately undergo senescence(7,45). We found that cell cycle
arrest of HSPCs by CKS1i could prevent DA reduction of normal cells *in vivo* (Figure
4D-G), and in the context of AML could rescue the reduction in HSCs induced by
chemotherapy (Figure 5I). Importantly, CKS1i treatment also induced changes in
fundamental HSPC signalling pathways known to be involved in stem cell potency
and response to stress. Indeed, we have previously shown that activating NFκB
signalling can reduce intracellular ROS and improve HSC outgrowth *in vitro*(36). The
overall suppression of key growth and activation cellular markers lead to an opposite
phenotype to that seen in AML cells, with a reduction in intracellular ROS and an
increase in normal HSC frequency (Figure 4N-P). These divergent phenotypes
between normal and malignant hematopoietic cells indicate that CKS1-dependent
protein degradation is required for the growth of normal cells and that CKS1
suppression pushes cells towards quiescence while retaining stem cell functions. On
the other hand, inhibiting CKS1-dependent protein degradation in AML can lead to
incorrect regulation of signalling, ultimately causing cellular toxicity, as shown before
for more broader regulators of the SCF complex(46).

The backbone of clinical chemotherapeutic protocols has largely remained
unchanged over the last 20 years, with induction chemotherapy reducing AML blasts
to prolong survival(47). Whilst reduction of AML burden is beneficial, classical
chemotherapy actually maintains or even enriches for LSCs and results in relapsed
AML, often with increased mutational burden(48). The addition of CKS1i to DA
results in a significant reduction in LSC frequency. Considering that resistant LSCs

can be traced as the origin of relapse(25,26), and constitute a key route to
chemoresistance in AML, CKS1 inhibition has excellent promise for eradicating
these cells.

The inhibition of CKS1-dependent protein degradation holds excellent promise for
AML therapy, both as a targeted agent towards *CKS1B^high* AML, and in combination
with induction chemotherapy where protection of healthy cells is key. Reports of
CKS1B overexpression correlating with outcome in other solid cancer
types(20,22,49) and novel ways to modulate CKS1 function(50), together with our
findings of a dual role for CKS1 inhibition in AML and the development of more
clinically ready molecules to target CKS1(51), indicate that proteostatic targeting,
through the CKS1/CKS2 axis, holds much hope for future cancer therapy.

Methods (online unlimited)

<u>Primary AML and UCB samples</u>

AML samples were obtained after informed consent at St Bartholomew's Hospital
(London, U.K.) at the time of diagnosis as part of the Bart's Cancer Institute Poor-
Risk AML consortium. Full details of patient information are provided in
Supplementary Table 1. Live mononuclear cells (MNCs) were isolated by density
centrifugation using Ficoll-Paque (GE healthcare). Prior to culture or
xenotransplantation AML cells were depleted for T-cells using the Easysep T-cell
depletion kit (StemCell Technologies). Umbilical Cord Blood (UCB) was obtained
from full term donors after informed consent at the Royal London Hospital (London,
U.K.). MNCs were isolated by density centrifugation using Ficoll-Paque (GE
healthcare). Cells were selected for CD34$^+$ using the Easysep CD34$^+$ enrichment kit
(StemCell Technologies). Purity was confirmed by flow cytometry.

<u>Drug sensitivity and resistance testing (DSRT)</u>

Single drug DSRT was performed as described previously(52). In brief, 35 different
compounds, each with 7 different concentrations (Supp. Table. 2), were pre-plated
using an acoustic liquid handling Echo 550 (Labcyte) to 384-well plates. Drug plate
well annotations and drug concentrations are presented in Supp. Table. 3. Primary

AML cells were suspended in conditioned medium (RPMI 1640 supplemented with
10% foetal bovine serum, 2mM L-glutamine, penicillin-100U/ml, streptomucing-
100ug/ml and 12.5% conditioned medium from HS-5 human bone marrow stromal
cells), DNase I treated for 4h (Promega), filtered through a 70μm cell strainer (Fisher
Scientific) to remove possible cell clumps, and viable cells were counted. Pre-plated
compounds in each 384-well plate were dissolved in 5ul of conditioned medium
using a MultiDrop Combi peristaltic dispenser (Thermo Scientific) and shaken for 5
minutes to dissolve the compounds. AML cells were plated at 5,000 cells/well in
20ul, leading to a final volume of 25ul/well. Plates were gently shaken for 5 minutes
to mix the cells with the compounds and incubated for 72 hours at 37C, 5% CO_2.
Cell viability was measured using the CellTiter-Glo assay (Promega) with a
PHERAstar microplate reader (BMG-labtech). Data was normalised to negative
(DMSO only) and positive control wells (100uM benzethonium chloride) and dose
response curves calculated.
Ex vivo drug sensitivity of AML cells to the tested drugs was calculated using a drug
sensitivity score (DSS), a modified form of the area under the inhibition curve
calculation that integrates multiple dose response parameters for each of the tested
drugs, as previously described(53).
447
448 <u>AML cell line, UCB CD34[+] and MS-5 culture</u>
449
All AML cell lines and MS-5 stromal cells were originally obtained from the ATCC
and maintained by the Francis Crick Cell Services. All AML cell lines were cultured in
RMPI 1640, 10% heat-inactivated FBS and 1% penicillin/streptomycin (Life
Technologies) at 37C, 5% CO_2. Umbilical cord blood CD34[+] cells were cultured in
StemSpan SFEMMII (StemCell Technologies) supplemented with Human SCF
(150ng/ml), Human FLT3 ligand (150ng/ml) and Human TPO (20ng/ml; all
Peprotech) at $2x10^5$ cells/ml at 37C, 5% CO_2. For relative viability, apoptosis and
IC_{50} calculations cell lines were seeded in 96 well plates at a concentration of $2x10^5$
cells/ml with the indicated dose of drug. Measurements of viability (% reduction O_2)
or apoptosis (Annexin V positivity) were taken at 48 hours post treatment. MS-5
stromal cells were cultured in IMDM, 10% heat-inactivated FBS and 2%
penicillin/streptomycin (Life Technologies) at 37C, 5% CO_2.

 <u>Publicly available datasets</u>

CKS1B expression in normal and malignant hematopoiesis was obtained through

Bloodspot.eu. Overall survival and stratification for *CKS1B* expression was

calculated from data obtained from The Cancer Genome Atlas (TCGA). AML cell line

RNA sequencing data was obtained from the EBI Expression Atlas (RNA-seq of 934

Human cancer cell lines from the Cancer Cell Line Encyclopedia).

<u>Leukemic/Normal Long-term culture initiating cell (L-LTC-IC) assay</u>

These experiments were performed as originally published by our group(27).

For all co-culture experiments, MS-5 stromal cells were seeded two days prior to

AML/UCB cell addition at $4x10^5$ cells/ml to reach confluence at the time of irradiation.

One day prior to AML/UCB addition, MS-5 stromal cells were irradiated with 7Gy and

culture media was exchanged. On the day of starting co-culture, AML cells were

plated at $2x10^5$ cells/ml in meylocult H5100 (StemCell Technologies) supplemented

with IL-3, G-CSF and TPO (all 20ng/ml; Peprotech). UCB cells were plated at $2x10^5$

cells/ml in myelocult H5100 (StemCell Technologies). Half media changes were

performed once per week without disrupting the feeder layer. At the start of week

two, indicated drug treatments were added at 2x concentration in the half media

change.

For LTC-CAFC assays, all cells were harvested at day 14 and sorted for live

hCD45$^+$mSca-1$^-$ cells. Resulting cells were seeded in co-culture with fresh MS-5

stromal cells in a 96 well plate in a limiting dilution range (200,000 to 1,000) in 10

replicates and cultured for a further 5 weeks. At the end of the co-culture period

cobblestone area forming cells were scored and L-LTC-IC frequency was calculated

using the ELDA (Extreme Limiting Dilution Analysis) function in the Statmod R

package.

For LTC-IC assays, media was continuous changed each week until week five

Cultures were harvested and sorted for live hCD45$^+$mSca-1$^-$ cells. Resulting cells

were seeded in co-culture with fresh MS-5 stromal cells in a 96 well plate in a limiting

dilution range (10,000 to 100) in 10 replicates and cultured for a further three weeks.

At week eight, myelocult H5100 was replaced with Methocult methycellulose

(StemCell Technologies H4434) for a further two weeks, after which wells were
scored for colony-forming units and LTC-IC frequency was calculated using the
ELDA (Extreme Limiting Dilution Analysis) function in the Statmod R package.
499

Patient derived xenografts (PDX) and *in vivo* drug treatment

Primary AML samples (1×10^6 – 5×10^6 cells total) or UCB-CD34$^+$ (5×10^4 cells total)
were injected intravenously (I.V.) into unconditioned 10-12 week old female or male
NOD-SCID IL2Rynull (NSG) mice (The Jackson laboratory). After 4 weeks
engraftment was assessed by bone marrow aspiration from long bones whilst mice
were under isoflurane anaesthesia. Mice were stratified according to engraftment
and sex and assigned to treatment and control groups accordingly. Mice were
treated as indicated with 10mg/kg CKS1i (Skp2-Cks1 E3 ligase inhibitor, Merck
Millipore) intraperitoneal injection (I.P.) for 5 days, DA (doxorubicin/cytarabine,
1.5mg/kg/10mg/kg respectively, Sigma Aldrich), doxorubicin on days 1-3, cytarabine
on days 1-5 co-injected I.V.(35). Mice were scored for engraftment over the
experimental course by bone marrow aspiration and for overall survival according to
U.K. home office license protocols and following CRUK guidance (>20% peak body
weight loss, overt signs of sickness/mortality).

AML cell line *in vivo* experimentation

AML cell lines were transduced with GFP-Luciferase containing vectors as per our
previous reports(24). For both cell lines (THP-1 and HL60) 2×10^6 cells were injected
I.V. into unconditioned 10-12 weeks old female or male NOD-SCID IL2Rynull (NSG)
mice (The Jackson laboratory). After 7 days engraftment was assessed by
bioluminescene imaging. Isofluorane anaesthetised mice were imaged 5-10 minutes
post D-luciferin injection I.P. (15mg.kg; Caliper life sciences) using the Xenogen IVIS
imaging system. Photons emitted were expressed as Flux (photons/s/cm^2), and
quantified and analysed using "living image" software (Caliper life sciences).

RNA extraction, reverse transcription and real time quantitative PCR (RT-qPCR)

Total RNA was isolated from patient samples after thawing, density centrifugation
and T-cell depletion, using a RNeasy mini kit (Qiagen). Resulting RNA was reverse
transcribed to produce cDNA using the Superscript III reverse transcriptase kit
(Thermo Scientific) with oligoDT$_{20}$ primers (Sigma Aldrich). RT-qPCR experiments
were performed with an ABI-7500 FAST Thermal Cycler (Applied Biosystems) using
SYBR Green (Thermo Scientific). RNA abundance was quantified by the
Comparative CT method with two independent control genes (*GAPDH* and *B-ACTIN*,
GAPDH presented). The CT values used for each patient sample were the result of
three technical triplicates. Primers are described in the resources table.
538
539 <u>Mass Spectrometry</u>
540
THP-1 AML cell lines and UCB CD34^{+} cells were cultured as per culture and drug
treatment above. Cells were recovered for 24 hours in their respective media
followed by sub-lethal AML doses of CKS1i (1μM) for 12 hours. All cells were
retrieved from wells, washed three times in ice-cold PBS and snap frozen in liquid
nitrogen as dry pellets.
Cell pellets were lysed in 100 μL of urea buffer (8 M urea in 20 mM HEPES, pH:
8.0), lysates were further homogenized by sonication (30 cycles of 30s on 30s off;
Diagenode Bioruptor® Plus) and insoluble material was removed by centrifugation.
Protein amount was quantified using BCA (Thermo Fisher Scientific). Then, 100 and
20 μg of protein for THP-1 and CD34^{+} samples, respectively, were diluted in urea
buffer to a final volume of 300 μL and subjected to cysteine alkylation using
sequential incubation with 10 mM dithiothreitol (DDT) and 16.6 mM iodoacetamide
(IAM) for 1 h and 30 min, respectively, at 25 °C with agitation. Trypsin beads (50%
slurry of TLCK-trypsin; Thermo-Fisher Scientific; Cat. #20230) were equilibrated with
3 washes with 20 mM HEPES (pH 8.0), the urea concentration in the protein
suspensions was reduced to 2 M by the addition of 900 μL of 20 mM HEPES (pH
8.0), 100 μL of equilibrated trypsin beads were added and samples were incubated
overnight at 37°C. Trypsin beads were removed by centrifugation (2000 xg at 5°C for
5 min) and the resulting peptide solutions were desalted using carbon C18 spin tips
(Glygen; Cat. # TT2MC18). Briefly, spin tips were activated twice with 200 μL of
Elution Solution (70% ACN, 0.1% TFA) and equilibrated twice with 200 μL of Wash

Solution (1% ACN, 0.1% TFA). Samples were loaded and spin tips were washed

twice with 200 µL of Wash Solution. Peptides were eluted into fresh tubes from the

spin tips with 4 times with 50 µl of Elution Solution. In each of the desalting steps,

spin tips were centrifuged at 1,500xg at 5C for 3 min. Finally, samples were dried in

a SpeedVac and peptide pellets were stored at –80°C.

For mass spectrometry identification and quantification of proteins, samples were run

twice in a LC-MS/MS platform. Briefly, peptide pellets were resuspended in 100 µL

and 20 µL of reconstitution buffer (20 fmol/µL enolase in 3% ACN, 0.1% TFA) for

THP-1 and CD34$^+$ samples, respectively. Then, 2 µL were loaded onto an LC-

MS/MS system consisting of a Dionex UltiMate 3000 RSLC coupled to a Q

Exactive™ Plus Orbitrap Mass Spectrometer (Thermo Fisher Scientific) through an

EASY-Spray source (Cat. # ES081, Thermo Fisher Scientific). Mobile phases for the

chromatographic separation of the peptides consisted in Solvent A (3% ACN: 0.1%

FA) and Solvent B (99.9% ACN; 0.1% FA). Peptides were loaded in a micro-pre-

column (Acclaim™ PepMap™ 100 C18 LC; Cat. # 160454, Thermo Fisher Scientific)

and separated in an analytical column (Acclaim™ PepMap™ 100 C18 LC; Cat. #

164569, Thermo Fisher Scientific) using a gradient running from 3% to 23% over 120

min. The UPLC system delivered a flow of 2 µL/min (loading) and 300 nL/min

(gradient elution). The Q-Exactive Plus operated a duty cycle of 2.1s. Thus, it

acquired full scan survey spectra (m/z 375–1500) with a 70,000 FWHM resolution

followed by data-dependent acquisition in which the 15 most intense ions were

selected for HCD (higher energy collisional dissociation) and MS/MS scanning (200–

2000 m/z) with a resolution of 17,500 FWHM. A dynamic exclusion period of 30s was

enabled with a m/z window of ±10 ppms.

Peptide identification from MS data was automated using a Mascot Daemon 2.5.0

workflow in which Mascot Distiller v2.5.1.0 generated peak list files (MGFs) from

RAW data and the Mascot search engine (v2.5) matched the MS/MS data stored in

the MGF files to peptides using the SwissProt Database (SwissProt_2016Oct.fasta).

Searches had a FDR of ~1% and allowed 2 trypsin missed cleavages, mass

tolerance of ±10 ppm for the MS scans and ±25 mmu for the MS/MS scans,

carbamidomethyl Cys as a fixed modification and PyroGlu on N-terminal Gln and

oxidation of Met as variable modifications. Identified peptides were quantified using

Pescal software in a label free procedure based on extracted ion chromatograms

(XICs). Thus, the software constructed XICs for all the peptides identified across all
samples with mass and retention time windows of ±7 ppm and ±2 min, respectively
and calculated the area under the peak. Individual peptide intensity values in each
sample were normalized to the sum of the intensity values of all the peptides
quantified in that sample. Data points not quantified were given a peptide intensity
value equal to the minimum intensity value quantified in the sample divided by 10.
Protein intensity values were calculated by adding the individual normalized
intensities of all the peptides comprised in a protein and values of 2 technical
replicates per sample were averaged. Protein score values were expressed as the
maximum Mascot protein score value obtained across samples. The mass
spectrometry proteomics data have been deposited to the ProteomeXchange
Consortium via the PRIDE partner repository (PXD022754 and
10.6019/PXD022754).

<u>Flow Cytometry, apoptosis and cell cycle assays</u>

Flow cytometry analysis was performed using a BD Fortessa flow cytometer (BD
biosciences). Cells were prepared by washing in PBS + 1% FBS three times before
staining in the same media with the indicated cell surface antibodies (resources
table) for 1 hour at 4C. For apoptosis assays, cells were incubated with annexin V
binding buffer in addition to the washing media (BD biosciences), washed three
times in PBS + 1% FBS + 1x annexin V binding buffer and incubated with 0.1μg/ml
DAPI prior to flow cytometry analysis. For cell cycle analysis, cells were washed
three times in PBS + 1% FBS and fixed in BD fix/perm buffer (BD biosciences) for 20
minutes at room temperature. Cells were washed three times in BD perm/wash
buffer (BD biosciences) and incubated with anti-Ki67 antibody for 4 hours at 4C.
Cells were washed three times in BD perm/wash buffer and 0.5 μg/ml DAPI was
added for 15 minutes prior to analysis. For all flow cytometry, cells were initially
identified based on forward and side scatter.

<u>Viability assays</u>

Relative cell viability was assessed by % reduction O_2 in culture wells using the

Alamar blue cell viability reagent (Life Technologies). Cells were seeded in 96 well

plates at $2x10^5$ cells/ml and the indicated dose of drugs were added on top and

incubated for 48 hours. Alamar blue reagent was added on top of cells and they

were continued to be incubated for 4 hours under the same conditions (37C, 5%

CO_2). Plates were read on a spectramax plate reader (Biostars) at 570nm and

600nm and % reduction O_2 was calculated as per the manufacturer's instructions.

<u>NADP/NADPH assays</u>

Total $NADP^+$ and NADP/H were measured using the NADP/NADPH colorimetric

assay kit (Abcam). AML cell lines were seeded at $2x10^5$ cells/ml one day prior to

treatment with the indicated drugs (day 0). The following day (day 1), the indicated

concentration of drugs were added to culture wells. The next day (day 2) all cells

were collected from the wells and washed three times in ice-cold PBS. Cells were

lysed in NADP/NADPH extraction buffer by performing two freeze/thaw cycles (20

mins on dry ice followed by 10 mins at room temperature). Lysates were centrifuged

at 13,000g for 10minutes and the supernatant was retained. Lysate supernatant was

split in half, with one half remaining on ice and the other half incubated at 60C for

30mins to remove $NADP^+$. Total NADP/H (NADPt) and NADPH only lysates were

run in 96 well plates with freshly made standards as per the manufacturers'

instructions. NADP/NADPH ratio was calculated as (NADPt-NADPH)/NADPH.

<u>Intracellular ROS staining</u>

Intracellular reactive oxygen species were assayed using the CellRox deep red

reagent (Life Technologies). AML cell lines were seeded at $2x10^5$ cells/ml one day

prior to treatment with the indicated drugs (day 0). The following day (day 1), the

indicated concentration of drugs were added to culture wells. The next day (day 2),

CellRox deep red was added to each well at a final concentration of 5uM and

verapamil was added at a final concentration of 50 μM. Cells were continued to be

incubated in the same conditions (37C, 5% CO_2) for 1hr. After incubation, cells were

collected from wells and washed three times in PBS + 1%FBS + 50 μM verapamil

and finally resuspended in PBS + 1% FBS + 50 μM verapamil + DAPI (0.1μg/ml)

before analysis on a BD Fortessa FACS analyser.

Mass Cytometry

CyTOF preparation and analysis was carried out as per our previous publication[36].

Cultured cells were washed in ice-cold PBS three times and incubated with 5uM

Cisplatin (Fluidigm) to mark dead cells. Cells were washed three times in ice-cold

PBS and fixed in 1.6% formaldehyde (Sigma Aldrich). Fixed cells were surface

stained with the relevant antibodies (resources table) for two hours at room

temperature followed by three washes with PBS. Cells were permeabilised in 1ml

Perm buffer III (BD biosciences) on ice for 30mins, washed three times in ice-cold

PBS and incubated with the relevant intracellular antibodies (resources table)

overnight at 4C with gentle rotation. Resulting cells were wash three times in ice-cold

PBS and stained with 100nM Iridium in PBS + 0.1% Saponin (Riedel-de Haen)

overnight before analysis on a Helios Mass Cytometer (Fluidigm).

Protein translation assays

Protein translation was measured using the OP-Puromycin protein translation kit

(Life Technologies). AML cell lines were seeded at 2×10^5 cells/ml one day prior to

treatment with the indicated drugs (day 0). The following day (day 1), the indicated

concentration of drugs were added to culture wells. The next day (day 2), 10 μM OP-

Puromycin was added to culture wells for one hour under culture conditions (37C,

5% CO_2). Cells were washed three times in ice-cold PBS and fixed in 4%

paraformaldehyde (Sigma Aldrich) at room temperature for 15 mins in the dark. Cells

were washed three times in PBS and permeabilised in PBS + 0.5% Triton X-100

(Sigma Aldrich) for 15 mins. Cells were washed twice in Click-IT reaction buffer wash

solution and stained as per the manufacturer's instructions (Life Technologies).

Abundance of OP-Puromycin was assessed using flow cytometry.

Colony forming units

For resident mouse hematopoietic cell response to 5-FU', CKS1i, DA and DAC,

colony forming ability was assessed in methylcellulose (StemCell Technologies

M3434-GF). 10^4 mCD45$^+$ cells were sorted from PDX mice at the indicated points

and seeded in methylcellulose and scored to colony forming units after 7 days.

Cultures were dissolved in PBS, counted and 10^4 cells were re-seeded for passage

2 and passage 3.

<u>Statistics and data interpretation</u>

Results shown are +/-SEM unless otherwise indicated. To compare treatment versus

control in all *in vitro* and *in vivo* experiments, a Student's *t*-test was used as indicated

in the figure legend with N number indicated. For all comparisons, unpaired *t*-tests

were undertaken unless otherwise indicated. All repeat samples presented are from

biological replicates of distinct samples/xenotransplantations.

Survival analyses were carried out using the "survminer" package on R to calculate

significance between Kaplan-Meier curves and Hazard ratios. Kaplan Meier graphs

were plotted using Graphpad Prism.

Correlation analyses were carried out using the "performance analytics" and

"corrplot" packages in R. Multiple DSS comparisons with *CKS1B* expression were

carried out with pairwise complete observations using Spearman, Pearson and

Kendall correlation coefficients. Individual correlations for *CKS1B* vs DSS or IC$_{50}$

were plotted using Graphpad Prism.

Stem cell frequency was calculated using the extreme limiting dilution analysis

(ELDA) function in the "statmod" R package(54).

Pathway analysis and enrichment was run through MetaCore (genego.com) and

network interactions produced on String (string-db.org)

Acknowledgements

We would like to acknowledge the Francis Crick core flow cytometry and biological

research facility STPs. The Francis Crick Institute receives its core funding from

Cancer Research UK (FC001115), the UK Medical Research Council (FC001115)

and the Wellcome Trust (FC001115). We would like to acknowledge Drs E.

Grönroos, R. Hynds, S Ali and H Wood & Prof. P. Parker for their critical feedback on
the manuscript.
727
Author Contributions
W.G. Conceived the study, designed and carried out experiments, analyzed data
and wrote the manuscript. A.R-M. Analyzed patient data. P.C-I. Carried out mass
spectrometry analyses. J.J.M. Designed and carried out experiments. F.C. Analyzed
patient data. A.P. Designed and carried out experiments. C.A.H. Supervised drug
screening. P.C. Supervised mass spectrometry analyses. J.G. Provided patient
samples and data. J.F. Provided patient samples and data. D.B. Conceived, write
manuscript and supervised the study. All authors provided critical feedback on the
manuscript pre-submission.
737

738   *References*
739
1.   Döhner H, Weisdorf DJ, Bloomfield CD. Acute Myeloid Leukemia. Longo DL,
editor. N Engl J Med 2015;373:1136–52. Available from:
http://www.nejm.org/doi/10.1056/NEJMra1406184
2.   DiNardo CD, Tiong IS, Quaglieri A, MacRaild S, Loghavi S, Brown FC, et al.
Molecular patterns of response and treatment failure after frontline venetoclax
combinations in older patients with AML. Blood 2020;135:791–803. Available
from: http://ashpublications.org/blood/article-
pdf/135/11/791/1718665/bloodbld2019003988.pdf
3.   DiNardo CD, Jonas BA, Pullarkat V, Thirman MJ, Garcia JS, Wei AH, et al.
Azacitidine and Venetoclax in Previously Untreated Acute Myeloid Leukemia.
N Engl J Med 2020;383:617–29. Available from:
https://www.nejm.org/doi/full/10.1056/NEJMoa2012971
4.   Jones CL, Stevens BM, Pollyea DA, Culp-Hill R, Reisz JA, Nemkov T, et al.
Nicotinamide Metabolism Mediates Resistance to Venetoclax in Relapsed
Acute Myeloid Leukemia Stem Cells. Cell Stem Cell. 2020
5.   Stevens BM, Jones CL, Pollyea DA, Culp-Hill R, D'Alessandro A, Winters A, et
al. Fatty acid metabolism underlies venetoclax resistance in acute myeloid
leukemia stem cells. Nat Cancer 2020;1–12. Available from:

http://www.nature.com/articles/s43018-020-00126-z

6.   Estey E, Levine RL, Löwenberg B. Current challenges in clinical development

of "targeted therapies": The case of acute myeloid leukemia. Blood. 2015.

7.   Shao L, Wang Y, Chang J, Luo Y, Meng A, Zhou D. Hematopoietic stem cell

senescence and cancer therapy-induced long-term bone marrow injury. Transl

Cancer Res. 2013;

8.   Vyas P, Appelbaum FR, Craddock C. Allogeneic hematopoietic cell

transplantation for acute myeloid leukemia. Biol. Blood Marrow Transplant.

2015.

9.   Grey W, Ivey A, Milne TA, Haferlach T, Grimwade D, Uhlmann F, et al. The

Cks1/Cks2 axis fine-tunes Mll1 expression and is crucial for MLL-rearranged

leukaemia cell viability. Biochim Biophys Acta - Mol Cell Res 2018;1865:105–

16. Available from:

http://linkinghub.elsevier.com/retrieve/pii/S0167488917302525

10.   Frontini M, Kukalev A, Leo E, Ng Y-MM, Cervantes M, Cheng C-WW, et al.

The CDK Subunit CKS2 Counteracts CKS1 to Control Cyclin A/CDK2 Activity

in Maintaining Replicative Fidelity and Neurodevelopment. Dev Cell

2012;23:356–70. Available from:

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dop

t=AbstractPlus&list_uids=22898779

11.   Wolthuis R, Clay-Farrace L, van Zon W, Yekezare M, Koop L, Ogink J, et al.

Cdc20 and Cks direct the spindle checkpoint-independent destruction of cyclin

A. Mol Cell 2008;30:290–302. Available from:

http://www.ncbi.nlm.nih.gov/pubmed/18471975

12.   Martinsson-Ahlzen HS, Liberal V, Grunenfelder B, Chaves SR, Spruck CH,

Reed SI, et al. Cyclin-dependent kinase-associated proteins Cks1 and Cks2

are essential during early embryogenesis and for cell cycle progression in

somatic cells. Mol Cell Biol 2008;28:5698–709. Available from:

http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2546922&tool=pmc

entrez&rendertype=abstract

13.   Morris MC, Kaiser P, Rudyak S, Baskerville C, Watson MH, Reed SI. Cks1-

dependent proteasome recruitment and activation of CDC20 transcription in

budding yeast. Nature. 2003;423:1009–13. Available from:

http://www.ncbi.nlm.nih.gov/pubmed/12827207

14.   Del Rincón S V, Widschwendter M, Sun D, Ekholm-Reed S, Tat J, Teixeira LK,

et al. Cks overexpression enhances chemotherapeutic efficacy by overriding

DNA damage checkpoints. Oncogene 2014 1–7. Available from:

http://www.ncbi.nlm.nih.gov/pubmed/24858038

15.   Liberal V, Martinsson-Ahlzén H-S, Liberal J, Spruck CH, Widschwendter M,

McGowan CH, et al. Cyclin-dependent kinase subunit (Cks) 1 or Cks2

overexpression overrides the DNA damage response barrier triggered by

activated oncoproteins. Proc Natl Acad Sci U S A 2012;109:2754–9. Available

from:

http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3286935&tool=pmc

entrez&rendertype=abstract

16.   Patra D, Wang SX, Kumagai a, Dunphy WG. The xenopus Suc1/Cks protein

promotes the phosphorylation of G(2)/M regulators. J Biol Chem

1999;274:36839–42. Available from:

http://www.ncbi.nlm.nih.gov/pubmed/10601234

17.   Patra D, Dunphy WG. Xe-p9, a Xenopus Suc1/Cks homolog, has multiple

essential roles in cell cycle control. Genes Dev 1996;10:1503–15. Available

from: http://www.genesdev.org/cgi/doi/10.1101/gad.10.12.1503

18.   McGrath D a, Balog ERM, Kõivomägi M, Lucena R, Mai M V, Hirschi A, et al.

Cks confers specificity to phosphorylation-dependent CDK signaling pathways.

Nat Struct Mol Biol 2013;20:1407–14. Available from:

http://www.ncbi.nlm.nih.gov/pubmed/24186063

19.   Spruck C, Strohmaier H, Watson M, Smith APL, Ryan A, Krek W, et al. A

CDK-independent function of mammalian Cks1: Targeting of SCFSkp2 to the

CDK inhibitor p27Kip1. Mol Cell. 2001;7:639–50.

20.   Shen D-YY, Fang Z-XX, You P, Liu P-GG, Wang F, Huang C-LL, et al. Clinical

significance and expression of cyclin kinase subunits 1 and 2 in hepatocellular

carcinoma. Liver Int 2010;30:119–25. Available from:

http://www.ncbi.nlm.nih.gov/pubmed/19845855

21.   Kitajima S, Kudo Y, Ogawa I, Bashir T, Kitagawa M, Miyauchi M, et al. Role of

Cks1 overexpression in oral squamous cell carcinomas: cooperation with Skp2

in promoting p27 degradation. Am J Pathol. 2004;165:2147–55.

22.   Masuda T-A, Inoue H, Nishida K, Sonoda H, Yoshikawa Y, Kakeji Y, et al.

Cyclin-dependent kinase 1 gene expression is associated with poor prognosis

in gastric carcinoma. Clin Cancer Res. 2003;9:5693–8.

23. Passaro D, Di Tullio A, Abarrategi A, Rouault-Pierre K, Foster K, Ariza-

McNaughton L, et al. Increased Vascular Permeability in the Bone Marrow

Microenvironment Contributes to Disease Progression and Drug Response in

Acute Myeloid Leukemia. Cancer Cell 2017;32:324-341.e6. Available from:

http://www.ncbi.nlm.nih.gov/pubmed/28870739

24. Tullio A Di, Rouault-Pierre K, Abarrategi A, Mian S, Grey W, Gribben J, et al.

The combination of CHK1 inhibitor with G-CSF overrides cytarabine resistance

in human acute myeloid leukemia. Nat Commun 2017;8:1679. Available from:

http://www.ncbi.nlm.nih.gov/pubmed/29162833

25. Boyd AL, Aslostovar L, Reid J, Ye W, Tanasijevic B, Porras DP, et al.

Identification of Chemotherapy-Induced Leukemic-Regenerating Cells Reveals

a Transient Vulnerability of Human AML Recurrence. Cancer Cell. 2018;

26. Galen P Van, Hovestadt V, Ii MHW, Aster JC, Lane AA, Bernstein BE, et al.

Single-Cell RNA-Seq Reveals AML Hierarchies Relevant to Disease

Progression and Immunity Article Single-Cell RNA-Seq Reveals AML

Hierarchies Relevant to Disease Progression and Immunity. Cell. 2019;

27. Griessinger E, Anjos-Afonso F, Pizzitola I, Rouault-Pierre K, Vargaftig J,

Taussig D, et al. A niche-like culture system allowing the maintenance of

primary human acute myeloid leukemia-initiating cells: a new tool to decipher

their chemoresistance and self-renewal mechanisms. Stem Cells Transl Med

2014;3:520–9. Available from: http://www.ncbi.nlm.nih.gov/pubmed/24493855

28. Michaelson D, Abidi W, Guardavaccaro D, Zhou M, Ahearn I, Pagano M, et al.

Rac1 accumulates in the nucleus during the G2 phase of the cell cycle and

promotes cell division. J Cell Biol 2008;181:485–96. Available from:

/pmc/articles/PMC2364699/?report=abstract

29. Murga C, Zohar M, Teramoto H, Gutkind JS. Rac1 and RhoG promote cell

survival by the activation of PI3K and Akt, independently of their ability to

stimulate JNK and NF-kB. Oncogene. 2002;21:207–16. Available from:

www.nature.com/onc

30. Chen S, Li H, Li S, Yu J, Wang M, Xing H, et al. Rac1 GTPase Promotes

Interaction of Hematopoietic Stem/Progenitor Cell with Niche and Participates

in Leukemia Initiation and Maintenance in Mouse. Stem Cells 2016;34:1730–

41. Available from: http://doi.wiley.com/10.1002/stem.2348

31.   Sallmyr A, Fan J, Datta K, Kim KT, Grosu D, Shapiro P, et al. Internal tandem
861   duplication of FLT3 (FLT3/ITD) induces increased ROS production, DNA
862   damage, and misrepair: Implications for poor prognosis in AML. Blood
863   2008;111:3173–82. Available from: www.amaxa.com
864   32.   Jeanne M, Lallemand-Breitenbach V, Ferhi O, Koken M, Le Bras M, Duffort S,
865   et al. PML/RARA Oxidation and Arsenic Binding Initiate the Antileukemia
866   Response of As2O3. Cancer Cell. Cell Press; 2010;18:88–98.
867   33.   Lagadinou ED, Sach A, Callahan K, Rossi RM, Neering SJ, Minhajuddin M, et
868   al. BCL-2 inhibition targets oxidative phosphorylation and selectively
869   eradicates quiescent human leukemia stem cells. Cell Stem Cell. 2013;
870   34.   Pavlides SC, Huang K-T, Reid DA, Wu L, Blank S V, Mittal K, et al. Inhibitors
871   of SCF-Skp2/Cks1 E3 ligase block estrogen-induced growth stimulation and
872   degradation of nuclear p27kip1: therapeutic potential for endometrial cancer.
873   Endocrinology 2013;154:4030–45. Available from:
874   /pmc/articles/PMC3800755/?report=abstract
875   35.   Wunderlich M, Mizukawa B, Chou FS, Sexton C, Shrestha M, Saunthararajah
876   Y, et al. AML cells are differentially sensitive to chemotherapy treatment in a
877   human xenograft model. Blood. 2013;
878   36.   Grey W, Chauhan R, Piganeau M, Huerga Encabo H, Garcia-Albornoz M,
879   McDonald NQ, et al. Activation of the receptor tyrosine kinase, RET, improves
880   long-term hematopoietic stem cell outgrowth and potency. Blood. 2020;
881   37.   Luis TC, Ichii M, Brugman MH, Kincade P, Staal FJT. Wnt signaling strength
882   regulates normal hematopoiesis and its deregulation is involved in leukemia
883   development. Leukemia 2012;26:414–21. Available from:
884   http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3378318&tool=pmc
885   entrez&rendertype=abstract
886   38.   Ludin A, Gur-Cohen S, Golan K, Kaufmann KB, Itkin T, Medaglia C, et al.
887   Reactive oxygen species regulate hematopoietic stem cell self-renewal,
888   migration and development, as well as their bone marrow microenvironment.
889   Antioxidants Redox Signal. 2014.
890   39.   Ye H, Adane B, Khan N, Stranahan AW, Park CY, Jordan CT, et al. Leukemic
891   Stem Cells Evade Chemotherapy by Metabolic Adaptation to an Adipose
892   Tissue Niche Article Leukemic Stem Cells Evade Chemotherapy by Metabolic
893   Adaptation to an Adipose Tissue Niche. Cell Stem Cell. 2016;

894   40.   Kadia T, Kantarjian H, Garcia-Manero G, Borthakur G, Wang X, Patel K, et al.
895   Prognostic significance of the Medical Research Council cytogenetic
896   classification compared with the European LeukaemiaNet risk classification
897   system in acute myeloid leukaemia. Br. J. Haematol. 2015.

898   41.   Raffel S, Klimmeck D, Falcone M, Demir A, Pouya A, Zeisberger P, et al.
899   Quantitative proteomics reveals specific metabolic features of Acute Myeloid
900   Leukemia stem cells. Blood. 2020;

901   42.   Aasebø E, Brenner A, Berven F, Bruserud Ø, Selheim F, Hernandez-
902   Valladares M. Proteomic Profiling of Primary Human Acute Myeloid Leukemia
903   Cells Does Not Reflect Their Constitutive Release of Soluble Mediators.
904   Proteomes. 2018;7:1.

905   43.   Kukalev A, Ng Y-M, Ju L, Saidi A, Lane S, Mondragon A, et al. Deficiency of
906   Cks1 Leads to Learning and Long-Term Memory Defects and p27 Dependent
907   Formation of Neuronal Cofilin Aggregates. Cereb Cortex. 2017;27:11–23.
908   Available from: https://academic.oup.com/cercor/article/2573019/Deficiency

909   44.   Mulloy JC, Cancelas JA, Filippi MD, Kalfa TA, Guo F, Zheng Y. Rho GTPases
910   in hematopoiesis and hemopathies. Blood. 2010.

911   45.   Ford CD, Green W, Warenski S, Petersen FB. Effect of prior chemotherapy on
912   hematopoietic stem cell mobilization. Bone Marrow Transplant. 2004;

913   46.   Lin JJ, Milhollen MA, Smith PG, Narayanan U, Dutta A. NEDD8-targeting drug
914   MLN4924 elicits DNA rereplication by stabilizing Cdt1 in S phase, triggering
915   checkpoint activation, apoptosis, and senescence in cancer cells. Cancer Res.
916   2010;

917   47.   Burnett AK, Russell NH, Hills RK, Kell J, Cavenagh J, Kjeldsen L, et al. A
918   randomized comparison of daunorubicin 90 mg/m2 vs 60 mg/m2in AML
919   induction: Results from the UK NCRI AML17 trial in 1206 patients. Blood.
920   2015;

921   48.   Ding L, Ley TJ, Larson DE, Miller CA, Koboldt DC, Welch JS, et al. Clonal
922   evolution in relapsed acute myeloid leukaemia revealed by whole-genome
923   sequencing. Nature. 2012;

924   49.   Kitajima S, Kudo Y, Ogawa I, Bashir T, Kitagawa M, Miyauchi M, et al. Role of
925   Cks1 Overexpression in Oral Squamous Cell Carcinomas. Am J Pathol.
926   2004;165:2147–55.

50.   Hamdi A, Lesnard A, Suzanne P, Robert T, Miteva M a, Pellerano M, et al.

Tampering with cell division by using small-molecule inhibitors of CDK-CKS

protein interactions. Chembiochem 2015;16:432–9. Available from:

http://www.ncbi.nlm.nih.gov/pubmed/25619419

51. Lough L, Hao B, Sherman D, Ni E, Young LM. ubiquitylation and degradation.

Royal Society of Chemistry; 2018;1.

52. Pemovska T, Kontro M, Yadav B, Edgren H, Eldfors S, Szwajda A, et al.

Individualized systems medicine strategy to tailor treatments for patients with

chemorefractory acute myeloid leukemia. Cancer Discov; 2013;3:1416–29.

Available from: https://pubmed.ncbi.nlm.nih.gov/24056683/

53. Yadav B, Pemovska T, Szwajda A, Kulesskiy E, Kontro M, Karjalainen R, et al.

Quantitative scoring of differential drug sensitivity for individually optimized

anticancer therapies. Sci Rep 2014;4. Available from:

https://pubmed.ncbi.nlm.nih.gov/24898935/

54. Hu Y, Smyth GK. ELDA: Extreme limiting dilution analysis for comparing

depleted and enriched populations in stem cell and other assays. J Immunol

Methods 2009;347:70–8. Available from:

http://www.ncbi.nlm.nih.gov/pubmed/19567251

Figure Legends

Figure 1. *CKS1B* expression status can predict sensitivity to inhibition of

CKS1-dependent protein degradation in AML. A. Expression of *CKS1B* (relative

to *GAPDH*) in a poor risk AML cohort. FAB and p53 status are indicated for each

patient (FAB color coded, p53 status: white = WT; black = mutant; N=32). **B.**

Pearson's correlation of *CKS1B* expression versus DSS for a panel of clinically

available and investigational compounds. Red indicates significant correlations

above R^2=0.5. Blue indicates CKS1i (N=21). **C.** Diagram of action for CKS1i binding

and inhibition of the $SCF^{SKP2-CKS1}$ ubiquitin ligase complex **D.** Percentage of viability

of example patient AML samples cultured for 72 hours with indicated doses of

CKS1i. **E.** Allocation of patient AML in high or low *CKS1B* expression (50^{th}

percentile) and compared for CKS1i DSS (N=21). **F.** Percentage viability of AML cell

lines cultured for 48 hours with indicated doses of CKS1i (N=3 for all cell lines on

graph). **G.** Correlation between AML cell line CKS1i IC_{50} and *CKS1B* expression.

95% confidence intervals presented. Pearson's correlation coefficient was calculated

for correlation (R^2). **H.** Representative FACS plots for induction of apoptosis in the
indicated AML cell lines by presence of annexin V at the cell surface in response to
CKS1i (5μM) at 48 hours. Fold change *in vivo* leukemic burden of **I.** THP-1 (Ctrl N=6,
CKS1i N=7) and **J.** HL60 (Ctrl N=5, CKS1i N=6) cells day 21 (9 days post-CKS1i)
versus day 7 (pre-CKS1i) expressed as bioluminescent total flux intensity. **K.** Overall
survival of xenografts carrying THP-1 and HL60 cell lines control or treated with
CKS1i. A Student's *t*-test was used to calculate significance of difference for all
graphs unless otherwise stated. ** $p<0.005$; **** $p<0.0001$.

Figure 2. Inhibition of CKS1-dependent protein degradation selectively kills
bulk AML *in vivo* and depletes leukemic stem cell potential. A. *CKS1B*
expression (relative to *GAPDH*) for patient AMLs tested *in vivo* (N=5)*.* **B.** Illustration
of *in vivo* engraftment of patient AMLs indicating bone marrow aspiration time points
and treatment interval. Each arrow for CKS1i treatment refers to one day. **C.**
Percentage of human CD45$^+$ cells of total CD45$^+$ cells in mouse bone marrow
aspirations one week after chemotherapy (week 6). **D.** Kaplan-Meier survival curve
and p value calculated for each individual PDX Control and CKS1i treated. Each
data point represents one mouse. **E.** *CKS1B* expression (relative to *GAPDH*) for
patient AMLs tested under L-LTC-LIC conditions (N=7). **F.** Illustration of treatment
time points, treatment and readout for *ex vivo* L-LTC-IC. **G.** Fold change L-LTC-IC
frequency, CKS1i treatment versus control, after 7 weeks of co-culture. **H.** Individual
1/L-LTC-IC frequencies with upper and lower limits for each patient tested. A
Student's *t*-test was used to calculate significance of difference for all graphs unless
otherwise stated.
* $p<0.05$; **$p<0.01$; *** $p< 0.001$.

Figure 3. Divergent responses to CKS1i lead to RAC1-dependent induction of
lethal ROS in AML. A. Workflow for timescale of cell preparation for mass
spectrometry analysis. **B.** UNIPROT keywords for significantly enriched proteins in
THP-1 (purple) and CD34$^+$ (Green) cells in CKS1i treated conditions versus control.
C. Venn diagram depicting overlap of differentially expressed proteins between THP-
1 and CD34$^+$ cells. Volcano plots for changes in protein abundance (CKS1i vs
Control) in **D.** THP-1 and **E.** CD34$^+$ cells. Red dots indicate significantly upregulated

proteins, blue dots indicate significantly downregulated proteins (N=4 per cell line

and treatment). **F.** String network analysis (Red = increased abundance, Blue =

decreased abundance) and **G.** Protein abundance of key RAC1 pathway members

differentially abundant in THP-1 but not CD34$^+$ cells. **H.** Illustration of

RAC1/NADPH/ROS pathway, indicating mode of action for RAC1 inhibitor

NSC23766 (NSC). **I.** Fold change total NADP/NADPH and **J.** fold change NADPH in

THP-1 cells treated with the indicated doses of CKS1i (+ = 1μM, ++ = 5μM) or NSC

(+ = 0.1μM) versus control for 24 hours (N=4 per cell line and treatment). **K.**

Representative flow plots and **L.** quantified mean fluorescence intensity of

intracellular reactive oxygen species (ROS) in the indicated cell lines in response to

CKS1i (+ = 1μM) and NSC (+ = 0.1μM) treatment (N=3 per cell line and treatment).

A Student's *t*-test was used to calculate significance of difference for all graphs. *

p<0.05; **p<0.01; *** p< 0.001; **** p<0.0001.

1008

1009

Figure 4. CKS1i protects normal hematopoietic cells from chemotherapeutic

toxicity by suppressing key signalling pathways. A. Live cell count (N=6); **B.**

Quantified cell cycle proportions of CD34$^+$ cells grown for 48 hours in control

conditions or treated with CKS1i. **C.** Illustration of CD34$^+$ engraftment and

chemotherapeutic treatment in NSG mice. Each dot corresponds to one mouse.

Change in percentage human CD45$^+$ of total CD45 at the indicated time points for **D.**

Control, **E.** Doxorubicin/Cytarabine (DA) and **F.** Doxorubicin/Cytarabine plus CKS1i

(DAC) treatments. **G.** Fold change % human CD45 cells week 4 to 6 for the indicated

treatments. **H.** Representative flow plots and **I.** % total cells annexin V positive after

6 weeks *in vivo* for human CD45 cells with the indicated treatment conditions (Ctrl

N=5, DA N=3, DAC N=3). **J.** Representative flow plots and **K.** quantified mean

fluorescence intensity for non-phosphorylated β-catenin in CD34$^+$ cells grown for 48

hours in control conditions or treated with CKS1i (N=4). **L.** Representative flow plots

(including cells grown without OP-Puromycin; -OPP) and **M.** % total OP-Puromycin

incorporation in CD34$^+$ cells grown for 48 hours in control conditions or treated with

CKS1i. OP-Puromycin was added 1hr prior to collection and fixation of cells (N=4).

N. Representative flow plots and **O.** quantified mean fluorescence intensity of

intracellular reactive oxygen species (ROS) in CD34$^+$ cells grown for 48 hours in

control conditions or treated with CKS1i (N=4). **P.** ELDA quantification plot for CD34$^+$
cells grown in LTC-IC under control (grey) and CKS1i (blue) treated conditions.
Linear model and confidence interval are shown with estimated stem cell frequency
indicated. A Student's *t*-test was used to calculate significance of difference unless
otherwise stated. * $p<0.05$; ** $p<0.01$; *** $p< 0.001$; **** $p<0.0001$.

1033

Figure 5. Combination induction chemotherapy and CKS1i reduces AML
burden and leukemic stem cell potential whilst protecting resident
hematopoietic cells. A. *CKS1B* expression (relative to *GAPDH*) for patient AMLs
tested *in vivo* (N=4). **B.** Illustration of *in vivo* engraftment of patient AMLs indicating
bone marrow aspiration time points and treatment interval. Each arrow for DA and
CKS1i treatment refers to one day. **C.** Percentage of human CD45$^+$ cells of total
CD45$^+$ cells in mouse bone marrow aspirations one week after chemotherapy (week
6). **D.** Colony forming units per 10,000 mouse CD45$^+$ cells seeded from week 6 bone
marrow aspirations. **E-H.** Kaplan-Meier survival curve and p value calculated for
each individual PDX Control and treated mice. Each data point represents one
mouse. **I.** Total number of murine Long-term HSCs obtained from bone marrow of
mice at the final survival time point (Ctrl N=8, DA N=5, DAC N=5). **J.** Serial colony
forming units per 10,000 mouse CD45$^+$ cells obtained from BM of mice at the final
survival time point (Ctrl N=6, DA N=5, DAC N=6). **K.** *CKS1B* expression (relative to
GAPDH) for patient AMLs tested in L-LTC-IC (N=7). **L.** Fold change live human
CD45$^+$ cells, indicated treatments versus control, after two weeks of co-culture. **M.**
Fold change of L-LTC-IC frequency, indicated treatment versus control, after 7
weeks of co-culture. **N-O.** LSC frequency in secondary transplanted mice injected
with AML26 **N.** and AML32 **O.** at limiting dilution 6 weeks post-transplantation. **P.**
Kaplan-Meier survival curve for all AML32 secondary mice up to 120 days. A
Student's *t*-test was used to calculate significance of difference for all graphs unless
otherwise stated. * $p<0.05$; ** $p<0.01$; *** $p< 0.001$; **** $p<0.0001$.

Figure 1.

Figure 2.

Figure 3.

Figure 4

Figure 5.