<u>**Title page**</u>

Title:

Feeding intervention potentiates the effect of mechanical loading to induce
new bone formation in mice

Running head:

Feeding intervention potentiates new bone formation

Hasmik Jasmine Samvelyan[1,2,3], John Cummings Mathers[1,4], Timothy Michael
Skerry[1,2,*]

[1] MRC-Versus Arthritis Centre for Integrated Research into Musculoskeletal
Ageing

[2] Mellanby Centre for Bone Research, Department of Oncology and
Metabolism, The Medical School, The University of Sheffield, Beech Hill
Road, Sheffield, , UK

[3] Present address: Centre for Stress and Age-Related Disease, School of
Pharmacy and Biomolecular Studies, University of Brighton, Brighton, BN2
4GJ, UK. E-mail: h.samvelyan@brighton.ac.uk

[4] Human Nutrition Research Centre, Centre for Healthier Lives, Population
Health Sciences Institute, Newcastle University, Framlington Place,
Newcastle upon Tyne, NE2 4HH, UK. E-mail: john.mathers@newcastle.ac.uk

*Corresponding author; Timothy M Skerry (Department of Oncology and
Metabolism, The Medical School, The University of Sheffield, Beech Hill
Road, Sheffield, S10, 2RX, UK, E-mail: t.skerry@sheffield.ac.uk, Tel: +44 (0)
114 215 9026)

<u>**Author contributions**</u>

HJS conducted the research study. TMS conceived the original idea for the study,

HJS, JCM and TMS all contributed to the study design, analysis and interpretation of

the data, drafted, reviewed, edited and approved the version of the manuscript for

publication.

Abstract

The benefits of increased human lifespan depend upon duration of healthy, independent living; the healthspan. Bone-wasting disorders contribute significantly to loss of independence, frailty and morbidity in older people. Therefore, there is an unmet need globally for lifestyle interventions to reduce the likelihood of bone fractures with age. Although many mechanisms are involved in disorders of bone loss, there is no single regulatory pathway and, therefore, there is no single treatment available to prevent their occurrence. Our aim in these studies was to determine whether fasting/feeding interventions alter the effect of mechanical loading on bone anabolic activities and increase bone mass. In young 17-week-old mice, 16-hour fasting period followed by reintroduction of food for 2 hours increased markedly the potency of mechanical loading, that mimics the effect of exercise, to induce new cortical bone formation. Consistent with this finding, fasting and re-feeding increased the response of bone to a loading stimulus that, alone, does not stimulate new bone formation in ad-lib fed mice. Older mice (20-months) experienced no potentiation of loading-induced bone formation with the same timing of feeding interventions. Interestingly, the pre-, prandial and postprandial endocrine responses in older mice were different from those in young animals. The hormones that change in response to timing of feeding have osteogenic effects that interact with loading-mediated effects. Our findings indicate associations between timing of food ingestion and bone adaptation to loading. If translated to humans, such non-pharmacological lifestyle interventions may benefit skeletal health of humans throughout life-course and in older age.

Introduction

The dramatic increase in human life expectancy in the last 60 years has been driven

by reduced deaths of people in their 60s and 70s, leading to demographic shifts so

that most populations globally are characterised by increasing numbers of older old

people[1]. However, an increasing number of these extra years of life are spent in

poor health with more years spent in care facilities due to reduced ability to live

independently. Osteoporosis is a major contributor to loss of independence due to

bone fractures, and resulting hospital treatments lead to significant morbidity. Even

after successful fracture treatment, independent living is compromised in many

patients[2]. While drug treatments reduce consequences of osteoporosis significantly,

there is a pressing need for non-pharmacological interventions to improve bone

health across the life-course and to reduce likelihood of age-related bone disease.

Physical activity is beneficial for the skeleton acting through the process of functional

adaptation, a process that is less effective in older animals and humans than young

ones[3–6]. In addition, older people and those with bone loss, who are already at

increased risk of fractures, may be unable to perform exercise that is sufficiently

vigorous to strengthen their bones. Since dietary interventions such as energy

(caloric) restriction are associated with healthy ageing, a combination of exercise

and nutrition may benefit bone health in ageing by utilising synergies between

osteotropic influences.

Increased physical activity is beneficial for the skeleton by reducing the rate of age-

related loss of bone mass through shifting the balance of bone remodelling in favour

of bone formation[7]. However, during ageing skeletal resorption rates are increased

with decrease in bone formation rates, resulting in overall net bone loss[8]. Therefore,

it is hard for elderly adults and patients with bone wasting disorders to perform

vigorous exercise safely, and there is a need to identify effective ways for those

people to exercise which produce maximum benefits for the musculoskeletal system

without increased risk of injury.

There is complex and coordinated relationship between the endocrine regulation of

energy metabolism, adipose tissue and bone homeostasis[9–12]. Specifically,

concentrations of hormones including ghrelin, leptin, insulin, amylin, incretins, GIP,

GLP1 and GLP2 change profoundly in anticipation of, during or after eating

(preprandial, prandial and postprandial responses). These hormones have anabolic

effects on bone homeostasis, therefore, synergies between endocrine and

mechanical effects could alter with changes in concentrations of these hormones.

Indeed, previous studies in model organisms provide evidence of synergistic effects

of one of the osteoregulatory hormones; parathyroid hormone and mechanical

loading on bone. Parathyroid hormone administration alone induces bone formation

but, in combination with mechanical loading, enhances mechanically-induced bone

formation synergistically[13–17].

In this study, we have investigated the synergistic effect of feeding interventions

following an overnight fast on the response of bone to mechanical loading in mice

and the concurrent role of candidate osteotropic hormones in bone homeostasis.

Using a well-established tibial axial loading model[18,19], we performed experiments to

determine osteogenic effects of different mechanical loading regimens after periods

of withholding food (fasting) or feeding in young (17-19 week-old) and aged (20-

month-old) male C57BL/6 mice. We measured new bone formation in these mice

using micro-computed tomography and dynamic histomorphometry. To explore

possible mechanisms by which fasting/feeding interventions could have their effect, we measured changes in serum concentrations of a range of candidate osteotropic hormones in respone to feeding in fed and fasted fed young and aged mice using a multiplex assay.

<u>Results</u>

The effect of 16-hour food restriction on feeding behaviour of male mice

To establish a robust model to study the effect of timing of feeding on bone's response to loading, we first assessed the effect of overnight, 16-hour food removal on subsequent feeding behaviour of young (17-18 weeks old) and aged (20-month-old) male C57BL/6 mice. The results provided insights into dynamics of the homeostatic feeding response and showed that both young adult (Fig. 1a) and aged (Fig. 1b) male mice exhibited a similar 2-hour hyperphagic response following fasting before returning to the same food intake as mice that had unrestricted access to a normal chow ad libitum for the whole period of experiment (Fig. 1a, b).

Feeding intervention potentiates adaptive response of the bones to mechanical loading in young mice

Next we compared bone formation responses in ad-lib fed animals and in mice fasted overnight and then allowed free access to maintenance diet for 1, 2 or 3 hours before loading. To mimic osteogenic exercise, compressive forces were applied between the hock (ankle) and stifle (knee) joints to induce deformation of the tibiae. Different loading regimens comprised 40 cycles of compressive force at high physiological strain rate to peak strain magnitudes of 2200, 1300 or 1100 microstrain, followed by a brief hold of 0.25 seconds, and a rapid symmetrical reduction in loading, with a 9 second rest period before the next compression. The

three strain magnitudes corresponded to maximal, sub-maximal and sub-threshold

stimuli respectively. Loading was applied on Monday, Wednesday and Friday of 2

successive weeks, at 1, 2 or 3 hours after food had been re-introduced to 16-hour

fasted mice, or at the same time of day in ad-lib fed animals (Supplementary Fig.

S1).

Using a high resolution µCT analysis of the cortical bone at the mid-shaft region of

loaded tibiae, we found the expected responses to loading at high physiological

strains with significant adaptive response in the mice fed ad-lib. However, in mice

that were fasted then fed for 2 hours, the change in cortical bone thickness induced

by maximal loading was increased by 36% compared with responses of mice fed ad-

lib before the loading (0.049 ± 0.007mm in fasted/fed loaded tibiae compared with

that in ad-lib fed mice 0.036 ± 0.004mm, $p < 0.05$) (Fig. 2a, c, d). The

histomorphometric data were consistent with the µCT measurements (Fig. 2b, Table

1). Colour-coded analysis revealed anatomical variation in cortical bone thickness

between loaded and non-loaded tibiae of fasted fed and ad-lib fed mice (Fig. 2d).

Differences in the cortical bone thickness of loaded tibiae were particularly apparent

between fasted/fed and ad-lib fed mice and correlated with µCT analysis. There was

no significant induction of bone formation in non-loaded legs of either ad-lib or

fasted/fed mice over the period of the experiment.

In animals fed for 1 or 3 hours after the end of fasting, the mean cortical thickenss

increased compared with ad lib-fed controls, but the difference was not significant.

Trabecular bone parameters were improved by loading in both fasted and ad-lib fed

groups and at all 3 times after re-introduction of food, but were not significantly

different between feeding interventions.

Load magnitude-related adaptive response of the cortical bone is potentiated by feeding intervention in young mice

We then determined effects of fasting and 2-hour feeding regimen on bone response to loading which, in ad-lib fed mice, induced less than maximal bone formation. In mice, whose bones were loaded to induce peak strain magnitudes of 1300 microstrain, a sub-maximal stimulus, loading after fasting and feeding increased cortical thickness by 13% compared with ad-lib fed mice (fasted/fed: 0.27 ± 0.007mm loaded limb versus 0.24 ± 0.003mm control limb, $p < 0.05$; ad-lib fed: 0.26 ± 0.004mm versus 0.24±0.004mm, $p < 0.05$) (Fig. 3b). Finally, we investigated the effects of a sub-threshold stimulus (1100 microstrain), not capable of inducing new bone formation in ad-lib fed mice. Remarkably, that previously ineffective stimulus together with imposition of the fasting/re-feeding regimen resulted in an 8% increase in cortical thickness (fasted/fed: 0.26 ± 0.008mm loaded limb versus 0.24 ± 0.008mm control limb, $p < 0.05$; ad-lib fed: 0.26 ± 0.002mm versus 0.25 ± 0.004mm, $p > 0.05$) (Fig. 3a). We did not detect significant changes in trabecular bone parameters in these studies.

Adaptive response of bones to mechanical loading in aged mice is not potentated by the same feeding intervention that is effective in young mice

We then compared effects of fasting/feeding regimens on load-induced bone formation in older 20-month-old mice, but observed no potentiation effect of loading of the fasting and 2 hour feeding regimen that we had seen in younger mice (cortical bone thickness fasted/fed: 0.23 ± 0.002mm loaded limb versus 0.21 ± 0.004mm control limb, $p < 0.05$; ad-lib fed: 0.24 ± 0.01mm versus 0.21 ± 0.007mm, $p < 0.05$) (Fig. 4a-c). Colour-coded analysis revealed no anatomical variation in cortical bone

thickness between loaded and non-loaded tibiae of fasted fed and ad-lib fed aged

mice (Fig. 4c).

Despite deterioration of the trabecular bone in these aged mice, trabecular bone

pattern factor decreased significantly in both fed and fasted 2-hour fed mice

(fasted/fed: -2.3 ± 4.3mm^{-1} loaded limb versus 20.8 ± 3.3mm^{-1} control limb, $p <$

0.001; ad-lib fed: 0.7 ± 3.7mm^{-1} versus 17.3±1.5mm^{-1}, $p < 0.01$) (Fig. 4d). Similarly,

there was no significant change of bone formation in non-loaded legs of either ad-lib

or fasted/fed aged mice over the period of the experiment.

Regulation of bone homeostasis by endocrine hormones differ between young

and aged mice

Next we assessed changes in circulating concentrations of gastro-entero-pancreatic

hormones in response to fasting and feeding, which may contribute to observed

potentiating effects. We measured concentrations of a range of hormones with

known effects on bone homeostasis in young and old mice, in both ad-lib fed and

fasted/fed groups. Since both young and aged mice previously exhibited a similar 2-

hour hyperphagic response following fasting before returning to the same food

intake, serum concentrations of ghrelin, leptin, pancreatic hormone insulin, and

intestinal peptides GIP and GLP1 were measured at 3 time points (overnight 16-hour

fasted at 12:00 noon on the day following the fast, then 2 hours and 3 hours after

initiating food intake at 14:00 and 15:00 respectively) using a multiplex metabolic

assay. Endocrine profiles differed between young and aged mice. Fasting and

feeding changed circulating serum concentrations of ghrelin, leptin, insulin and GLP-

1, but not GIP in young mice (Table 2). In older mice, absolute concentrations of

hormones were different, but differences between ad-lib fed and fasted fed mice

were not significant (Table 3).

Ghrelin concentrations fell significantly 2 hours after initiating food intake in young

mice, but remained high in older animals (Fig. 5a, b). In fasted fed young and aged

mice, serum leptin concentrations increased significantly at 14:00 and 15:00

compared with those at 12:00, whereas in ad-lib fed mice remained unchanged.

Fasting serum concentration of leptin was significantly lower at 12:00 and

significantly higher at 15:00 in fasted fed young mice compared to those in ad-lib fed

mice (Fig. 5c and d). In fasted fed young mice serum insulin concentration

significantly increased from 12:00 to 14:00 and in comparison to that in ad-lib fed

mice, then significantly decreased at 15:00 (Fig. 5e). In fasted fed aged mice, serum

insulin concentration increased but not significantly from 12:00 to 14:00, then

significantly decreased at 15:00 compared to that at 14:00, whereas serum insulin

concentrations remained unchanged in ad-lib fed aged mice (Fig. 5f). In fasted fed

young mice, serum GIP and GLP1 concentrations significantly increased at 14:00

and 15:00 compared with those at 12:00 and remained unchanged in ad-lib fed mice

(Fig. 5g, i). Serum GLP1 concentration was significantly higher at 15:00 in fasted fed

mice compared to that in ad-lib fed mice (Fig. 5i). There were no significant changes

in serum GIP and GLP1 concentrations within and between fed and fasted fed aged

mice (Fig. 5h, j).

Discussion

These data provide compelling evidence for a link between variations in timing of

feeding and adaptive responses of bone to loading. Age-related bone loss across

multiple sites within the skeleton suggests that systemic regulation plays an essential

role in bone homeostasis. Concentrations of a number of gastro-entero-pancreatic

hormones change before, during and after feeding, which have direct or indirect

effects on bone cells. We interpret the data from this study to provide proof-of-

principle that the effects of anabolic hormones in increasing bone formation in

response to mechanical loading may be manipulated by changing timing of feeding

in relation to exercise imposition. It has been previously shown that parathyroid

hormone (an osteotropic hormone) administration potentiates effects of loading[14–16],

but this study is the first report of a non-pharmacological intervention with such

effects. The ability of feeding interventions to amplify the effects of loading stimuli is

compelling, but its ability to turn a sub-threshold stimulus (one which alone does not

stimulate any adaptive bone responses) into one causes significant new bone

formation has potentially pervasive implications.

Our study of the fasting/2-hour feeding before mechanical loading in older mice did

not recapitulate results seen in younger animals. Measurement of hormones in our

studies gives clues to possible explanations for differences in adaptive responses of

bones between young and aged mice as in the data on 5 different hormones, we

determined clear differences in changes over the fasting/feeding/loading period in

the two age groups. Age-related impairment of the function of mechanostat and

decline of the skeletal robustness may have also reduced sensitivity of the cortical

bone to mechanical stimulation in aged male mice. Further, the altered cortical bone

response to the strain-related stimulus in aged male mice compared to young ones

may reflect the deficiencies in osteoblast recruitment, differentiation and function.

However, one limitation of this study is that we did not examine different timings of

fasting and feeding in old mice as we had in young mice.

We were conscious that repeated blood sampling from mice could be stressful, and

that such stress might affect levels of the hormones we wished to measure.

However, assessments of serum cortisol concentrations during those blood sampling

studies showed no significant changes suggesting minimal impact of the procedure

on stresses experienced by the mice (Supplementary Fig. S2).

Currently, although the effects we demonstrate are potent and robust, we have only

association between the changes in bone responses and the hormonal alterations

before, during and after feeding. There are however, compelling data on the way

that such hormonal changes occur, and it is appropriate to consider some of the

possible candidates in this connection.

Leptin is well recognised as a bone anabolic hormone, with its own effects mediated

by central and peripheral mechanisms as well as those through other central

relays[20]. We found like others before, that leptin concentrations are lower in fasted

mice and that they increase following feeding[21–23]. It is known that leptin increases

bone formation[24,25], although to our knowledge, there are no reports of concurrent

administration of leptin and loading of bones.

Ghrelin is thought to be antagonistic to leptin, has direct effects on bone

homeostasis both *in vivo* and *in vitro*[26,27] and changes during fasting and in advance

of feeding[28,29]. The high circulating ghrelin concentrations during fasting and then low

concentrations after food intake would be consistent with an influence of the

leptin/ghrelin axis on the loading related bone formation that we demonstrate.

Insulin increases bone formation and decreases bone resorption both *in vivo* and *in*

vitro[30–32]. During fasting circulating insulin concentrations are low (basal condition)

and as eating commences, plasma insulin concentrations rise to regulate blood

glucose concentrations. These elevated levels would be consistent with a

potentiated anabolic bone response to loading. Specifically, we demonstrate that 2-

hours of feeding after fasting increased serum insulin concentrations in young mice.

The lack of similar change in insulin in the older mice is consistent with our

demonstration that the same potentiation did not occur in them. As insulin has

promiscuous effects on IGF receptors, which are expressed on osteoblasts[33], it is

likely that any insulin-mediated effect is complex.

GIP is a bone anabolic *in vitro*[34,35] and mice overexpressing GIP have increased

bone mass[36]. Similarly, GLP1 receptors are expressed on osteoblasts, bone marrow

stromal cells[37] and on thyroid C cells[38], so GLP1 has the potential for direct and

indirect actions on bone. However, as the concentrations of both hormones were

unchanged in aged mice by our interventions we conclude that impairment of GIP or

GLP1 signalling in bone cells are likely to be involved in attenuated bone response

we observed. The roles of these different hormones could be clarified by

interventional studies where hormones were administered in order to mimic the

fasting/feeding changes, or in knockout or over-expressing mice, but those studies

are beyond the scope of this research and any single report.

While fasting mice for 16 hours is a much more significant period of food deprivation

than overnight fasting in humans, it is possible that similar biological interactions

occur in humans, that could be exploited to benefit bone health. Future studies may

allow us to determine whether such potentiating effects are observed in humans and

what recommendations can be made to humans with respect to timing, amount and

composition of food ingestion to obtain the maximal benefits from exercise. If we can

identify interventions that potentiate bone's response to exercise in young humans,

then such interventions would increase adult bone mass so that with ageing, the

additional strength of bones would extend time before osteoporotic "fracture

thresholds" were reached. If similar interventions were identified in older mice and

then humans too, then they could be translatable into a non-pharmacological way to

improve bone health of older people. There are already widespread fasting and

caloric restriction interventions in use in humans for weight control and for healthy

ageing[39]. In particular, the 5:2 diet in which food intake is restricted to 500 – 600

calories on two days per week, which could be exploited to provide the equivalent of

fast experienced by mice in this study. The likelihood that effects we show here

could be translatable to humans has been increased by a recent study[40] that showed

in postmenopausal women eating only twice daily, positive effects of treadmill

exercise on serum biomarkers of bone formation, one hour after eating but not 1

hour before eating.

Materials and methods

Animals

Male C57BL/6 wild type mice at 16 weeks of age (young adult) and 20 months of

age (aged) were obtained from Charles River Laboratories Inc. (Margate, UK). The

mice were caged separately and acclimatised to their surroundings for 7 days. All

mice were allowed free access to water and a maintenance diet ad libitum (2018C

Teklad Global 18% Protein Rodent Diet; Madison, WI, USA, Supplementary Table

S1) in a 12-hour light/dark cycle at a room temperature of 21 ± 2^0C and relative

humidity of $55 \pm 10\%$. Cages contained wood shavings, bedding and metal rings for

mice to play. All procedures complied with the United Kingdom Animals (Scientific

Procedures) Act 1986 (ASPA) and were reviewed and approved by the University of

Sheffield Research Ethics Committee (Sheffield, UK). 3Rs principle followed as the

ethical framework for conducting all the scientific experiments.

Assessment of feeding behaviour of fed or fasted young and aged mice

Groups of young adult male mice at 17 weeks of age and aged male mice at 20

months of age were either allowed access to a maintenance diet ad libitum or fasted

overnight for 16 hours and were assessed for subsequent feeding behaviour. The

nesting material, bedding and wood shavings were removed from the cages prior to

the experiment. All mice were allowed free access to water. Food intake was

measured by monitoring the consumed food and body weights of animals every 30

minutes for the first 3 hours then every hour for another 4 hours twice weekly for one

week. After the experiments mice were placed in new cages with wood shavings and

bedding.

In vivo non-invasive axial mechanical loading of knee joints

The right tibia of each mouse was subjected to non-invasive, dynamic axial

mechanical loading under the isofluorane-induced anaesthesia (liquid isofluorane

was vaporised to a concentration of 5% and maintained at a concentration of 3%

with oxygen) for 7min/day, 3 alternate days a week for 2 weeks according to the

protocols described in the previous studies[16,18]. The left tibiae were non-loaded

internal controls. To apply mechanical loading, mice were anaesthetised, and the

knee and ankle joints of the right limbs were fixed in customised concave cups. The

knee was positioned into the upper cup, which was attached to the activator arm

displacement transducer and the ankle in the lower cup attached to the dynamic load

cell. The tibia was held in place by continuous static preload of 0.5N onto which

dynamic loads were superimposed in a series of 40 trapezoidal shaped waveform

cycles with steep up and down ramps with a high dwell time of 0.25s and 9s low

dwell 'rest interval' between each cycle. The load was applied to engender the

required magnitude of strain on the bones of animals in each of the age groups[41].

The engendered average strain rate was 30000με/s, which is shown to be equivalent

to physiological strain rates[42].

Calcein administration

150mg of calcein powder was added to 50ml dH$_2$O. Then 0.1g of NaHCO$_3$ was

added to neutralise the pH of the solution. The solution was protected from light with

aluminium foil. After the solution was fully dissolved, it was filter-sterilized at 0.22μm

into autoclaved vials also protected from light with aluminium foil. The pH of the

solution was measured with a bench pH-meter and corrected to neutral pH 7 with

HCl or NaOH. The calcein solutions were made on the same days prior to the

experiments ensuring no loss of fluorescence from the fluorochrome label. Calcein

was administered at a concentration of 30mg/kg in 0.2% NaHCO$_3$ solution

subcutaneously on the first and last days of mechanical loading (days 1 and 12)

using insulin syringes[43].

Micro-computed tomography (μCT) analysis

The tibiae were collected after mice were culled, and loading responses of bones

determined by μCT using a SkyScan 1172 (Bruker MicroCT, Kontich, Belgium).

Since mouse bones are small, and axial loading related osteogenesis is site-specific,

high-resolution μCT analysis was used primarily to quantify three-dimensional (3D)

bone microarchitecture at comparable sites (proximal and mid-shaft regions) of

loaded and contralateral non-loaded control tibiae. The tibiae were stored in 70%

EOH and mounted in a plastic tube wrapped in cling film to prevent drying during

scanning. The high-resolution scans were imaged with a pixel size of 4.3μm. The

applied X-ray voltage was 50kV, X-ray intensity 200μA with a 0.5mm aluminium

filtration. The scans were taken over 180 degrees with a 0.7-degree rotation step.

The images were reconstructed and binarised with a threshold of 0 to 0.16, ring

artefact reduction was set at 10 and beam hardening correction at 0% using the

SkyScan NRecon software package (version 1.6.9.4, Bruker MicroCT, Kontich,

Belgium). The images then were realigned vertically using DataViewer software

(version 1.5.1.2 64-bit) before the 3D quantification.

3-dimensional (3D) and colour coded analysis

Structural parameters were calculated using SkyScan Ct.An software (Bruker

MicroCT, Kontich, Belgium) for cortical bone 1-mm-long section at the mid-shaft of

the tibia with an offset of 0.5mm, and for trabecular bone - secondary spongiosa,

1mm distal to the tibial proximal growth plate with an offset of 0.2mm ensuring the

growth plate was not included in the analysis. In the trabecular region, an irregular,

anatomic region of interest (ROI) adjacent to the endocortical boundary was

analysed. Two ROIs were drawn to analyse the cortical region including 'doughnut'

shaped ROI and 'crude' ROI around the outside of the bone.

The bone mass, architecture and changes due to the mechanical loading were

evaluated in cortical and trabecular bone regions according to the guidelines for the

assessment of bone microstructure in rodents using μCT[44]. 3D representations of

the bones were created using the Voxler software (Golden Software Inc., 2006). The

local bone thickness (Th; voxel) was determined and colour-coded thickness images

were generated using the Avizo® (version 8.0, VSG, Burlington, VT, USA) software.

Bone dynamic histomorphometry

After μCT scanning and being fixed in 70% EOH for a minimum of 48hrs, the mouse

bones were infiltrated in increasing concentrations of alcohol (80% IMS alcohol, 80%

IMS alcohol, 100% IMS alcohol, 100% IMS alcohol, 100% EOH, 100% EOH and

100% EOH) on a shaker at room temperature for 48hrs at each concentration.

Bones were then infiltrated in glass bottles with medium grade LR White resin at 4^0C

for 6 days, resin was changed every 48hrs. The sample bottles were agitated gently

and continuously on a shaker for a duration of the infiltration at 4^0C. Next, the bones

were placed longitudinally and diagonally in plastic moulds fully filled with fresh resin

and covered with labelled plastic embedding stubs at room temperature. Resin was

then allowed to polymerise at 55^0C overnight for up to 24hrs until it was hard. After

polymerisation, blocks were removed from the moulds and protected from the light

during the storage and subsequent use. The bones were protected from the light at

all stages of the processing to prevent the loss of fluorescence from the labels.

Sectioning and visualisation

The bones were reoriented vertically, then 8μm thick transverse sections were

obtained from the cortical ROI corresponding to the mid-shaft of the tibiae and as

determined by μCT, using the tungsten carbide-tipped 16cm microtome knife on the

rotary microtome. Sections were unrolled and floated onto a drop of the dH$_2$O on a

labelled Superfrost Plus glass slides using fine paintbrush and forceps. The slides

were covered with Saran wrap and wrapped with blotting paper making sure the

Saran wrap-covered sections were flat without creases or residual water. The slides

then were dried at 50^0C overnight stacked in the slide clump to aid adherence of

bones sections. After peeling off the Saran wraps slides were mounted in the xylene

for 2-3min and then covered with glass coverslip using DPX mountant. Images of

calcien labelled transverse bone sections were visualised using Leica fluorescent

microscope at 4x and 10x magnifications. All slides were stored protected from the

light until time for histomorphometry analysis.

Measurements of bone formation parameters

The histological assessment of bone phenotypes was carried out by dynamic

histomorphometry measuring its derived kinetic indices[45]. A Leica microscope was

set up for fluorescence before placing the slides with calcein labeled sections under

the microscope. To reach its full output, the fluorescent lump was allowed to warm

up for 5min before measurements were made. Bone formation parameters including

mineralizing surface (MS), mineral apposition rate (MAR) and bone formation rate

(BFR) were measured using the OsteoMeasure software (version 3.3.0.2).

Blood collection

Small blood volumes were obtained from the mice by tail vein sampling technique

with assistance from the Biological Services Unit at the Royal Hallamshire Hospital

of The University of Sheffield. To avoid hyperthermia and dehydration and reduce

stress levels, this technique was performed without warming mice in the warming

cabinet prior to taking the blood samples. Mice were gently placed in the restraint

tube and lateral tail veins were accessed by making incisions in approximately one-

third from the proximal end along the length of the tails[46]. Repeated blood micro

samples (typically ≤50μl) were collected from the lateral mouse-tail veins 3 times in

any one 24-hour period.

Mouse multiplex metabolic hormone assay

Serum ghrelin, leptin, insulin, GIP and GLP1 concentrations were assayed in mouse

sera using mouse multiplex assay (Bio-Rad, Watford, UK) following manufacturer's

instructions. The mouse assay is magnetic bead-based multiplex assay that allows

measurement of multiple proteins in small volumes of sera. Briefly, a 96-well plate

was pre-coated with 50μl of beads coupled with the capture antibodies. After

washing twice with wash buffer, 50μl of standards, blank and serum samples were

added into the wells, incubated at room temperature with shaking for 1hr, washed

and followed by addition of the detection antibodies and streptavidin-phycoerythrin

(SA-PE) conjugates. Following incubation at room temperature with shaking, the

plate was washed three times with wash buffer and beads were resuspended in

125μl assay buffer with shaking at 850rpm for 30s. The plate was then read on Bio-

Plex system using Bio-Plex Manager software (v 6.0). Samples (pg/ml) were

assessed in duplicate.

Serum cortisol analysis

Cortisol concentrations were measured in mouse sera by a solid phase competitive

enzyme linked immunosorbent assay (ELISA) (Sigma-Aldrich Co Ltd, USA)

according to the manufacturer's protocol. Prior to the assay the frozen sera were

thawed at a room temperature and mixed well using vortexer. Briefly, 25μl of mouse

serum was added into each microwell of the plate, coated with the anti-cortisol MAb,

along with the standards for 6-point standard curve (1-80ng/ml), blank and internal

control. After incubation at room temperature with shaking for 60min, wells were

washed three times with wash buffer and treated with 3,3',5,5' -

Tetramethylbenzidine (TMB) substrate for 15min. The colorimetric reaction was then

stopped and absorbance was measured at 450nm on ELISA reader. A 4-parametre

logistic regression was applied to the standard curve for the calculation of cortisol

concentrations in the samples (ng/ml). Samples were assessed in duplicate.

Statistical analysis

All analyses were performed using GraphPad Prism software 6.0f version

(GraphPad Inc, La Jolla, CA, USA). The results were presented as the mean ±

standard error (SEM). The Normal distribution of data was assessed using the

Shapiro-Wilk normality test. Statistical significance was tested using parametric

statistical tests. For comparing two groups, two-tail Student's t-test (paired or

unpaired) was used. For comparing more than two groups, two-way ANOVA

(analysis of variance) was used with Tukey post-hoc test to assess the effect of

loading, fasting/feeding and interactions within and across the experimental groups.

The significance was set at $p < 0.05$.

Data availability

The data generated and analysed during the current study are available from the corresponding author upon reasonable request.

Acknowledgements

This study was supported by CIMA (The MRC – Versus Arthritis Centre for Integrated Research into Musculoskeletal Ageing) (to TMS and JCM; MR/K006312/1) performed as a collaborative project between the Universities of Sheffield, Newcastle and Liverpool. The authors thank the members of the SkeletAL laboratory, The University of Sheffield, for tissue processing, Dr Carmen Martin-Ruiz (NIHR Newcastle Biomedical Research Centre, Ageing Research laboratories) for the analysis of serum cortisol concentrations in young adult and aged mice.

Conflict of interests

The authors declare no conflict of interests.

Author contributions

HJS conducted the research study. HJS, JCM and TMS contributed to the study design, analysis and interpretation of the data, drafted, reviewed, edited and approved the version of the manuscript for publication.

References

1. He, W., Goodkind, D. & Kowal, P. An Aging World⬜: 2015 International Population Reports. *Aging (Albany. NY).* 165 (2016) doi:P95/09-1.

2. Schemitsch, E. H. *et al.* Loss of Independence After Operative Management of Femoral Neck Fractures. *J. Orthop. Trauma* **33**, 292–300 (2019).

3. Wolff, J. Ueber die innere Architectur der Knochen und ihre Bedeutung für die

Frage vom Knochenwachsthum. *Arch. Für Pathol. Anat. Physiol. Für Klin. Med*

50, 389–450 (1870).

4.   Frost, H. M. On the strength-safety factor (SSF) for load-bearing skeletal

organs. *J. Musculoskelet. Neuronal Interact.* **3**, 136–140 (2003).

5.   Bonnet, Nicolas and Ferrari, S. L. Exercise and the Skeleton: How It Works

and What It Really Does. *IBMS Bonekey* **7**, 235–248 (2010).

6.   Konda, N. N. *et al.* A comparison of exercise interventions from bed rest

studies for the prevention of musculoskeletal loss. *npj Microgravity* **5**, (2019).

7.   Gonzalo-Encabo, P., McNeil, J., Boyne, D. J., Courneya, K. S. & Friedenreich,

C. M. Dose-response effects of exercise on bone mineral density and content

in post-menopausal women. *Scand. J. Med. Sci. Sport.* **29**, 1121–1129 (2019).

8.   Raisz, L. G. & Seeman, E. Causes of age-related bone loss and bone fragility:

An alternative view. *J. Bone Miner. Res.* **16**, 1948–1952 (2001).

9.   Perrini, S. *et al.* The GH/IGF1 axis and signaling pathways in the muscle and

bone: Mechanisms underlying age-related skeletal muscle wasting and

osteoporosis. *J. Endocrinol.* **205**, 201–210 (2010).

10.   D'Amour, P. Acute and chronic regulation of circulating PTH: Significance in

health and in disease. *Clin. Biochem.* **45**, 964–969 (2012).

11.   Govoni, K. E. Insulin-like growth factor-I molecular pathways in osteoblasts:

potential targets for pharmacological manipulation. *Curr. Mol. Pharmacol.* **5(2)**,

143–152 (2012).

12.   De Paula, F. J. A. & Rosen, C. J. Bone Remodeling and Energy Metabolism:

New Perspectives. *Bone Res.* **1**, 72–84 (2013).

13.   Ma, Y. *et al.* Parathyroid hormone and mechanical usage have a synergistic

effect in rat tibial diaphyseal cortical bone. *J. Bone Miner. Res.* **14**, 439–448

(1999).

14. Kim, C. H. *et al.* Trabecular Bone Response to Mechanical and Parathyroid

Hormone Stimulation: The Role of Mechanical Microenvironment. *J. Bone*

Miner. Res. **18**, 2116–2125 (2003).

15. Li, J., Duncan, R. L., Burr, D. B., Gattone, V. H. & Turner, C. H. Parathyroid

hormone enhances mechanically induced bone formation, possibly involving L-

type voltage-sensitive calcium channels. *Endocrinology* **144**, 1226–1233

(2003).

16. Sugiyama, T. *et al.* Mechanical loading enhances the anabolic effects of

intermittent parathyroid hormone (1-34) on trabecular and cortical bone in

mice. *Bone* **43**, 238–248 (2008).

17. Shibamoto, A. *et al.* Effect of high-frequency loading and parathyroid hormone

administration on peri-implant bone healing and osseointegration. *Int. J. Oral*

Sci. **10**, 1–7 (2018).

18. De Souza, R. L. *et al.* Non-invasive axial loading of mouse tibiae increases

cortical bone formation and modifies trabecular organization: A new model to

study cortical and cancellous compartments in a single loaded element. *Bone*

37, 810–818 (2005).

19. Sugiyama, T. *et al.* Bones' adaptive response to mechanical loading is

essentially linear between the low strains associated with disuse and the high

strains associated with the lamellar/woven bone transition. *J. Bone Miner. Res.*

27, 1784–1793 (2012).

20. Upadhyay, J., Farr, O. M. & Mantzoros, C. S. The role of leptin in regulating

bone metabolism. *Metabolism.* **64**, 105–113 (2015).

21. Ahima, Rexford S., Prabakaran, Daniel, Mantzoros, Christos, Qu, Daqing,

Lowell, Bradford, Maratos-Flier, Eleftheria, Flier, J. S. Role of leptin in the

neuroendocrine response to fasting. *Nature* **382**, 250–252 (1996).

22. Ahrén, B., Månsson, S., Gingerich, R. L. & Havel, P. J. Regulation of plasma

leptin in mice: Influence of age, high-fat diet, and fasting. *Am. J. Physiol. -*

Regul. Integr. Comp. Physiol. **273**, (1997).

23. Trayhurn, P., Hoggard, N., Mercer, J. G. & Rayner, D. V. Hormonal and

neuroendocrine regulation of energy balance - The role of leptin. *Arch. Anim.*

Nutr. **51**, 177–185 (1998).

24. Thomas, T. *et al.* Leptin acts on human marrow stromal cells to enhance

differentiation to osteoblasts and to inhibit differentiation to adipocytes.

Endocrinology **140**, 1630–1638 (1999).

25. Hamrick, M. W. & Ferrari, S. L. Leptin and the sympathetic connection of fat to

bone. *Osteoporos. Int.* **19**, 905–912 (2008).

26. Fukushima, N. *et al.* Ghrelin directly regulates bone formation. *J. Bone Miner.*

Res. **20**, 790–798 (2005).

27. Van Der Velde, M. *et al.* An age-dependent interaction with leptin unmasks

Ghrelin's bone-protective effects. *Endocrinology* **153**, 3593–3602 (2012).

28. Asakawa, A. *et al.* Ghrelin is an appetite-stimulatory signal from stomach with

structural resemblance to motilin. *Gastroenterology* **120**, 337–345 (2001).

29. Kojima, M. & Kangawa, K. Ghrelin: Structure and function. *Physiol. Rev.* **85**,

495–522 (2005).

30. Thomas, D. M. *et al.* Insulin receptor expression in primary and cultured

osteoclast-like cells. *Bone* **23**, 181–186 (1998).

31. Cornish, J., Callon, K. E. & Reid, I. R. Insulin increases histomorphometric

indices of bone formation in vivo. *Calcif. Tissue Int.* **59**, 492–495 (1996).

32.   Pun, K. K., Lau, P. & Ho, P. W. M. The characterization, regulation, and function of insulin receptors on osteoblast-like clonal osteosarcoma cell line. *J. Bone Miner. Res.* **4**, 853–862 (1989).

33.   Fulzele, K. *et al.* Disruption of the insulin-like growth factor type 1 receptor in osteoblasts enhances insulin signaling and action. *J. Biol. Chem.* **282**, 25649–25658 (2007).

34.   Bollag, R. J. *et al.* Osteoblast-derived cells express functional glucose-dependent insulinotropic peptide receptors. *Endocrinology* **141**, 1228–1235 (2000).

35.   Zhong, Q. *et al.* Effects of glucose-dependent insulinotropic peptide on osteoclast function. *Am. J. Physiol. - Endocrinol. Metab.* **292**, 543–548 (2007).

36.   Xie, D. *et al.* Glucose-dependent insulinotropic peptide-overexpressing transgenic mice have increased bone mass. *Bone* **40**, 1352–1360 (2007).

37.   Sanz, C. *et al.* Signaling and biological effects of glucagon-like peptide 1 on the differentiation of mesenchymal stem cells from human bone marrow. *Am. J. Physiol. - Endocrinol. Metab.* **298**, 634–643 (2010).

38.   Yamada, C. *et al.* The murine glucagon-like peptide-1 receptor is essential for control of bone resorption. *Endocrinology* **149**, 574–579 (2008).

39.   Redman, L. M. *et al.* Metabolic Slowing and Reduced Oxidative Damage with Sustained Caloric Restriction Support the Rate of Living and Oxidative Damage Theories of Aging. *Cell Metab.* **27**, 805-815.e4 (2018).

40.   Borer, K. T., Zheng, Q., Jafari, A., Javadi, S. & Kernozek, T. Nutrient intake prior to exercise is necessary for increased osteogenic marker response in diabetic postmenopausal women. *Nutrients* **11**, (2019).

41.   Meakin, L. B., Galea, G. L., Sugiyama, T., Lanyon, L. E. & Price, J. S. Age-

related impairment of bones' adaptive response to loading in mice is

associated with sex-related deficiencies in osteoblasts but no change in

osteocytes. *J. Bone Miner. Res.* **29**, 1859–1871 (2014).

42.   Fritton, S. P. & Rubin, C. T. In vivo measurement of bone deformations using

strain gauges. *Bone Mech. Handbook, Second Ed.* 8-1-8–42 (2001)

doi:10.1201/b14263-11.

43.   Hillam, R. A. & Skerry, T. M. Inhibition of bone resorption and stimulation of

formation by mechanical loading of the modeling rat ulna in vivo. *J. Bone*

Miner. Res. **10**, 683–689 (1995).

44.   Bouxsein, M. L. *et al.* Guidelines for assessment of bone microstructure in

rodents using micro-computed tomography. *J. Bone Miner. Res.* **25**, 1468–

1486 (2010).

45.   Dempster, D. W. *et al.* Standardized nomenclature, symbols, and units for

bone histomorphometry: A 2012 update of the report of the ASBMR

Histomorphometry Nomenclature Committee. *J. Bone Miner. Res.* **28**, 2–17

(2013).

46.   Sadler, A. M. & Bailey, S. J. Validation of a refined technique for taking

repeated blood samples from juvenile and adult mice. *Lab. Anim.* **47**, 316–319

(2013).

Figure legends

Fig. 1 Feeding behaviour of young and aged male C57BL/6 mice

a In total 14 mice were randomly divided into two groups; control ad-lib fed ($n = 7$) or

overnight 16-hour fasted ($n = 7$) at 17 weeks of age. **b** In total 16 male mice were

randomly divided into two groups; ad-lib fed ($n = 8$) or 16-hour fasted ($n = 8$) at 20

months of age. The body weights of mice were measured for 7 hrs following

overnight fast to monitor food intake of the animals. Following overnight food

restriction, fasted mice consumed more food compared with the mice in the control

group that had free access to food ad libitum. The body weights of both young adult

and aged mice in the fasted group increased significantly for 2 hrs compared with

their pre-fasted body weights indicating rapid ingestion of food (and water) in the

early re-feeding period. The proportions of the starting pre-fasted body weights (%) ±

SEM are shown. * indicates significant differences between fed and fasted fed

groups; * $p < 0.05$, ** $p < 0.01$, **** $p < 0.0001$ by two-way ANOVA with Tukey post-

hoc test

Fig. 2 Cortical bone phenotype of control (non-loaded) and loaded tibiae with

2200 microstrain (maximal) loading regimen in 19-week-old fed or overnight

fasted 2-hour fed male C57BL/6 mice

a Three-dimensional images of a 1.0-mm-thick cortical bone region 0.9mm below the

growth plate from the mid-shaft of control non-loaded (left) and loaded (right) tibiae

of fed or fasted 2-hour fed mice. **b** Transverse cortical bone confocal microscope

images from the mid-shaft of left non-loaded and right loaded tibiae of fed (top) or

fed for 2 hours following overnight fast (bottom) male mice. Double calcein labels

can be visualised on endocortical and periosteal surfaces of the loaded tibiae

confirming new bone formation at these sites. **c** The change in cortical bone

thickness (Ct.Th) of tibiae loaded at 13N to induce peak strain magnitude of 2200

microstrain compared with non-loaded controls of fed or fasted 2-hour fed young

adult male mice measured using μCT at tibial mid-shaft. Fasted 2-hour fed mice had

significantly greater response in Ct.Th of the right tibiae compared with ad-libitum fed

mice. Scale bar = 200μm. **d** Three-dimensional visualisation and representative

colour-coded images of the cortical bone thickness of non-loaded and loaded tibiae

of fed and fasted 2-hour fed young adult mice.

All data are means ± SEM. $n = 7$ fed mice, $n = 6$ fasted 2-hour fed mice. **c** Two-tailed

Student's *t*-test; * $p < 0.05$

Fig. 3 Cortical bone phenotype of control (non-loaded) and loaded tibiae with

1100 microstrain (sub-threshold) and 1300 microstrain (submaximal) loading

regimens in 19-week-old fed or overnight fasted 2-hour fed male C57BL/6 mice

a The effect of sub-threshold and **b** submaximal mechanical loadings with peak force

of 8N to induce peak strain magnitude of 1100 microstrain and 11N to induce peak

strain magnitude of 1300 microstrain respectively on Ct.Th of fed or fasted 2-hour

fed young adult mice. All data are means ± SEM, $n = 7$. **a, b** Two-way ANOVA with

post-hoc Tukey comparisons; * $p < 0.05$, ** $p < 0.01$

Fig. 4 Cortical and trabecular bone phenotypes of control (non-loaded) and

loaded tibiae with 2200 microstrain osteogenic loading regimen in 20-month-

old fed or overnight fasted 2-hour fed male C57BL/6 mice

a Three-dimensional images of a 1.0-mm-thick cortical bone region 0.9mm below the

growth plate from the mid-shaft of control non-loaded (left) and loaded (right) tibiae

of fed or fasted 2-hour fed aged mice; scale bar = 200μm. **b** The change in cortical

bone thickness (Ct.Th) of tibiae loaded at 10N to induce peak strain magnitude of

2200 microstrain compared with non-loaded controls of fed or fasted 2-hour fed aged

male mice measured using μCT at tibial mid-shaft. There was no significant

difference in Ct.Th of the loaded tibiae of fasted 2-hour fed mice compared with ad-

libitum fed mice. **c** Three-dimensional visualisation and representative colour-coded

images of the cortical bone thickness of non-loaded and loaded tibiae of fed and

fasted 2-hour fed aged adult mice. **d** Trabecular bone pattern factor (Tb.Pf) of the

right tibiae loaded at 10N to induce peak strain magnitude of 2200 microstrain

compared with non-loaded left controls of fed or fasted 2-hour fed aged male mice

measured at tibial proximal region using μCT. Box-plots represent means ± SEM. *n*

= 8. **b** Two-tailed Student's *t*-test **d** Two-way ANOVA with post-hoc Tukey

comparisons; ** $p < 0.01$, *** $p < 0.001$

Fig. 5 Serum concentrations of ghrelin, leptin, insulin, GIP and GLP1 in young

adult 17-week-old and aged 20-month-old fed or overnight fasted 2-hour fed

male C57BL/6 mice

a Serum ghrelin concentration significantly increased from 12:00 to 14:00 in young

fed mice and declined at 14:00 in young fasted fed mice compared with that in fed

mice. **b** Serum ghrelin concentrations were significantly higher at 14:00 and 15:00

compared to those at 12:00 in aged fed but not fasted fed mice. **c, d** In young and

aged fasted fed mice, serum leptin concentrations significantly increased at 14:00

and 15:00 compared with fasting serum concentration at 12:00 and remained

relatively unchanged in ad-lib fed mice. Concentrations of serum leptin was

significantly lower at 12:00 and significantly higher at 15:00 in young fasted fed mice

compared to those in ad-lib fed mice (c). **e** Serum insulin concentrations significantly

increased from 12:00 to 14:00 in young fasted fed mice and in comparison to that in

ad-lib fed mice, then significantly decreased at 15:00. **f** In aged fasted fed mice,

serum insulin concentrations increased but not significantly from 12:00 to 14:00, then

decreased significantly at 15:00 compared with that at 14:00 and remained

unchanged in fed mice. **g, i** In young fasted fed mice, serum GIP and GLP1

concentrations significantly increased at 14:00 and 15:00 compared with fasting

concentrations at 12:00 and remained unchanged in fed mice. Serum GLP1

concentrations were significantly higher at 15:00 in fasted fed mice compared to that

in ad-lib fed mice (i). **h, j** There were no significant changes in serum GIP and GLIP1

concentrations within and between aged fed and fasted fed mice. All data are means

± SEM. *n* = 6 ad-libitum fed mice, *n* = 7 overnight (16-hour) fasted 2-hour fed mice.

Two-way ANOVA with post-hoc Tukey comparisons; * $p < 0.05$, ** $p < 0.01$, *** $p <$

0.001, **** $p < 0.0001$

698   **Tables**

699   **Table 1 Effect of 2-hour feeding following overnight fast on load-induced**

700   **cortical bone formation in young adult mice**

	Fed			Fasted Fed 2hrs		
	Loaded	**Non-loaded**	*p* value	**Loaded**	**Non-loaded**	*p* value
MS (%)	1.03 ±0.02	0.55±0.06	< 0.01	1.35±0.15	0.6±0.05	< 0.0001
MAR (μm/day)	2.0±0.13	0.33±0.15	< 0.05	4.9±0.8	0.07±0.06	< 0.0001
BFR/BS ($\mu m^3/\mu m^2$/day)	1.04± 0.07	0.12±0.06		3.57±0.8	0.02±0.01	< 0.0001

702   MS = mineralising surface; MAR = mineral apposition rate; BFR/BS = bone formation

703   rate. Values are mean ± SEM.

704   **Table 2 Serum concentrations of hormones altered during fasting and feeding**

705   **in young mice**

	Fed			
	12:00	**14:00**	**15:00**	*p* value
Hormone				
Ghrelin	703±162	1493±156	1326±119	<0.05[a]
Leptin	298±0.1	269±25	276±25	

Hormone	12:00	14:00	15:00	p value
Insulin	929±94	1023±133	1291±110	
GIP	54±3	50±3	51±2	
GLP1	8±1	7±1	7±1	

Fasted Fed

Hormone				
Ghrelin	1233±251	757±95[1]	1143±48	
Leptin	95±15[1]	437±62	601±76[3]	<0.0001[b]
Insulin	552±0.1	2175±303[2]	1319±107	<0.05[c]
GIP	44±2	58±2	56±3	0.01[b]
GLP1	7±1	10±1	11±1.[1]	0.05[b]

Two-way ANOVA used to assess the differences within and between the groups.

Comparisons of time points within the same group are indicated as [a]12:00 < 14:00;

[b]12:00 < 14:00,15:00; [c]12:00 < 14:00 > 15:00. Comparisons of same time points

between the groups are indicated as [1]$p < 0.05$; [2]$p < 0.01$; [3]$p < 0.001$. Samples were

assessed in duplicate. Results are expressed as pg/ml, mean ± SEM.

Table 3 Serum concentrations of hormones altered during fasting and feeding

in aged mice

	Fed			
	12:00	**14:00**	**15:00**	**p value**
Hormone				
Ghrelin	803±129	2272±359	2330±209	0.01[a]
Leptin	686±123	681±115	528±84	
Insulin	1000±99	914±87.1	1123±131	
GIP	46±2	43.1±1	45±2	
GLP1	6±1	7±1	6.±1	

Fasted Fed

Ghrelin	944±256	1862±277	1934±327	
Leptin	292±32	832±135	935±186	<0.05[a]
Insulin	570±0.1	1458±208	800±57	<0.05[b]
GIP	35±3	63±12	48±16	
GLP1	4±1	6±1	5±1	

713   Two-way ANOVA used to assess the differences within and between the groups.

714   Comparisons of time points within the same group are indicated as [a]12:00 <

715   14:00,15:00; [b]14:00 > 15:00. Samples were assessed in duplicate. Results are

716   expressed as pg/ml, mean ± SEM.

a

Fed

Fasted Fed 2hrs

Non-loaded   Loaded

b

Fed

Fasted Fed 2hrs

Non-loaded   Loaded

c

2200 microstrain

Ct.Th (mm)
change over non-loaded control

Fed   Fasted Fed 2hrs

d

Fed

Fasted Fed 2hrs

local thickness (voxel)
0   36   73

Non-loaded   Loaded

a

Fed — Non-loaded | Loaded
Fasted Fed 2hrs — Non-loaded | Loaded

b

2200 microstrain

Ct.Th (mm) change over non-loaded control

Fed | Fasted Fed 2hrs

c

Fed — Non-loaded | Loaded
Fasted Fed 2hrs — Non-loaded | Loaded

0 — 28 — 56
local thickness (voxel)

d

Left Control
Right Loaded

Tb.Pf (1/mm)

**

***

Fed | Fasted Fed 2hrs