Quantitative proteome and PTMome analysis of *Arabidopsis thaliana* root responses to persistent osmotic and salinity stress.

Rodriguez, MC.[1*]., Mehta, D[1*]., Tan, M[1], and Uhrig, RG[1]

[#]Corresponding Authors:

*Denotes equal contribution

Dr. R. Glen Uhrig

Department of Biological Sciences
University of Alberta
Edmonton, Alberta
Canada
T6G 2E9

Phone: +1 780 492 3088
E-Mail: ruhrig@ualberta.ca

[1]Department of Biological Sciences, University of Alberta, Edmonton, Alberta, Canada

Running Title: Impact of Salt and Osmotic Stress on Root Proteome

ABSTRACT

Environmental conditions contributing to abiotic stress such as drought result in large annual economic losses around the world. As sessile organisms, plants cannot escape the environmental stresses they encounter, but instead must adapt to survive. Studies investigating plant responses to osmotic and/or salt stress have largely focused on short-term systemic responses, leaving our understanding of intermediate to longer-term adaptation (24 h - days), less well understood. In addition to protein abundance and phosphorylation changes, evidence suggests reversible protein acetylation may also be important for abiotic stress responses. Therefore, to characterize protein-level effects of osmotic and salt stress, we undertook a label-free proteomic analysis of *Arabidopsis thaliana* roots exposed to 300 mM Mannitol and 150 mM NaCl for 24 hours. We assessed protein phosphorylation, acetylation and changes in abundance, detecting significant changes in the status of 106, 66 and 447 proteins, respectively. Comparison with available transcriptome data from analogous treatments, indicate that transcriptome- and proteome-level changes occur in parallel. Furthermore, significant changes in abundance, phosphorylation and acetylation involve different proteins from the same networks, indicating a concerted, multifaceted regulatory approach to prolonged osmotic and/or salt stress. Lastly, our quantitative proteomic approach uncovered a new root elongation protein, Armadillo repeat protein 2 (ARO2), which exhibits a salt stress dependent phenotype. Collectively, our findings indicate dynamic protein-level changes continue to occur in plant roots 24 hours from the onset of osmotic and salt stress and that these changes differ across multiple levels of the proteome.

Key Words: ***Arabidopsis thalian****a,* **Protein Acetylation, Protein Phosphorylation, Post-Translational Modifications, Quantitative Proteomics**

INTRODUCTION

Environmental conditions contributing to abiotic stresses such as drought and salinity result in large annual economic losses around the world. Between 2005 and 2015, drought was responsible for 30 % of agricultural loss, totaling to USD 19.5 billion (Food and Agriculture Organization of the United Nations (FAO), 2017). The combined result of human activities and climate change represent two of the major environmental factors currently threatening global food security (Ghosh and Xu, 2014), with rising global temperatures extending drought periods and lengthening plant drought recovery time (Schwalm et al., 2017). More frequent and severe droughts have left farmers to depend on progressively unsustainable irrigation and fertilization practices to maintain crop yield, placing further stress on drainage patterns and salinity levels of farmland (Kaushal et al., 2018). Both drought and salinity stress create a hyperosmotic condition typically referred to as osmotic stress, while increased salinity has the additive effect of concurrently increasing ionic strength. These osmotic and ionic effects massively impede the ability of roots to take up water, forcing molecular, cellular and/or morphological changes in order to adapt and survive.

All plant organs must adapt to drought and salt stress; however, roots are at the frontlines and are thus responsible for adjusting their molecular program as necessary to ensure whole plant survival. In particular, under water-limiting conditions, roots restructure their investment from lateral root formation to axile roots to forage deeper soil layers (Paez-Garcia et al., 2015). For example, overexpression of the cell wall expansion protein EXPA5, stimulates root cell elongation by lengthening the root tip rather than increasing cell number (Xu et al., 2014). While DEEPER ROOTING 1 (DRO1), a rice quantitative trait locus, is capable of increasing root angle in response to gravity, creating a steeper and deeper root system architecture, in addition to a higher photosynthetic capability, less wilting, and a >30 % increase in filled grain under severe drought (Uga et al., 2013). Thus, root system changes help define the growth and structure of aboveground biomass.

Within all plant cells, proteins are at the heart of sensing and initiating signaling events that then launch stress-adaptive responses. While the majority of our understanding of drought responses has been driven by transcriptomic responses (Kreps et al., 2002; Killan et al 2007), understanding a plant's environmental stress response and adaptation over time at the protein-level through quantitative proteomics, offers new opportunities for both fundamental understanding and biotechnological application.

Osmotic and salt stress have a major influence over crop productivity worldwide (Ashraf and Akram, 2009), making it essential that we continue to refine our understanding of how plants respond to these perturbations in order to breed and/or engineer more climate resilient crops. Emphasizing this need are the multiple studies that have examined the effects of osmotic and/or salt stress across a range of photosynthetic eukaryotes using both global and targeted analyses to understand the range of molecular, biochemical, physiological and morphological responses these stresses elicit (Zhang et al., 2012). This compendium of osmotic and salt stress data encompass a vast multitude of plant systems, time-frames and applied technologies, ranging from minutes to days of stress exposure (Jiang et al., 2007), followed by transcriptomic (Kilian et al., 2007; Kreps et al., 2002; Rest et al., 2016; Yin et al., 2019), proteomic (Jiang et al., 2007; Koh et al., 2015; Luo et al., 2015; Zhang et al., 2012) and metabolomic (Shelden et al., 2016) analyses to assess how the root environment responds to either abiotic stress. However, given the large variability in sampling time frames and root growth approaches, coupled with the: 1) limited in-parallel examination of multiple stresses and proteoforms within the

same study, 2) known disconnect between changing transcripts and proteins (Baerenfaller et al., 2012; Petricka et
al., 2012; Reiland et al., 2009; Seaton et al., 2018; Uhrig et al., 2019) and 3) limited investigations of the role post-
translational modifications (PTMs) play in root osmotic and salt adaptation (Koh et al., 2015; Wu et al., 2016; Zhu,
2016), further quantitative proteomic analysis are required.

PTMs are a pillar of properly functioning eukaryotic cells. Of the known PTMs, protein phosphorylation
and acetylation are the two most abundant (Choudhary et al., 2009; Henriksen et al., 2012; Sharma et al., 2014;
Uhrig et al., 2019; Weinert et al., 2011), with plants maintaining a comparable phosphoprotein landscape to
humans   (PhosPhat,   http://phosphat.uni-hohenheim.de/;   (Heazlewood   et   al.,   2008),   ATHENA;
http://athena.proteomics.wzw.tum.de/;   (Mergner   et   al.,   2020);   Plant   PTM   Viewer,
https://www.psb.ugent.be/webtools/ptm-viewer/; (Willems et al., 2019). More recently, global analysis of protein
acetylation in Arabidopsis (Finkemeier et al., 2011; Hartl et al., 2017; Liu et al., 2018; Uhrig et al., 2019; Wu et al.,
2011) and multiple crop species (Nallamilli et al., 2014; Smith-Hammond et al., 2014) have similarly identified
thousands of acetylation events occurring on proteins with diverse functions and subcellular localizations. These
two highly abundant PTMs have also been found to intersect on proteins involved in the same cellular processes
(Rao et al., 2014; Soufi et al., 2012; Tran et al., 2012; Uhrig et al., 2019) as well as occur on different proteins
involved in the same metabolic pathways (Uhrig et al., 2019), suggesting protein phosphorylation and acetylation
are functionally intertwined on multiple scales.

To better understand osmotic and salt stress responses in roots, we endeavored to examine the
intermediate-term responses (24 h) of *Arabidopsis thaliana* (Arabidopsis) roots subjected to either 300 mM
Mannitol or 150 mM NaCl. Then using label-free quantitative (LFQ) proteomics, we quantified the changes in the
Arabidopsis root proteome, phosphoproteome, and acetylome. Our findings demonstrate how protein abundance
and PTMs change relative to one another and to the transcriptome, in addition to understanding intersections
within, and between, each abiotic stress condition to reveal osmotic and/or salt-specific responses.

RESULTS

Quantification and localization of mannitol- and salt-induced protein phosphorylation, acetylation and
abundance changes
Twenty-one day old Arabidopsis seedlings grown in liquid MS media were subjected to either osmotic
(300 mM mannitol)- or salt (150 mM NaCl)-stress for 24 hours and assessed for quantifiable changes in protein
phosphorylation and/or acetylation status as well as protein abundance (Figure 1A). We quantified a total of
3,564 phosphorylated peptides and 281 acetylated peptides across 1,225 and 209 proteins, respectively, with a
PTM localization ≥ 0.75 (Supplemental Table 1), while a total of 3,113 proteins were quantified for changes in
abundance. Of these, 83 phospho- and 35 acetyl-peptides corresponding to 82 and 33 proteins, respectively,
exhibited significant changes under osmotic stress (*pval* ≤ 0.05; FC ≥ 1.5), while 56 phosphorylated and 59
acetylated peptides found on 55 and 56 proteins, respectively, exhibited significant change (*pval* ≤ 0.05; FC ≥ 1.5)
under salt stress (Figure 1B-D; Table 1; Supplemental Table 1). A total of 38 and 71 proteins exhibited significant
changes in abundance (*pval* ≤ 0.05; FC ≥ 1.5) under osmotic and salt stress (Figure 1B-D; Table 1; Supplemental
Table 1).

Next, we compared the $\log_2$ fold-change ($\log_2$FC) distribution of significantly changing proteins and PTMs
between osmotic- and salt-treated roots. This revealed both up- and down-regulation of phosphorylation events
and protein abundance, while finding only increases in protein acetylation (Figure 1B-D). Further analysis of
proteins exhibiting a significant change in either abundance, phosphorylation or acetylation found a total of 10
(10.2 %), 29 (27.4 %) and 27 (44.3 %) proteins overlapping between stress conditions (Figure 1B-D). Lastly, to
understand where intermediate-term changes in the proteome are occurring, we assessed the predicted
subcellular localization for all proteins exhibiting a significant change in PTM status or abundance under either
osmotic- or salt-stress. Here, we found differences in the subcellular distribution of significantly changing proteins
between each treatment (Figure 1E), with mannitol predominantly inducing changes in nuclear, mitochondrial and
plasma membrane proteins and salt eliciting greatest change in the plastid and extracellular / ER / Golgi localized
proteins (Figure 1E).

Overlap between changing PTMome and proteome is limited

Often, PTMs are quantified independent of assessing changes in proteome abundance. Therefore,
reported changes in PTM levels may also be due to fluctuations in protein abundance. To better understand this
here, we examined proteins exhibiting PTM changes for concurrent changes in protein abundance. Under osmotic
stress 21 (26 %) and 26 (79 %) proteins exhibiting significant changes in either phosphorylation or acetylation
were also quantified at the proteome level, however no PTM modified proteins concurrently exhibited a significant
change in PTM and protein abundance (Table 1). Similarly, under salt stress, 19 (35 %) and 43 (78 %)
phosphorylated and acetylated proteins, respectively, were quantified in our proteome data (Table 1), with only a
single acetylated protein exhibiting a concurrent change in protein abundance. Our concurrent assessment of
proteome and PTM abundance changes provides increased confidence that observed phosphorylation and
acetylation changes are related to PTM-level changes, and not changes in proteome abundance. Moreover, since
assessing changes in PTMs requires enrichment of modified peptides, it is likely that the remainder of PTM
modified proteins not observed in our protein abundance measurements are simply below detection.

**Gene Ontology (GO) analysis of proteome and PTM fluctuations reveals differences between osmotic-
and salt-induced stress.**

To better understand what cell processes are changing at the proteome-level after 24 h of osmotic or salt
stress, we performed a gene ontology (GO) enrichment analysis for biological process GO terms (Supplemental
Table 2). In response to salt stress, we found a number of significantly enriched GO terms including: sulfur
compound biosynthetic process (GO:0044272), sulfate reduction (GO:0019419), sulfur compound metabolic
processes (GO:0006790), seed coat development (GO:0010214), S-glycoside metabolic processes
(GO:0016143), glucosinolate metabolic processes (GO:0019757), in addition to metabolic activities related to
adenylyl-sulfate reductase activity (GO:0009973) (Table 2).
We then performed a KEGG pathway and protein domain analysis using Thalemine
(https://bar.utoronto.ca/thalemine/). Here, we found that proteins involved in primary metabolism, carbon fixation,
and amino acid metabolism exhibited significant change in their acetylation status in response to either osmotic or
salt stress (Supplemental Table 3). Alternatively, proteins exhibiting a significant change in abundance in
response to salt stress alone were enriched in proteins involved in 2-oxocarboxylic acid metabolism or

glucosinolate biosynthesis, supporting our previous GO enrichment analysis (Table 2). Furthermore, we queried
our data for enrichment of protein domains and found an enrichment of protein domains amongst proteins
significantly changing in abundance under salt stress (Supplemental Table 3). This includes multiple chaperone
and chaperonin domains, in addition to glycoside hydrolase domain 35, which supports our GO and KEGG
findings of augmented S-glycoside metabolic processes (GO:0016143) in salt stressed roots.

Comparative analysis of proteome and transcriptome changes reveals osmotic and salt-response
disconnects.
It is well established that transcriptome and proteome changes are often disconnected in their temporal
peak expression patterns (Baerenfaller et al., 2012; Graf et al., 2017; Seaton et al., 2018). Using publicly available
microarray transcriptome data from Arabidopsis plants subjected to analogous treatments (Kilian et al., 2007), we
performed a comparative analysis to better understand the temporal dynamics of transcripts that correspond to
proteins exhibiting phosphorylation, acetylation or abundance changes upon 24 h of osmotic- or salt-stress.
Further, we sought to cluster corresponding transcripts based on their expression dynamics and investigate how
these dynamics were reflected at the level of protein abundance as well as acetylation or phosphorylation status
(Figure 2-3; Supplemental Table 4).
Under both osmotic- and salt-stress conditions, the majority of proteins exhibiting a significant change in
abundance, phosphorylation and/or acetylation 24 h post-stress maintained a minimal log_2FC transcriptional
change (Cluster 1 - Figure 2-3). We also saw protein clusters emerge under both osmotic- and salt-stress
whereby an increasing trend in their transcript abundance from stress onset (0 h) to 24 h post-stress paralleled
our observed end-point proteome changes at 24 h (Cluster 2 - Figure 2-3). In the case of osmotic stress, the
parallel increase in transcript and protein abundance is also reflected in the protein phosphorylation levels
(Cluster 2 - Figure 2).
We then evaluated each cluster for the enrichment of biological process GO terms to uncover how
transcriptome and proteome responses may be related. Here, we found enriched GO terms for Cluster 1 of
osmotic-stressed roots and Cluster 1 and 2 of salt-stressed roots (Figure 2-3). Under osmotic stress GO terms
corresponding to Cluster 1, included multiple metabolic terms (gluconeogenesis; glycolytic process; oxidative
phosphorylation) and stress response terms (response to metal ion and temperature) (Figure 2). Comparatively,
Cluster 1 and 2 of salt stressed roots were similarly enriched in metabolic processes (gluconeogenesis
GO:0006094, glycolysis GO:0006096), in addition to salt-specific terms related to sulfate assimilation (Figure 3).
By clustering the significantly changing proteome and PTMome based on their corresponding
transcriptional expression profile from a comparably acquired, publicly available dataset, we find that transcript
changes are generally a good indicator of proteome changes 24 h post-osmotic or -salt stress initiation in
Arabidopsis roots. Conversely, we find that transcriptional changes are a poor indicator of PTM-level changes.
Overall, our data finds that osmotic and salt-stress induced proteomes, when clustered by transcriptional change
profile (0 h to 24 h), reveals key differences in how Arabidopsis roots respond to these stresses after 24 h.

Network analysis of proteins exhibiting osmotic- and salt-induced changes in PTM status and/or abundance reveals how the proteomes intersect

Using a combination of STRING-DB association network analyses (https://string-db.org/) and SUBA4 (https://suba.live/) protein subcellular localization information, we assembled subcellular localization-resolved protein association networks to elucidate the degree of interconnectedness between osmotic- and salt-induced PTM and protein abundance changes (Figure 4-5). Proteins with an edge score threshold of 0.5 or below were excluded from visualization. Here, we find that changes in the PTM status and/or protein abundance of metabolic proteins were most common between both osmotic and salt stress conditions (Figure 4-5). This is highlighted by proteins involved in both the transport and assimilation of sulfur as well as carbon metabolism (Figure 4-5). Similarly, changes in RNA processing and protein translation were also common between osmotic and salt treatments, indicating a likely adjustment to transcribed RNAs and a corresponding diversification of translated proteoforms. Conversely, clusters of significantly changing proteins related to sulfur and sucrose metabolism, as well as glucosinolate metabolism, were exclusively found to be salt-induced (Figure 4).

Our association networks further validate earlier GO and KEGG enrichment analyses in addition to visualizing the strength of these relationships using compiled meta-association data between proteins. These networks also provided a means by which to integrate additional data types including: the type and degree of change in acetylation, phosphorylation and/or abundance (outer ring), in addition to the subcellular localization of each protein within the plant cell (inner circle). Together, this data highlights the multi-dimensional relationships that exist between proteins exhibiting significant changes in either abundance or PTM status in response to osmotic and/or salt stress.

Root phenotyping identifies a potential new regulator of root elongation

To validate the findings of our quantitative proteomics workflow, we undertook root phenotypic analysis of an Armadillo repeat protein 2 (ARO2; AT5G66200) insertional mutant (*aro2-1*; SALK_003387). ARO2 exhibited the largest fold-change in abundance (Log$_2$FC= -2.8) of any protein in our NaCl treatment, rendering it a logical candidate (Supplemental Table 1; Figure 1B; Figure 4). Phenotypic analysis of *aro2-1* root length after growth on 300 mM mannitol and 150 mM NaCl revealed *aro2-1* possessed significantly shorter roots under salt stress by day 11 (n = 100; Student t-test *pval ≤ 0.05*) (Figure 6). This is consistent with our hypothesis that by 24 h we are measuring protein-level changes that relate to longer-term adjustments being made by Arabidopsis roots in response to these stresses.

DISCUSSION

Protein acetylation shows unique upward trend under salt and drought stress

Protein acetylation is almost universally upregulated under both osmotic and salt stress (Figure 1B). Many of the acetylated proteins we find here represent core metabolic proteins central to plant survival. For example, under both osmotic and salt treatments, key metabolic enzymes, such as UGP1 and PGK1, were acetylated (Figure 4-5). Protein acetylation has been shown to be a regulatory PTM (Zhao et al., 2010). Catalyzed by acetyltransferases, protein acetylation requires the primary metabolite acetyl-CoA. Acetyl CoA is a central intermediate metabolite between catabolic and anabolic pathways and as such, it is uniquely placed to exert either feedback or feedforward regulation through reversible protein acetylation. Multiple studies have found that

N$^\varepsilon$ protein acetylation is an abundant PTM across plants (De Boor et al., 2015; Finkemeier et al., 2011; Hartl et al., 2017; Li et al., 2018; Uhrig et al., 2019; Wu et al., 2011). In particular, it was found that the glycolytic and specialized metabolic proteins were extensively acetylated under nitrogen stress in tea leaves (Jiang et al., 2018). N$^\alpha$ protein acetylation has also been connected to drought resistance and is suggested to be regulated by ABA stress response (Linster et al., 2015; Linster et al., 2020). However, since both N$^\alpha$ and N$^\varepsilon$ protein acetylation rely on acetyl-CoA as an acetate donor, there are likely some higher-level intersections between these two forms of protein acetylation. Highlighting this possibility are the recent discovery of dual-specificity GNAT Acetyltransferases in Arabidopsis (Bienvenut et al., 2020).

Sulfur metabolism needed for short- and intermediate-term drought response

During drought stress, abscisic acid (ABA) is synthesized in the roots and then transported to the shoots to induce stomatal closure (Agurla et al., 2018). However, ABA biosynthesis in the roots requires sulfur. Plant root sulfate assimilation is primarily driven by adenosine 5′-phosphosulfate reductase (APR), which reduces sulfate to sulfite followed by further reduction to sulfide by ferredoxin dependent sulfite reductase (SiR) (Takahashi et al., 2011). Sulfide is then fixed by O-acetylserine(thiol)lyase into Cysteine (Cys) (Chan et al., 2013) to act as the sulfur donor for the sulfuration of molybdenum cofactor (Moco) sulfurase (Xiong et al., 2001), which is then necessary to initiate abscisic aldehyde oxidase 3 (AAO3) (Mendel, 2002; Schwarz and Mendel, 2006) and the ultimate production of ABA.

Studies have determined that Cys levels attune ABA biosynthesis (Batool et al., 2018; Cao et al., 2014), which in turn, is ultimately limited by sulfur assimilation at the level of APR (Takahashi et al., 2011). APR enzymatic activity is driven by the demand for sulfur, meaning that the uptake and assimilation of sulfur during drought is likely foundational to the short-term drought response. At 24 h of NaCl stress, we observed a number of key sulfur assimilating enzymes decrease in their protein abundance, including: ATP sulfurylase 1 (APS1) (Log$_2$FC = -1.37), adenosine 5′-phosphosulfate reductase (APR1; Log$_2$FC = -0.64), APR2 (Log$_2$FC = -1.69) and APR3 (Log$_2$FC = -1.75) (Figure 3). This suggests that by 24 h post salt-stress initiation plants have transitioned from utilizing short-term adaptations, to implementing longer term solutions. Transcriptionally, APR is regulated by sulfur availability (Hirai et al., 2003), demand for Cys (Davidian and Kopriva, 2010), and also in response to environmental stressors such as high salt (Koprivova et al., 2008). Specifically, Koprivova and colleagues (2008) reported a 3-fold increase in APR mRNA after 5 h of 150 mM NaCl in Arabidopsis roots. Using the AtGenExpress Arabidopsis stress expression dataset, we calculated a peak expression of APR transcripts at 6 h post 150 mM NaCl stress initiation, which then was found to continually decrease through to 24 h (Kilian et al., 2007; Supplemental Table 4).

Interestingly, an opposing trend in shoot APR expression is observed (Kilian et al., 2007). While APR in roots peaks soon after high salt exposure, APR in shoots remains low until it eventually peaks at 24 h. This points to APR and sulfur driven drought-responses acting as a short-term responder in the roots and an intermediate responder in the shoots. Given this, a sharp increase in root APR levels may facilitate the sulfur assimilation required for ABA production, which then leads to prompt stomatal closure. A later peak in shoot APR may be useful for intermediate drought response mechanisms, such as reducing reactive oxygen species (ROS).

Salt stress is also known to increase the production of oxidative enzymes (e.g. NADPH oxidase), which
are necessary for signaling and response during abiotic stress (Mittova et al., 2003), with subsequent reduction in
ROS required as the salt stress persists (Hasegawa and Bressan, 2000). To mitigate the damaging effects of
ROS, plants can produce glutathione (GSH), an antioxidant derived from Glu, Cys, and Gly. Stress-induced GSH
production requires sulfur-containing Cys, with high levels of Cys found to enhance synthesis of GSH in plants
(Takahashi et al., 2011). Demand for both, Cys and GSH, reciprocally regulates sulfur assimilation through APR
(Davidian and Kopriva, 2010), with depletion of GSH pools under prolonged drought stress conditions driving the
production of APR (Chan et al., 2013). Thus, upregulation of APR activity in the shoots may contribute to the
production of antioxidants such as GSH. Further time-course experimentation is required to fully elucidate the
interplay between salt-induced sulfur metabolism changes at the protein level in roots versus shoots, and is
beyond the scope of this study.
310

Acetylation of ANNAT1 annexin may increase peroxide activity during osmotic stress
Annexins are Ca^{2+} signaling membrane-binding proteins that possess peroxidase activity and are
involved in plant drought responses (Mortimer et al., 2008). Overexpression of ANNAT1 in Arabidopsis shoots
resulted in enhanced drought tolerance (Konopka-Postupolska et al., 2009) and while under salt stress, ANNAT1
was found to increase in a Ca^{2+} dependent manner (Lee et al., 2004). Furthermore, *annat1* plants are
hypersensitive to salt stress and ABA (Konopka-Postupolska et al., 2009; Lee et al., 2004). Under both osmotic
and salt treatment we find ANNAT1 to be acetylated ($Log_2FC = 1.801$). Interestingly, Arabidopsis ANNAT1 protein
expressed in, and purified from, *Nicotiana benthamiana* had three times more activity than bacterially expressed
AtANNAT1, suggesting that PTMs such as protein acetylation or some other binding partner specific to plants,
may make the observed peroxidase activity physiologically relevant (Gorecka et al., 2005). Previously, ANNAT1
peroxidase activity was thought to be dependent on phosphorylation events (Kim et al., 2015); however, our
evidence suggests that under osmotic stress, acetylation may be a primary regulatory PTM of ANNAT1. Overall,
the annexin-peroxidase connection could mean that annexins play a role in limiting the exposure and/or
accumulation of ROS during salt or osmotic stress-induced oxidative bursts in plants.
325

Osmotic and salt stress induce high-levels of UDP-glucose pyrophosphorylase (UGP1) acetylation
We find that UGP1 maintained the second largest increase in acetylation, with salt ($Log_2FC = 4.14$) and
osmotic ($Log_2FC = 3.33$) stress both eliciting large changes (Figure 2-3). UGP1 activity has previously been found
to be regulated by PTMs. For example, phosphorylation by Per-Arnt-Sim (PAS) kinase downregulates UGP1
activity resulting in a net decrease of glycogen in yeast (Rutter et al., 2002). Analysis of available osmotic- and
salt-stress induced transcript data found limited changes in UGP1 transcript levels under mannitol (-0.35) and salt
(-0.15) treatments 24 h post stress initiation (Kilian et al., 2007); however, UGP1 transcripts are also known to be
a poor indicator of protein-level or enzyme activity changes (Ciereszko et al., 2001).
UGP1 is an ubiquitous enzyme found in plants, animals, and bacteria that produces UDP-glucose (Meng
et al., 2009), which is the direct substrate for trehalose synthesis (Schluepmann and Matthew, 2009). Thus,
activation of UGP1 may be linked to increased trehalose biosynthesis in osmotic- and salt-stressed roots. In *E.*
coli, trehalose stabilizes proteins and prevents protein aggregation during osmotic stress (Moruno Algara et al.,

2019). This is also true for resurrection plants such as *Myrothamnus flabellifolius* where trehalose functions as a
protectant to stabilize membranes and proteins, allowing them to survive during dehydration-rehydration cycles
(Figueroa and Lunn, 2016; O'Hara et al., 2013). There is evidence that trehalose synthesis in Arabidopsis may
serve a similar function during drought stress (Lin et al., 2019). Furthermore, knockdown of UGP1 in *S. cerevisiae*
results in a reduction in trehalose as well as a reduction in oxidative stress resistance (Yi and Huh, 2015),
suggesting that UGP1 involvement in mitigating osmotic stress is an evolutionary conserved mechanism.

In plants, stachyose and raffinose family oligosaccharides (RFO) are degraded by UGP1 (PWY-6527;
https://biocyc.org/), releasing diphosphate. Thus an upregulation of UGP1 protein-levels and/or adjustments to its
activity could serve as a homeostatic mechanism for adjusting the nutritional status of a plant under drought
stress. Alternatively, if UGP1 acetylation decreases functionality, it may increase RFO content in the cell. RFOs
are suggested to have a dual role as osmoprotectants and compatible solutes (Elsayed et al., 2014). Other
groups have found higher RFO content in desiccated plant tissues (Obendorf and Gorecki, 2012) and that RFOs
stabilize membranes during dehydration (Pluskota et al., 2015). Overall, our data suggests that UGP1
functionality may be regulated by reversible acetylation under both osmotic and salt stress conditions. However,
whether UGP1 acetylation increases or decreases UPG1 activity, and how that activity confers osmotic
protection, is beyond the scope of this study.

Acetylation of GTP-Proteins may support root tip mitotic division
In Arabidopsis, GTP hydrolysis by Ran and Ran binding protein complexes are important in the regulation
of auxin-induced mitotic progression in root tips (Kim and Roux, 2003; Xu et al., 2016). In particular, both AtRAN1
and AtRAN3 are known to both be induced by salt treatment, with *ran1/ran3* double mutants showing higher
sensitivity to salt treatment, displaying significantly reduced root growth after being treated for 5 d with 100 mM
NaCl. GTPases are also known to directly intersect with ABA response and drought tolerance (Luo et al., 2013).

Our data supports the involvement of RAN-GTPases in osmotic and salt stress responses. In particular,
we find SIRANBP and RAN3 to be acetylated under salt ($Log_2FC = 2.01$ SIRANBP; $Log_2FC = 2.46$ RAN3) and
mannitol ($Log_2FC = 2.09$ SIRANBP; $Log_2FC = 1.55$ RAN3) treatment (Figure 2-3). Phylogenetic analysis of RAN-
GTPases across Eukaryotes finds a high sequence conservation between higher plants and vertebrates (Coppola
et al., 2018), rendering it interesting that Ran GTPases in humans and mice are also regulated by reversible
protein acetylation (De Boor et al., 2015). In these cases, increased acetylation of Ran-GTPases prevents the
degradation of Ran-GTP, leading to a higher supply of active Ran-GTP that more effectively induces mitosis (De
Boor et al., 2015). In particular, the acetylation of RAN-GTPases induces more efficient GTP loading to Ran,
which in turn increases Ran-GTPase activity for accurate chromosome-kinetochore attachment (Bao et al., 2018).
Therefore, our observation of increased AtRAN-GTPase acetylation may be a mechanism by which the plants are
trying to maintain root elongation under drought stress conditions.

Salt-dependent decreases in ARO2 protein abundance conditionally limit root elongation.
ARO2 is a member of the Armadillo repeat-containing proteins (ARCPs), which are conserved across all
eukaryotic proteomes, and are typically characterized by domains composed of multiple 42 amino acid Arm
repeats (Saito, 2006). ARCPs are involved in diverse cellular processes such as cytoskeletal regulation, signal
transduction, and transcription regulation (Mei et al., 2012). Previous research found ARCPs were essential for

proper microtubule assembly needed in areas of active mitosis, such as root tips. CSI1, another member of the
ARCPs, stabilizes microtubule assembly required for proper anther formation and root tip elongation (Mei et al.,
2012). Microtubular cytoskeleton regulation is not unique to CSI1 but a core characteristic of ARCPs (Sakai et al.,
2008; Sapiro et al., 2000). The relationship between microtubular regulation and ARPC was further exemplified by
ARO1, an orthologous relative of ARO2. ARO1 plays a key role in the organization of actin cytoskeleton, growth
polarization, and pollen tube growth (Gebert et al., 2008), rendering it likely that ARO2, like its ARCP relatives,
possess a similar function, but in root tip elongation. Under drought and salinity stress, water acquisition favours
deeper root systems (Wasson et al., 2012). Therefore, redistributing root branch density from near-surface to
deeper depths by elongating primary roots is essential (Uga et al., 2013). Interestingly, our data indicates that loss
of ARO2 is detrimental to root growth under salt-stress conditions, suggesting that ARO2 is a key component of
the salt-stress root elongation response in Arabidopsis (Figure 6).
388
CONCLUSION
Here, we describe proteomic and PTMomic changes occurring in Arabidopsis roots exposed to 300 mM
mannitol and 150 mM NaCl for 24 h. By examining intermediate proteome responses, we have uncovered a
number of potential regulatory candidates and PTM events that may be key to longer-term survival under osmotic
and/or salt stress conditions. This is highlighted by the identification of a new salt-stress dependent root
elongation regulator ARO2. Furthermore, protein acetylation has emerged as a potentially interesting PTM in the
ability of plants to mount responses to either osmotic- or salt-stress conditions. In undertaking a quantitative
analysis of the proteome, phosphoproteome and acetylome, we were able to capture a wide swath of the
intermediate term protein responses. Moving forward, understanding the intermediate to long-term responses of
either osmotic or salt stress at the protein level through time-course experimentation, is especially important for
the development of new climate resilient crops.
400
MATERIALS & METHODS
Plant Growth and Harvesting
Arabidopsis thaliana Col-0 seeds were imbibed 2 days at 4 $^{\circ}$C in the dark prior to growth in Magenta
boxes containing 0.5 x MS media containing 0.5 % (w/v) sucrose. Plants were grown at 22 $^{\circ}$C under a 12 h light :
12 h dark photoperiod for 21 days prior to the application of 0.5 x MS media containing 300 mM mannitol
(osmotic) or 150 mM NaCl (salt). Seedlings were grown in either experimental or control media for an additional
24 h. Roots were manually separated from shoots and immediately frozen in liquid N_2. Roots were then ground in
a mortar and pestle under liquid N_2 and stored at -80 $^{\circ}$C until further use.

Sample Preparation and PTM Affinity Enrichments
Root tissue (800 mg) was extracted using 1:1.5 w/v extraction buffer (4 % SDS, 50 mM HEPES-KOH pH
8.0, 50 mM DTT). Samples were then processed and digested with trypsin (Promega) as previously described
(Uhrig et al., 2019). Subsequent enrichment of phosphorylated or acetylated peptides was performed using
titanium dioxide (NP-Ti02; Sachtopore; SNX 030S 005 #9205/1639) and anti-acetylated lysine IgG-coupled
agarose (ICP0388; ImmuneChem, https://www.immunechem.com) as previously described (Uhrig et al., 2019),
prior to analysis by mass spectrometry.

Liquid Chromatography MS/MS

A total of 1 µg of re-suspended peptide was injected by an Easy-nLC 1000 system (Thermo Scientific)

and separated on a Thermo Scientific™ EASY-Spray™ HPLC Column (75 µm x 500 mm; particle size 2.0 µm;

E803A). The column was equilibrated with 100 % solvent A (0.1 % formic acid (FA) in water). Peptides were

eluted using the following gradient of solvent B (0.1 % FA in ACN): 0-50 min; 0-25 % B, 50-60 min; 25-32 % B,

60-70 min; 32-98 % B at a flow rate of 0.3 µl/min. High accuracy mass spectra were acquired with an Orbitrap

Fusion (Thermo Scientific) that was operated in data dependent acquisition mode. All precursor signals were

recorded in the Orbitrap using quadrupole transmission in the mass range of 300-1500 m/z. Spectra were

recorded with a resolution of 120 000 at 190 m/z, a target value of 4E5 and the maximum cycle time was set to 3

seconds. Data dependent MS/MS were recorded in the linear ion trap using quadrupole isolation with a window of

2 Da and HCD fragmentation with 30 % fragmentation energy. The ion trap was operated in rapid scan mode with

a target value of 3E4 and a maximum injection time of 100 ms. Precursor signals were selected for fragmentation

with a charge state from +2 to +7 and a signal intensity of at least 1E4. A dynamic exclusion list was used for 30

seconds and maximum parallelizing ion injections was activated.

Mass Spectrometry Data Analysis

Raw data were processed using MaxQuant software (version 1.4.1.2, http://www.maxquant.org/)(Cox and

Mann, 2008) and searched against The Arabidopsis Information Resource (TAIR10) protein database

concatenated with a decoy database supplemented with contaminants using the Andromeda search engine.

Fixed modifications included carbamidomethylation of cysteine residues, while methionine oxidation (all

searches), lysine acetylation (acetylated peptide enrichments) and phosphorylated serine / threonine and tyrosine

(phosphopeptide enrichments) were set as variable modifications, respectively. One missed cleavage for

phosphoproteomic and proteomic measurements and two missed cleavages for lysine-acetylated peptides were

permitted. A protein and PSM false discovery rates (FDR) threshold of 1 % was used, while match between runs

and re-quantify options were enabled. Further downstream analysis was performed using Perseus version 1.5.5.3

(Tyanova et al., 2016). For all data reverse hits and contaminants were removed, followed by assembly into

treatment groups, filtering based on the presence of measured data in at least 2 replicates per group and log_2-

transformation with missing values replaced based on the normal distribution function set to default parameters.

All PTM analyses utilized a PTM site localization score threshold ≥ 0.75. Significant changes in either protein or

PTM peptide levels between each treatment condition and the control samples were assessed by Student's T-test

(*pval* ≤ 0.05). Mass spectrometry data acquired for this study can be found at the EMBL proteomic repository

Proteomics IDEntifications (PRIDE; https://www.ebi.ac.uk/pride/archive/; PXD019139).

Bioinformatics

To identify the biological functions of proteins changing in abundance, phosphorylation or acetylation, a

gene ontology (GO) analysis of biological processes was performed using the ONTOLOGIZER (http://ontologizer.de;

(Bauer et al., 2008). A parent–child intersection analysis approach was used (Benjamini–

Hochberg *FDR* correction *pval* ≤ 0.05). Clustering of Arabidopsis microarray time-course data was performed

using a trajectory-based longitudinal clustering algorithm implemented in the kml-shape package in R (Genolini et

al., 2016). For inputs, microarray counts were obtained from (Kilian et al., 2007) and fold changes were calculated

for each time point comparing treated samples versus the controls. The resulting dataset was filtered based on

the set of 619 proteins that were found to significantly change in either abundance, phosphorylation or acetylation

status at 24 h. The resulting transcript fold-changes over the time-course were used to create four clusters based

on the transcript trajectories over the 1, 3, 6, 12 and 24 h time points previously obtained (Kilian et al., 2007).

Clustered trajectories and corresponding protein data were plotted using the *ggplot2* and *cowplot* packages in R

version 3.3.1 (R Core Team 2016). Further characterization of connections between quantified and significantly

changing phospho- and acetyl-proteins was performed through association network analyses using the

Cytoscape StringDB plugin StringApp (http://apps.cytoscape.org/apps/stringapp;(Szklarczyk et al., 2017) with a

minimum association threshold of 0.5. Cytoscape 3.4 (http://www.cytoscape.org/) visualized networks were then

supplemented with subcellular localization data derived from SUBAcon (http://suba.live/; (Hooper et al., 2014)

Arabidopsis *aro2-1* root assay

A. thaliana wild type *Col-0* and insertional mutant (*aro2-1*; SALK_003387) seeds were imbibed for 2 days

at 4 °C. Seeds were then germinated for 4 days under a 12 h light : 12 h dark photoperiod. At the end of the fourth

day, seedlings were transferred to either control (0.5 X MS), salt stress (0.5 X MS + 150 mM NaCl), or mannitol

stress (0.5 X MS + 300 mM mannitol) plates for vertical growth and phenotypic analysis. Each plate contained 8

WT and *aro2-1* mutant seedlings. The seedlings were grown vertically on a custom, blacked-out 3D printed

vertical plate holders to minimize root light exposure. Root length was then measured every day for the following

7 days.

ACKNOWLEDGEMENT

The authors would like to thank the Functional Genomics Center Zurich for helpful discussions and the

National Science and Engineering Research Council of Canada (NSERC) for funding this research.

476
LITERATURE CITED

Agurla, S., Gahir, S., Munemasa, S., Murata, Y., and Raghavendra, A.S. (2018) Mechanism of Stomatal Closure
in Plants Exposed to Drought and Cold Stress BT - Survival Strategies in Extreme Cold and Desiccation:
Adaptation Mechanisms and Their Applications.
Ashraf, M., and Akram, N.A. (2009) Improving salinity tolerance of plants through conventional breeding and
genetic engineering: An analytical comparison. *Biotechnol Adv*. 27: 744–752.
Baerenfaller, K., Massonnet, C., Walsh, S., Baginsky, S., Bühlmann, P., Hennig, L., et al. (2012) Systems-based
analysis of Arabidopsis leaf growth reveals adaptation to water deficit. *Mol Syst Biol*. 8.
Bao, X., Liu, H., Liu, X., Ruan, K., Zhang, Y., Zhang, Z., et al. (2018) Mitosis-specific acetylation tunes Ran
effector binding for chromosome segregation. *J Mol Cell Biol*. 10: 18–32.
Batool, S., Uslu, V.V., Rajab, H., Ahmad, N., Waadt, R., Geiger, D., et al. (2018) Sulfate is incorporated into
cysteine to trigger ABA Production and stomatal closure. *Plant Cell*. 30: 2973–2987.
Bauer, S., Grossmann, S., Vingron, M., and Robinson, P.N. (2008) Ontologizer 2.0 - A multifunctional tool for GO
term enrichment analysis and data exploration. *Bioinformatics*. 24: 1650–1651.
Bienvenut, W. V, Brünje, A., Boyer, J., Mühlenbeck, J.S., Bernal, G., Lassowskat, I., et al. (2020) Dual lysine and
N-terminal acetyltransferases reveal the complexity underpinning protein acetylation. *Mol Syst Biol*. 16: 1–
23.
De Boor, S., Knyphausen, P., Kuhlmann, N., Wroblowski, S., Brenig, J., Scislowski, L., et al. (2015) Small GTP-
binding protein Ran is regulated by posttranslational lysine acetylation. *Proc Natl Acad Sci U S A*. 112:
E3679–E3688.
Cao, M.J., Wang, Z., Zhao, Q., Mao, J.L., Speiser, A., Wirtz, M., et al. (2014) Sulfate availability affects ABA
levels and germination response to ABA and salt stress in Arabidopsis thaliana. *Plant J*. 77: 604–615.
Chan, K.X., Wirtz, M., Phua, S.Y., Estavillo, G.M., and Pogson, B.J. (2013) Balancing metabolites in drought□:
the sulfur assimilation conundrum. *Trends Plant Sci*. 18: 18–29.

Choudhary, C., Kumar, C., Gnad, F., Nielsen, M.L., Rehman, M., Walther, T.C., et al. (2009) Lysine acetylation
targets protein complexes and co-regulates major cellular functions. *Science (80-)*. 325: 834–840.
Ciereszko, I., Johansson, H., Hurry, V., and Kleczkowski, L.A. (2001) Phosphate status affects the gene
expression, protein content and enzymatic activity of UDP-glucose pyrophosphorylase in wild-type and pho
mutants of Arabidopsis. *Planta*. 212: 598–605.
Coppola, U., Caccavale, F., Scelzo, M., Holland, N.D., Ristoratore, F., and D'Aniello, S. (2018) Ran GTPase, an
eukaryotic gene novelty, is involved in amphioxus mitosis. *PLoS One*. 13: 1–13.
Cox, J., and Mann, M. (2008) MaxQuant enables high peptide identification rates, individualized p.p.b.-range
mass accuracies and proteome-wide protein quantification. *Nat Biotechnol*. 26: 1367–1372.
Davidian, J.C., and Kopriva, S. (2010) Regulation of sulfate uptake and assimilation - The same or not the same?
Mol Plant. 3: 314–325.
Elsayed, A.I., Rafudeen, M.S., and Golldack, D. (2014) Physiological aspects of raffinose family oligosaccharides
in plants: Protection against abiotic stress. *Plant Biol*. 16: 1–8.
Figueroa, C.M., and Lunn, J.E. (2016) A tale of two sugars: Trehalose 6-phosphate and sucrose. *Plant Physiol*.
172: 7–27.
Finkemeier, I., Laxa, M., Miguet, L., Howden, A.J.M., and Sweetlove, L.J. (2011) Proteins of diverse function and
subcellular location are lysine acetylated in Arabidopsis. *Plant Physiol*. 155: 1779–1790.
Food and Agriculture Organization of the United Nations (FAO) (2017) The Impact of disasters and crises on
agriculture and Food Security.
Gebert, M., Dresselhaus, T., and Sprunck, S. (2008) F-actin organization and pollen tube tip growth in
Arabidopsis are dependent on the gametophyte-specific armadillo repeat protein ARO1. *Plant Cell*. 20:
2798–2814.
Genolini, C., Ecochard, R., Benghezal, M., Driss, T., Andrieu, S., and Subtil, F. (2016) kmlShape: An efficient
method to cluster longitudinal data (Time-Series) according to their shapes. *PLoS One*. 11: 1–24.
Ghosh, D., and Xu, J. (2014) Abiotic stress responses in plant roots: A proteomics perspective. *Front Plant Sci*. 5:
1–13.
Gorecka, K.M., Konopka-Postupolska, D., Hennig, J., Buchet, R., and Pikula, S. (2005) Peroxidase activity of
annexin 1 from Arabidopsis thaliana. *Biochem Biophys Res Commun*. 336: 868–875.
Graf, A., Coman, D., Uhrig, R.G., Walsh, S., Flis, A., Stitt, M., et al. (2017) Parallel analysis of *Arabidopsis*
circadian clock mutants reveals different scales of transcriptome and proteome regulation. *Open Biol*. 7:
160333.
Hartl, M., Füßl, M., Boersema, P.J., Jost, J., Kramer, K., Bakirbas, A., et al. (2017) Lysine acetylome profiling
uncovers novel histone deacetylase substrate proteins in Arabidopsis . *Mol Syst Biol*. 13: 949.
Hasegawa, P.M., and Bressan, R.A. (2000) Plant Cellular and Molecular Responses to High Salinity. *Annu Rev*
Plant Biol. 51: 463–499.
Heazlewood, J.I., Durek, P., Hummel, J., Selbig, J., Weckwerth, W., Walther, D., et al. (2008) PhosPhAt□: A
database of phosphorylation sites in Arabidopsis thaliana and a plant-specific phosphorylation site predictor.
Nucleic Acids Res. 36: 1015–1021.
Henriksen, P., Wagner, S.A., Weinert, B.T., Sharma, S., Bačinskaja, G., Rehman, M., et al. (2012) Proteome-wide
analysis of lysine acetylation suggests its broad regulatory scope in saccharomyces cerevisiae. *Mol Cell*
Proteomics. 11: 1510–1522.
Hirai, M.Y., Fujiwara, T., Awazuhara, M., Kimura, T., Noji, M., and Saito, K. (2003) Global expression profiling of
sulfur-starved Arabidopsis by DNA macroarray reveals the role of O-acetyl-L-serine as a general regulator of
gene expression in response to sulfur nutrition. *Plant J*. 33: 651–663.
Hooper, C.M., Tanz, S.K., Castleden, I.R., Vacher, M.A., Small, I.D., and Millar, A.H. (2014) SUBAcon: A
consensus algorithm for unifying the subcellular localization data of the Arabidopsis proteome.
Bioinformatics. 30: 3356–3364.
Jiang, J., Gai, Z., Wang, Y., Fan, K., Sun, L., Wang, H., et al. (2018) Comprehensive proteome analyses of lysine
acetylation in tea leaves by sensing nitrogen nutrition. *BMC Genomics*. 19: 1–13.
Jiang, Y., Yang, B., Harris, N.S., and Deyholos, M.K. (2007) Comparative proteomic analysis of NaCl stress-
responsive proteins in Arabidopsis roots. *J Exp Bot*. 58: 3591–3607.
Kaushal, S.S., Likens, G.E., Pace, M.L., Utz, R.M., Haq, S., Gorman, J., et al. (2018) Freshwater salinization
syndrome on a continental scale. *Proc Natl Acad Sci U S A*. 115: E574–E583.
Kilian, J., Whitehead, D., Horak, J., Wanke, D., Weinl, S., Batistic, O., et al. (2007) The AtGenExpress global
stress expression data set: Protocols, evaluation and model data analysis of UV-B light, drought and cold
stress responses. *Plant J*. 50: 347–363.
Kim, D., Ntui, V.O., Zhang, N., and Xiong, L. (2015) Arabidopsis Yak1 protein (AtYak1) is a dual specificity protein
kinase. *FEBS Lett*. 589: 3321–3327.
Kim, S.H., and Roux, S.J. (2003) An Arabidopsis Ran-binding protein, AtRanBP1c, is a co-activator of Ran

GTPase-activating protein and requires the C-terminus for its cytoplasmic localization. *Planta*. 216: 1047–1052.

Koh, J., Chen, G., Yoo, M.J., Zhu, N., Dufresne, D., Erickson, J.E., et al. (2015) Comparative Proteomic Analysis of Brassica napus in Response to Drought Stress. *J Proteome Res*. 14: 3068–3081.

Konopka-Postupolska, D., Clark, G., Goch, G., Debski, J., Floras, K., Cantero, A., et al. (2009) The role of annexin 1 in drought stress in Arabidopsis. *Plant Physiol*. 150: 1394–1410.

Koprivova, A., North, K.A., and Kopriva, S. (2008) Complex signaling network in regulation of adenosine 5′-phosphosulfate reductase by salt stress in arabidopsis roots. *Plant Physiol*. 146: 1408–1420.

Kreps, J.A., Wu, Y., Chang, H.S., Zhu, T., Wang, X., and Harper, J.F. (2002) Transcriptome changes for Arabidopsis in response to salt, osmotic, and cold stress. *Plant Physiol*. 130: 2129–2141.

Lee, S., Lee, E.J., Yang, E.J., Lee, J.E., Park, A.R., Song, W.H., et al. (2004) Proteomic of identification of annexins, calcium-dependent membrane binding proteins that mediate osmotic stress and abscisic acid signal transduction in arabidopsis. *Plant Cell*. 16: 1378–1391.

Li, X., Ye, J., Ma, H., and Lu, P. (2018) Proteomic analysis of lysine acetylation provides strong evidence for involvement of acetylated proteins in plant meiosis and tapetum function. 142–154.

Lin, Q., Yang, J., Wang, Q., Zhu, H., Chen, Z., Dao, Y., et al. (2019) Overexpression of the trehalose-6-phosphate phosphatase family gene AtTPPF improves the drought tolerance of Arabidopsis thaliana. *BMC Plant Biol*. 19: 1–15.

Linster, E., Layer, D., Bienvenut, W. V., Dinh, T. V., Weyer, F. A., Wiebke, L., et al. (2020) The Arabisopsis N-acetyltransferase NAA60 locates to the plasma membrane and is vital for the high salt stress response. *New Phytol*. 2: 554-546.

Linster, E., Stephan, I., Bienvenut, W. V., Maple-Grødem, J., Myklebust, L.M., Huber, M., et al. (2015) Downregulation of N-terminal acetylation triggers ABA-mediated drought responses in Arabidopsis. *Nat Commun*. 6.

Liu, S., Yu, F., Yang, Z., Wang, T., Xiong, H., Chang, C., et al. (2018) Establishment of dimethyl labeling-based quantitative acetylproteomics in Arabidopsis. *Mol Cell Proteomics*. 17: 1010–1027.

Luo, J., Tang, S., Peng, X., Yan, X., Zeng, X., Li, J., et al. (2015) Elucidation of cross-talk and specificity of early response mechanisms to salt and PEG-simulated drought stresses in brassica napus using comparative proteomic analysis. *PLoS One*. 10: 1–24.

Luo, Y., Wang, Z., Ji, H., Fang, H., Wang, S., Tian, L., et al. (2013) An Arabidopsis homolog of importin β1 is required for ABA response and drought tolerance. *Plant J*. 75: 377–389.

Mei, Y., Gao, H.B., Yuan, M., and Xue, H.W. (2012) The Arabidopsis ARCP protein, CSI1, which is required for microtubule stability, is necessary for root and anther development. *Plant Cell*. 24: 1066–1080.

Mendel, R.R. (2002) Molybdoenzymes and molybdenum cofactor in plants. *J Exp Bot*. 53: 1689–1698.

Meng, M., Geisler, M., Johansson, H., Harholt, J., Scheller, H. V., Mellerowicz, E.J., et al. (2009) UDP-glucose pyrophosphorylase is not rate limiting, but is essential in arabidopsis. *Plant Cell Physiol*. 50: 998–1011.

Mergner, J., Frejno, M., List, M., Papacek, M., Chen, X., Chaudhary, A., et al. (2020) Mass-spectrometry-based draft of the Arabidopsis proteome. *Nature*. 579: 409–414.

Mittova, V., Tal, M., Volokita, M., and Guy, M. (2003) Up-regulation of the leaf mitochondrial and peroxisomal antioxidative systems in response to salt-induced oxidative stress in the wild salt-tolerant tomato species Lycopersicon pennellii. *Plant, Cell Environ*. 26: 845–856.

Mortimer, J.C., Laohavisit, A., Macpherson, N., Webb, A., Brownlee, C., Battey, N.H., et al. (2008) Annexins: Multifunctional components of growth and adaptation. *J Exp Bot*. 59: 533–544.

Moruno Algara, M., Kuczyńska-Wiśnik, D., Dębski, J., Stojowska-Swędrzyńska, K., Sominka, H., Bukrejewska, M., et al. (2019) Trehalose protects Escherichia coli against carbon stress manifested by protein acetylation and aggregation. *Mol Microbiol*. 112: 866–880.

Nallamilli, B.R.R., Edelmann, M.J., Zhong, X., Tan, F., Mujahid, H., Zhang, J., et al. (2014) Global analysis of lysine acetylation suggests the involvement of protein acetylation in diverse biological processes in rice (Oryza sativa). *PLoS One*. 9: 1–12.

O'Hara, L.E., Paul, M.J., and Wingler, A. (2013) How do sugars regulate plant growth and development? new insight into the role of trehalose-6-phosphate. *Mol Plant*. 6: 261–274.

Obendorf, R.L., and Gorecki, R.J. (2012) Soluble carbohydrates in legume seeds. *Seed Sci Res*. 22: 219–242.

Paez-Garcia, A., Motes, C.M., Scheible, W.R., Chen, R., Blancaflor, E.B., and Monteros, M.J. (2015) Root traits and phenotyping strategies for plant improvement. *Plants*. 4: 334–355.

Petricka, J.J., Schauer, M.A., Megraw, M., Breakfield, N.W., Thompson, J.W., Georgiev, S., et al. (2012) The protein expression landscape of the Arabidopsis root. *Proc Natl Acad Sci U S A*. 109: 6811–6818.

Pluskota, W.E., Szablińska, J., Obendorf, R.L., Górecki, R.J., and Lahuta, L.B. (2015) Osmotic stress induces genes, enzymes and accumulation of galactinol, raffinose and stachyose in seedlings of pea (Pisum sativum L.). *Acta Physiol Plant*. 37.

619   Rao, R.S.P., Thelen, J.J., and Miernyk, J.A. (2014) Is Lys-Ne{open}-acetylation the next big thing in post-
620   translational modifications? *Trends Plant Sci*. 19: 550–553.
621   Reiland, S., Messerli, G., Baerenfaller, K., Gerrits, B., Endler, A., Grossmann, J., et al. (2009) Large-scale
622   arabidopsis phosphoproteome profiling reveals novel chloroplast kinase substrates and phosphorylation
623   networks1[w]. *Plant Physiol*. 150: 889–903.
624   Rest, J.S., Wilkins, O., Yuan, W., Purugganan, M.D., and Gurevitch, J. (2016) Meta-analysis and meta-regression
625   of transcriptomic responses to water stress in Arabidopsis. *Plant J*. 85: 548–560.
626   Rutter, J., Probst, B.L., and McKnight, S.L. (2002) Coordinate regulation of sugar flux and translation by PAS
627   kinase. *Cell*. 111: 17–28.
628   Saito, K. (2006) Sulfur Assimilatory Metabolism. The Long and Smelling Road. 136: 2443–2450.
629   Sakai, T., Honing, H. Van Der, Nishioka, M., Uehara, Y., Takahashi, M., Fujisawa, N., et al. (2008) Armadillo
630   repeat-containing kinesins and a NIMA-related kinase are required for epidermal-cell morphogenesis in
631   Arabidopsis. *Plant J*. 53: 157–171.
632   Sapiro, R., Tarantino, L.M., Velazquez, F., Kiriakidou, M., Hecht, N.B., Bucan, M., et al. (2000) Sperm antigen 6 is
633   the murine homologue of the Chlamydomonas reinhardtii central apparatus protein encoded by the PF16
634   locus. *Biol Reprod*. 62: 511–518.
635   Schluepmann, H., and Matthew, P. (2009) Trehalose Metabolites in Arabidopsis — elusive , active and central
636   Published By⬚: The American Society of Plant Biologists Trehalose Metabolites in Arabidopsis — elusive ,
637   active and central. 7.
638   Schwalm, C.R., Anderegg, W.R.L., Michalak, A.M., Fisher, J.B., Biondi, F., Koch, G., et al. (2017) Global patterns
639   of drought recovery. *Nature*. 548: 202–205.
640   Schwarz, G., and Mendel, R.R. (2006) Molybdenum Cofactor Biosynthesis and Molybdenum Enzymes. *Annu Rev*
641   *Plant Biol*. 57: 623–647.
642   Seaton, D.D., Graf, A., Baerenfaller, K., Stitt, M., Millar, A.J., and Gruissem, W. (2018) Photoperiodic control of
643   the Arabidopsis proteome reveals a translational coincidence mechanism . *Mol Syst Biol*. 14: 1–19.
644   Sharma, K., D'Souza, R.C.J., Tyanova, S., Schaab, C., Wiśniewski, J.R., Cox, J., et al. (2014) Ultradeep Human
645   Phosphoproteome Reveals a Distinct Regulatory Nature of Tyr and Ser/Thr-Based Signaling. *Cell Rep*. 8:
646   1583–1594.
647   Shelden, M.C., Dias, D.A., Jayasinghe, N.S., Bacic, A., and Roessner, U. (2016) Root spatial metabolite profiling
648   of two genotypes of barley (Hordeum vulgare L.) reveals differences in response to short-term salt stress. *J*
649   *Exp Bot*. 67: 3731–3745.
650   Smith-Hammond, C.L., Swatek, K.N., Johnston, M.L., Thelen, J.J., and Miernyk, J.A. (2014) Initial description of
651   the developing soybean seed protein Lys-Nε-acetylome. *J Proteomics*. 96: 56–66.
652   Soufi, B., Soares, N.C., Ravikumar, V., and Macek, B. (2012) Proteomics reveals evidence of cross-talk between
653   protein modifications in bacteria: Focus on acetylation and phosphorylation. *Curr Opin Microbiol*. 15: 357–
654   363.
655   Szklarczyk, D., Morris, J.H., Cook, H., Kuhn, M., Wyder, S., Simonovic, M., et al. (2017) The STRING database in
656   2017: Quality-controlled protein-protein association networks, made broadly accessible. *Nucleic Acids Res*.
657   45: D362–D368.
658   Takahashi, H., Kopriva, S., Giordano, M., Saito, K., and Hell, R. (2011) Sulfur Assimilation in Photosynthetic
659   Organisms: Molecular Functions and Regulations of Transporters and Assimilatory Enzymes. *Annu Rev*
660   *Plant Biol*. 62: 157–184.
661   Tran, H.T., Nimick, M., Uhrig, R.G., Templeton, G., Morrice, N., Gourlay, R., et al. (2012) Arabidopsis thaliana
662   histone deacetylase 14 (HDA14) is an α-tubulin deacetylase that associates with PP2A and enriches in the
663   microtubule fraction with the putative histone acetyltransferase ELP3. *Plant J*. 71: 263–272.
664   Tyanova, S., Temu, T., Sinitcyn, P., Carlson, A., Hein, M.Y., Geiger, T., et al. (2016) The Perseus computational
665   platform for comprehensive analysis of (prote)omics data. *Nat Methods*. 13: 731–740.
666   Uga, Y., Sugimoto, K., Ogawa, S., Rane, J., Ishitani, M., Hara, N., et al. (2013) Control of root system architecture
667   by DEEPER ROOTING 1 increases rice yield under drought conditions. *Nat Genet*. 45: 1097–1102.
668   Uhrig, R.G., Schläpfer, P., Roschitzki, B., Hirsch-Hoffmann, M., and Gruissem, W. (2019) Diurnal changes in
669   concerted plant protein phosphorylation and acetylation in Arabidopsis organs and seedlings. *Plant J*. 99:
670   176–194.
671   Wasson, A.P., Richards, R.A., Chatrath, R., Misra, S.C., Prasad, S.V.S., Rebetzke, G.J., et al. (2012) Traits and
672   selection strategies to improve root systems and water uptake in water-limited wheat crops. *J Exp Bot*. 63:
673   3485–3498.
674   Weinert, B.T., Wagner, S.A., Horn, H., Henriksen, P., Liu, W.R., Olsen, J. V., et al. (2011) Proteome-wide
675   mapping of the Drosophila acetylome demonstrates a high degree of conservation of lysine acetylation. *Sci*
676   *Signal*. 4: 1–12.
Willems, P., Horne, A., Van Parys, T., Goormachtig, S., De Smet, I., Botzki, A., et al. (2019) The Plant PTM

Viewer, a central resource for exploring plant protein modifications. *Plant J.* 99: 752–762.
Wu, X., Gong, F., Cao, D., Hu, X., and Wang, W. (2016) Advances in crop proteomics: PTMs of proteins under
abiotic stress. *Proteomics.* 16: 847–865.
Wu, X., Oh, M.H., Schwarz, E.M., Larue, C.T., Sivaguru, M., Imai, B.S., et al. (2011) Lysine acetylation is a
widespread protein modification for diverse proteins in Arabidopsis. *Plant Physiol.* 155: 1769–1778.
Xiong, L., Ishitani, M., Lee, H., and Zhu, J.K. (2001) The Arabidopsis LOS5/ABA3 locus encodes a molybdenum
cofactor sulfurase and modulates cold stress- and osmotic stress-responsive gene expression. *Plant Cell.*
13: 2063–2083.
Xu, P., Cai, X.T., Wang, Y., Xing, L., Chen, Q., and Xiang, C. Bin (2014) HDG11 upregulates cell-wall-loosening
protein genes to promote root elongation in Arabidopsis. *J Exp Bot.* 65: 4285–4295.
Xu, P., Zang, A., Chen, H., and Cai, W. (2016) The small G protein AtRAN1 regulates vegetative growth and
stress tolerance in arabidopsis thaliana. *PLoS One.* 11: 1–15.
Yi, D.G., and Huh, W.K. (2015) UDP-glucose pyrophosphorylase Ugp1 is involved in oxidative stress response
and long-term survival during stationary phase in Saccharomyces cerevisiae. *Biochem Biophys Res*
Commun. 467: 657–663.
Yin, H., Li, M., Li, D., Khan, S.A., Hepworth, S.R., and Wang, S.M. (2019) Transcriptome analysis reveals
regulatory framework for salt and osmotic tolerance in a succulent xerophyte. *BMC Plant Biol.* 19: 1–15.
Zhang, H., Han, B., Wang, T., Chen, S., Li, H., Zhang, Y., et al. (2012) Mechanisms of plant salt response:
Insights from proteomics. *J Proteome Res.* 11: 49–67.
Zhao, S., Xu, W., Jiang, W., Yu, W., Lin, Y., Zhang, T., et al. (2010) Regulation of Cellular Metabolism by Protein
Lysine Acetylation. *Science (80-).* 1000–1004.
Zhu, J.K. (2016) Abiotic Stress Signaling and Responses in Plants. *Cell.* 167: 313–324.
700
701

 TABLES

704 Table 1: Summary results of proteomic and PTMomics analyses of Arabidopsis roots after 24 hours of salt and
705 osmotic stress.

Stress	Proteome Assessed	*Quantified		**Significantly Changing under Stress		% Sig PTM Quantified not changing in Proteome	% Sig PTM Quantified changing in Proteome
		Proteins	Peptides	Proteins	Peptides	Proteins	Proteins
300mM Mannitol (Osmotic)	Acetylproteome (Ac)	209	281	33	35	26 (79%)	0 (0%)
	Phosphoproteome (P)	1225	3564	81	86	21 (26%)	0 (0%)
	Proteome (TP)	3113	n/a	38	n/a	n/a	n/a
150mM NaCl (Salinity)	Acetylproteome (Ac)	209	281	55	59	43 (78%)	1 (1.8%)
	Phosphoproteome (P)	1225	3564	54	56	19 (35%)	0 (0%)
	Proteome (TP)	3113	n/a	71	n/a	n/a	n/a

707 *Quantified PTM (Ac & P) = Localization ≥ 0.75
708 *Quantified Proteome (TP) = min 2 peptides per protein
709 **Significantly changing under stress (Ac, P & TP) = *p-val* ≤ 0.05 and ≥ 1.5 fold-change in PTM abundance

711 Table 2: Enriched GO terms for significant protein abundance changes upon 24 hours of 150 mM salt (NaCl)
712 treatment

GO Term	GO ID	pvalue	BH pvalue
sulfur compound biosynthetic process	GO:0044272	2.56E-05	0.009996086447
sulfate reduction	GO:0019419	3.33E-05	0.009996086447
sulfur compound metabolic process	GO:0006790	5.79E-05	0.01302706954
adenylyl-sulfate reductase activity	GO:0009973	7.55E-05	0.0135910601
seed coat development	GO:0010214	1.26E-04	0.01736891571
S-glycoside metabolic process	GO:0016143	1.35E-04	0.01736891571
glycosinolate metabolic process	GO:0019757	1.80E-04	0.02021935016
sulfate assimilation	GO:0000103	4.64E-04	0.04180239927

Pathway Description	AGI	Pathway	BH pvalue
2-Oxocarboxylic acid metabolism	AT1G23310, AT1G65930, AT2G20610, AT2G31810, AT2G42790, AT3G19710, AT3G58610, AT4G13770, AT5G11520, AT5G14200, AT5G23010, AT5G23020	1210	0.001175924641
Glucosinolate biosynthesis	AT1G18590, AT2G20610, AT3G19710, AT4G13770, AT5G23010, AT5G23020	966	0.002252925854

Protein Domain Name	AGI	Domain	BH pvalue
Chaperonin TCP-1, conserved site	AT1G24510, AT3G02530, AT3G11830,AT3G20050, AT5G20890, AT5G26360	IPR002194	0.00000773058189
GroEL-like equatorial domain	AT1G24510, AT3G02530, AT3G11830, AT3G20050, AT5G20890, AT5G26360, AT5G56500	IPR027413	0.00001576681032
Chaperonin Cpn60/TCP-1 family	AT1G24510, AT3G02530, AT3G11830, AT3G20050, AT5G20890, AT5G26360, AT5G56500	IPR002423	0.00005815154064
GroEL-like apical domain	AT1G24510, AT3G02530, AT3G11830, AT3G20050, AT5G20890, AT5G26360, AT5G56500	IPR027409	0.00005815154064
Glycoside hydrolase superfamily	AT1G47600, AT1G78060, AT2G32810, AT3G09260, AT3G13750, AT3G18080, AT3G60130, AT4G19810, AT5G08380, AT5G11720, AT5G26000, AT5G49360, AT5G56870, AT5G63800, AT5G63810	IPR017853	0.001488229715
Thioredoxin-independent 5'-adenylylsulphate reductase	AT1G62180, AT4G04610, AT4G21990	IPR004508	0.003244018163
Phosphoadenosine phosphosulphate reductase	AT1G62180, AT4G04610, AT4G21990	IPR002500	0.009327460679
Alkaline-phosphatase-like, core domain	AT1G09780, AT3G03530, AT4G29690, AT4G29700	IPR017850	0.036840083

FIGURES

Figure 1:

Proteome and PTMome changes under osmotic or salt stress. (A) Experimental workflow (B-D) Venn diagram and Log$_2$ fold-change (FC) scatterplots for significantly changing (B) acetylome, (C) phosphoproteome and (D) proteome. (E) Subcellular localization distribution of significantly changing proteins (*pval* ≤ 0.05; FC ≥ 1.5).

Figure 2:

Comparison of transcript-level changes with corresponding protein abundance, phosphorylation, and acetylation changes upon 24 hours of osmotic stress. Transcript-level changes corresponding to significantly changing proteins were calculated from previously described microarray experiments (KIllan et al., 2007) and clustered by longitudinal shape. Significantly enriched GO terms (Biological process) for each cluster were obtained from geneontology.org.

Figure 3:

Comparison of transcript-level changes with corresponding protein abundance, phosphorylation, and acetylation changes upon 24 hours of salt stress. Transcript-level changes corresponding significantly changing proteins were calculated from previously described microarray experiments (Killan et al., 2007) and clustered by longitudinal shape. Significantly enriched GO terms (Biological process) for each cluster were obtained from geneontology.org.

Figure 4:

Association network of Arabidopsis root under 24 h 150 mM salt (NaCl). STRING database association network shown for proteins significantly changing in their total protein, phosphorylation, or acetylation status. The maximal log$_2$FC change ranges from -2.5 (Max log$_2$FC EN) to +2.5 (Max log$_2$FC ED). Edge thickness equates to confidence in association. Inner circle represents subcellular localization (SUBA 4; https://suba.live/): cytosol (red), mitochondria (purple), nucleus (blue), peroxisome (green), plasma membrane (beige), plastid (amber), vacuole (lavender), endoplasmic reticulum/extracellular/Golgi (grey). Outer ring represents protein status: total protein change (1st division), phosphorylation (2nd division), Acetylation (3rd division). Grey clusters depict proteins belonging to a common cellular process. Proteins with an edge score threshold of 0.5 or below were excluded from visualization.

Figure 5:

Association network of Arabidopsis root under 24 h 300 mM mannitol. STRING database association network shown for proteins significantly changing in their total protein, phosphorylation, or acetylation status. The maximal log$_2$FC change ranges from -2.5 (Max log$_2$FC EN) to +2.5 (Max log$_2$FC ED). Edge thickness equates to confidence in association. Inner circle represents subcellular localization (SUBA 4; https://suba.live/): cytosol (red), mitochondria (purple), nucleus (blue), peroxisome (green), plasma membrane (beige), plastid (amber), vacuole (lavender), endoplasmic reticulum/extracellular/Golgi (grey). Outer ring represents protein status: total protein change (1st division), phosphorylation (2nd division), Acetylation (3rd division). Grey clusters depict proteins belonging to a common cellular process. Proteins with an edge score threshold of 0.5 or below were excluded from visualization.

Figure 6:

Root phenotypes of *aro2-1* and wild-type Arabidopsis under osmotic and salt stress. Root length (cm) of WT and *aro2-1* was measured every day for 11 days. (*:Unpaired t-test *pval* ≤ 0.05 with Holms-Bonferroni correction).

SUPPLEMENTAL INFORMATION

Supplemental Table 1: Results of proteome and PTMome analysis, including proteins with significantly changing protein abundance, phosphorylation and acetylation.

Supplemental Table 2: Gene Ontology enrichment for proteins with significantly changing abundance upon osmotic and salt stress.

Supplemental Table 3: KEGG pathway and protein domain analysis

Supplemental Table 4: Transcript data clustering analysis

A.

0.5x MS media (21 days) → Change Media →
0.5x MS Media +300mM Mannitol
Control
+150mM NaCl (24 h)
Harvest Roots & Freeze in liq'd N₂ → -80°C →

Tissue Extraction — Grind under liq'd N₂
FASP Digestion — Trypsin
Phosphopeptide Enrichment — TiO₂
Acetylpeptide Enrichment — Anti-AcetylLysine IgG
Mass Spectrometry — LC-MS/MS

B. Acetylation

Venn diagram: Mannitol 6, overlap 27, Salt 28

Acetylation scatter plot: Salt Log₂FC vs Mannitol Log₂FC. Labeled points: CRB, UGP1, NAT.

C. Phosphorylation

Venn diagram: Mannitol 52, overlap 29, Salt 25

Phosphorylation scatter plot: Salt Log₂FC vs Mannitol Log₂FC. Labeled points: BZIP16, ABF3, RSZ32, FH2.

D. Protein Abundance

Venn diagram: Mannitol 27, overlap 10, Salt 61

Protein Abundance scatter plot: Salt Log₂FC vs Mannitol Log₂FC. Labeled points: PEX11D, AGAL1, ARO2.

E. Bar chart of Percent (%) by localization for Mannitol and Salt: nucleus, cytosol, plastid, mitochondrion, plasma membrane, extracellular / golgi / ER, peroxisome, vacuole.

Gene Ontology (Biological Process)	FDR
Response to cadmium ion	5.58E-02
Glucosinolate biosynthetic process	4.75E-02
Response to oxygen containing compound	4.36E-02
Pentose phosphate shunt	3.76E-02
Actin-filament severing	3.61E-02
Ribonucleoprotein complex biogenesis	3.37E-02
Cellular component organisation	2.07E-02
Glycolytic process	2.00E-02
Gluconeogenesis	1.89E-02

Gene Ontology (Biological Process)	FDR
Sulfate assimilation	5.48E-04
Sulfate reduction	4.28E-04