1

Coupling between ribotypic and phenotypic traits of protists across life-cycle stages and

temperatures

Songbao Zou[1, 2]**, Rao Fu**[1, 2, 3]**, Qianqian Zhang**[1, 2]**, Eleni Gentekaki**[4]**, Jun Gong**[5, 6*]

[1] *Yantai Institute of Coastal Zone Research, Chinese Academy of Sciences, Yantai, China,*

[2] *University of Chinese Academy of Sciences, Beijing, China,*

[3] *Shandong Institute of Sericulture, Shandong Academy of Agricultural Sciences, Yantai, China,*

[4] *School of Science, Mae Fah Luang University, Chiang Rai, 57100, Thailand,*

[5] *School of Marine Sciences, Sun Yat-Sen University (Zhuhai Campus), Zhuhai, China,*

[6] *Southern Marine Science and Engineering Guangdong Laboratory (Zhuhai), Zhuhai, China*

*Correspondence:

Jun Gong, School of Marine Sciences, Sun Yat-sen University, Zhuhai Campus, Tangjiawan, Zhuhai

519082, China, gongj27@mail.sysu.edu.cn

Running title: Coupling ribotypic to phenotypic traits in protists

ABSTRACT

Relationships between ribotypic and phenotypic traits of protists across life-cycle stages remain largely unknown. Herein, we assessed the effect of life-cycle stage (lag, log, plateau, cyst) and temperature on quantity and polymorphisms of ribosomal (r)RNA gene (rDNA) and rRNA transcripts by using single cells of ciliate species. *Colpoda inflata* and *C. steinii* demonstrated allometric relationships between 18S rDNA copy number per cell (CNPC), cell volume (CV), and macronuclear volume across all life-cycle stages. Integrating previously reported data of *Euplotes vannus* and *Strombidium sulcatum* indicated taxon-dependent rDNA CNPC–CV functions. Ciliate and prokaryote data analysis revealed that the rRNA CNPC followed a unified power-law function, only if the rRNA-deficient resting cysts were not considered. Hence, a theoretical framework was proposed to estimate quantity of resting cysts of a protistan population. Using rDNA CNPC was a better predictor of growth rate at a given temperature than rRNA CNPC and CV, suggesting replication of redundant rDNA operons is a key factor that slows cell division. Single-cell high throughput sequencing revealed hundreds to thousands of rDNA and rRNA variants. The number of rDNA variants corresponded to the amount of cellular rDNA. Despite intra-individual divergence reaching up to 9%, operational taxonomic units (OTUs) clustered in a species-specific manner and independently of life-cycle stage or temperature. This indicates limited influence of said divergence on distributional pattern of OTU richness among samples.

IMPORTANCE

Based on phenotypic traits, traditional surveys usually characterize organismal richness, abundance, biomass, and growth potential to describe diversity, organization and function of protistan populations and communities. The ribosomal RNA gene (rDNA) and its transcripts have been widely used as molecular markers in ecological studies of protists. Nevertheless, the manner in which these molecules relate to cellular (organismal) and physiological traits remains poorly understood, which could lead to misinterpretations. The current research supports the idea that cellular ribosomal RNA transcript quantity reflects cell volume (biomass) of protists better than rDNA quantity, across multiple life-cycle stages except for resting cysts. We demonstrate that

quantity of resting cysts and maximum growth rate of a population can be theoretically estimated

using ribotypic trait-based formulas. Assessment of rDNA and rRNA sequence polymorphisms at

single-cell level indicates that intra-individual divergence does not affect recognizing variational

patterns of protistan diversity across time and space.

KEYWORDS: Cell size; Copy number variation; Growth rate; Life-cycle; Ribosome

heterogeneity.

INTRODUCTION

Protists are single-celled eukaryotes of astounding morphological and functional diversity inhabiting virtually every life-harboring niche on this planet (Caron et al., 2009). These microorganisms have complex life cycles comprising stages of distinct phenotypic traits (e.g., morphology, cell size and growth rate). Several of these traits undergo changes in response to fluctuating environmental variables (Corliss, 2001; Padilla and Savedo, 2013; Mulot et al., 2017). Cyst formation is a widespread trait among protists and is commonly induced under stressful conditions. Protists can form different types of cysts (e.g., resting, division and unstable cysts), which are typically nonmotile and dormant. Cysts play important roles in the ecology of protists, as they are resistant to unfavorable conditions, e.g., food depletion, overpopulation, drying, temperature and barometric pressure extremes, ultraviolet irradiation, and changing chemical factors (McQuoid and Hobson, 1996; Corliss, 2001; Persson, 2001; Ross and Hallock, 2016). Cyst cells comprise an essential life cycle stage of many protists promoting dispersal to new locations, thus increasing their population in a short period of time (Corliss and Esser, 1974; Corliss, 2001). Protistan diversity, community composition, rare-to-abundant transition, seasonality, biogeographical patterns, dynamic responses to perturbations, as well as, ecological and biogeochemical functions in soil, planktonic and benthic systems are all affected by dormancy of cysts (Corliss, 2001; Persson, 2001; Fistarol et al., 2004; Anderson and Rengefors, 2006), highlighting its functional and ecological significance in natural systems. However, cysts are highly reduced in size and lack most diagnostic morphological features in comparison with vegetative cells. As such, protistan cysts are usually difficult to morphologically identify and enumerate in environmental samples (Corliss, 2001).

The difficulties in morphological identification of protistan species in natural ecosystems have been largely circumvented by metabarcoding analyses using ribosomal (r)RNA genes (rDNA) and/or rRNA transcripts as molecular markers. Nevertheless, recent studies have shown that the number of rDNA copies in protists is highly variable raising concerns in regards to its effect on estimates of species richness and evenness in a community (reviewed in Fu and Gong, 2017). The majority of these investigations have not taken into account variability of rDNA copy number (CN)

of various life stages, particularly in the cystic forms. So far, cystic rDNA CN data are only

available for a limited number of protistan species, e.g., bloom-forming dinoflagellates (e.g., Erdner

et al., 2010) and medically important amoebae (e.g., Rivière et al., 2006). Hence, fundamental

knowledge on cystic rDNA and rRNA transcripts and their effect on taxon richness and evenness is

still lacking. Moreover, CN of rRNA (a proxy of ribosome number) in a cyst has seldom been

quantified, probably due to the common assumption that rDNA is sparsely or not at all transcribed

in the dormant stages. Nevertheless, it has been noted that, apart from phenotypic changes

(reduction of cell volume and nuclear size), encystment also induces substantial genetic alterations,

including loss of genomic DNA and altered nuclear DNA methylation patterns (Gutiérrez et al.,

1998). Previous studies have demonstrated variation in rDNA CN in a foraminiferan species across

life cycle stages (Parfrey et al., 2012), and have provided support for scaling relationships among

rDNA, rRNA CNs per cell, growth rate, and cell volume (CV) during the exponential growth stage

of ciliate species (Fu and Gong, 2017). Collectively considering these data, it is reasonable to

expect a substantial decrease of the CNs of both cellular rDNA and rRNA during encystment due to

shrinkage of the whole cell. However, the following points have yet to be empirically investigated:

(1) the extent to which the ribotypic CNs and their ratios are reduced in cysts; (2) the quantitative

relation of these ribotypic traits to phenotypic traits across different growth stages (e.g., lag, log,

plateau and cystic phases); and (3) whether previous models of ribotypic CN-CV scaling

relationships hold for cysts.

The available data suggest that the rDNA operons of diverse protistan groups have a high

degree of intra-individual polymorphisms (Pillet et al., 2012; Gong et al., 2013; Decelle et al., 2014;

Weber and Pawlowski, 2014; Kudryavtsev and Gladkikh, 2017; Wang et al., 2017), but the

variability of protistan rDNA polymorphisms has rarely been linked to life cycle stages and/or to

environmental conditions. Hence, if the cellular rDNA CN oscillates during the life cycle (Parfrey et

al., 2012), it is conceivable that some rDNA variants may be differentially eliminated or amplified,

resulting in altered ribotypic diversity across developmental stages and variable environmental (e.g.,

temperature) conditions. More importantly, the intracellular rRNA landscape could be further

altered, resulting in ribosomal structural heterogeneity, due to selective transcription of specific

rDNA operons, as has been previously shown for several prokaryotes and an apicomplexan protist (e.g., Gunderson et al., 1987; López-López et al., 2007; Genuth and Barna, 2018; Kurylo et al., 2018). However, the dynamics of both rDNA operons and rRNA molecules (ribosomes) within a protistan cell at different life cycle stages or under varying conditions have been little explored.

In this study, we used two ubiquitously distributed soil ciliate species of *Colpoda* to investigate both rDNA and rRNA CNs across lag, log (exponential), plateau (stationary), unstable and resting cyst stages. *Colpoda* has been used as a model for the study of cyst-related biology and ecology for a long time (e.g., Stout, 1955; Gutiérrez et al., 1998; Müller et al., 2010). The two species were reared at two temperatures (18°C and 28°C) and their phenotypic features (cell size, macronuclear size, nucleocytoplasmic ratio) were determined. The environmental and growth-stage effects on intra-individual polymorphisms of ribotypic sequences were explored by comparing the distributional patterns and nucleotide contents of rDNA and rRNA variants in actively growing cells vs. resting cysts at both temperatures. We also extended previous work on the marine ciliates *Euplotes vannus* and *Strombidium sulcatum*, which were reared at three different temperatures by analyzing rDNA and rRNA sequence diversity in actively growing cells (Fu and Gong, 2017). Single ciliate cells were used for quantitative PCR and high throughput sequencing of rDNA and rRNA (cDNA). Our main expectations were that: (1) in line with reduction of cell and macronuclear size, the quantity of both rDNA operons and rRNA transcripts would be highly reduced in the cystic forms; (2) the scaling relationships between ribotypic CN and CV would generally hold across life cycle stages; and (3) the intra-individual polymorphic traits of both rDNA operons and rRNA molecules (ribosome composition) would be life stage- and temperature-dependent.

MATERIALS AND METHODS

Source organisms, cultivation and treatments

Two cyst-forming, soil ciliate species, *Colpoda steinii* and *C. inflata*, were collected from a coastal apple orchard in Yantai, Shandong, China. The protists were cultivated and isolated using the non-flooded Petri dish method (Omar et al., 2017). *Euplotes vannus* and *Strombidium sulcatum* were originally isolated from the coastal waters off Qingdao (Shandong, China), and have been

maintained in the laboratory for a long time. For the purpose of this study, cultures of these four

species were initiated from a single cell supplied with bacterial food.

Cultivation of these ciliates and monitoring of their abundances were as previously described

(Fu and Gong, 2017). In brief, ten cells of each species were picked with a micropipette, washed

several times to remove other microorganisms, and then inoculated into a tissue-culture flask with

50 ml water. All ciliates were fed with heat-killed *Escherichia coli* (at 70 ℃ for 10 min) cells, which

were cultured in LB-medium, pelleted and resuspended in ~20 ml sterilized water. The *Colpoda*

species grew well and underwent distinct phases (lag, log, and plateau) during an incubation period

of ten days at 18 ℃ and 28 ℃. Following the plateau phase, both species consistently formed resting

cysts, probably due to overpopulation and food depletion. Phase-specific cells (including the resting

cysts) were individually isolated to determine both phenotypic and ribotypic traits at the single-cell

level. The two marine species *E. vannus* and *S. sulcatum* were reared at 16 ℃, 21 ℃ and 25 ℃. For

these two species, only cells in log phase were investigated. In order to explore possible footprint

effect of the higher temperature on ribotypic traits, we set an additional "touchdown" treatment

(designated with "*") as previously described (Fu and Gong, 2017): briefly, a few individuals of the

treatments maintained at 25 ℃ for 30 d were transferred and re-cultured at 16 ℃ for another month.

All treatments were done in triplicate.

Many ciliate species become thin-walled, partially dehydrated, and division inhibited when

temperature shifts beyond certain limitations in a short time period, thus form unstable cysts (Stout,

1955). These unstable cysts are generally larger than the resting cysts, and undergo complete ciliary

dedifferentiation returning to vegetative cells in a few hours. We tried obtaining unstable cysts of

Colpoda by chilling cells for various periods of time. Tubes containing actively growing vegetative

cells were transferred into three separate conditions: ice-water mixture (0 ℃) and chemostats at 4 ℃

and 10 ℃. Twenty μL of culture solution were used to identify and count vegetative cells and

unstable cysts, all of which were subsequently observed using a stereomicroscope (XLT-400,

Guiguang, China) at 45× magnification. The dynamics of the unstable cysts and vegetative cells

were monitored every 0.5 to 2 h to determine the optimal time period for achieving the highest yield

of unstable cysts. All chilling treatments were performed in triplicate. Since unstable cysts reflect

high phenotypic plasticity, relevant phenotypic and ribotypic traits of this temporary stage were also

determined and analyzed.

Determination of growth rate, cell volume and macronuclear size

The cell abundances were determined once a day (Fig. 1A). Growth rates of the two *Colpoda*

species were calculated as the slope of *ln* individual versus time during the logarithmic phase. At

least 18 cells from each growth stage were randomly selected and fixed with Lugol's solution (final

concentration 2%). This fixation approach is identical to the one we previously used for

characterizing cell sizes of *E. vannus* and *S. sulcatum*, and allows direct comparisons of relevant

data between studies. The cell width and length were individually measured at 400x magnification

using a compound microscope (Olympus BX51, Tokyo, Japan). Fixed vegetative cells of both

Colpoda species were ellipsoid, with cell thickness approximately equal to cell width. Cell volume

(CV) was calculated according to the formula:

$CV = 4/3 * \pi * (length/2) * (width/2) * (thickness/2)$.

Both resting and unstable cysts were spheroid, thus their length, width and thickness were

considered equal.

Macronuclear and cell sizes of two *Colpoda* species were determined based on specimens fixed

with Bouin's solution and stained using protargol impregnation (Wilbert, 1975). This approach

reveals morphological features of ciliates (e.g., nuclear size, infraciliature) in detail, thus has been

commonly used for their identification and taxonomy (Warren et al., 2017). Nevertheless, the

fixative may alter the volume of the cells compared with the Lugol's fixation. To circumvent this

issue, volumes of both protargol impregnated specimens and those fixed with Lugol's were

determined. Measurements of length and width of the cells and their macronuclei were carried out

using a compound microscope (Olympus BX51) at a magnification of 1000X. The volumes of oval

to ellipsoidal cells and macronuclei were calculated as above described.

Cell lysis, DNA extraction and cDNA synthesis

Cell lysis, DNA extraction and RNA transcription of single-cells followed Fu and Gong (2017).

In brief, ciliate cells/cysts were picked and washed with autoclaved double distilled water. A single cell with a minimum amount of water was then transferred with a micropipette into the inner wall surface of a nuclease-free PCR tube. This process was carried out for more than 18 cells per stage. Before adding lysis solution, cell presence was verified using a stereoscope. Only tubes with intact cells were kept. Tubes containing broken cells were discarded to minimize RNA degradation. Cell lysis and nucleic acid extraction were performed for all 18 replicates on an ultra-clean bench pre-treated with RNaseZap (Ambion). Each cell was suspended in 1 μL of distilled water, with addition of 2 μL of mixed solution containing 1.9 μL 0.2% Triton-X-100 (Solarbio, Beijing, China) and 0.1 μL of recombinant RNase inhibitor (Takara, Biomedicals, Japan). The tubes were gently flicked and centrifuged briefly, then incubated at room temperature for 10 min. Nuclease-free PCR tubes, microtubes and micropippette tips (Axygen Scientific, CA, USA) were used for all manipulations. One μL of the cell suspension was used to extract genomic DNA, while 2 μL were used for RNA reverse transcription. Genomic DNA extraction was performed with a REDExtract-N-Amp Tissue PCR Kit (Sigma, St. Louis, MO), according to the manufacturer's instructions, except the suggested volume of each reagent was modified to 1/50, as previously described (Gong et al., 2014).

The RNA reverse transcription of single cells was as previously described (Fu and Gong, 2017). The remaining 2 μL of cell suspension were mixed with 1 μL random hexamers B0043 (10 μM, Sangon, China) and 1 μL of dNTP mix (10 μM). After incubation at 72 ℃ for 3 min the tube was immediately transferred onto ice. RNA was reverse transcribed using a SuperScript III Reverse Transcriptase Kit (Invitrogen, CA, USA). The total volume used for PCR was 12.42 μL, which was made up of 4 μL rRNA template, 3 μL 5× first strand buffer, 0.3 μL SuperScript III reverse transcriptase, 1 μL 0.1M DTT, 2 μL Betaine (Sigma–Aldrich), 0.12 μL 0.5M $MgCl_2$, 1 μL RNase inhibitor (Roche, Germany), and 1 μL DEPC-treated water. The following program was run on a thermal cycler (Biometra, Gottingen, Germany): an initial temperature of 50 ℃ for 90 min, followed by 10 cycles at 55 ℃ for 2 min, with a final step at 70 ℃ for 15 min. Both the DNA and synthesized cDNA were stored at -80 ℃ for subsequent assays.

Quantitative real-time PCR (qPCR)

For quantifying 18S rDNA and rRNA (cDNA) copy numbers in *Colpoda* cells, the newly

designed species-specific primers CsQ207f (5'-TAACCCTGGCAACAGGA-3') and CsQ459r

(5'-TGCAATCTCGCAACCCCA-3') were used in qPCR assays to amplify a 252-bp fragment of

the 18S rRNA gene of *C. steinii*. The eukaryote-specific primers EUK345f

(5'-AAGGAAGGCAGCAGGCG-3') and EUK499r (5'-CACCAGACTTGCCCTCYAAT-3') (Zhu

et al., 2005) were used to amplify a 149-bp fragment of *C. inflata*. Two standard curves were

constructed with the sequences of the two *Colpoda* species as previously described by Gong et al.

(2013). In brief, serial 10-fold dilutions (10^{-3} to 10^{-10}) were used to generate the standard curves.

The qPCR reaction was performed using SYBR Premix Ex Taq™ II (Takara, Dalian, China). The

reaction solution (20 μL) contained 0.4 μL of 10 μM primers, 0.4 μL ROX Reference Dye II, 10 μL

Premix Ex Taq II, 7.8 μL milli-Q water and 1 μL template. All the qPCR reactions were performed

on a 7500 Fast Real-Time PCR System (Applied Biosystems) with the following program: 95 ℃ for

7 min, followed by 45 cycles of 95 ℃ for 30 s, 55 ℃ (60 ℃ for *C. inflata*) for 1 min, and 72 ℃ for 1

min (77 ℃ for 25 s for *C. inflata*). The data were collected at 72 ℃ and 77 ℃ for *C. steinii* and *C.*

inflata, respectively; all reactions ended with a melt curve stage from 60 ℃ to 95 ℃ (gradually

increasing by 0.3 ℃). Standard curves were generated using $\log_{10}$ number of copies vs. the threshold

cycle (Ct). The linear correlation coefficient (R^2) ranged from 0.997 to 1.000. The amplification

efficiency ranged from 99% to 103%. As the samples of rRNA also contained genomic rDNA, the

rRNA (cDNA) copy numbers were calculated by subtracting the rDNA CNs from the sum of rDNA

and cDNA copy numbers.

248   Single-cell high throughput sequencing

Genomic DNA and synthesized cDNA from single cells of each of the four ciliate species was

used for high throughput sequencing. These included cells of *Colpoda steinii* and *C. inflata* at two

typical stages (i.e., vegetative cells in log-phase and resting cysts) and under two temperatures

(18 ℃ and 28 ℃), plus *Euplotes vannus* and *Strombidium sulcatum* during log phase grown at four

temperature treatments (i.e., 16 ℃, 21 ℃, 25 ℃ and 16* ℃; Fu and Gong, 2017). In the samples for

cDNA sequencing, genomic DNA was completely digested at 37 ℃ for 1 h using a TURBO

DNA-free Reagent Kit (Invitrogen), in which the digestion mixture consisted of 1 μl cell lysate, 1

μl 10×TURBO DNase buffer, 1 μl TURBO DNase (2 units μl^{-1}), and 7 μl nuclease-free water.

The V4 hypervariable regions of 18S rDNA and cDNA in *C. steinii* and *C. inflata* were

amplified with the eukaryotic specific primers TAReuk454FWD1

(5'-CCAGCASCYGCGGTAATTCC-3'), TAReukREV3 (5'-ACTTTCGTTCTTGATYRA-3')

(Stoeck et al., 2010). The V1-V3 regions of 18S rDNA and cDNA in *E. vannus* and *S. sulcatum*

were amplified with primers Euk82F (5'-GAADCTGYGAAYGGCTC-3') and Euk516R

(5'-ACCAGACTTGCCCTCC-3') (Díez et al., 2001; Dawson and Pace, 2002). A 6-bp barcode was

added to the sequences of the forward and reverse primers aimed to identifying each sample. The

PCR reaction solution (30 μL) contained 10 ng of DNA or cDNA, 0.2 μM of each primer, and 15

μL of 2× Phusion High-Fidelity PCR Master Mix (New England Biolabs). All PCR amplification

reactions were carried out on a T100 Thermal Cycle (Bio-Rad) with the following program: an

initial pre-denaturation at 98 ℃ for 1 min, followed by 30 cycles of denaturation at 98 ℃ for 10 s,

annealing at 50 ℃ (56 ℃ for two marine ciliates) for 30 s, and elongation at 72 ℃ for 30 s, with a

final extension step of 72 ℃ for 5 min. The libraries were prepared using a NEB Next Ultra DNA

Library Prep kit. Paired-end 250 bp and 300 bp sequencing was executed on HiSeq and MiSeq

platforms (Illumnina, USA) for the soil and marine species, respectively. A total of 80 single-cell

rDNA and rRNA pools of the two *Colpoda* species were sequenced. For each *Colpoda*, 20 cells

were used to sequence DNA and another 20 for cDNA. There were five biological replicates for

each treatment. Four samples failed to be amplified, resulting in 39 samples of logarithmic phase

and 37 samples of resting cysts. A total of 48 single cells of *E. vannus* and *S. sulcatum* were

sequenced. For each species, 24 cells were used to sequence DNA and 24 for RNA pools. Each

treatment was run in triplicate.

Data processing and analysis

A total of 8,478,407 reads were generated for rDNA and cDNA of the four ciliate species

(Table S1). Data processing and analysis was done using a combination of Prinseq-lite v.0.20.4

(Schmieder and Edwards, 2011), QIIME v.1.8.0 (Caporaso et al., 2010) and MOTHUR v.1.34.4

(Schloss et al., 2009). Quality control was conducted and high-quality reads were retained according to the following criteria: length of the reads including primers and barcode sequences ≥ 380 and ≥ 400 bp for the soil and marine species, respectively; quality score > 20; no ambiguous bases; no primer sequence mismatches; and homopolymers ≤ 6. Reads of replicates of the same species were merged in a single file. USEARCH v.61 was used to screen for putative chimeras (Edgar, 2010) based on Silva database (release 132). Primer sequences were trimmed off and amplicons were assigned to their closest hits in the Silva database using UCLUST v.1.2.2 (Edgar, 2010) with a minimum 90% identity threshold. Based on taxonomic assignment, the reads assigned to each species were extracted by MOTHUR. Singletons (the unique variants with a single read across all samples of the same species) were excluded prior to further analysis. For treatments of each species, the filtered reads were clustered using UCLUST at various similarity thresholds ranging from 90% to 100%, and ribotypic polymorphisms between rDNA and rRNA sequences and across treatments were assessed.

Characterizations of single-cell ribotype sequence polymorphisms

Nucleotide polymorphisms were identified by aligning all rDNA and rRNA (cDNA) reads against the dominant haplotype of each species, which was implemented using Burrows-Wheeler Aligner v.0.7.17 with default parameters (Li and Durbin, 2009). The alignments were converted into BAM format, sorted and indexed using SAMtools v.0.1.19 (Li et al., 2009). The ribotype nucleotide polymorphisms of each species were then called using the *mpileup* function and summarized using an in-house *perl* script (printBasesBwa2.pl).

Statistical analysis

Student's *t*-test (two-tailed) was performed to test the significance ($\alpha = 0.05$) of the growth phase-wise differences in phenotypic (i.e., cell volume, macronuclear volume, cell to macronuclear volume ratio) and ribotypic traits (per-cell rDNA and rRNA CNs and concentrations, rDNA: rRNA CN ratio). The maximum growth rates of two *Colpoda* species at two different temperatures were also statistically compared using *t*-test. Pearson correlation and linear regression analyses were carried out to link these phenotypic traits with ribotypic traits or to explore the association between

ribotypic traits. All these analyses were executed using the statistical software SPSS v.13.0 (SPSS,

Chicago, IL).

Nucleotide sequence accession numbers

The reads from the high throughput sequencing of eukaryotic 18S rRNA genes and transcripts

are available under accession numbers SRR10589967 - SRR10590042 (*Colpoda steinii* and

Colpoda inflata) and SRR10590394 - SRR10590441 (*Euplotes vannus* and *Strombidium sulcatum*).

RESULTS

Cell volume, macronuclear volume, and nucleocytoplasmic ratio

The lag, logarithmic, plateau and resting cystic phase of *Colpoda steinii* and *C. inflata* were

distinguishable when cultured with replete bacterial prey at both 18 ℃ and 28 ℃ (Fig. 1A). At 28 ℃,

the two species grew approximately 2 and 6 times faster than at the 18 ℃ treatments, respectively.

At both temperatures, the smaller species (*C. steinii*) grew consistently faster than the larger species

(*C. inflata*): 7.2 ± 0.87 d^{-1} (mean $\pm$ SE) vs. 5.1 ± 0.09 d^{-1} at 28 ℃, and 3.6 ± 0.32 d^{-1} vs. 0.9 ± 0.32

d^{-1} at 18 ℃ (Fig. 1B). Abundant unstable cysts were successfully obtained by chilling actively

growing cells of *C. steinii* at 0 ℃ for 4 h and *C. inflata* at 4 ℃ for 10 h (Fig. S1).

The cell volume (CV) of both *Colpoda* species progressively reduced through lag, log, and

plateau to resting cyst phases at both temperatures (Fig. 1C, D; 2A). In the 18 ℃ treatment of *C.*

inflata, the largest CV (about 2.14×10^5 μm^3) was recorded for the lag phase cells. Upon entering

the exponential phase, the cells became smaller (on average 8.9×10^4 μm^3), and further shrank by

46% in plateau phase to about 4.8×10^4 μm^3. Relative to the log-phase cells, the resting cysts of this

species greatly reduced by 83% in size to 1.5×10^4 μm^3. *Colpoda inflata* cells were significantly

smaller in the 28 ℃ treatment (lag, 1.3×10^5 μm^3; log, 7.7×10^4 μm^3; plateau, 3.9×10^4 μm^3,

resting cysts, 7.5×10^3 μm^3) than in the 18 ℃ treatment ($P < 0.05$ in all cases; Fig. 2A). Similarly,

the small-sized *C. steinii* lost 85% and 69% of its CV during encystment relative to the log phase

cells in 18 ℃ and 28 ℃ treatments, respectively. At a given vegetative phase the cell volume was

always larger in the cultures reared at 18 ℃ than at 28 ℃ ($P < 0.05$), except for the resting cysts,

whose size (about 2.0×10^3 μm^3) was not significantly different between these two temperature

treatments (Fig. 2A).

The chilling-induced unstable cysts of both *Colpoda* species were generally larger than the

resting cysts, but smaller than the plateau phase cells (Fig. 2A). In the treatment of 18 ℃, the

volumes of unstable cysts of *C. steinii* and *C. inflata* were on average 5.3×10^3 and 0.21×10^5 μm^3,

respectively (Fig. 1C, 2A). A similar pattern of unstable cyst size was observed for the treatments of

28 ℃ (Fig. 2A). Assuming that the *C. steinii* and *C. inflata* cells were globular in body shape, the

equivalent spherical diameter (ESD) of the cells ranged from 13 to 29 μm and from 25 to 74 μm,

respectively, with cross-phase patterns similar to these of the cell volume (Fig. 2B).

Macronuclear volume (MV) exhibited a similar trend to the cell volume in both *Colpoda*

species, i.e., decreasing gradually through lag, log, and plateau phases to resting cyst, regardless of

cultivation temperatures (Fig. 2C). *Colpoda inflata* had consistently a larger MV than *C. steinii* at

any specific phase ($P < 0.05$). When the MV was relatively smaller (e.g., MV < 500 μm^3, or

macronuclear diameter < 10 μm), the shrink of MV of both species became consistently significant

in the 18 ℃ treatments relative to the 28 ℃ treatments. However, when the macronucleus was larger

(i.e., macronuclear diameter > 10 μm), the effect of temperature increase on MV was insignificant

($P > 0.05$; Fig. 2C).

The nucleocytoplasmic ratio of two *Colpoda* species ranged from approximately 1% to 8%, and

tended to progressively increase from lag, log, to plateau phases and resting cysts (Fig. 2D). The

log-phase cells of *C. steinii* had the lowest nucleocytoplasmic ratios (on average 2.92% to 2.61% at

18 ℃ and 28 ℃, respectively), which were about 1.13 to 1.32 times lower than these in lag and

plateau phases, respectively. Culturing at the higher temperature generally did not induce significant

decreases in the nucleocytoplasmic ratio in most growth phases of either species ($P > 0.05$), except

for resting cysts of *C. steinii*, which had a much higher nucleocytoplasmic ratio (on average 7.93%)

at 18 ℃ than that at 28 ℃ (3.82%). Compared with the small species, *Colpoda inflata* usually had

lower cytoplasmic volume ratios in all phases and at both temperatures (Fig. 2D). Regression

analysis showed that macronuclear volume scaled well with CV$^{0.83}$ (Fig. 2E); the nucleocytoplasmic

ratio fit well to a power-lower function of *ESD*$^{-0.522}$ (Fig. 2F):

$\lg (MV) = 0.83 \lg (CV) - 0.87$, or $MV = 0.136\ CV^{0.83}$ (1)

$R^2 = 0.90$, $P < 0.001$, $n = 733$;

$MV/CV = 0.136\ CV^{-0.17} = 0.151\ ESD^{-0.52}$ (2)

$CV = 5.14\ ND^{3.63}$ (3)

Where ND is the mean macronuclear diameter of *Colpoda* spp.

Variations in single-cell rDNA and rRNA copy numbers and concentrations

The per-cell rDNA and rRNA copy numbers varied significantly among growth phases

(ANOVA, $P < 0.001$), showing a consistently decreasing trend from the lag, log, and plateau phases

to the resting cyst stage, for both *Colpoda* species and at both temperatures. The resting cysts of

both *Colpoda* species lost 50% ~ 90% of rDNA and over 89% ~ 99.99% of rRNA copies relative to

the cells in log phase (Fig. 3A, B). In the 18 ℃ treatment, a lag-phase cell of *C. steinii* had on

average $(10.2 \pm 1.22) \times 10^3$ rDNA copies. The CN was about 2 and 4.6 times higher than that in the

log [$(5.4 \pm 0.7) \times 10^3$ copies] and plateau phases [$(2.2 \pm 0.3) \times 10^3$ copies], respectively. The rDNA

amount reduced greatly in the resting cysts of this species, with only about 1000 copies per cyst

(Fig. 3A). Although the unstable cysts of *C. steinii* were smaller than the plateau-phase cells, the

former had a similar copy number of rDNA [$(2.0 \pm 0.2) \times 10^3$] as the latter. Compared with the

lower temperature treatments, *C. steinii* cultured at 28 ℃ had lower per-cell rDNA copy numbers,

particularly for the cells at log and plateau phases ($P < 0.05$); however, the number of rDNA copies

in the resting cysts of this species did not change significantly at the warmer condition ($P = 0.69$).

The large-sized *Colpoda* species had many more rDNA copies than the small-sized congener at

any life cycle stages (Fig. 3B). A single cell of *C. inflata* reared at 18 ℃ had [4.48×10^4], [$1.98 \times$

10^4], and [1.33×10^4] rDNA copies at the lag, log, and plateau phases, respectively. Phase-specific

comparisons of the cells between these two temperature treatments showed that the effect of

warming was not significant for this species ($P > 0.05$). The resting and unstable cysts of *C. inflata*

contained about 2500 and 8100 rDNA copies, respectively, which were about 3.3 and 4.2 times than

those of *C. steinii* (Fig. 3B).

The cellular rRNA copies were far more abundant, and varied more dramatically across growth

stages, than the rDNA copies in both *Colpoda* spp. (Fig. 3A, B). In the 18 ℃ treatments, a single cell of *C. steinii* had about 3.0×10^7, 1.0×10^7, 5.7×10^6, and 3.4×10^4 rRNA copies at the lag, log, plateau and resting cyst stages, respectively (Fig. 3A). At the same temperature, the rRNA molecules were much more abundant in *C. inflata*, with about 1.7×10^8, 1.3×10^8, 3.2×10^7 and 3.1×10^4 across these phases (Fig. 3B). Although smaller than the plateau-phase cells, the unstable cysts had relatively similar (in *C. steinii*) or even higher number of rRNA copies (in *C. inflata*) (Fig. 3A, B). Comparing the two temperature treatments, warming always led to a significant decrease in cellular rRNA amount of both species and at all the vegetative stages plus the unstable cysts ($P < 0.05$). Nevertheless, there were no significant differences in the cellular rRNA abundance in resting cysts at either temperature (Fig. 3A, B). A cell at any vegetative growth phase of these two species had approximately 200 to 5000 times more rRNA copies than resting cysts (Fig. 3A, B).

When per-cell copy numbers of rDNA and rRNA were standardized to cell volume, the concentrations of these two ribotypes remained relatively stable across most growing phases and cystic stages (Fig. 3C). In both temperature treatments, the mean cellular rDNA concentrations ranged mostly between 0.4 and 0.8 copies μm^{-3} in *C. steinii*, and were slightly lower (from 0.2 to 0.3 copies μm^{-3}) in *C. inflata*, showing no significant differences across lag, log, plateau and resting cystic phases in either species ($P > 0.05$). The only exception lied in a significantly lower concentration (0.44 copies μm^{-3}) at the log phase (relative to the lag phase) of *C. steinii* in the warmer treatment (Fig. 3C); and there was a significant difference of concentrations (on average 0.20 vs. 0.35 copies μm^{-3}) in resting cysts of *C. steinii* between two temperature treatments. The culturing temperature prior to chilling affected the rDNA concentrations in subsequently formed unstable cysts of both species: the cellular rDNA concentrations in the unstable cysts of both species at 28 ℃ were twice these at 18 ℃ (Fig. 3D).

Generally speaking, the rRNA concentration varied within relative larger ranges in the vegetative cells (1000 - 2200 copies μm^{-3} in *C. steinii*, and 560 - 1500 copies μm^{-3} in *C. inflata*), but the difference at a given growth phase was mostly insignificant (Fig. 3C, D). Nevertheless, differing from the rDNA molecules that still sustained reasonably high cellular concentrations during formation of resting cysts via the plateau phase, the rRNA concentrations became significantly

lower in resting cysts of both *C. steinii* (27 copies μm^{-3} at 18 ℃, and 380 copies μm^{-3} at 28 ℃; Fig.

3C) and *C. inflata* (0.13 copies μm^{-3} at 18 ℃, and 2.1 copies μm^{-3} at 28 ℃; Fig. 3D). Culturing

temperature seemed to not affect rRNA concentrations in the vegetative cells most of the time (Fig.

3C, D). The unstable cysts of *C. steinii* (1120 copies μm^{-3} at 18 ℃, and 1895 copies μm^{-3} at 28 ℃)

had similar concentrations of rRNA with the vegetative cells, while *C. inflata* (5336 copies μm^{-3} at

18 ℃, and 3734 copies μm^{-3} at 28 ℃) had many more rRNA molecules than its vegetative forms.

Furthermore, rRNA concentrations in unstable cysts were significantly different between these two

treatments (Fig. 3C, D).

There were minor differences between two *Colpoda* species in terms of the patterns of rRNA to

rDNA CN ratio across growth phases and between two temperature treatments (Fig. 3E, F). The

rRNA: rDNA CN ratios in *C. steinii* mostly ranged from 2000 to 4000, without statistical

differences among the lag, log, plateau and unstable cystic stages ($P > 0.05$; Fig. 3E). Nevertheless,

the rRNA: rDNA ratios in *C. inflata* decreased from 10000 to 2000 when the population growth at

18 ℃ entered into plateau phase; and this species also exhibited a significantly high rRNA: rDNA

ratio in unstable cysts induced from the actively growing cells in the 18 ℃ treatment (Fig. 3F).

Furthermore, while the rRNA: rDNA ratios in log-phase cells, unstable, and resting cysts of *C.*

inflata significantly dropped at the higher culturing temperature (Fig. 3F), the same was not

observed in *C. steinii* (Fig. 3E). Nevertheless, a trait shared by both species was that the rRNA:

rDNA ratios (150 – 400 in *C. steinii*, and 2 – 16 in *C. inflata*) were much lower in resting cysts than

in the cells at all other phases (Fig. 3E, F).

Linking ribotypic with phenotypic traits

Based on all CN data of vegetative and cystic cells of these two *Colpoda* spp., correlation and

linear regression analyses showed that both ribotypic CN traits (rDNA and rRNA copy number per

cell, $rDNA_{CNPC}$ and $rRNA_{CNPC}$) were positively and significantly related to CV ($P < 0.01$). Ribotype

CNs could well fit to linear regression curves (Fig. 4A):

$\lg(rDNA_{CNPC}) = 0.76 \lg(CV) + 0.61$, or $rDNA_{CNPC} = 4.07 \, CV^{0.76}$ (4)

$R^2 = 0.91, P < 0.001, n = 20$;

Assuming the cell is globular in shape, with an equivalent spherical diameter *(ESD)*, then:

$rDNA_{CNPC} = 2.49\ ESD^{2.28}$ (5)

$\lg (N \cdot rDNA_{CNPC}) = \lg N + 2.28 \lg ESD + 0.396$ (6)

Where N is the number of cells of identical size; $N \cdot rDNA_{CNPC}$ is the total rDNA copy number of

the population that can be experimentally determined (for example, by using qPCR). The scaling

relationships and R^2 scores indicated that the $rDNA_{CNPC}$ of these two *Colpoda* species could be

accurately predicted using biovolume (equation 4) or *ESD* (equations 5 and 6) of cells, no matter

whether the cells were at lag, log, plateau, or in unstable or resting cystic forms. Furthermore,

equation (6) clearly indicates that the total rDNA CN of a population is determined not only by cell

abundance (*N*), but also by cell size (*ESD*).

As for rRNA, the correlation between $rRNA_{CNPC}$ and CV across all life cycle phases including

resting cysts (+RC) was significant. However, the R^2 score (37%) was only moderate (Fig. 4A):

$\lg (rRNA_{CNPC}) = 1.24 \lg (CV) + 1.69,\ R^2 = 0.37,\ P = 0.005,\ n = 20$ (7)

Apparently, the rRNA amount in resting cysts of *Colpoda* species was extremely lower and not

easily predicted when using cell size. Nevertheless, when the rRNA CN data of the resting cysts

(i.e., the four blue full diamonds that are discrete from other points in Fig. 4A) were excluded from

the analysis (-RC), we obtained:

$\lg (rRNA_{CNPC}) = 0.87 \lg (CV) + 3.67,$ or $rRNA_{CNPC} = 4677\ CV^{0.87}$ (8)

$R^2 = 0.84,\ P < 0.001,\ n = 16;$

$rRNA_{CNPC} = 2664\ ESD^{2.61}$ (9)

$\lg (N \cdot rRNA_{CNPC}) = \lg N + 2.61 \lg ESD + 3.43$ (10)

By removing resting cystic data, the re-modeled rRNA CN and cell size relationship became much

more robust ($R^2 = 0.84$). Therefore, based on a combination of equations (6) and (10), which do not

take resting cysts into account, we have:

$\lg (N \cdot rRNA_{CNPC}) - \lg (N \cdot rDNA_{CNPC}) = 0.33 \lg ESD + 3.034$

Or, $\lg (rRNA_{CNPC} / rDNA_{CNPC}) = 0.33 \lg ESD + 3.034$ (11)

Equation (11) indicates that, given that all cells in a population are non-resting cysts, the rRNA:

rDNA copy number ratio depends on a single factor, *ESD* or cell size.

Since the total rDNA pool ($rRNA_{total} = N \cdot rRNA_{CNPC}$) and the total rDNA pool ($rDNA_{total} =$

$N \cdot rDNA_{CNPC}$) can be experimentally quantified (e.g., qPCR), then:

$$\lg [(rRNA_{total} - rRNA_{rc}) / (rDNA_{total} - rDNA_{rc})] = 0.33 \lg ESD + 3.034, \tag{12}$$

Where $rRNA_{rc}$ and $rDNA_{rc}$ are the total cellular rRNA and rDNA CNs in the resting cysts existing in

the population. Assuming that the proportion of resting cysts in the rDNA pool is $cyst\%$, and their

fraction in the rRNA pool is negligible (about 0.02% to 0.5% relative to a vegetative cell of

Colpoda spp.), then:

$$\lg [(rRNA_{total})/(rDNA_{total} - rDNA_{total} \cdot cyst\%)] = 0.33 \lg ESD + 3.034 \tag{13}$$

Thus, the rDNA proportion of non-resting cells ($NRC\%$) in a population can be estimated as

follows:

$$NRC\% = 1 - cyst\% = (rRNA_{total}/rDNA_{total}) / (1081 \cdot ESD^{0.33}) \tag{14}$$

The correlations between MV of cells at all life cycle stages and ribotype CNs exhibited

patterns similar to the observed for CV (Fig. 4B):

$$\lg (rDNA_{CNPC}) = 0.90 \lg (MV) + 1.44 \text{ or } rDNA_{CNPC} = 27.54 \, MV^{0.9} \tag{15}$$

$R^2 = 0.93, P < 0.001, n = 16$;

$$\lg (rRNA_{CNPC}) = 1.70 \lg (MV) + 2.28 \text{ or } rRNA_{CNPC} = 190.55 \, MV^{1.7} \tag{16}$$

$R^2 = 0.50, P = 0.002, n = 16$;

The rRNA CN data did not fit the model very well. However, when the rRNA data of resting cysts

were not considered, the explanation of variance in rRNA CNPC could be greatly improved:

$$\lg (rRNA_{CNPC}) = 1.05 \lg (MV) + 4.47 \text{ or } rRNA_{CNPC} = 29512 \, MV^{1.05} \tag{17}$$

$R^2 = 0.83, P < 0.001, n = 12$.

According to Fu and Gong (2017), the per-cell rRNA CNs in the log phase cells of *E. vannus*

and *S. sulcatum* were at similar levels as these in *Colpoda* cells of similar cell sizes [the $\lg(CV)$

ranging from 3.3 to 4.1; Fig. 4C]. However, these two marine species had about 30 ~ 80 times more

rDNA copies (109,000 ~ 263,000 per cell) than the *C. steinii* cells (1300 ~ 9540 copies per cell)

within the same cell size range (Fig. 4C). In other words, the previous data of *E. vannus* and *S.*

sulcatum data largely followed the scaling relationship between rRNA and CV of non-resting

Colpoda ($R^2 = 0.86$, $P < 0.001$), but was not consistent with the rDNA-CV relationship described in

the present study ($R^2 = 0.05$, $P < 0.001$).

Interestingly, based on a combination of datasets comprising both these previously reported for

Euplotes vannus and *Strombidium sulcatum* ($n = 8$; Fu and Gong, 2017) and the present data for

Colpoda spp. ($n = 4$), the maximum growth rate (r) of these species at a given culturing temperature

(t) can be well predicted by $rDNA_{CNPC}$, but not $rRNA_{CNPC}$ ($n = 12$), as follows:

$\ln r = 0.051\, t - 0.474 \ln (rDNA_{CNPC}) + 4.192$ (18)

$R^2 = 0.911$, adjusted $R^2 = 0.891$, $P < 0.001$, $n = 12$ (Fig. 4D)

$\ln r = 0.131\, t + 0.229 \ln (rRNA_{CNPC}) - 6.323$ (19)

$R^2 = 0.457$, adjusted $R^2 = 0.336$, $P = 0.064$, $n = 12$

$\ln r = 0.084\, t + 0.289 \ln (rRNA_{CNPC} : rDNA_{CNPC}) - 3.121$ (20)

$R^2 = 0.799$, adjusted $R^2 = 0.755$, $P < 0.001$, $n = 12$

These predictions of maximum growth rate based on $rDNA_{CNPC}$ or $rRNA_{CNPC}$: $rDNA_{CNPC}$

outperformed the ones based on CV:

$\ln r = 0.126\, t + 0.206 \ln (CV) - 4.420$ (21)

$R^2 = 0.45$, adjusted $R^2 = 0.33$, $P = 0.069$

Dominance of a single haplotype in single-cell rDNA and rRNA pools

A total of 124 rDNA and cDNA samples derived from 124 single cells of the four ciliate

species were successfully sequenced. After quality filtering, removal of chimeric and singleton

sequences, the cleaned reads of 18S rDNA or cDNA fragments were subjected to statistical analyses

to characterize the distributional patterns of dominant haplotypes or variants under different

conditions (Table S1). In each species, there was a single haplotype, which was numerically

dominant (referred to as dominant haplotype, DH), and many variants in the rDNA (or rRNA) pool

(Table S1).

The variant sequences differed from the DH by having many single nucleotide substitutions.

The relative abundance of the DH in a single cell varied widely, depending on species, rDNA or

rRNA pools, life cycle stage, and temperature (Fig. 5A - D). The DHs accounted for 68% - 86% in *Colpoda* spp., the proportion of which was higher than that of *E. vannus* and *S. sulcatum* (37% to 46%). The DH percentages in the rDNA pools appeared to be higher than in the rRNA pools in both *Colpoda* species (*C. steinii:* 84.44% ±0.52% vs. 79.92% ±0.80%; $P < 0.001$, $n = 20$; and *C. inflata*: 82.55% ± 1.13% vs. 74.97% ± 2.10%; $P = 0.006$, $n = 20$). At a given life cycle stage and temperature ($n = 3$), however, the differences in DH percentage between rDNA and rRNA pools were frequently not significant with some exceptions. The DH had much lower proportions in the rRNA pools ($P < 0.05$) in the 28 ℃ treatments of resting cysts of *C. steinii* (Fig. 5A), and all the treatments of *C. inflata* (Fig. 5A, B), while DH proportion was higher ($P < 0.05$) in the rDNA pools than in the touchdown treatments (16*) of *E. vannus* (Fig. 5C).

The resting cysts of *Colpoda* spp. were frequently comparable to the log-phase cells in terms of DH proportions in both rDNA and rRNA pools (Fig. 5A, B). Nevertheless, significant differences in DH proportions between these two life cycle stages were found for *C. inflata* reared at 28 ℃, in which the resting cysts had lower levels in both rDNA (81.70% vs. 85.02%) and rRNA pools (78.85% vs. 82.73%) (Fig. 5B).

The effect of temperature on DH proportion was significant in several pairwise comparisons (Fig. 5B, C). Compared with the treatments of 18 ℃, the DH proportion in the rDNA pool significantly decreased ($P < 0.05$), but the opposite was observed in the rRNA pool of resting cysts of *C. inflata* ($P < 0.05$; Fig. 5B). In the actively growing cells of *E. vannus*, the lowest DH proportion in rRNA pool was found in the treatments of 21 ℃. This pattern, however, was not observed for the DH in the rDNA pool of this species (Fig. 5C).

Single-cell landscape of rDNA and rRNA variants

Despite many more quality reads having been obtained, fewer rDNA variants were detected in single cells of *C. steinii* (300 ~ 700 cell^{-1}) and *C. inflata* (400 ~ 800 cell^{-1}), compared with those in *E. vannus* (3200 ~ 3400 cell^{-1}) and *S. sulcatum* (2200 ~ 2700 cell^{-1}) (Fig. 5E; Table S1). In contrast, the single-cell rRNA pools of *C. steinii* (600 ~ 900 cell^{-1}), *C. inflata* (850 ~ 950 cell^{-1}), *E. vannus* (3000 ~ 3600 cell^{-1}) and *S. sulcatum* (2100 ~ 3000 cell^{-1}) generally exhibited higher numbers than

their rDNA variants (Fig. 5F; Table S1). Regression analysis indicated that the number of rDNA variants was positively related with rDNA CN per cell ($R^2 = 0.97$, $P < 0.001$; Fig. 5E), whereas the number of rRNA variants in a single cell was not significantly related to cellular rRNA CN of these species ($R^2 = 0.093$, $P > 0.21$; Fig. 5F). These patterns remained even when the variants were called by randomly sampling an equal number (1600) of rDNA and rRNA reads per sample (Table S1).

The hypotheses that life cycle stage and temperature affect the numbers of rDNA and rRNA variants were also tested (Fig. 6A). To standardize and minimize the uneven number of reads obtained from each sample, the variants were called by randomly sampling an equal number (1600) of rDNA or rRNA reads per sample (Table S1). Significant differences in numbers of rDNA and rRNA variants were exclusively found among certain treatments of the *Colpoda* species, but never among the temperature treatments of log-phase *E. vannus* and *S. sulcatum*. More specifically, contrasting numbers of rRNA variants (4 out of 6 cases) occurred more often than those of rDNA variants (2 out of 6 cases). For example, at 18 ℃, a resting cyst of both *C. steinii* (118 ± 6) and *C. inflata* (116 ± 2) was significantly richer in rRNA variants than a log-phase cell of these species (102 ± 2 and 103 ± 4, respectively; $P < 0.05$). The cell phase effect on rDNA variant numbers was significant only in the treatments of *C. inflata* reared at 28 ℃ ($P < 0.05$). The number of rDNA (and rRNA) variants differed between the two temperature treatments ($P < 0.05$) in resting cysts of *C. inflata*; whereas a temperature effect on the number of ribotype variants in log cells was detected only in *C. steinii* (Fig. 6A).

By rarefying at 1600 reads and clustering operational taxonomic units (OTUs) at thresholds ranging from 90% to 99%, we identified the maximum differences among the rDNA and rRNA variants in single cells of these ciliate species (Fig. 6B). The highest divergence of both rDNA and rRNA variants was 9%, which was observed in *C. steinii*. The lowest divergence was observed in *S. sulcatum*, in which intra-individual differences of rDNA and rRNA variants did not exceed 1%. Both *C. inflata* and *E. vannus* exhibited intermediate levels of divergence in both rDNA (6% and 7%) and rRNA (8% and 4%) (Fig. 6B).

The obtained numbers of OTUs ranged from 1 to 5 at a cutoff of 99%, and gradually decreased with lower cutoffs. Typically, at the 97% cutoff that has been popularly used for metabarcoding

surveys, on average of 1 to 2 OTUs remained for the *Colpoda* species. The resting cysts usually

appeared to have more OTUs than the log-phase cells, however, this was significant ($P < 0.05$) only

in 3 out of 8 pairwise comparisons for *Colpoda* spp. The temperature effect on OTU numbers in

single cells was even less frequent, observed only in 1 out of 8 comparisons for *Colpoda*, and in

none for *E. vannus* and *S. sulcatum* ($P > 0.05$) (Fig. 6B).

Nucleotide polymorphisms

The intra-individual sequence variations of rDNA and rRNA variants in the single cells of the

four ciliate species were exclusively single nucleotide polymorphisms (SNPs). By setting the

consensus sequence (the DH) as the reference, the number of polymorphic sites and types of

nucleotide substitutions were identified (Figs 7 and 8). The frequency of polymorphic sites (SNP

density) ranged from 4.0×10^{-4} to 2.9×10^{-3} in rDNA, and from 4.9×10^{-4} to 6.3×10^{-3} in rRNA.

Nucleotide replacements occurred in all positions along the sequenced rDNA and rRNA fragments,

most of which had a frequency lower than 3.0×10^{-3}. Fifteen sites were highly polymorphic (with a

replacement frequency of $> 3.0 \times 10^{-3}$), and their positions were conserved in the two *Colpoda*

species and in both ribotypes. Likewise, 20 positions exhibited high nucleotide replacement

frequencies in both rDNA and rRNA in *E. vannus* and *S. sulcatum* (Fig. 7A). Relative to the log

phase cells of *Colpoda*, the resting cysts often had significantly higher frequencies of nucleotide

substitutions (SNP density) in rRNA, but not rDNA. The effect of temperature was only significant

for *E. vannus*, in which the SNP density of rDNA was much higher in the touchdown (16* ℃) than

in the treatment at 16 ℃, and even higher at 21 ℃ than all other treatments of this species (Fig. 7B).

Overall, the transition: transversion (Ts: Tv) ratio in rDNA and rRNA was conserved within

species, but varied from 1.1 to 3.0 among the four ciliate species (Fig. 8). Temperature and

encystment had no significant effects on the Ts: Tv ratio within a species either ($P > 0.05$). However,

individual nucleotide substitution categories showed different patterns. For example, transitional

A/G substitutions were more frequent than C/T substitutions in both rDNA and rRNA transcripts of

all ciliate species, except for *C. steinii*, which showed the opposite pattern, especially between two

life cycle stages. In the 18 ℃ treatments of *C. steinii*, the frequency of A/G substitutions in rDNA

was significantly lower in resting cysts (on average 10.3%) than in log-phase cells (22.4%), whereas the reverse was observed for transitional C/G substitutions (43.4% for resting cysts and 29.6% for log-phase cells). Transversions in the rDNA and in the expressed transcripts appeared to be frequently higher in log phase cells vs. resting cysts of *Colpoda* spp. in both temperature treatments ($P < 0.05$). In the rDNA of *C. inflata*, G/C substitutions occurred more frequently in resting cysts (on average 5.0%) than in log cells (3.5%) at 28 ℃. In contrast, in rRNA transcripts of *C. inflata*, the cell phase-wise differences were observed in 5 of the 6 nucleotide substitution categories, especially in the 28 ℃ treatments (Fig. 8).

There were only a few cases in which the frequency of nucleotide substitution was significantly different between temperature treatments (Fig. 8). Higher frequencies in transversional G/T (17.8% vs. 13.6% in *C. steinii*) and G/C substitutions (5.0% vs. 2.9% in *C. inflata*) occurred in resting cysts at 28 ℃ (vs. at 18 ℃). Nevertheless, the variants with G/T substitutions in log-phase cells of *C. steinii* were more highly expressed at the lower temperatures (17.1% vs. 15.2%).

DISCUSSION

Allometric scaling relationship of nuclear volume to cell volume in *Colpoda*

Single cells of two *Colpoda* species in all growing phases and cystic forms obtained from two temperature treatments were fixed, stained with protargol and their cell and nuclear volumes measured. Both volumes progressively reduced from lag, log, and plateau phases to resting cysts, the latter being a phenotypic trait common in many protists (e.g., a scuticociliate, Pérez-uz and Guinea, 2001) a dinoflagellate, Anderson and Lindquist, 1985). In parallel, the N/C ratio in both *Colpoda* species significantly increased, especially after resting cyst formation. The increase was about a 3-fold difference in *C. steinii* reaching an N/C ratio of 7.31% and in *C. inflata* 4.35% (Fig. 2D, F). These findings are in contrast to the "karyoplasmic" ratio theory, which postulates that eukaryotic cells of the same type or species maintain a constant N/C ratio. For example, despite fission and budding yeasts spanning a large range of cell sizes ($10 \sim 300\ \mu m^3$), their N/C ratio is approximately 7% to 8% (Jorgensen et al., 2007; Neumann and Nurse, 2007). Our data indicate that the N/C volume ratio of the two *Colpoda* species examined herein inversely scales with cell volume to a power of -0.174. Thus, both species display allometry similar to that previously described for

amoebae (Rogerson et al., 1994) and *Xenopus* cells during early embryonic development (Jevtić and

Levy, 2015), rather than isometry of N/C ratio with CV, which has been commonly observed in

many other eukaryotes (Reber and Goehring, 2015). In this study, the *Colpoda* cells covered a

broad size range (from 7500 μm³ to 2.14 ×10⁵ μm³) at all life cycle stages allowing for a good fit of

the scaling relationship with power-law functions. Linear models might fit the data equally well (so

that missed the curved trajectory corresponding to a large size range) with relatively narrow ranges

of cell size or very large cells. According to the ciliate-based equation 2, calculations based on the

average CV of wild-type fission and budding yeasts yield the same N/C ratios as experimentally

determined, indicating that the allometry between N/C ratio and CV could be widely applicable

within and among eukaryotic species. Given that it is relatively easy to measure the volume of

nucleus, which is usually globular or ellipsoid in living and/or fixed materials, the N/C ratio

allometry provides a tool to calculate biovolume (biomass) of amorphous cells (e.g., amoebae)

using nuclear size (Rogerson et al., 1994).

Ribotype copy number variations across different phases of growth and cystic forms

Due to macronuclear polyploidy and micronuclear diploidy in ciliates, most rDNA copies

quantified and sequenced in this study were of macronuclear origin. Substantial variations in rDNA

and/or rRNA quantities have been previously recorded in randomly selected individuals of ciliates

(Gong et al., 2013; Wang et al., 2019), log phase populations (Fu and Gong, 2017), and among cells

of different phases of growth or physiological states in a range of protistan species (e.g., a

Tetrahymena ciliate, Engberg and Pearlman, 1972; a raphidophyte, Main et al., 2014; a foraminifer

strain, Parfrey et al., 2012; and dinoflagellates, Galluzzi et al., 2010; Eckford-Soper and Daugbjerg,

2016; Nishimura et al., 2016). Nevertheless, our study provides the first evidence of linkage

between ribotypes and phenotypes across life cycle stages of protists. Specifically, several ribotypic

traits (i.e., the per-cell CN, cellular concentrations of rRNA and rDNA molecules, along with their

CN ratio) are quantitatively linked with cell and macronuclear size, across all growing phases and

dormant cystic forms. This finding extends our understanding obtained by observations of log phase

cells of *E. vannus* and *S. sulcatum*, and reinforces previous recommendations, that in molecular

protistan ecology, ribotype CNs have to be interpreted to reflect biomass rather than solely cell

abundance (Fu and Gong, 2017).

Chilling of actively growing *Colpoda* cells at 0 ~ 4 ℃ for a few hours artificially induced

formation of unstable cysts (Fig. S1), an event that was accompanied by significant reduction in

per-cell rDNA and rRNA CNs, as well as, cell size (Fig. 2A; 3A, B). Freezing raw environmental

samples (e.g. raw soil, sediment or water samples) temporarily to preserve microbiomes is

commonly used in molecular ecology. Herein, we show that this practice might artificially alter both

phenotypic and ribotypic traits of certain protistan populations and thus the whole community

structure. This could lead to underestimation of their biomass and proportions of rDNA and rRNA

quantities in the ribotypic pools. Nevertheless, chilling treatment is a simple, non-destructive

approach that arrests growing cells at the unstable cyst stage, thus narrowing down the cell size

spectrum of a population consisting of different-sized cells in various growth phases, while

preserving cell numbers in the original samples. Thus, chilling treatment has the potential to

enhance the accuracy in estimating population abundances using rDNA-based qPCR or

hybridization methods. Abundance estimates derived from either method tend to be inconsistent due

to variable rRNA gene content across growth phases in natural populations of harmful marine

microalgae (Galluzzi et al., 2010; Main et al., 2014; Eckford-Soper and Daugbjerg, 2016;

Nishimura et al., 2016). Second, we provide evidence that both types of *Colpoda* cysts follow the

rDNA CN-CV relationship (Fig. 4; Equations 4 and 5), so their biomass can be reasonably predicted

in rDNA-based molecular surveys. Third, the two types of cysts have variable amount of rRNA or

ribosomes. In the unstable cysts, the amount of rRNA copies corresponds well with cell size,

whereas a resting cyst has much fewer rRNA molecules. This is a unique ribotypic trait of resting

cysts that cannot readily be explained by their small cell size (Fig. 4). These results clarify the

ribotypic distinctness between unstable and resting cysts. Herein, we demonstrate a theoretical

framework that makes use of the experimentally determined rRNA:rRNA ratio of a given *Colpoda*

species. The framework can be adopted to estimate the proportion of resting cysts in the rDNA pool

of a population. Importantly, our proposed framework takes into account cell size (see Equation 14),

indicating that it affects estimation of resting cysts. This implies that caution should be taken in

using rRNA: rDNA ratios to compare within and between species/groups of contrasting cell sizes.

Thus, qPCR could be applied in future studies for quantifying cysts of *Colpoda*, and/or probably

also other protists, in environmental samples.

709 Cellular rRNA content scales inversely with cell size, but maximum growth rate depends on

710 rDNA content

Previously, a scaling relationship between rRNA content and cell size of log-phase cells of two

protist species and five species of bacteria and archaea was proposed (Fu and Gong, 2017). This

study provides evidence that the scaling relationship generally holds, irrespective of life cycle

stages. These findings collectively support the universality of the scaling law between rRNA

content and cell/body size applying to micro- to macro-organisms (Gillooly et al., 2005).

Nevertheless, the scaling relationship did not apply to resting cysts, as they exhibited significantly

lower rRNA than expected, indicating that rRNA is a suitable molecular marker for not only

revealing "active" components, but also for providing a biomass-based view of community

structure of microbial communities. Previous studies of mock protistan communities showed

consistent patterns using rRNA and cell counting approaches (Weber and Pawlowski, 2013).

Unlike rRNA, the scaling relationship between rDNA and cell volume was not uniform for any

of the four ciliate species examined. Instead, two separate functions were applied; one for *Euplotes*

and *Strombidium* (rDNA ~ $CV^{0.44}$; Fu and Gong, 2017) and another for *Colpoda* spp. (rDNA ~

$CV^{0.76}$; this study). This is primarily due to the much higher cellular rDNA content in *E. vannus* and

S. sulcatum (23 - 59 copies per μm^3) than in the similar sized cells of *Colpoda steinii* and *C. inflata*

(0.2 - 0.8 copies per μm^3) (Fig. 4C). The disproportion between per-cell rDNA copy number and

cell size was also pointed out by Wang et al. (2019). The variable rDNA content between

Spirotrichea and Colpodea may stem from their genomic distinctness. Spirotrichean ciliates are well

known for their highly polyploidized genomic macronuclear DNA (Prescott, 1994). The higher

rDNA content also implies higher genomic DNA content and larger macronuclear volumes

(Cavalier-Smith, 1978). In support of this, both *E. vannus* and *S. sulcatum* have disproportionately

large macronuclei (N/C ratios about 4.71% and 2.73%, respectively) that do not correspond to the

predicted values (1.57% and 2.43%) of the *Colpoda*-based model (Equations 1 and 2) and their

actual average cell volumes (2.34 $\times$ 10^5 and 1.93 $\times$ 10^4 μm³). This indicates that larger

macronuclear size does contribute to high rDNA content in these two spirotricheans. Even if the

"enlarged" macronuclear sizes of *E. vannus* and *S. sulcatum* are taken into consideration, their

calculated per-cell rDNA copy numbers according to the *Colpoda*-based function (Equation 15),

will be only about 1/2 and 1/16 of these experimentally determined. This suggests that the high

rDNA content of spirotricheans is not just due to their unusual phenotype (enlarged macronuclei),

but that certain, as yet undiscovered, mechanisms for genomic innovation may also be at play.

Interestingly, *Colpoda steinii* grew 7 to 12 times faster than *E. vannus* and *S. sulcatum*, despite

these three species having comparable cell size and cellular rRNA content. The model

parameterized with rDNA performed much better than the one with rRNA in predicting growth rate

(Equations 18 and 19). This indicates that, besides temperature (which modulates cell size via the

temperature-size rule), cellular rDNA content is an important factor in determining generation time

of these protists. This observation seems to violate the grow rate hypothesis (GRH) that has been

repeatedly validated in diverse organisms, which postulates that a high growth rate depends on a

high rRNA or ribosomal concentration, or phosphorus (P) content in biomass (Elser et al., 2000). To

reconcile the two, we propose that growth rate is determined by rRNA content, but can be modified

by rDNA contents. In other words, growth rate depends mainly on rDNA, when the cells have

similar cell sizes or rRNA contents (see Fig. 4C).

Existing studies on genomic content and growth rates of various organisms can shed light in

understanding the mechanisms underlying modulation of cell growth rate by rDNA content. For

instance, in a previous study, haploid cells of budding yeast with a smaller cell size grew more

rapidly than their diploid counterparts of larger cell size. The observed fitness difference was

attributed to their cell surface to volume ratio. Specifically, growth rate was dependent on the

transfer of products into cytoplasm generated upon hydrolysis of organic phosphorous by a cell

surface-bearing acid phosphatase (Weiss et al., 1975; Mable and Otto, 2001). The

surface-to-volume-ratio theory does not seem to apply to phagotrophic ciliates, which graze on food

particles through their cytostome and digest them inside cytoplasmic food vacuoles. Nevertheless,

the lower nuclear to cell volume ratio of *C. steinii* may provide a relatively larger cytoplasmic space

to accommodate food vacuoles, enabling digestion of more bacterial prey and maintenance of a

higher rate of material supply for growth than *E. vannus* and *S. sulcatum*. Spirotricheans have a

high content of rDNA and genomic DNA, both of which are particularly P- and N-rich. Thus, these

ciliates need more element-bearing resources and invest a longer time for DNA replication,

synthesis of associated proteins, and cell division (Hessen et al. (2010). Conversely, the similarly

sized colpodean ciliate *C. steinii* with fewer rDNA copies and a more streamlined genome could be

able to save more P and N to synthesize rRNA and assemble ribosomes for rapid growth.

Intra-individual rDNA and rRNA sequence variations across life cycle stages and at different

temperatures

In the present study, the intragenomic polymorphisms of 18S rDNA were examined and a level

of divergence of up to 9% was noted. This is comparable to the divergence uncovered using high

throughput sequencing in other surveys (13% in radiolarians, Decelle et al., 2014; up to about 16%

in a *Protocruzia* ciliate, Zhao et al., 2019). Previous studies of rDNA polymorphism using clone

libraries and Sanger sequencing yielded lower intragenomic divergence (< 2%) in ciliates (Fu and

Gong, 2017; Wang et al., 2017; Wang et al., 2019) and diatom species (0.57%-1.81%, Alverson and

Kolnick, 2005), while divergence was moderate (about 5%) in foraminifera (Weber and Pawlowski,

2014). These results suggest that estimation of intra-individual variability in rDNA sequences

depends on sequencing depth, and that variability is likely much higher than previously recognized

for many protistan groups. Use of polymerases of varying fidelity also affects polymorphism

estimations (Wang et al., 2017). Intragenomic polymorphisms in rDNA along with experimental

bias could inflate species richness estimations (Gong et al., 2013; Caron and Hu, 2019). Fortunately,

the observed variation seemed to have negligible effect on the rDNA-based OTU number of a

protistan species, at least under the conditions examined in this study (temperature gradients; and

cystic vs. rapidly growing life cycle stages), provided that the OTUs were defined at a proper

sequence similarity threshold (e.g., 97%; Fig. 6). These results indicate that, even though the OTU

richness tends to be inevitably higher than the species richness in a multi-species community, the

variation pattern in rDNA-based OTU richness is, to some extent, still informative to reflect

organizational changes among a series of communities collected across time and space.

Although intra-individual rDNA polymorphisms have been previously recognized, our study is

the first to provide evidence that multiple rDNA variants are extensively expressed in protists,

resulting in hundreds to several thousands of rRNA variants (Fig. 6E, F). The variants could

combine with ribosomal proteins, giving rise to extremely high ribosomal heterogeneity in these

ciliates. Such a high diversity of ribosomes in single-celled eukaryotes is surprising, because

ribosomes have long been thought of as homogeneous cellular machinery. A notable exception is

the parasitic protist, *Plasmodium berghei*, in which two structurally distinct 18S rRNA transcripts

were found (Gunderson et al., 1987). Recent studies have suggested that ribosomal variants could

preferentially bind different mRNAs to regulate gene expression and corresponding phenotypes of

bacteria under stress conditions (Genuth and Barna, 2018; Kurylo et al., 2018; Song et al., 2019).

Given the high intracellular rDNA diversity in diverse protistan groups, the significance of

ribosomal heterogeneity in the context of physiological and ecological adaptations of protists

inhabiting various ecosystems would be an interesting topic to explore.

The lower Variants% in both rDNA and rRNA pools of the soil species *Colpoda* could be

related to their lower ribotypic copy numbers than these of the marine species *Euplotes* and

Strombidium (Fig. 5E). Analogous to the evolutionary significance of polyploidy (Van de Peer et al.,

2017), it is possible that many of the ribotypic sequences are redundant in single cells of *Euplotes*

and *Strombidium*, allowing for increased mutational robustness. This would enhance cellular

ribosome heterogeneity thus facilitating adaptation to rapidly changing littoral environments.

Interestingly, the number of rRNA variants remained stable when comparing the log and resting

cystic stages of *Colpoda* species (Fig. 5E), indicating that shrinkage of the cellular rRNA pool in

resting cysts is mainly due to the dominant haplotype being expressed less (Fig. 5A, B), rather than

the variants. Keeping a large number of rRNA variants (and by extension, a high diversity of

ribosomes) in the resting cysts may be ecologically significant for soil ciliates in maintaining a high

metabolic potential to readily respond to favorable environmental conditions.

Previous characterization using clone libraries and Sanger sequencing, detected limited

numbers of polymorphic sites, resulting in random distributions of SNPs in rDNA and rRNA

variants (Alverson and Kolnick, 2005; Gong et al., 2013; Weber and Pawlowski, 2014). In this

study, undersampling was largely overcome by using high throughput sequencing, thus additional

attributes of SNPs in rDNA and rRNA fragments could be assessed: (1) within a species, the

distributional patterns of polymorphic sites in rDNA and their transcripts were almost identical

between rDNA and rRNA; (2) the positions of highly polymorphic sites were conserved between *C.*

steinii and *C. inflata*, and between *E. vannus* and *S. sulcatum*; and (3) both rDNA and rRNA had

hotspot regions, which exhibited much higher substitution frequencies than other regions (Fig. 7A).

Furthermore, the rDNA sequences of all four ciliate species exhibited a higher number of transitions

than transversions. This finding is consistent with previous clone-library-based results for four other

ciliates (Gong et al., 2013), six diatom strains (Alverson and Kolnick, 2005) and the model plant

Arabidopsis thaliana (Mentewab et al., 2011), and generally comparable with protein-coding

sequences of many other species (Stoltzfus and Norris, 2015). Whole genome mutation

accumulation rates of the four ciliate species in this study are not available yet. Nevertheless,

compared with previously estimated genomic transition: transversion (Ts: Tv) ratios in *Paramecium*

biaurelia (1.90) and *P. sexaurelia* (1.67) (Long et al., 2018), the Ts: Tv ratios of 18S rDNA

fragments were slightly lower in *C. steinii* (1.1~1.2) and *C. inflata* (1.3 ~ 1.6), but higher in *E.*

vannus (2.8~ 3.0) and *S. sulcatum* (2.1 ~ 2.6) (Fig. 7B). All these new findings are in support of the

observed nucleotide replacements in the ribotype variants not being random and likely reflecting the

non-concerted evolution nature of rDNA in protists and many eukaryotic organisms.

To the best of our knowledge, this is the first study investigating nucleotide substitution

patterns of rDNA and rRNA at different life cycle stages and temperatures. Contrasting frequencies

of certain nucleotide substitution categories were exclusively detected between resting cysts and

log-phase cells of *Colpoda* species (Fig. 8). This could be a consequence of losing variant copies

during the biological process of encystment, during which dramatic changes in nucleolar

morphology occur and a large portion of macronuclear chromatin is degenerated via cell autophagy

(Gutiérrez et al., 1998). Resting cysts have specialized mRNA and proteins, which have been

previously been associated with loss of RNA synthetic capability (Benfez and Gutiérrez, 1997;

Gutiérrez et al., 1998). Notably, *C. inflata* undergoes demethylation of macronuclear DNA during encystment (Palacios et al., 1994). Nevertheless, the link between changes in nucleotide polymorphisms and rRNA-derived ribosomal heterogeneity remains unknown. Further investigations are needed to explore the mechanistic links between DNA methylation, rRNA-derived ribosomal heterogeneity, and mRNA translation towards a better understanding of the biology and ecology of protistan adaptations.

Acknowledgments

This work was supported by the grants from the Natural Science Foundation of China (Nos. 31572255 and 41976128), the Marine S & T Fund of Shandong Province for Pilot National Laboratory for Marine Science and Technology (Qingdao) (No. 2018SDKJ0406-4), and the Key Research Program of Frontier Sciences (QYZDB-SSW-DQC013). Thanks are due to Prof. Hongan Long (OUC) for suggestions on discussing nucleotide mutations.

REFERENCES

1. Caron DA, Countway PD, Savai P, Gast RJ, Schnetzer A, Moorthi SD, Dennett MR, Moran DM, Jones AC. 2009. Defining DNA-based operational taxonomic units for microbial-eukaryote ecology. Appl Environ Microbiol 75:5797-5808. https://doi.org/10.1128/aem.00298-09.

2. Corliss JO. Protozoan cysts and spores[M]. eLS, 2001.

3. Padilla DK, Savedo MM. A systematic review of phenotypic plasticity in marine invertebrate and plant systems[M]. In: Lesser M, editor. Adv Mar Biol. 65. Academic Press, 2013, pp. 67-94.

4. Mulot M, Marcisz K, Grandgirard L, Lara E, Kosakyan A, Robroek BJM, Lamentowicz M, Payne RJ, Mitchell EAD. 2017. Genetic determinism vs. phenotypic plasticity in protist morphology. J Eukaryot Microbiol 64:729-739. https://doi.org/10.1111/jeu.12406.

5. McQuoid MR, Hobson LA. 1996. Diatom resting stages. J Phycol 32:889-902. https://doi.org/10.1111/j.0022-3646.1996.00889.x.

6. Persson A. 2001. Proliferation of cryptic protists and germination of resting stages from untreated sediment samples with emphasis on dinoflagellates. Ophelia 55:151-166. https://doi.org/10.1080/00785326.2001.10409482.

7. Ross BJ, Hallock P. 2016. Dormancy in the foraminifera: A review. J Foraminifer Res 46:358-368. https://doi.org/10.2113/gsjfr.46.4.358.

8. Corliss JO, Esser SC. 1974. Comments on the role of the cyst in the life cycle and survival of free-living protozoa. Trans Am Microsc Soc 93:578-593. https://doi.org/10.2307/3225158.

9. Fistarol GO, Legrand C, Rengefors K, Granéli E. 2004. Temporary cyst formation in phytoplankton: A response to allelopathic competitors? Environ Microbiol 6:791-798. https://doi.org/10.1111/j.1462-2920.2004.00609.x.

10. Anderson DM, Rengefors K. 2006. Community assembly and seasonal succession of marine dinoflagellates in a temperate estuary: The importance of life cycle events. Limnol Oceanogr 51:860-873. https://doi.org/10.4319/lo.2006.51.2.0860.

886   11.   Fu R, Gong J. 2017. Single cell analysis linking ribosomal (r)DNA and rRNA copy numbers
887   to cell size and growth rate provides insights into molecular protistan ecology. J Eukaryot
888   Microbiol 64:885-896. https://doi.org/10.1111/jeu.12425.

889   12.   Erdner DL, Percy L, Keafer B, Lewis J, Anderson DM. 2010. A quantitative real-time PCR
890   assay for the identification and enumeration of *Alexandrium* cysts in marine sediments.
891   Deep   Sea   Res   Part   II   Top   Stud   Oceanogr   57:279-287.
892   https://doi.org/10.1016/j.dsr2.2009.09.006.

893   13.   Rivière D, Szczebara FM, Berjeaud J-M, Frère J, Héchard Y. 2006. Development of a
894   real-time PCR assay for quantification of *Acanthamoeba* trophozoites and cysts. J Microbiol
895   Methods 64:78-83. https://doi.org/10.1016/j.mimet.2005.04.008.

896   14.   Gutiérrez JC, Martin-González A, Callejas S. 1998. Nuclear changes, macronuclear
897   chromatin reorganization and DNA modifications during ciliate encystment. Eur J Protistol
898   34:97-103. https://doi.org/10.1016/S0932-4739(98)80018-7.

899   15.   Parfrey LW, Grant J, Katz LA. 2012. Ribosomal DNA is differentially amplified across
900   life-cycle stages in the foraminifer *Allogromia Laticollaris* strain CSH. J Foraminifer Res
901   42:151-155. https://doi.org/10.2113/gsjfr.42.2.151.

902   16.   Pillet L, Fontaine D, Pawlowski J. 2012. Intra-genomic ribosomal RNA polymorphism and
903   morphological variation in *Elphidium macellum* suggests inter-specific hybridization in
904   foraminifera. PloS One 7:e32373. https://doi.org/10.1371/journal.pone.0032373.

905   17.   Gong J, Dong J, Liu X, Massana R. 2013. Extremely high copy numbers and
906   polymorphisms of the rDNA operon estimated from single cell analysis of oligotrich and
907   peritrich ciliates. Protist 164:369-379. https://doi.org/10.1016/j.protis.2012.11.006.

908   18.   Decelle J, Romac S, Sasaki E, Not F, Mahé F. 2014. Intracellular diversity of the V4 and V9
909   regions of the 18S rRNA in marine protists (Radiolarians) assessed by high-throughput
910   sequencing. PLoS One 9:e104297. https://doi.org/10.1371/journal.pone.0104297.

911   19.   Weber AA-T, Pawlowski J. 2014. Wide occurrence of SSU rDNA intragenomic
912   polymorphism in foraminifera and its implications for molecular species identification.
913   Protist 165:645-661. https://doi.org/10.1016/j.protis.2014.07.006.

914   20.   Kudryavtsev A, Gladkikh A. 2017. Two new species of *Ripella* (Amoebozoa, Vannellida)

915   and unusual intragenomic variability in the SSU rRNA gene of this genus. Eur J Protistol

916   61:92-106. https://doi.org/10.1016/j.ejop.2017.09.003.

917   21.   Wang C, Zhang T, Wang Y, Katz LA, Gao F, Song W. 2017. Disentangling sources of

918   variation in SSU rDNA sequences from single cell analyses of ciliates: Impact of copy

919   number variation and experimental error. Proc R Soc Lond B Biol Sci 284:20170425.

920   https://doi.org/10.1098/rspb.2017.0425.

921   22.   Gunderson JH, Sogin ML, Wollett G, Hollingdale M, de la Cruz VF, Waters AP, McCutchan

922   TF. 1987. Structurally distinct, stage-specific ribosomes occur in Plasmodium. Science

923   238:933. https://doi.org/10.1126/science.3672135.

924   23.   López-López A, Benlloch S, Bonfá M, Rodríguez-Valera F, Mira A. 2007. Intragenomic 16S

925   rDNA divergence in *Haloarcula marismortui* Is an adaptation to different temperatures. J

926   Mol Evol 65:687-696. https://doi.org/10.1007/s00239-007-9047-3.

927   24.   Genuth NR, Barna M. 2018. The discovery of ribosome heterogeneity and its implications

928   for   gene   regulation   and   organismal   life.   Mol   Cell   71:364-374.

929   https://doi.org/10.1016/j.molcel.2018.07.018.

930   25.   Kurylo CM, Parks MM, Juette MF, Zinshteyn B, Altman RB, Thibado JK, Vincent CT,

931   Blanchard SC. 2018. Endogenous rRNA sequence variation can regulate stress response

932   gene   expression   and   phenotype.   Cell   Rep   25:236-248.e6.

933   https://doi.org/10.1016/j.celrep.2018.08.093.

934   26.   Stout JD. 1955. Environmental factors affecting the life history of three soil species of

935   *Colpoda* (Ciliata). Trans Roy Soc New Zealand 82:1165-1188. https://doi.org/.

936   27.   Müller H, Achilles-Day UEM, Day JG. 2010. Tolerance of the resting cysts of *Colpoda*

937   *inflata* (Ciliophora, Colpodea) and *Meseres corlissi* (Ciliophora, Spirotrichea) to desiccation

938   and freezing. Eur J Protistol 46:133-142. https://doi.org/10.1016/j.ejop.2009.12.004.

939   28.   Omar A, Zhang Q, Zou S, Gong J. 2017. Morphology and phylogeny of the soil ciliate

940   *Metopus yantaiensis* n. sp. (Ciliophora, Metopida), with identification of the intracellular

941   bacteria. J Eukaryot Microbiol 64:792-805. https://doi.org/10.1111/jeu.12411.

29. Wilbert N. 1975. Eine verbesserte technik der protargo-limpragnation fur ciliaten.
Mikrokosmos 64:171-179.

30. Warren A, Patterson DJ, Dunthorn M, Clamp JC, Achilles-Day UEM, Aescht E, Al-Farraj
SA, Al-Quraishy S, Al-Rasheid K, Carr M, Day JG, Dellinger M, El-Serehy HA, Fan Y, Gao
F, Gao S, Gong J, Gupta R, Hu X, Kamra K, Langlois G, Lin X, Lipscomb D, Lobban CS,
Luporini P, Lynn DH, Ma H, Macek M, Mackenzie-Dodds J, Makhija S, Mansergh RI,
Martń-Cereceda M, McMiller N, Montagnes DJS, Nikolaeva S, Ong'ondo GO, Pérez-Uz B,
Purushothaman J, Quintela-Alonso P, Rotterová J, Santoferrara L, Shao C, Shen Z, Shi X,
Song W, Stoeck T, La Terza A, Vallesi A, Wang M, Weisse T, et al. 2017. Beyond the
"Code": A guide to the description and documentation of biodiversity in ciliated protists
(Alveolata, Ciliophora). J Eukaryot Microbiol 64:539-554.
https://doi.org/10.1111/jeu.12391.

31. Gong J, Qing Y, Guo X, Warren A. 2014. "*Candidatus* Sonnebornia yantaiensis", a member
of candidate division OD1, as intracellular bacteria of the ciliated protist *Paramecium*
bursaria (Ciliophora, Oligohymenophorea). Syst Appl Microbiol 37:35-41.
https://doi.org/10.1016/j.syapm.2013.08.007.

32. Zhu F, Massana R, Not F, Marie D, Vaulot D. 2005. Mapping of picoeucaryotes in marine
ecosystems with quantitative PCR of the 18S rRNA gene. FEMS Microbiol Ecol 52:79-92.
https://doi.org/10.1016/j.femsec.2004.10.006.

33. Stoeck T, Bass D, Nebel M, Christen R, Jones MDM, Breiner H-W, Richards TA. 2010.
Multiple marker parallel tag environmental DNA sequencing reveals a highly complex
eukaryotic community in marine anoxic water. Mol Ecol 19:21-31.
https://doi.org/10.1111/j.1365-294X.2009.04480.x.

34. Díez B, Pedrós-Alió C, Marsh TL, Massana R. 2001. Application of denaturing gradient gel
electrophoresis (DGGE) to study the diversity of marine picoeukaryotic assemblages and
comparison of DGGE with other molecular techniques. Appl Environ Microbiol
67:2942-2951. https://doi.org/10.1128/aem.67.7.2942-2951.2001.

35. Dawson SC, Pace NR. 2002. Novel kingdom-level eukaryotic diversity in anoxic

environments. Proc Natl Acad Sci U S A 99:8324. https://doi.org/10.1073/pnas.062169599.

36.   Schmieder R, Edwards R. 2011. Quality control and preprocessing of metagenomic datasets.

Bioinformatics 27:863-864. https://doi.org/10.1093/bioinformatics/btr026.

37.   Caporaso JG, Kuczynski J, Stombaugh J, Bittinger K, Bushman FD, Costello EK, Fierer N,

Pe ña AG, Goodrich JK, Gordon JI, Huttley GA, Kelley ST, Knights D, Koenig JE, Ley RE,

Lozupone CA, McDonald D, Muegge BD, Pirrung M, Reeder J, Sevinsky JR, Turnbaugh PJ,

Walters WA, Widmann J, Yatsunenko T, Zaneveld J, Knight R. 2010. QIIME allows analysis

of high-throughput community sequencing data. Nat Methods 7:335-336.

https://doi.org/10.1038/nmeth.f.303.

38.   Schloss PD, Westcott SL, Ryabin T, Hall JR, Hartmann M, Hollister EB, Lesniewski RA,

Oakley BB, Parks DH, Robinson CJ, Sahl JW, Stres B, Thallinger GG, Van Horn DJ, Weber

CF. 2009. Introducing mothur: Open-source, platform-independent, community-supported

software for describing and comparing microbial communities. Appl Environ Microbiol

75:7537. https://doi.org/10.1128/AEM.01541-09.

39.   Edgar RC. 2010. Search and clustering orders of magnitude faster than BLAST.

Bioinformatics 26:2460-2461. https://doi.org/10.1093/bioinformatics/btq461.

40.   Li H, Durbin R. 2009. Fast and accurate short read alignment with Burrows–Wheeler

transform. Bioinformatics 25:1754-1760. https://doi.org/10.1093/bioinformatics/btp324.

41.   Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin

R, Genome Project Data Processing S. 2009. The sequence alignment/map format and

SAMtools. Bioinformatics 25:2078-2079. https://doi.org/10.1093/bioinformatics/btp352.

42.   P érez-uz B, Guinea A. 2001. Morphology and infraciliature of a marine Scuticociliate with a

polymorphic life cycle: *Urocryptum tortum* n. gen., n. comb. J Eukaryot Microbiol

48:338-347. https://doi.org/10.1111/j.1550-7408.2001.tb00323.x.

43.   Anderson DM, Lindquist NL. 1985. Time-course measurements of phosphorus depletion

and cyst formation in the dinoflagellate *Gonyaulax tamarensis* Lebour. J Exp Mar Bio Ecol

86:1-13. https://doi.org/10.1016/0022-0981(85)90039-5.

44.   Jorgensen P, Edgington NP, Schneider BL, Rupeš I, Tyers M, Futcher B. 2007. The size of

the nucleus increases as yeast cells grow. Mol Biol Cell 18:3523-3532.

https://doi.org/10.1091/mbc.e06-10-0973.

45. Neumann FR, Nurse P. 2007. Nuclear size control in fission yeast. J Cell Biol 179:593-600.

https://doi.org/10.1083/jcb.200708054.

46. Rogerson A, Butler HG, Thomason JC. 1994. Estimation of amoeba cell volume from

nuclear diameter and its application to studies in protozoan ecology. Hydrobiologia

284:229-234. https://doi.org/10.1007/BF00006692.

47. Jevtić P, Levy DL. 2015. Nuclear size scaling during *Xenopus* early development contributes

to midblastula transition timing. Curr Biol 25:45-52.

https://doi.org/10.1016/j.cub.2014.10.051.

48. Reber S, Goehring NW. 2015. Intracellular scaling mechanisms. Cold Spring Harb Perspect

Biol 7. https://doi.org/10.1101/cshperspect.a019067.

49. Wang Y, Wang C, Jiang Y, Katz LA, Gao F, Yan Y. 2019. Further analyses of variation of

ribosome DNA copy number and polymorphism in ciliates provide insights relevant to

studies of both molecular ecology and phylogeny. Sci China Life Sci 62:203-214.

https://doi.org/10.1007/s11427-018-9422-5.

50. Engberg J, Pearlman RE. 1972. The amount of ribosomal RNA genes in *Tetrahymena*

pyriformis in different physiological states. Eur J Biochem 26:393-400.

https://doi.org/10.1111/j.1432-1033.1972.tb01779.x.

51. Main CR, Doll C, Bianco C, Greenfield DI, Coyne KJ. 2014. Effects of growth phase, diel

cycle and macronutrient stress on the quantification of *Heterosigma akashiwo* using qPCR

and SHA. Harmful Algae 37:92-99. https://doi.org/10.1016/j.hal.2014.05.014.

52. Galluzzi L, Bertozzini E, Penna A, Perini F, Garcés E, Magnani M. 2010. Analysis of rRNA

gene content in the Mediterranean dinoflagellate *Alexandrium catenella* and *Alexandrium*

taylori: implications for the quantitative real-time PCR-based monitoring methods. J Appl

Phycol 22:1-9. https://doi.org/10.1007/s10811-009-9411-3.

53. Eckford-Soper LK, Daugbjerg N. 2016. A quantitative real-time PCR assay for identification

and enumeration of the occasionally co-occurring ichthyotoxic *Pseudochattonella farcimen*

and *P. verruculosa* (Dictyochophyceae) and analysis of variation in gene copy numbers

during the growth phase of single and mixed cultures. J Phycol 52:174-183.

https://doi.org/10.1111/jpy.12389.

54. Nishimura T, Hariganeya N, Tawong W, Sakanari H, Yamaguchi H, Adachi M. 2016.

Quantitative PCR assay for detection and enumeration of ciguatera-causing dinoflagellate

Gambierdiscus spp. (Gonyaulacales) in coastal areas of Japan. Harmful Algae 52:11-22.

https://doi.org/10.1016/j.hal.2015.11.018.

55. Gillooly JF, Allen AP, Brown JH, Elser JJ, del Rio CM, Savage VM, West GB, Woodruff

WH, Woods HA. 2005. The metabolic basis of whole-organism RNA and phosphorus

content. Proc Natl Acad Sci U S A 102:11923. https://doi.org/10.1073/pnas.0504756102.

56. Weber AAT, Pawlowski J. 2013. Can abundance of protists be inferred from sequence data:

A case study of Foraminifera. PLoS One 8:e56739.

https://doi.org/10.1371/journal.pone.0056739.

57. Prescott DM. 1994. The DNA of ciliated protozoa. Microbiol Rev 58:233.

https://doi.org/10.1128/mmbr.58.2.233-267.1994.

58. Cavalier-Smith T. 1978. Nuclear volume control by nucleoskeletal DNA, selection for cell

volume and cell growth rate, and the solution of the DNA C-value paradox. J Cell Sci

34:247.

59. Elser JJ, Sterner RW, Gorokhova E, Fagan WF, Markow TA, Cotner JB, Harrison JF, Hobbie

SE, Odell GM, Weider LW. 2000. Biological stoichiometry from genes to ecosystems. Ecol

Lett 3:540-550. https://doi.org/10.1111/j.1461-0248.2000.00185.x.

60. Weiss RL, Kukora JR, Adams J. 1975. The relationship between enzyme activity, cell

geometry, and fitness in *Saccharomyces cerevisiae*. Proc Natl Acad Sci U S A 72:794-798.

https://doi.org/10.1073/pnas.72.3.794.

61. Mable BK, Otto SP. 2001. Masking and purging mutations following EMS treatment in

haploid, diploid and tetraploid yeast (*Saccharomyces cerevisiae*). Genet Res 77:9-26.

https://doi.org/10.1017/S0016672300004821.

62. Hessen DO, Jeyasingh PD, Neiman M, Weider LJ. 2010. Genome streamlining and the

elemental   costs   of   growth.   Trends   Ecol   Evol   25:75-80.
https://doi.org/10.1016/j.tree.2009.08.004.

63.   Zhao F, Filker S, Xu K, Li J, Zhou T, Huang P. 2019. Effects of intragenomic polymorphism
in the SSU rRNA gene on estimating marine microeukaryotic diversity: A test for ciliates
using single-cell high-throughput DNA sequencing. Limnol Oceanogr Methods 17:533-543.
https://doi.org/10.1002/lom3.10330.

64.   Alverson AJ, Kolnick L. 2005. Intragenomic nucleotide polymorphism among small subunit
(18S) rDNA paralogs in the diatom genus *Skeletonema* (Bacillariophyta). J Phycol
41:1248-1257. https://doi.org/10.1111/j.1529-8817.2005.00136.x.

65.   Caron DA, Hu SK. 2019. Are we overestimating protistan diversity in nature? Trends
Microbiol 27:197-205. https://doi.org/10.1016/j.tim.2018.10.009.

66.   Song W, Joo M, Yeom J-H, Shin E, Lee M, Choi H-K, Hwang J, Kim Y-I, Seo R, Lee JE,
Moore CJ, Kim Y-H, Eyun S-i, Hahn Y, Bae J, Lee K. 2019. Divergent rRNAs as regulators
of   gene   expression   at   the   ribosome   level.   Nat   Microbiol   4:515-526.
https://doi.org/10.1038/s41564-018-0341-1.

67.   Van de Peer Y, Mizrachi E, Marchal K. 2017. The evolutionary significance of polyploidy.
Nat Rev Genet 18:411-424. https://doi.org/10.1038/nrg.2017.26.

68.   Mentewab AB, Jacobsen MJ, Flowers RA. 2011. Incomplete homogenization of 18S
ribosomal DNA coding regions in *Arabidopsis thaliana*. BMC Res Notes 4:93.
https://doi.org/10.1186/1756-0500-4-93.

69.   Stoltzfus A, Norris RW. 2015. On the causes of evolutionary transition: Transversion bias.
Mol Biol Evol 33:595-602. https://doi.org/10.1093/molbev/msv274.

70.   Long H, Doak TG, Lynch M. 2018. Limited mutation-rate variation within the *Paramecium*
aurelia   species   complex.   G3:   Genes|Genomes|Genetics   8:2523-2526.
https://doi.org/10.1534/g3.118.200420.

71.   Benítez L, Gutiérrez JC. 1997. Encystment - specific mRNA is accumulated in the resting
cysts   of   the   ciliate   *Colpoda   inflata*.   IUBMB   Life   41:1137-1141.
https://doi.org/10.1080/15216549700202221.

72.   Palacios G, Martin-Gonzalez A, Gutierrez JC. 1994. Macronuclear DNA demethylation is

1083   involved in the encystment process of the ciliate *Colpoda inflata*. Cell Biol Int 18:201-206.

1084   https://doi.org/10.1006/cbir.1994.1062.

1085

 Figure legends

Figure 1. Growth and morphology of the soil ciliates *Colpoda steinii* and *C. inflata* reared at 18 ℃
and 28 ℃. **(A)** Growth curves of *C. steinii* (arrows and arrowheads in red) and *C. inflata* (arrows
and arrowheads in blue), depicting cell isolation time-points at lag, log, plateau and resting cyst
phases. **(B)** Maximum growth rates of cells at log phase. Treatments not sharing common letters
indicate significant differences ($P < 0.05$). **(C)** Morphology of vegetative cells, unstable cysts (UC),
dividing cysts (DC) and resting cysts (RC) from life. **(D)** Microphotographs of fixed specimens
after protargol impregnation. Cell and macronuclear sizes across life-cycle stages, temperatures and
species. Scale bars = 20 μm.

Figure 2. Variation in phenotypic traits of *Colpoda steinii* and *C. inflata* across life-cycle stages and at two temperatures. Unconnected points in line graph panels indicate unstable cysts. (**A**) Cell volume progressively decreased from lag to resting cyst phases. In general, unstable cysts were larger than resting cysts. Cells at a given growth phase were consistently larger at 18 ℃ than at 28 ℃. (**B**) Equivalent spherical diameter (ESD) progressively decreased from lag to resting cyst phases. (**C**) Macronuclear volume changed similarly to cell volume across stages, but was mostly not significant between the two temperatures. (**D**) Ratio of macronuclear to cell volume (MV:CV) across life-cycle stages. (**E**) A linear log-log relationship between macronuclear and cell volumes across life-cycle stages. (**F**) A modeled scaling relationship between MV:CV ratio and ESD. Nucleocytoplasmic ratio varies dramatically in small cells [e.g., pico- (< 2 μm) and nano-sized protists (2 ~ 20 μm)], but changes only slightly in large cells [e.g., microplanktons (20 ~ 200 μm)]. Non-sharing of common letters indicates significant differences ($P < 0.05$). RC = resting cyst; UC = unstable cyst.

Figure 3. Variation in ribotypic traits of *Colpoda steinii* (A, C and E) and *C. inflata* (B, D and F) across lag, log, plateau, resting (RC) and unstable cyst (UC) stages and at two culturing temperatures. Unconnected points in line graph panels indicate unstable cysts. (**A, B**) Single-cell 18S rDNA and rRNA copy numbers progressively and significantly decreased from lag to RC phase. Cells in the UC stage generally had similar amounts of rDNA and rRNA as the vegetative cells in plateau phase. (**C, D**) Cellular concentrations of 18S rDNA and rRNA molecules were generally stable across lag, log, plateau and UC phases, but rRNA concentrations were significantly reduced in resting cysts of both species. (**E, F**) Copy number ratios of rRNA to rDNA in single cells were significantly lower in resting cysts of both species. The ratio was consistent in all other life-cycle phases of *C. steinii*, and lower in plateau and unstable cystic phases of *C. inflata* (relative to its log phase). Bars represent standard errors. Treatments not sharing common letters indicate significant differences ($P < 0.05$).

1122
Figure 4. Scaling relationships among ribotypic traits, growth rate, and cellular attributes. **(A, B)**
The per-cell rDNA and rRNA copy numbers (CNs) in two *Colpoda* species across life-cycle stages
was significantly related to cell volume **(A)** and macronuclear volume **(B)**. Nevertheless, the rRNA
CNs appeared to be much lower than similarly sized vegetative cells or unstable cysts, such that the
coefficients of determination (R^2) of regressions were much lower with resting cysts being taken
into account (+RC) than without (-RC). **(C)** Scaling relationships based on a combined dataset
consisting of two *Colpoda* species (present study), *Euplotes vannus* and *Strombidium sulcatum*, and
five prokaryotic species (Fu and Gong 2017). The per-cell rRNA CNs are better predicted using cell
volume when resting cysts (indicated by four symbols of solid diamond in blue) are excluded from
the regression analysis (-RC). The rDNA CNs of *E. vannus* and *S. sulcatum* were much higher than
those of *Colpoda* cells with similar size ranges, suggesting that rDNA CN and CV relationship may
be conserved among closely related taxa, but becomes inconsistent among distant taxa. **(D)**
Maximum growth rates (*r*) of *Colpoda steinii*, *C. inflata*, *E. vannus* and *S. sulcatum* can be well
predicted by rDNA CN per cell (CNPC) and temperature (*t*). Lines in blue denote predicted growth
rates at 0, 10, 20 and 30 ℃. The amount of rDNA copies in genomes is inversely related to the
growth potential of the cell, which might underline r- and K-selection of these organisms. RC =
resting cysts; UC = unstable cysts.

1140

Figure 5. Variation in ribotypic traits based on intra-individual sequence polymorphisms in four ciliate species. **(A - D)** Proportions of the most dominant haplotypes in single-cell rDNA and rRNA pools, showing the differences between resting cysts (RC) and log-phase vegetative cells (Log), and among different culturing temperatures (in °C). **(E, F)** The haplotype richness of rDNA depended on rDNA number within a cell **(E)**, but the haplotype richness of their transcripts was relatively stable and taxon-dependent, regardless of changes in cellular rRNA content **(F)**. Two treatments not sharing common letters indicate significant differences (multiple pairwise comparisons, $P < 0.05$). Log = cells in log phase of growth; RC = resting cysts.

1149

Figure 6. Variation in the numbers of variants and OTUs in rDNA and rRNA (cDNA) pools of single cells of four ciliated protists. **(A)** Comparisons of variant numbers in a single resting cyst (RC) and log-phase cell (Log) of *Colpoda steinii* and *C. inflata* at 18 ℃ and 28 ℃. **(B)** Comparisons of variant numbers in a single log-phase cell of *Euplotes vannus* and *Strombidium sulcatum* at four temperatures. The OTUs were defined at a sequence similarity threshold ranging from 90% to 99%. At the threshold of 97% (the typical OTU definition threshold applied in metabarcoding studies), the single cells of these species had on average > 1 (up to 2.8) OTU(s), except for *S. sulcatum*. The maximum intra-individual rDNA and rRNA sequence differences of the four species ranged from 1% (in *S. sulcatum*) to 9% (in *C. steinii*). Singletons were not included in these analyses. Two treatments not sharing common letters indicate significant differences (multiple pairwise comparisons, $P < 0.05$).

Figure 7. Single nucleotide polymorphisms in rDNA and rRNA (cDNA) fragments of four ciliate species identified using high throughput sequencing. **(A)** Nucleotide replacement frequencies were much higher at certain positions (polymorphic sites), which were consistent between rDNA and rRNA, between the two *Colpoda* species, and between *E. vannus* and *S. sulcatum*. **(B)** Comparisons of nucleotide replacement frequencies of rDNA and rRNA at log-phase cells vs. resting cysts, and at different temperatures of four ciliate species. Two treatments not sharing common letters indicate significant differences (multiple pairwise comparisons, $P < 0.05$).

1169

Figure 8. Distribution of transition (Ts) and transversion (Tv) categories in rDNA and rRNA
(cDNA) variants in resting cyst and log-phase cells of four ciliate species reared at different
temperatures. RC, resting cysts; Log, log phase cells. Two treatments not sharing common letters
indicate significant differences (multiple pairwise comparisons, $P < 0.05$).

 Supplementary Material

Figure S1. Chilling treatments of two *Colpoda* species induced unstable cyst formation. The chilling treatments at 0 ℃ and 4 ℃ showed that cold shock could induce formation of unstable cysts in actively growing *Colpoda* cells. Nevertheless, this effect was strongly dependent on chilling temperature and duration. Putting both *Colpoda* species at 10 ℃ did not induce formation of unstable cysts (data not shown). **(A)** Cell division of *Colpoda steinii* was completely inhibited upon ice water chilling, and nearly 100% of vegetative cells were transformed into unstable cysts in 4 hrs. Longer chilling duration led to a decrease in the number of unstable cysts. **(B)** In contrast, the population of *C. inflata* collapsed completely during ice water treatments within 3 hrs. **(C)** At 4 ℃, the small-sized *C. steinii* continued to grow and divide during the first 1.5 hrs. Afterwards, total cell abundance (= vegetative cells + unstable cysts) remained relatively stable, with approximately 76% of cells being transformed from vegetative status into unstable cysts following incubation for 9 to 12 hrs. Treating cells at 4 ℃ for a longer period of time damaged both types of cell. **(D)** Chilling at 4 ℃ completely inhibited cell division of *C. inflata*; unstable cysts progressively increased to 37% in 10 hrs, after which the number of unstable cysts remained stable for the next 8 hrs. All individuals transformed into unstable cysts in 20 hrs. Approximately, 36% of cells were lost during this incubation.