Identification of proximal SUMO-dependent interactors using SUMO-ID

Orhi Barroso-Gomila[1], Fredrik Trulsson[2], Veronica Muratore[1], Iñigo Canosa[1], Ana Rosa Cortazar[1,5], Coralia Perez[1], Mikel Azkargorta[1,3,4], Ibon Iloro [1,3,4], Arkaitz Carracedo[1,5,6,7], Ana M Aransay[1,3], Felix Elortza[1,3,4], Ugo Mayor[6,7], Alfred C. O. Vertegaal[2], Rosa Barrio[1*], James D. Sutherland[1*].

1. Center for Cooperative Research in Biosciences (CIC bioGUNE), Basque Research and Technology Alliance (BRTA), Bizkaia Technology Park, Building 801A, 48160 Derio, Spain.

2. Cell and Chemical Biology, Leiden University Medical Center (LUMC), 2333 ZA Leiden, The Netherlands

3. CIBERehd, Instituto de Salud Carlos III, C/ Monforte de Lemos 3-5, Pabellón 11, Planta 0, 28029 Madrid, Spain.

4. ProteoRed-ISCIII, Instituto de Salud Carlos III, C/ Monforte de Lemos 3-5, Pabellón 11, Planta 0, 28029 Madrid, Spain.

5. CIBERONC, Instituto de Salud Carlos III, C/ Monforte de Lemos 3-5, Pabellón 11, Planta 0, 28029 Madrid, Spain.

6. Ikerbasque, Basque Foundation for Science, 48011 Bilbao, Spain

7. Biochemistry and Molecular Biology Department, University of the Basque Country (UPV/EHU), E-48940, Leioa, Spain.

23   (*) Corresponding authors: jsutherland@cicbiogune.es, rbarrio@cicbiogune.es

24   R Barrio ORCID: 0000-0002-9663-0669

25   J D Sutherland ORCID: 0000-0003-3229-793X

27   Lead Contact: Rosa Barrio rbarrio@cicbiogune.es

ABSTRACT

The fast dynamics and reversibility of posttranslational modifications by the ubiquitin family pose significant challenges for research. Here we present SUMO-ID, a technology that merges proximity biotinylation by TurboID and protein-fragment complementation to find SUMO-dependent interactors of proteins of interest. We developed an optimized split-TurboID version and show SUMO interaction-dependent labelling of proteins proximal to PML and RANGAP1. SUMO-dependent interactors of PML are involved in transcription, DNA damage, stress response and SUMO modification and are highly enriched in SUMO Interacting Motifs, but may only represent a subset of the total PML proximal proteome. Likewise, SUMO-ID also allowed us to identify novel interactors of SUMOylated SALL1, a less characterized SUMO substrate. Thus, SUMO-ID is a powerful tool that allows to study the consequences of SUMO-dependent interactions, and may further unravel the complexity of the ubiquitin code.

INTRODUCTION

Ubiquitin-like (UbL) proteins belong to a superfamily of small proteins that attach covalently to target substrates in a transient and reversible manner. The UbL family includes Small Ubiquitin-like Modifiers (SUMOs). The mammalian SUMO family consists of at least three major *SUMO* paralogues (*SUMO1*,*-2*,*-3*). Human SUMO2 and SUMO3 share 97% sequence identity, whereas they share 47% of sequence identity with SUMO1[1]. Protein SUMOylation is a rigorously regulated cycle involving an enzymatic machinery that acts in a stepwise manner. Briefly, the C-terminal di-glycine motif of mature SUMOs binds lysines in substrates through the sequential action of E1 SUMO-activating enzyme SAE1/SAE2, E2 conjugating enzyme UBC9 and SUMO E3 ligases[2]. If required, SUMO as well as the substrate can be recycled by the action of SENPs that cleave the isopeptide bond. Like Ub, SUMO has internal lysines that can be further modified, extended as SUMO chains, modified by Ub chains to target degradation, or even modified by smaller moieties, like acetyl groups[3-5]. Together, these constitute the concept of the "SUMO code" and the ongoing challenge is to understand how these modifications drive distinct substrate outcomes and cellular fates.

SUMO plays crucial roles in nuclear processes underlying health and disease such as the DNA damage response, cell cycle regulation, transcription and proteostasis[6]. SUMO is known to control vital biological processes including development[7] and cholesterol homeostasis[8]. Improvements in mass spectrometry technology have led to the identification to date of more than 40,700 SUMO sites within 6,700 SUMO substrates[9]. While cell-wide proteomics approaches can help to understand global SUMO signaling[10], better tools are needed that allow the study of the cause and consequences of particular SUMOylation events for individual substrates.

SUMO can also interact non-covalently with SUMO interacting motifs (SIMs) found in some proteins. SIMs are β strands composed of an hydrophobic core motif that

interacts with the hydrophobic residues of the SIM-binding groove of SUMOs to form an intramolecular β-sheet[11]. A well characterized role of the SUMO-SIM interaction concerns the SUMO-targeted Ub ligases (STUbL). The two described human STUbLs, RNF4 and Arkadia/RNF111, recognize poly-SUMOylated substrates through their SIMs and ubiquitylate them, leading to their proteasomal degradation[12,13]. The SUMO-SIM interaction also plays critical roles in assembling protein complexes: interaction of the SIM1 of Ran Binding Protein 2 (RanBP2) with the SUMOylated version of Ran GTPase-activating protein 1 (RanGAP1) is crucial for the RanBP2/RanGAP1*SUMO1/UBC9 E3 ligase complex formation[14].

Another intriguing function of SUMO-SIM interaction is the targeting of proteins to Promyelocytic Leukemia Nuclear Bodies (PML NBs). PML NBs are membrane-less ring-like protein structures found in the nucleus. They are bound to the nuclear matrix, make contacts with chromatin fibers[15] and associate with transcriptionally active genomic regions[16]. They consist of a shell composed of PML proteins that surround an inner core in which client proteins localize. Due to the heterogeneity of client proteins, PML NBs have diverse nuclear functions (reviewed in[17,18]). The *PML* gene contains 9 exons and numerous splicing variants. All PML isoforms contain the N terminal TRIpartite Motif (TRIM) that is responsible for PML polymerization and NB formation[19], binding to Arsenic Trioxide (ATO)[20] and may act as an oxidative stress sensor[18]. PML also contains a phospho-SIM located at its exon 7 and shared by most PML isoforms[21]. Almost all PML isoforms contain three putative SUMO sites: K65, K160 and K490. PML SUMOylation is a well characterized signal for RNF4-mediated ubiquitylation and degradation[22].

Proximity-dependent labelling methods are based on promiscuous labeling enzymes that produce reactive molecules that covalently bind neighbor proteins. Labeled proteins can be then purified and identified using affinity-purification coupled to mass

spectrometry methods[23]. Proximity-dependent biotin identification (BioID)[24] uses a

promiscuously active *Escherichia coli* biotin ligase (BirA*) generated by a point mutation

(R118G) to biotinylate lysines in nearby proteins within an estimated range of 10 nm[25].

By fusing BirA* to specific proteins, BioID efficiently identifies interactors at

physiological levels in living cells[26] . It has been extensively used in the Ub field, for

instance, to identify substrates of E3 ligases[27,28]. Recently, a more efficient version of

BioID, termed TurboID, has been developed[29], being this more suitable for transient

protein-protein interaction (PPI) detection. Several studies have developed split-versions

and applied "protein fragment complementation" to BioID and TurboID, where proximal

biotinylation is dependent on the proximity of the fusion partners, opening new

opportunities for spatial and temporal identification of complex-dependent

interactomes[30,31].

To study how SUMOylation and SUMO-SIM interactions can lead to other roles

and fates for particular substrates poses particular challenges. SUMOylation occurs

transiently and often in a small percentage of a given substrate. Modified proteins can be

readily deSUMOylated and SUMO can be recycled and passed to other substrates.

SUMO-SIM interactions are also difficult to analyze due to their weak affinity (Kd 1-100

µM). To overcome those technical issues, we developed SUMO-ID, a new strategy based

on Split-TurboID to identify interactors of specific substrates dependent on SUMO

conjugation or interaction. Using PML as a model, we demonstrate that SUMO-ID can

enrich for factors that depend on PML-SUMO interaction. Importantly, the identified

proteins are represented among proximal interactors of PML identified using full-length

TurboID. We also applied SUMO-ID to a less-characterized SUMO substrate, Spalt Like

Transcription Factor 1 (SALL1), and identified both known and novel interactors that

depend on intact SUMOylation sites in SALL1. SUMO-ID is thus a powerful tool to study

transient and dynamic SUMO-dependent interaction events. The developed methodology

is generic and therefore widely applicable in the Ub and UbL field to identify readers of

these modifications for individual target proteins to improve our insight in non-covalent

signal transduction by Ub and UbL.

RESULTS

Identification of SUMO-dependent interactions: the SUMO-ID strategy

We posited that Split-TurboID, in which one fragment is fused to SUMO and the

complementary fragment to a protein of interest, could identify transient SUMO-

dependent interactors (Fig. 1). Upon covalent SUMOylation or non-covalent SUMO-SIM

interaction, both fragments are brought together, presumably close enough to allow

refolding of the TurboID enzyme. In the presence of biotin, the reconstituted TurboID

can then label proximal complexes, which can be purified by streptavidin pull-down and

identified by liquid chromatography-mass spectrometry (LC-MS). Due to the high

affinity of streptavidin-biotin interaction, harsh cell lysis and stringent washes that

significantly reduce unspecific protein binding can be applied. We named this approach

"SUMO-ID".

T194/G195 Split-TurboID enables SUMO-ID studies

We applied the previously described E256/G257 BioID split-point[30] to TurboID,

but found it unsuitable for SUMO-ID. While SUMO-dependent reconstitution of the

E256/G257 was observed in our pilot experiments, the NTurboID[256]-fusions had a

significant background biotinylating activity (Figure S1). This is likely due to residual

biotin binding and activation by the intact NTurboID[256] biotin-binding pocket. We

examined the BirA structure to identify a new TurboID split-point that would yield two

completely inactive fragments (see File S1). The biotin binding pocket of BirA is

composed of three β-strands (strands 5, 8 and 9), the N-terminus of helix E and the 110-

128 loop (Fig. 2A). We split TurboID at T194/G195, so that the resulting NTurboID[194]

fragment (NTbID) carries the principal 110-128 biotin binding loop and the β-strands 5

and 8, while the CTurboID[195] fragment (CTbID) carries the β-strand 9 necessary to the

formation of the biotin binding β-sheet.

We tested T194/G195 Split-TurboID for SUMO-ID. CTbID-SUMOs were

incorporated into substrates in a very efficient manner (Fig. 2B). NTbID and CTbID alone

were catalytically inactive and yielded no biotinylation. Combining NTbID-substrate and

CTbID-SUMOs resulted in a high-yield biotinylation activity of TurboID after 16 hours

of biotin exposure. Modification of NTbID-substrates by CTbID-SUMOs (Fig. 2B,

FLAG blot, black arrowheads) and its corresponding biotinylation activity (Fig. 2B,

biotin blot, black arrowheads) were efficiently detected, notably in the case of PML

protein. Free biotinylated CTbID-SUMOs, that might come from recycling of previously

labeled moieties, were observed (Fig. 2B, biotin blot, white squares). We also examined

by immunofluorescence, and confirmed that the streptavidin signal recognizing the

biotinylated substrates is dependent on fragment-complementation (Fig. 2C). Thus,

T194/G195 Split-TurboID biotinylation activity is dependent on fragment

complementation, with reduced or no leaky biotinylation of the two fragments, so it could

be useful for SUMO-ID and for studying other protein-protein interactions.

We applied the rapamycin-inducible dimerization system, based on FKBP (12-

kDa FK506-binding protein) and FRB (FKBP-rapamycin-binding domain)[32], which has

been used previously to evaluate the PPI dependency of Split-BioID reconstitution[30]. We

fused NTbID and CTbID to FRB and FKBP, respectively, and stably expressed the

constructs in HEK293FT cells. We tested short and long rapamycin treatments, together

with short biotin labeling times, to evaluate self-biotinylation activity of the reconstituted

TurboID. As previously shown, the NTbID, even though it contains the principal biotin

binding 110-128 loop, was catalytically inactive after 24 hours of biotin treatment (Fig.

2D). We observed that biotinylation activity of the reconstituted TurboID correlated well

with rapamycin and biotin treatments, showing its dependency on PPI and biotin labeling

times. 24 hours of rapamycin treatment led to a 16-fold higher FKBP/FRB PPI dependent

biotinylation activity at 2 hours of labeling time. Altogether, these data demonstrate that

T194/G195 Split-TurboID fragments have low intrinsic affinity and high biotinylation

activity at short biotin labeling times, making them suitable for SUMO-ID.

SUMO-ID detects both covalent and non-covalent SUMO-dependent interactions

using short biotin-labeling times

Interaction of a protein with SUMO can be via covalent SUMOylation or non-

covalent SUMO-SIM interaction. We used PML, which can both be SUMOylated and

has a well-characterized SIM domain, in conjunction with SUMO wild type (WT) or

mutants that lack the C-terminal di-glycine (ΔGG) necessary for covalent conjugation.

We used a stable HEK293FT cell line expressing NTbID-PML, into which CTbID-

SUMOs were transfected, using short biotin-labeling times (0.5-2 hours). We observed

that CTbID-SUMO1/2 transfections led to high SUMO-dependent biotinylation activity

after only 2 hours of biotin treatment (Fig. 3A, biotin blot, black arrowhead).

Additionally, ATO treatment, which induces PML SUMOylation, further enhanced the

SUMO-dependent biotinylation. With 30 minutes of biotin treatment, CTbID-

SUMO1/2$^{\Delta GG}$ induced biotinylation of unmodified NTbID-PML, likely through SUMO-

SIM interactions (Fig. 3A, biotin blot, white arrowhead). With longer biotin labeling (2

hours), biotinylation of endogenous SUMO modified NTbID-PML was also detected,

more strongly in the case of ATO treatment (Fig. 3A, biotin blot, black arrowhead).

Biotinylated free CTbID-SUMO1/2$^{\Delta GG}$ were detected, while the WT counterparts were

not biotynilated at 2 hours (Fig. 3A, biotin blot, white squares), supporting that recycling

of biotinylated SUMOs may be linked to longer labeling times. Indeed, additional

experiments showed that appearance of free biotinylated CTbID-SUMOs increased with

longer labelling times (Figure S2). Altogether, these results demonstrate that SUMO-

dependent biotinylation activity for specific targets, especially at short biotin labeling

times, may be a useful strategy for identifying specific SUMO-dependent interactors of

those proteins.

Reduced labeling times and lower expression levels reduced SUMO recycling,

but still allowed some degree of recycling (and therefore loss-of-specificity) to occur, so

we incorporated two further modifications. First, we designed SUMO isopeptidase-

resistant versions of CTbID-SUMOs (SUMO non-cleavable, or SUMOnc, Fig. 1B)[33].

This could also reduce the target identification derived from SUMO-SIM interactions

involving free unincorporated CTbID-SUMOs, since most CTbID-SUMOncs would be

incorporated into substrates. The same strategy was applied to Ub (CTbID-Ubnc).

Secondly, we transferred non-cleavable CTbID-SUMOs into pTRIPZ, an all-in-one

doxycycline-inducible (Dox) lentiviral vector (Fig. 1B). Regulated expression would

offset any deleterious effects stemming from non-cleavable SUMO isoforms, and provide

useful experimental control (i.e. non-induced *versus* induced). Inducible TRIPZ-CTbID-

SUMOnc showed enhanced SUMOylation compared to the constitutive WT SUMO

versions (Fig. 3B). Free non-incorporated versions of SUMO1/2nc and Ubnc were not

detectable (Fig. 3B, MYC blot, white squares). Stable cell populations were established

(HEK293FT, U2OS and RPE-1 cells) for each (SUMO1nc, SUMO2nc, Ubnc). Validation

of TRIPZ-CTbID-SUMO2nc by WB and immunofluorescence is shown (Figure S3). We

then introduced constitutively-expressed NTbID-PML into TRIPZ-CTbID⁻SUMO1nc or

-SUMO2nc cells and proved that biotinylation occurs in PML NBs as expected, in a

doxycycline dependent manner (Figure S4). These data show that the use of regulated

SUMOnc versions leads to both high activity and specificity needed for the SUMO-ID

approach.

To further validate the specificity of SUMO-dependent biotinylation activity with

PML, we generated control cells carrying NTbID-PML[3MAS], a mutated version of PML

lacking the three principal SUMOylation sites (K65, K160 and K490) and the best-

characterized SIM domain. While strong SUMO-ID biotinylation activity was observed

with the WT version of PML, this biotinylation activity was completely abrogated in the

case of PML[3MAS] (Fig. 3C). This lack of biotinylation activity was specific to SUMO, as

ubiquitylation-dependent biotinylation activity was observed in TRIPZ-CTbID-Ubnc /

NTbID-PML[3MAS] double stable cell line (Figure S5). NTbID-PML[WT] forms true NBs,

while NTbID-PML[3MAS] forms NB-like bodies, as reported previously[34] (Fig. 3D).

Biotinylation driven by SUMO-ID was observed in NBs containing NTbID-PML[WT], and

it was enhanced after 2 hours of ATO treatment, but not in the NB-like structures

containing NTbID-PML[3MAS]. Thus, these results show that SUMO-ID biotinylation

activity is dependent on substrate-SUMO interaction.

SUMO-ID identifies SUMO-dependent interactors of PML

Since PML NBs are known hubs of SUMO-dependent signaling[17,18], we

wondered which interactions in NBs via PML are SUMO-dependent, so we performed

SUMO-ID using NTbID-PML[WT] compared to NTbID-PML[3MAS], each combined with

TRIPZ-CTbID-SUMO2nc. Biotinylated proteins were purified by streptavidin pull-down

and sequenced by LC-MS (Table S1). 59 high-confidence SUMO-dependent interactors

of PML were enriched in PML[WT] SUMO-ID compared to PML[3MAS] SUMO-ID (Fig. 4A).

Among those, SUMO E3 ligases (PIAS1, PIAS2, PIAS4, TRIM28), transcriptional

regulators (TRIM22, TRIM24, TRIM33, GTF2I, IRF2BP2, IFI16, ZNF280B, MED23,

MEF2D, SNW1, RPAP3), and DNA repair proteins (RMI1, BLM, SLX4, XAB2) were

identified. Of note, PIAS1 is known to induce PML SUMOylation[35] and SUMO-SIM

interaction of BLM is necessary for its targeting to PML bodies[36], which highlights the

specificity of the SUMO-ID strategy. Of particular interest, GTF2I and IRF2BP2,

identified here by SUMO-ID, form fusion proteins with RARA and cause Acute

Promyelocytic Leukaemia (APL, see Discussion)[37,38]. We validated these two proteins,

as well as TRIM33 and UBC9, as SUMO-dependent interactors of PML by WB (Fig.

4B).

STRING networking of SUMO-dependent interactors of PML shows a highly

interconnected cluster related to protein SUMOylation, DNA damage response and

transcriptional regulation (Fig. 4C), while GO enrichment also highlighted protein

SUMOylation and transcriptional regulation, as well as DNA repair and stress response

pathways (Fig. 4D; Table S2). Collectively, this data show that the SUMO-ID strategy

can efficiently identify SUMO-dependent interactors of PML, and that SUMO interaction

with PML reinforces essential processes.

PML SUMO-ID hits localize to PML NBs

We checked whether some of the SUMO-dependent interactors of PML localize

to NBs. We generated a YFP-PML cell line by inserting YFP into the endogenous PML

locus in U2OS cells (Figure S6), and looked for co-localization of selected SUMO-

dependent PML interactors by confocal microscopy. Within individual cells, we observed

frequent and multiple co-localization events for PIAS4, TRIM24, TRIM33 and UBC9 in

PML NBs (Fig. 5), whereas PIAS2, GTF2I and IRF2BP2 colocalizations were less

frequent, suggesting heterogeneity in PML NB composition that may depend on different

factors (including, but not limited to SUMOylation density, subnuclear localization, cell

cycle stage, other PTMs, or contrastingly, technical limitations with antibodies or

fixations).

SUMO-dependent interactions are a subset of PML proximal proteome

PML NBs are membraneless structures thought to behave as phase-separated liquids and with high heterogeneity in composition[39]. These characteristics make their purification very challenging, and no proteomic data are nowadays available. Therefore, to compare the obtained PML SUMO-ID specific sub-proteome with the regular PML interactome, we decided to characterize a comprehensive PML and PML^{3MAS} proximity interactome using standard full-length TurboID (FLTbID). We generated stable U2OS cell lines for FLTbID-PML^{WT}, FLTbID-PML^{3MAS} and FLTbID alone, and treated them or not with ATO to induce PML SUMOylation. High confidence PML proximal proteome was composed of 271 proteins that were enriched in FLTbID-PML^{WT} samples compared to FLTbID alone (Fig. 6A, Table S3). STRING networking showed a main core cluster composed of 73.6% of the identified proteins (Fig. 6B). The most representative subclusters were composed of 1) RNA splicing and mRNA processing proteins, 2) transcription, RNA biosynthesis and DNA damage response proteins and 3) replication and SUMOylation related proteins. This largely aligned with Gene Ontology (GO) enrichment analysis, which revealed that PML proximal interactors participate in replication, transcription, RNA splicing, DNA damage response, cell cycle regulation, SUMOylation and ubiquitylation, and telomere maintenance, consistent with fact that PML in U2OS regulates the ALT mechanism[40] (alternative lengthening of telomeres; Fig. 6C, Table S4).

SUMOylation of PML is thought to be a controlling factor for composition and dynamics of NBs. Are all NB interactions linked to PML dependent on SUMO? To answer this question, we subdivided the PML interactome into SUMO-dependent or -independent interactors, by comparing FLTbID-PML^{WT} and FLTbID-PML^{3MAS} samples. We observed some proteins that likely localize to PML NBs[41], such as NCOR-1, STAT3,

JUN, BRCA2 and HDAC9, were also enriched in TurboID-PML^{3MAS}, suggesting SUMO-

independent targeting to PML NBs (Table S3). Importantly, many of SUMO-dependent

interactors identified by SUMO-ID are part of SUMO-dependent PML NBs proteome

using standard TurboID, including PIAS2, PIAS4, TRIM24, TRIM33 and IRF2BP2 (Fig.

7A; Table S3), supporting the validity of SUMO-ID to identify SUMO-dependent

interactors. Interestingly, scores of some PML interactors decreased after ATO treatment

(TRIM24, TRIM33, SENP5), suggesting that those proteins may rapidly undergo

dissociation or degradation in response to PML SUMOylation. We confirmed such effect

for TRIM24 by WB (Fig. 7B). Altogether, these data confirm that SUMO-ID identified

hits are a subset of the SUMO-dependent PML proximal proteome.

SUMO-dependent interactors of PML are enriched in SIMs

We expected that many of the SUMO-dependent PML interactors might do so via

SUMO-SIM interactions and, therefore, should contain or be enriched in SIMs. To test

this, we designed and executed an *in-silico* SIM enrichment analysis. We generated 1000

random lists of 59 proteins (the size of the SUMO-ID identified protein list) and evaluated

the presence of SIMs (Table S5). The median of single SIM and multiple SIM presence

in the random lists were 45.76% and 23.73% respectively (Fig. 7C). SUMO-ID identified

proteins showed a much higher content of SIMs, with single SIM and multiple SIM

presence values of 96.61% and 89.83% respectively. It is noteworthy that around 83% of

the identified SIMs in PML SUMO-ID list were preceded or followed within the first 4

amino-acids by acidic residues (D, E) or a Serine. Since longer proteins are expected to

have more SIMs, we then normalized the SIM content with the size of proteins on the

lists to obtain the value of "SIMs per thousand of amino acids" (STAA) (Table S5). The

values obtained with the random lists showed a Gaussian distribution (d'Agostino and

Pearson normality test, K2 value 3.836, *p*-value 0.15) (Fig. 7D). The median of the values

obtained with the random lists was 4.85 (Log2 = 2.28) STAA, while for PML SUMO-ID

list was 18.42 (Log2 = 4.20) STAA, which translates to a SIM enrichment value of 3.8

times higher than the random lists (empirical *p*-value < 0.001). These results show that

SUMO-dependent interactors of PML are highly enriched in SIMs.

SUMO-ID identifies interactors of SUMOylated SALL1

324   To test the sensitivity and discovery potential of the SUMO-ID, we applied this

technique to SALL1, a transcriptional repressor that is SUMOylated[42,43], but of which

nothing is known about the causes or consequences of this modification. Using TRIPZ-

CTbID–SUMO1nc or SUMO2nc HEK293FT stable cell lines, we introduced NTbID–

SALL1^{WT} or SALL1$^{\Delta SUMO}$ (with mutations in 4 major SUMOylation consensus sites) and

evaluated SUMO-ID by WB. Efficient SUMO-ID biotinylation activity was observed

when using SUMO2nc (Fig. 8A, black arrowhead). NTbID–SALL1^{WT} localizes to the

nucleus, forming nuclear bodies with high SUMO-ID activity, and NTbID–SALL1$^{\Delta SUMO}$

also forms aggregates in the cytoplasm (Fig. 8B). Specificity of SALL1 SUMO-ID was

confirmed in cells, as biotinylation occurs only in SALL1^{WT} upon doxycycline induction

and biotin supplementation. SALL1 SUMO-ID identified potential SUMO-dependent

interactors of SALL1 such as the transcription factors TLE3, DACH1/2 and ARID3B, as

well as NuRD complex proteins GATAD2A/B, MTA1/2 and RBBP4/7 (Fig. 8C; Table

S6), already known as SALL1 interactors[44]. We also identified components of the

SUMOylation machinery, such as PIAS1. We confirmed that SUMOylated SALL1 was

biotinylated and purified via SUMO-ID (Fig. 8D, black arrowheads) as well as NuRD

complex proteins GATAD2B, MTA2 and RBBP4 (Fig. 8D). MCODE subclustering of

the STRING interaction network showed a highly interconnected cluster composed of

NuRD complex proteins (Fig. 8E) that was also enriched as GO term (*p*-value $2.40 \cdot 10^{-4}$).

Thus, SUMO-ID is sensitive and specific, allowing the study of SUMO-dependent

interactors for proteins of interest, opening new avenues of understanding how SUMO

can affect their function.

DISCUSSION

The fast dynamics and reversibility of SUMOylation, and the low affinity of

SUMO-SIM interactions pose significant challenges not only for SUMO research, but for

respective studies of Ub and other UbLs. The use of His-tagged K0-SUMO to isolate

substrates and map SUMOylation sites has been instrumental to show the widespread

presence of this modification in the human proteome[9,45]. Direct purification of

SUMOylated proteins using immunoprecipitation is a gold standard and can be applied

to cells and tissues[46], but is also challenging because SUMOylation might affect a small

proportion of low abundance proteins, and perhaps only under certain conditions (e.g a

discrete cell cycle phase or upon DNA damage). Recently, the NanoBiT-based ubiquitin

conjugation assay (NUbiCA) was described that uses a split-luciferase approach to allow

a quantitative assessment of Ub-modified proteins[47]. Bimolecular fluorescence

complementation (BiFC) approaches employ a split fluorescent protein that enables the

localization of UbL-modified proteins in yeast or human cells to be monitored[48-50]. If

applied to UbLs and substrates, the BiCAP method[51], which allows purification of

reconstituted GFP using GFP nanobodies, could likely enrich modified substrates and

perhaps interactors. However, none of these approaches captures the dynamic

environment of specific UbL-modified proteins, often characterized by weak and

transient interactions.

Here we describe SUMO-ID, a powerful technique that allows the study of the

causes and consequences of SUMO-dependent interactions for specific proteins of

interest. The fast biotinylation activity of TurboID and the specificity obtained with

"protein fragment complementation" permit SUMO-ID to specifically biotinylate

interactors of substrates in a SUMO-dependent manner. Combined with sensitive

proteomic methods, SUMO-ID allows the identification of specific interactors,

potentially revealing enzymatic machinery responsible for the SUMOylation as well as

interactors that may be stabilized or recruited as a consequence of the modification. Like

approaches using BiFC, the subcellular localization of SUMO-modified substrates using

SUMO-ID can also be inferred, through the use of fluorescent streptavidin. However,

caution should be taken with the mentioned factors in order to maintain specificity, such

is the use of non-cleavable forms of UbLs or the application of short biotin labelling

times. This strategy might compromise the identification of SUMO isopeptidases since

their binding to SUMOylated substrate is likely affected.

At the core of SUMO-ID is Split-TurboID, which individual halves should ideally

have no activity, as with all split-protein approaches. For SUMO-ID, we initially applied

the E256/G257 split point described for Split-BioID[30] to the fast-labelling TurboID

derivative, but found that the N-terminal half (1-256) retained substantial biotinylation

capacity. We speculate that this is because the "biotin pocket" is still intact and might

allow leaky release of biotinoyl-AMP. Leaky biotinylation of TurboID 1-256 was also

observed by Cho and colleagues in their recently published report on Split-TurboID[31].

Their final design was based on a L73/G74 split point which showed efficient proximity-

dependent reconstitution and biotinylation, but still leaves the biotin pocket intact in C-

terminal 74-321 half, opening the possibility of leaky biotinylation during longer labelling

times or in stable cell lines. To avoid this, we developed and validated T194/G195 Split-

TurboID that separates the β-strands 5 and 8 from the β-strand 9, completely abrogating

any residual biotinylation activity of the fragments.

Here we used SUMO-ID to unravel the role of PML SUMOylation in PML NBs

function. We identified 59 proteins as SUMO-dependent PML interactors that participate

in essential nuclear processes such as protein SUMOylation, transcriptional regulation, DNA repair and stress response. There is growing evidence that PML interaction with SUMO might allow partners to localise into PML NBs through SUMO-SIM interactions[36,52]. We demonstrated that most of the proteins identified by SUMO-ID are indeed part of the proteome of SUMO-dependent PML NBs and that they are enriched in SIMs, suggesting SUMO-SIM interaction dependency. It has been proposed that, after such partner recruitment, proteins might undergo SUMOylation by the PML NB-localized SUMO machinery that reinforces their sequestration[53]. In fact, PML NBs are, together with the nuclear rim, the major targets of active SUMOylation[54]. Our data reinforce this enzyme/substrate co-concentration model as we observed that SUMOylation machinery enzymes (UBC9, PIAS1, PIAS2, PIAS4 and TRIM28) localize to PML NBs in a SUMO-dependent manner and 80% of the SUMO-dependent PML interactors (47 out of 59) are SUMO substrates[9,10].

To compare our list of SUMO-dependent *versus* general interactors of PML, we performed a TurboID assay for PML, with cells alone or treated with ATO, and identified 271 proteins. ATO induces PML NB formation, subsequent PML SUMOylation, partner recruitment and finally PML degradation[22,55]. It is used to treat APL, a type of Acute Myelocytic Leukaemia (AML), which is mainly caused by the t(15;17) translocation that fuses PML to RARA. Interestingly, two of our SUMO-ID hits, IRF2BP2 and GTF2I, also form fusion proteins with RARA and cause APL, albeit less commonly than PML fusions[37,38]. We validated that both localize to PML NBs. While many of the SUMO-ID candidates show increased peptide intensity in PML NBs after ATO treatment, we observed that some of them decreased. IRF2BP2 and TRIM24, which has also been linked to AML[56,57], showed reduced levels after ATO treatment, suggesting that they might undergo degradation. In line with this idea, the 11S proteasome components are recruited

into mature PML NBs and their localization is enhanced with ATO treatment[58],

suggesting that mature PML NBs may also act as proteolytic sites. In fact, inhibition of

ubiquitylation accumulates SUMOylated proteins within PML NBs[59], suggesting that

many clients that are targeted to PML NBs and that are SUMOylated, might undergo

ubiquitylation and degradation. Altogether, these data provide further insight into the role

of PML SUMOylation in NB biology and open new ways of looking at the mechanisms

of ATO in APL treatment.

The successful application of SUMO-ID to SALL1, a poorly characterized SUMO

substrate, illustrates the sensitivity and utility of SUMO-ID. Although SALL1

SUMOylation levels are vanishingly low under physiological conditions, SUMO-ID

revealed SUMO-dependent enrichment of the NuRD complex proteins GATAD2A/B,

MTA1/2 and RBBP4/7. The association between SALL1, a transcriptional repressor, and

the NuRD complex, a repressive histone deacetylase complex, has been previously

described[44]. The interaction is mediated by an N-terminal 12 amino acid motif of

SALL1[44]. Once recruited, SUMOylation of SALL1 might serve to stabilize the repressor

complex via SUMO-SIM interactions, with predicted SIMs present in multiple NuRD

complex subunits. As histone SUMOylation is also linked to transcriptional repression[60],

SUMO-SIM interactions might further reinforce the SALL1/NuRD complex and drive

histone deacetylation at SALL1 targets. In addition, we also found TLE3, DACH1/2 and

ARID3B transcription factors as SUMO-dependent interactors of SALL1. TLE3, a

transcriptional repressor of the Groucho/TLE family, interacts with HDAC2 (another

NuRD complex component) and can regulate acetylation levels[61]. Both TLE3 and the

tumor suppressor DACH1 are negative regulators of Wnt signaling[62,63]. Interestingly,

SALL1 has been shown to enhance Wnt signaling[64]. Perhaps interaction with

SUMOylated SALL1 serves to counteract these negative effects.

In summary, we demonstrate here that SUMO-ID, based on the 194/195 Split-

TurboID reconstitution, can facilitate the identification of SUMO-dependent interactions

with a protein of interest. It has little or no background, with high biotinylation activity

when expressed at low levels and with short biotin incubation time. We believe that this

technique improves sensitivity and selectivity when applied to infrequent SUMOylation

events and low-affinity of SUMO-SIM interactions. This strategy can be applied to other

UbL modifications (e.g Ub-ID shown in Figure S5), and the 194/195 Split-TurboID may

be useful for other applications in cell biological studies.

METHOD DETAILS

Cell Culture

U2OS (ATCC HTB-96) and HEK293FT (Invitrogen) were cultured at 37°C and 5% CO_2

in Dulbecco's modified Eagle Medium (DMEM) supplemented with 10% fetal bovine

serum (FBS, Gibco) and 1% penicillin/streptomycin (Gibco). Human telomerase reverse

transcriptase immortalized retinal pigment epithelial cells (hTERT-RPE1, ATCC CRL-

4000) were cultured in DMEM:F12 (Gibco) supplemented with 10% FBS, 2 mM L-

Glutamine and 1% penicillin and streptomycin. Cultured cells were maintained through

20 passages maximum and tested negative for mycoplasma.

Cloning

TurboID was a kind gift of A. Ting (Addgene #107171)[29]. PMLIVa[WT] and PMLIVa[3MAS]

were previously described[21]. SUMO1, SUMO2, Ub, RANGAP1 and UBC9 ORFs were

amplified from U2OS cell cDNA by high-fidelity PCR (Platinum SuperFi DNA

Polymerase; Invitrogen). All constructs were generated by standard cloning or by Gibson

Assembly (NEBuilder HiFi Assembly, NEB) using XL10-Gold bacteria (Agilent).

Depending on the construction, plasmid backbones derived from EYFP-N1

(Clontech/Takara), Lenti-Cas9-blast (a kind gift of F. Zhang; Addgene #52962) or TRIPZ

(Open Biosystems/Horizon) were used. After assembly, all vectors were validated by

sequencing. Additional details for constructs are described in Table S7. Cloning details

are available upon request.

Lentiviral transduction

Lentiviral expression constructs were packaged in HEK293FT cells using calcium

phosphate transfection of psPAX2 and pMD2.G (kind gifts of D. Trono; Addgene

#12260, 12259) and pTAT (kind gift of P. Fortes; for TRIPZ-based vectors). Transfection

media was removed after 12-18 hours and replaced with fresh media. Lentiviral

supernatants were collected twice (24 hours each), pooled, filtered (0.45 μm), and

supplemented with sterile 8.5% PEG6000, 0.3 M NaCl, and incubated 12-18 hours at

4°C. Lentiviral particles were concentrated by centrifugation (1500 x g, 45 minutes, 4°C).

Non-concentrated virus (or dilutions thereof) were used to transduce HEK293FT, and 8x

concentrated virus was used for U2OS and hTERT-RPE1 cells. Drug selection was

performed as follows: 1 μg/ml puromycin (Santa Cruz) for U2OS and HEK293FT cells,

5 μg/ml for hTERT-RPE1 cells; 5 μg/ml blasticidin (Santa Cruz) for U2OS, HEK293FT

and hTERT-RPE1 cells.

CRISPR-Cas9 genome editing

Human PML encodes multiple isoforms, but most differ at the 3' end. To target EYFP

into the first coding exon, shared by most PML isoforms, an sgRNA target site was chosen

(CTGCACCCGCCCGATCTCCG) using Broad institute GPP sgRNA Designer[65].

Custom oligos were cloned into px459v2.0 (a kind gift of F. Zhang; Addgene #62988).

A targeting vector was made by amplifying 5' and 3' homology arms using U2OS

genomic DNA, as well as the EYFP ORF (see Table S7 for oligo details). These fragments

were assembled by overlap extension using high fidelity PCR and the resulting amplicon

was TOPO-cloned and sequence-confirmed. Lipofectamine 2000 (Invitrogen) was used

to transfect U2OS with linear targeting vector and px459 encoding SpCas9, puromycin

resistance, and the PML-targeting sgRNA. 24 hours post-transfection, cells were selected

for additional 24-36 hours with 2 μg/ml puromycin. Cells were plated at low density and

clones were examined by fluorescence microscopy. Clones with clear nuclear body

signals were manually picked and expanded. YFP-PML insertions and copy number were

validated by PCR, sequencing, and Western blotting.

Transient transfections and drug treatments

HEK293FT cells were transiently transfected using calcium phosphate method. U2OS

cells were transiently transfected using Effectene Transfection Reagent (Qiagen). After

24 hours of transfection, cells were treated with biotin (50 μM; Sigma-Aldrich) for

indicated exposure times. For stably transduced TRIPZ cell lines, induction with

doxycycline (1 μg/ml; 24 hours; Sigma-Aldrich) was performed prior to biotin treatment.

ATO (1 μM; 2 hours; Sigma-Aldrich) treatments were performed (with or without biotin,

depending on experiment) prior to cell lysis or immunostaining.

Western blot analysis

Cells were washed 2x with PBS to remove excess biotin and lysed in highly stringent

washing buffer 5 (WB5; 8 M urea, 1% SDS in 1X PBS) supplemented with 1x protease

inhibitor cocktail (Roche) and 50 μM NEM. Samples were then sonicated and cleared by

centrifugation (25000 x g, 30 minutes, RT). 10-20 μg of protein was loaded for SDS-

PAGE and transferred to nitrocellulose membranes. Blocking was performed in 5% milk

in PBT (1x PBS, 0.1% Tween-20). Casein-based blocking solution was used for anti-

biotin blots (Sigma). Primary antibodies were incubated over-night at 4°C and secondary

antibodies 1 hour at room temperature (RT). Antibodies used: anti-biotin-HRP (1/1000),

anti-Myc (1/1000), anti-alpha-Actinin (1/5000) (Cell Signaling Technology); anti-Flag

(1/1000), anti-GTF2I (1/1000) (Sigma-Aldrich); anti-BirA (1/1000; SinoBiological);

anti-IRF2BP2 (1/1000), anti-UBC9 (1/1000), anti-TRIM24 (1/1000), anti-TRIM33

(1/1000), anti-PIAS2 (1/1000), anti-PIAS4 (1/1000), anti-GATAD2B (1/1000), anti-

MTA2 (1/1000), anti-RBBP4 (1/1000), anti-PML (1/1000), anti-GAPDH (1/5000), anti-

beta-Actin (1/5000), anti-alpha-Tubulin (1/5000) (Proteintech); anti-PML (1/1000)

(Bethyl); anti-GFP (1/1000) (SantaCruz); anti-Mouse-HRP, anti-Rabbit-HRP (1:5000)

(Jackson ImmunoResearch). Proteins were detected using Clarity ECL (BioRad) or Super

Signal West Femto (ThermoFisher). Quantification of bands was performed using ImageJ

(v2.0.0-rc-69/1.52n) software and normalized against loading controls (GAPDH, actin,

tubulin or alpha-actinin depending on experiments).

Immunostaining and confocal microscopy

U2OS and HEK293FT cells were seeded on 11 mm coverslips (25,000 cells per well; 24

well plate). HEK293FT coverslips were pre-treated with poly-L-lysine. After washing 3

times with 1x PBS, cells were fixed with 4% PFA supplemented with 0.1% Triton X-100

in PBS for 15 minutes at RT. Then, coverslips were washed 3 times with 1x PBS.

Blocking was performed for 30 minutes at RT in blocking buffer (2% fetal calf serum,

1% BSA in 1x PBS). Primary antibodies were incubated for 2 hours at 37°C and cells

were washed with 1x PBS 3 times. Primary antibodies used: anti-BirA (1/500;

SinoBiological); anti-Myc (1/200; Cell Signaling Technology); anti-GTF2I (1/100;

Sigma-Aldrich); anti-IRF2BP2, anti-UBC9, anti-TRIM24, anti-TRIM33, anti-PIAS2,

anti-PIAS4, anti-CBX4 (all 1/100; Proteintech); anti-B23 (1/100) (Santa Cruz); anti-

SC35 (1/100) (BD biosciences); anti-SUMO2/3 (1/100) (DSHB). Then secondary

antibodies (together with fluorescent streptavidin) were incubated for 1 hour at 37°C,

followed by nuclear staining with DAPI (10 minutes, 300 ng/ml in PBS; Sigma Aldrich).

Antibodies used: anti-Rabbit Alexa Fluor 488, anti-Mouse Alexa Fluor 568, anti-Rabbit

Alexa Fluor 568 (all 1/200; ThermoFisher). Streptavidin Alexa Fluor 594 (1/200, Jackson

ImmunoResearch) was used. Fluorescence imaging was performed using confocal

microscopy (Leica SP8 Lightning) with 63x Plan ApoChromat NA1.4. To obtain the

signal histograms for co-localization studies in Fig. 5, we used the plot profile tool in

ImageJ (v2.0.0-rc-69/1.52n).

Pulldown of biotinylated proteins

Cleared lysates from WB5 lysis buffer were adjusted to the same protein concentration

before incubating them with 1/50 (v_{beads}/v_{lysate}) equilibrated NeutrAvidin-agarose beads

(ThermoFisher) over-night at RT. Due to the high-affinity interaction between biotin and

streptavidin, beads were subjected to stringent series of washes, using the following WBs

($v_{WB}/2v_{lysate}$), all made in 1x PBS: 2x WB1 (8 M urea, 0.25% SDS); 3x WB2 (6 M

Guanidine-HCl); 1x WB3 (6.4 M urea, 1 M NaCl, 0.2% SDS); 3x WB4 (4 M urea, 1 M

NaCl, 10% isopropanol, 10% ethanol and 0.2% SDS); 1x WB1; 1x WB5; and 3x WB6

(2% SDS). Biotinylated proteins were eluted in 80 µl of Elution Buffer (4x Laemmli

buffer, 100 mM DTT) through heating at 99°C for 5 minutes and subsequent vortexing.

Beads were separated by centrifugation (25000 x g, 2 minutes).

Liquid Chromatography Mass Spectrometry (LC-MS/MS)

A stable TRIPZ–MYC-CTurboID[195]–SUMO2nc U2OS cell line was transduced with

either EFS–FLAG-NTurboID[194]–PML[WT] or EFS–FLAG-NTurboID[194]–PML[3MAS] for

PML SUMO-ID experiments. Selection in blasticidin (5 µg/ml) and puromycin (1 µg/ml)

was performed. Expression was validated by Western blot and immunostaining prior to

scale-up for mass spectrometry. The TurboID-PML experiments used U2OS stable cell

lines expressing low and equivalent levels of PML[WT]–TurboID, PML[3MAS]–TurboID and

TurboID alone, selected by blasticidin (5 µg/ml), and treated or not with ATO for 2 hours.

For SALL1 SUMO-ID, a HEK293FT stable cell line expressing low levels of TRIPZ–

MYC-CTurboID[195]–SUMO2nc (selected with puromycin, 1 µg/ml) was transiently

transfected with EFS-FLAG-NTurboID^{194}–SALL1^{WT} or EFS-FLAG-NTurboID^{194}–

SALL1$^{\Delta SUMO}$ (a mutant carrying Lys>Arg mutations at K571, K592, K982, K1086).

Three independent pulldown experiments (8 x 10^7 cells per replicate, 8 ml of lysis) were

analyzed by LC-MS/MS.

Samples eluted from the NeutrAvidin beads were separated in SDS-PAGE (50% loaded)

and stained with Sypro Ruby (Invitrogen) according to manufacturer's instructions. Gel

lanes were sliced into 3 pieces as accurately as possible to guarantee reproducibility. The

slices were subsequently washed in milli-Q water. Reduction and alkylation were

performed using dithiothreitol (10 mM DTT in 50 mM ammonium bicarbonate) at 56°C

for 20 min, followed by iodoacetamide (50 mM chloroacetamide in 50 mM ammonium

bicarbonate) for another 20 minutes in the dark. Gel pieces were dried and incubated with

trypsin (12.5 μg/ml in 50 mM ammonium bicarbonate) for 20 minutes on ice. After

rehydration, the trypsin supernatant was discarded. Gel pieces were hydrated with 50 mM

ammonium bicarbonate, and incubated overnight at 37ºC. After digestion, acidic peptides

were cleaned with TFA 0.1% and dried out in a RVC2 25 speedvac concentrator (Christ).

Peptides were resuspended in 10 μl 0.1% formic acid (FA) and sonicated for 5 min prior

to analysis.

PML samples were analyzed in a novel hybrid trapped ion mobility spectrometry –

quadrupole time of flight mass spectrometer (timsTOF Pro with PASEF, Bruker

Daltonics) coupled online to a nanoElute liquid chromatograph (Bruker). This mass

spectrometer takes advantage of a novel scan mode termed parallel accumulation – serial

fragmentation (PASEF). Sample (200 ng) was directly loaded in a 15 cm Bruker

nanoelute FIFTEEN C18 analytical column (Bruker) and resolved at 400 nl/min with a

100 minutes gradient. Column was heated to 50ºC using an oven.

For the analysis of SALL1 samples peptides were eluted from stage-tips in a solvent

composed of deionized water/acetonitrile/FA at a ratio of 50/50/0.1 v/v/v. Peptides were

lyophilized and dissolved in solvent A composed of deionized water/FA at a ratio of

100/0.1 v/v and subsequently analyzed by on-line C18 nano-HPLC MS/MS with a system

consisting of an Ultimate 3000 nano gradient HPLC system (ThermoFisher), and an

Exploris 480 mass spectrometer (ThermoFisher). Fractions were loaded onto a cartridge

precolumn (5 mm x ID 300 μm, C18 PepMap 100 A, 5 μm particles (ThermoFisher)),

using solvent A at a flow of 10 μl/minute for 3 minutes and eluted via a homemade

analytical nano-HPLC column (50 cm × ID 75 μm; Reprosil-Pur C18-AQ 1.9 μm, 120 A;

Dr. Maisch GmbH). The gradient was run from 2% to 40% solvent B

(water/acetonitrile/FA at a ratio of 20/80/0.1 v/v/v) in 40 minutes. The nano-HPLC

column was drawn to a tip of ~10 μm and acted as the electrospray needle of the MS

source. The temperature of the nano-HPLC column was set to 50°C (Sonation GmbH).

The mass spectrometer was operated in data-dependent MS/MS mode for a cycle time of

3 seconds, with a HCD collision energy at 28 V and recording of the MS2 spectrum in

the orbitrap, with a quadrupole isolation width of 1.6 Da. In the master scan (MS1) the

resolution was set to 60,000, and the scan range was set to 300-1600, at an Automatic

Gain Control (AGC) target of 3×10^6 with automated fill time. A lock mass correction on

the background ion m/z=445.12 was used. Precursors were dynamically excluded after

n=1 with an exclusion duration of 30 seconds, and with a precursor range of 10 ppm.

Charge states 2-6 were included. For MS2 the first mass was set to 120 Da, and the MS2

scan resolution was 30,000 at an AGC target of 75,000 with automated fill time.

Mass Spectrometry data analysis

Raw MS files were analyzed using MaxQuant (version 1.6.17)[66] matching to a human

proteome (Uniprot filtered reviewed *H. sapiens* proteome, UP000005640) with a

maximum of 4 missed cleavages and with precursor and fragment tolerances of 20 ppm

and 0.05 Da. Label-Free Quantification was enabled with default values except for a ratio

count set to 1. Slices corresponding to same lanes were considered as fractions.

Biotinylation on lysine and on protein N-term was included as a variable modification for

SALL1 SUMO-ID data, and biotinylated peptides were set to be included for

quantification. Matching between runs and matching unidentified features were enabled.

Only proteins identified with at least one peptide at FDR<1% were considered for further

analysis. Data were loaded onto the Perseus platform (version 1.6.14)[67] and further

processed (Log2 transformation, imputation, median normalization when needed). A t-

test was applied in order to determine the statistical significance of the differences

detected. Proteins detected with at least 2 peptides (except when otherwise specified) and

in at least 2 of the 3 replicates were included.

Network analysis was performed using the STRING app version 1.4.2[68] in Cytoscape

version 3.7.2[69], with a high confidence interaction score (0.7). Transparency and width of

the edges were continuously mapped to the String score (text mining, databases,

coexpression, experiments, fusion, neighborhood and cooccurrence). The Molecular

COmplex DEtection (MCODE) plug-in version 1.5.1[70] was used to identify highly

connected subclusters of proteins (degree cutoff of 2; Cluster finding: Haircut; Node score

cutoff of 0.2; K-Core of 2; Max. Depth of 100). Gene ontology analysis was performed

using g:Profiler web server[71].

The mass spectrometry proteomics data corresponding to PML SUMO-ID, PML-

TurboID and SALL1 SUMO-ID experiments have been deposited to the

ProteomeXchange Consortium via the PRIDE partner repository[72] with the dataset

identifiers PXD021770, PXD021809 and PXD021923, respectively.

SIM enrichment analysis

A thousand lists with the same number of proteins as PML SUMO-ID list (59) have been randomly generated from the human proteome (Uniprot filtered reviewed *H. sapiens* proteome, UP000005640). All those lists have been analyzed by adapting the script from[73] and running it on Python version 2.7.5, to obtain the content and number of SIM motifs per protein (ψ-ψ-X-ψ; ψ-X-ψ-ψ; ψ-ψ-ψ; where ψ is either a L, I or V and X is any amino acid) and the number of SIMs per thousand of amino acids (STAA). After removing three outliers (lists 46, 782, 794; ROUT method, Q=1%), STAA values from the random lists were normalized to Log2 and validated for Gaussian distribution (d'Agostino and Pearson normality test). Enrichments were computed using R software v3.6.0 and calculated as the ratio between PML SUMO-ID STAA value and the median of STAA values from the random lists. Empirical p-values have been calculated by counting the number of random lists whose STAA values were as extreme as the PML SUMO-ID STAA value. The raw data from the SIM enrichment analysis and the script can be found in Table S5 and Supplementary Source 1, respectively.

ACKNOWLEDGEMENTS

We are thankful to Iraide Escobes for her work in the Proteomics Platform at CIC bioGUNE and Arnoud de Ru in the Center for Proteomics and Metabolomics at the LUMC. RB and ACOV acknowledge funding by the grant 765445-EU (UbiCODE Program). RB acknowledges funding by grants BFU2017-84653-P (MINECO/FEDER, EU), SEV-2016-0644 (Severo Ochoa Excellence Program), SAF2017-90900-REDT (UBIRed Program) and IT1165-19 (Basque Country Government). Additional support was provided by the Department of Industry, Tourism, and Trade of the Basque Country Government (Elkartek Research Programs) and by the Innovation Technology Department of the Bizkaia County. FE is at Proteomics Platform, member of ProteoRed-

ISCIII (PT13/0001/0027). FE and AMA acknowledge CIBERehd. AC acknowledges the

Basque Department of education (IKERTALDE IT1106-16), the MCIU (PID2019-

108787RB-I00 (FEDER/EU), Severo Ochoa Excellence Accreditation SEV-2016-0644,

Excellence Networks RED2018-102769-T), the AECC (GCTRA18006CARR), La Caixa

Foundation (ID 100010434), under the agreement LCF/PR/HR17, and the European

Research Council (Starting Grant 336343, PoC 754627, Consolidator grant 819242).

CIBERONC was co-funded with FEDER funds. UM acknowledges the Basque

Government Department of Education (IT1165-19) and the Spanish MCIU (SAF2016-

76898-P (FEDER/EU)).

AUTHOR CONTRIBUTIONS

O.B.-G., J.D.S. and R.B designed experiments, analyzed data and wrote the manuscript.

O.B.-G., F.T., V.M., I.C., A.R.C., C.P., M.A., I.I. and J.D.S. developed experimental

protocols and performed experiments. A.C., A.M.A., F.E., U.M. and A.C.O.V. provided

scientific resources.

COMPETING INTERESTS

The authors declare no competing interests.

REFERENCES

1. Flotho, A. & Melchior, F. Sumoylation: a regulatory protein modification in

health and disease. *Annu Rev Biochem* **82**, 357-385 (2013).

2. Gareau, J.R. & Lima, C.D. The SUMO pathway: emerging mechanisms that

shape specificity, conjugation and recognition. *Nat Rev Mol Cell Biol* **11**, 861-

871 (2010).

3.   Ullmann, R., Chien, C.D., Avantaggiati, M.L. & Muller, S. An acetylation

switch regulates SUMO-dependent protein interaction networks. *Mol Cell* **46**,

759-770 (2012).

4.   Vertegaal, A.C. SUMO chains: polymeric signals. *Biochem Soc Trans* **38**, 46-49

(2010).

5.   Aillet, F. *et al.* Heterologous SUMO-2/3-ubiquitin chains optimize

IkappaBalpha degradation and NF-kappaB activity. *PLoS One* **7**, e51672 (2012).

6.   Pichler, A., Fatouros, C., Lee, H. & Eisenhardt, N. SUMO conjugation - a

mechanistic view. *Biomol Concepts* **8**, 13-36 (2017).

7.   Talamillo, A. *et al.* The role of SUMOylation during development. *Biochem Soc*

Trans **48**, 463-478 (2020).

8.   Talamillo, A. *et al.* SUMOylation in the control of cholesterol homeostasis.

Open Biol **10**, 200054 (2020).

9.   Hendriks, I.A. *et al.* Site-specific mapping of the human SUMO proteome

reveals co-modification with phosphorylation. *Nat Struct Mol Biol* **24**, 325-336

(2017).

10.   Hendriks, I.A. & Vertegaal, A.C. A comprehensive compilation of SUMO

proteomics. *Nat Rev Mol Cell Biol* **17**, 581-595 (2016).

11.   Kerscher, O. SUMO junction-what's your function? New insights through

SUMO-interacting motifs. *EMBO Rep* **8**, 550-555 (2007).

12.   Kumar, R., Gonzalez-Prieto, R., Xiao, Z., Verlaan-de Vries, M. & Vertegaal,

A.C.O. The STUbL RNF4 regulates protein group SUMOylation by targeting

the SUMO conjugation machinery. *Nat Commun* **8**, 1809 (2017).

13.   Sriramachandran, A.M. *et al.* Arkadia/RNF111 is a SUMO-targeted ubiquitin

ligase with preference for substrates marked with SUMO1-capped SUMO2/3

chain. *Nat Commun* **10**, 3678 (2019).

14.   Werner, A., Flotho, A. & Melchior, F. The RanBP2/RanGAP1*SUMO1/Ubc9

complex is a multisubunit SUMO E3 ligase. *Mol Cell* **46**, 287-298 (2012).

15.   Eskiw, C.H., Dellaire, G. & Bazett-Jones, D.P. Chromatin contributes to

structural integrity of promyelocytic leukemia bodies through a SUMO-1-

independent mechanism. *J Biol Chem* **279**, 9577-9585 (2004).

16.   Wang, J. *et al.* Promyelocytic leukemia nuclear bodies associate with

transcriptionally active genomic regions. *J Cell Biol* **164**, 515-526 (2004).

17.   Bernardi, R. & Pandolfi, P.P. Structure, dynamics and functions of

promyelocytic leukaemia nuclear bodies. *Nat Rev Mol Cell Biol* **8**, 1006-1016

(2007).

18.   Lallemand-Breitenbach, V. & de The, H. PML nuclear bodies: from architecture

to function. *Curr Opin Cell Biol* **52**, 154-161 (2018).

19.   Jensen, K., Shiels, C. & Freemont, P.S. PML protein isoforms and the

RBCC/TRIM motif. *Oncogene* **20**, 7223-7233 (2001).

20.   Zhang, X.W. *et al.* Arsenic trioxide controls the fate of the PML-RARalpha

oncoprotein by directly binding PML. *Science* **328**, 240-243 (2010).

21.   Shen, T.H., Lin, H.K., Scaglioni, P.P., Yung, T.M. & Pandolfi, P.P. The

Mechanisms of PML-Nuclear Body Formation. *Mol Cell* **24**, 805 (2006).

22.   Tatham, M.H. *et al.* RNF4 is a poly-SUMO-specific E3 ubiquitin ligase required

for arsenic-induced PML degradation. *Nat Cell Biol* **10**, 538-546 (2008).

23.   Kim, D.I. & Roux, K.J. Filling the Void: Proximity-Based Labeling of Proteins

in Living Cells. *Trends Cell Biol* **26**, 804-817 (2016).

24.   Choi-Rhee, E., Schulman, H. & Cronan, J.E. Promiscuous protein biotinylation

738   by Escherichia coli biotin protein ligase. *Protein Sci* **13**, 3043-3050 (2004).

739   25.   Kim, D.I. *et al.* Probing nuclear pore complex architecture with proximity-

740   dependent biotinylation. *Proc Natl Acad Sci U S A* **111**, E2453-2461 (2014).

741   26.   Roux, K.J., Kim, D.I., Raida, M. & Burke, B. A promiscuous biotin ligase

742   fusion protein identifies proximal and interacting proteins in mammalian cells. *J

743   Cell Biol* **196**, 801-810 (2012).

744   27.   Coyaud, E. *et al.* BioID-based Identification of Skp Cullin F-box (SCF)beta-

745   TrCP1/2 E3 Ligase Substrates. *Mol Cell Proteomics* **14**, 1781-1795 (2015).

746   28.   Dho, S.E. *et al.* Proximity interactions of the ubiquitin ligase Mind bomb 1

747   reveal a role in regulation of epithelial polarity complex proteins. *Sci Rep* **9**,

748   12471 (2019).

749   29.   Branon, T.C. *et al.* Efficient proximity labeling in living cells and organisms

750   with TurboID. *Nat Biotechnol* **36**, 880-887 (2018).

751   30.   Schopp, I.M. *et al.* Split-BioID a conditional proteomics approach to monitor the

752   composition of spatiotemporally defined protein complexes. *Nat Commun* **8**,

753   15690 (2017).

754   31.   Cho, K.F. *et al.* Split-TurboID enables contact-dependent proximity labeling in

755   cells. *Proc Natl Acad Sci U S A* **117**, 12143-12154 (2020).

756   32.   Banaszynski, L.A., Liu, C.W. & Wandless, T.J. Characterization of the

757   FKBP.rapamycin.FRB ternary complex. *J Am Chem Soc* **127**, 4715-4721

758   (2005).

759   33.   Bekes, M. *et al.* The dynamics and mechanism of SUMO chain deconjugation

760   by SUMO-specific proteases. *J Biol Chem* **286**, 10238-10247 (2011).

761   34.   Shen, T.H., Lin, H.K., Scaglioni, P.P., Yung, T.M. & Pandolfi, P.P. The

762   mechanisms of PML-nuclear body formation. *Mol Cell* **24**, 331-339 (2006).

763   35.   Rabellino, A. *et al.* The SUMO E3-ligase PIAS1 regulates the tumor suppressor

764   PML and its oncogenic counterpart PML-RARA. *Cancer Res* **72**, 2275-2284

765   (2012).

766   36.   Zhu, J. *et al.* Small ubiquitin-related modifier (SUMO) binding determines

767   substrate recognition and paralog-selective SUMO modification. *J Biol Chem*

768   **283**, 29405-29415 (2008).

769   37.   Li, J. *et al.* GTF2I-RARA is a novel fusion transcript in a t(7;17) variant of acute

770   promyelocytic leukaemia with clinical resistance to retinoic acid. *Br J Haematol*

771   **168**, 904-908 (2015).

772   38.   Yin, C.C. *et al.* Identification of a novel fusion gene, IRF2BP2-RARA, in acute

773   promyelocytic leukemia. *J Natl Compr Canc Netw* **13**, 19-22 (2015).

774   39.   Banani, S.F. *et al.* Compositional Control of Phase-Separated Cellular Bodies.

775   *Cell* **166**, 651-663 (2016).

776   40.   Chung, I., Osterwald, S., Deeg, K.I. & Rippe, K. PML body meets telomere: the

777   beginning of an ALTernate ending? *Nucleus* **3**, 263-275 (2012).

778   41.   Van Damme, E., Laukens, K., Dang, T.H. & Van Ostade, X. A manually curated

779   network of the PML nuclear body interactome reveals an important role for

780   PML-NBs in SUMOylation dynamics. *Int J Biol Sci* **6**, 51-67 (2010).

781   42.   Netzer, C., Bohlander, S.K., Rieger, L., Muller, S. & Kohlhase, J. Interaction of

782   the developmental regulator SALL1 with UBE2I and SUMO-1. *Biochem*

783   *Biophys Res Commun* **296**, 870-876 (2002).

784   43.   Pirone, L. *et al.* A comprehensive platform for the analysis of ubiquitin-like

785   protein modifications using in vivo biotinylation. *Sci Rep* **7**, 40756 (2017).

44.   Lauberth, S.M. & Rauchman, M. A conserved 12-amino Acid motif in Sall1

recruits the nucleosome remodeling and deacetylase corepressor complex. *The*

Journal of biological chemistry **281**, 23922-23931 (2006).

45.   Hendriks, I.A. & Vertegaal, A.C. A high-yield double-purification proteomics

strategy for the identification of SUMO sites. *Nat Protoc* **11**, 1630-1649 (2016).

46.   Becker, J. *et al.* Detecting endogenous SUMO targets in mammalian cells and

tissues. *Nat Struct Mol Biol* **20**, 525-531 (2013).

47.   Le Boulch, M., Brossard, A., Le Dez, G., Leon, S. & Rabut, G. Sensitive

detection of protein ubiquitylation using a protein fragment complementation

assay. *J Cell Sci* **133** (2020).

48.   Sung, M.K. *et al.* Genome-wide bimolecular fluorescence complementation

analysis of SUMO interactome in yeast. *Genome Res* **23**, 736-746 (2013).

49.   Fang, D. & Kerppola, T.K. Ubiquitin-mediated fluorescence complementation

reveals that Jun ubiquitinated by Itch/AIP4 is localized to lysosomes. *Proc Natl*

Acad Sci U S A **101**, 14782-14787 (2004).

50.   Komiya, M. *et al.* A genetic screen to discover SUMOylated proteins in living

mammalian cells. *Sci Rep* **7**, 17443 (2017).

51.   Croucher, D.R. *et al.* Bimolecular complementation affinity purification

(BiCAP) reveals dimer-specific protein interactions for ERBB2 dimers. *Sci*

Signal **9**, ra69 (2016).

52.   Lin, D.Y. *et al.* Role of SUMO-interacting motif in Daxx SUMO modification,

subnuclear localization, and repression of sumoylated transcription factors. *Mol*

Cell **24**, 341-354 (2006).

53.   Sahin, U. *et al.* Oxidative stress-induced assembly of PML nuclear bodies

controls sumoylation of partner proteins. *J Cell Biol* **204**, 931-945 (2014).

54.   Saitoh, N. *et al.* In situ SUMOylation analysis reveals a modulatory role of

RanBP2 in the nuclear rim and PML bodies. *Exp Cell Res* **312**, 1418-1430

(2006).

55.   Lallemand-Breitenbach, V. *et al.* Arsenic degrades PML or PML-RARalpha

through a SUMO-triggered RNF4/ubiquitin-mediated pathway. *Nat Cell Biol* **10**,

547-555 (2008).

56.   Gechijian, L.N. *et al.* Functional TRIM24 degrader via conjugation of

ineffectual bromodomain and VHL ligands. *Nat Chem Biol* **14**, 405-412 (2018).

57.   Li, C., Xin, H., Shi, Y. & Mu, J. Knockdown of TRIM24 suppresses growth and

induces apoptosis in acute myeloid leukemia through downregulation of

Wnt/GSK-3beta/beta-catenin signaling. *Hum Exp Toxicol* **39**, 1725-1736 (2020).

58.   Lallemand-Breitenbach, V. *et al.* Role of promyelocytic leukemia (PML)

sumolation in nuclear body formation, 11S proteasome recruitment, and As2O3-

induced PML or PML/retinoic acid receptor alpha degradation. *J Exp Med* **193**,

1361-1371 (2001).

59.   Sha, Z., Blyszcz, T., Gonzalez-Prieto, R., Vertegaal, A.C.O. & Goldberg, A.L.

Inhibiting ubiquitination causes an accumulation of SUMOylated newly

synthesized nuclear proteins at PML bodies. *J Biol Chem* **294**, 15218-15234

(2019).

60.   Shiio, Y. & Eisenman, R.N. Histone sumoylation is associated with

transcriptional repression. *Proc Natl Acad Sci U S A* **100**, 13225-13230 (2003).

61.   Jangal, M. *et al.* The transcriptional co-repressor TLE3 suppresses basal

signaling on a subset of estrogen receptor alpha target genes. *Nucleic Acids Res*

42, 11339-11348 (2014).

62. Flack, J.E., Mieszczanek, J., Novcic, N. & Bienz, M. Wnt-Dependent

Inactivation of the Groucho/TLE Co-repressor by the HECT E3 Ubiquitin

Ligase Hyd/UBR5. *Mol Cell* **67**, 181-193 e185 (2017).

63. Yan, W. *et al.* Epigenetic regulation of DACH1, a novel Wnt signaling

component in colorectal cancer. *Epigenetics* **8**, 1373-1383 (2013).

64. Sato, A. *et al.* Sall1, a causative gene for Townes-Brocks syndrome, enhances

the canonical Wnt signaling by localizing to heterochromatin. *Biochem Biophys*

Res Commun **319**, 103-113 (2004).

65. Doench, J.G. *et al.* Optimized sgRNA design to maximize activity and minimize

off-target effects of CRISPR-Cas9. *Nat Biotechnol* **34**, 184-191 (2016).

66. Cox, J. & Mann, M. MaxQuant enables high peptide identification rates,

individualized p.p.b.-range mass accuracies and proteome-wide protein

quantification. *Nat Biotechnol* **26**, 1367-1372 (2008).

67. Tyanova, S. *et al.* The Perseus computational platform for comprehensive

analysis of (prote)omics data. *Nat Methods* **13**, 731-740 (2016).

68. Snel, B., Lehmann, G., Bork, P. & Huynen, M.A. STRING: a web-server to

retrieve and display the repeatedly occurring neighbourhood of a gene. *Nucleic*

Acids Res **28**, 3442-3444 (2000).

69. Shannon, P. *et al.* Cytoscape: a software environment for integrated models of

biomolecular interaction networks. *Genome Res* **13**, 2498-2504 (2003).

70. Bader, G.D. & Hogue, C.W. An automated method for finding molecular

complexes in large protein interaction networks. *BMC Bioinformatics* **4**, 2

(2003).

71. Reimand, J. *et al.* g:Profiler-a web server for functional interpretation of gene

lists (2016 update). *Nucleic Acids Res* **44**, W83-89 (2016).

72.   Vizcaino, J.A. *et al.* 2016 update of the PRIDE database and its related tools.

Nucleic Acids Res **44**, 11033 (2016).

73.   Sun, H. & Hunter, T. Poly-small ubiquitin-like modifier (PolySUMO)-binding

proteins identified through a string search. *J Biol Chem* **287**, 42071-42083

(2012).

74.   Weaver, L.H., Kwon, K., Beckett, D. & Matthews, B.W. Corepressor-induced

organization and assembly of the biotin repressor: a model for allosteric

activation of a transcriptional regulator. *Proc Natl Acad Sci U S A* **98**, 6045-

6050 (2001).

75.   Sehnal, D., Rose, A.S., Koča, J., Burley, S.K. & Velankar, S. in Proceedings of

the Workshop on Molecular Graphics and Visual Analysis of Molecular Data

29–33 (Eurographics Association, Brno, Czech Republic; 2018).

A

B

<u>**Figure 1:**</u> **Identification of SUMO-dependent interactions: the SUMO-ID strategy.**

Schematic representation of the SUMO-ID strategy **(A)** and the constructs used **(B)**.

BLAST^R, blasticidin resistant cassette; CTurboID, C-terminal TurboID; DOX,

doxycycline; NTurboID, N-terminal TurboID; PURO^R, puromycin resistant cassette; S,

SUMO; SIM: SUMO Interacting Motif; SDI: SUMO-dependent interactor; SUMOnc,

SUMO non-cleavable; Strep, streptavidin.

<u>Figure 2:</u> T194/G195 Split-TurboID is suitable for SUMO-ID studies. (A) Structure

of the *E. coli* BirA with bound biotin (PDB ID: 1HXD[74,75]), depicting the T194/G195

split point and the BirA-Biotin interaction. T194/G195 split point breaks the 193-199 loop

that connects the biotin-interacting β8 and β9 strands (see File S1). **(B)** Western blot of

HEK293FT cells that were transiently transfected with combinations of the FLAG-

NTurboID[194] (N) or MYC-CTurboID[195] (C) fused to PMLIVa, RANGAP1 or SUMO1/2

and treated with 50 μM of biotin for 16 hours. Black arrowheads indicate SUMO-ID

activity derived from MYC-CTurboID[195]-SUMOylated FLAG-NTurboID[194]-substrates.

White squares indicate biotinylated free MYC-CTurboID[195]-SUMOs. Neither FLAG-

NTurboID[194] nor MYC-CTurboID[195] showed any detectable background biotinylating

activity. **(C)** Immunostainings of transiently transfected U2OS cells treated with 50 μM

of biotin for 16 hours, showing the fragment-complementation dependency of SUMO-ID

and its correct localization within the cell, enriched at PML NBs as expected for

SUMOylated PML. Nuclei are stained with DAPI (blue) and biotinylated material with

fluorescent streptavidin (Strep, magenta). BirA antibody recognizes both NTurboID[194]

and CTurboID[195] (green). Black and white panels show the single green and magenta

channels. Scale bar: 5 μm. **(D)** Western blot of HEK293FT stable cells for NTurboID[194]-

FRB alone or in combination with CTurboID[195]-FKBP, treated or not with 1 μg/mL of

rapamycin and 50 μM of biotin at indicated time-points. BirA antibody recognizes both

NTurboID[194] and CTurboID[195]. White squares and circles indicate NTurboID[194]-FRB and

CTurboID[195]-FKBP, respectively. Self-biotinylating activity of the reconstituted TurboID

was measured and normalized to expression levels (BirA blot). Molecular weight markers

are shown to the left of the blots in kDa.

Figure 3: SUMO-ID is specific for SUMO-dependent interactions. (A) Western blot

of HEK293FT FLAG-NTurboID[194]-PMLIVa stable cell line transfected with different

combinations of MYC-CTurboID[195] (C) and SUMO[WT] or SUMO[ΔGG]. Cells were treated

or not with 1 μM of ATO for 2 hours and 50 μM of biotin at indicated time points. White

square indicates biotinylation of unconjugated MYC-CTurboID[195]- SUMO[ΔGG]. White

arrowhead points to SUMO-SIM interaction mediated SUMO-ID. Black arrowhead

shows PML-SUMOylation derived SUMO-ID. **(B)** Western blot of HEK293FT

transfected with constitutive MYC-CTurboID[195]-SUMO1/2 or doxycycline-inducible

and isopeptidase-cleavage resistant (nc) MYC-CTurboID[195]-SUMO1/2nc or MYC-

CTurboID[195]-Ubnc. Doxycycline was added or not at 1 μg/mL for 24 hours. White

squares point to free/unconjugated MYC-CTurboID[195]-SUMOs. **(C)** Western blot of

U2OS double stable cell lines for FLAG-NTurboID[194]-PMLIVa^{WT} or the SUMO/SIM

mutant FLAG-NTurboID[194]-PMLIVa^{3MAS} together with doxycycline-inducible TRIPZ-

MYC-CTurboID[195]-SUMO2nc. Doxycycline was added or not at 1 μg/mL for 24 hours.

50 μM of biotin was added at indicated time-points. PML SUMO-ID showed a high

PML/SUMO interaction dependency. **(D)** Confocal microscopy of the same cells as in

(C), treated or not with doxycycline (1 μg/mL, 24 hours), biotin (50 μM, 2 hours) and

ATO (1 μM, 2 hours). Nuclei are stained with DAPI (blue) and biotinylated material with

fluorescent streptavidin (Strep, magenta). BirA antibody shows NTurboID[194]-PML

staining (green). Black and white panels show the single green and magenta channels.

Colocalization of the streptavidin and NTurboID[194]-PMLIVa^{WT} signal is observed within

PML NBs, that depends on PML-SUMO interaction. Scale bar: 5 μm. Molecular weight

markers are shown to the left of the blots in kDa in (A-C).

9__

Figure 4: SUMO-ID identifies SUMO-dependent interactors of PML. (A) Volcano

plot of LC-MS analysis comparing streptavidin pull-downs of U2OS double stable cell

lines for TRIPZ-MYC-CTurboID[195]-SUMO2nc together with FLAG-NTurboID[194]-

PMLIVa[WT] or FLAG-NTurboID[194]-PMLIVa[3MAS]. Cells were treated with 1 μg/mL of

doxycycline for 24 hours and 50 μM of biotin for 2 hours. 59 high-confidence SUMO-

dependent PML interactors were defined. Asterisk (*) indicates that IRF2BP2 was

detected with one peptide but further validated by Western blot and immunofluorescence.

(B) Western blot validations of PML SUMO-dependent interactors identified by SUMO-

ID in (A). Blots 1/2 represent 2 independent experiments. UBC9 was added as an

expected positive control. Dots indicate endogenous carboxylases. Black arrowheads

point to specific PML SUMO-ID biotinylating activity. IN: input; St PD: streptavidin

pull-down. Molecular weight markers are shown to the left of the blots in kDa. **(C)**

STRING network analysis of the 59 SUMO-dependent interactors of PML identified in

(A). A highly interconnected cluster related to protein SUMOylation/ubiquitylation,

transcriptional regulation, DNA repair and RNA stability proteins is depicted. Color,

transparency and size of the nodes were discretely mapped to the Log2 enrichment value

as described. The border line type was discretely mapped to the p-value as described. **(D)**

Gene ontology analysis of the 59 SUMO-dependent interactors of PML identified in (A).

Biological processes and molecular functions related to SUMOylation/ubiquitylation,

stress response, DNA repair, transcription and RNA stability were significantly enriched.

Dotted line represents the threshold of the p-value (0.05).

Figure 5: **SUMO-dependent interactors of PML localize to PML NBs.** Confocal

microscopy analysis of PML SUMO-ID identified proteins in U2OS YFP-PML *knock-in*

958   cell line. UBC9 was added as an expected positive control. Yellow arrowheads indicate

959   colocalization events. Dotted line-squares show the selected colocalization events for

960   digital zooming and the signal profile plotting shown to the right. Nuclei are stained with

961   DAPI (blue), YFP-PML is shown in green and the indicated proteins in magenta. Black

962   and white panels show the single green and magenta channels. Scale-bar: 5 μm.

Figure 6: Characterization of the whole PML NBs proteome. (A) Volcano plot of LC-MS analysis comparing streptavidin pull-downs of U2OS stable cell lines for TurboID-PMLIVa[WT] or TurboID alone. Cells were treated with 50 μM of biotin for 2 hours. High-

confidence PML proteome composed of 267 proteins is shown as blue dots. **(B)** STRING

network analysis of the whole PML NBs proteome defined in (A) shows a high

interconnected network composed of the 73.6% of the proteins. Highly interconnected

sub-clusters were characterized using MCODE. Color, transparency and size of the nodes

were discretely mapped to the Log2 enrichment value as described. The border line type

was discretely mapped to the p-value as described. **(C)** Gene ontology analysis of the

whole PML NBs proteome defined in (A). Depicted biological processes, molecular

functions and cellular components were significantly enriched. Dotted line represents the

threshold of the p-value (0.05).

A

B

C

D

Figure 7: Proteins identified by PML SUMO-ID are a subset of the SUMO-dependent PML NBs proteome and are enriched in SIMs. (A) Volcano plot of LC-MS analysis comparing streptavidin pull-downs of U2OS stable cell lines for TurboID-PMLIVa^{WT} or TurboID-PMLIVa^{3MAS}. Cells were treated with 50 μM of biotin for 2 hours. Proteins enriched in TurboID alone compared to TurboID-PMLIVa^{WT} were previously removed for the comparison. PML SUMO-ID identified proteins (including 1 peptide identified proteins) are highlighted in orange. LC-MS data on the effect of the ATO treatment (1 μM; 2 hours) for TurboID-PMLIVa^{WT} enriched proteins is represented with symbols as described. **(B)** WB validation of the effect of ATO treatment (1 μM; 2 hours) on TRIM24 by PML SUMO-ID. Cells were treated with 1 μg/mL of doxycycline for 24 hours and 50 μM of biotin for 2 hours. After streptavidin pulldown, decreased levels of SUMO-PML interacting TRIM24 upon ATO treatment is observed. **(C)** Violin plots comparing the percentage of single SIM and multiple SIM presence in 1000 random lists and PML SUMO-ID list. The 1000 random lists contain the same number of proteins (59) as the SUMO-ID list. **(D)** SIM presence was normalized by the length of the proteins to obtain the value of SIMs per thousand of amino acids (STAA). Gaussian distribution of STAA median values of the random lists was validated (d'Agostino and Pearson normality test, p-value = 0.15), and PML SUMO-ID SIM enrichment factor was calculated. The dotted black line represents the median STAA value of random lists. The dotted blue line represents the STAA value of the PML SUMO-ID list.

Figure 8: SUMO-ID identifies interactors of SUMOylated SALL1. (A) WB of

HEK293FT stable cell lines for TRIPZ-MYC-CTurboID[195]-SUMO1nc/SUMO2nc

transfected with FLAG-NTurboID[194]-SALL1[WT] or the SUMO site mutant FLAG-

NTurboID^{194}-SALL1$^{\Delta SUMO}$. Cells were treated or not with 1 μg/mL of doxycycline for

24 hours and 50 μM of biotin at indicated time points. Efficient SALL1 SUMO-ID

biotinylating activity was detected for SUMO2nc (black arrowhead). Dot indicates

endogenous carboxylase. **(B)** Confocal microscopy of U2OS stable cell line for TRIPZ-

MYC-CTurboID^{195}-SUMO2nc transfected with FLAG-NTurboID^{194}-SALL1^{WT} or the

SUMO site mutant FLAG-NTurboID^{194}-SALL1$^{\Delta SUMO}$. Cells were treated or not with 1

μg/mL of doxycycline for 24 hours and 50 μM of biotin for 4 hours. Nuclei are stained

with DAPI (blue) and biotinylated material with fluorescent streptavidin (Strep,

magenta). BirA antibody shows NTurboID^{194}-SALL1 staining (green). Black and white

panels show the single green and magenta channels. Nuclear colocalization of FLAG-

NTurboID^{194}-SALL1^{WT} and streptavidin signal was observed. **(C)** Volcano plot of LC-

MS analysis comparing streptavidin pull-downs of HEK293FT TRIPZ-MYC-

CTurboID^{195}-SUMO2nc stable cell line transfected with FLAG-NTurboID^{194}-SALL1^{WT}

or the SUMO site mutant FLAG-NTurboID^{194}-SALL1$^{\Delta SUMO}$. Cells were treated with 1

μg/mL of doxycycline for 24 hours and 50 μM of biotin for 4 hours. Potential interactors

of SUMOylated SALL1 are depicted. **(D)** Western blot validations of SUMOylated

SALL1 interactors found in (C). NuRD complex proteins GATAD2B, MTA2 and RBBP4

were confirmed. Black arrowheads point to SUMOylated SALL1 signal. Dots indicate

endogenous carboxylases. **(E)** STRING network analysis of the SALL1 SUMO-ID list

and MCODE clustering identifies the NuRD complex as a highly interconnected

subcluster. Gene ontology analysis also identified the NuRD complex as an enriched

term. Color, transparency and size of the nodes were discretely mapped to the Log2

enrichment value as described. The border line type was discretely mapped to the p-value

as described. Molecular weight markers are shown to the left of the blots in kDa in (A,

D).