Neural dysfunction at the upper thermal limit in the zebrafish

Anna H. Andreassen[*] & Petter Hall[*], Pouya Khatibzadeh, Fredrik Jutfelt, Florence Kermen[#]

[*] These authors contributed equally.

[#] Corresponding author: florence.kermen@ntnu.no

Authors affiliation

Department of Biology, Faculty of Natural Sciences, Norwegian University of Science and Technology, 7491 Trondheim, Norway.

Keywords: heat stress, spreading depolarization, CT_{max}, thermal tolerance, central nervous system, calcium imaging, zebrafish

Highlights

- Larval zebrafish reach their critical thermal limit (CT_{max}) at similar temperature as adult zebrafish
- Acute heat stress causes a brain-wide spreading depolarization near the upper thermal limit
- CT_{max} precedes brain-wide depolarization
- Heart rate declines at high temperatures but is maintained during CT_{max} and brain depolarization
- Neural activity is impaired prior to CT_{max} and brain-wide depolarization
- Oxygen availability in the water affects both CT_{max} and neural activity

ABSTRACT

Understanding animal thermal tolerance is crucial to predict how animals will respond to increasingly warmer temperatures, and to mitigate the impact of the climate change on species survival. Yet, the physiological mechanisms underlying animal thermal tolerance are largely unknown. In this study, we developed a method for measuring upper thermal limit (CT_{max}) in larval zebrafish (*Danio rerio*) and found that it occurs at similar temperatures as in adult zebrafish. We discovered that CT_{max} precedes a transient, heat-induced brain-wide depolarization during heat ramping. By monitoring heart rate, we established that cardiac function is sub-optimal during the period where CT_{max} and brain depolarization occur. In

addition, we found that oxygen availability affects both locomotor neural activity and CT_{max}
during a heat stress. The findings of this study suggest that neural impairment due to limited
oxygen availability at high temperatures can cause CT_{max} in zebrafish.

INTRODUCTION

Periods of abnormally high temperatures are becoming more numerous and severe as climate
change progresses (Seneviratne *et al.*, 2014; Stillman, 2019). As biological rates are highly
affected by temperature, animals that hold similar body temperatures as the surrounding
environment (i.e. ectotherms) might be particularly vulnerable to extreme heat (Deutsch *et al.*,
2008; Schulte, 2015; Pinsky *et al.*, 2019). Under acute heating, ectotherms reach a critical
thermal limit (CT_{max}), a temperature where their equilibrium is lost and their movement
becomes disorganized (Friedlander *et al.*, 1976; Morgan *et al.*, 2018; Jørgensen *et al.*, 2020).
The thermal tolerance of ectotherms correlates with their geographical distributions (Perry *et*
al., 2005; Sunday *et al.*, 2012), suggesting that limits in thermal tolerance partly determines
distribution ranges. Moreover, rapid onset of heat waves can cause mass mortality in fishes,
which suggests that thermal tolerance can be an increasingly important trait in the future (Genin
et al., 2020).

Knowledge about the mechanisms controlling thermal tolerance is therefore important to
understand how ectotherms may respond to increasingly warmer temperatures and predict
ecosystem impacts. Although the physiological mechanisms of thermal tolerance are largely
unknown, two major bottlenecks have been proposed; oxygen limitation and neural
dysfunction (Friedlander *et al.*, 1976; Pörtner *et al.*, 2017; Jutfelt *et al.*, 2019).

The oxygen limitation view (i.e. the oxygen- and capacity-limited thermal tolerance (OCLTT)
hypothesis) suggests that tissue oxygen shortage limits thermal tolerance under acute heat
stress (Pörtner & Knust, 2007; Ern, 2019). According to this hypothesis, the cardiorespiratory
system in ectotherms is unable to match the increasing oxygen requirement with temperature,
leading to tissue hypoxia and thus setting the limits for thermal tolerance. Insufficient
cardiorespiratory function at high temperature, or near upper thermal limit, has been reported
in insects and fish (Pörtner, 2002; Farrell, 2007, 2009; Ern *et al.*, 2014), but it is maintained in
a number of other ectotherms (Ern *et al.*, 2015). In addition, experimental manipulation of
oxygen availability during heat stress does not always affect thermal tolerance (McArley *et al.*,
2020). These contrasting results suggest that oxygen limitation can only partly explain thermal

tolerance, which is likely limited by different mechanisms in various species and contexts (Ern

et al., 2016; Lefevre, 2016; Verberk *et al.*, 2016; Jutfelt *et al.*, 2018, 2019). Furthermore, the

OCLTT hypothesis does not specify which physiological mechanisms fail first due to tissue

hypoxia (Jutfelt *et al.*, 2018).

Efficient locomotion (which is impaired at CT_{max}) requires normal neural function that is

disrupted at high temperature. Therefore, neural control of locomotion is a heat-sensitive

physiological mechanism that could underlie upper thermal tolerance, either via direct thermal

effects on neurons (Friedlander *et al.*, 1976; Jutfelt *et al.*, 2019) or via indirect thermal effects

from oxygen limitations (Schulte, 2015). Examples of severe heat-induced neural dysfunctions

include spreading depolarizations in fruit fly (Jørgensen *et al.*, 2020), loss of rhythmic neural

activity in the digestive system of the crab (Marder *et al.*, 2015), and thermogenic seizures in

vertebrates (Dube *et al.*, 2000; Shinnar & Glauser, 2002; Hunt *et al.*, 2012). Yet, very few

studies have directly investigated how neural dysfunction relates to measures of thermal

tolerance. In the fruit fly, spreading depolarizations measured in head-restrained flies occur at

similar temperatures as heat-induced coma in freely moving conspecifics (Jørgensen *et al.*,

2020). Similarly, goldfish lose equilibrium at temperatures that induce hyperexcitability in

cerebellar neurons of anesthetized conspecifics (Friedlander *et al.*, 1976), and a study on

Atlantic cod (*Gadus morhua*) found that cooling the brain marginally increased CT_{max},

suggesting a causal link between brain function and thermal tolerance (Jutfelt *et al.*, 2019).

The relative importance of cardiorespiratory and oxygen limitations, versus direct thermal

impacts on neuronal function, as the mechanisms limiting thermal tolerance during warming

remains unclear (Clark *et al.*, 2013a, 2013b; Jutfelt *et al.*, 2014, 2018, 2019). This is partly due

to the challenge of recording brain activity and CT_{max} in the same freely moving animal. In this

study, we solved that challenge by simultaneously measuring brain activity and locomotion

during heating using non-invasive fluorescence imaging in transparent larval zebrafish (*Danio*

rerio) expressing a calcium indicator in the brain. We also recorded heart rate simultaneously

with neural activity during heating in a separate group of larval zebrafish. Finally, we measured

CT_{max} and neural activity in hypo- and hyperoxic conditions. This allowed us to test two

predictions. First, if dysfunction of the central nervous system limits thermal tolerance, neural

dysfunction should briefly precede or coincide with CT_{max}. Second, if neural dysfunction is

caused by a cardiorespiratory limitation, we predict that failure of cardiac function precedes

neural malfunction, and that water oxygen manipulation affects neural function and CT_{max}.

RESULTS

CT_{max} is similar in larval and adult zebrafish

The activity and upper thermal limit (CT_{max}) were recorded in freely swimming five-day-old larval zebrafish in a custom-designed glass chamber (**Figure 1A**, **B**), in which water temperature increased from 28°C at a rate of 0.3°C/min (heat ramp fish), or was maintained at 28°C (control fish). To measure the activity level of larval zebrafish in our set-up, we recorded swimming speed (**Figure S1**), episodes of disorganized swimming (spiral swimming, **Movie S1**) and loss of equilibrium (**Movie S2**) at selected temperature ranges during the heat ramp. The swimming speed of five-day-old zebrafish decreased on average from 2.9±0.6 mm/s to 1.7±0.3 mm/s during the assay, at a similar rate in the control and the heat ramp fish (linear mixed-effects model, time: β±S.E.=0.3±0.1, $t_{(103)}$=-2.5, p=0.013; treatment: β±S.E.=0.8±0.6, $t_{(24)}$=1.2, p=0.23, R^2=0.1, **Figure S1**). Ten of the 12 heat ramp fish, but none of the 14 control fish, displayed episodes of disorganized circular swimming prior to CT_{max} (Chi-square: $X^2_{(df=1,N=26)}$=15.6, p<0.001; **Figure 1C**). Animals close to their upper thermal limit often lose equilibrium, a behavioural event that is commonly used to determine CT_{max} across species (Moyano *et al.*, 2017; Morgan *et al.*, 2018). Here, we found that the heat ramp fish were unable to maintain equilibrium 15.3% of the time, more frequently than control fish, which lost equilibrium 0.8% of the time (Wilcoxon rank sum test, heat ramp: median±IQR=15.3±13.7%, control: 0.8±6.8%, r=0.58, p=0.004, **Figure 1D**). Yet, half of the control fish lost equilibrium during the assay (**Figure 1D**), likely because five-day-old zebrafish do not completely master postural stability.

We therefore measured CT_{max} as the temperature at which zebrafish larvae repeatedly failed to escape touches to the trunk (loss of response, see **Methods**). Using this loss of response criterion, five-day-old heat ramp fish reached CT_{max} at 41.4±0.1°C (**Figure 1E**). Nine-day-old fish tested in the same setup reached CT_{max} at similar temperatures of 41.3±0.2°C, indicating that the loss of response criterium is stable during larval development (linear regression model, nine-day-old: β±S.E.=-0.1±0.2°C, $t_{(36)}$=-0.5, p=0.7, **Figure 1E**). Furthermore, adults from the same line tested using the loss of equilibrium protocol (Morgan *et al.*, 2018), reached CT_{max} at similar temperatures (41.0±0.1°C) as larvae (adult: β±S.E.=-0.4±0.2, $t_{(36)}$=-1.9, p=0.07; $F_{(2,36)}$=1.9, R^2=0.1, **Figure 1E**).

Warming causes a transient brain-wide depolarization near the upper thermal limit

Neurons operate best within a certain thermal range (Robertson & Money, 2012; Tang *et al.*,

2012; Marder *et al.*, 2015). Therefore, we hypothesized that a neural malfunction would occur

as the fish's temperature approaches the upper thermal limit. To record neural activity in the

entire zebrafish brain during a heat ramp, zebrafish larvae, expressing the calcium indicator

GCaMP6s in all neurons, were mounted in agarose under an epifluorescence microscope

(**Figure 2A**, **B**). Calcium events were detected in the brainstem of heat ramp and control fish
(**Figure 2C**, **D**). The frequency of the brainstem calcium events remained stable over time in
the control fish (**Figure 2F**). In heat ramp fish, the frequency of brainstem calcium events
increased with temperature, before abruptly declining (**Figure 2F**) during the 15 minute-period
preceding a transient brain-wide depolarization (**Figure 2D**). The telencephalon and the
brainstem, which are normally weakly co-active, both displayed a sharp increase in activity
during the depolarization (**Figure 2D**). The brain-wide spread of the depolarization is further
illustrated in laterally mounted fish (**Figure 2E**, **Movie S3**): it spread relatively slowly, and
reached the dorsal brain regions 10-12 seconds after the onset in the ventral diencephalon
(**Figure 2E**). The heat-induced brain-wide depolarizations started on average at 40.5±0.4°C in
agar-embedded fish (**Figure 2G**), 0.8°C below the average CT_{max} temperature recorded in
freely swimming age-matched fish (41.4±0.1°C, **Figure 1E**). Our results show that a transient
brain-wide depolarization occurs during the heat ramp at temperatures close to CT_{max} in
zebrafish.

Figure 2. Embedded zebrafish larvae develop brain-wide depolarizations near the upper thermal limit.
Schematic overview (**A**) and image (**B**) from the experimental setup for whole brain neural activity measurement

in agar-embedded five-day-old *Tg(elavl3:GCaMP6s)* zebrafish larva. **B.** Raw fluorescence image of the larva brain highlighting the right telencephalon (dashed green line) and brainstem (dashed magenta line; A=anterior, L=lateral). **C**, **D** Change in fluorescence (% $\Delta F/F_0$, left y-axis) in the telencephalon and brainstem of a representative control larva (**C**) and a representative heat ramp larva (**D**). The water temperature (grey) was maintained at 28°C throughout the recording (right y-axis, **C**) for the control fish and was increased during a heat ramp treatment (0.3°C/min) until a brain-wide depolarization was detected, and then rapidly adjusted to 28°C until the end of the recording (right y-axis, **D**). **E.** Heatmap illustrating the temporal spread of the depolarization throughout the brain in a representative heat ramp fish mounted laterally (see **Methods**; A=anterior, D=dorsal). **F.** Frequency of brainstem calcium events in heat ramp fish (n=11; magenta line) during the 50 minutes preceding the brain-wide depolarization, and during the corresponding period in control fish (n=8, magenta dashed line). The depolarization onset in heatramp fish is indicated by a dashed vertical line. **G.** Temperature at depolarization onset in heat ramp fish (40.5°C ±0.4, n=11). Data are presented with mean (solid line) and S.E. (shaded area) in **F** and with a bar and error bars in **G**.

CT$_{max}$ precedes the brain-wide depolarization

Since agar-embedded larvae experienced a brain-wide depolarization at temperatures close to the upper thermal limit of zebrafish, we asked whether the depolarization preceded, or co-occurred with, the loss of response observed during heat ramp in freely swimming fish. To do so, we simultaneously measured CT$_{max}$ and neural activity in freely swimming zebrafish during a heat ramp, using epifluorescence imaging (**Figure 3A**, **B**, (Muto *et al.*, 2017)). CT$_{max}$ (40.9±0.2°C) preceded the brain-wide depolarization (41.4±0.2°C, **Figure 3C**) in all individuals by 0.5°C on average (linear mixed-effects model, β±S.E.=0.5±0.1, $t_{(6)}$=4.5, p=0.004, R^2=0.3, **Figure 3D**). These results from freely swimming fish show that brain-wide depolarizations were not an artefact due to agarose embedding in the previous experiment (**Figure 2**) and that the brain-wide depolarization does not precede CT$_{max}$.

Figure 3. CT$_{max}$ shortly precedes the brain-wide depolarization in freely swimming zebrafish larvae.
A. Schematic overview of the experimental setup for calcium imaging in freely swimming five-day-old
Tg(elavl3:GCaMP6s) zebrafish larva. **B.** Top view of setup with colour-coded time projection of a heat ramp
fish's movements in the set-up for three minutes (34-35°C). **C.** Representative images of brain fluorescence in the
same fish as in **B**, before (left) and during (right) the brain-wide depolarization. **D.** CT$_{max}$ and depolarization onset
temperatures measured in heat ramp fish (black circles, n=7; linear mixed effects model: β±S.E.=0.5±0.1, $t_{(6)}$=4.5,
p=0.004, R^2=0.3, **Table S3**). The bars and error bars indicate the mean and S.E. of the CT$_{max}$ and depolarization
measurements.

Cardiac function declines before CT$_{max}$ and the brain-wide depolarization

Hypoxic brain tissues due to a cardiorespiratory limitation, could result in neural dysfunction
during heat ramping. We therefore simultaneously measured neural activity and heart-beat
frequency in the same fish during a heat ramp (**Figure 4A**, **B**) to test if the heart rate was altered
prior to the depolarization onset. When the temperature was increased from 28°C to 34°C, the
heart rate increased with a mean Q$_{10}$ value of 1.9, from 4.4±0.1Hz to 5.7±0.1Hz (linear mixed-
effects model: β±S.E.=1.3±0.2Hz, $t_{(36)}$=6.9, p<0.001, R^2=0.8 **Figure 4C**). From 34 to 37°C,
the heart rate decreased by 16% to 4.7±0.2Hz (β±S.E.=0.9±0.2, $t_{(36)}$=-4.9, p<0.001) and then
remained stable during the period preceding the brain-wide depolarization (β±S.E.=-0.1±0.2,
$t_{(36)}$=0.6, p=0.6, **Figure 4C**). We also examined whether heart rate was affected during the

brain-wide depolarization but it was unchanged (**Figure 4D**, **E**). When the fish were returned
to normal holding temperature after the brain-wide depolarization the heart rate decreased by
34% from 4.7±0.2Hz to 3.1±0.3 Hz (β±S.E.=1.6±0.2 Hz, $t_{(36)}$=-8.5, $p<0.001$, **Figure 4C**).
Altogether, our data shows a decline in cardiac activity during the heat ramp, but no abrupt
change in cardiac function prior to the depolarization onset, during the period when CT_{max}
occurs.

Figure 4. Heart rate before and during brain-wide depolarization in embedded larval zebrafish.
A. Top panel: representative example of simultaneous neural activity and heartbeat imaging in a five-day-old
Tg(elavl3:GCaMP6s) zebrafish larva embedded laterally in agar, using epifluorescence microscopy. The brain is
outlined with black dashed lines and the heart with a dashed grey square (top panel). Bottom panel: The heart
region (dashed yellow line) is shown in inverted grayscale (D=dorsal, A=anterior) during a beating cycle. **B**. Top
panel: change in brain fluorescence (% $\Delta F/F_0$,) in the same fish as in **A**, at 31°C. Calcium events are indicated by
grey arrows. Bottom panel: change in luminosity within a region of interest in the heart during the same period,
which is used to calculate the heartbeat frequency. **C**. Average heartbeat frequency in embedded heat ramp fish
(n=10), recorded at different temperature intervals. The x-axis shows ranges of 1°C elevation during which the
heartbeat was measured. The fifth temperature category (brain depolarization) corresponds to heartbeat frequency
during the 60 seconds pre- and the 30 seconds post-peak brain depolarization. The sixth category corresponds to
heartbeat frequency after fish were rapidly returned to holding temperature. Horizontal bars and error bars indicate
the mean and S.E. of the heartbeat value for each temperature range. Individual fish are colour-coded (linear
mixed effects model: R^2= 0.8, **Table S4**). Change in neural activity (**D**) and heartbeat frequency (**E**) during the
brain-wide depolarization (same period as "Brain depolarization" in **C**). Thin grey lines represent individual fish
and the thick black line represent the mean. All traces are aligned with respect to the peak fluorescence (vertical
dashed line).

Oxygen availability modulates temperature for CT$_{max}$ and brain-wide depolarization

Since the heart rate is only 4-5 Hz at the time of the brain-wide depolarization (**Figure 4C** and **4E**), it is possible that the cardiorespiratory system cannot match the increasing oxygen demand of the neural tissue. If oxygen availability limits thermal tolerance, we expect that oxygen manipulations during heat ramping would alter CT$_{max}$ and neural function.

To test the effect of oxygen availability on CT$_{max}$, we first measured the temperature at loss of response in freely swimming larvae zebrafish exposed to hypoxic and hyperoxic conditions during heat ramping, at 60 and 150% oxygen of air-saturated water respectively. Both treatments had on average a lower CT$_{max}$ than in the previous experiment (**Figure 1E**) likely due to experimental differences. The average CT$_{max}$ occurred 0.9°C lower in hypoxia (39.2±0.1°C) than in hyperoxia (40.1±0.3°C; linear regression model, β±S.E.=0.9±0.3°C, $t_{(26)}$=3.2, p=0.004; $F_{(1,26)}$=10.2, R^2=0.3; **Figure 5A**), which indicates that oxygen availability affects behavioural thermal tolerance in larval zebrafish.

To test the effect of oxygen availability on neural activity, agar-embedded fish were placed under an epifluorescence microscope and subjected to three oxygen treatments during heat ramping: hypoxia, normoxia and hyperoxia at 60, 100 and 150% oxygen of air saturated water, respectively. The brain-wide depolarization occurred at 40.7±0.4°C in normoxia. It occurred 1.8°C lower in the hypoxia (38.8±0.2°C; linear regression model, β±S.E.=-1.8±0.4, $t_{(20)}$=-4.8, p<0.001; $F_{(2,20)}$=40.4, R^2=0.8; **Figure 5B**). On the contrary, the hyperoxia treatment during heat ramping increased the depolarization temperature by 1.3°C compared to normoxia (42.0±0.2°C; β±S.E.=1.3±0.4, $t_{(20)}$=3.3, p=0.003). These results indicate that oxygen availability strongly influences the onset of brain-wide depolarization in larval zebrafish.

To further determine the effect of oxygen availability on neural activity when zebrafish reach their upper thermal limits, we calculated the frequency of brainstem calcium events during the 15 minutes before the brain-wide depolarization. Neural activity in the locomotor brain center sharply decreased at high temperature in all treatments: almost no calcium events were detected during the minutes preceding the depolarization (**Figure 5C**). The hyperoxia group retained a higher event frequency than the hypoxia and control groups at the start of this period (linear mixed-effects model, time: β±S.E.=0.08±0.02, $t_{(216)}$=3.9, p<0.001; hyperoxia*time: β±S.E.=0.09±0.03, $t_{(216)}$=3.3, p=0.001; R^2=0.3, **Figure 5C**). This indicates that increased oxygen availability partially rescued neural function near zebrafish upper thermal limit.

We also observed an effect of oxygen availability on the temporal dynamic of the brain depolarization. Oxygen availability did not change the amplitude of the brain-wide depolarization (normoxia: $88.2\pm17.6\%\Delta F/F_0$; linear regression model, hypoxia: $\beta\pm$S.E.$=$-3.1 ± 17.1, $t_{(20)}=$-0.2, $p=0.86$; hyperoxia: $\beta\pm$S.E.$=$-15.4 ± 17.4, $t_{(20)}=$-0.9, $p=0.39$; $F_{(2,20)}=0.5$, $R^2=$-0.05, data not shown). However, oxygen availability significantly improved the depolarization recovery time. Compared to the normoxia group, the hyperoxia group returned to baseline activity after the depolarization 1.3 minutes faster, and the hypoxia group 2 minutes slower (recovery time in normoxia: 3.0 ± 0.4 min; linear regression model, hypoxia: $\beta\pm$S.E.$=2.0\pm0.5$, $t_{(19)}=4.1$, $p<0.001$; hyperoxia: $\beta\pm$S.E.$=$-1.3 ± 0.5, $t_{(19)}=$-2.5, $p=0.02$; $F_{(2,19)}=0.5$, $R^2=0.7$, **Figure 5D-E**).

Figure 5. Oxygen availability affects both CT_{max} and depolarization onset temperatures.
A. Effect of oxygen availability on CT_{max} measured in freely swimming fish (set-up illustrated in left panel) during heat ramping with water oxygen set at 60% (hypoxia, magenta circles, n=16) or 150% (hyperoxia, cyan circles, n=12) of air saturated water (right panel; linear regression model: $\beta\pm$S.E.$=0.9\pm0.3$, $F_{(1,26)}=10.2$, $p=0.004$, $R^2=0.3$, **Table S5**). **A-E.** All results for hypoxia (60%, magenta), normoxia (100%, black) and hyperoxia (150%, cyan) are presented with mean and S.E. (**A**, **B**, **E**: bars and error bars; **C**, **D**: solid lines and shaded area). **B.** Brain-wide depolarization onset temperatures measured in agar-embedded *Tg(elavl3:GCaMP6s)* five-day-old fish (set-up illustrated in left panel) during heat ramping in hypoxia (n=9), normoxia (n=6) and hyperoxia (n=8; linear regression model, hypoxia: $\beta\pm$S.E.$=$-$1.8\pm0.4°C$, $t_{(20)}=$-4.8, $p<0.001$; hyperoxia $\beta\pm$S.E.$=1.3\pm0.4°C$, $t_{(20)}=3.3$, $p=0.003$; $F_{(2,20)}=40.4$, $R^2=0.8$, **Table S6**). **C.** Frequency of brainstem calcium events recorded during the 30 minutes preceding the brain-wide depolarization in hypoxia (n=5), normoxia (n=5) and hyperoxia (n=6; linear mixed-effects model for the last 15 minutes, time: $\beta\pm$S.E.$=0.08\pm0.02$, $t_{(216)}=3.9$, $p<0.001$; hyperoxia*time: $\beta\pm$S.E.$=0.09\pm0.03$, $t_{(216)}=3.3$, $p=0.001$; $R^2=0.3$, **Table S7**). **D.** Change in brain fluorescence aligned to the peak

296   depolarization in hypoxia (n=9), normoxia (n=6) and hyperoxia (n=8). **E.** Recovery time between peak
297   depolarization and the return to baseline fluorescence in hypoxia (n=9), normoxia (n=6) and hyperoxia (n=7;
298   linear regression model, hypoxia: $\beta\pm$S.E.$=2.0\pm0.5$, $t_{(19)}=4.1$, $p<0.001$; hyperoxia: $\beta\pm$S.E.$=-1.3\pm0.5$, $t_{(19)}=-2.5$,
299   $p=0.02$; $F_{(2,19)}=0.5$, $R^2=0.7$, **Table S7**).
300

DISCUSSION

A massive heat-induced global depolarization arose in the larval brain, at similar temperatures as CT_{max} occurred in freely swimming fish. These brain depolarizations were 4-5 times the magnitude of brainstem locomotor calcium events, and lasted much longer (**Figure 2**). This abnormally high neural activity in response to heat stress is reminiscent of heat-induced seizures (Dube *et al.*, 2000; Shinnar & Glauser, 2002; Hunt *et al.*, 2012), and of heat-induced spreading depolarizations (Jørgensen *et al.*, 2020), which are slow propagating waves of brain depolarization (Spong *et al.*, 2016). The temporal dynamic of the events measured here is characteristic of the latter. First, the depolarizations spread slowly across the brain (**Figure 2E**), at a speed consistent with that reported for spreading depolarizations (2-9 mm/min, (Woitzik *et al.*, 2013; Spong *et al.*, 2016)), and slower than seizures (Wenzel *et al.*, 2017; Liu & Baraban, 2019). Second, the long post-depolarization recovery time (**Figure 5E**) is consistent with that seen after spreading depolarizations (Sawant-Pokam *et al.*, 2016; Spong *et al.*, 2016), and it is much longer than that measured after brain seizures (a few seconds, (Diaz Verdugo *et al.*, 2019; Liu & Baraban, 2019)). Overall, the temperatures at which these spreading depolarizations occurred, as well as their transient nature, made it a plausible candidate to explain the prolonged, yet reversible unresponsive state that we observed at CT_{max}.

If upper thermal tolerance in zebrafish larvae is caused by a spreading depolarization shutting down central nervous system function, the depolarization should precede or coincide with CT_{max} measured in the same animal. Surprisingly, larval zebrafish reached CT_{max} *before* they developed a brain-wide depolarization. This is in contrast with a series of studies in insects in which measures of upper thermal tolerance correlated with the onset of spreading depolarization in the central nervous system (Andersen *et al.*, 2018; Jørgensen *et al.*, 2020). In these studies, upper thermal limits and spreading depolarization onsets were measured in separate individuals, respectively freely moving flies and central nervous system preparations. Using a similar approach in the first part of this article, we also found overlapping temperature ranges for CT_{max} and spreading depolarization onset. Only when measuring both parameters within the same individual could we resolve the small but consistent difference between these consecutive events (**Figure 3D**), and determine that CT_{max} preceded spreading depolarization. We thus conclude that spreading depolarization is not the cause of CT_{max} in zebrafish.

The fact that CT_{max} precedes the spreading depolarization does not rule out neural impairment

as a possible mechanism determining the upper thermal limit. Previous studies suggested that

neural function is impaired during acute heat stress due to either a direct effect of temperature

(Friedlander et al. 1975; Jutfelt et al. 2019), or due to heat-induced tissue hypoxia. Extracellular

recordings of the pyloric neuron in the crab stomatogastric nervous system show that the

characteristic rhythmic firing crashes at high temperatures (Marder *et al.*, 2015). Similarly,

extracellular recordings of cerebellar neurons in anesthetized goldfish indicate a sharp decrease

in spontaneous and sensory-evoked firing rates at high temperatures (Friedlander *et al.*, 1976).

In addition, a strong reduction in spontaneous neural activity occurs in ischemic brain tissue in

rodents (Buzsaki *et al.*, 1989; Barth & Mody, 2011) and in oxygen-deprived cultures of human

cortical neurons (le Feber *et al.*, 2016). Such findings indicate that both direct effects of

temperature and tissue hypoxia are plausible candidate mechanisms. In line with these previous

studies, we found that neural activity was strongly reduced in locomotor brain regions during

the minutes preceding the depolarization (**Figure 2 and 5C**), which we interpret as an early

indication of neural malfunction that underlies the degradation of zebrafish locomotion

observed during the corresponding period (**Figure 1C-D**). As this period of neural inactivity

in the minutes leading up to CT_{max} could be due to both direct impacts of temperature and/or

oxygen deficit on neural function, we manipulated oxygen water concentrations to test if

oxygen availability alters CT_{max} and neural function.

We found that hyperoxia (150% air saturation) improved both behavioural and neural thermal

resilience compared to hypoxia. The fact that neural activity in the locomotor brain centre was

more resilient to high temperature in hyperoxia than in hypoxia and normoxia provides a likely

mechanism for the improved behavioural resilience of zebrafish in hyperoxic conditions.

Furthermore, similarly to earlier measurements in normoxia, neural activity was strongly

silenced in the brainstem of fish during the very last minutes preceding the brain depolarization,

when fish reach their upper thermal limit (**Figure 3D**). This shows that central nervous function

impairment coincided with the temperatures where CT_{max} occurs, further suggesting that neural

impairment due to a cumulative lack of oxygen and/or accumulation of anaerobic metabolites

contributes to the unresponsive state observed at the upper thermal limit.

A few studies have found a positive effect of increased oxygen availability on thermal

tolerance, while most fail to find an effect of hyperoxia (McArley *et al.*, 2020). Hyperoxia

improved the upper thermal limit by 1.1 °C (loss of equilibrium) in the European perch

(Ekström *et al.*, 2016) and in the Common triplefin (McArley *et al.*, 2018). Increased aquatic

oxygen availability also extended the survival of several ectotherm species (arthropods,
chordates, echinoderms, molluscs, and fish) at extreme high temperatures (Verberk *et al.*, 2018;
Giomi *et al.*, 2019). However, hyperoxia did not improve thermal tolerance for a large number
of other fish species (McArley *et al.*, 2020). Moreover, the upper thermal limit of the red drum
and lumpfish was only reduced under severe hypoxic conditions (Ern *et al.*, 2016).
Additionally, the aerobic scope, the aerobic capacity available for non-maintenance activities,
remains high in some ectotherms at thermal limits, indicating surplus oxygen transport capacity
(Overgaard *et al.*, 2012; Grans *et al.*, 2014; Wang *et al.*, 2014; Brijs *et al.*, 2015; Ekström *et
al.*, 2016). Taken together these studies suggest that oxygen limitation is not a general
mechanism limiting thermal tolerance across all species and contexts. Our results suggest that
inter-species differences in neural tissue resilience to heat and oxygen deprivation might be a
critical physiological factor explaining these different reports. It will also be interesting to
investigate whether the sensitivity of CT_{max} to oxygen availability in zebrafish is limited to
early life stages or persists throughout ontogeny.

To test whether heat ramping causes cardiac dysfunction we examined heart rate and brain
activity simultaneously. The heart rate initially (from 28-34°C) increased as expected with
temperature, similar to previous reports (Gollock *et al.*, 2006; Farrell, 2007). This initial
increase was followed by a decrease to to 270 beats per minute, a heart rate then sustained upon
further heating and throughout the brain depolarization. Similarly, a decrease or plateau in heart
rate with increasing temperatures has been reported for salmonids (Farrell, 2009), and
European perch (Ekström *et al.*, 2016). The lack of heart rate increase when approaching
thermal limits might contribute to mismatch between oxygen delivery and metabolic rate. Such
a mismatch may have caused developing tissue hypoxia and accumulation of metabolites,
which could have contributed to causing the loss of responses at CT_{max}.

Animals are adapted to widely different climates, and because of this, the temperatures at which
heat-induced behavioral alterations occur are strongly species-specific. Yet, the sequence and
nature of heat-induced behavioral alterations seem relatively well conserved across species.
For example, a similar sequence of events occur in fruit fly and goldfish subjected to rapidly
increasing temperatures: hyperactivity, then loss of coordinated movement, followed by loss
of equilibrium, before entering into a heat-coma indicated by an animal lying still and
unresponsive to mechanical stimulations (Friedlander *et al.*, 1976; Jørgensen *et al.*, 2020).

To enable comparison with previous studies we quantified the behavioral changes that occurred
during heat ramping. Although the swimming velocity of heat ramp fish tended to be higher

than that of controls between 31-38 degrees, there was no clear period of heat-induced
hyperactivity, possibly due to the high interindividual variability in locomotion in larval
zebrafish. Loss of motor coordination (spiral swimming) and loss of equilibrium occurred in
most heat ramp larval zebrafish, followed by a state of unresponsiveness that was similar to the
heat-coma documented in goldfish and fruit flies. The loss of response endpoint used here was
stable throughout larval development and yielded results within the range recorded using the
loss of equilibrium criterion in adults of the same *Tg(elavl3:GCaMP6s)* line (**Figure 1E**), or
in wild-type adult zebrafish (Morgan *et al.*, 2018), thus supporting its use a criterion for CT_{max}
in larvae. In addition, spiral swimming could be used as a novel, specific and early sign of heat-
induced locomotor dysfunction before the upper thermal limit is reached.

In conclusion, we show that the acute thermal limits of larval zebrafish are not caused by global
brain depolarization, but are instead related to a drop in neural acivity preceding CT_{max}.
Furthermore, in concordance with OCLTT predictions, tissue oxygen availability appears to
constrains both brain function as well as the whole animal thermal limits during thermal
ramping.

METHODS
Animals and housing

Experiments were conducted on five- to ten-day-old larvae and seven- to ten-month-old
zebrafish (*Danio rerio*). The transgenic line *Tg(elavl3:GCaMP6s)* (Vladimirov *et al.*, 2014) in
nacre/mitfa background (Lister *et al.*, 1999) was used. Eggs were collected in the morning and
kept at a density of one per mL in fish water (0.2g of marine salt and 0.04 L AquaSafe per litre
of carbon-filtrated water, used for all experiments). After hatching at three days post-
fertilization, the larvae were kept in small nursery tanks and fed twice a day with larval food
(Tetramin, Tetra) after 5 days post-fertilization. Fish were maintained under standard
laboratory conditions (26.8 ± 0.1°C; 12/12-hour light/dark cycle). On the day of the
experiment, the fish were fed once after the experiment was finished. All experimental
procedures performed on zebrafish were in accordance with the 2010/63/EU directive and
approved by the Norwegian Animal Research Authority (Food and Safety Authority; Permit
number: 8578).

The experiments consisted of five parts performed on different fish: 1) measuring CT_{max} in
freely swimming zebrafish larvae, 2) recording brain activity during warming in agar-

embedded zebrafish larvae, 3) simultaneous recording of CT_{max} and depolarization in freely

swimming zebrafish larvae, 4) simultaneous recording of heart rate and brain activity during

heat ramping and 5) recording of CT_{max} and neural function under oxygen manipulation during

warming.

CT_{max} in freely swimming larvae

CT_{max} setup

Larval zebrafish swam in the behavioural arena (central compartment) of a double-walled glass

heating-mantle made in the NTNU glass workshop. The dimensions were as follows: outer

diameter= 42 mm, inner diameter= 29 mm, outer height= 32 mm, inner height= 19 mm. The

fish movement was recorded using a webcam (Logitech C270, 720p) positioned above the

arena. The arena was placed above a background illumination light source to enhance contrast.

Two outlets connected the glass heating mantle to a water bath. Adjustment of water

temperature of the arena was achieved by pumping water from an external heating bath through

the heating-mantle surrounding the arena. The heating rate in the arena was 0.3°C per minute

as described in Morgan et al., (2018). The water temperature inside the central chamber of the

arena was recorded at 1 Hz using two thermocouples (type K, Pico Technology) connected to

a data logger (TC-08, Pico Technology).

CT_{max} assay

The arena was filled with three mL of 28°C water supplied with air bubbling through a

modified hypodermic needle fixed to the side wall of the compartment. At the beginning of a

CT_{max} recording, a single larva was carefully transferred to the arena. All individuals were

given 15 minutes to habituate to the arena before the recording started. The CT_{max} assay lasted

up to 50 minutes, during which the temperature within the arena increased from 28°C by

0.3°C/min until the larvae reached CT_{max}. The temperature was kept constant at 28°C for

recordings of control fish. The fish behaviour was recorded at 4-7 Hz during four intervals: 28-

29°C, 31-32°C, 34-35°C and 37°C-CT_{max}. For control larvae, recordings of matching durations

were taken at corresponding time points. Methods commonly used to determine CT_{max} in fish

were unsuitable for 5 dpf larvae zebrafish: loss of equilibrium (Morgan *et al.*, 2018) also

occurred in control fish without heat ramp; and muscular spasms (Lutterschmidt & Hutchison,

2011) could not be reliably quantified due to the larvae's small size. Thus, CT_{max} was

determined as the temperature at which the animal became unresponsive (Friedlander *et al.*,

1976; Sherman & Levitis, 2003; Jørgensen *et al.*, 2020), which was defined as the first of three

consecutive tactile stimulations that did not elicit an escape response (loss of response). The
stimulations were applied to the larva's trunk using the tip of a capillary micro-ladder (VWR),
with a minimum of three seconds between consecutive stimulations. Upon CT_{max}, the fish was
transferred to 28°C water and visually monitored. Fish that did not recover normal locomotor
activity were rapidly euthanized and were not included in the analysis (this happened in only
four out of 44 fish, similar to previous reports (Morgan *et al.*, 2018; Åsheim *et al.*, 2020)).

CT_{max} assay in adult zebrafish

To compare CT_{max} throughout development, 16 adult Tg(elavl3:GCaMP6s) zebrafish, aged
seven- to ten-month-old, were tested in the CT_{max} setup described by Morgan et al., (2018).
The fish were tested in groups of eight fish in a rectangular acrylic tank (25 cm long, 20 cm
wide, 18 cm deep) filled with nine litres of water. Individual CT_{max} was measured at the
temperature of loss of equilibrium determined as two seconds of inability to maintain postural
stability (Morgan *et al.*, 2018). The fish were removed immediately after the criterion was
reached. All animals survived the test.

CT_{max} assay data analysis

The fish position was manually detected in MATLAB (MathWorks 2018) at 28-29°C, 31-32°C,
34-35°C, 37-38°C and during the 1°C elevation preceding CT_{max}. The average speed was
calculated during these intervals. Loss of equilibrium and spiral swimming events were
manually labelled by an experimenter during the 12 min preceding CT_{max} in heat ramp fish,
and during the corresponding period in control fish. Loss of equilibrium was recorded when
the fish tilted to the side for at least one second (**Movie S2**). Spiral swimming event was
recorded when the fish rapidly performed a minimum of two consecutive revolutions with
diameter below 5 mm (**Movie S1**).

Brain activity during warming in agar-embedded larvae

486 The calcium imaging to record brain activity during warming was executed using the same
487 double-walled glass heating mantle as described for the CT_{max} assay above.

Epifluorescence calcium imaging in agar-embedded zebrafish larvae

For the imaging of neural activity in embedded fish (**Figure 2A**, **B**), a five-day-old larva was
embedded at the bottom of the central glass compartment in 2% low-gelling temperature
agarose (Merck). The agarose was carefully removed around the eyes and mouth upon
hardening and the arena was filled with 3 mL water. Calcium fluorescence was recorded with
a custom-made epifluorescence microscope equipped with a 10x water-immersion objective

(UMPLFLN, Olympus), a set of GFP emission-excitation filters (FGL400, MD498, MF525-9,

Thorlabs) and a mounted blue LED controlled by a driver (MWWHL4, LEDD1B, Thorlabs).

Images were collected at 5 Hz via a custom-written Python script using the Pymba wrapper for

interfacing with the camera (Mako G319B, Allied vision).

Simultaneous recording of CT_{max} and depolarization in freely swimming larvae

Calcium imaging in freely swimming fish was used to record CT_{max} and brain-wide

depolarization in the same individuals during one trial. For the imaging of freely swimming

fish, the double-walled glass heating-mantle was placed under an epifluorescence microscope

(AxioImager, ZEISS) equipped with a megapixel camera (AxioCam 506, ZEISS). The larva

was swimming at the centre of the arena, within a 13 mm diameter region delimited with a

nylon mesh, to match the limited field of view of the microscope (14 x 14 mm). Time-lapse of

fluorescent images were recorded at 5 Hz using the Zen software (ZEN Lite Blue, ZEISS).

Since the recordings were made in the dark, the experimenter watched the live recording

display of the recording on the computer screen to visually guide the pokes to the fish tail. The

nylon mesh created a small thermal gradient in the arena from the central part to the outer area

outside the mesh. During the experimental trials, the temperature was recorded outside the

mesh to ensure a full view of the fish. The temperature gradient was thoroughly quantified in

an additional experiment over six heat ramps, by recording the temperature in the arena centre

and outside the mesh near the outer wall of the arena with two thermocouples at each location.

The gradient was accounted for to calculate the water temperature the fish was exposed to.

Calcium imaging data analyses

Calcium imaging recordings on embedded fish were corrected for slow drift due to agarose

expansion using the Fiji's (Schindelin *et al.*, 2012) Template matching plugin (*Align slices in*

stack). The regions of interest corresponding to the whole brain, the telencephalon and the

brainstem were manually segmented. The raw calcium fluorescence signal was calculated by

averaging all pixels within a region. To reproducibly detect the brain-wide depolarization onset

time across fish, the whole brain raw fluorescence signal was first processed to filter out the

calcium events using a 2^{nd} order Butterworth filter (low pass, 0.01 Hz cut-off frequency). The

average value and standard deviation of the filtered trace's time derivative were calculated

during the 17 min preceding the approximate onset of the depolarization. The depolarization

onset was set when the time derivative exceeded the average baseline value by more than five

standard deviation. For calcium event detection during the period preceding the brain-wide

depolarization, the fluorescence change was calculated for each brain region using a sliding

window of the previous two minutes. The calcium events were detected automatically using
the MATLAB function *findpeak*.

Measuring brain-wide fluorescence during the freely swimming fluorescence assay was
cumbersome due to the fish moving and tilting. Thus, the depolarization onset temperature was
determined by two investigators, who independently selected the time at which the
fluorescence increased abruptly within the brainstem/diencephalon. The final depolarization
onset value was obtained by averaging both estimates (agreement between experimenters: 19
±4s out of 55 minutes).

Cardiac function and brain activity during heat ramping in agar-embedded larvae

Heat rate and neural activity were simultaneously recorded at 20 Hz in laterally mounted larvae,
under an epifluorescence microscope (AxioImager, ZEISS) equipped with a megapixel camera
(AxioCam 506, ZEISS). Recordings were aligned when needed using Fiji's Template matching
plugin. Whole brain and heart regions of interest were selected in Fiji. Fish whose heart was
out of focus during more than two recording intervals (1 fish out of 10), and frames where
movement artefacts could not be corrected were not included in the rest of the analysis.
Heartbeat frequency during a recording interval was calculated using Matlab's continuous
wavelet transform function (cwt).

Oxygen effects on CT_{max} and neural function during warming

CT_{max} assay with oxygen level manipulation

The CT_{max} setup described earlier was used to record the effect of oxygen manipulation on
CT_{max}. The arena was intermittently bubbled with pure oxygen or pure nitrogen to increase or
decrease oxygen levels, respectively. The bubbling flow rate was manually adjusted using
tubing clamps. A fibre optic oxygen probe (OXROB10, PyroScience) and a temperature probe
(TSUB21, PyroScience) connected to an oxygen and temperature meter (FireSting O2,
PyroScience) were placed in the chamber to monitor and record the oxygen levels during the
assay. The oxygen saturation level was displayed in real time using an oxygen logger software
(pyro oxygen logger, PyroScience) and kept at 150 %, or 60%, of air saturation during the
whole CT_{max} assay by manually adjusting the bubbling intensity.

Recording neural activity with oxygen level manipulation

The epifluorescence calcium imaging setup for embedded fish was used to measure neural
activity during heat ramping with oxygen manipulation. Fish were visually stimulated using
pulses of red light (2 second long, every 30 second) using a red LED positioned in front of the

fish. For control treatment, the water was bubbled with air through a modified hypodermic
needle and maintained at 100% oxygen of air saturation during the trials. To increase the
oxygen level during oxygen manipulation trials, oxygen was gently blown on the surface. To
decrease oxygen saturation, the water was bubbled with nitrogen. A bare fibre microsensor
(OXB430, PyroScience) and a temperature probe (TDIP15, PyroScience) were placed in the
chamber to monitor and record oxygen levels. Oxygen levels were adjusted to either 150%, or
60%, of air saturated water during the high and low oxygen treatments, respectively.

Calcium imaging data analyses
Depolarization onset (**Figure 5B**) and brainstem calcium events (**Figure 5C**) were calculated
as described above. For calcium events, fish in which the brainstem signal was weak and noisy
due to uneven mounting were not included in the analysis. One fish in the hyperoxic group
could not be included in the calculation of the recovery time (**Figure 5E**) because the recording
ended before the fluorescence reached baseline after the depolarization.

Statistical analyses

Data are reported in the text as mean and standard error (S.E.), unless stated otherwise. All
statistical analyses were conducted in R version 4.0.2 (R Core Team, 2020). We used linear
regression models on independent measurements and the assumptions of normality for
homoscedasticity of residual variance were assessed visually using residual plots. Alternative
models were considered if assumptions were violated. For responses recorded as repeated
measures on individual fish linear mixed-effects models were created using the *lmer* function
from the *lme4* package v.1.1-23 to account for fish identity as random effect. Akaike
information criterion was used to consider interactions in models with two predictor variables
and model assumptions were visually assessed using residual plots. The effects of predictor
variables are presented with effect size (β), their S.E. estimated by the respective models and
the R^2 for the fixed effects is presented. The significance of effects was considered with a *p*-
value criterion of $p<0.05$.

CT_{max} and behavioural response to heat ramping
To analyse the swimming velocity during heat ramping and in the control group a linear mixed-
effects model was performed with treatment and time step as predictor variables and the fish
identity as random factor. No interaction was included (ΔAIC=2). **Table S1** is a summary of
the model results and **Figure S1** displays the datapoints. The proportion of spiral swimming

events was analysed using a Chi-square test to handle the lack of variance in the control group.

A Wilcoxon rank sum test was used to test the treatment effect on loss of equilibrium response.

To test the difference in CT_{max} throughout development, a linear regression model was used

with CT_{max} as a function of the age. The CT_{max} value of one adult fish was an extreme outlier

and was not included in the analysis. Results from models with and without this fish are

included in the supplements (**Table S2**).

Temperature of CT_{max} and depolarization

The relationship between depolarization and CT_{max} response in freely swimming fish was

analysed with a linear mixed-effects model with temperature as response variable, a categorical

fixed effect of response type (CT_{max} or depolarization) and fish identity as a random factor

(**Table S3**).

Change in cardiac function during heat ramping

Effect of temperature on heart rate was tested with a linear mixed-effects model where fish

identity was defined as a random factor. The model was releveled to check for differences

between the different temperature steps and the *p*-value significance criterion was accordingly

reduced to p<0.001 for this analysis. Model results are presented in **Table S4**.

Effect of oxygen manipulation during on CT_{max} and depolarization

The effect of oxygen level on CT_{max} was analysed with a linear regression with CT_{max} as a

function of the hypoxia and hyperoxia treatments. Two outliers and two fish who were

euthanized after the assay were removed from the hyperoxic group. **Table S5** displays model

results including these outliers. The effect of oxygen on neural activity was assessed using

linear regression analyses. Depolarization temperature, amplitude of depolarization and

recovery after depolarization were tested as functions of the treatment groups hypoxia,

normoxia and hyperoxia. Frequency of calcium events for the 15 minutes preceding the brain-

wide depolarization was analysed with an interaction between time and treatment. One extreme

outlier in the hyperoxic group was excluded from the analyses. Results from models made with

and without this fish are presented in **Table S6**.

Authors contributions

AHA, PH, FJ & FK conceived the project. FJ & FK jointly supervised. AHA, PH & FK built

the experimental setups. AHA, PH, PK collected the data. FK wrote the software for data

acquisition and analyses. AHA, PH & FK analysed the data, AHA and FK prepared the figures

and AHA performed the statistical analyses. AHA & FK wrote the manuscript, with
contributions from all authors.

Acknowledgements

The authors wish to thank the staff working at the animal facility for fish care, the members of
the Animal Physiology section at the Department of Biology for scientific discussions, and
Marius Mæhlum for the heartbeat frequency analysis code.

Funding

This work was supported by the Research Council of Norway (FK: FRIPRO grant 262698, FJ:
262942)

SUPPLEMENT

Supplementary Figure S1. Average swimming speed of (**A**) five-day-old control (n=14, empty circles) and (**B**)
heat ramp fish (n=12, black circles) as a function of temperature The swimming speed decreased from 2.9±0.4 to
1.7±0.5 mm/s during the assay. There was no difference in the change in swimming speed between the control
and the heat ramp treatment (linear mixed-effects model, time: $\beta\pm$S.E.$=0.3\pm0.1$, $t_{(103)}=-2.5$, $p=0.013$; treatment:
$\beta\pm$S.E.$=0.8\pm0.6$, $t_{(24)}=1.2$, $p=0.23$, **Table S1**). Bars and error bars represent mean and S.E, respectively.

Supplementary Table S1. Results of the statistical test for swimming speed in larval zebrafish (**Figure S1**) during
heat ramping (0.3°C/min) and over an equally long period for control fish (without heat ramping), testing the
effect of treatment and of time/temperature. The mixed-effects model includes fish identity as a random factor.
The control treatment is in the intercept and the units are in mm/s.

	Swimming speed			
Parameter	*Estimate (β)*	*S.E.*	*t-value*	*p-value*
Intercept (Control)	2.84	0.53	5.31	**<0.001**
Treatment (Heat ramp)	0.76	0.62	1.22	0.221
Time or temperature interval	-0.28	0.11	-2.53	**0.012**
Random Effects				
σ^2	3.08			
τ_{00} FishID	1.88			
N FishID	26			
Observations	130			
Marginal R^2 / Conditional R^2	0.056 / 0.414			

Supplementary Table S2. Results for the statistical test for CT_{max} in five-day-old, nine-day-old and adult zebrafish (**Figure 1E**) testing the effect of life-stage on the upper thermal limit. Model 1 is a linear regression model including all fish whereas in Model 2 one extreme outlier (<Q1-3*IQR) was removed from the adult group. The five-day-old group is in the intercept and units are in °C.

	Model 1				Model 2			
Parameter	*Estimate (β)*	*S.E.*	*t-value*	*p-value*	*Estimate (β)*	*S.E.*	*t-value*	*p-value*
Intercept (5dpf)	41.36	0.19	219.54	**<0.001**	41.36	0.17	250.19	**<0.001**
Age (9dpf)	-0.10	0.27	-0.39	0.698	-0.10	0.23	-0.45	0.659
Age (Adult)	-0.54	0.25	-2.16	**0.038**	-0.41	0.22	-1.84	0.073
Observations	40				39			
R^2 / R^2 adjusted	0.129 / 0.082				0.095 / 0.045			

Supplementary Table S3. Results of the statistical test on the relationship between CT_{max} and brain-wide depolarization (**Figure 3D**) testing the effect of event type on the recorded temperature. The mixed-effects model includes fish identity as a random factor. CT_{max} is in the intercept and units are in °C.

	CTmax and brain-wide depolarization			
Parameter	*Estimate (β)*	*S.E.*	*t-value*	*p-value*
Intercept (CTmax)	40.86	0.17	240.01	**<0.001**
Brain-wide depolarization	0.50	0.11	4.49	**<0.001**
Random Effects				
σ^2	0.04			
τ_{00} FishID	0.16			
N FishID	7			
Observations	14			
Marginal R^2 / Conditional R^2	0.249 / 0.840			

661 **Supplementary Table S4**. Results of the statistical test for the effect of temperature on heart rate in larval
662 zebrafish (**Figure 4C**) recorded for 1 min intervals during heat ramping. The mixed-effects model includes fish
663 identity as a random factor. The first temperature interval is in the intercept and the units are in Hz.

	Heart rate			
Parameter	*Estimate (β)*	*S.E.*	*t-value*	*p-value*
Intercept (28°C)	4.39	0.16	27.05	**<0.001**
Temperature (31°C)	0.91	0.18	4.92	**<0.001**
Temperature (34°C)	1.27	0.18	6.89	**<0.001**
Temperature (37°C)	0.36	0.19	1.91	0.057
Temperature (Depolarizaton)	0.26	0.18	1.40	0.161
Temperature (After)	-1.38	0.20	-6.98	**<0.001**
Random Effects				
σ^2	0.14			
$\tau_{00 \text{ FishID}}$	0.07			
N_{FishID}	9			
Observations	50			
Marginal R^2 / Conditional R^2	0.753 / 0.836			

666 **Supplementary Table S5.** Results of the statistical test for CT_{max} in hypo- and hyperoxia in larval zebrafish
667 (**Figure 5A**) testing the effect of oxygen level on the upper thermal limit. Model 1 is a linear regression model
668 excluding two extreme outliers (<Q1-3*IQR) from the hyperoxia group whereas Model 2 includes all fish. The
669 hypoxia treatment is in the intercept and units are in °C.

	Model 1				**Model 2**			
Parameter	*Estimate (β)*	*S.E.*	*t-value*	*p-value*	*Estimate (β)*	*S.E.*	*t-value*	*p-value*
Intercept (Hypoxia)	39.19	0.19	209.48	**<0.001**	39.19	0.40	97.26	**<0.001**
Hyperoxia	0.91	0.29	3.19	**0.004**	0.08	0.59	0.14	0.888
Observations	28				30			
R^2 / R^2 adjusted	0.282 / 0.254				0.001 / -0.035			

672 **Supplementary Table S6.** Results of the statistical tests on 1) brain-wide depolarization onset, 2) amplitude of
673 depolarization, and 3) recovery time after depolarization in norm-, hypo- and hyperoxia in larval zebrafish (**Figure
674 5B, D, E**) testing the effect of oxygen level on attributes of the brain-wide depolarization. The normoxic treatment
675 (control) is in the intercept and units are in °C.

Parameter	Depolarization onset				Amplitude of depolarization				Recovery time			
	Estimate (β)	S.E.	t-value	p-value	Estimate (β)	S.E.	t-value	p-value	Estimate (β)	S.E.	t-value	p-value
Intercept (Control)	40.65	0.29	138.14	**<0.001**	88.17	13.22	6.67	**<0.001**	2.95	0.38	7.76	**<0.001**
Hypoxia	-1.83	0.38	-4.81	**<0.001**	-3.06	17.07	-0.18	0.860	2.02	0.49	4.10	**0.001**
Hyperoxia	1.30	0.39	3.34	**0.003**	-15.42	17.49	-0.88	0.389	-1.29	0.52	-2.49	**0.022**
Observations	23				23				22			
R^2 / R^2 adjusted	0.802 / 0.782				0.045 / -0.050				0.729 / 0.701			

Supplementary Table 7. Results of the statistical test for frequency of calcium events in norm-, hypo- and hyperoxia in larval zebrafish (**Figure 5C**) testing the effect of oxygen level on neural activity. The mixed-effects model includes an interaction between treatment and time and fish identity as a random factor. In Model 1 one fish from the hyperoxia treatment was excluded due to extremely high values during the last 15 minutes (>Q3-3*IQR) whereas Model 2 includes all fish. The normoxic treatment (control) is in the intercept and the units are in event/min.

Parameter	Model 1				Model 2			
	Estimate (β)	S.E.	t-value	p-value	Estimate (β)	S.E.	t-value	p-value
Intercept (Control)	0.30	0.23	1.30	0.195	0.30	0.34	0.89	0.371
Time	0.08	0.02	3.90	**<0.001**	0.08	0.03	3.05	**0.002**
Treatment (Hypoxia)	0.11	0.32	0.34	0.737	0.11	0.46	0.24	0.814
Treatment (Hyperoxia)	-0.07	0.33	-0.21	0.835	0.03	0.46	0.08	0.940
Time*Treatment (Hypoxia)	-0.00	0.03	-0.02	0.982	-0.00	0.03	-0.02	0.985
Time*Treatment (Hyperoxia)	0.09	0.03	3.25	**0.001**	0.12	0.03	3.46	**0.001**
Random Effects								
σ^2	0.53				0.87			
τ_{00}	0.11 FishID				0.30 FishID			
N	16 FishID				17 FishID			
Observations	235				250			
Marginal R^2 / Conditional R^2	0.332 / 0.447				0.307 / 0.484			

Supplementary Movie S1. Spiral swimming in a five-day-old zebrafish heat ramp larva. Speed 1X.

Supplementary Movie S2. Loss of equilibrium in the same heat ramp larva as in Supplementary Movie 1. Speed 1X.

Supplementary Movie S3. Representative illustration of the depolarization spreading in the brain of a five-day-old *Tg(elavl3:GCaMP6s)* larva exposed to a heat ramp (same fish as in Figure 2E). Speed 1X.

REFERENCES

Andersen, M.K., Jensen, N.J.S., Robertson, R.M., & Overgaard, J. (2018) Central nervous system shutdown underlies acute cold tolerance in tropical and temperate *Drosophila* species. *The Journal of Experimental Biology*, **221**, jeb179598.

Åsheim, E., Andreassen, A., Morgan, R., & Jutfelt, F. (2020) Rapid-warming tolerance correlates with tolerance to slow warming but not growth at non-optimal temperatures in zebrafish. *Journal of Experimental Biology*,.

Barth, A.M.I. & Mody, I. (2011) Changes in Hippocampal Neuronal Activity During and After Unilateral Selective Hippocampal Ischemia In Vivo. *Journal of Neuroscience*, **31**, 851–860.

Brijs, J., Jutfelt, F., Clark, T.D., Gräns, A., Ekström, A., & Sandblom, E. (2015) Experimental manipulations of tissue oxygen supply do not affect warming tolerance of European perch. *Journal of Experimental Biology*, **218**, 2448–2454.

Buzsaki, G., Freund, T.F., Bayardo, F., & Somogyi, P. (1989) Ischemia-induced changes in the electrical activity of the hippocampus. *Experimental Brain Research*, **78**.

Clark, T.D., Sandblom, E., & Jutfelt, F. (2013a) Aerobic scope measurements of fishes in an era of climate change: respirometry, relevance and recommendations. *Journal of Experimental Biology*, **216**, 2771–2782.

Clark, T.D., Sandblom, E., & Jutfelt, F. (2013b) Response to Farrell and to Pörtner and Giomi. *Journal of Experimental Biology*, **216**, 4495–4497.

Deutsch, C.A., Tewksbury, J.J., Huey, R.B., Sheldon, K.S., Ghalambor, C.K., Haak, D.C., & Martin, P.R. (2008) Impacts of climate warming on terrestrial ectotherms across latitude. *PNAS*, **105**, 6668–6672.

Diaz Verdugo, C., Myren-Svelstad, S., Aydin, E., Van Hoeymissen, E., Deneubourg, C., Vanderhaeghe, S., Vancraeynest, J., Pelgrims, R., Cosacak, M.I., Muto, A., Kizil, C., Kawakami, K., Jurisch-Yaksi, N., & Yaksi, E. (2019) Glia-neuron interactions underlie state transitions to generalized seizures. *Nature Communications*, **10**.

Dube, C., Chen, K., Eghbal- Ahmadi, M., Brunson, K., Soltesz, I., & Baram, T.Z. (2000) Prolonged febrile seizures in the immature rat model enhance hippocampal excitability long term. *Annals of Neurology*, **47**, 336–344.

Ekström, A., Brijs, J., Clark, T.D., Gräns, A., Jutfelt, F., & Sandblom, E. (2016) Cardiac oxygen limitation during an acute thermal challenge in the European perch: effects of chronic environmental warming and experimental hyperoxia. *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology*, **311**, R440–R449.

Ern, R. (2019) A mechanistic oxygen- and temperature-limited metabolic niche framework. *Philosophical Transactions of the Royal Society B: Biological Sciences*, **374**, 20180540.

Ern, R., Huong, D.T.T., Phuong, N.T., Madsen, P.T., Wang, T., & Bayley, M. (2015) Some like it hot: Thermal tolerance and oxygen supply capacity in two eurythermal crustaceans. *Scientific Reports*, **5**, 10743.

Ern, R., Huong, D.T.T., Phuong, N.T., Wang, T., & Bayley, M. (2014) Oxygen delivery does not limit thermal tolerance in a tropical eurythermal crustacean. *The Journal of Experimental Biology*, 6.

Ern, R., Norin, T., Gamperl, A.K., & Esbaugh, A.J. (2016) Oxygen dependence of upper thermal limits in fishes. *Journal of Experimental Biology*, **219**, 3376–3383.

Farrell, A. p (2007) Cardiorespiratory performance during prolonged swimming tests with salmonids: a perspective on temperature effects and potential analytical pitfalls. *Philosophical Transactions of the Royal Society B: Biological Sciences*, **362**, 2017–2030.

Farrell, A.P. (2009) Environment, antecedents and climate change: lessons from the study of temperature physiology and river migration of salmonids. *Journal of Experimental Biology*, **212**, 3771–3780.

Friedlander, M.J., Kotchabhakdi, N., & Prosser, C.L. (1976) Effects of cold and heat on behavior and cerebellar function in goldfish. *J. Comp. Physiol.*, **112**, 19–45.

Genin, A., Levy, L., Sharon, G., Raitsos, D.E., & Diamant, A. (2020) Rapid onsets of warming events trigger mass mortality of coral reef fish. *PNAS*, **117**, 25378–25385.

Giomi, F., Barausse, A., Duarte, C.M., Booth, J., Agusti, S., Saderne, V., Anton, A., Daffonchio, D., & Fusi, M. (2019) Oxygen supersaturation protects coastal marine fauna from ocean warming. *Science Advances*, **5**, eaax1814.

Gollock, M.J., Currie, S., Petersen, L.H., & Gamperl, A.K. (2006) Cardiovascular and haematological responses of Atlantic cod (Gadus morhua) to acute temperature increase. *Journal of Experimental Biology*, **209**, 2961–2970.

Grans, A., Jutfelt, F., Sandblom, E., Jonsson, E., Wiklander, K., Seth, H., Olsson, C., Dupont, S., Ortega-Martinez, O., Einarsdottir, I., Bjornsson, B.T., Sundell, K., & Axelsson, M. (2014)

Aerobic scope fails to explain the detrimental effects on growth resulting from warming and elevated CO2 in Atlantic halibut. *Journal of Experimental Biology*, **217**, 711–717.

Hunt, R.F., Hortopan, G.A., Gillespie, A., & Baraban, S.C. (2012) A novel zebrafish model of hyperthermia-induced seizures reveals a role for TRPV4 channels and NMDA-type glutamate receptors. *Experimental Neurology*, **237**, 199–206.

Jørgensen, L.B., Robertson, R.M., & Overgaard, J. (2020) Neural dysfunction correlates with heat coma and CT $_{max}$ in *Drosophila* but does not set the boundaries for heat stress survival. *J Exp Biol*, **223**, jeb218750.

Jutfelt, F., Gräns, A., Jönsson, E., Wiklander, K., Seth, H., Olsson, C., Dupont, S., Ortega-Martinez, O., Sundell, K., Axelsson, M., & Sandblom, E. (2014) Response to 'How and how not to investigate the oxygen and capacity limitation of thermal tolerance (OCLTT) and aerobic scope – remarks on the article by Gräns et al.' *Journal of Experimental Biology*, **217**, 4433–4435.

Jutfelt, F., Norin, T., Ern, R., Overgaard, J., Wang, T., McKenzie, D.J., Lefevre, S., Nilsson, G.E., Metcalfe, N.B., Hickey, A.J.R., Brijs, J., Speers-Roesch, B., Roche, D.G., Gamperl, A.K., Raby, G.D., Morgan, R., Esbaugh, A.J., Gräns, A., Axelsson, M., Ekström, A., Sandblom, E., Binning, S.A., Hicks, J.W., Seebacher, F., Jørgensen, C., Killen, S.S., Schulte, P.M., & Clark, T.D. (2018) Oxygen- and capacity-limited thermal tolerance: blurring ecology and physiology. *Journal of Experimental Biology*, **221**.

Jutfelt, F., Roche, D.G., Clark, T.D., Norin, T., Binning, S.A., Speers-Roesch, B., Amcoff, M., Morgan, R., Andreassen, A.H., & Sundin, J. (2019) Brain cooling marginally increases acute upper thermal tolerance in Atlantic cod. *Journal of Experimental Biology*, **222**.

le Feber, J., Tzafi Pavlidou, S., Erkamp, N., van Putten, M.J.A.M., & Hofmeijer, J. (2016) Progression of Neuronal Damage in an In Vitro Model of the Ischemic Penumbra. *PLOS ONE*, **11**, e0147231.

Lefevre, S. (2016) Are global warming and ocean acidification conspiring against marine ectotherms? A meta-analysis of the respiratory effects of elevated temperature, high CO2 and their interaction. *Conserv Physiol*, **4**.

Lister, J.A., Robertson, C.P., Lepage, T., Johnson, S.L., & Raible, D.W. (1999) nacre encodes a zebrafish microphthalmia-related protein that regulates neural-crest-derived pigment cell fate. *Development*, **126**, 3757–3767.

Liu, J. & Baraban, S.C. (2019) Network Properties Revealed during Multi-Scale Calcium Imaging of Seizure Activity in Zebrafish. *eneuro*, **6**, ENEURO.0041-19.2019.

Lutterschmidt, W.I. & Hutchison, V.H. (2011) The critical thermal maximum: history and critique. *Canadian Journal of Zoology*,.

Marder, E., Haddad, S.A., Goeritz, M.L., Rosenbaum, P., & Kispersky, T. (2015) How can motor systems retain performance over a wide temperature range? Lessons from the crustacean stomatogastric nervous system. *J Comp Physiol A*, **201**, 851–856.

McArley, T.J., Hickey, A.J.R., & Herbert, N.A. (2018) Hyperoxia increases maximum oxygen consumption and aerobic scope of intertidal fish facing acutely high temperatures. *The Journal of Experimental Biology*, **221**, jeb189993.

McArley, T.J., Sandblom, E., & Herbert, N.A. (2020) Fish and hyperoxia—From cardiorespiratory and biochemical adjustments to aquaculture and ecophysiology implications. *Fish and Fisheries*,.

Morgan, R., Finnøen, M.H., & Jutfelt, F. (2018) CTmax is repeatable and doesn't reduce growth in zebrafish. *Scientific Reports*, **8**, 7099.

Moyano, M., Candebat, C., Ruhbaum, Y., Álvarez-Fernández, S., Claireaux, G., Zambonino-Infante, J.-L., & Peck, M.A. (2017) Effects of warming rate, acclimation temperature and ontogeny on the critical thermal maximum of temperate marine fish larvae. *PLOS ONE*, **12**, e0179928.

Muto, A., Lal, P., Ailani, D., Abe, G., Itoh, M., & Kawakami, K. (2017) Activation of the hypothalamic feeding centre upon visual prey detection. *Nature Communications*, **8**.

Overgaard, J., Andersen, J.L., Findsen, A., Pedersen, P.B.M., Hansen, K., Ozolina, K., & Wang, T. (2012) Aerobic scope and cardiovascular oxygen transport is not compromised at high temperatures in the toad Rhinella marina. *Journal of Experimental Biology*, **215**, 3519–3526.

Perry, A.L., Low, P.J., Ellis, J.R., & Reynolds, J.D. (2005) Climate Change and Distribution Shifts in Marine Fishes. *Science*, **308**, 1912–1915.

Pinsky, M.L., Eikeset, A.M., McCauley, D.J., Payne, J.L., & Sunday, J.M. (2019) Greater vulnerability
to warming of marine versus terrestrial ectotherms. *Nature*, **569**, 108–111.
Pörtner, H.O. (2002) Climate variations and the physiological basis of temperature dependent
biogeography: systemic to molecular hierarchy of thermal tolerance in animals. *Comparative*
Biochemistry and Physiology Part A: Molecular & Integrative Physiology, **132**, 739–761.
Pörtner, H.-O., Bock, C., & Mark, F.C. (2017) Oxygen- and capacity-limited thermal tolerance:
bridging ecology and physiology. *J Exp Biol*, **220**, 2685–2696.
Pörtner, H.O. & Knust, R. (2007) Climate Change Affects Marine Fishes Through the Oxygen
Limitation of Thermal Tolerance. *Science*, **315**, 95–97.
R Core Team (2020) *R: A Language and Environment for Statistical Computing*. R Foundation for
Statistical Computing, Vienna, Austria.
Robertson, R.M. & Money, T.G. (2012) Temperature and neuronal circuit function: compensation,
tuning and tolerance. *Current Opinion in Neurobiology*, **22**, 724–734.
Sawant-Pokam, P.M., Suryavanshi, P., Mendez, J.M., Dudek, F.E., & Brennan, K.C. (2016)
Mechanisms of Neuronal Silencing After Cortical Spreading Depression. *Cerebral Cortex*, **27**.
Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S.,
Rueden, C., Saalfeld, S., Schmid, B., Tinevez, J.-Y., White, D.J., Hartenstein, V., Eliceiri, K.,
Tomancak, P., & Cardona, A. (2012) Fiji: an open-source platform for biological-image
analysis. *Nature Methods*, **9**, 676–682.
Schulte, P.M. (2015) The effects of temperature on aerobic metabolism: towards a mechanistic
understanding of the responses of ectotherms to a changing environment. *Journal of*
Experimental Biology, **218**, 1856–1866.
Seneviratne, S.I., Donat, M.G., Mueller, B., & Alexander, L.V. (2014) No pause in the increase of hot
temperature extremes. *Nature Climate Change*, **4**, 161–163.
Sherman, E. & Levitis, D. (2003) Heat hardening as a function of developmental stage in larval and
juvenile Bufo americanus and Xenopus laevis. *Journal of Thermal Biology*, **28**, 373–380.
Shinnar, S. & Glauser, T.A. (2002) Febrile Seizures. *J Child Neurol*, **17**, S44–S52.
Spong, K.E., Andrew, R.D., & Robertson, R.M. (2016) Mechanisms of spreading depolarization in
vertebrate and insect central nervous systems. *J Neurophysiol*, **116**, 1117–1127.
Stillman, J.H. (2019) Heat Waves, the New Normal: Summertime Temperature Extremes Will Impact
Animals, Ecosystems, and Human Communities. *Physiology*, **34**, 86–100.
Sunday, J.M., Bates, A.E., & Dulvy, N.K. (2012) Thermal tolerance and the global redistribution of
animals. *Nature Climate Change*, **2**, 686–690.
Tang, L.S., Taylor, A.L., Rinberg, A., & Marder, E. (2012) Robustness of a Rhythmic Circuit to Short-
and Long-Term Temperature Changes. *Journal of Neuroscience*, **32**, 10075–10085.
Verberk, W.C.E.P., Leuven, R.S.E.W., Velde, G., & Gabel, F. (2018) Thermal limits in native and alien
freshwater peracarid Crustacea: The role of habitat use and oxygen limitation. *Functional*
Ecology, **32**, 926–936.
Verberk, W.C.E.P., Overgaard, J., Ern, R., Bayley, M., Wang, T., Boardman, L., & Terblanche, J.S.
(2016) Does oxygen limit thermal tolerance in arthropods? A critical review of current
evidence. *Comparative Biochemistry and Physiology Part A: Molecular & Integrative*
Physiology, **192**, 64–78.
Vladimirov, N., Mu, Y., Kawashima, T., Bennett, D.V., Yang, C.-T., Looger, L.L., Keller, P.J.,
Freeman, J., & Ahrens, M.B. (2014) Light-sheet functional imaging in fictively behaving
zebrafish. *Nature Methods*, **11**, 883–884.
Wang, T., Lefevre, S., Iversen, N.K., Findorf, I., Buchanan, R., & McKenzie, D.J. (2014) Anaemia only
causes a small reduction in the upper critical temperature of sea bass: is oxygen delivery the
limiting factor for tolerance of acute warming in fishes? *Journal of Experimental Biology*, **217**,
4275–4278.
Wenzel, M., Hamm, J.P., Peterka, D.S., & Yuste, R. (2017) Reliable and Elastic Propagation of Cortical
Seizures In Vivo. *Cell Reports*, **19**, 2681–2693.
Woitzik, J., Hecht, N., Pinczolits, A., Sandow, N., Major, S., Winkler, M.K.L., Weber-Carstens, S.,
Dohmen, C., Graf, R., Strong, A.J., Dreier, J.P., Vajkoczy, P., & For the COSBID study group
(2013) Propagation of cortical spreading depolarization in the human cortex after malignant
stroke. *Neurology*, **80**, 1095–1102.