Growth-mediated negative feedback shapes quantitative antibiotic response

S. Andreas Angermayr[1,2,#], Guillaume Chevereau[3], Tobias Bollenbach[1,4*]

[1]Institute for Biological Physics, University of Cologne, Cologne, Germany

[2]Institute of Science and Technology Austria, Klosterneuburg, Austria

[3]INSA de Strasbourg, Strasbourg, France

[4]Center for Data and Simulation Science, University of Cologne, Cologne, Germany

[#]Present address: CeMM Research Center for Molecular Medicine of the Austrian Academy of Sciences, Vienna, Austria.

[*]Corresponding author, t.bollenbach@uni-koeln.de

Abstract

Dose-response relationships are a general concept for quantitatively describing biological systems across multiple scales, from the molecular to the whole-cell level. A clinically relevant example is the bacterial growth response to antibiotics, which is routinely characterized by dose-response curves. The shape of the dose-response curve varies drastically between antibiotics and plays a key role for treatments, drug interactions, and resistance evolution. However, the mechanisms shaping the dose-response curve remain largely unclear. Here, we show in *Escherichia coli* that the distinctively shallow dose-response curve of the antibiotic trimethoprim is caused by a negative growth-mediated feedback loop: Trimethoprim slows growth, which in turn weakens the effect of this antibiotic. At the molecular level, this feedback is caused by the upregulation of the drug target dihydrofolate reductase (FolA/DHFR). We show that this upregulation is not a specific response to trimethoprim but follows a universal trend line that depends only on growth rate, irrespective of its cause. Rewiring the feedback loop alters the dose-response curve in a predictable manner, which we corroborate with a mathematical model. Our results indicate that growth-mediated feedback loops shape drug responses and could be exploited to design evolutionary traps that enable selection against drug resistance.

Introduction

Dose-response curves are a central concept in systems biology and essential for understanding emergent nonlinear phenomena at different scales. A prime example is bacterial gene regulation where cooperativity of transcription factor binding to promoter regions governs the steepness of dose-response curves that characterize gene expression as a function of transcription factor concentration (Bintu *et al*, 2005). The steepness of such transcription factor dose-response curves ultimately determines whether feedback loops in genetic circuits can produce biologically relevant functions such as bistability or oscillations (Elowitz & Leibler, 2000; Gardner *et al*, 2000). At the population level, the bacterial response to antibiotics is captured by similar dose-response curves that quantify the dependence of growth rate on drug concentration. Antibiotic dose-response curves are routinely measured to characterize antibiotic susceptibility via the minimal inhibitory concentration (MIC) or the concentration leading to 50% growth inhibition (IC_{50}), two classic quantities to describe antibiotic efficacy. However, the quantitative shape of the antibiotic dose-response curve – especially its steepness – and its implications are underappreciated.

The steepness of the dose-response curve varies drastically between antibiotics. For many antibiotics, the growth rate drops gradually from high to low as the drug concentration is increased (Fig. 1A); in particular, this is the case for antibiotics targeting DNA replication at the gyrase (e.g. ciprofloxacin) or antibiotics targeting translation at the ribosome (e.g. tetracycline). Beta-lactams like mecillinam (an antibiotic targeting cell wall biosynthesis at a penicillin binding protein) have extremely steep dose-response curves where just a slight relative increase in drug concentration – by about two-fold – causes an abrupt transition from full-speed growth to near-zero net growth (Fig. 1A). At the other end of the spectrum, the folic acid synthesis inhibitor trimethoprim (TMP) has an extremely shallow dose-response curve (Palmer & Kishony, 2014; Rodrigues *et al*, 2016; Russ & Kishony, 2018; Chevereau *et al*, 2015): Reducing growth from full speed to zero with TMP requires a more than 100-fold increase in drug concentration (Fig. 1A). In general, dose-response curves are well approximated by Hill functions and the Hill slope n ("dose-sensitivity") is a quantitative measure of their steepness (Regoes *et al*, 2004; Chou & Talalay, 1983; Chevereau *et al*, 2015): TMP has $n \approx$ 1.1, while most antibiotics fall in the range $1.8 \leq n \leq 3.5$, and $n > 6$ (Fig. 1A).

Fig. 1| Trimethoprim exhibits an extremely shallow dose response curve and its efficacy correlates strongly with growth rate. (A) Dose-response curves (normalized growth rate as a function of drug concentration) for different antibiotics. Growth rate was measured via optical density measurements over time (Methods). Antibiotics used: Trimethoprim (TMP), tetracycline (TET), chloramphenicol (CHL), ciprofloxacin (CPR), lincomycin (LIN), nitrofurantoin (NIT), and mecillinam (MEC). The TMP dose-response curve (dark blue) is by far the shallowest. Lines are fits of the Hill function $g(c) = \frac{1}{1+\left(\frac{c}{IC_{50}}\right)^n}$ to the data. Drug concentrations were arbitrarily rescaled to better visualize dose-response curve steepness. Error bars show standard deviation of 12 replicates. (B) Dose-response curves calculated from a mathematical model that captures growth-mediated feedback. Negative feedback (blue) renders the dose-response curve shallower than in the absence of feedback (gray); positive feedback (red) steepens the dose-response curve. Parameters are $n = 2$, $\alpha = -1$ for negative feedback and $\alpha = 5$ for positive feedback (see main text); drug concentrations are normalized to the IC_{50} in the absence of growth mediated feedback ($\alpha = 0$). (C) Density scatterplot showing growth response to TMP versus normalized drug-free growth rate for genome-wide gene deletion strains (Baba et al, 2006). These gene deletion strains exhibit diverse growth rates, offering an unbiased way to test the relation between the drug-free growth rate and the response to antibiotics. Response is defined as growth rate in the presence of TMP normalized to the drug-free growth rate of the respective deletion strain. TMP was used at a fixed concentration that inhibits wild type growth by about 30% (Chevereau et al, 2015). Spearman correlation coefficient ρ_S is shown. (D) Bar chart showing negative Spearman correlation coefficients $-\rho_S$ compared across antibiotics (Supplementary Fig. 1). Error bars show bootstrap standard error of ρ_S. TMP (blue) exhibits by far the strongest negative correlation, indicating a growth-mediated negative feedback loop.

The steepness of the dose-response curve strongly affects the evolution of resistance by spontaneous mutations (Hermsen *et al*, 2012; Chevereau *et al*, 2015). Resistance mutations that slightly increase the MIC provide greater fitness benefits for drugs with a steep dose-response curve compared to drugs with a shallow curve, implying a greater chance to fix in the population. Thus, all else being equal, the rate of resistance evolution for a drug increases with the steepness of its dose-response curve – a trend that is observed in evolution experiments (Chevereau *et al*, 2015). This effect is strongest for drug concentrations near the IC_{50}, which occur for example when populations

of motile bacteria evolve resistance in spatial drug gradients (Hol *et al*, 2016; Baym *et al*, 2016).

Despite their fundamental relevance for resistance evolution and bacterial responses to antibiotics,

the mechanisms that shape the dose-response curve are largely unknown.

Feedback loops mediated by growth rate may play a key role in shaping the dose-response curve

(Deris *et al*, 2013; Greulich *et al*, 2015). The action of antibiotics affects bacterial growth but the

inverse is also true: Slower growing bacteria are less rapidly killed by antibiotics targeting cell wall

biosynthesis (β-lactams) (Tuomanen *et al*, 1986; Lee *et al*, 2018) and non-growing (persister) cells

are fully protected from many antibiotics (Balaban *et al*, 2004), offering a possibility to evade

antibiotic treatments. Slower growth caused by nutrient limitation also affects the bacterial

susceptibility to ribosome-targeting antibiotics (Greulich *et al*, 2015). In engineered strains

expressing a constitutive resistance gene, a positive feedback loop leads to high dose-sensitivity and

even bistability (i.e. co-existence of growing and non-growing cells) in the presence of the ribosome-

targeting antibiotic chloramphenicol (Deris *et al*, 2013). Positive feedback occurs as faster growth

leads to the upregulation of the resistance enzyme, which in turn enables even faster growth.

Growth-mediated feedback loops could more generally explain the drastic differences in dose-

sensitivity between antibiotics (Fig. 1A) with positive feedback producing higher (Deris *et al*, 2013)

and negative feedback lower dose-sensitivity. However, such feedback loops shaping the dose-

response curve of sensitive wild-type bacteria have not yet been characterized.

Here, we establish that negative growth-mediated feedback produces an extremely shallow drug

dose-response curve. Focusing on TMP, we vary bacterial growth rates by diverse environmental and

genetic perturbations and show that slower growth generally lowers the susceptibility of *Escherichia*

coli to this antibiotic. The molecular origin of this phenomenon lies in the expression of the drug

target, which is upregulated in response to TMP but also when growth rate is lowered by other

means: TMP lowers growth, which in turn reduces susceptibility to TMP. We show that synthetically

reversing this feedback loop can drastically steepen the dose-response curve. The negative feedback

loop leads to a seemingly paradoxical situation where adding the antibiotic can even enhance

growth under extreme nutrient limitation. Such growth-mediated feedback loops in drug responses

could be used to design evolutionary traps that invert selection for resistance.

Results

A minimal mathematical model shows that growth-mediated feedback loops affect the dose-

sensitivity of drugs

We hypothesized that a growth-mediated negative feedback loop could explain the extreme

shallowness of the dose-response curve of TMP (Fig. 1A). As an antibiotic, TMP lowers bacterial

growth (by inhibiting dihydrofolate reductase, DHFR, encoded by *folA*). If a lower growth rate in turn

protects bacteria from TMP, the dose-response curve should become shallower. To test this idea, we

began by developing a mathematical model of bacterial growth in the presence of an antibiotic. We

studied a minimal model in which growth-mediated feedback is introduced by making drug efficacy

dependent on growth rate. The dose-response curve of antibiotics is captured by a Hill function

$g(c) = \frac{g_0}{1+(c/c_0)^n}$, where g_0 is the growth rate in the absence of the drug, n is the dose-sensitivity, c_0

is the drug concentration that inhibits growth by 50%, and c is the drug concentration, which is

controlled in experiments. To capture effects of growth rate on drug efficacy, we replace c with an

effective drug concentration c_{eff} that depends on the externally controlled drug concentration c_{ext}

and on the growth rate g, i.e. $c = c_{\text{eff}}(c_{\text{ext}}, g)$. This effective drug concentration captures that the

same external concentration can have different effects on the cell, e.g. if the intracellular drug

concentration or the expression level of the target is different. The resulting dose-response curve

$g(c_{\text{ext}})$ is then implicitly given by

$$g = \frac{g_0}{1+\left[\frac{c_{\text{eff}}(c_{\text{ext}}, g)}{c_0}\right]^n}. \tag{1}$$

Growth-mediated feedback is captured by a function c_{eff} that increases or decreases with the

growth rate. For simplicity, we assume that this dependence is linear, i.e. $c_{\text{eff}} = [1 + \alpha (1 -$

$g/g_0)]c_{\text{ext}}$, where $-1 \leq \alpha < 0$ corresponds to negative and $\alpha > 0$ to positive feedback. Solving

equation (1) confirms that negative growth-mediated feedback leads to a shallow dose-response

curve (Fig. 1B). In contrast, positive growth-mediated feedback leads to ultrasensitivity (Fig. 1B) and

can even produce bistability as previously reported (Elf *et al*, 2006; Deris *et al*, 2013). In this model,

the IC_{50} also changes with α: $\text{IC}_{50} = c_0/(1 + \alpha/2)$. These theoretical results show how growth-

mediated feedback loops can affect the dose-sensitivity of drugs in general.

Slower growth generally lowers susceptibility to TMP and steepens its dose-response curve

To test experimentally whether negative growth-mediated feedback underlies the shallow TMP

dose-response curve, we varied growth rate in several independent ways and investigated its effect

on TMP susceptibility. We first made use of a purely genetic way of varying growth. Specifically, we

exploited the growth rate variability resulting from genome-wide gene deletions to expose global

trends that are independent of the specific effects of individual gene deletions. Non-essential gene

deletions often reduce the drug-free growth rate – some by up to ~50% (Chevereau *et al*, 2015).

We re-analyzed a dataset of growth rates of ~4,000 *E. coli* gene deletion mutants under different

antibiotics representing common modes of action (Chevereau & Bollenbach, 2015); growth rates

were measured at concentrations that inhibit the reference strain by ~30%. While each gene can

have specific effects for each antibiotic (Nichols *et al*, 2011; Chevereau *et al*, 2015), most genes

should be unrelated to the drug mode of action. The global trend of drug susceptibility across all

gene deletion strains can thus reveal general consequences of growth inhibition, independent of the

specific cellular limitation causing the growth rate reduction.

Non-specific growth rate changes caused by gene deletions indicate that slower growth protects

E. coli from TMP. By correlating the drug-free growth rate of deletion strains with their growth rate

in the presence of drugs, we revealed dependencies of drug susceptibility on the drug-free growth

rate. The clearest trend emerged for TMP: Its relative effect on growth was weaker in gene deletion

strains that had lower growth rates in the absence of drugs (Spearman correlation $\rho_s = -0.6$;

Fig. 1C). Compared to other antibiotics, this effect was most pronounced for TMP (Fig. 1D;

Supplementary Fig. 1). Slower-growing mutants can even grow at increased TMP concentrations: Full

dose-response curve measurements for 78 arbitrary gene deletion mutants showed that the IC_{50} is

negatively correlated with the growth rate in the absence of drug for TMP ($\rho_s = -0.27, p = 0.019$)

but not significantly correlated for other antibiotics (Supplementary Fig. 2). Thus, TMP represents an

extreme case, both in terms of dose-sensitivity and in terms of susceptibility-dependence on growth

rate. Overall, these results suggest that slower growth generally lowers the susceptibility to TMP.

Slow growth also protects *E. coli* from other antibiotics, but to a far lesser extent. For the prodrug

nitrofurantoin (NIT) and the translation inhibitors tetracycline (TET) and chloramphenicol (CHL),

there was a weak negative correlation between the drug-free growth rate and that in the presence

of the drug ($\rho_s = -0.31$ for NIT, $\rho_s = -0.26$ for TET, $\rho_s = -0.21$ for CHL; Fig. 1D; Supplementary

Fig. 1). For the beta-lactam mecillinam (MEC), this trend was even weaker ($\rho_s = -0.14$) and for

ciprofloxacin (CPR) almost entirely absent ($\rho_s = -0.05$). Notably, the magnitude of this negative

correlation roughly decreases with increasing dose-sensitivity when compared across drugs

(Fig. 1A,D). This observation supports the notion that growth-mediated feedbacks are an important

contributor to the shape of the dose-response curve for TMP (Fig. 1B) and possibly also for other

antibiotics.

Reducing growth rate by other means like nutrient limitation or imposing a protein burden also

protects *E. coli* from TMP. First, we used glucose limitation in batch culture by adding a non-

metabolizable structural analog of glucose, α-methyl glucoside, in varying concentrations to the

growth medium. This analog competes with glucose for uptake into the cell, but unlike glucose it

cannot be utilized for growth (Hansen *et al*, 1975). Second, we used different carbon sources

(glucose, fructose, mannose, glycerol, and galactose) in the growth medium, which is a classic

strategy to test for growth-dependent effects (Bremer & Dennis, 2008). Third, we overexpressed a

gratuitous protein from an inducible promoter to burden the cells (Dong *et al*, 1995; Scott *et al*,

2010). These approaches have different physiological consequences, but they all reduce growth rate

in a gradual and controlled manner, while the maximal growth rate and the accessible dynamic

range of relative growth inhibition varies between them (Fig. 2). Collectively, they enable us to vary

growth rate over a wide range and identify general effects of growth rate, which occur

independently of the exact cause of the growth rate reduction. TMP inhibits growth less under

glucose limitation: Lowering growth rate by glucose limitation enabled bacteria to grow at slightly

increased TMP concentrations (Fig. 2B,C). This trend was even more pronounced when growth was

lowered by overexpressing a gratuitous protein – a truncated and inactive version of *tufB* (Dong *et*

al, 1995) expressed from a synthetic promoter $P_{LlacO-1}$ (Lutz & Bujard, 1997) induced by addition of

isopropyl β-D-1-thiogalactopyranoside (IPTG) (Fig. 2D,E,F). Reducing growth by using different

carbon sources in a minimal medium could also slightly protect bacteria from TMP, in particular for

glycerol (Fig. 2G,H,I). Changing carbon sources has modest effects, presumably because even the

highest growth rate (achieved with glucose only) is relatively low and the fold-change in growth is

considerably smaller than for glucose limitation (Fig. 2A,D,G). These effects did not occur to a

comparable extent for other antibiotics representing common modes of action (Fig. 2C,F,I);

however, gratuitous protein overexpression also lowered the susceptibility to mecillinam (MEC),

albeit to a lesser extent (Fig. 2F, Supplementary Fig. 6). The effects of growth rate changes were

clearly drug-specific and strongest for TMP.

Under severe glucose limitation (high ratios of α-methyl glucoside over glucose), which does not

support growth, the addition of TMP even rescued bacteria and enabled them to grow again

(Fig. 2B). This seemingly paradoxical phenomenon indicates that, under extreme nutrient limitation,

the antibiotic TMP can facilitate bacterial growth – perhaps the most drastic illustration of the close

interplay between growth rate and TMP susceptibility we observed.

Fig. 2|Slower growth generally lowers the efficacy of trimethoprim. (A) Growth rate under glucose limitation achieved by adding the non-metabolizable structural glucose analog α-methyl glucoside (αMG) at different ratios to glucose in a minimal medium (Methods). **(B)** Normalized growth rate (gray scale) from a checkerboard assay in a two-dimensional concentration gradient of TMP and αMG. Dashed black line shows contour line of 90% growth inhibition (IC_{90} line). Red arrow shows increase in IC_{90} as growth is lowered. **(C)** Fold-change in IC_{90} at αMG/glucose ratio 2.5 in assays as in B for different antibiotics (Supplementary Fig. 5). Lowering growth rate increases IC_{90} for TMP but not for other antibiotics. **(D)** Growth rate in rich medium (LB) under different levels of overexpression of a gratuitous protein from a T5-lac promoter; overexpression burden is controlled by IPTG concentration (Methods). **(E)** As B but for growth rate reduction by protein overexpression in a two-dimensional concentration gradient of TMP and IPTG. **(F)** Fold-change in IC_{90} at 1.25 mM IPTG in assays as in E for different antibiotics (Supplementary Fig. 6). Overexpression of unnecessary protein increases IC_{90} for TMP by almost five-fold; no comparable increase occurs for other antibiotics. **(G)** Growth rate in minimal medium containing different carbon sources (Methods): Glucose (GLU), fructose (FRU), mannose (MAN), galactose (GAL), and glycerol (GLY). **(H)** Normalized growth rates (gray scale) on different carbon sources (x-axis) at different TMP concentrations (y-axis). **(I)** Fold-change in IC_{90} in assays as in H for different antibiotics (Supplementary Fig. 7). Error bars in A, D show standard deviation from eight replicates, those in G from three replicates; day-to-day reproducibility of growth rate measurements is high (Supplementary Fig. 3). Error bars in C,F show standard deviation from three neighboring αMG/glucose ratios and IPTG concentrations centered at 2.5 and 1.25 mM, respectively. IPTG alone has no detectable effect on growth at these concentrations (Supplementary Fig. 4). Error bars in I show standard deviation from three replicates. Antibiotic abbreviations are as in Fig. 1. CHL was not used in the protein overexpression assay in F since the plasmid used for overexpression has a CHL-resistance marker (Methods). Sample growth curves are in Supplementary Fig. 11.

Lowering growth rate by changing temperature does not show a similar effect: The relative growth reduction by antibiotics remains the same at different temperatures (Supplementary Fig. 8). This is expected since the key physiological parameters that determine growth are invariant under temperature changes (Bremer & Dennis, 2008). This observation indicates that the observed effect has a biological – rather than an elementary physical – origin.

Slower growth increases the steepness of the TMP dose-response curve. We noticed that the extremely shallow dose-response curve of TMP (Fig. 1A) became steeper when growth was slowed

by glucose limitation (Fig. 3A): Halving the growth rate increased the dose-sensitivity from $n \approx 1.1 \pm$

0.2 to $n \approx 1.6 \pm 0.3$ (Fig. 3B). This steepening occurred similarly when growth was slowed by

gratuitous protein overexpression (Fig. 3C,D) or by changing the carbon source in the growth

medium (Fig. 3E,F). In addition to this change in steepness, the concentration at which TMP starts to

have an effect on growth is higher for slower-growing bacteria. However, once the effect of TMP

kicks in, the growth rate drops more rapidly with increasing TMP concentration. To better

understand this unexpected increase in dose-sensitivity resulting from slower growth, we next

aimed to elucidate the underlying mechanism of TMP's growth-rate-dependent action.

Fig. 3|Slower growth increases the steepness of the trimethoprim dose-response curve. (A) TMP dose-response curve in minimal medium with glucose as carbon source and at lower drug-free growth rate due to glucose limitation, achieved by increasing the αMG /glucose ratio from 0 (black) to 5 (gray). Glucose limitation results in a steeper dose-response curve. Lines show Hill function fits (cf. Fig. 1A). (B) Steepness of TMP dose-response curves (dose-sensitivity n) versus drug-free growth rate at different αMG concentrations (Methods). Numbers next to data points show αMG/glucose ratio. (C,D) As A,B but for growth limitation by gratuitous protein overexpression in rich growth medium (Methods). Inducing overexpression with IPTG at 10 mM (light gray) steepens the dose-response curve compared to no induction (black). Numbers next to data points in D show IPTG concentration in mM. (E,F) As A,B but for growth limitation by varying the carbon source in a minimal medium (Methods). Carbon sources as in Fig. 2. Growth rate error bars show standard deviation of three replicates; vertical error bars in B,D,F show standard deviation of parameter estimates from Hill function fit.

Growth-dependent regulation of the TMP drug target leads to a negative feedback loop that flattens the dose-response curve

Regulation of the drug target DHFR could cause the growth-rate-dependent efficacy of TMP. The

abundance of the target of TMP (DHFR/FolA) correlates with growth (Bershtein *et al*, 2013);

increasing its expression, e.g. by overexpressing *folA* from a plasmid, alleviates the effect of TMP on

growth (Palmer & Kishony, 2014). Accordingly, TMP resistance in the lab and in the clinic often

evolves by overexpressing *folA*, e.g. by mutating its promoter or by increasing gene copy number

(Toprak *et al*, 2012; Nyerges *et al*, 2018; Rood *et al*, 1980; Flensburg & Sköld, 1987; Baym *et al*,

2016). These phenomena suggest a plausible mechanism for the reduced susceptibility to TMP at

lower growth rates: We hypothesized that slower growth generally leads to increased *folA*

expression, which in turn partially protects bacteria from TMP (Soo *et al*, 2011; Palmer & Kishony,

2014) – a buffering mechanism against inhibition of FolA.

DHFR expression increases similarly in response to TMP and to other means of reducing growth rate.

Using a promoter-GFP reporter (Methods), we confirmed that *folA* expression increases in response

to TMP (Fig. 4A), as previously observed in whole populations (Bollenbach *et al*, 2009; Bershtein *et*

al, 2015; Rodrigues *et al*, 2016) and single cells (Mitosch *et al*, 2017). However, we noticed that *folA*

expression increases similarly when growth is slowed by glucose limitation (Fig. 4B). This observation

suggests that the upregulation of *folA* under TMP is not a specific response to target inhibition, but

rather a general response to reduced growth rate. Expression levels of constitutive genes are

generally expected to increase when the quality of the nutrient environment is lowered (Scott *et al*,

2010). While the *folA* promoter can be regulated by two transcription factors (TyrR (Yang *et al*, 2007)

and IHF (Keseler, 2004)) under certain conditions, it behaved similarly to a constitutive promoter in

these experiments. Indeed, *folA* expression across a two-dimensional concentration gradient of TMP

and the glucose analog varied (Supplementary Fig. 9) but was largely determined by growth rate

alone (Fig. 4C). Like constitutively expressed genes (Scott *et al*, 2010), *folA* expression followed a

general, approximately linear increase with decreasing growth rate, approaching a fixed maximum

level at zero growth (Fig. 4C). Since increased *folA* expression protects bacteria from TMP (Palmer &

Kishony, 2014), this mode of regulation results in a negative growth-mediated feedback loop: TMP

inhibits growth, which leads to the upregulation of its target, thereby attenuating its own efficacy.

Saturating growth-dependent regulation of the drug target can explain the steepening of the dose-

response curve at lower growth rates. Higher drug target expression at lower growth rates can

compensate for some of the target inhibition caused by TMP. This offers a plausible explanation as

to why the effect of TMP becomes apparent only at higher concentrations when the drug-free

growth rate is lower (Fig. 2). But how does slower growth steepen the TMP dose-response curve

(Fig. 3)? We noticed that *folA* expression at different drug-free growth rates converges to a fixed

value when TMP is added (Fig. 4D). In other words, the relative upregulation of *folA* in response to

TMP gets weaker with decreasing drug-free growth rate; it even disappears completely at the

highest glucose-analog concentrations (Fig. 4D). This convergence of *folA* expression in different

conditions may reflect that the promoter reaches its maximal induction level. At lower drug-free

growth rates, the promoter is already near its maximum expression level without TMP and saturates

quickly when TMP is added, resulting in weaker relative upregulation than at higher growth rates.

Consistent with this scenario, increasing *folA* expression at low growth rates is deleterious

(Supplementary Fig. 10). Thus, lower drug-free growth rates weaken – or even break – the growth-

mediated negative feedback loop, resulting in steeper dose-response curves.

Fig. 4 | Slower growth increases folA expression, irrespective of whether growth is reduced by trimethoprim or by nutrient
limitation. (A) *Dependence of growth rate (black) and folA expression (green) on TMP concentration. Schematic: FolA*
expression was measured using a promoter-GFP reporter inserted at a neutral site in the genome (Methods). ***(B)*** *Growth*
rate (black) and folA expression (green) in the absence of TMP at different growth rates achieved by different ratios of
αMG/glucose. ***(C)*** *Scatterplot of folA expression level with growth rate across all combinations of TMP concentrations and*
αMG/glucose ratios shown in A and B (Supplementary Fig. 9). Pearson's correlation coefficient ρ and p-value are shown. ***(D)***
Dependence of folA expression on TMP concentration at four different αMG/glucose ratios (0, 2.5, 5, and 10 as shown).
Darker green indicates greater αMG/glucose ratio. FolA expression converges approximately to the same level at high TMP
concentrations. Black line in A shows Hill function fit as in Fig. 1A; other lines show polynomial fits of first (C) or second
order (A,B) to guide the eye. Horizontal dotted green line shows folA expression level in the absence of TMP. Error bars
show standard deviation of three replicates.

Artificially breaking the growth-mediated feedback loop steepens the TMP dose-response curve

To corroborate that the shallowness of the TMP dose-response curve is due to a growth-mediated

negative feedback loop, we aimed to break this loop even under nutrient conditions that support

high drug-free growth rates. To this end, we constructed a synthetic strain in which the expression of

folA from its endogenous locus is controlled by an inducible promoter. We used a strain allowing

IPTG-mediated induction of *folA* and *folA-gfp,* respectively, with an expression level in the same

range as wild-type *folA* (Methods). Note that this alone does not eliminate the feedback loop since,

at constant inducer levels, expression from the inducible promoter can change with growth rate, similar to expression from the endogenous *folA* promoter. Nevertheless, we can use this synthetic strain to infer the shape of the TMP dose-response curve at constant *folA* expression by continuously varying the inducer concentration and measuring FolA levels. Specifically, we measured growth rate and *folA* expression using a FolA-GFP fusion protein across a two-dimensional concentration gradient of TMP and inducer (Fig. 5A; Methods). We then determined the growth rate as a function of TMP concentration on a path through this two-dimensional concentration space along which *folA* expression is constant. The resulting TMP dose-response curve at constant *folA* expression is steeper than in wild type ($n = 2.0 \pm 0.3$; Fig. 5B,C). It becomes even steeper for a positive feedback loop, which is inferred from a path through the two-dimensional concentration space along which *folA* expression decreases with increasing TMP concentration ($n = 5.0 \pm 0.9$; Fig. 5B,C). These results provide direct evidence that a negative growth-mediated feedback loop implemented by the regulation of the drug target causes the exceptional shallowness of the TMP dose-response curve.

Fig. 5|Breaking the growth-mediated negative feedback loop steepens the trimethoprim dose-response curve. (A) *Schematic: FolA expression is controlled by varying the IPTG concentration and measured by flow cytometry using a GFP fusion to FolA. Shades of green indicate different FolA expression levels. Wells encircled in red indicate how the effect of FolA down-regulation with increasing TMP concentration can be inferred; wells encircled in black illustrate the same for constant FolA expression.* ***(B)*** *Growth rate as a function of TMP concentration for different paths through IPTG-TMP concentration space as illustrated in A (Methods). Constant FolA is shown in black and FolA down-regulation in red. Wild type dose-response curve (blue line; fit from Fig. 4A) is shown for comparison.* ***(C)*** *Steepness of the dose response curve (quantified as dose-sensitivity n) for the three cases in B. Inset: Normalized FolA expression level as a function of TMP concentration for the three cases in B; WT (blue) shows fit from Fig. 4A; colors as in the bar chart and in B. Error bars in B show standard deviation of the measured growth rates used for interpolating the values shown (Methods). Error bars in C show standard deviation of parameter estimates from Hill function fit.*

Discussion

We showed that slower-growing bacteria are generally less affected by TMP, largely regardless of what causes their slower growth (Fig. 2). This phenomenon implies a growth-mediated negative feedback loop, which causes TMP's extremely shallow dose-response curve (Fig. 1A): TMP lowers growth, which in turn weakens the inhibitory effect of the drug. Mechanistically, this feedback loop

is rooted in the expression level of the drug target DHFR, which is upregulated with decreasing

growth rate (Fig. 4). Elimination or inversion of this feedback loop from negative to positive

drastically steepens the dose-response curve (Fig. 5). Together with recent work on ribosome-

targeting antibiotics (Deris *et al*, 2013), these results support a general role of growth-mediated

feedbacks in shaping antibiotic dose-response curves (Fig. 1B).

Consistent with this view, the steepness of the dose-response curve of antibiotics representing

different modes of action correlates roughly with the decrease in drug susceptibility under slower

growth (Fig. 1A,D). In particular, while their effect is less extreme than for TMP, the ribosome

inhibitors CHL and TET also exhibit relatively low dose-sensitivity and slightly reduced susceptibility

under slower growth (Fig. 1D, Supplementary Fig. 1). The mechanism underlying this weaker growth-

mediated negative feedback for CHL and TET resembles that for TMP since their drug target, the

ribosome, is upregulated in response to these drugs (Scott *et al*, 2010)—similar to DHFR in response

to TMP. The prodrug NIT is an outlier: It has a relatively steep dose-response curve (Fig. 1A) despite

being less susceptible under slower growth (Fig. 1D). This may be caused by additional (unknown)

mechanisms acting on top of the growth-mediated feedback we focus on here. Identifying the

molecular mechanisms underlying feedback loops or other phenomena that shape dose-response

curves will likely require detailed studies for each antibiotic or antibiotic class.

By using a mathematical model in which the effective drug concentration is growth-rate dependent,

we showed that growth-mediated feedbacks generally affect the shape of the dose-response curve.

For TMP, the effective drug concentration is reduced due to the upregulation of DHFR: The same

concentration of TMP has a weaker effect on growth at higher intracellular concentrations of DHFR.

For simplicity, we assumed that the effective drug concentration changes linearly with the growth

rate. This is in line with the experimental observation that DHFR is upregulated linearly with

decreasing growth rate, approaching a maximum at zero growth (Fig. 4C). Decoupling the DHFR level

from the growth rate by forcing it to a constant value, or even to decrease in response to TMP,

results in a steeper dose-response curve (Fig. 1B) in agreement with experimental observations

(Fig. 5B,C). The model further helps to rationalize why dose-response curves become steeper when

the drug-free growth rate is decreased, corresponding to a poorer nutrient environment. At very low

drug-free growth rates, the DHFR level becomes almost constant (Fig. 4D), effectively breaking the

negative feedback loop and thus steepening the dose-response curve, as observed experimentally

(Fig. 3). Note, that our model can explain relative changes in dose-sensitivity but it remains

challenging to explain its absolute value. Indeed, all antibiotics we studied have dose-sensitivities

$n > 1$ and most have $n > 2$ (Fig. 1A), which requires additional nonlinearities beyond the growth-

mediated one we focused on here, e.g., in the kinetics of antibiotic transport and target binding.

Explaining the absolute values of the dose-sensitivities for different drugs remains a challenge for
future work.

We observed that an artificial nutrient limitation that results in no or extremely slow growth can be
alleviated by adding the antibiotic TMP (Fig. 2B). This indicates that bacteria may not regulate DHFR
expression in a way that maximizes growth under extreme nutrient limitation. Indeed, slow growth
in these conditions coincides with high *folA* expression (Fig. 4D). High *folA* expression is deleterious
(Bhattacharyya *et al*, 2016) as cellular resources are diverted toward excessive folic acid synthesis.
Consistent with this view, TMP facilitated bacterial growth when *folA* was overexpressed to levels
that were deleterious in the absence of TMP (Supplementary Fig. 10). Together, these observations
show that DHFR level is the main driver of TMP susceptibility and suggest that costly overproduction
of DHFR, which can be rescued by adding TMP, occurs under extreme nutrient limitation. Since TMP
increases the fitness of bacteria that evolve under extreme nutrient limitation, the usual selection
pressure for antibiotic resistance is inverted under such conditions: Mutations that usually enhance
TMP action (e.g. increased drug uptake) can be selected. Similar to certain drug combinations (Chait
et al, 2007), this situation provides an opportunity to select against antibiotic-resistant bacteria. One
potential advantage of creating such conditions with a sugar analog instead of a second drug is that
bacteria can hardly evolve resistance to such an analog, as impaired sugar uptake would come at a
massive fitness cost.

Methods

Growth conditions and growth rate measurements

Unless otherwise noted the chemicals used were from Sigma-Aldrich (Steinheim, DE). The growth
medium used was either LB Broth Lennox (L3022), pH set to 7.0 with NaOH before autoclaving, or
M9 minimal medium made from $Na_2HPO_4.7H_2O$ (Fisher Scientific Acros Organics 206515000), KH_2PO_4
(P9791), NaCl (S3014), and NH_4Cl (A9434)) supplemented with 0.1 mM CaCl2 (Fluka 223506), 2mM
MgSO4 (M7506), and 0.001% (v/v) Triton-X 100 (T8787). Triton-X was added to flatten the meniscus
that forms in 96-well pates (Mitosch *et al*, 2017). Carbon sources in the M9 medium were glucose
(G8270), glycerol (VWR 0854), mannose (Carl Roth 4220.2), fructose (F0127), galactose (G0750), all
of which were added at 0.4% (w/v) and prepared as filter sterilized 20% (w/v) stock solutions stored
at room temperature in the dark. Experiments were started from a frozen glycerol stock. Bacteria
were streaked on an LB agar (L2897) plate (containing antibiotics as appropriate) incubated
overnight at 37°C and a single colony was inoculated in 2 ml of the appropriate growth medium

(containing antibiotics if appropriate) and grown for about 20 h to obtain a pre-culture that has

reached stationary phase. We inoculated experimental cultures with a 1000-fold dilution from a

stationary phase culture when growth was determined by optical density measurements at 600 nm

(OD_{600}). For the experiments with luminescence-readout, the pre-culture was grown in 20 ml LB

medium in a 250 ml flask until stationary phase; 100 μl aliquots were transferred to the wells of a

96-well plate, supplemented with glycerol to 15% and frozen at -80°C. To start a luminescence-based

experiment the plate was thawed, and dilutions were performed in 96-well plates with fresh

medium using pin tools (VP407 and VP408, V&P Scientific Inc., CA, USA), which transfer 1.5 μl and

0.2 μl per well, respectively. Subsequent use resulted in a 10^7-fold dilution from a stationary phase

culture. In all cases, the pre-cultures were incubated at 30°C with a shaking speed of 250 rpm

(Innova 44, Eppendorf New Brunswick, DE).

Pre-cultures carrying plasmids and cultures needed for molecular cloning procedures were prepared

with antibiotics at the following concentrations: chloramphenicol 35 μg/ml (C0378), kanamycin

25 μg/ml (K4000), ampicillin 50 μg/ml (A9518), spectinomycin 100 μg/ml (S6501).

Unless otherwise noted antibiotics were dissolved in ethanol (32221). Stock solutions in water were

filter-sterilized. Aliquots of stocks were stored at -20°C in the dark. The antibiotics used were

trimethoprim (92131), nitrofurantoin (N7878), chloramphenicol (C0378), lincomycin (dissolved in

water, 62143), mecillinam (dissolved in water, 33447), tetracycline (268054), and ciprofloxacin

(dissolved in water, 17850). IPTG (VWR 437144N) was added to cultures to control expression from

IPTG-responsive promoters (P_{T5-lac}, $P_{LlacO-1}$) (Kitagawa *et al*, 2005; Lutz & Bujard, 1997). A filter-

sterilized solution of 1M IPTG in water served as stock solution. IPTG was stored at -20°C in the dark

and aliquots were thawed at room temperature before use. For the non-metabolizable glucose

analog α-methyl glucoside (M9376), which competes for glucose uptake and essentially imposes

glucose limitation (Hansen *et al*, 1975), a filter sterilized solution of 50% (w/v) in M9 salts served as

stock solution.

The experiments shown in Fig. 1-3 were performed using a robotic system as described

previously (Chevereau *et al*, 2015) and have a day-to-day variability (coefficient of variation, CV) of

growth rate for unperturbed cultures of less than 5% (Ref. (Chevereau *et al*, 2015) and

Supplementary Fig. 3). The experiments shown in Fig. 4 and 5 were performed using two plate

readers: A Synergy Neo2 and a Synergy H1 (both from Biotek Inc., VT, USA). Both were set to 30°C

with continuous shaking at an orbital displacement of 1 mm and a speed of 807 rpm, and after a

settling period of 10 seconds the optical density at 600 nm and GFP fluorescence were measured

every 10 min. Flat transparent microtiter plates (Nunc Thermo Scientific FT 96-well, 236105) with

lids were used. The experiments presented in Supplementary Fig. 8 were performed using an Infinite

M1000 Pro plate reader (Tecan Inc., CH) equipped with an integrated stacking module. The stack

was housed in a custom-built (IST Austria Miba Machine Shop, Klosterneuburg, AT) acrylic glass box

equipped with a custom-built heating block, a thermostat and strong ventilation to assure a

homogenous temperature over the plates and the stack (Kavčič *et al*, 2020). For these experiments

(Supplementary Fig. 8), the wild-type strain used here (*E. coli* BW25113) was transformed with a

kanamycin resistance-bearing plasmid (pCS-λ) carrying luciferase genes used to determine the

growth rate (Kishony & Leibler, 2003; Chait *et al*, 2007). For the actual growth experiments

kanamycin was omitted; however this was not a problem as the plasmid is retained throughout the

duration of such an experiment (Kavčič *et al*, 2020). Luminescence assays were performed using flat

white microtiter plates (Nunc Thermo Scientific FW 96-well, 260860). These plates were sealed with

a transparent foil (TopSeal-A Plus, PerkinElmer) and about ten plates were used per stack.

Luminescence was measured every 10 to 20 min. Before each measurement, plates were shaken for

10 sec at 582 rpm with a 1 mm amplitude. The culture volume per well was 150 µl. The day-to-day

CV for unperturbed cultures for the growth rate in the luminescence-setup was 3%.

The growth rate was determined by a linear fit of the log-transformed and background-subtracted

OD_{600} from the exponential growth phase of the cultures using custom Matlab (R2016b, MathWorks

Inc.) scripts. To capture the exponential growth phase for cultures in LB we used background-

subtracted OD_{600} windows of 0.02 to 0.2 and for minimal medium 0.03 to 0.12; these windows cover

one order of magnitude and at least two doublings and take the lower growth yield in minimal

medium into account. The lowest accepted growth rate for LB was 0.1 h^{-1} and for minimal medium

0.03 h^{-1}, both corresponding to about 10% of the respective unperturbed maximal growth rate. For

the Hill function fits in Fig. 3C,D, growth rates below 0.2 h^{-1} were ignored because too many data

points fell in this range at higher IPTG concentrations (Fig. 2E) – these data points would thus

dominate the fit, which is undesirable since they contain less information about the shape of the

dose-response curve (i.e. the dose-sensitivity). The duration of experiments for LB cultures was

about 22 h, for minimal medium about 46 h. For all experiments performed in LB medium, data after

~1,000 min were discarded to avoid the inclusion of faster growing mutants which occurred

sporadically in the presence of antibiotics; this was not necessary for experiments in minimal

medium. To capture the growth rate strictly during exponential phase from the luminescence-based

experiments, the rate of luminescence increase was determined by a linear fit of the log-

transformed data between 10^2 cps and 10^5 cps.

Expression level measurements using plate readers

Two plate readers Synergy Neo2 and a Synergy H1 (see section *Growth conditions* for further details)

were used for GFP fluorescence measurements. The filter set used in the Neo2 provided excitation

at 485nm (BW20) and emission at 516nm (BW20) (Biotek fluorescent filter #105). The settings for

the monochromator-based H1 model were 485 nm for excitation and 528 nm for emission. Both

readers produced consistent values and results. The measured values for experiments where both

plate readers were used in parallel were adjusted accordingly (i.e. simply normalized by a constant

obtained from measuring the same sample on both readers). The expression level was determined

essentially as described (Zaslaver *et al*, 2006; Mitosch *et al*, 2017). Briefly, for each GFP-expressing

strain, a similar strain without GFP-expression was grown in parallel in the same conditions (see

section *Strain construction* for further details). For both strains the exponential growth phase was

determined and the background subtracted GFP-signal from the GFP-less strain was subtracted from

the GFP-carrying strain for cultures with similar growth rates and at the same OD_{600}. As the exact

same OD_{600} values were mostly not met, linear interpolation (Matlab function *interp1*) was used to

generate an interpolated GFP-value between the two GFP values of the two nearest OD_{600} values.

The expression level is obtained from the slope of a linear fit (Matlab function *fit*) to the GFP over

OD_{600} data during exponential growth. In the experiments using the strains with the reporter

construct with the native promoter (BWAA01, Fig. 4), fast-folding GFP (Zaslaver *et al*, 2006) was used

whereas in the experiments with the synthetic IPTG-inducible promoter construct (BWAA29R1, Fig.

5) the GFP from the ASKA-library (Kitagawa *et al*, 2005) was used.

Expression level measurements using flow cytometry

For the expression level determination of the strains with IPTG-induced *folA-gfp* expression (Fig. 5)

we used a combination of plate readers (Biotek Synergy H1) for optical density measurements for

growth rate determination (see section growth rate measurements for details) and flow cytometry

(Beckman Coulter CytoFLEX B2-RQ-V2 with 96-well plate module) for fluorescence measurements.

Flow cytometry was used because of its higher signal-to-noise ratio compared to fluorescence

measurements on plate readers. Strains were grown in the plate readers and growth was monitored

by measuring optical density every 10 min. When strains were in mid-exponential growth phase (OD

~ 0.1), they were diluted 1,000-fold in ISOTON II (Beckman Coulter) and measured immediately on

the flow cytometer. Gating in SSC-A and GFP FITC-A channels in the flow cytometry analysis software

(Beckman Coulter Cytexpert 2.3.0.84) allows finding of (fluorescent) cells and determination of the

mean and relative coefficient of variation of fluorescence intensity. Strains used were BWAA11,

BWAA 12, BWAA 19R1, and BWAA 20R1 (see section *Strain construction* for details) and TMP and

IPTG gradients starting at 0.9 μg/ml and 2.5 mM were applied, respectively. Growth rates at

constant or decreasing FolA expression level were calculated by linear interpolation of the growth

rates measured at different IPTG and TMP concentrations as illustrated in Fig. 5A.

Strains and strain construction

We used *E. coli* BW25113 and several derivatives thereof. BW25113 is the parent strain of the KEIO

collection, a widely used whole-genome deletion mutant collection (Baba *et al*, 2006). For the

overexpression experiments, BW25113 was transformed with the necessary plasmids (Table 1)

which stem from the ASKA-library, a plasmid-based whole-genome overexpression

collection (Kitagawa *et al*, 2005). To reduce growth rate by gratuitous protein expression we used a

truncated elongation factor Tu (EF-Tu, *tufB*) as previously done for a similar purpose (Dong *et al*,

1995). Briefly, starting with the ASKA-library plasmid carrying *tufB*, the SmaI restriction fragment of

243 bp in length was cut out and the blunt-ended DNA fragment was closed by ligation to form a

plasmid again, named pAAtufB here. This deletion results in a shortened, non-functional gene

(*ΔtufB*), which can be used to provide gratuitous protein expression, resulting in a burden that slows

down growth (Dong *et al*, 1995; Scott *et al*, 2010). The plasmids from the ASKA-library (Kitagawa *et*

al, 2005) use the P$_{T5-lac}$ promoter, which allows for a graded control of expression by the addition of

the inducer IPTG (which works sufficiently well in a *lac*-operon compromised strain like *E. coli*

BW25113). As control, we used pAA30 which is the empty ASKA plasmid modified to not contain a

gene to prevent any expression; we created this plasmid since the original empty ASKA plasmid does

in fact encode a short coding sequence in frame with the promoter. Briefly, through a PCR with

overlapping primers (for general strategy see (Heckman & Pease, 2007; Hansson *et al*, 2008)) (Fw:

CATTAAAGAGGAGAAATTAACTGGGTCGACCTGCAG,   Rv:

CTGCAGGTCGACCCAGTTAATTTCTCCTCTTTAATG) a short stretch of pCA24N(-) encompassing start

codon over the His-Tag and until the stop codon, was eliminated. The elimination was confirmed by

sequencing the resulting plasmid with primers flanking the gene insertion site (Fw:

CAACAGTTGCCTAAGAAACCAT, Rv: TGAGGTCATTACTGGATCTATCAAC). For the strong *folA*

overexpression the ASKA plasmid pCA24N(-)folA was used.

We generated reporter strains and a strain with inducible *folA* regulation. To construct the first *gfp*-

reporter and corresponding *gfp*-less control pair integrated into the chromosome (BWAA01 and

BWAA02), the promoter-reporter construct for P$_{folA}$ and the corresponding region from the empty

plasmid pUA66 from the reporter library (Zaslaver *et al*, 2006) were integrated into a neutral site

(*phoA*) in the genome, respectively. To this end, P1 transduction was used to move the construct

from an MG1655 strain carrying the reporter constructs (Bollenbach *et al*, 2009) into the BW25113

background. The insertion was confirmed by sequencing PCR products generated using primers

binding outside the *phoA* locus (Fw: GGCGCTGTACGAGGTAAAG, Rv: GGGTTAAAGTTCTCTCGGCA).

The other reporters were based on *folA-gfp* fusion constructs from the ASKA-library (Kitagawa *et al*,

2005). Again, pairs of strains were made where each pair consists of a strain with and a strain

without the *gfp* fused to *folA*. We generated several pairs to induce and thereby control expression

level by an IPTG-responsive promoter ($P_{LlacO-1}$ (Lutz & Bujard, 1997)) and one pair with the native

regulation through P_{folA}. This was done to approximately match the induced expression with the level

of the native regulation. We generated strains with the inducible $P_{LlacO-1}$ promoter with five different

ribosome binding sites of different strength with the sequences (RBS1, 3, 4, 5, and 6) originating

from (Deris *et al*, 2013). Those strains were compared to strains with the native regulation (BWAA11

and BWAA12) which were made in parallel according to a similar strategy as described below. Based

on similar expression levels for BWAA19R1 and BWAA11, BWAA19R1 with RBS1, was chosen for the

experiment shown in Fig. 5. To create the *gfp* fusion strains, *folA-chlR* and *folA-gfp-chlR* fragments

were PCR-amplified from the *folA*-carrying ASKA-library plasmids (using as template the respective

plasmids from the library, with and without *gfp* (Kitagawa *et al*, 2005)) as a first step and were used

for recombineering (Datsenko & Wanner, 2000) into the plasmid pKD13-gfpmut3 (a derivative of

pKD13 (Datsenko & Wanner, 2000); gift from Bor Kavčič). Primers used were

CAGCAGGACGCACTGACCGAATTCATTAAAGAGGAGAAAGGTACCGCATGATCAGTCTGATTGCGGCGTTAG

and GACTGAGCCTTTCGTTTTATTTGATGCCTCTAGACTCAGCTAATTAAGCGTAGCACCAGGCGTTTAAGG.

This resulted in pAA39 and pAA40 where an FRT-flanked kanamycin resistance cassette, the

promoter $P_{LlacO-1}$ driving *folA* and the *folA-gfp* fusion, respectively, and a chloramphenicol resistance

cassette are present (in this order). These plasmids first served as source for the promoter-*folA* and

folA-gfp fusion with a chloramphenicol resistance cassette to be inserted into the genome at the

folA locus to generate the strains with the native regulation (BWAA11 and BWAA12). PCR-fragments

for recombineering were obtained with the previously used forward primer

CAGCAGGACGCACTGACCGAATTCATTAAAGAGGAGAAAGGTACCGCATGATCAGTCTGATTGCGGCGTTAG

and AAGACGCGACCGGCGTCGCATCCGGCGCTAGCCGTAAATTCTATACAAAACTAGACTCAGCTAATTAAGC

serving as reverse primer to get the *folA* gene with and without *gfp,* respectively, and the

chloramphenicol resistance cassette (but not the synthetic promoter) were inserted into the

genome of BW25113 replacing the *folA* gene (but not the promoter on the genome). Next, by

recombineering with PCR-fragments containing the kanamycin resistance cassette only obtained

with   the   primers

GTGATACGCCTATTTTTATAGGTTAATGTCATGATAATAATGGTTTCTTAGAAGTTCCTATTCTCTAGAAAGTAT

AGG   and

AAGACGCGACCGGCGTCGCATCCGGCGCTAGCCGTAAATTCTATACAAAAGTGTAGGCTGGAGCTGCTTC

from pKD13-gfpmut3 the chloramphenicol resistance cassette was replaced with the FRT-flanked

kanamycin resistance cassette. Next, to obtain a marker-less strain the kanamycin resistance

cassette was removed using the plasmid pCP20, as described (Cherepanov & Wackernagel, 1995).

For the strains with the IPTG-inducible regulation (BWAA19R1 and BWAA20R1) a similar strategy

applied.   Primers

AACATACCGGCAACATGGCGGATGAACCGGAAACGAAACCCTCATCCTAAGTGTAGGCTGGAGCTGC   and

TCCATGCCGATAACGCGATCTACCGCTAACGCCGCAATCAGACTGATCATGCGGTACCTTT<u>CT</u>CCTCTTT

(putative RBS1 sequence from (Baba *et al*, 2006) underlined) were used to amplify the FRT-flanked

kanamycin resistance cassette, the promoter P*LlacO-1* driving *folA* and the *folA-gfp* respectively of

pAA39 and pAA40 (but not the chloramphenicol resistance cassette). Next, to obtain a marker-less

strain the kanamycin resistance cassette was removed using pCP20. All four marker-less strains were

further modified by P1 transduction from a MG1655 strain carrying the *lacI* gene under the

promoter P*lacO1* and a FRT-flanked kanamycin resistance cassette at the neutral insertion site *intS*

(based on the strain from (Garcia *et al*, 2011) (HG105) and a gift from Bor Kavčič). The insertion was

confirmed by sequencing PCR products generated using primers binding outside the *intS* locus (Fw:

GTACTTACCCCGCACTCCAT, Rv: TGTTCAGCACACCAATAGAGG). Next, that kanamycin resistance

cassette was removed using pCP20. The resulting markerless strains were further modified by P1

transduction with the *lacI* knock-out strain from the KEIO collection (Baba *et al*, 2006) replacing the

lacI gene with the kanamycin resistance cassette. The deletion was confirmed by sequencing PCR

products generated using primers binding outside the *lacI* locus (Fw: CGGCTCATGGATGGTGTT, Rv:

CGAAGCGGCATGCATTTAC). We reasoned, that here a P*LlacO-1*-driven *lacI* allows a better control of

the P*LlacO-1*-driven *folA* based on observations in (Kavčič *et al*, 2020; Klumpp *et al*, 2009) dealing with

growth rate independent negative autoregulation. Moreover, with the combination of RBS1 and the

Lac-repressor driven by P*lacO1* we achieved expression close to wild-type levels. For the generation of

the strains with modified RBS3-6 from BWAA19R1 and BWAA20R1, the kanamycin resistance

cassette was removed using pCP20. We again used long reverse primers RBS3-Rv:

TCCATGCCGATAACGCGATCTACCGCTAACGCCGCAATCAGACTGATCATGCGGTACCTTT<u>AGGACT</u>CCTCTTT

aatgaattcggtcag,   RBS4-Rv:

TCCATGCCGATAACGCGATCTACCGCTAACGCCGCAATCAGACTGATCATGCGGTACCTTT<u>GT</u>CCTCTTTaatga

attcggtcag,   RBS5-Rv:

TCCATGCCGATAACGCGATCTACCGCTAACGCCGCAATCAGACTGATCATGCGGTACCTTT<u>AGGAGT</u>CCTCTTT

aatgaattcggtcag,   RBS6-Rv:

TCCATGCCGATAACGCGATCTACCGCTAACGCCGCAATCAGACTGATCATGCGGTACCTTT<u>AGGAGG</u>CCTCTTT

aatgaattcggtcag, each modified such that the RBS sequence was changed according to (Deris *et al*,

2013). To aid primer binding (compare (Liu & Naismith, 2008)) to the template (which carries RBS1

sequence) and change of the RBS-sequence (underlined) additional 15 nt were added to the reverse

primers   (lower   case   letters)   and   used   in   combination   with   Fw:

AACATACCGGCAACATGGCGGATGAACCGGAAACGAAACCCTCATCCTAAGTGTAGGCTGGAGCTGC   on

pAA39 and pAA40. The respective PCR products were sequenced and used for recombineering of the

markerless intermediate versions of BWAA19R1 and BWAA20R1. After integration, all promoter

regions were confirmed by sequencing a PCR product obtained with primers targeting the *folA* locus

with the primers (Fw: CCAGCGCGATGTTAAGTGA, Rv: GATTGATTCCCAGGTATGGCG).

For the recombineering procedure (Datsenko & Wanner, 2000) the temperature-inducible system

from pSIM19 (Sharan *et al*, 2009) was used. Chloramphenicol at 10 µg/ml and kanamycin at

25 µg/ml were used. During the whole strain construction procedure wherever *folA* was driven by

PL$_{lacO1}$, 1 mM IPTG was added as this inducer controls expression from PL$_{lacO1}$ and *folA* is an essential

gene.

Mathematical model

Solutions of the mathematical model in Fig. 1B were numerically calculated using Python (function

fsolve from *scipy* version 1.2.1). As dose-sensitivity without growth-mediated feedback, we used $n =$

2 since this is a typical value for many antibiotics (Fig. 1A).

Strains and plasmids used in this study

Name	Information	Source
E. coli BW25113	KEIO collection parent strain (Baba *et al*, 2006)	Lab strain collection
BWAA01	*ΔphoA:kan:P$_{folA}$:gfp* based on P$_{folA}$-gfp plasmid from (Zaslaver *et al*, 2006)	This study, based on (Bollenbach *et al*, 2009)
BWAA02	*ΔphoA:kan:P$_{folA}$* based on pUA66 plasmid from (Zaslaver	This study, based on (Bollenbach *et al*, 2009)

	et al, 2006)	
BWAA11	Native regulation *folA:gfp:kan*	This study
BWAA12	Native regulation *folA:kan*	This study
BWAA19R1	IPTG-inducible regulation ΔP_{folA}*:kan*$:P_{LlacO-1}$*:folA:gfp, intS*$:P_{LlacO-1}$*:lacI, ΔlacI:kan*	This study
BWAA19R3	As BWAA19R1 but with RBS3 from (Deris *et al*, 2013)	This study
BWAA19R4	As BWAA19R1 but with RBS4 from (Deris *et al*, 2013)	This study
BWAA19R5	As BWAA19R1 but with RBS5 from (Deris *et al*, 2013)	This study
BWAA19R6	As BWAA19R1 but with RBS6 from (Deris *et al*, 2013)	This study
BWAA20R1	IPTG-inducible regulation ΔP_{folA}*:kan*$:P_{LlacO-1}$*:folA, intS*$:P_{LlacO-1}$*:lacI, ΔlacI:kan*	This study
BWAA20R3	As BWAA20R1 but with RBS3 from (Deris *et al*, 2013)	This study
BWAA20R4	As BWAA20R1 but with RBS4 from (Deris *et al*, 2013)	This study
BWAA20R5	As BWAA20R1 but with RBS5 from (Deris *et al*, 2013)	This study
BWAA20R6	As BWAA20R1 but with RBS6 from (Deris *et al*, 2013)	This study
BW25141	Cloning strain CGSC#: 7633 with *pir*[+] from (Datsenko & Wanner, 2000)	Lab strain collection
pSIM19	Recombineering plasmid (Datta *et al*, 2006)	Lab strain collection
pCP20	Source of Flp for excision (Cherepanov & Wackernagel, 1995)	Lab strain collection
pKD13-mutgfp	FRT-flanked *kan*, P_{LlacO1}	Bor Kavcic

pCA24N(-)folA	Source of *folA:chl* (Kitagawa *et al*, 2005)	Lab strain collection
pCA24N(-)	Empty ASKA plasmid (Kitagawa *et al*, 2005)	Lab strain collection
pAAtufB	Modified pCA24N(-)tufB	This study
pAA30	Modified pCA24N(-)	This study
pAA39	*folA:gfp:chl* in pKD13-mutgfp	This study
pAA40	*folA:chl* in pKD13-mutgfp	This study
pCS-λ	Luciferase genes for luminescence measurements, *kan*	Ref. (Kishony & Leibler, 2003)

Acknowledgements

This work was in part supported by Human Frontier Science Program Grant RGP0042/2013, Marie Curie Career Integration Grant 303507, Austrian Science Fund (FWF) Grant P 27201-B22, and German Research Foundation (DFG) Collaborative Research Center (SFB) 1310 to Tobias Bollenbach. S. Andreas Angermayr was supported by the European Union's Horizon 2020 research and innovation program under the Marie Skłodowska-Curie grant agreement No 707352. We would like to thank the Bollenbach group for regular fruitful discussions. We are particularly thankful for technical assistance of Booshini Fernando and for discussions of the theoretical aspects with Gerrit Ansmann. We are indebted to Bor Kavčič for invaluable advice, help with setting up the luciferase-based growth monitoring system, and for sharing plasmids. We are grateful to Rosalind Allen, Bor Kavčič , Karin Mitosch and Dor Russ for feedback on the manuscript.

We declare that there are no conflicts of interest.

SAA, GC, and TB conceived the study. SAA and TB designed the experiments. SAA performed the experiments and analyzed the data. GC analyzed data presented in Fig. 1C and Supplementary Fig. 1. SAA and TB wrote the manuscript with input from GC.

References

Baba T, Ara T, Hasegawa M, Takai Y, Okumura Y, Baba M, Datsenko KA, Tomita M, Wanner BL, Mori H, Anderson R, Roth J, Baudin A, Ozier-Kalogeropoulos O, Denouel A, Lacroute F, Cullin C, Blattner F, Plunkett G, Bloch C, et al (2006) Construction of Escherichia coli K-12 in-frame, single-gene knockout mutants: the Keio collection. *Mol. Syst. Biol.* **2:** 2006.0008

Balaban NQ, Merrin J, Chait R, Kowalik L & Leibler S (2004) Bacterial persistence as a phenotypic
switch. *Science* **305:** 1622–5 Available at:
http://www.sciencemag.org/cgi/doi/10.1126/science.1099390 [Accessed January 20, 2014]

Baym M, Lieberman TD, Kelsic ED, Chait R, Gross R, Yelin I & Kishony R (2016) Spatiotemporal
microbial evolution on antibiotic landscapes. *Science* **353:** 1147–51 Available at:
http://www.sciencemag.org/cgi/doi/10.1126/science.aag0822

Bershtein S, Choi J-M, Bhattacharyya S, Budnik B & Shakhnovich E (2015) Systems-level response to
point mutations in a core metabolic enzyme modulates genotype-phenotype relationship. *Cell
Rep.* **11:** 645–56 Available at: http://dx.doi.org/10.1016/j.celrep.2015.03.051

Bershtein S, Mu W, Serohijos AWR, Zhou J & Shakhnovich EI (2013) Protein Quality Control Acts on
Folding Intermediates to Shape the Effects of Mutations on Organismal Fitness. *Mol. Cell* **49:**
133–144 Available at: http://dx.doi.org/10.1016/j.molcel.2012.11.004

Bhattacharyya S, Bershtein S, Yan J, Argun T, Gilson AI, Trauger SA & Shakhnovich EI (2016) Transient
protein-protein interactions perturb E. coli metabolome and cause gene dosage toxicity. *Elife* **5:**
1–22 Available at: http://www.ncbi.nlm.nih.gov/pubmed/27938662

Bintu L, Buchler NE, Garcia HG, Gerland U, Hwa T, Kondev J, Phillips R, Kuhlman T & Phillips R (2005)
Transcriptional regulation by the numbers: models. *Curr. Opin. Genet. Dev.* **15:** 116–24
Available at: http://linkinghub.elsevier.com/retrieve/pii/S0959437X05000298

Bollenbach T, Quan S, Chait R & Kishony R (2009) Nonoptimal microbial response to antibiotics
underlies suppressive drug interactions. *Cell* **139:** 707–18 Available at:
http://dx.doi.org/10.1016/j.cell.2009.10.025 [Accessed January 24, 2014]

Bremer H & Dennis PP (2008) Modulation of Chemical Composition and Other Parameters of the Cell
at Different Exponential Growth Rates. *EcoSal Plus* **3:** 765–77 Available at:
http://www.ncbi.nlm.nih.gov/pubmed/26443740

Chait R, Craney A & Kishony R (2007) Antibiotic interactions that select against resistance. *Nature*
446: 668–71 Available at: http://www.ncbi.nlm.nih.gov/pubmed/17410176 [Accessed January
24, 2014]

Cherepanov PP & Wackernagel W (1995) Gene disruption in Escherichia coli: TcR and KmR cassettes
with the option of Flp-catalyzed excision of the antibiotic-resistance determinant. *Gene* **158:** 9–
14 Available at: http://www.ncbi.nlm.nih.gov/pubmed/7789817

Chevereau G & Bollenbach T (2015) Systematic discovery of drug interaction mechanisms. *Mol. Syst.
Biol.* **11:** 807 Available at: http://www.ncbi.nlm.nih.gov/pubmed/25924924

Chevereau G, Dravecká M, Batur T, Guvenek A, Ayhan DH, Toprak E & Bollenbach T (2015)
Quantifying the Determinants of Evolutionary Dynamics Leading to Drug Resistance. *PLoS Biol.*
13: e1002299 Available at: http://dx.plos.org/10.1371/journal.pbio.1002299

Chou T-C & Talalay P (1983) Analysis of combined drug effects: a new look at a very old problem.
Trends Pharmacol. Sci. **4:** 450–454 Available at:
https://linkinghub.elsevier.com/retrieve/pii/016561478390490X

Datsenko KA & Wanner BL (2000) One-step inactivation of chromosomal genes in Escherichia coli K-
12 using PCR products. *Proc. Natl. Acad. Sci. U. S. A.* **97:** 6640–5

Datta S, Costantino N & Court DL (2006) A set of recombineering plasmids for gram-negative
bacteria. *Gene* **379:** 109–15 Available at: http://www.ncbi.nlm.nih.gov/pubmed/16750601

Deris JB, Kim M, Zhang Z, Okano H, Hermsen R, Groisman A & Hwa T (2013) The innate growth

bistability and fitness landscapes of antibiotic-resistant bacteria. *Science* **342:** 1237435
Available at: http://www.sciencemag.org/cgi/doi/10.1126/science.1237435 [Accessed January
20, 2014]

Dong H, Nilsson L & Kurland C (1995) Gratuitous overexpression of genes in Escherichia coli leads to
growth inhibition and ribosome destruction. *J. Bacteriol.* **177:** 1497–1504

Elf J, Nilsson K, Tenson T & Ehrenberg M (2006) Bistable bacterial growth rate in response to
antibiotics with low membrane permeability. *Phys. Rev. Lett.* **97:** 1–4 Available at:
http://link.aps.org/doi/10.1103/PhysRevLett.97.258104 [Accessed January 24, 2014]

Elowitz MB & Leibler S (2000) A synthetic oscillatory network of transcriptional regulators. *Nature*
403: 335–338 Available at: http://www.ncbi.nlm.nih.gov/pubmed/10659856

Flensburg J & Sköld O (1987) Massive overproduction of dihydrofolate reductase in bacteria as a
response to the use of trimethoprim. *Eur. J. Biochem.* **162:** 473–6 Available at:
http://www.ncbi.nlm.nih.gov/pubmed/3549289

Garcia HG, Lee HJ, Boedicker JQ & Phillips R (2011) Comparison and Calibration of Different
Reporters for Quantitative Analysis of Gene Expression. *Biophys. J.* **101:** 535–544

Gardner TS, Cantor CR & Collins JJ (2000) Construction of a genetic toggle switch in Escherichia coli.
Nature   **403:**   339–342   Available   at:
http://www.nature.com/nature/journal/v403/n6767/full/403339a0.html

Greulich P, Scott M, Evans MR & Allen RJ (2015) Growth-dependent bacterial susceptibility to
ribosome-targeting   antibiotics.   *Mol.   Syst.   Biol.*   **11:**   796   Available   at:
http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=4380930&tool=pmcentrez&rende
rtype=abstract [Accessed March 21, 2015]

Hansen MT, Pato ML, Molin S, Fill NP & von Meyenburg K (1975) Simple downshift and resulting lack
of correlation between ppGpp pool size and ribonucleic acid accumulation. *J. Bacteriol.* **122:**
585–91 Available at: http://www.ncbi.nlm.nih.gov/pubmed/1092659 [Accessed September 6,
2018]

Hansson MD, Rzeznicka K, Rosenbäck M, Hansson M & Sirijovski N (2008) PCR-mediated deletion of
plasmid   DNA.   *Anal.   Biochem.*   **375:**   373–5   Available   at:
http://www.ncbi.nlm.nih.gov/pubmed/18157935

Heckman KL & Pease LR (2007) Gene splicing and mutagenesis by PCR-driven overlap extension. *Nat.*
Protoc. **2:** 924–32 Available at: http://www.ncbi.nlm.nih.gov/pubmed/17446874

Hermsen R, Deris JB & Hwa T (2012) On the rapidity of antibiotic resistance evolution facilitated by a
concentration gradient. *Proc. Natl. Acad. Sci.* **109:** 10775–10780

Hol FJH, Hubert B, Dekker C & Keymer JE (2016) Density-dependent adaptive resistance allows
swimming bacteria to colonize an antibiotic gradient. *ISME J.* **10:** 30–38 Available at:
http://www.nature.com/doifinder/10.1038/ismej.2015.107

Kavčič B, Tkačik G & Bollenbach T (2020) Mechanisms of drug interactions between translation-
inhibiting   antibiotics.   *Nat.   Commun.*   **11:**   4013   Available   at:
https://www.nature.com/articles/s41467-020-17734-z [Accessed November 17, 2019]

Keseler IM (2004) EcoCyc: a comprehensive database resource for Escherichia coli. *Nucleic Acids Res.*
33: D334–D337 Available at: http://nar.oxfordjournals.org/lookup/doi/10.1093/nar/gki108
[Accessed January 20, 2014]

Kishony R & Leibler S (2003) Environmental stresses can alleviate the average deleterious effect of

mutations. *J. Biol.* **2:** 14 Available at: http://www.jbiol.com/content/2/2/14 [Accessed February
21, 2014]

Kitagawa M, Ara T, Arifuzzaman M, Ioka-Nakamichi T, Inamoto E, Toyonaga H & Mori H (2005)
Complete set of ORF clones of Escherichia coli ASKA library (A complete set of E. coli K-12 ORF
archive): unique resources for biological research. *DNA Res.* **12:** 291–299

Klumpp S, Zhang Z & Hwa T (2009) Growth rate-dependent global effects on gene expression in
bacteria.   *Cell*   **139:**   1366–75   Available   at:
http://linkinghub.elsevier.com/retrieve/pii/S0092867409015050 [Accessed January 20, 2014]

Lee AJ, Wang S, Meredith HR, Zhuang B, Dai Z & You L (2018) Robust, linear correlations between
growth rates and β-lactam-mediated lysis rates. *Proc. Natl. Acad. Sci. U. S. A.* **115:** 4069–4074
Available at: http://www.pnas.org/lookup/doi/10.1073/pnas.1719504115

Liu H & Naismith JH (2008) An efficient one-step site-directed deletion, insertion, single and
multiple-site plasmid mutagenesis protocol. *BMC Biotechnol.* **8:** 91 Available at:
http://bmcbiotechnol.biomedcentral.com/articles/10.1186/1472-6750-8-91

Lutz R & Bujard H (1997) Independent and tight regulation of transcriptional units in Escherichia coli
via the LacR/O, the TetR/O and AraC/I1-I2 regulatory elements. *Nucleic Acids Res.* **25:** 1203–10

Mitosch K, Rieckh G & Bollenbach T (2017) Noisy Response to Antibiotic Stress Predicts Subsequent
Single-Cell Survival in an Acidic Environment. *Cell Syst.* **4:** 393-403.e5 Available at:
http://www.ncbi.nlm.nih.gov/pubmed/28342718

Nichols RJ, Sen S, Choo YJ, Beltrao P, Zietek M, Chaba R, Lee S, Kazmierczak KM, Lee KJ, Wong A,
Shales M, Lovett S, Winkler ME, Krogan NJ, Typas A & Gross CA (2011) Phenotypic landscape of
a   bacterial   cell.   *Cell*   **144:**   143–56   Available   at:
http://linkinghub.elsevier.com/retrieve/pii/S0092867410013747 [Accessed January 23, 2014]

Nyerges Á, Csörgő B, Draskovits G, Kintses B, Szili P, Ferenc G, Révész T, Ari E, Nagy I, Bálint B,
Vásárhelyi BM, Bihari P, Számel M, Balogh D, Papp H, Kalapis D, Papp B & Pál C (2018) Directed
evolution of multiple genomic loci allows the prediction of antibiotic resistance. *Proc. Natl.*
Acad.   Sci.   U.   S.   A.   **115:**   E5726–E5735   Available   at:
http://www.ncbi.nlm.nih.gov/pubmed/29871954%0Ahttp://www.pnas.org/lookup/doi/10.107
3/pnas.1801646115

Palmer AC & Kishony R (2014) Opposing effects of target overexpression reveal drug mechanisms.
Nat. Commun. **5:** 4296 Available at: http://www.ncbi.nlm.nih.gov/pubmed/24980690
[Accessed July 2, 2014]

Regoes RR, Wiuff C, Zappala RM, Garner KN, Baquero F & Levin BR (2004) Pharmacodynamic
functions: a multiparameter approach to the design of antibiotic treatment regimens.
Antimicrob.   Agents   Chemother.   **48:**   3670–6   Available   at:
http://www.ncbi.nlm.nih.gov/pubmed/15388418

Rodrigues J V, Bershtein S, Li A, Lozovsky ER, Hartl DL & Shakhnovich EI (2016) Biophysical principles
predict fitness landscapes of drug resistance. *Proc. Natl. Acad. Sci. U. S. A.* **113:** E1470-8
Available at: http://www.pnas.org/lookup/doi/10.1073/pnas.1603613113

Rood JI, Laird AJ & Williams JW (1980) Cloning of the Escherichia coli K-12 dihydrofolate reductase
gene   following   mu-mediated   transposition.   *Gene*   **8:**   255–65   Available   at:
http://www.ncbi.nlm.nih.gov/pubmed/6444603

Russ D & Kishony R (2018) Additivity of inhibitory effects in multidrug combinations. *Nat. Microbiol.*
3: 1339–1345 Available at: http://www.ncbi.nlm.nih.gov/pubmed/30323252

Scott M, Gunderson CW, Mateescu EM, Zhang Z & Hwa T (2010) Interdependence of cell growth and
gene expression: origins and consequences. *Science* **330:** 1099–102 Available at:
http://science.sciencemag.org/content/330/6007/1099.long

Sharan SK, Thomason LC, Kuznetsov SG & Court DL (2009) Recombineering: a homologous
recombination-based method of genetic engineering. *Nat. Protoc.* **4:** 206–223

Soo VWC, Hanson-Manful P & Patrick WM (2011) Artificial gene amplification reveals an abundance
of promiscuous resistance determinants in Escherichia coli. *Proc. Natl. Acad. Sci. U. S. A.* **108:**
1484–9

Toprak E, Veres A, Michel J-B, Chait R, Hartl DL & Kishony R (2012) Evolutionary paths to antibiotic
resistance under dynamically sustained drug selection. *Nat. Genet.* **44:** 101–105 Available at:
http://www.nature.com/doifinder/10.1038/ng.1034 [Accessed January 20, 2014]

Tuomanen E, Cozens R, Tosch W, Zak O & Tomasz A (1986) The rate of killing of Escherichia coli by
beta-lactam antibiotics is strictly proportional to the rate of bacterial growth. *J. Gen. Microbiol.*
132: 1297–304 Available at: http://www.ncbi.nlm.nih.gov/pubmed/3534137

Yang J, Ogawa Y, Camakaris H, Shimada T, Ishihama A & Pittard AJ (2007) folA, a New Member of the
TyrR Regulon in Escherichia coli K-12. *J. Bacteriol.* **189:** 6080–6084 Available at:
https://jb.asm.org/content/189/16/6080

Zaslaver A, Bren A, Ronen M, Itzkovitz S, Kikoin I, Shavit S, Liebermeister W, Surette MG & Alon U
(2006) A comprehensive library of fluorescent transcriptional reporters for Escherichia coli.
Nat. Methods **3:** 623–628 Available at: http://www.nature.com/doifinder/10.1038/nmeth895

Supplementary figures

Supplementary Fig. 1|Correlation of antibiotic response and drug-free growth rate for genome-wide gene deletion
strains. *Density scatterplots showing growth response to different antibiotics versus normalized growth rate in the absence*
of drug for genome-wide gene deletion strains (Baba *et al*, 2006) *as in Fig. 1C. Response is defined as growth rate in the*
presence of the respective drug normalized to the drug-free growth rate of the respective deletion strain. Each drug was
used at a fixed drug concentration that inhibits wild type growth by about 30% (Chevereau *et al*, 2015)*. Spearman*
correlation coefficient ρ_S is shown.

Supplementary Fig. 2|Correlation of IC$_{50}$ and drug-free growth rate in gene deletion strains. *Scatterplots of the IC$_{50}$ of gene deletion mutants versus the growth rate of these mutants in the absence of drug; each panel shows a different antibiotic as labeled. ρ_s is the Spearman correlation; p-values of this correlation are shown. The only significant (negative) correlation occurs for TMP, consistent with growth-mediated negative feedback for this drug. IC$_{50}$s were determined from dose-response curve measurements of 78 arbitrary gene deletions strains (Chevereau et al, 2015).*

Supplementary Fig. 3| Day-to-day reproducibility of growth rate measurements. *Scatterplot showing comparison of growth rate data from αMG-TMP two-dimensional concentration gradient experiment (Fig. 2B) performed on two different days.*

Supplementary Fig. 4| IPTG alone at the concentrations used here has no effect on growth rate. *Black data points show*
growth rate versus IPTG concentration for a control strain with an empty expression vector; data from Fig. 2C is shown in
gray for comparison. Error bars show standard deviation from eight replicates.

Supplementary Fig. 5| Effect of growth rate reduction by glucose limitation on susceptibility to diverse antibiotics. *As Fig.*
2B, for nitrofurantoin (NIT), tetracycline (TET), lincomycin (LIN), ciprofloxacin (CPR), mecillinam (MEC), and chloramphenicol
(CHL). Lowering growth rate by glucose limitation via αMG does not lower susceptibility to these antibiotics as for TMP (cf.
Fig. 2C).

Supplementary Fig. 6| Effect of growth rate reduction by gratuitous protein overexpression on susceptibility to antibiotics. As Fig. 2E, for trimethoprim (TMP), nitrofurantoin (NIT), tetracycline (TET), lincomycin (LIN), ciprofloxacin (CPR), and mecillinam (MEC). Lowering growth rate by gratuitous protein overexpression lowers susceptibility to TMP and, to a lesser extent, to MEC (bottom right), but not for the other antibiotics (cf. Fig. 2F).

847 ***Supplementary Fig. 7| Effect of growth rate reduction by changing carbon source on susceptibility to antibiotics.*** *As Fig.*
848 *2H, for nitrofurantoin (NIT), tetracycline (TET), ciprofloxacin (CPR), and chloramphenicol (CHL). Lowering growth rate via*
849 *poorer carbon sources does not lower susceptibility to other antibiotics than TMP (cf. Fig. 2I). Data shown is the mean of*
850 *three replicates. We also performed this assay for mecillinam (MEC), but excluded it from further analysis because – for*
851 *unknown reasons – it consistently showed extremely noisy dose-response curves in this assay.*

Supplementary Fig. 8|Lowering growth rate by changing temperature does not affect the shape of antibiotic dose-response curves. *Left column: Growth rate versus drug concentration for eight different antibiotics at eight different temperatures as shown. Right column: Growth rate normalized to drug-free growth rate.*

Supplementary Fig. 9|Expression level of folA as a function of trimethoprim and αMG/glucose ratio. *Data from Fig. 2 in two-dimensional checkerboard plot.*

Supplementary Fig. 10|Growth rate decrease due to DHFR overexpression is rescued by trimethoprim. *Normalized*
growth rate (gray scale) in a two-dimensional concentration gradient of IPTG and TMP. IPTG controls overexpression of folA
(Methods). Dotted lines are contour lines at 50% growth inhibition. DHFR overexpression lowers growth rate but adding
TMP at high IPTG concentrations partially rescues this phenotype: Growth rate increases with increasing TMP
concentration. The increase in TMP IC_{50} resulting from DHFR overexpression confirms previous reports (Palmer & Kishony,
2014)*.*

Supplementary Fig. 11| Bacterial growth curves from optical density measurements. *Representative data for different ways of lowering the growth rate in different growth media.* ***(A)*** *TMP in LB medium. Growth curves underlying the data shown in the leftmost column of Fig. 2E; TMP concentration increases from cyan to magenta.* ***(B)*** *Gratuitous protein overexpression in LB medium; data correspond to Fig. 2D; IPTG concentration increases from cyan to magenta.* ***(C)*** *TMP in glucose (minimal M9) medium; data correspond to the leftmost column of Fig. 2B.* ***(D)*** *αMG in glucose (minimal M9) medium; data correspond to Fig. 2A; αMG/glucose ratio increases from cyan to magenta.*