1

Plasma mass spectrometry fingerprints induced by bacterial endotoxins in murine models as markers of pathophysiological evolution in sepsis

Author names and affiliations:

Martín Ledesma[1, 7], María Florencia Todero[2], Lautaro Maceira[2], Monica Prieto[3], Carlos Vay[1], Marcelo Galas[4], Beatriz López[5], Noemí Yokobori[6, 7], Bárbara Rearte[2, 7]

Laboratorio de Bacteriología, Departamento de Bioquímica Clínica, Hospital de Clínicas José de San Martín; Facultad de Farmacia y Bioquímica, UBA, Buenos Aires, Argentina

Instituto de Medicina Experimental (IMEX) – CONICET -Academia Nacional de Medicina, Buenos Aires, Argentina.

Servicio Bacteriología Especial. Instituto Nacional de Enfermedades Infecciosas (INEI), ANLIS "Dr. C. G. Malbrán", Buenos Aires, Argentina.

Special Program of AMR, Communicable Diseases and Environmental Determinants of Health Department, Panamerican Health Organization, Washington, D.C., USA .

Departamento de Bacteriología. Instituto Nacional de Enfermedades Infecciosas (INEI), ANLIS "Dr. C. G. Malbrán", Buenos Aires, Argentina.

Instituto Nacional de Enfermedades Infecciosas (INEI), ANLIS "Dr. C. G. Malbrán" Buenos Aires, Argentina.

7. Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Argentina.

ABSTRACT

Sepsis constitutes a major cause of death in intensive care units. Patients present a dysregulated response to infection, which progresses through different pro/anti-inflammatory phases. The lack of tools to monitor their response constrains the therapeutic approaches. Here, we evaluated a test based on plasma protein fingerprints acquired by MALDI-TOF-mass spectrometry and supervised/unsupervised algorithms to discriminate the different immunological stages of sepsis in lipopolysaccharide-induced murine models with encouraging results. Moreover, our predictive models through machine learning algorithms were able to discriminate the different groups with a sensitivity of up to 95.7% and a specificity of 90.9% depending on the selected peaks number. Potential individual biomarkers associated with each phase were also analysed. Our data reveal the potential of plasma peptidome analysis by MALDI-TOF-mass spectrometry as a highly relevant strategy for sepsis patient stratification that could contribute to therapeutic decisions, depending on the immunological phase that the patient is undergoing.

INTRODUCTION

Sepsis continues to be one the major causes of death in intensive care units (ICU)[1,2]. Estimations indicate a worldwide incidence of 31.5 million cases annually with approximately 7-9 million deaths[3] The precise impact of sepsis in Latin America is elusive due to the limited number of prevalence studies. However, in view of the high incidence rate of infectious diseases along with the inequities in access to health care prevailing there, sepsis can be regarded as an alarming health problem for the region[4-6]. A recent study in Brazil indicates a prevalence of sepsis in ICUs of 30% with a mortality rate of 55%[7], while in Argentina a mortality ranging from of 41.6% to 51%[4] has been ascribed to septic shock.

Considering this critical situation, in its 70[th] Assembly held in 2017, the World Health Organization reinforced the recommendations to improve the prevention, early diagnosis, follow-up and recording of sepsis cases, along with the promotion of research for the development of new treatments and diagnostic methods[8,9].

Despite the steady advances made in our understanding of sepsis pathophysiology in the last decades, this knowledge has not been fully translated into a substantial improvement in the therapeutic approach. Both diagnosis and treatment of sepsis still present numerous challenges[10]. A comprehensive discussion in 2016 led to the redefinition of sepsis (Sepsis-3) based on consensus recommendations, as a "life-threatening organic dysfunction caused by a dysregulated host response to infection"[11]. This concept highlights the need to integrate the information on the aetiological agent and the host response status, with a strong emphasis in the latter.

In this line, from the immunological point of view, sepsis is characterized by a dysregulated or dysfunctional host immune response to infection that varies over time; clinically, it shows the predominance of an overwhelming pro-inflammatory response during the early stages. As sepsis progresses, an anti-inflammatory and immunosuppressive response becomes predominant, which impacts directly on patient outcome[12] since the most deaths occur during this late stage due to opportunistic reinfections[12-14]. The dynamic nature of sepsis constrains the potential immunotherapy approaches due to the lack of tools to monitor the host immune status promptly.

In spite of the efforts put in biomarker search in the past decades, the application of clinically useful biological molecules or signatures for the diagnosis and the monitoring of sepsis is still finite[15-18]. FDA-approved biomarkers, such as procalcitonin, have a limited application as a guide for antibiotic treatment and as a prognostic marker under certain conditions[16,18-20]. Moreover, none of these approaches focused in the stages of the disease based on its quickly changing immunological status[12-14,16,18-20]. In this light, a fast and readily transferrable test is urgently needed.

Over the past decades, the emergence of "omics" technologies have proven to be a new source of promising results in the study of many diseases including sepsis. Matrix-assisted light desorption ionization-time of flight-mass spectrometry (MALDI-TOF-MS) is a powerful tool for studying protein

molecules present in complex biological samples, with high fidelity and in a few minutes[21]. Notably,
this technology proved to be extremely cost-effective in clinical microbiology for the identification of
bacteria[22], and it is becoming increasingly available in the clinical diagnosis setting, even in resource-
constrained countries. The field of proteome and peptidome analysis of diverse biological fluids has
also been dominated by mass spectrometry, and top-down analysis is progressively gaining ground.
This approach can provide information about the physiological state of an organism and contribute
to disease diagnosis[23], using for this purpose, analysis strategies such as artificial intelligence
algorithms. Machine learning (ML) is a concept in which generic algorithms are used to learn about
certain sample dataset. The algorithms are trained with these data to look for unique features among
groups. This knowledge is then applied to classify or cluster new data and make predictions[24].

Our approach based on the analysis of plasma biomolecular fingerprints has the advantage of
allowing the simultaneous evaluation of multiple markers, thus overriding costly and time-consuming
purification and immunolabeling steps. In addition, this method would be readily transferrable,
considering that MALDI-TOF-MS technology is currently available for microbiological biotyping in
several health care facilities and that analysis were performed using the open source software R. Our
final goal is to develop a fast, simple and inexpensive tool that can be available at health care centers
and can serve as a guide in the management of sepsis.

Around 50% of all sepsis cases are caused by Gram-negative bacteria[25,26] whose membrane
lipopolysaccharide (LPS) are one of the main agents associated both inflammatory and anti-
inflammatory processes[27,28]. Our group has intensively worked in LPS-based models that emulate
these phases[29-35]. Thus, in this study we tested the potential of our approach by developing well
defined animal models emulating pro- and anti-inflammatory phases that occur during a septic
process in LPS-induced murine models. The final goal of the current work is to develop predictive
models based on machine-learning algorithms that would allow the differentiation between the two
different phases (pro/anti-inflammatory) of sepsis through the analysis of the mass spectra of plasma
specimens generated by MALDI-TOF-MS.

RESULTS

Simulation of pro- and anti-inflammatory sepsis phases

The inflammatory phase was induced with a high dose of LPS (LPS group), whereas the anti
inflammatory/immunosuppression phase was reached with an scheme of increasing doses of LPS
(IS group) (inoculation and sample collection schemes are described in detail in Suppl. Fig.1). In order
to validate the two models, immunological, inflammatory and tissue damage markers were studied in
the two experimental groups as well in the control group (CTL group). In accordance with our previous
results, the pro-inflammatory phase was characterized by a marked leukopenia (Suppl. Fig 2a), high
levels of pro-inflammatory cytokines such as TNF-α in plasma, but also of the anti-inflammatory
cytokines IL-10 and TGF-β (Suppl. Fig 3a, b). Tissue enzymes indicative of damage were also
elevated (Suppl. Fig 4a-c). On the other hand, in the anti-inflammatory/immunosuppressed (IS) mice,

a leukocytosis due to increased numbers of circulating polymorphonuclear neutrophils (PMN) (Suppl.
Fig 2b) along with high plasma levels of TGF- β and low levels of pro-inflammatory cytokines were
observed (Suppl. Fig 3c). Tissue damage markers were unaltered in this group (Suppl. Fig 4d-f). As
previously described[32,34,35], IS mice had a profound immunological impairment, accompanied by a
marked lymphopenia and an increased number of PMN in the spleen, which expressed higher levels
of the inhibitory receptor PDL-1 (Suppl. Fig 5a, 5b). Moreover, IS group showed significantly low
antibody titres upon immunization with sheep red blood cells (SRBCs) (Suppl. Fig 5c). Collectively,
these results indicate that the pro- and anti-inflammatory/immunosuppression phases were emulated
in our murine models.

MALDI-TOF MS data analysis

Plasma samples of the different groups (CTL, IS and LPS) were analyzed by MALDI-TOF MS and
the spectra were obtained. Representative spectra of each experimental group are shown in Suppl
Fig. 6 and the analysis pipeline is detailed in Figure 1. A statistical analysis set was implemented to
test if MALDI-TOF MS data were useful to discriminate the three experimental groups. First, a
supervised classification algorithm was performed, searching for peaks that best differentiated the
CTL, LPS, and IS groups. The 20 best discriminant peaks identified by binary discriminant analysis
(BDA) and random forest (RF) algorithms are shown in Suppl. Fig. 7, which partially overlapped. Of
the 20 best peaks selected by the BDA algorithm, the 10 most discriminat peaks were chosen for a
unsupervised statistical analysis to plot the hierarchical k-means clustering-PCA clusters (Figure 2a).
Clusters appeared as three non-superimposed groups, and the first two principal components (PC)
explained 76.6% of the variation with good intra cluster homogeneity. Specifically, cluster 2 achieved
100% (16/16) of homogeneity for spectra corresponding to CTL plasma and 64% (16/25) of CTL
spectra fell in this cluster (Figure 2b), while for cluster 1 an 85% (29/34) homogeneity was achieved
for spectra corresponding to the LPS group and 100% (29/29) of LPS spectra fell in this cluster. Lastly,
homogeneity of cluster 3 was 77% (20/26) mostly represented by spectra of the IS group, with 91%
of the IS spectra (20/22) included in this cluster. This result show that the immunosuppression/anti-
inflammatory and pro-inflammatory phases have distinctive signatures that allow their discrimination.
139

Figure 1. Analysis Pipeline

Abbreviations. CTL: control group; IS: immunosuppression/anti-inflammatory group; LPS: inflammatory group; BDA: binary discriminant analysis; RF: random forest.

Figure 2

a

202 b

Figure 2. Unsupervised statistical analysis. (a) Hierarchical k-means clustering (Hkmc)-Principal Component Analysis (PCA) cluster plot using the top ten peaks selected by the binary discriminant analysis (BDA) algorithm. Labels contain the mice ID and immunological sepsis status. Principal components 1 (Dim1, x axis) and 2 (Dim2, y axis) are depcted. Spectra were clustered into three groups using the Hierarchical k-means clustering algorithm, which were represented with three different colours. 95% confidence ellipses were added around cluster means, assuming a multivariate normal distribution. (b) Hierarchical k-means clustering-PCA cluster composition. The green color represents the CTL mice group, the blue represents the IS group, and the red color represents the LPS group.

Machine learning analysis

Machine learning (ML) algorithms were used to develop a predictive model based on the plasma fingerprints, using a training data set (14 CTL, 13 IS and 15 LPS samples) challenged with a test data set (11 CTL, 9 IS and 14 LPS samples;. Figure 1). To find the most refined model, a dimensionality reduction was performed through the selection of different numbers of distinctive peaks for each physiological phase to train the algorithms, and the performances on the test set were computed for each condition. Specifically, the top five, ten, fifteen, and twenty discriminant peaks for BDA and RF algorithms were evaluated by cross-validation (CV; 5 folds and 20 repetitions) and percentages of accuracy, sensibility, specificity, negative predictive value (NPV) and positive predictive value (PPV) in the test data set were obtained (Table 1). Among models, RF using the best 20 peaks showed the most consistent values, with sensibility and specificity values above 90%. The BDA model trained with the best five peaks showed the best performance in terms of detection of positive cases, with a sensitivity above 95%; it strategy constitutes a more simple method in terms of both the number of required peaks and from theoretical viewpoint, being an approach more economical in terms of computational demands. These results display the potential of plasma fingerprints to find predictive models.

Table 1

Peaks	Algorithm	A (%)	S (%)	E (%)	PPV (%)	NPV (%)
5	BDA	90.6	95.7	80	90.9	89.8
10	BDA	85.6	92	72.3	87.4	81.1
15	BDA	87.6	93.9	74.5	88.5	85.4
20	BDA	90.1	94.3	81.4	91.4	87.3
5	RF	88.2	95.7	72.7	88	88
10	RF	91.2	91.3	90.9	95.5	83.3
15	RF	88.2	91.3	81.8	91.3	81.8
20	RF	94.1	95.7	90.9	95.7	90.9

Table 1. Performance of classification models. Performance values obtained with the top 5, 10, 15, and 20 peaks were presented. A = accuracy, S = sensitivity, E = specificity, PPV = positive predictive value, and NPV = negative predictive value.

Bioinformatic peak assignment

The 20 discriminant peaks (total = 25, 15 accredited by both methods, 5 accredited only by BDA and 5 accredited only by RF) were bioinformatically assigned to proteins/peptides matching entries in the Uniprot database according to their m/z listed in Suppl. Table 1. Most hits corresponded to fragments of bigger proteins (Suppl. Table 1). The only peptide identified in its native form was the antimicrobial peptide β-defensin-2, which was selected by both BDA and RF approaches as highly discriminant.

Other peptides identified in this analysis were β-enolase (7,643 Da peak), BRCA1 gene-1 protein, and lysosomal α-mannosidase (4,870 Da peak), among others (Suppl. Table 1). These peptides were also found in human plasma by Dufresne et al.[36]. Among fragments of the most abundant plasma proteins decribed by these authors, a fragment that matched the joining region of the immunoglobulin heavy chain was also found in this analysis.

DISCUSSION

Sepsis constitutes a highly heterogeneous syndrome of complex pathophysiology. The underlying immunological alterations that include pro-inflammatory, anti-inflammatory and immunosuppressive components[12] are reflected in septic plasma protein composition and varies over time according to disease severity[37]. Thus, plasma represents a favoured biological sample to search for biochemical signals that provide information on the physiological state at a given time. In the last decades, several circulating markers, both soluble as well as cell surface molecules with well recognized functions in sepsis were identified. However, none of these markers, nor a combination of them, showed the specificity and sensitivity required for its clinical use[15,18] .

Furthermore, methods that rely on immunolabelling as ELISA or flow cytometry, are highly specific and sensitive for the detection of low abundance markers, but requires costly equipment and consumables, as well as specific technical knowhow. In addition, these techniques are time consuming, limiting their clinical usefulness for early decision making.

Regarding biomarker discovery, proteomics proved to be an important source of information. However, in the classical approaches the plasma samples are subjected to depletion of abundant proteins[23] and /or fractionation to identify individual biomarkers[37,38] which requires more analysis time, specific equipment and higher costs.

These constraints led us to explore a new top-down strategy that could overcome these difficulties. Thus, having in mind that we aimed to design a readily transferrable method, our approach was meant to be as simple as possible at the beginning, to eventually introduce technical improvements as the analysis demanded. For this reasons pre-processing of the sample was avoided and whole plasma were directly read by MALDI-TOF-MS.

Surprisingly, the results described herein showed that the performance of the test with whole plasma and standard experimental settings used in routine microbiological biotyping were excellent in terms

of discriminatory power. First, an ensemble between supervised and unsupervised algorithms allowed
the overall discrimination of plasma samples from the three experimental groups, not only the ones
representing the immunosuppressive and the inflammatory phases from the control group, but also
between them. These results showed that each phase comprise a distinctive MALDI-TOF-MS pattern,
which turns this approach into a promising tool to determine different phases in sepsis. A recent study
reported a top-down plasma analysis showing that differences between samples from healthy donors
and cancer patients could be appreciated without the depletion of large abundant proteins by setting
a cutoff at 30KDa[39], supporting our results.

Furthermore, the performance of our machine learning algorithms challenged with the test dataset,
was able to identify basal, inflammatory, and immunosuppressive phases in a very satisfactory way,
exceeding 90% of accuracy, even supassing 95% in some of the tested coditions. The cross validation
strategy used herein allows a fine tuning of the models according to particular needs. Thus depending
on the particular clinical and epidemiological settings, algorithms with higher sensitivity, like our 5-
peaks BDA model, would be preferred over those with a good overall accuracy. In addition, this model
would be more economical in terms of computational demands.

The peaks found in the 2-20KDa spectrum range mostly correspond to small peptides. Plasma
peptidome has been proposed as an interesting source of information for diagnostic purposes[36,39].
These peptides are thought to be fragments of larger proteins derived from digestion by proteases or
non-enzymatic degradation[40]. Accordingly, recent reports demonstrate the importance of plasma
peptidome as a useful tool for clinical monitoring in septic and hemorragic shock[41,42]. Some of the
potential peptides identified in our analysis (Suppl. Table 1), were also found by other authors in
human plasma[36]. The only peptide assigned in its native form was the β-defensin-2, and was selected
by both BDA and RF approaches. This antimicrobial peptide has been proposed as an useful
biomarker in sepsis[37]. The identification of our discriminant peaks should be confirmed by more
specific methods, and the lack of comprehensive digestome/degradome databases is a well
recognized problem in the field[43] that could have conditioned our results.

A major limitation of this study is related to the animal models chosen. Although LPS-induced
inflammation/immunosuppression models are widely used, they do not accurately represent the
dynamic physiological changes that occur in sepsis[44]. Endotoxin challenge promotes a more quick
and transient release of pro/anti-inflammatory mediators compared to the septic processes. However
these models allowed us to define two discrete phases that actually occur in a septic process. For
this reason, we considered this was the most appropriate approach to test the potential of MALDI-
TOF-MS analysis in the proof-of concept phase. We are currently validating our results in a dynamic
model of sepsis, the cecal ligation and puncture model, with very encouraging results.

However, our approach has shown numerous advantages. First, MALDI-TOF-MS is a simple and
speedy method, which is a critical feature taking into account the dynamism of sepsis. This technology
is spreading in the healcare system, particularly in centers were patients with sepsis are usually
treated. In addition, we used an accessible sample as plasma, without complex pre-processing steps.
In addition, we performed our ML analysis in the open source softwere R. This language, that also

supports user friendly web-based interface integration, is another feature that guarantees the transferability even in resource contrained settings.

In the light of the results presented herein, we consider that plasma peptidome analysis by MALDI-TOF-MS has the potential to be a highly relevant strategy for sepsis patient stratification. Ideally, it would constitute a powerful tool for therapeutic decision making in sepsis depending on the pro- or anti-inflammatory phase the patient is undergoing.

MATERIALS AND METHODS

Animals

Female and male BALB/c mice (8–12 weeks old) were provided by the IMEX-CONICET-Academia Nacional de Medicina, Buenos Aires, Argentina. Animals were maintained under a 12 h light–dark cycle at 22 ± 2 °C and were fed with standard diet and water ad libitum. Animals were bred and housed in accordance with the NIH Guide and Use of Laboratory Animals[45]. Experimental designs were approved by the Committee for the Care and Use of Laboratory Animals (CICUAL) of IMEX-CONICET, Academia Nacional de Medicina de Buenos Aires.

Murine Models

Inoculation schemes are depicted in Supplementary Figure 1.

Inflammatory phase model (LPS group): BALB/c mice were inoculated with an LD_{50} of lipopolysaccharide of *Escherichia coli* O111: B4 (LPS) (100 ug/ mouse) (Sigma-Aldrich, St Louis, MO, USA) intraperitoneally (i. p.)[29]. Plasma samples were obtained 1.5 h and 6 h after the injection.

Anti-inflammatory phase model (IS group): BALB/c mice were daily inoculated with LPS for 10 consecutive days. The inoculation scheme consisted of 5 μg/mouse i.p. for the first 4 days, followed by 50 μg/mouse i.p. for 3 days, and then 100 μg/mouse i.p. for the last 3 days[34]. 24 h after the last LPS dose, plasma samples were obtained.

A third group of mice inoculated with vehicle (saline solution) served as control *(CTL group)*. The plasma in this group was collected at the same time as indicated in the experimental groups .

The submandibular bleeding method was used to obtain plasma in order to maximize the quality of the sample. The heparinized samples were centrifuged twice (400xg, 10 min at 4ºC) and plasma were stored at -20ºC until analysis.

Immunological, hematological and biochemical parameters

Peripheral blood leukocytes count: blood cells obtained by a submandibular puncture in heparinized
tubes were analyzed by Coulter hematology analyzer (Diatron Abacus Junior Vet, Budapest,
Hungary) at 1.5 h and 24 h post LPS for the LPS and the IS groups respectively. *Cytokine*
determination: pro-inflammatory (TNF-α; IL-12p70; IFN-γ; IL-6) and anti-inflammatory cytokines (IL-
10, TGF-β) were determined in plasma by ELISA at the indicated time points (OptEIA set; BD
Biosciences, San Diego, CA, USA) according to the manufacturer's instructions. *Tissue damage*
markers: creatinine (Cre), alanine aminotransferase (ALT) and aspartate transaminase (AST) levels
were determined in plasma, at 6 and 24 h in the LPS and IS groups respectively, using a kit from
BioTecnica (Varginha, Minas Gerais, Brazil) and the MINDRAY BS-200E autoanalyzer according to
the manufacturer's instructions. *Flow cytometry*: twenty four hours after the last LPS dose splenocytes
suspension was obtained and the cells were immunolabeled as described previously[34,35] for the
following cell types: T lymphocytes (FITC)-CD4 (clone RM4-5) and (PECy5)-CD8 (clone 53-6.7), B
lymphocytes (FITC)-CD45R-B220 (clone RA3-6B2), neutrophils (FITC)-CD11b (clone M1/70)/ (PE)-
Ly6G clone 1A8), and the expression levels of (PE)-PDL-1 (clone MIH5) on CD11b gate of myeloid
lineage cells. Antibodies were obtained from Invitrogen and BD Pharmingen™ . The acquisition was
carried out by a Becton Dickinson FACScan using Cell Quest software (Becton Dickinson, San Jose,
CA,USA). *Antibodies titre:* primary antibody response to sheep red blood cells (SRBC) by immunizing
animals 24 h after the last dose of LPS (SRBC: 5×10^8/mouse, 0.1ml i.p.). The anti-SRBC antibody
titre was evaluated in serum through a hemagglutination assay seven days after immunization as
described previously[34,35].

Acquisition of MALI-TOF-MS spectra

Plasma samples were analyzed with the MALDI Biotyper System (Bruker Daltonik GmbH, Bremen,
Germany). The total number of plasma samples analyzed per group were the following: LPS
group=n:29; IS group=n: 22 ; CTL group=n: 25. The standard method used for routine microbial
biotyping was used to analyse the plasma samples. Briefly, one µl of plasma was loaded onto each
spot of the metal plate in duplicate. After the sample was allowed to dry, one µl of the matrix (alpha-
cyano-4-hydroxycinnamic acid matrix in 50% acetonitrile and 2.5% trifluoroacetic acid) was added to
each spot and was allowed to dry again. Duplicates were acquired seeding two spots per sample.

Continuous mass spectra were obtained with a Microflex LT/SH MALDI-TOF mass spectrometer (with
flexControl software version 3.4.135.0; ionization mode: LD+; acquisition method: MBT_FC.par;
acquisition mode: qsim; tofMode: linear; acquisition Operator Mode: linear; digitizerType: Bruker
BD0G5; instrument ID: 269944.00510; instrument type: Microflex) within a mass range of 2,000-
20,000 Da. Spectra were obtained in the manual mode, using 60% of laser intensity with 40 laser
repetition in each shot and reaching between 400-500 spectra by acquisition. The rationale of
choosing these acquisition conditions were 1) to use the acquisition range set for bacterial biotyping
to avoid a specific calibration step and 2) to avoid the >20,000 Da range where highly abundant
proteins such as albumin mask other proteins with lower abundance. Spectra were internally

calibrated and controlled every day using *Escherichia coli* ribosomal proteins (bacterial test standard; Bruker Daltonik GmbH).

MALDI-TOF-MS data pre-processing

We generated a dataset containing 152 mass spectra from mice plasma samples (50 belonged to the CTL group, 44 to the IS group and 58 to the LPS group), including duplicates. Mass spectra were read as fid/aqus files with MALDIquantForeign (v0.10)[46] and they were processed using MALDIquant (v1.16.2)[47] R package. Briefly, spectra were square root-transformed and smoothed using the Savitzky-Golay algorithm. Spectra were baseline-corrected, applying the SNIP process across 100 iterations. The peak detection was carried out by a function that estimates the noise of mass spectrometry data by calculating the median absolute deviation (MAD). The *s*ignal-to-*n*oise-*r*atio (SNR) was set up in 4, with a halfWindowSize of 40 and a tolerance of 0.2 for peak binning. One replicate corresponding to the CTL group was removed due to low quality. After averaging the replicates, 76 averaged mass spectra were obtained, of which 25 belonged to the Control group, 22 to the IS group and 29 to LPS group. Peaks that occured in less than 33% of spectra were removed. MALDI-TOF-MS data were transformed by a categorization of the peak intensity, performed with the Binda R package[48]. The algorithm compares the intensity value with the peak group average, returning 1 when it was equal or higher and 0 when it was lower. The analysis workflow is depicted in the Figure 1.

Statistical analysis

Immunological, hematological and biochemical parameters. Statistical analysis and plotting were achieved using Graph Pad Prism 6 software (GraphPad Software, San Diego, CA). Values are expressed as the mean ± standard error of the mean (SEM) of n samples. The number of samples (n) is refers to number of mice or biological repeats analyzed each time. These biological repeats were used for the statistical analysis. No outliers were removed. The assumption test to determine the Gaussian distribution was performed by the Kolmogorov and Smirnov method. For parameters with a Gaussian distribution, the differences between 2 experimental groups were assessed by unpaired Student's t test. For multiple group comparisons, as the peripheral blood leukocytes differences, were analyzed using one-way ANOVA followed by a Tukey's multiple comparison test. For parameters with a non-Gaussian distribution, comparisons between 2 experimental groups were performed using Mann–Whitney U test. All statistical tests were interpreted in a two-tailed fashion and a $P<0.05$ are considered statistically significant; the level of significance is indicated by asterisk numbers as follows: *, $P<0.05$; **, $P<0.01$; ***, $P<0.001$, ****, $P<0.0001$.The statistical analysis used as also the number of samples (n) and the repetitions of each experimental design are reported in each figure legend.

MALDI-TOF-MS data. Statistical analysis and plotting were performed using Rstudio Software. Programmed peaks selection was performed in the entire dataset seeking for biomarkers of each class representing the different sepsis phases (LPS and IS groups) by the Binary Discriminant Analysis (BDA) algorithm. The algorithm outputs the t.score (Class means vs. Pooled mean) of each peak. The sign of the t.score indicates the presence (positive t.score) or absence (negative t.score) of that peak in each group. A significance level of 95% was achieved if the t.score was equal or higher than 2.5 and equal or less than -2.5. The best-extracted features were then used to perform a hierarchical k-means clustering-principal component analysis (PCA) with the factoextra R package[49]. The binary distance was used to measure dissimilarity between observations, ward.D2 was the agglomeration method used, and k was set to 4. In addition, spectra were analyzed with the random forest (RF) classifier. Afterwards, two machine learning (ML) methods were applied in the dataset to train two classifiers: BDA and random forest RF[50] also with RStudio software.

These classification models, used for the feature extraction and ranking, performe their selection based on distinctive criteria. For instance, BDA seeks for differences by measuring the t-scores between the class mean and pooled mean, while RF computes the total decrease in node impurities from splitting on the variable, averaged over all decision trees, and the Gini index measures the node impurity. Initially, the dataset was randomly partitioned into a training set (60% of plasma samples) and a test set (40% of plasma samples). Programmed feature (peaks) selection was performed with the respective algorithms in the training set seeking for discriminant peaks corresponding to each sepsis phase. The extracted features were then used to train several ML models. Metrics such as accuracy, sensitivity, and specificity, were used to evaluate the performances of all resulting models in the test set by a cross-validation strategy. The entire dataset analyzed in this work together with the R sample code used for the analysis is available at: https://github.com/MarManLed/SepsisData.

Bioinformatic peak assignment

Uniprot database was used to assign the best 20 discriminant peaks. The search was performed on the *Mus musculus* data base (ID) and within a 15 Da interval around each peak.

Acknowledgements

We thank Dr. Martín Isturiz for his eternal accompaniment and support for the development of this study. His differential contribution stems from his singular thought. Forever in our hearts, Dr. Isturiz. We thank Dr. Viviana Ritacco for her thoughtful comments on the manuscript.

Funding

This study was supported by the Agencia Nacional de Promoción Científica y Tecnológica [grant number PICT 2015-0412] Buenos Aires, Argentina.

Author contribution

ML designed and performed the bioinformatic analysis. BR conceived and designed the experiments. MFT, LM and BR performed the experiments with mice and analyzed data. MP and CV provided materials and technical support. MG and BL provided materials and collaborated in study design. ML and NY contributed in data curation. NY and BR analyzed, discussed and realized the study. BR was responsible for funding acquisition and study supervision. Manuscript was written by NY and BR, and was corrected by ML and BL. All the authors revised and approved the final version of the manuscript.

Competing interests

The authors declare that no competing interests exist.

References

Novosad, S. A., Sapiano, M. R., Grigg, C., Lake, J., Robyn, M., Dumyati, G., Felsen, C., Blog, D., Dufort, E., Zansky, S., Wiedeman, K., Avery, L., Dantes, R. B., Jernigan, J. A., Magill, S. S., Fiore, A. & Epstein, L. Vital Signs: Epidemiology of Sepsis: Prevalence of Health Care Factors and Opportunities for Prevention. *MMWR Morb Mortal Wkly Rep* **65**, 864-869, doi:10.15585/mmwr.mm6533e1 (2016).

Vincent, J. L., Rello, J., Marshall, J., Silva, E., Anzueto, A., Martin, C. D., Moreno, R., Lipman, J., Gomersall, C., Sakr, Y. & Reinhart, K. International study of the prevalence and outcomes of infection in intensive care units. *JAMA* **302**, 2323-2329, doi:10.1001/jama.2009.1754 (2009).

Fleischmann, C., Scherag, A., Adhikari, N. K., Hartog, C. S., Tsaganos, T., Schlattmann, P., Angus, D. C. & Reinhart, K. Assessment of Global Incidence and Mortality of Hospital-treated Sepsis. Current Estimates and Limitations. *Am J Respir Crit Care Med* **193**, 259-272, doi:10.1164/rccm.201504-0781OC (2016).

Estenssoro, E., Kanoore Edul, V. S., Loudet, C. I., Osatnik, J., Rios, F. G., Vazquez, D. N., Pozo, M. O., Lattanzio, B., Palizas, F., Klein, F., Piezny, D., Rubatto Birri, P. N., Tuhay, G., Diaz, A., Santamaria, A., Zakalik, G. & Dubin, A. Predictive Validity of Sepsis-3 Definitions and Sepsis Outcomes in Critically Ill Patients: A Cohort Study in 49 ICUs in Argentina. *Crit Care Med* **46**, 1276-1283, doi:10.1097/CCM.0000000000003208 (2018).

Estenssoro, E., Loudet, C. I., Edul, V. S. K., Osatnik, J., Rios, F. G., Vasquez, D. N., Pozo, M. O., Lattanzio, B., Palizas, F., Klein, F., Piezny, D., Rubatto Birri, P. N., Tuhay, G., Diaz, A., Santamaria, A., Zakalik, G. & Dubin, A. Health inequities in the diagnosis and outcome of sepsis in Argentina: a prospective cohort study. *Crit Care* **23**, 250, doi:10.1186/s13054-019-2522-6 (2019).

Azevedo, L. C. P., Cavalcanti, A. B., Lisboa, T., Pizzol, F. D. & Machado, F. R. Sepsis is an important healthcare burden in Latin America: a call to action! *Rev Bras Ter Intensiva* **30**, 402-404, doi:10.5935/0103-507X.20180061 (2018).

Machado, F. R., Cavalcanti, A. B., Bozza, F. A., Ferreira, E. M., Angotti Carrara, F. S., Sousa, J. L., Caixeta, N., Salomao, R., Angus, D. C. & Pontes Azevedo, L. C. The epidemiology of sepsis in Brazilian intensive care units (the Sepsis PREvalence Assessment Database, SPREAD): an observational study. *Lancet Infect Dis* **17**, 1180-1189, doi: 10.1016/S1473-3099(17)30322-5 (2017).

512   8   Organization, W. H. WHO: WHA70.7 Improving the Prevention, Diagnosis and Clinical Management of Sepsis.
513   doi:http://apps.who.int/gb/e/e_wha70.html. Accessed October 2, 2017 (2017).
514
515   9   Reinhart, K., Daniels, R., Kissoon, N., Machado, F. R., Schachter, R. D. & Finfer, S. Recognizing Sepsis as a Global
516   Health Priority - A WHO Resolution. *N Engl J Med* **377**, 414-417, doi:10.1056/NEJMp1707170 (2017).
517
10   Rhodes, A., Evans, L. E., Alhazzani, W., Levy, M. M., Antonelli, M., Ferrer, R., Kumar, A., Sevransky, J. E., Sprung,
C. L., Nunnally, M. E., Rochwerg, B., Rubenfeld, G. D., Angus, D. C., Annane, D., Beale, R. J., Bellinghan, G. J.,
Bernard, G. R., Chiche, J. D., Coopersmith, C., De Backer, D. P., French, C. J., Fujishima, S., Gerlach, H., Hidalgo, J.
L., Hollenberg, S. M., Jones, A. E., Karnad, D. R., Kleinpell, R. M., Koh, Y., Lisboa, T. C., Machado, F. R., Marini, J.
J., Marshall, J. C., Mazuski, J. E., McIntyre, L. A., McLean, A. S., Mehta, S., Moreno, R. P., Myburgh, J., Navalesi,
P., Nishida, O., Osborn, T. M., Perner, A., Plunkett, C. M., Ranieri, M., Schorr, C. A., Seckel, M. A., Seymour, C.
W., Shieh, L., Shukri, K. A., Simpson, S. Q., Singer, M., Thompson, B. T., Townsend, S. R., Van der Poll, T., Vincent,
J. L., Wiersinga, W. J., Zimmerman, J. L. & Dellinger, R. P. Surviving Sepsis Campaign: International Guidelines for
Management of Sepsis and Septic Shock: 2016. *Intensive Care Med* **43**, 304-377, doi:10.1007/s00134-017-4683-
6 (2017).
11   Singer, M., Deutschman, C. S., Seymour, C. W., Shankar-Hari, M., Annane, D., Bauer, M., Bellomo, R., Bernard,
G. R., Chiche, J. D., Coopersmith, C. M., Hotchkiss, R. S., Levy, M. M., Marshall, J. C., Martin, G. S., Opal, S. M.,
Rubenfeld, G. D., van der Poll, T., Vincent, J. L. & Angus, D. C. The Third International Consensus Definitions for
Sepsis and Septic Shock (Sepsis-3). *JAMA* **315**, 801-810, doi: 10.1001/jama.2016.0287 (2016).

12   Rubio, I., Osuchowski, M. F., Shankar-Hari, M., Skirecki, T., Winkler, M. S., Lachmann, G., La Rosee, P., Monneret,
G., Venet, F., Bauer, M., Brunkhorst, F. M., Kox, M., Cavaillon, J. M., Uhle, F., Weigand, M. A., Flohe, S. B.,
Wiersinga, W. J., Martin-Fernandez, M., Almansa, R., Martin-Loeches, I., Torres, A., Giamarellos-Bourboulis, E.
J., Girardis, M., Cossarizza, A., Netea, M. G., van der Poll, T., Scherag, A., Meisel, C., Schefold, J. C. & Bermejo-
Martin, J. F. Current gaps in sepsis immunology: new opportunities for translational research. *Lancet Infect Dis*,
doi: 10.1016/S1473-3099(19)30567-5 (2019).

13   van der Poll, T., van de Veerdonk, F. L., Scicluna, B. P. & Netea, M. G. The immunopathology of sepsis and
potential therapeutic targets. *Nat Rev Immunol* **17**, 407-420, doi: 10.1038/nri.2017.36 (2017).

14   Venet, F., Rimmele, T. & Monneret, G. Management of Sepsis-Induced Immunosuppression. *Crit Care Clin* **34**,
97-106, doi: 10.1016/j.ccc.2017.08.007 (2018).

15   Parlato, M., Philippart, F., Rouquette, A., Moucadel, V., Puchois, V., Blein, S., Bedos, J. P., Diehl, J. L., Hamzaoui,
O., Annane, D., Journois, D., Ben Boutieb, M., Esteve, L., Fitting, C., Treluyer, J. M., Pachot, A., Adib-Conquy, M.,
Cavaillon, J. M., Misset, B. & Captain Study, G. Circulating biomarkers may be unable to detect infection at the
early phase of sepsis in ICU patients: the CAPTAIN prospective multicenter cohort study. *Intensive Care Med* **44**,
1061-1070, doi:10.1007/s00134-018-5228-3 (2018).
547
16   Gunsolus, I. L., Sweeney, T. E., Liesenfeld, O. & Ledeboer, N. A. Diagnosing and Managing Sepsis by Probing the
Host Response to Infection: Advances, Opportunities, and Challenges. *J Clin Microbiol* **57**, doi:
10.1128/JCM.00425-19 (2019).

17   Pierrakos, C. & Vincent, J. L. Sepsis biomarkers: a review. *Crit Care* **14**, R15, doi: 10.1186/cc8872 (2010).

18   van Engelen, T. S. R., Wiersinga, W. J., Scicluna, B. P. & van der Poll, T. Biomarkers in Sepsis. *Crit Care Clin* **34**,
139-152, doi: 10.1016/j.ccc.2017.08.010 (2018).

19 Al Jalbout, N., Troncoso, R., Jr., Evans, J. D., Rothman, R. E. & Hinson, J. S. Biomarkers and Molecular Diagnostics
for Early Detection and Targeted Management of Sepsis and Septic Shock in the Emergency Department. *J Appl*
Lab Med **3**, 724-729, doi: 10.1373/jalm.2018.027425 (2019).

20 Schenz, J., Weigand, M. A. & Uhle, F. Molecular and biomarker-based diagnostics in early sepsis: current
challenges and future perspectives. *Expert Rev Mol Diagn*, 1-10, doi:10.1080/14737159.2020.1680285 (2019).

21 Angeletti, S. Matrix assisted laser desorption time of flight mass spectrometry (MALDI-TOF MS) in clinical
microbiology. *J Microbiol Methods* **138**, 20-29, doi: 10.1016/j.mimet.2016.09.003 (2017).

22 Hou, T. Y., Chiang-Ni, C. & Teng, S. H. Current status of MALDI-TOF mass spectrometry in clinical microbiology. *J*
Food Drug Anal **27**, 404-414, doi: 10.1016/j.jfda.2019.01.001 (2019).

23 Ludwig, K. R. & Hummon, A. B. Mass spectrometry for the discovery of biomarkers of sepsis. *Mol Biosyst* **13**,
648-664, doi:10.1039/c6mb00656f (2017).

24 López Fernández, H., Reboiro-Jato, M., Pérez Rodríguez, J. A., Fdez-Riverola, F. & Glez-Peña, D. Implementing
effective machine learning-based workflows for the analysis of mass spectrometry data. *Journal of Integrated*
OMICS **6**, doi:10.5584/jiomics.v6i1.196 (2016).

25 Dolin, H. H., Papadimos, T. J., Chen, X. & Pan, Z. K. Characterization of Pathogenic Sepsis Etiologies and Patient
Profiles: A Novel Approach to Triage and Treatment. *Microbiol Insights* **12**, doi:10.1177/1178636118825081
(2019).

26 van Engelen, T. S. R., Joost Wiersinga, W. & van der Poll, T. Phatogenesis of sepsis. *Handbook of Sepsis* Ch.
Chapter 3, 31-43 doi: 10.1007/978-3-319-73506-1_3 (2018).

27 Genga, K. R., Shimada, T., Boyd, J. H., Walley, K. R. & Russell, J. A. The Understanding and Management of
Organism Toxicity in Septic Shock. *J Innate Immun* **10**, 502-514, doi:10.1159/000487818 (2018).

28 Opal, S. M., Scannon, P. J., Vincent, J. L., White, M., Carroll, S. F., Palardy, J. E., Parejo, N. A., Pribble, J. P. &
Lemke, J. H. Relationship between plasma levels of lipopolysaccharide (LPS) and LPS-binding protein in patients
with severe sepsis and septic shock. *J Infect Dis* **180**, 1584-1589, doi:10.1086/315093 (1999).

29 Cordoba-Moreno, M. O., Todero, M. F., Fontanals, A., Pineda, G., Daniela, M., Yokobori, N., Ramos, M. V.,
Barrientos, G., Toblli, J. E., Isturiz, M. A. & Rearte, B. Consequences of the Lack of IL-10 in Different Endotoxin
Effects and its Relationship With Glucocorticoids. *Shock* **52**, 264-273, doi:10.1097/SHK.0000000000001233
(2019).

30 Landoni, V. I., Chiarella, P., Martire-Greco, D., Schierloh, P., van-Rooijen, N., Rearte, B., Palermo, M. S., Isturiz,
M. A. & Fernandez, G. C. Tolerance to lipopolysaccharide promotes an enhanced neutrophil extracellular traps
formation leading to a more efficient bacterial clearance in mice. *Clin Exp Immunol* **168**, 153-163,
doi:10.1111/j.1365-2249.2012.04560.x (2012).

31 Martire-Greco, D., Landoni, V. I., Chiarella, P., Rodriguez-Rodrigues, N., Schierloh, P., Rearte, B., Isturiz, M. A. &
Fernandez, G. C. all-trans-retinoic acid improves immunocompetence in a murine model of lipopolysaccharide-
induced immunosuppression. *Clin Sci (Lond)* **126**, 355-365, doi:10.1042/CS20130236 (2014).

32 Montagna, D. R., Duarte, A., Todero, M. F., Ruggiero, R. A., Isturiz, M. & Rearte, B. Meta-tyrosine modulates the
immune response induced by bacterial endotoxins. *Immunobiology* **225**, 151856,
doi:10.1016/j.imbio.2019.10.005 (2020).
603

604 33 Rearte, B., Landoni, V., Laborde, E., Fernandez, G. & Isturiz, M. Differential effects of glucocorticoids in the
605   establishment and maintenance of endotoxin tolerance. *Clin Exp Immunol* **159**, 208-216, doi:10.1111/j.1365-
606   2249.2009.04052.x (2010).
607

608 34 Rearte, B., Maglioco, A., Balboa, L., Bruzzo, J., Landoni, V. I., Laborde, E. A., Chiarella, P., Ruggiero, R. A.,
609   Fernandez, G. C. & Isturiz, M. A. Mifepristone (RU486) restores humoral and T cell-mediated immune response
610   in endotoxin immunosuppressed mice. *Clin Exp Immunol* **162**, 568-577, doi:10.1111/j.1365-2249.2010.04262.x
611   (2010).
612

613 35 Rearte, B., Maglioco, A., Machuca, D., Greco, D. M., Landoni, V. I., Rodriguez-Rodrigues, N., Meiss, R., Fernandez,
614   G. C. & Isturiz, M. A. Dehydroepiandrosterone and metyrapone partially restore the adaptive humoral and
615   cellular immune response in endotoxin immunosuppressed mice. *Innate Immun* **20**, 585-597,
616   doi:10.1177/1753425913502243 (2014).
617

618 36 Dufresne, J., Bowden, P., Thavarajah, T., Florentinus-Mefailoski, A., Chen, Z. Z., Tucholska, M., Norzin, T., Ho, M.
619   T., Phan, M., Mohamed, N., Ravandi, A., Stanton, E., Slutsky, A. S., Dos Santos, C. C., Romaschin, A., Marshall, J.
620   C., Addison, C., Malone, S., Heyland, D., Scheltens, P., Killestein, J., Teunissen, C., Diamandis, E. P., Siu, K. W. M.
621   & Marshall, J. G. The plasma peptidome. *Clin Proteomics* **15**, 39, doi:10.1186/s12014-018-9211-3 (2018).
622

623 37 Hayashi, N., Yamaguchi, S., Rodenburg, F., Ying Wong, S., Ujimoto, K., Miki, T. & Iba, T. Multiple biomarkers of
624   sepsis identified by novel time-lapse proteomics of patient serum. *PLoS One* **14**, e0222403,
625   doi:10.1371/journal.pone.0222403 (2019).
626

627 38 Harberts, E., Liang, T., Yoon, S. H., Opene, B. N., McFarland, M. A., Goodlett, D. R. & Ernst, R. K. Toll-like Receptor
628   4-Independent Effects of Lipopolysaccharide Identified Using Longitudinal Serum Proteomics. *J Proteome Res*
629   **19**, 1258-1266, doi:10.1021/acs.jproteome.9b00765 (2020).
630

631 39 Cheon, D. H., Nam, E. J., Park, K. H., Woo, S. J., Lee, H. J., Kim, H. C., Yang, E. G., Lee, C. & Lee, J. E. Comprehensive
632   Analysis of Low-Molecular-Weight Human Plasma Proteome Using Top-Down Mass Spectrometry. *J Proteome*
633   *Res* **15**, 229-244, doi:10.1021/acs.jproteome.5b00773 (2016).
634

635 40 Greening, D. W. & Simpson, R. J. Characterization of the Low-Molecular-Weight Human Plasma Peptidome.
636   *Methods Mol Biol* **1619**, 63-79, doi:10.1007/978-1-4939-7057-5_6 (2017).
637

638 41 Bauza-Martinez, J., Aletti, F., Pinto, B. B., Ribas, V., Odena, M. A., Diaz, R., Romay, E., Ferrer, R., Kistler, E. B.,
639   Tedeschi, G., Schmid-Schonbein, G. W., Herpain, A., Bendjelid, K. & de Oliveira, E. Proteolysis in septic shock
640   patients: plasma peptidomic patterns are associated with mortality. *Br J Anaesth* **121**, 1065-1074,
641   doi:10.1016/j.bja.2018.05.072 (2018).
642

643 42 Aletti, F., Maffioli, E., Negri, A., Santamaria, M. H., DeLano, F. A., Kistler, E. B., Schmid-Schonbein, G. W. &
644   Tedeschi, G. Peptidomic Analysis of Rat Plasma: Proteolysis in Hemorrhagic Shock. *Shock* **45**, 540-554,
645   doi:10.1097/SHK.0000000000000532 (2016).
646

647 43 Shen, Y., Liu, T., Tolić, N., Petritis, B. O., Zhao, R., Moore, R. J., Purvine, S. O., Camp, D. G. & Smith, R. D. Strategy
648   for Degradomic-Peptidomic Analysis of Human Blood Plasma. *Journal of Proteome Research* **9**, 2339-2346,
649   doi:10.1021/pr901083m (2010).
650
651 44 Lewis, A. J., Seymour, C. W. & Rosengart, M. R. Current Murine Models of Sepsis. *Surg Infect (Larchmt)* **17**, 385-
652   393, doi:10.1089/sur.2016.021 (2016).
653
654 45 Institute of Laboratory Animal Resources (U.S.). Committee on Care and Use of Laboratory Animals and National
655   Institutes of Health (U.S.). Division of Research Resources. Guide for the care and use of laboratory animals.
656   Bethesda, MD: US Department Health and Human Services, Public Health Service, National Institutes of Health..
657   (1985).
658
659 46 MALDIquantForeign: Import/Export Routines for MALDIquant. v. 0.12 (2019).
660
660 47 Gibb, S. & Strimmer, K. MALDIquant: a versatile R package for the analysis of mass spectrometry data.
661   *Bioinformatics* **28**, 2270-2271, doi:10.1093/bioinformatics/bts447 (2012).
662
663 48 Gibb, S., Strimmer, K. binda: Multi-Class Discriminant Analysis using Binary Predictors. R package version 1.0.3.
664   (2015).
665
666 49 factoextra: Extract and Visualize the Results of Multivariate Data Analyses v. R package version 1.0.6 (2019).
667
668 50 Liaw, A. & Wiener, M. Classification and Regression by randomForest. *R News* **2**, 18-22, doi:
669   10.4236/ijcns.2016.95010 (2002).
670