Elucidation of Japanese pepper （*Zanthoxylum piperitum* De Candolle)

domestication using RAD-Seq

Maddumage Dona Ginushika Priyadarshani Premarathne[1,2], Nami Fukutome[3, #], Kazuaki

Yamasaki[4], Fumiyo Hayakawa[5], Atsushi J. Nagano[6], Hisataka Mizuno[7], Nobuo Ibaragi[8],

Yukio Nagano[1, 2, #, *]

[1] Analytical Research Center for Experimental Sciences, Saga University

[2] Graduate School of Advanced Health Science, Saga University

[3] Department of Food Sciences, Tokyo Seiei College

[4] Faculty of Health Science, Hyogo University

[5] Food Research Institute, National Agriculture and Food Research Organization

[6] Faculty of Agriculture, Ryukoku University

[7] Gifu Agriculture and Forestry Office

[8] Association for Conservation of Asakura Sanshō

[#]These authors contributed equally to this work

*Correspondence to Y. N. (nagano@cc.saga-u.ac.jp)

Abstract

Japanese pepper, *Zanthoxylum piperitum*, is native to Japan and has four well-known

lineages (Asakura, Takahara, Budou, and Arima), which are named after their production

area or morphology. Restriction-site associated DNA sequencing (RAD-Seq) was used to

analyse 93 accessions from various areas, including these four lineages. Single nucleotide

variant analysis was used to classify the plants into eight groups: the Asakura and Arima

lineages each had two groups, the Takahara and Budou lineages each had one group, and

two additional groups were present. In one Asakura group and two Arima groups, the

plants were present in agricultural fields and mountains, thus representing the early stage

of domestication of the Japanese pepper. The second Asakura lineage group was closely

related to plants present in various areas, and this represents the second stage of

domestication of this plant because, after early domestication, genetically related lineages

with desirable traits spread to the periphery. These results demonstrate that domestication

of Japanese pepper is ongoing. In addition, this study shows that spineless plants are

polyphyletic, despite the spineless lineage being considered a subspecies of Japanese

pepper.

Introduction

Zanthoxylum species belongs to the family Rutaceae, commonly known as the citrus family. Many *Zanthoxylum* species are distributed worldwide and are used for culinary and medicinal purposes. For example, the Āyurvedic traditional medicine of India and Sri Lanka uses *Z. tetraspermum* and *Z. rhetsa* ("*Katu Keena*")[1]. Sichuan pepper is an essential ingredient in the cuisines of the Sichuan province of China and contains several species of *Zanthoxylum*, including *Z. simulans*. *Z. simulans* is used in traditional Chinese medicine for its stomachic, analgesic, and anthelmintic properties[2].

Zanthoxylum piperitum is native to Japan (except for Ryukyu) and southern Korea. In Japan, it is mainly used for culinary purposes. In English, it is known as Japanese pepper; in Japanese, it is called sanshō. This plant is considered to be highly medicinal in various uses[3,4,5,6,7,8,9,10]. Although *Zanthoxylum piperitum* is widely consumed and distributed throughout Japan, lineages have been formed based on the production area and morphology of this plant. The Asakura, Arima, and Takahara lineages are classified based on the place of production, while the Budou lineage is classified based on its morphology.

The Asakura lineage is believed to have originated in the Asakura district of Yabu City in central Hyōgo Prefecture. A typical feature of this lineage is the absence of spines. For this reason, the spineless plant is called the Asakura lineage in Japan. This spineless plant has been identified as a subspecies of *Z. piperitum* and named *Z. piperitum* (L.) DC forma *inerme* (Makino) Makino[11]. However, there is no evidence that all spineless plants are either from Asakura or are monophyletic. In Yabu city, *Zanthoxylum piperitum* grow as wild plants in mountainous areas and as cultivated plants in agricultural fields and gardens.

The Arima lineage, also known as the Rokko lineage, is another Japanese pepper lineage found in Kobe City in southern Hyōgo Prefecture. The flowers of these plants are typically used as ingredients in Arima cuisine.

The Takahara lineage is grown in Takayama City in northern Gifu Prefecture. These

plants are grown at high altitudes 800 m above sea level, in contrast to the other lineages,

which are cultivated on the plains. These plants bloom in mid-May, whereas other

lineages bloom in late April. The wild Takahara lineage is difficult to find in the

mountains of this region.

The Budou lineage (Budou means "grape" in Japanese) is categorised based on its

morphology. This lineage is cultivated mainly in Kainan City, Wakayama Prefecture. It

bears clusters of fruits that resemble grapes. According to oral tradition, it appeared as a

chance seedling in a garden about 200 years ago and has been cultivated ever since.

Therefore, the wild lineage is probably absent.

Many studies have investigated the genetic relationships among *Zanthoxylum*

species[12,13,14,15,16,17,18,19,20,21,22,23]. There are no reports that compare the intraspecific

diversity of *Zanthoxylum piperitum* in detail. However, three important studies have

examined the DNA sequences of *Zanthoxylum piperitum* growing in Korea. One is the

complete sequencing of the chloroplast genome[12], and the others involved the

development of markers to identify *Zanthoxylum piperitum*[13,14]. Many other studies have

used the genetic information for *Zanthoxylum piperitum* as a reference for

comparison[15,16,17,18,19,20,22,23]. There are two important studies on the intraspecific genetic

diversity of *Zanthoxylum* species, which focus on: (1) the relationship and diversity

between wild and cultivated plants[22] and (2) the genetic structure of cultivated plants[23].

A previous study[22] compared wild plants to cultivated plants using DNA sequence data

and detected genetic similarities between cultivated and wild *Z. armatum*. Although this

finding may be relevant to the domestication of *Z. armatum*, the study only used a small

number of markers and was unable to examine the species domestication details.

Studies on intraspecific genetic diversity of *Zanthoxylum* species have been based on

the analysis of a small number of markers, such as sequence-related amplified

polymorphism markers[22] and simple sequence repeat (SSR) markers[23]. Compared to

genetic studies using a small number of markers, high-throughput sequencing can

generate many single nucleotide variants (SNVs) that are suitable for analysing

intraspecific differences[24,25]. Restriction site-associated DNA sequencing (RAD-Seq)[26] is

a high-throughput sequencing method. Initially, RAD-Seq cleaves individual genomes

using a selected site-specific restriction enzyme. Then, DNA fragments adjacent to the

restriction site are sequenced using a high-throughput sequencer. Double-digest RAD-

Seq (ddRAD-Seq)[27] is a variation of RAD-Seq that uses two restriction enzymes to cleave

the genome. These methods allow for reduced representation of individual genomes.

Moreover, these methods do not require prior knowledge of the genome sequence[27]. Our

group has previously used the original RAD-Seq and ddRAD-Seq methods to analyse

citrus[28,29,30], loquat[31], and firefly[32].

The four lineages are well known, and *Zanthoxylum piperitum* is cultivated all over

Japan, except for Ryukyu. Some lineages are present in both the mountains and

agricultural fields. Therefore, the domestication of Japanese pepper may be an ongoing

process. In the present study, ddRAD-Seq was used to elucidate the intraspecific genetic

diversity and occurrence of domestication in Japanese pepper.

Results

Variant detection by *de novo* mapping of RAD-Seq data

Double digest RAD-Seq generated more than 11.9 gigabases of data for a total of 235.2

million raw single-end 51-bp reads. Quality-based filtering resulted in an average of 2.5

million reads (maximum of 4.3 million and a minimum of 0.4 million) among 93 samples

(Supplementary Table 1). The Stacks program built loci *de novo* with an average coverage

depth of 27.32 times (Supplementary Table 2). We performed the following analysis

using data from 4,334 variant sites.

Correlation of genetic similarity with geographic locations

Based on the principal component analysis (PCA) (Figure 1) and geographical

information (Figure 2, Table 1), 93 samples were classified into seven groups. PC1

divided group A from the other groups (Figure 1a); PC2 divided groups D and E from the

other groups (Figure 1a); PC3 (Figure 1b) identified groups C and F and confirmed the

presence of group E; PC4 identified group B (Figure 1c) and confirmed the presence of

groups C and F; PC5 reconfirmed the presence of groups B and F (Figure 1d); and PC6

identified group G (Figure 1e). Thus, probably because the genetic relationships among

plants were complex, it was necessary to use PC1 through PC6 for classification.

Group A may be an Asakura lineage group because the area across which the group A

plants were distributed coincided with the production area for the Asakura lineage in

Yabu City, central Hyogo Prefecture (Figure 2, Table 1). All plants in group A were

spineless. Group A was widely distributed and cultivated in other areas, including Kobe

and Kami in Hyogo Prefecture, Ayabe, and Kyōtamba in Kyoto Prefecture, and Gojō in

Nara Prefecture (Table 1). Interestingly, plants 42, 43, 44, 45, 46, 47, 48, 49, 50, and 51

formed tight cluster, and are from Yabu, Ayabe, Gojō, Kyōtamba, Yabu, Kobe, Yabu,

Gojō, Gojō, and Kami, respectively. The other plants in group A (35, 36, 37, 38, 39, 40,

and 41) were from Yabu, Kyōtamba, Kobe, Yabu, Yabu, Kyōtamba, and Kyōtamba,

respectively.

The analysis identified group D as the second Asakura lineage group. Group D plants

were from Yabu City only (Figure 1a, Figure 2, Table 1), except for plant 74, which was

from Kami. Unlike group A, which had more cultivated plants, group D contained many wild plants found in mountainous areas. Most, but not all, of the group D plants were spineless. Plant 58 was 100 years old and was growing in an agricultural field. When we observed plant 58 in 2017, this plant was spineless. As of 2020, this plant was near death, and its spines were visible.

The analysis identified group E as the Takahara lineage. Group E (Figure 1a) was an isolated group with plants only from Takayama city, located in the highland region of northern Gifu Prefecture (Figure 2, Table 1). This group consisted of only cultivated plants. Most of the group E plants were spineless (Table 1).

The analysis identified group C as the first group of the Arima lineage. Group C plants were from Kobe and Asago cities in southern Hyogo Prefecture (Figure 2, Table 1), where the Arima lineage was produced. There is an agricultural field in Asago City for the preservation of the Arima lineage. All plants in group C were spiny and transplanted from Mt. Rokko (Table 1).

The analysis identified group F as the second Arima lineage group. All plants in group F (Table 1) were from the Arima lineage production area in Asago and Kobe (Figure 2). Group F contained spiny plants. Three plants cultivated at the Fruit Tree Research Institute came from Mt. Inariyama, located near Arima. PCA did not clearly separate group F from the other plants.

The analysis identified group B as the Budou lineage. Group B was from Kainan city in Wakayama Prefecture (Figure 2), where the Budou lineage is produced (Table 1). Group B is the grape-like Budou lineage, which has a different phenotype than the other lineages. The plants in group B were all cultivated and had spines.

Group G was designated as an unclassified lineage. This plant group included plants from different locations across Japan (Table 1), including Adachi, Akita, Agamachi,

160 Ayabe, Kami, Kobe, Nagahama, Nangoku, Noto, Toyooka, and Tsuruoka. The plants in

161 group G were both cultivated and wild (Table 1).

Tight cluster and additional groups revealed by cluster analysis

Cluster analysis was performed using a pairwise distance matrix. The cluster analysis

identified the same seven groups that were identified in the PCA analysis and showed

that plants 42, 43, 44, 45, 46, 47, 48, 49, 50, and 51 formed tight cluster.

In addition, cluster analysis identified an eighth group (Figure 3), named group H.

Group H (Figure 3) contained the plants from Kochi Prefecture (Figure 2, Table 1). This

was named "unforeknown Kochi group". The plants in this group were spiny. Plants 1

and 87 were cultivated, and plant 2 was wild. These plants are used as ingredients to make

rice cakes in this region (Table 1). The other plant (number 92) from Kochi Prefecture

did not belong to group H.

Regional groups and their exceptions elucidated through phylogenetic analyses

To further establish the above classifications, we performed phylogenetic analyses

using the maximum likelihood (ML) method (Figure 4) and a method based on the

coalescent model (Figure 5). Both phylogenetic trees classified plants into eight groups.

The phylogenetic trees (Figures 4 and 5) also showed a tight cluster in group A,

containing plants 42, 43, 44, 45, 46, 47, 48, 49, 50, and 51, present as a subclade within

clade A with a high bootstrap value. The branch lengths of the ML tree (Figure 4)

indicated that these nine plants had a high degree of genetic similarity.

There were some differences in these eight groups between the two phylogenetic trees.

Two trees separated plant 14 from the other plants in group E. The ML tree separated

plant 13 from other plants in group D. Indeed, in PCA, these two plants were closer to the

plants in the other groups (Figure 1a and 1b). The discrepancies between the two

phylogenetic analyses, including these ones, are related to the low bootstrap values.

The cluster analysis and both phylogenetic analyses showed that group G had two

regional groups in common: (1) plants 78 and 79, and (2) plants 81, 82, and 83. Plants 78

and 79 in the first group were from Agamachi, Niigata Prefecture, and were wild spineless

plants. In the second group, plant 81 was from Yamagata Prefecture, plant 82 was from

Akita Prefecture, and plant 83 was from Tsuruoka, Yamagata Prefecture, which are

adjacent to each other. However, plant 76 from Tsuruoka, Yamagata Prefecture, did not

belong to this regional group.

In both phylogenetic trees, group G plants 88, 92, and 93 were isolated from the other

group G plants. In group G in the coalescent tree (Figure 5), plants 77 and 80 were

separated from other plants in the same group. This observation suggests that group G

may not be a single isolated group. Thus, the grouping of group G is ambiguous, probably

because plants in various locations belonged to group G.

Admixture history of the eight groups and the tight cluster within group A

Admixture analysis was used to estimate ancestral history and calculated cross-

validation (CV) errors to estimate possible values of K (number of ancestral populations)

= 1–12 (Supplementary Figure 1). $K = 2$ was the most likely value because the error value

of the cross-validation was minimal. We presented the results of the admixture analysis

from $K = 2$ to $K = 7$ (Figure 6).

In the case of $K = 2$ (Figure 6), except for plant 42, the tight cluster within group A

(plants 43, 44, 45, 46, 47, 48, 49, 50, and 51) formed a single population, shown in yellow.

The red colour corresponds to plants other than those in group A. The remaining group A

plants were admixed groups of the two populations. Thus, at $K = 2$, group A members

were clearly separated from the other plants.

The analysis of $K = 3–7$ separated the six groups A, B, C, D, E, and H (Figure 6).

However, the analysis did not separate group F, which is consistent with the results of

PCA, which also did not clearly separate group F from other plants.

The admixture analysis also showed that the grouping of group G was ambiguous, and

group G may not be a single isolated group. Reflecting this observation, the pattern of

colours in group G was not consistent across several K values. Thus, the plants in group

G plants may be admixtures of a variety of groups.

Conservation of heterozygosity

Asexual reproduction was confirmed by checking for heterozygosity conservation

using pairwise alignments. Plants 8 and 9 of group F were grafted from the same tree

(Table 1). Between these two plants, 2288 of 3699 sites were conserved heterozygous

sites (61.9%) (Supplementary Fig. 2). Similarly, plants 33 and 34 of group C were grafted

from the same tree (Table 1). Between these two plants, 1947 of the 3097 sites were

conserved heterozygous sites (62.9%) (Supplementary Fig. 3). Grafting is a common

method of propagating the Budou lineage, namely, group B plants. Between plants 3 and

4 of group B, 2198 of 3879 sites were conserved heterozygous sites (56.7%)

(Supplementary Fig. 4). Thus, this degree of heterozygosity conservation was found

among asexually reproduced plants when analysing the data using *de novo* mapping.

Between plants 40 and 41 of group A, growing nearby, 2140 of 3414 sites were

conserved heterozygous sites (62.7%) (Supplementary Fig. 5); between plants 38 and 39

of group A, growing nearby, 1700 of 3502 sites were conserved heterozygous sites

(48.5%) (Supplementary Fig. 6). Thus, plant pairs "40 and 41" and "38 and 39" were

234 identified as asexually propagated plants, although no records have been kept regarding

235 their propagation. Based on the analysis of these five pairs, we observed a conservation

236 of heterozygosity of approximately 50% or above between asexually propagated plants

237 when the RAD-Seq data were analysed *de novo*.

238  We examined the conservation of heterozygosity for the tight cluster in group A (plants

239 42, 43, 44, 45, 46, 47, 48, 49, 50, and 51). The degree of conservation ranged from 50.4%

240 to 68.9% (Supplementary Fig. 7, Supplementary Table 3). By contrast, the plants from

241 the tight clusters within group A shared lower levels of conservation (from 29.5% to

242 36.5%) with the other group A plants (plants, 35, 38, and 40) (Supplementary Fig. 7,

243 Supplementary Table 3). Therefore, the members of the tight cluster in group A were

244 asexually propagated plants derived from the same tree.

246 **Statistical interpretations of the classifications**

247  In our statistical analysis (Table 2 and 3, Supplementary Table 4, 5, and 6), we

248 removed the plants (13 and 14) for which we obtained uncertain results and the plants in

249 group G that caused the ambiguity. Although groups C and F belonged to the Arima

250 lineages, the Fst value between them was high (0.110, Table 2). Thus, the difference in

251 the pairwise Fst values between these two groups may indicate two places of origin.

252  Similarly, the pairwise Fst value between groups A and D of the Asakura lineage was

253 high (0.0803, Table 2). The reason for the higher Fst value may be the presence of a tight

254 cluster in group A. Therefore, we performed a second statistical analysis by modifying

255 the grouping. In the new grouping, we designated the tight cluster in group A as A2. We

256 denoted the plants that remained in group A as A1. As expected, the pairwise Fst values

257 were high between group A2 and each of the other groups (Table 3). Table 3 shows that

group A1 was genetically closer to group D (0.0539). Thus, the high genetic distance

between groups A and D in Table 2 was due to the tight cluster within group A.

We performed tests for f3-statistics and f4-statistics (Supplementary Tables 5 and 6,

respectively). Negative values of f3-statistics with A1 as the outgroup (Supplementary

Table 5) showed that group A1 is an admixture population of group A2 and the remaining

plants. This result is in good agreement with the results of the admixture analysis (Figure

6). The f4-statistics test (Supplementary Table 6) also supports this result with high or

low Z-scores. As for source populations other than A2, groups C and E were more likely

than group D. Although groups C and F belonged to the Arima lineages, positive values

of f3-statistics with A2 as the outgroup showed that group C may have more gene flow

from groups E, D, and H than from group F.

Discussion

Our study classified *Zanthoxylum piperitum* into several groups based on DNA

sequences and geographic information. Each group showed interesting characteristics,

some of which may have been related to domestication. The domestication of

Zanthoxylum piperitum can be described in two steps. The first step is the cultivation of

wild plants in agricultural fields, in which case the wild and cultivated plants are

genetically similar. In the second step, the lineages with desirable traits spread from their

place of origin to the surrounding areas, in which case the genetic diversity of individuals

within the lineages should be low. Grafting is a means of spreading crops such as fruit

trees.

Group D plants reflect the first step of domestication, because the cultivated plants

probably originated in the Yabu Mountains. Group D comprised both wild plants in

mountainous areas and cultivated plants in agricultural fields (Table 1).

The spineless Asakura lineage (*Z. piperitum* (L.) DC forma *inerme* (Makino) Makino),

first described 90 years ago[11], may be placed in group D. This 90-year-old

documentation[11], based on a much older book, states that the origin of the Asakura lineage

was on a cliff near the Konryuji temple. Plant 90, belonging to group D, grows in a place

similar to the place of origin and is spineless. In addition, spineless wild plants 53, 54, 60,

and 69 were collected near the Konryuji temple.

Group A plants reflect the second domestication step because they include lineages

with desirable traits and appear to have been spread. Group A contained a tight cluster

comprising plants 42, 43, 44, 45, 46, 47, 48, 49, 50, and 51. This cluster showed minimal

genetic diversity (Supplementary Table 4). Furthermore, the amount of conservation of

heterozygosity showed that these plants were propagated by grafting (Supplementary Fig.

7, Supplementary Table 3). These plants were also found to be widely cultivated in many

cities (Yabu, Ayabe, Gojo, Kyotamba, Kobe, and Kami), indicating that they have been

spread. Plants 42 and 48 are highly productive trees in Yabu City, which supplies scions

for grafting. Thus, the plants in this tight cluster exhibited a variety of desirable traits.

There are two potential places of origin for group A: Yabu City, where the Asakura

district is located and another distant location. The following reasons may explain the

first possibility: (1) many of the group A plants grow in Yabu City (plants 35, 38, 39, 42,

46, and 48); (2) a 90-year-old document[11] describes the spineless plants originating from

Asakura District, Yabu City; and (3) all of the group A plants are spineless. Interestingly,

we collected plants 35, 38, and 39 near the Konryuji temple, a potential place of origin

for the Asakura lineage. Therefore, we do not exclude the possibility that the spineless

Asakura lineage described 90 years ago[11] may represent group A plants. However, plants

35, 38, and 39 were cultivated. Thus, a serious problem for the first possibility is the

absence of wild plants in group A in the mountains of Yabu City.

The second possibility is that the group A plants are from another distant location; that

is, the plants in the tight cluster of group A of unknown origin hybridised with the group

D plants to form the rest of the group A plants. This possibility is consistent with the

results of admixture analysis for $K = 2$. However, it is worth noting that the results of the

admixture analysis may have been artificial since the second step of domestication formed

the ancestral population containing the plants in the tight cluster. In addition, the f3-

statistics test results (Supplementary Table 5) do not necessarily indicate that the source

population of group A is group D. If the second possibility is true, the origin of the plants

in the tight cluster remains unknown, although the origin of the rest of the group A plants

can be explained by hybridisation.

In groups C and F of the Arima lineage, we observed the first domestication step based

on documentation. According to the records, the group F plants (plants 7, 8, and 9) were

transplanted from Mt. Inariyama to Asago city (Table 1). Group C plants (plants 29, 30,

31, 32, 33, and 34) were transplanted from Mt. Rokko to Asago and Kobe cities. The

other group F plants (plants 5, 6, 10, 11, and 12) should have the same origin. According

to the PCA results, the group F plants (plants 5, 6, 10, 11, and 12) were more closely

related to the plants in the other groups.

We did not find any records describing the origins of the group E plants. In addition,

we could not find wild representatives of this group in the mountains. If the group E plants

have adaptations to high altitudes, the mountainous areas near Takayama city may be

where group E originated. Spineless plants are present in group E (plants 16, 17, 18,19,

20, 21, 22, 23, 24, 25, and 26) and are genetically different from the plants in groups A

and D in Yabu City, where the Asakura district is located.

In addition to these observations, the results for group G also suggest that

domestication is underway in many areas. Ongoing domestication is associated with the

cultivation of wild lineages on nearby agricultural lands and gardens. Group G was

genetically different from the other plant groups undergoing domestication. Therefore,

these plants may have been domesticated at their respective locations.

Recent reports have stated that the domestication of other crop species, such as olives

in Morocco[33] and dragon fruits/pitaya in Mexico[34], is also ongoing. Both olives and

dragon fruits undergo two common steps to become domesticated: first, the *in situ*

conservation of wild plants, and second, the introduction of wild plants into agricultural

fields. These plants were selected as the desirable plants for cultivation. The ongoing

domestication of Japanese pepper is not entirely similar to that of olives and dragon fruit.

In the first step of olive domestication, farmers continue to disturb the natural habitat of

the olive when selecting wild populations[33]. For Japanese pepper, wild plants are

harvested from their natural habitats and transplanted into agricultural fields or gardens

while leaving other wild plants intact in the mountains. The second step of the

domestication of olives and dragon fruits occurs through the continuous introduction of

wild plants into agricultural fields to further select for preferred traits[33,34]. In Japanese

pepper, wild plants are not frequently reintroduced into cultivated plant sites.

On the other hand, the ongoing domestication of Japanese pepper is similar to that of

olives and dragon fruit in which domestication syndrome has been observed.

Domestication syndrome changes the genomic, physiological, and morphological

characteristics of cultivated plants compared to those of wild plants[35]. The genomic

characteristics of the plants in the tight cluster in group A were different from those of

the wild plants. Similarly, the morphological characteristics of the plants in group B were

different from those of wild plants.

Spineless plants are known as the Asakura lineage and are classified as a single

subspecies, *Z. piperitum* (L.) DC forma *inerme* (Makino) Makino[11]. However, it is

inappropriate to refer to the spineless plant as the Asakura lineage or *Z. piperitum* (L.)

DC forma *inerme* (Makino) Makino. One Asakura lineage group (group A) contained a

tight cluster of spineless plants of unknown origin. Although the second Asakura lineage

group (group D) had a plant growing in a recorded place of origin[11], it contained some

spiny plants. In addition, the spineless trait is not unique to the Asakura lineage, as some

of the plants in the Takahara lineage (group E) are also spineless. Some group G plants

were spineless. Importantly, these spineless plants were from many locations. Therefore,

our study does not support the notion that the spineless lineage is either from Asakura or

monophyletic. Our results indicate that the spineless lineage is not a single subspecies.

Some Fis values were negative (Supplementary Table 4). Notably, the tight cluster

plants in group A (group A2) had the highest negative Fis value. Therefore, we speculated

that some of these plants had been asexually reproduced by grafting, as asexual

reproduction may cause negative Fis values[36]. Pairwise alignments support this

possibility (Supplementary Fig. 7 and Supplementary Table 3).

In this study, we elucidated the genetic diversity of Japanese pepper, while tracing the

process of domestication. Our results demonstrate that the conservation of plant genetic

resources is an important agricultural practice. Several cultivated lineages of Japanese

pepper were found to be closely related to plants growing in the mountains, in contrast to

other crops. On the other hand, several cultivated lineages, groups A, B, and E, had no

close relatives growing in the mountains, which suggests that wild plants of this group no

longer exist. Therefore, there is a need to conserve both plants growing in the mountains

in situ and plants growing in agricultural fields or preservation facilities *ex situ*. The

results presented in this study will be useful for developing a plan for the conservation

and breeding of Japanese peppers.

Methods

Plant materials

Ninety-three *Zanthoxylum piperitum* samples collected from various parts of Japan were

used (Table 1). Photographs of the materials were taken, and the leaves were stored in a

freezer (−80°C). At each location, where possible, plants growing in the mountains and

agricultural fields were collected. Any lineages of Budou or Takahara that grew in the

mountains were not found. All methods involving plants were carried out in accordance

with relevant guidelines and regulations. Plants were collected with permission from their

owners.

DNA extraction and double-digest restriction site amplified DNA sequencing

(ddRAD-Seq)

DNA extraction and purification were performed using previously described methods[30].

The library for ddRAD-Seq was prepared based on the original protocol[27] with some

modifications[37]. The first restriction site near the primer binding site, which reads the

single-ended DNA sequence, was *Bgl*II. The second restriction site, adjacent to the

binding site for reading the index sequence, was *Eco*RI. The library was sequenced with

51-bp single-end reads at Macrogen (Seoul, Korea) on two lanes of an Illumina

HiSeq2000 (Illumina, San Diego, CA, USA).

Processing and quality control of ddRAD-Seq data

The Stacks package[38] was used to analyse the ddRAD-Seq reads. The

process_shortreads program of the Stacks package (version 2.4) was used to clean reads

using -c (clean data, remove any read with an uncalled base), -q (discard reads with low-

quality scores), and -r (rescue barcodes) options. The cutadapt program (version 2.6)[39]

removed adaptor sequences from reads using the -b (remove adapter anywhere, the

sequence of an adapter that may be ligated to the 5′ or 3′ end) and -e 0.1 (maximum

allowed error rate by keeping the default value 0.1 = 10%) options. Any low-quality bases

in the single-end reads were trimmed using the trimmomatic program (version 0.39)[40]

with the following settings LEADING:19 (cut bases off the start of a read, if below a

threshold quality), TRAILING:19 (cut bases off the end of a read, if below a threshold

quality), AVGQUAL:20 (drop the read if the average quality is below a specified level),

MINLEN:51 (drop the read if it is below a specified length), and

SLIDINGWINDOW:20:20 (perform sliding window trimming, clipping once the average

quality within the window falls below a threshold).

***De novo* mapping and variant calling**

The denovo_map.pl of the Stacks package (version 2.5), a wrapper script for ustacks,

cstacks, sstacks, tsv2bam, and gstacks, was used to map reads *de novo* without using a

reference genome. The options for denovo_map.pl were -M 4 (number of mismatches

allowed between stacks within individuals [for ustacks]), -n 4 (number of mismatches

allowed between stacks between individuals [for cstacks]), and -m 3 (number of identical

reads required to initiate a new putative allele [for ustacks]). Genotyping data were

created to specify all the samples under analysis by assigning one individual per

population. After performing denovo_map.pl, the populations program of the Stacks

package (version 2.5) created the vcf (variant call format)[41], plink[42], phylip, and treemix

files using the -R 0.5 (minimum percentage of individuals across populations required to

process a locus), --write-single-snp (restrict data analysis to only the first SNP per locus),

--min-maf 0.05 (minimum minor allele count required to process a SNP), --vcf, --plink, -

-phylip-var-all --treemix options. The populations program was also used to create

pairwise alignments between the two individuals. In this case, the -R option was set to

one.

Principal component analysis (PCA) and cluster analysis

Principal component analysis was performed based on the vcf file generated by the

populations program. The SNPRelate program[43] in the R software environment (version

3.6.2)[44] converted the vcf file to a gds (genomic data structure) file, drew the PCA

diagrams, and calculated the contribution ratios for each principal component. This step

used only bi-allelic loci. The SNPRelate program plotted the dendrogram by considering

the identity by state (IBS) pairwise distances. Images of the results were generated using

the basic functions of the R software environment.

Admixture analysis

The plink program (plink 2, version 1.90p)[42] was used to create the input files for the

admixture program. The admixture program (version 1.3)[45] was used to determine the

admixture history and the cross-validation (CV) error for the hypothetical runs from K

(number of ancestral populations) = 1–12. The CV error plot was drawn using the

received log data to determine the optimal K value. R software (version 3.6.2) was used

to draw admixture plots using the Q estimate files created by the admixture programs.

Images of the results were generated using the basic functions of the R software

environment.

Maximum likelihood phylogenetic tree analysis

ModelTest-NG was used to select the model[46]. The maximum likelihood (ML)

phylogenetic tree was created using the raxmlHPC-PTHREADS-SSE3 program (version

8.2.12)[47]. The phylip file created by the populations program was used as the input file.

The parameters for raxml were -f a (rapid bootstrap analysis and search for the best-

scoring ML tree in one program run), -x 12345 (an integer number [random seed] and

turn on rapid bootstrapping), -p 12345 (a random number seed for parsimony inferences),

-N 1000 (bootstrap value), and -m GTRGAMMAX (model of binary [morphological],

nucleotide, multi-state, or amino acid substitution). Group H was used as the root. Image

of the phylogenetic tree was generated using Dendroscope (version 3.6.3)[48] and TreeView

X (version 0.5.0, https://treeview-x.en.softonic.com/).

Phylogenetic tree analysis under the coalescent model

Phylogenetic tree analysis was performed using SVDquartets[49] integrated into PAUP

(version 4.0a) software[50]. The input file for the PAUP was created by converting a phylip

file containing all sites to the nexus file format. The parameters used for SVDquartets

were 'quartet evaluation', 'evaluate all possible quartets', 'tree inference', 'select trees'

'using QFM quartet assembly', 'tree model', 'multi-species coalescent', 'write quartets

file' 'QMC format', 'handling of ambiguities' and 'distribute'. The number of bootstrap

analyses was 1,000 replicates. Group A was used as the root. Image of the phylogenetic

tree was generated using FigTree (version 1.4.4, http://tree.bio.ed.ac.uk/).

Statistical analysis

Groups separated by PCA and cluster analysis were assigned as separate populations

in the population map data required for the Stacks package. The denovo_map.pl was re-

performed after modifying the data for the population map. The populations program was

run with the options -R 0.5, --write-single-snp, --min-maf 0.05, --fstats (SNP and

haplotype-based F statistics), and --Fst_correction (a correction to be applied to Fst values

with p-value [default p-value to keep an Fst measurement: 0.05]). The program treemix

(version 1.12)[51] inferred gene flow using the f3- and f4-statistics with the option -k 500

(standard errors in blocks of 500 SNPs).

Acknowledgments

We are grateful to those who helped us collect samples from various locations. We thank

Ms. Yuki Sano, Mr. Atushi Shirakawa, and Mr. Kazuma Kubo for preparing the DNA

samples. This work was supported by the Urakami Foundation for Food and Food Culture

Promotion.

Author contributions

N.F., K.Y., F.H., and Y.N. designed the study. N.F., H.M., N.I., and Y.N. collected the

samples. Y.N. performed DNA extraction. M.D.G.P.P. and Y.N. performed the

bioinformatic analysis. A.J.N. performed the RAD-Seq analysis. M.D.G.P.P., N.F., K.Y.,

F.H.H.M., N.I., and Y.N. analysed the data. M.D.G.P.P. and Y.N. wrote the manuscript.

Competing interests

The authors declare no competing interests.

Data availability

Sequences are available at the DNA Data Bank of Japan Sequence Read Archive

(https://www.ddbj.nig.ac.jp/dra/index-e.html; Accession no. DRA011046).

References

1.   Ayurvedic Medicinal Plants of Sri Lanka, *Ayurvedic plants of Sri Lanka: details*

508 http://www.instituteofayurveda.org/plants (2020).

2. Zhou, J., Xie, G. & Yan, X. *Encyclopedia of Traditional Chinese Medicines - Molecular Structures, Pharmacological Activities, Natural Sources and Applications* (Berlin: Springer, 2011).

3. Donald, G. R., Fernandes, P. D. & Boylan, F. Antinociceptive Activity of *Zanthoxylum piperitum* DC. Essential Oil. *Evid. Based Complement Alternat. Med.* 2016, 3840398 (2016).

4. Gwon, S. Y., Ahn, J. Y., Kim, T. W. & Ha, T. Y. *Zanthoxylum piperitum* DC ethanol extract suppresses fat accumulation in adipocytes and high fat diet-induced obese mice by regulating adipogenesis. *J. Nutr. Sci. Vitaminol.* **58**, 393–401 (2012).

5. Ha, S. Y. *et al.* Antiviral effect of flavonol glycosides isolated from the leaf of *Zanthoxylum piperitum* on influenza virus. *J. Microbiol.* **52**, 340–344 (2014).

6. Jang, K. H. *et al.* New polyunsaturated fatty acid amides isolated from the seeds of *Zanthoxylum piperitum*. *Arch. Pharm. Res.* **31**, 569–572 (2008).

7. Kim, M. H., Lee, H. J., Park, J. C., Hong, J. & Yang, W. M. *Zanthoxylum piperitum* reversed alveolar bone loss of periodontitis via regulation of bone remodeling-related factors. *J. Ethnopharmacol.* **195**, 137–142 (2017).

8. Kusuda, M. *et al.* Polyphenolic constituent structures of *Zanthoxylum piperitum* fruit and the antibacterial effects of its polymeric procyanidin on methicillin-resistant *Staphylococcus aureus*. *Biosci. Biotechnol. Biochem.* **70**, 1423–1431 (2006).

9. Nozaki, R. *et al. Zanthoxylum* fruit extract from Japanese pepper promotes autophagic cell death in cancer cells. *Oncotarget.* **7**, 70437–70446 (2016).

10. Song, S., Oh, S. & Lim, K. T. Bioactivity of proteins isolated from *Lactobacillus plantarum* L67 treated with *Zanthoxylum piperitum* DC glycoprotein. *Lett. Appl. Microbiol.* **60**, 597–604 (2015).

11. Makino, T. The origin of Asakura sanshō. *J. Jpn. Bot.* **5**, 378–380 (1928).

12. Lee, J., *et al.* The complete chloroplast genome sequence of *Zanthoxylum piperitum*.

Mitochondrial DNA A **27**, 3525–3526 (2016).

13. Sun, Y. L., Park, W. G., Kwon, O. W. & Hong, S. K. The internal transcribed spacer

rDNA specific markers for identification of *Zanthoxylum piperitum Afr. J.*

Biotechnol. **9**, 6027–6039 (2010).

14. Kim, Y., Shin, J., Cho, S. S., Hwang, Y. P., & Choi, C. Development and application

of InDel markers for authentication of the Korean herbs *Zanthoxylum schinifolium*

and *Zanthoxylum piperitum. Foods*, **8**, 658 (2019).

15. Feng, S., *et al.* Phylogenetic relationships among cultivated *Zanthoxylum* species in

China based on cpDNA markers. *Tree Genet. Genomes* **12**, 45 (2016).

16. Appelhans, M. S., Reichelt, N., Groppo, M., Paetzold, C., & Wen, J. Phylogeny and

biogeography of the pantropical genus *Zanthoxylum* and its closest relatives in the

proto-Rutaceae group (Rutaceae). *Mol. Phylogenetics Evol.* **126**, 31–44 (2018).

17. Liu, Y., & Wei, A. The complete chloroplast genome sequence of an economically

important plant, *Zanthoxylum bungeanum* (Rutaceae). *Conserv. Genet. Resour.* **9**,

25–27 (2017).

18. Zhao, *et al.* Comparative analyses of chloroplast genomes from 14 *Zanthoxylum*

species: Identification of variable DNA markers and phylogenetic relationships

within the genus. *Front. Plant Sci.* **11**, 2226 (2021).

19. Lee, H. J., *et al.* (2017). Authentication of *Zanthoxylum* species based on integrated

analysis of complete chloroplast genome sequences and metabolite profiles. *J. Agric.*

Food Chem. **65**, 10350–10359 (2017).

20. Li, H. L., Zhou, N., & Guo, D. Q. The complete chloroplast genome of *Zanthoxylum*

acanthopodium DC.(Rutaceae) and its phylogenetic analysis. *Mitochondrial DNA B*

5, 3636–13637 (2020).

21. Gupta, D. Das & Mandi, S. Sen. Species specific AFLP markers for authentication

of *Zanthoxylum acanthopodium* & *Zanthoxylum oxyphyllum*. *J. Med. Plants Stud.* **1**,

1–9 (2013).

22. Feng, S. *et al.* Genetic diversity and relationships of wild and cultivated *Zanthoxylum*

germplasms based on sequence-related amplified polymorphism (SRAP) markers.

Genet. Resour. Crop Evol. **62**, 1193–1204 (2015).

23. Hu, Y. *et al.* Genetic structure of cultivated *Zanthoxylum* species investigated with

SSR markers. *Tree Genet. Genomes* **14**, 89 (2018).

24. Henry, R. J. Next-generation sequencing for understanding and accelerating crop

domestication. *Brief. Funct. Genomics* **11**, 51–56 (2012).

25. Grover, A. & Sharma, P. C. Development and use of molecular markers: past and

present. *Crit. Rev. Biotechnol.* **36**, 290–302 (2016).

26. Baird, N. A. *et al*. Rapid SNP discovery and genetic mapping using sequenced RAD

markers. *PLOS ONE* **3**, e3376 (2008).

27. Peterson, B. K., Weber, J. N., Kay, E. H., Fisher, H. S. & Hoekstra, H. E. Double

digest RADseq: an inexpensive method for de novo SNP discovery and genotyping

in model and non-model species. *PLOS ONE* **7**, e37135 (2012).

28. Penjor, T. *et al.* RAD-Seq analysis of typical and minor *Citrus* accessions, including

Bhutanese varieties. *Breed. Sci.* **66**, 797–807 (2016).

29. Nagano, Y. *et al.* Phylogenetic relationships of Aurantioideae (Rutaceae) based on

RAD-Seq. *Tree Genet. Genomes* **14**, 6 (2018).

30. Penjor, T., Mimura, T., Matsumoto, R., Yamamoto, M. & Nagano, Y.

Characterization of limes (*Citrus aurantifolia*) grown in Bhutan and Indonesia using

high-throughput sequencing. *Sci. Rep.* **4**, 4853 (2014).

31. Fukuda, S. *et al*. Construction of a high-density linkage map for bronze loquat using

RAD-Seq. *Sci. Hortic.* **251**, 59–64 (2019).

32. Kato, D. *et al*. Evaluation of the population structure and phylogeography of the

Japanese Genji firefly, *Luciola cruciata*, at the nuclear DNA level using RAD-Seq

analysis. *Sci. Rep.* **10**, 1533 (2020).

33. Aumeeruddy-Thomas, Y., Moukhli, A., Haouane, H. & Khadari, B. Ongoing

domestication and diversification in grafted olive–oleaster agroecosystems in

northern Morocco. *Reg. Environ. Change* **17**, 1315–1328 (2017).

34. Parra, F. *et al*. Evolution under domestication: ongoing artificial selection and

divergence of wild and managed *Stenocereus pruinosus* (Cactaceae) populations in

the Tehuacán Valley, Mexico. *Ann. Bot.* **106**, 483–496 (2010).

35. Gepts, P. *Domestication of Plants in Encyclopedia of Agriculture and Food Systems*

(ed. Van Alfen, N. K.) 474–486 (Academic Press, 2014).

36. Kusuma, Y. W. C. *et al*. Seedling selection using molecular approach for *ex situ*

conservation of critically endangered tree species (*Vatica bantamensis* (Hassk.)

Benth. & Hook. ex Miq.) in Java, Indonesia. *Trop. Conserv. Sci.* **12**, 1–12 (2019).

37. Sakaguchi, S. *et al*. High-throughput linkage mapping of Australian white cypress

pine (*Callitris glaucophylla*) and map transferability to related species. *Tree Genet.*

Genomes **11**, 121 (2015).

38. Rochette, N. C., Rivera-Colón, A. G. & Catchen, J. M. Stacks 2: analytical methods

for paired-end sequencing improve RADseq-based population genomics. *Mol. Ecol.*

28, 4737–4754 (2019).

39. Martin, M. Cutadapt removes adapter sequences from high-throughput sequencing

reads. *EMBnet J.* **17**, 10–12 (2011).

40. Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina

sequence data. *Bioinformatics* **30**, 2114–2120 (2014).

41. Danecek, P. *et al.* The variant call format and VCFtools. *Bioinformatics* **27**, 2156–

2158 (2011).

42. Purcell, S. *et al.* PLINK: a tool set for whole-genome association and population-

based linkage analyses. *Am. J. Hum. Genet.* **81**, 559–575 (2007).

43. Zheng, X. *et al.* A high-performance computing toolset for relatedness and principal

component analysis of SNP data. *Bioinformatics* **28**, 3326–3328 (2012).

44. R Core Team. *R: A Language and Environment for Statistical Computing. R*

Foundation for Statistical Computing Vienna, Austria. https://www.R-project.org/

(2019).

45. Alexander, D. H., Novembre, J. & Lange, K. Fast model-based estimation of ancestry

in unrelated individuals. *Genome Res.* **19**, 1655–1664 (2009).

46. Darriba, D., *et al.* ModelTest-NG: a new and scalable tool for the selection of DNA

and protein evolutionary models. *Mol. Biol. Evol.* **37**, 291–294 (2020).

47. Stamatakis, A. RAxML version 8: a tool for phylogenetic analysis and post-analysis

of large phylogenies. *Bioinformatics* **30**, 1312–1313 (2014).

48. Huson, D. H. & Scornavacca, C. Dendroscope 3: an interactive tool for rooted

phylogenetic trees and networks. *Syst. Biol.* **61**, 1061–1067 (2012).

49. Chifman, J. & Kubatko, L. Quartet inference from SNP data under the coalescent

model. *Bioinformatics* **30**, 3317–3324 (2014).

50. Swofford, D. L. PAUP* phylogenetic analysis using parsimony (* and other

methods) version 4. (Sinauer, Sunderland, MA, 2003).

51. Pickrell, J. K. & Pritchard, J. K. Inference of Population Splits and Mixtures from

Genome-Wide Allele Frequency Data. *PLOS Genet*. **8**, e1002967 (2012).

Figure 1. Principal component (PC) analysis of *Zanthoxylum* accessions used in the study. The six major component data are shown by the five two-dimensional data sets: (**a**) first and second PCA axes, (**b**) first and third PCA axes, (**c**) first and fourth PCA axes, (**d**) first and fifth PCA axes, and (**e**) first and sixth PCA axes. Group A of the Asakura lineage is shown in blue; group B of the Budou lineage is shown in brown; group C of the Arima lineage is shown in orange; group D of the Asakura lineage is shown in violet; group E of the Takahara lineage is shown in maroon; group F of the Arima lineage is shown in green; group G is shown in violet. The letters indicating the sample number are also shown in the same colour. The contribution rate of each principal component is shown in parentheses. Figure was generated using R software (version 3.6.2)[44].

Figure 2. Sampling points and locations of the lineages. The colour scheme is the same as in Figure 1.

Figure 3. Cluster analysis of *Zanthoxylum* accessions. The new group, H group, is shown in light blue. The colour scheme is the same as in Figure 1, with an additional colour. Figure was generated using R software (version 3.6.2)[44].

Figure 4. Maximum likelihood phylogenetic analysis of *Zanthoxylum* accessions. The colour scheme is the same as in Figure 3. The numbers at the nodes indicate bootstrap values (% over 1,000 replicates). The nodes with ≥50% bootstrap value are labeled. Figure was generated using Dendroscope (version 3.6.3)[48] and TreeView X (version 0.5.0, https://treeview-x.en.softonic.com/)

Figure 5. Coalescent model phylogenetic tree of *Zanthoxylum* accessions. The colour

scheme is the same as in Figure 3. The numbers at the nodes indicate bootstrap values

(% over 1,000 replicates). The nodes with ≥50% bootstrap value are labeled. Figure was

generated using FigTree (version 1.4.4, http://tree.bio.ed.ac.uk/).

Figure 6. *K* = 2-7 admixture plots in admixture analysis of *Zanthoxylum* accessions used

in this study. The horizontal axis shows the group names and the sample numbers. The

colour scheme for the group names and the sample numbers is the same as in Figure 3.

The colour for the genetic cluster is different from that used to denote the group names

and sample numbers. Figure was generated using R software (version 3.6.2)[44].

Tables

Table 1. Information on *Zanthoxylum piperitum* accessions used in the study.

Plant Number	Prefecture	City/Town	Spiny/Spineless	Information	Group	Lineage
1	Kochi	Sakawa	Spiny	Agricultural field, cultivated	H	Unforeknown Kochi group
2	Kochi	Nangoku	Spiny	Mountain, wild tree	H	Unforeknown Kochi group
3	Wakayama	Kainan	Spiny	Agricultural field, cultivated, Nagamine agricultural cooperative	B	Budou
4	Wakayama	Kainan	Spiny	Agricultural field, cultivated, Nagamine agricultural cooperative	B	Budou
5	Hyogo	Kobe	Spiny	Garden, cultivated	F	Arima
6	Hyogo	Kobe	Spiny	Garden, cultivated	F	Arima

7	Hyogo	Asago	Spiny	Agricultural field in fruit tree research institute for preservation purpose, cultivated, originally transplanted from Mt. Inariyama, agricultural field	F	Arima
8	Hyogo	Asago	Spiny	Agricultural field in fruit tree research institute for preservation purpose, cultivated, originally transplanted from Mt. Inariyama, agricultural field, grafted from the same tree as plant 9	F	Arima
9	Hyogo	Asago	Spiny	Agricultural field in fruit tree research institute for preservation purpose, cultivated, originally transplanted from Mt. Inariyama, agricultural field, grafted from the same tree as plant 8	F	Arima
10	Hyogo	Kobe	Spiny	Garden, cultivated	F	Arima
11	Hyogo	Kobe	Spiny	Garden, cultivated	F	Arima
12	Hyogo	Kobe	Spiny	Garden, cultivated	F	Arima
13	Hyogo	Yabu	Spineless	Garden, cultivated	D	Asakura
14	Gifu	Takayama	Spiny	Garden, cultivated	E	Takahara
15	Gifu	Takayama	Spiny	Agricultural field, cultivated	E	Takahara

16	Gifu	Takayama	Spineless	Agricultural field, cultivated	E	Takahara
17	Gifu	Takayama	Spineless	Agricultural field, cultivated	E	Takahara
18	Gifu	Takayama	Spineless	Agricultural field, cultivated	E	Takahara
19	Gifu	Takayama	Spineless	Agricultural field, cultivated	E	Takahara
20	Gifu	Takayama	Spineless	Garden, cultivated, used as mother tree for grafting	E	Takahara
21	Gifu	Takayama	Spineless	Agricultural field, cultivated, used as mother tree for grafting	E	Takahara
22	Gifu	Takayama	Spineless	Agricultural field, cultivated	E	Takahara
23	Gifu	Takayama	Spineless	Agricultural field, cultivated	E	Takahara
24	Gifu	Takayama	Spineless	Garden, cultivated	E	Takahara
25	Gifu	Takayama	Spineless	Agricultural field, cultivated, about 30 years old	E	Takahara
26	Gifu	Takayama	Spineless	Agricultural field, cultivated, about 25-26 years old	E	Takahara
27	Gifu	Takayama	Spiny	Agricultural field, cultivated	E	Takahara
28	Gifu	Takayama	Spiny	Agricultural field, cultivated	E	Takahara
29	Hyogo	Kobe	Spiny	Garden, cultivated, originally transplanted from Mt. Rokko	C	Arima
30	Hyogo	Kobe	Spiny	Garden, cultivated, originally transplanted from Mt. Rokko	C	Arima

31	Hyogo	Asago	Spiny	Agricultural field in fruit tree research institute for preservation purpose, cultivated, originally transplanted from Yubunedani of Mt. Rokko	C	Arima
32	Hyogo	Asago	Spiny	Agricultural field in fruit tree research institute for preservation purpose, cultivated, originally transplanted from Yubunedani of Mt. Rokko	C	Arima
33	Hyogo	Asago	Spiny	Agricultural field in fruit tree research institute for preservation purpose, cultivated, originally transplanted from Yubunedani of Mt. Rokko, grafted from the same tree as plant 34	C	Arima
34	Hyogo	Asago	Spiny	Agricultural field in fruit tree research institute for preservation purpose, cultivated, originally transplanted from Yubunedani of Mt. Rokko, grafted from the same tree as plant 33	C	Arima
35	Hyogo	Yabu	Spineless	Agricultural field, cultivated, near the Konryuji temple	A	Asakura
36	Kyoto	Kyōtamba	Spineless	Agricultural field, cultivated	A	Asakura

37	Hyogo	Kobe	Spineless	Agricultural field, cultivated, purchased as nursery stock of high-yielding Asakura sanshō	A	Asakura
38	Hyogo	Yabu	Spineless	Agricultural field, cultivated, near the Konryuji temple	A	Asakura
39	Hyogo	Yabu	Spineless	Agricultural field, cultivated, near the Konryuji temple	A	Asakura
40	Kyoto	Kyōtamba	Spineless	Agricultural field, cultivated	A	Asakura
41	Kyoto	Kyōtamba	Spineless	Agricultural field, cultivated	A	Asakura
42	Hyogo	Yabu	Spineless	Agricultural field, cultivated, used as high-yielding mother tree for grafting	A	Asakura
43	Kyoto	Ayabe	Spineless	Agricultural field, cultivated	A	Asakura
44	Nara	Gojō	Spineless	Agricultural field, cultivated	A	Asakura
45	Kyoto	Kyōtamba	Spineless	Agricultural field, cultivated	A	Asakura
46	Hyogo	Yabu	Spineless	Agricultural field, cultivated	A	Asakura
47	Hyogo	Kobe	Spineless	Agricultural field, cultivated, purchased as nursery stock of Asakura sanshō	A	Asakura
48	Hyogo	Yabu	Spineless	Agricultural field, cultivated, used as mother tree for grafting	A	Asakura
49	Nara	Gojō	Spineless	Agricultural field, cultivated	A	Asakura
50	Nara	Gojō	Spineless	Agricultural field, cultivated	A	Asakura

51	Hyogo	Kami	Spineless	Agricultural field, cultivated	A	Asakura
52	Hyogo	Yabu	Spineless	Mountain, wild tree	D	Asakura
53	Hyogo	Yabu	Spineless	Mountain, wild tree, near the Konryuji temple	D	Asakura
54	Hyogo	Yabu	Spineless	Mountain, wild tree, near the Konryuji temple	D	Asakura
55	Hyogo	Yabu	Spiny	Agricultural field, cultivated, near the Konryuji temple	D	Asakura
56	Hyogo	Yabu	Spineless	Mountain, wild tree	D	Asakura
57	Hyogo	Yabu	Spineless	Mountain, wild tree	D	Asakura
58	Hyogo	Yabu	Spineless in 2017, spiny in 2020	Agricultural field, cultivated, 100 years old tree	D	Asakura
59	Hyogo	Yabu	Spineless	Mountain, wild tree	D	Asakura
60	Hyogo	Yabu	Spineless	Mountain, wild tree, near the Konryuji temple	D	Asakura
61	Hyogo	Yabu	Spiny	Possible abandoned tree near an abandoned house	D	Asakura
62	Hyogo	Yabu	Spineless	Mountain, wild tree	D	Asakura
63	Hyogo	Yabu	Spineless	Mountain, wild tree	D	Asakura
64	Hyogo	Yabu	Spiny	Mountain, wild tree	D	Asakura
65	Hyogo	Yabu	Spineless	Mountain, wild tree	D	Asakura

66	Hyogo	Yabu	Spiny	Mountain, wild tree, near the Konryuji temple	D	Asakura
67	Hyogo	Yabu	Spiny	Mountain, wild tree	D	Asakura
68	Hyogo	Yabu	Spiny	Mountain, wild tree	D	Asakura
69	Hyogo	Yabu	Spineless	Mountain, wild tree	D	Asakura
70	Hyogo	Yabu	Spineless	Mountain, wild tree	D	Asakura
71	Hyogo	Yabu	Spineless	Mountain, wild tree	D	Asakura
72	Hyogo	Yabu	Spiny	Mountain, wild tree	D	Asakura
73	Hyogo	Yabu	Spiny	Mountain, wild tree	D	Asakura
74	Hyogo	Kami	Spineless	Garden, spontaneously grown tree	D	Asakura
75	Ishikawa	Noto	Spineless	Mountain, wild tree	G	Unclassified
76	Yamagata	Tsuruoka	Spiny	Garden, spontaneously grown tree	G	Unclassified
77	Shiga	Nagahama	Spineless	Agricultural field, cultivated	G	Unclassified
78	Niigata	Agamachi	Spineless	Mountain, wild tree	G	Unclassified
79	Niigata	Agamachi	Spineless	Mountain, wild tree	G	Unclassified
80	Hyogo	Kami	Spineless	Agricultural field, cultivated	G	Unclassified
81	Yamagata	Tsuruoka	Spineless	Garden, spontaneous grown tree, growing adjacent to plant 83	G	Unclassified

82	Akita	Akita	Spiny	Transplanted from mountain, wild tree	G	Unclassified
83	Yamagata	Tsuruoka	Spineless	Garden, spontaneous grown tree, growing adjacent to plant 81	G	Unclassified
84	Tokyo	Adachi	Spiny	Garden, cultivated from the seed of an old tree	G	Unclassified
85	Ishikawa	Noto	Spineless	Mountain, wild tree	G	Unclassified
86	Ishikawa	Noto	Spineless	Mountain, wild tree	G	Unclassified
87	Kochi	Sakawa	Spiny	Agricultural field, cultivated	H	Unforeknown Kochi group
88	Kyoto	Ayabe	Spineless	Agricultural field, cultivated	G	Unclassified
89	Hyogo	Yabu	Spineless	Garden, cultivated, near the Konryuji temple	D	Asakura
90	Hyogo	Yabu	Spineless	Mountain, wild tree, near the Konryuji temple, located in the recorded place of the origin of Asakura lineage	D	Asakura
91	Hyogo	Toyooka	Spiny	Mountain, wild tree	G	Unclassified
92	Kochi	Nangoku	Spiny	Agricultural field, cultivated	G	Unclassified
93	Hyogo	Kobe	Spineless	Garden, spontaneously grown tree	G	Unclassified

Table 2. Fst values for each pair of groups for A, B, C, D, E, F, H.

Group	B	F	E	C	A	D
H	0.307	0.120	0.0847	0.165	0.1080	0.0490
B		0.130	0.0950	0.191	0.125	0.0529
F			0.0689	0.110	0.103	0.0457
E				0.0874	0.0994	0.0464
C					0.126	0.0572
A						0.0803

Table 3. Fst values for each pair of groups for A1, A2, B, C, D, E, F, H.

Group	B	F	E	C	A1	A2	D
H	0.268	0.109	0.0760	0.156	0.132	0.196	0.0464
B		0.127	0.0950	0.189	0.159	0.226	0.0531
F			0.0697	0.111	0.100	0.154	0.0454
E				0.0878	0.0807	0.132	0.0464
C					0.134	0.195	0.0582
A1						0.0990	0.0539
A2							0.0993

Ancestry

Plant Number

H B F D E C A D G H G D G

K=7
K=6
K=5
K=4
K=3
K=2