Title: Multiple adjoining word- and face-selective regions in ventral temporal cortex exhibit

distinct dynamics

Abbreviated title: Dynamic word- and face-selective networks

Authors: Matthew J. Boring*[1,2,3], Edward H. Silson[4,5], Michael J. Ward[3], R. Mark

Richardson[3,6,7], Julie A. Fiez[1,2,8], Chris I. Baker[4], Avniel Singh Ghuman[1,2,3,8]

1. Center for Neuroscience at the University of Pittsburgh, University of Pittsburgh, A210

Langley Hall, Pittsburgh, PA 15260, USA.

2. Center for the Neural Basis of Cognition, 4400 Fifth Avenue, Pittsburgh, PA 15213.

3. Department of Neurological Surgery, University of Pittsburgh Medical Center, 200 Lothrop

Street, Pittsburgh, PA 15213, USA.

4. National Institute of Mental Health, National Institutes of Health, Magnuson Clinical Center,

Bethesda, MD 20814 USA;

5. School of Philosophy, Psychology and Language Sciences, University of Edinburgh, 7

George Square, Edinburgh EH8 9JZ, UK;

6. Department of Neurosurgery, Massachusetts General Hospital, 55 Fruit Street, Boston, MA

02144, USA.

7. Harvard Medical School, 25 Shattuck St., Boston, MA 02115, USA.

8. Department of Psychology, University of Pittsburgh, 210 South Bouquet St, Pittsburgh, PA

15260, USA.

Corresponding author email address: mjb200[at]pitt.edu

Number of pages: 52

Number of Figures: 8, **Number of tables:** 1

Number of words (total: 12,448, abstract: 247, introduction: 647, discussion: 1500)

Conflict of interest statement: The authors declare no competing financial interests.

Acknowledgements: We would like to thank the patients, their families, and the clinical staff at

the Epilepsy Monitoring Unit at the University of Pittsburgh Medical Center, without whom this

study would not be possible. We would also like to thank Marlene Behrmann, David Plaut,

Michael Tarr, and the anonymous reviewers for helpful suggestions regarding the analysis and

interpretations of the intracranial EEG results; Sean Walls, Ellyanna Kessler, Roma Konecky,

and Ashley Whiteman for assistance in intracranial EEG data collection. This study was

supported by the National Institutes of Health under R01MH107797 and R21EY030297 and

National Science Foundation under 1734907 to A.S.G., National Institutes of Health under

T32NS007433-20 to M.J.B., and National Institutes of Health under ZIAMH002909 to C.I.B.

and E.H.S.

Abstract

The map of category-selectivity in human ventral temporal cortex (VTC) provides organizational

constraints to models of object recognition. One important principle is lateral-medial response

biases to stimuli that are typically viewed in the center or periphery of the visual field. However,

little is known about the relative temporal dynamics and location of regions that respond

preferentially to stimulus classes that are centrally viewed, like the face- and word-processing

networks. Here, word- and face-selective regions within VTC were mapped using intracranial

recordings from 36 patients. Partially overlapping, but also anatomically dissociable patches of

face- and word-selectivity were found in VTC. In addition to canonical word-selective regions

along the left posterior occipitotemporal sulcus, selectivity was also located medial and anterior

to face-selective regions on the fusiform gyrus at the group level and within individual subjects.

These regions were replicated using 7 Tesla fMRI in healthy subjects. Left hemisphere word-

selective regions preceded right hemisphere responses by 125 ms, potentially reflecting the left

hemisphere bias for language; with no hemispheric difference in face-selective response latency.

Word-selective regions along the posterior fusiform responded first, then spread medially and

laterally, then anteriorly. Face-selective responses were first seen in posterior fusiform regions

bilaterally, then proceeded anteriorly from there. For both words and faces, the relative delay

between regions was longer than would be predicted by purely feedforward models of visual

processing. The distinct time-courses of responses across these regions, and between

hemispheres, suggest a complex and dynamic functional circuit supports face and word

perception.

Significance Statement: Representations of visual objects in the human brain have been shown

to be organized by several principles, including whether those objects tend to be viewed centrally

or peripherally in the visual field. However, it remains unclear how regions that process objects

that are viewed centrally, like words and faces, are organized relative to one another. Here,

invasive and non-invasive neuroimaging suggests there is a mosaic of regions in ventral temporal

cortex that respond selectively to either words or faces. These regions display differences in the

strength and timing of their responses, both within and between brain hemispheres, suggesting

they play different roles in perception. These results illuminate extended, bilateral, and dynamic

brain pathways that support face perception and reading.

Introduction

Investigations into the spatial organization of category-selectivity in ventral temporal cortex

(VTC) have been instrumental in establishing several organizational principles of the visual

system. Functional magnetic resonance imaging (fMRI) studies have helped identify lateral-

medial biases in ventral stream responses to objects depending on where they typically appear in

the visual field (retinotopic eccentricity) (Hasson et al., 2002; Konkle and Caramazza, 2013;

Grill-Spector and Weiner, 2014). Specifically, lateral regions of VTC are selective for objects

that tend to be viewed centrally (foveated), like words and faces, whereas more medial regions

are selective for objects that tend to fall on the periphery of the retina, like navigationally

relevant information such as buildings (Haxby et al., 1996; Aguirre et al., 1998; Cohen et al.,

2000; Hasson et al., 2002). This broad principle of organization by eccentricity fails to inform us

about how representations of different stimuli that are foveated, like words and faces, are

organized in VTC relative to one another.

Despite sharing similar typical retinotopic eccentricity, word and face stimuli are highly

distinct along several axes that are also hypothesized to influence where they are processed in

VTC (Op de Beeck et al., 2019). Word- and face-processing operate on very different low-level

visual properties (Kay and Yeatman, 2017), follow different developmental trajectories (Saygin

et al., 2016), and feed into distinct networks that support either language or social interactions

(Stevens et al., 2015, 2017), respectively. Despite this, the cortical localizations for word- and

face-processing in VTC are remarkably close together, and it remains debated whether or not

there are regions in VTC that independently encode word or face information at all (Behrmann

and Plaut, 2013).

Neuroimaging studies have separately mapped word- and face-processing networks in VTC. Printed word recognition is thought to be carried out in part by a network of regions along the left occipitotemporal sulcus, that differ in the complexity of their responses and are thought to be hierarchically organized (Halgren et al., 1994; Cohen et al., 2000; Vinckier et al., 2007; Dehaene and Cohen, 2011; Lerma-Usabiaga et al., 2018). Face-processing is thought to be carried out in part by a network of regions distributed bilaterally along the midfusiform sulcus (Tsao et al., 2008; Weiner and Grill-Spector, 2010). However, few studies have investigated VTC's responses to word and face stimuli within the same participants (Allison et al., 1994; Haxby et al., 1994; Puce et al., 1996; Matsuo et al., 2015; Harris et al., 2016). Those that have, have relied on low sample sizes or imaging modalities with differential sensitivity to different aspects of neural activity (like high and low-frequency neural activity (Engell et al., 2012; Jonas et al., 2016)). Therefore, much remains unknown about how visual word- and face-processing networks organize relative to one another, and to what degree they overlap (Haxby et al., 1994; Puce et al., 1996; Dehaene et al., 2010; Matsuo et al., 2015; Harris et al., 2016).

Further, word- and face-selective regions have primarily been mapped using methods lacking high spatiotemporal resolution. Therefore, it is unclear if the nodes within these processing networks differ in the temporal dynamics of their responses, although previous studies have suggested that different regions may contribute to distinct stages of word- and face-processing (Federmeier and Kutas, 1999; Vinckier et al., 2007; Li et al., 2018). Further, category-selective maps derived from BOLD responses may be incomplete due to BOLD's increased sensitivity to early stimulus evoked activity (100-300 ms after stimulus presentations) relative to later responses (Jacques et al., 2016; Ghuman and Martin, 2019) and greater

correlation with high frequency broadband activity in invasive neural recordings compared to

lower-frequency electrical potentials (Engell et al., 2012; Jacques et al., 2016).

In the present study, we characterized the spatial organization and functional dynamics of

word- and face-processing networks within VTC using intracranial electroencephalography

(iEEG) data collected from 36 patients with pharmacologically intractable epilepsy and 7 T

fMRI data collected from eight healthy participants.

Materials and Methods

Intracranial EEG data collection and preprocessing

Participants

38 patients (14 males, ages 19-65, 32 righthanded) had intracranial surface and/or depth

electrodes implanted for the treatment of pharmacologically intractable epilepsy. Depth

electrodes were produced by Ad-Tech Medical and PMT Corporation and were 0.86 and 0.8 mm

in diameter, respectively. Grid electrodes were produced by PMT Corporation and were 4 mm in

diameter. Because depth electrode contacts are cylindrical, the surface area of the recording site

was similar across grid and strip electrode contacts. To be concise, "electrode contacts" are

referenced to as "electrodes" throughout the manuscript. No consistent differences in neural

responses were observed between grid and depth electrodes. Only electrodes implanted in ventral

temporal cortex, defined as below the inferior temporal gyrus and anterior to the posterior tip of

the fusiform in the participant-centered space, were considered in this study. Two patients did

not have any electrodes within this region of interest, therefore only data from 36 participants

were analyzed for this study. Electrodes identified as belonging to the seizure onset zone based

on the clinical report or showing epileptiform activity during the tasks were excluded from the

analysis. All participants gave written informed consent. The study was approved by the University of Pittsburgh Institutional Review Board. Patients were monetarily compensated for their time.

Electrodes were localized via either post-operative magnetic resonance imaging (MRI) or computed tomography scans co-registered to the pre-operative MRI using Brainstorm (Tadel et al., 2011). Surface electrodes were projected to the nearest point on the pre-operative cortical surface automatically parcellated via Freesurfer (Dale et al., 1999) to correct for brainshift (Hermes et al., 2010). Electrode coordinates were then coregistered via surface-based transformations to the fsaverage template using Freesurfer cortical reconstructions.

Experimental Design

All participants underwent a category localizer task where they viewed grayscale images presented on a computer screen positioned two meters from their face. Images occupied approximately 6 x 6 degrees of visual angle and were presented for 900 ms with 1500 ms inter-stimulus interval with random 400 ms jitter. Participants were instructed to press a button every time an image was presented twice in a row (1/6 of the trials). These repeat trials were excluded from the analysis yielding 70 trials per stimulus category left for analysis. Several participants underwent multiple runs of this task and therefore had 140-210 trials per stimulus category.

31 of the participants saw pictures of faces, words, bodies, hammers, houses and phase-scrambled faces. The remaining participants viewed a modified set of stimuli with the same viewing parameters described above. One participant viewed pictures of consonant-strings and pseudowords instead of hammers, two viewed shoes instead of words, one viewed consonant-

strings and pseudowords instead of hammers and houses, and one viewed general tools and animals instead of hammers.

A subset of the participants that underwent the category localizer task also participated in word and/or face individuation tasks (Table 1). These tasks shared identical presentation parameters as the category-localizer task (i.e. inter-stimulus interval, stimulus-on time, and viewing angle) but contained different images. Twelve underwent a word individuation task that included pictures of real words, pseudowords, and consonant-strings or false-fonts. Participants again were instructed to respond if a given stimulus was repeated twice in a row. Every stimulus (i.e. individual word) was presented sixty times. Twenty underwent a face individuation task where they viewed individuals of varying identity and emotions. Participants were instructed to indicate if each face was male or female during this task. Each identity was repeated 60 times.

Local field potentials were recorded via a GrapeVine Neural Interface (Ripple, LLC) sampling at 1 kHz. Notch filters at 60/120/180 Hz were applied online. Data was subsequently filtered from 0.1-115 Hz to isolate single trial potentials (stP) or decomposed via Morlet wave convolution to determine the power from 40-100 Hz to isolate single trial high frequency broad-band activity (stHFBB). These stHFBB responses were then Z-scored based on the baseline period from 500-0 ms proceeding stimulus onsets. It has been previously shown that these two aspects of the local-field potential, stP and stHFBB, contain complementary information (Miller et al., 2016), though also potentially arise from different neurophysiological generators (Engell et al., 2012; Hermes et al., 2012; Jacques et al., 2016; Leszczyński et al., 2020). Therefore, to assess the overall selectivity across VTC we use both as features in the classifiers described in *Multivariate temporal pattern analysis* (Figures 1B, 2-4, 6-8). We also investigated the independent contributions of these signal components to our category-selectivity maps (Figure

6). Trials where the stHFBB or stP exceeded 5 standard deviations from the mean were thought

to contain noise and therefore excluded from further analysis.

Determining Language Laterality

Records from preclinical magnetoencephalography (MEG) language mapping sessions were

used to determine the laterality of language function for 30 of the 36 iEEG participants.

Language mapping records for the remainder of the participants could not be located. The

preclinical language mapping records contained laboratory technician notes indicating whether

MEG activity during reading, listening, and word-repetition tasks was lateralized to the left or

right hemisphere. The original data from these sessions was not available to conduct more

precise analyses of language laterality for these participants.

Multivariate temporal pattern analysis

To determine which electrodes contained information about word and face categories, leave-one

trial out cross-validated Gaussian Naïve Bayes classifiers were used to predict the category of

object participants were viewing given a sliding 100 ms of neural activity from one iEEG

electrode during the category-localizer task (six-way classification). Signals from stP and

stHFBB were both fed in as features to a single classifier for the main selectivity maps. This

procedure was repeated from 100 ms prior to 900 ms after stimulus onset with 10 ms time-step to

derive a time-course of decoding at each VTC electrode. We also ran separate classifiers on only

features from stP or stHFBB to investigate the independent sources of information contained

within these signal components. We ensured the number of features fed into these two types of

classifiers was consistent by averaging 10 ms bins of stP, since stHFBB was sampled only every

10 ms, before classification.

Face-selective iEEG electrodes were defined as those that achieved a peak sensitivity (d')

of decoding for faces greater than the chance at the $p < 0.05$ level, Bonferroni corrected for

multiple comparisons in time and across the total number of electrodes within a participant.

Sensitivity (d') describes the separation between a classifier's noise and signal distributions and

is defined as the inverse normal cumulative distribution function (Z') of the true positive rate

(TPR) minus the inverse normal cumulative distribution function of the false positive rate (FPR),

$$Z'(TPR) - Z'(FPR).$$

The Bonferroni corrected d' sensitivity threshold was found by performing a binomial test on a

null distribution of 1 million d' sensitivities that were obtained by randomly classifying

permutations of the trial labels. A small number of electrodes responded to all categories *except*

faces, which resulted in above-chance face classification, since the distribution of responses to

faces was significantly different from the responses to other object categories. Therefore, we

imposed an additional criterion to determine selectivity: face-selective channels had to

demonstrate a maximum peak event-related potential or event-related broadband response to

faces relative to the other object categories. An identical procedure was done to define word- and

house-selective electrodes.

To determine the independence of word- and face-selectivity within electrodes, we

repeated the above multivariate pattern analysis for word- and face-selective electrodes after

removing trials from the category they were most selective to. Word-selective electrodes were

determined to also be selective for face stimuli if, after removing trials when words were

presented, we could reliably predict trials where faces were presented from the other object

categories (d' sensitivity corresponding to $p < 0.05$, Bonferroni corrected for multiple temporal

and electrode comparisons within participants using the same permutation test described above).

1   Further, we stipulated that this d' for faces must be greater than the d' for all the remaining

2   object categories. An identical procedure was used to define face-selective electrodes that were

3   also selective for words.

4   To determine if word- and face-selective electrodes contained exemplar-level information

about either faces or words, we performed pairwise classification of the face and word

individuation stimuli for the electrodes on which we had data (Table 1). Specifically, in the case

of word individuation, we used three-fold cross-validated Gaussian Naïve Bayes classifiers to

predict which of two real words a participant was viewing based on sliding 100 ms of data from

the word-selective electrodes. Three-fold cross-validation was used instead of leave-one-out

cross validation (which was used for assessing category-level selectivity) to save computational

time as there were many more models (stimulus pairs) tested with the exemplar classifier. We

repeated this procedure across all pairs of real-words of the same length and averaged the time-

courses of this pairwise decoding (56 pairs of words). We determined the $p < 0.05$ chance-level

of this average pairwise decoding by repeating this procedure 1,000 times on data with shuffled

trial labels in a subset of the word-selective electrodes (Maris and Oostenveld, 2007). These

global null distributions were similar across the randomly subsampled electrodes and therefore

we chose a d' threshold corresponding to the highest $p < 0.05$ level obtained from this randomly

chosen subset. We ran similar pairwise decoding and threshold definition on real-word versus

pseudowords of the same length (36 pairs) and real-word versus false-font stimuli (136 pairs) to

determine if electrodes that could not individuate real-words could perform these finer

discriminations compared to those tested in the category localizer task.

22   Similarly, for face individuation we performed pairwise decoding of face stimuli during

23   sliding 100 ms time-windows of face-selective electrode activity. We then averaged these time-

courses across all 120 pairwise face classifications and calculated the $p < 0.05$ corrected level by repeating the permutation analysis described for the word individuation task on a random subset of face-selective electrodes.

Spatiotemporal k-means clustering

We used a spatiotemporal variant of k-means clustering to determine if spatially contiguous word- or face-selective regions demonstrated distinct temporal dynamics. For word- and face-selective electrodes, we separately standardized the d' sensitivity time-courses derived from the category-level multivariate classifiers of left and right hemisphere electrodes from 100 to 600 ms post stimulus onset. We then concatenated this matrix with the electrodes' MNI-coordinate, which was multiplied by a constant (spatial weighting parameter) that modulated the weight of the spatial versus temporal components of the signal to the clustering algorithm. We then performed k-means clustering using Euclidean distances and 100 repeats with random initializations to determine clusters of nearby word- or face-selective electrodes within each hemisphere that demonstrated correlated dynamics. Because the d' time-courses were standardized, Euclidean distances were equivalent to correlation distance for the temporal data and Euclidean distance for the spatial data.

To determine the optimal weighting of spatial and temporal signal components and optimal number of clusters, we calculated the total spatial and temporal variance explained by the clustering solutions run with several spatial weighting parameters. This was performed for k = 1 to 10 clusters per hemisphere per faces or words. The elbow method was used to determine the optimal number of clusters per hemisphere. The optimal number of clusters was 4 for right hemisphere face-selective electrodes, 3 for right hemisphere word-selective electrodes, 3 for left hemisphere face-selective electrodes, and 4 for left hemisphere word-selective electrodes. We

chose the spatial weighting parameter that explained the maximum amount of variance across k

= 3 to 4 clusters per hemisphere per category (spatial weight = 300). Small deviations in the

spatiotemporal weighting parameter did not strongly affect the overall organization of

spatiotemporal clusters. The dynamics of these electrode clusters were then determined by

averaging the selectivity time courses (d' derived using *multivariate temporal pattern analysis*)

across the electrodes belonging to each cluster.

Statistical analyses

Two sample t-tests were used to compare peak d' sensitivity, peak latency, and onset latency for

right versus left word- and face-selective electrodes. Onset latency was defined as the first time

point that the d' sensitivity reached a $p < 0.001$ threshold, which was non-parametrically defined

using the d' sensitivities of all object-selective electrodes from 500-0 ms prior to stimulus onset.

Spearman's rank-order correlations were used to test for relationships between peak d'

sensitivities and latency. We used linear mixed effects models to compare face and real word

individuation in the category-selective clusters identified by the spatiotemporal k-means

algorithm. Linear mixed effects models allowed us to determine if there were differences in peak

individuation d' or latency across these clusters while correcting for cross-subject differences.

We only compared spatiotemporal clusters with greater than 10 electrodes with individuation

data. The Satterthwaite approximation was used to estimate the degrees of freedom in these

linear mixed effects models to compute the reported p-values. The time points corresponding to

the leading edge of the classification window were used for all temporal statistical analyses.

fMRI data collection and preprocessing

Participants

Eight participants (six females, mean age 25 years) participated in the fMRI experiment. All

participants were right-handed, had normal or corrected to normal vision and gave written

informed consent. The National institutes of Health Institutional review Board approved the

consent and protocol (protocol 93 M-0170, clinical trials #NCT00001360). Participants were

monetarily compensated for their time.

fMRI scanning parameters

All fMRI scans were conducted on a 7 T Siemens Mangetom scanner at the Clinical Research

Center on the National Institutes of Health campus. Partial volumes of the occipital and temporal

cortices were acquired using a 32-channel head-coil (42 slices, 1.2x1.2x1.2 mm; 10% interslice

gap; TR=2s, TE=27ms; matrix size=170x170).

Experimental Paradigm

Participants fixated centrally whilst images of words, faces and houses were presented in blocks

(16s per block). These images were taken from the same category localizer task presented to

iEEG patients. In each block 20 exemplar stimuli were presented (300ms with a 500ms ISI).

Participants performed a one-back task, responding, via MRI compatible response box, whenever

the same image appeared twice in a row. Participants completed 10 runs of the localizer.

fMRI data preprocessing

All data were analyzed using the Analysis of Functional NeuroImages (AFNI) software package

(Cox, 1996). Prior to statistical analysis, all images were motion corrected to the first volume of

the first run. Post motion-correction data were detrended.

Statistical analysis

To identify word-, face- and house-selective regions, we performed a general linear model (GLM) analysis using the AFNI functions 3ddeconvolve and 3dREMLfit. The data at each time-point were treated as the sum of all effects thought to be present at that time point and the time series was compared against a Generalized Least Squares Regression model fit with REML estimation of the temporal auto-correlation structure. Responses were modelled by convolving a standard gamma function with a 16 second square wave for each condition (words, faces & houses). Estimated motion parameters were included as additional regressors of no-interest and fourth-order polynomials were included to account for any slow drifts in the MRI signal over time. Significance was determined by comparing the beta estimates for each condition (normalized by the grand mean of each voxel for each run) against baseline.

Split-half analysis

For each participant, the ten localizer runs were divided into odd and even splits. In each split, we performed the same GLM analysis as described above and looked for significant voxels for the contrast of words versus faces. Despite having only half of the data, we observed significant word-selectivity that was medial of face-selectivity consistently across participants. In order to quantify this selectivity in an independent manner, we first defined medial word-selective regions within a split (e.g. odd) and then sampled the data from the other half (e.g. even). ROIs were defined using data spatially smoothed with a 2 mm Gaussian kernel to generate spatially contiguous clusters, whereas the test data was not spatially smoothed. To avoid any bias in node selection, this process was then reversed and the average computed. Within each ROI we calculated the average t-value for each condition versus baseline.

Results

From 1,396 intracranial electrode contacts implanted within or on the surface of VTC of 36

patients, we isolated those implanted in regions that were highly selective for either faces, words,

or houses. Highly face-selective electrodes were defined as those that had both (1) single-trial

responses that could significantly discriminate face presentations from presentations of five other

object categories (words, houses, bodies, hammers, and phase-scrambled objects; $p < 0.05$ level,

Bonferroni corrected for multiple spatial and temporal comparisons within participant; see

Materials and Methods) and (2) responded maximally to faces compared to all other object

categories on average. This ensured that electrodes designated as highly "face-selective" were

those that responded maximally and were significantly selective for faces compared to the five

other object categories. An identical procedure was used to define word- and house-selective

electrodes.

108 electrodes demonstrated primarily face-selective responses (80 in the left, 28 in the

right), 87 demonstrated primarily word-selective responses (64 in the left, 23 in the right), and 85

demonstrated primarily house-selective responses (44 in the left, and 41 in the right) (Figure 1).

Figure 2 and Table 1 illustrate the distribution of object-selective electrodes across participants.

The greater number of left versus right object-selective electrodes was comparable to the greater

coverage of left VTC relative to right VTC in our patient population (883 electrodes implanted in

the left, 513 in the right, Figure 1A). Although some word- and face-selective electrodes

demonstrated partial selectivity for the other object category, there were several examples of

electrodes that were strongly tuned to only words or faces (Figure 3). This suggests that the

neural circuits responsible for processing words and faces are, at least, partially dissociable

(Behrmann and Plaut, 2013; Susilo and Duchaine, 2013; Susilo et al., 2015).

To assess how word- and face-processing networks organize relative to one another, the

spatial topography of word-, face-, and house-selective electrodes was examined. At the group

level, selectivity to house stimuli was found primarily along the left and right parahippocampal

gyrus, with some cases where selectivity extended into the collateral sulcus and medial fusiform

gyrus. These patches were generally medial to word- and face-selective locations, consistent with

previous fMRI and iEEG studies (Halgren et al., 1994; Haxby et al., 1996; Aguirre et al., 1998;

Cohen et al., 2000; Kadipasaoglu et al., 2016). Face-selectivity was found primarily along the

left and right fusiform gyrus with some face-selective regions within the lingual gyrus, and

occipitotemporal sulcus (Figure 1B). Consistent with prior findings (Cohen et al., 2000), word-

selective regions were found on the lateral bank of the fusiform and into the occipitotemporal

sulcus in the left hemisphere. Word-selective regions were also found anterior to most prior

reports from fMRI, in locations that generally have poor signal due to susceptibility artifacts

(Devlin et al., 2000). In contrast to most maps of word- and face-selective regions obtained from

fMRI (Allison et al., 1994; Haxby et al., 1994; Puce et al., 1996; Harris et al., 2016; Saygin et al.,

2016; Dehaene-Lambertz et al., 2018; Gomez et al., 2018), a mosaic of word-selective regions

were also found medial to face-selective regions, on the medial bank of the fusiform and into the

collateral sulcus. Each of these face-, word-, and house-selective regions were found in multiple

participants (Figure 2), demonstrating relatively consistent localization of these regions at a

group level.

Interdigitation of word- and face-selective regions was seen in the left hemisphere of 5

out of 9 participants with at least two word-selective electrodes and one face-selective electrode

or vice-versa and in the right hemisphere of 3 out of 5 such participants (Table 1, see Figure 4 for

examples). Word-selective regions were found strictly medial to face-selective regions in the left

hemisphere of 7 out of 10 participants with at least one word- and one face-selective electrode

and in right hemisphere of 4 out of 5 participants (Table 1, see Figure 4 for an example). Thus,

highly word-selective regions medial to face-selective regions was not simply a consequence of

individual variability in a group-level map but instead was detected in the majority of

participants that had coverage of both face- and word-selective VTC.

Because word-selective patches were found medial to face-selective patches in the iEEG

data, which is generally not observed in 3 T fMRI studies (Haxby et al., 1994; Puce et al., 1996;

Dehaene et al., 2010), we sought to determine if a similar organization existed in healthy

participants using the higher resolution of 7 T fMRI. When contrasting responses to words and

faces in eight participants, face-selectivity was primarily centered on the midfusiform sulcus

while word-selectivity was greatest in the occipitotemporal sulcus (Figure 5). Consistent with the

iEEG results, six of the eight participants demonstrated left word-selective regions medial to

face-selective regions on the fusiform gyrus. In these medial word-selective patches, responses to

words were significantly greater than responses to both face and house stimuli ($p < 0.001$, split-

halves analysis). These medial word-selective regions were approximately 1/3rd the size of more

lateral word-selective regions (mean size of lateral word-selective regions: 398 voxels; std. error:

43 versus medial regions: 139 voxels; std. error: 29 voxels; $p < 0.01$). Also, 7 out of 8 of the

healthy participants demonstrated word-selective patches near the anterior tip of the fusiform,

despite susceptibility artifacts (Devlin et al., 2000), consistent with the iEEG data (Figure 1B).

Altogether, the map of word- and face-selective regions of the left hemisphere derived from 7 T

fMRI were consistent with those derived from iEEG, medial and anterior word-selective regions

are not seen in most maps drawn from 3 T fMRI (Haxby et al., 1994; Puce et al., 1996; Dehaene

et al., 2010).

The maps in Figures 1-3 were made by combining two key aspects of the iEEG signals,

the single trial potentials (stP) and the single trial high frequency broadband activity (stHFBB),

to examine the category-selectivity of the underlying VTC neural populations in aggregate across

these signal components. Studies have shown that while category-selectivity demonstrated in stP

and stHFBB often overlaps, they are not redundant (Engell and McCarthy, 2011; Engell et al.,

2012; Miller et al., 2016), suggesting that stP and stHFBB have at least partially distinct

physiological generators. To examine these signal components separately, we trained

multivariate classifiers solely on stP or stHFBB and isolated electrodes that were selective in

either signal component using the same criteria as before (single-trial discriminability and

highest signal amplitude for words, faces, or houses). 58 electrodes showed significant

selectivity in both stP and stHFBB (Figure 6A). Notably, the regions that demonstrated

selectivity in both stP and stHFBB were those most often identified in canonical maps of

category-selectivity based on fMRI (Cohen et al., 2000; Vinckier et al., 2007; Tsao et al., 2008;

Weiner and Grill-Spector, 2010; Lerma-Usabiaga et al., 2018). Specifically, house-selectivity

was restricted to the parahippocampal cortex, face-selectivity was primarily restricted to the

fusiform bilaterally, and word-selectivity was restricted primarily to the left posterior-lateral

fusiform and occipitotemporal sulcus. Regions that were less consistent with canonical fMRI

maps tended to be those that were not significantly selective in both stP and stHFBB. For

example, the medial word-selective patches were primarily seen in stP alone (Figure 6B),

whereas anterior and right hemisphere word-selectivity was prevalent in either stP or stHFBB

alone (Figures 6B and 6C). Broadly, more electrodes demonstrated selectivity in stP (232

electrodes from 32 participants, Figure 6B) compared to stHFBB (115 electrodes from 24

participants, Figure 6C). More widespread stP selectivity is consistent with a previous study

comparing stP and stHFBB responses for faces in VTC, though that study did not observe any

cases where selectivity for faces was demonstrated in stHFBB but not stP (Engell and McCarthy,

2011). The similarities and differences in selectivity demonstrated in stHFBB and stP are

consistent with the hypothesis that these signals have different physiological generators

(Lachaux et al., 2005), which may differ in their laminar distribution (Leszczyński et al., 2020)

and spatial signal-to-noise falloff (Engell and McCarthy, 2011). Additionally, different category-

selectivity across these iEEG signal components may also help explain differences between

category-selectivity maps drawn from iEEG and fMRI, as some studies suggest fMRI has

differential sensitivity to these aspects of the iEEG signal (Conner et al., 2011; Engell et al.,

2012; Jacques et al., 2016).

One question is whether word- and face-selective regions identified using iEEG

discriminate between individual face and word exemplars, respectively. Classifying at the

exemplar level also can address the potential concern that the word- and face-selective regions

identified using iEEG may be responding to low-level features that drastically differ between the

sampled image categories. A subset of the iEEG participants underwent independent word and

face individuation tasks (see *Materials and Methods,* Table 1). Activity from 85 out of 97

sampled face-selective electrodes in 13 participants could be used to reliably predict the identity

of a presented face. Similarly, activity from 40 out of 53 sampled word-selective electrodes from

10 participants could be used to discriminate single words of the same length from one another.

Of those 13 word-selective electrodes that could not reliably achieve word individuation, six

could reliably discriminate pseudowords from real words of the same length, seven could

reliably discriminate false-fonts from real words. Therefore, most of the word- and face-selective

regions mapped with iEEG contained reliable exemplar-level information specific to the categories they were selective to.

Peak word and face individuation was significantly correlated with peak category-selectivity in word and face-selective regions for which we had individuation data (word-selective: Spearman's $\rho(53) = 0.50$, $p < 0.0001$, face-selective: $\rho(97) = 0.48$, $p < 0.0001$). Note that correlations in peak category-selectivity and within-category individuation may arise due to similar differences in measurement noise across recording contacts (for example, due to the distance the electrode was placed from the underlying face or word selective neural groups), underlying neural/physiological factors, or some mix of both.

In addition to the medial band of word-selective regions, there were a high proportion of right word-selective electrodes in our iEEG population (Figure 1B, Table 1). Although this finding is consistent with some other fMRI (Ben-Shachar et al., 2007; White et al., 2019) and iEEG studies (Halgren et al., 1994; Lochy et al., 2018), right hemisphere word-selectivity is often not seen in neuroimaging (Cohen et al., 2000, 2002) and was not very strong in our 7 T fMRI data either (Figure 5). 23 word-selective electrodes were found across nine participants in right VTC, out of 21 participants with right VTC object-selectivity. This discrepancy between right word-selectivity observed in fMRI and iEEG was also not attributable to participant handedness, since no participant with right word-selective regions was lefthanded. Three out of nine of these participants demonstrated evidence for bilateral language function while the other six demonstrated left dominant language function determined by preclinical magnetoencephalography (MEG, see *Materials and Methods*). Across the entire participant population, seven out of 30 iEEG participants with preclinical MEG demonstrated bilateral language function, the others were considered left dominant. One participant with bilateral

language function and right hemisphere object-selectivity did not demonstrate right word-

selectivity. Overall, neither participant handedness nor language dominance sufficiently explains

the high proportion of word-selective regions found in right VTC.

While neither language laterality nor handedness explained right word-selectivity,

substantial differences were seen in the dynamics of neural activity recorded from left versus

right word-selective regions (Figure 7). Latency to word-selectivity onset and peak was shorter

in left compared to right hemisphere word-selective regions (mean onset latency difference +/-

95 % CI: -133 +/- 61 ms, T(85) = -4.4, p < 0.0001, mean peak latency difference: -138 +/- 63 ms,

T(85) = -4.3, p < 0.0001, Figure 7). These relationships held when taking into account potential

differences in posterior to anterior coordinate of word-selective regions across hemispheres

(onset: T(85) = -4.01, p = 0.0001, peak: T(85) = -3.97, p = 0.0002). There was no significant

difference between the latency to peak d' sensitivity or sensitivity onset for right and left face-

selective regions (mean onset latency difference: -29 +/- 53 ms, T(106) = -1.1, p = 0.28, mean

peak latency difference: 18 +/- 57 ms, T(106) = 0.63, p = 0.53, Figure 7). Additionally, the

amplitude of peak d' sensitivity for words was significantly greater in the left compared to right

hemisphere word-selective regions (mean peak d' sensitivity difference: 0.66 +/- 0.37, T(85) =

3.5, p = 0.0006). The amplitude of peak d' sensitivity to faces was also significantly greater in

the left compared to right hemisphere face-selective regions (mean peak d' sensitivity difference:

0.58 +/- 0.39, T(85) = 3.0, p = 0.0037). There was a significant correlation between peak latency

and peak magnitude within face-selective regions in the left ($\rho(80)$ = -0.61, p < 0.0001) and right

($\rho(28)$ = -0.79, p < 0.0001) hemisphere and word-selective regions in the left ($\rho(64)$ = -0.68, p <

0.0001), but not right ($\rho(23)$ = -0.15, p = 0.48) hemisphere, suggesting that longer peak latencies

were associated with smaller peak selectivity. These correlations were not significantly different

between face-selective regions in the left and right hemisphere (T(85) = -1.56, p = 0.058), but

there was a greater correlation between peak latency and magnitude in left compared to right

hemisphere word-selective regions (T(85) = 2.63, p = 0.004) Given that it was only true for

word-selective electrodes, the relatively slower response of right versus left word-selective

regions may potentially explain differences in word-selectivity maps derived from iEEG and

fMRI and may reflect the left hemisphere bias for language.

Finally, using the iEEG data, we sought to determine if there were any differences in the

temporal dynamics of neural responses across word or face-selective regions within the same

hemisphere. We used a spatiotemporal k-means clustering algorithm to find spatially contiguous

regions of left and right VTC which demonstrated correlated category-selective dynamics. After

optimizing the algorithm to capture the most spatiotemporal variance with the optimal number of

clusters (see *Materials and Methods*), we could compare the dynamics of distinct word- and

face-selective clusters within VTC.

Word-selective regions were clustered into 4 distinct left hemisphere clusters and 3 right

hemisphere clusters (Figure 8A). Word-selective regions on the left fusiform gyrus demonstrated

the earliest and strongest selectivity, peaking around 200 ms (Figure 8B, gray). Left hemisphere

medial word-selective regions and right hemisphere word-selective regions came next, peaking

around 300 ms (Figure 8B, green and cyan) followed by lateral regions around 350 ms (Figure

8B, red). Word-selective regions in left anterior VTC peaked around 400-450 ms (Figure 8B,

blue); right more anterior regions peaked around 600 ms (Figure 8B, magenta). When

considering word-selectivity dynamics exhibited independently in stP and stHFBB signal

components, word-selective electrodes on the fusiform demonstrated strong selectivity in both

signal components, whereas other regions displayed distinct dynamics across these signal

components (Figures 7C-D).

Face-selective regions were organized into 3 distinct clusters in the left hemisphere and 4

distinct clusters in the right hemisphere (Figure 8E). Face-selective regions of the left and right

fusiform gyrus demonstrated the earliest and largest peak selectivity around 200-250 ms (Figure

8F, gray and cyan). More anterior right hemisphere regions and a cluster of electrodes in left

posteromedial VTC (Figure 8F, yellow and green) peaked around 300 ms. Finally, more anterior

face-selective electrodes in left and right VTC peaked around 400 ms (Figure 8F, blue, black,

and magenta). When considering face-selectivity dynamics exhibited independently in stP and

stHFBB signal components, electrodes on the fusiform demonstrated strong selectivity in both

components, whereas other regions displayed distinct dynamics across these signal components

(Figures 7G-H).

From electrodes sampled in the word individuation task, we observed stronger word

individuation in left word-selective regions on the fusiform compared to the more medial word-

selective cluster illustrated in Figure 8A (peak d' of fusiform minus medial regions: $T(30) =$

3.62, $p = 0.001$, linear mixed-effects model). There was no significant difference between the

latency to peak word individuation across these clusters ($T(30) = 2.91$, $p = 0.68$). There were not

sufficient subjects with electrodes in the other word-selective clusters with word individuation

data to make comparisons between all clusters. Neither peak face individuation ($T(50) = 1.03$, p

$= 0.31$) nor latency to peak face individuation ($T(50) = -0.21$, $p = 0.84$) was significantly

different between face-selective regions along the left fusiform gyrus and the posteromedial face-

selective cluster observed in Figure 8E. There were not sufficient subjects with electrodes in the

other face-selective clusters with face individuation data to make comparisons between all

clusters.

Overall, for both faces and words, these results suggest a cascade of processing that

begins in the fusiform. Notably, the dynamics of these clusters suggest that they contribute to

distinct stages of face- and word-processing, since the latencies of their responses are far longer

than would be expected from feedforward visual transmission delays alone (Thorpe et al., 1996;

Kravitz et al., 2013), but not long enough to exclude them from being relevant to perceptual

behavior (Quian Quiroga et al., 2008; Tang et al., 2014) .

Discussion

In the current study, we found several VTC regions that demonstrated strong word-, face- and

house-selective responses. Although activity recorded from VTC electrodes often contained

information about multiple object categories, several selectively responded only to faces or

words (Figure 3). Electrodes which demonstrated preference to only words or faces suggests that

VTC word- and face-processing networks are not entirely overlapping (Behrmann and Plaut,

2013), but instead involve at least some independent nodes (Susilo and Duchaine, 2013; Susilo et

al., 2015), which is also supported by stimulation and lesion evidence (Hirshorn et al. 2016).

In both the iEEG and fMRI data, strong face-selectivity along the fusiform gyrus was

adjoining with highly word-selective regions in and around the occipitotemporal and collateral

sulci. House-selective regions were found primarily along the parahippocampal gyrus. This

organization of house- versus word- and face-selective regions supports that typical retinotopic

eccentricity is an important organizing principle of VTC (Grill-Spector and Weiner, 2014). The

word-selective regions around the occipitotemporal sulcus are consistent with prior studies

showing word-selectivity within lateral aspects of (Dehaene et al., 2002; Price and Devlin,

2003). Due to sparse and variable sampling across participants, the data cannot address the

question of whether there is a gradient of word-selectivity along the occipitotemporal sulcus

(Vinckier et al., 2007) or distinct patches (Lerma-Usabiaga et al., 2018; White et al., 2019).

Despite some similarities with previous neuroimaging work, the iEEG and 7 T fMRI data

here are inconsistent with a map of VTC wherein word-selective regions are strictly lateral to

face-selective regions (Haxby et al., 1994; Puce et al., 1996; Dehaene et al., 2010). While there

has been some mixed reporting of word-selectivity in anterior and medial VTC regions (Allison

et al., 1994; Haxby et al., 1994; Puce et al., 1996; Harris et al., 2016; Saygin et al., 2016;

Dehaene-Lambertz et al., 2018; Gomez et al., 2018), most models of orthographic-processing

within VTC consider only the more lateral, traditional "visual word form area" (Dehaene et al.,

2002; Price and Devlin, 2003). The disagreement between the observed organization of face- and

word-processing networks in VTC and most previous maps drawn from fMRI may be the

product of spatial smoothing commonly applied during fMRI data analysis (Geissler et al.,

2005), signal dropout induced by susceptibility artifacts (Devlin et al., 2000), or the inferior

sensitivity of 3 T fMRI relative to 7 T fMRI. Here, a mosaic of word-selective regions was found

medial and anterior to face-selective regions at the group level and within multiple iEEG patients

and in 7 T fMRI in healthy individuals. This evidence makes it unlikely that our observations are

the product of inter-participant variability or differences between healthy controls and patients

with intractable epilepsy (see also (Matsuo et al., 2015; Jonas et al., 2016; Kadipasaoglu et al.,

2016; Lochy et al., 2018)). This mosaic organization of visual word-selective regions is similar

to the mosaic organization of auditory language processing networks (Flinker et al., 2011),

suggesting this pattern of organization may not be specific to the visual system.

Medial word-selective regions may reflect differential mediolateral VTC selectivity to object rectilinearity, with more medial VTC being more responsive to straight over curvy objects (Srihasam et al., 2014; Bao et al., 2020). However, the interdigitation of word- and face-selective regions along the mediolateral axis is not well captured solely by this rectilinear model or the retinotopic model. Instead, medial and lateral word-selective regions with distinct dynamics may indicate an interaction between multiple representational axes in VTC (Konkle and Caramazza, 2013; Grill-Spector and Weiner, 2014) and possibly competition between faces and words for cortical space (Behrmann and Plaut, 2020).

Previous studies have used electrical stimulation to demonstrate that a large portion of VTC, sometimes termed the "basal temporal language area," plays a role in language processing (Krauss et al., 1996; Mani et al., 2008; Fonseca et al., 2009; Enatsu et al., 2017). However, generalized language deficits after lesions of the basal temporal language area are relatively minor (Krauss et al., 1996) and the relationship between reading deficits and VTC lesions, other than the visual word form area (Gaillard et al., 2006; Hirshorn et al., 2016), is unclear. A recent study reported differential language-related deficits during reading, repetition, and picture naming depending on the area of VTC stimulated (Forseth et al., 2018). Future studies are necessary to understand the precise relationship between medial, lateral, and anterior word-selective VTC dynamics and these regions' functional contribution to reading and/or language processing.

Category-selective regions most consistent with prior fMRI studies were those that demonstrated selectivity in both stHFBB and stP iEEG signal components. In contrast, we found that medial word-selectivity was primarily demonstrated in stP rather than stHFBB. Previous studies have suggested that fMRI BOLD have differential sensitivity to stHFBB versus stP

(Hermes et al., 2012), with some suggesting greater sensitivity to stHFBB (Engell et al., 2012;

Jacques et al., 2016). Differential sensitivity to stP and stHFBB may explain why previous fMRI

studies have only inconsistently observed medial word-selective regions. Our 7 T fMRI data

shows that, with adequate power, both lateral and medial word-selective regions are seen in the

left hemisphere using BOLD within individual participants. Future studies are necessary to fully

understand the functional characteristics and neurophysiological generators of stP and stHFBB

iEEG components (Miller, 2010; Ray and Maunsell, 2011; Leszczyński et al., 2020) and how

they relate to any differential roles that medial and lateral word-selective regions play in reading.

In addition to this complex organization of word- and face-selectivity within

hemispheres, our iEEG analyses suggest that right word-selective regions demonstrate longer

latencies and lower amplitudes of peak selectivity compared to left word-selective regions,

which may reflect the primary role the left, language dominant, hemisphere plays in word-

processing (Fiez and Petersen, 1998). Previous studies have demonstrated weaker correlations

between object-selectivity measured with iEEG and fMRI correlations at later time windows

(Jacques et al., 2016). This may explain why bilateral selectivity to words is inconsistent across

neuroimaging studies.

It has previously been suggested that right word-selective regions (along with left

posterior word-selective regions) are involved in relatively early visual processing of words and

then this information flows to left anterior word-selective regions (White et al., 2019). However,

the dynamics observed here do not support this hypothesis, because left word-selectivity

substantially preceded right word-selectivity. Instead, the time-course of right hemisphere

activation is coincident with P300 and N400 potentials observed during reading, suggesting that

right hemisphere word-selective regions may support the left hemisphere in later computations,

such as those involving word syntax, memory encoding, and/or semantic processing (Friedman et al., 1975; Kutas and Hillyard, 1980; Federmeier and Kutas, 1999; Otten and Donchin, 2000; Arbel et al., 2011).

Word- and face-selective regions within hemispheres also demonstrated distinct dynamics. Word-selective regions on the left fusiform gyrus demonstrated the earliest and strongest word-selective responses. This was followed by word-selective activity in left occipitotemporal and collateral sulcus as well as right posterior word-selective regions. Finally, word-selective activity spread to anterior VTC between 400-600 ms. Further, the relatively later responses of word-selective regions outside of the fusiform may also contribute to differences in category-selective maps drawn from iEEG and fMRI (Jacques et al., 2016).

Face-selective responses were strongest and earliest on the fusiform gyrus bilaterally. A cluster of posteromedial face-selective electrodes was found in early visual cortex. The slower time-course of these regions compared to face-selective regions on the fusiform suggests this posterior face-selectivity is a result of top-down attentional effects previously reported during face-viewing (Mo et al., 2018). Following fusiform responses, face-selectivity was then seen in more anterior VTC.

While delays in processing along the posterior-to-anterior VTC axis for both faces and words is somewhat consistent with feedforward models of visual processing, the relative latencies are far longer than would be expected in these models (Thorpe et al., 1996; Kravitz et al., 2013). These results instead suggest more extended dynamics, perhaps governed by recurrent processes (Kravitz et al., 2013), with different category-selective regions contributing differentially to multiple, temporally extended stages of face- and word-processing (Ghuman et al., 2014; Hirshorn et al., 2016; Li et al., 2018). Further studies are required to identify these

stages and link them to different spatiotemporal patterns of VTC activity. It is important to acknowledge that when analyzing the data at this fine granularity, between-participant variability in neural organization may influence the differences observed in dynamics across regions (Zhen et al., 2015; Gao et al., 2018).

The high-resolution maps of category-selectivity within VTC provided here suggest that in addition to more extensively studied word-selective patches within the occipitotemporal sulcus, additional patches of word-selectivity exist along the mid and anterior fusiform gyrus. These patches of word-selectivity differ in their temporal dynamics from word-selective patches along the occipitotemporal sulcus, but still contain information about word identity. How these word-selective regions differentially contribute to reading and the factors that lead to the development of adjoining patches of word- and face-selective regions remain as important outstanding questions. Understanding this complex and dynamic map of selectivity in VTC is necessary to fully understand the organizational and computational principles governing object recognition.

References

Aguirre GK, Zarahn E, D'Esposito M (1998) An area within human ventral cortex sensitive to "building" stimuli: Evidence and implications. Neuron 21:373–383.

Allison T, Mccarthy G, Nobre A, Puce A, Belger A (1994) Human extrastriate visual cortex and the perception of faces, words, numbers, and colors. Cereb Cortex 4:544–554.

Arbel Y, Spencer KM, Donchin E (2011) The N400 and the P300 are not all that independent. Psychophysiology 48:861–875.

Bao P, She L, McGill M, Tsao DY (2020) A map of object space in primate inferotemporal

cortex. Nature 583:103–108.

Behrmann M, Plaut DC (2013) Distributed circuits, not circumscribed centers, mediate visual recognition. Trends Cogn Sci 17:210–219.

Behrmann M, Plaut DC (2020) Hemispheric Organization for Visual Object Recognition: A Theoretical Account and Empirical Evidence. Perception 49:373–404.

Ben-Shachar M, Dougherty RF, Deutsch GK, Wandell BA (2007) Differential Sensitivity to Words and Shapes in Ventral Occipito-Temporal Cortex. Cereb Cortex 17:1604–1611.

Chen R, Hillis AE, Pawlak M, Herskovits EH (2008) Voxelwise Bayesian lesion-deficit analysis. Neuroimage 40:1633–1642.

Cohen L, Dehaene S, Naccache L, Lehéricy S, Dehaene-Lambertz G, Hénaff MA, Michel F (2000) The visual word form area. Spatial and temporal characterization of an initial stage of reading in normal subjects and posterior split-brain patients. Brain 123:291–307.

Cohen L, Lehéricy S, Chochon F, Lemer C, Rivaud S, Dehaene S (2002) Language-specific tuning of visual cortex? Functional properties of the Visual Word Form Area. Brain 125:1054–1069.

Conner CR, Ellmore TM, Pieters TA, di Sano MA, Tandon N (2011) Variability of the relationship between electrophysiology and BOLD-fMRI across cortical regions in humans. J Neurosci 31:12855–12865.

Cox RW (1996) AFNI: Software for Analysis and Visualization of Functional Magnetic Resonance Neuroimages. Comput Biomed Res 29:162–173.

Dale AM, Fischl B, Sereno MI (1999) Cortical Surface-Based Analysis. Neuroimage 9:179–194.

Dehaene-Lambertz G, Monzalvo K, Dehaene S (2018) The emergence of the visual word form: Longitudinal evolution of category-specific ventral visual areas during reading acquisition. PLoS Biol 16:e2004103.

Dehaene S, Cohen L (2011) The unique role of the visual word form area in reading. Trends Cogn Sci 15:254–262.

Dehaene S, Pegado F, Braga LW, Ventura P, Nunes Filho G, Jobert A, Dehaene-Lambertz G, Kolinsky R, Morais J, Cohen L (2010) How learning to read changes the cortical networks for vision and language. Science (80-) 330:1359–1364.

Dehaene SC, Le Clec'H G, Poline J-B, Le Bihan D, Cohen L (2002) The visual word form area: a prelexical representation of visual words in the fusiform gyrus. Neuroreport 13:321–325.

Devlin JT, Russell RP, Davis MH, Price CJ, Wilson J, Moss HE, Matthews PM, Tyler LK (2000) Susceptibility-Induced Loss of Signal: Comparing PET and fMRI on a Semantic Task. Neuroimage 11:589–600.

Enatsu R, Kanno A, Ookawa S, Ochi S, Ishiai S, Nagamine T, Mikuni N (2017) Distribution and Network of Basal Temporal Language Areas: A Study of the Combination of Electric Cortical Stimulation and Diffusion Tensor Imaging. World Neurosurg 106:1–8.

Engell AD, Huettel S, McCarthy G (2012) The fMRI BOLD signal tracks electrophysiological spectral perturbations, not event-related potentials. Neuroimage 59:2600–2606.

Engell AD, McCarthy G (2011) The relationship of gamma oscillations and face-specific ERPs recorded subdurally from occipitotemporal cortex. Cereb Cortex 21:1213–1221.

Federmeier KD, Kutas M (1999) Right words and left words: Electrophysiological evidence for

hemispheric differences in meaning processing. Cogn Brain Res 8:373–392.

Fiez JA, Petersen SE (1998) Neuroimaging studies of word reading. Proc Natl Acad Sci U S A
95:914–921.

Flinker A, Chang EF, Barbaro NM, Berger MS, Knight RT (2011) Sub-centimeter language
organization in the human temporal lobe. Brain Lang 117:103–109.

Fonseca AT Da, Bénar CG, Bartoloméi F, Régis J, Démonet JF, Chauvel P, Liégeois-Chauvel C
(2009) Electrophysiological study of the basal temporal language area: A convergence zone
between language perception and production networks. Clin Neurophysiol 120:539–550.

Forseth KJ, Kadipasaoglu CM, Conner CR, Hickok G, Knight RT, Tandon N (2018) A lexical
semantic hub for heteromodal naming in middle fusiform gyrus. Brain 141:2112–2126.

Friedman D, Simson R, Ritter W, Rapin I (1975) The late positive component (P300) and
information processing in sentences. Electroencephalogr Clin Neurophysiol 38:255–262.

Gaillard R, Naccache L, Pinel P, Clémenceau S, Volle E, Hasboun D, Dupont S, Baulac M,
Dehaene S, Adam C, Cohen L (2006) Direct Intracranial, fMRI, and Lesion Evidence for
the Causal Role of Left Inferotemporal Cortex in Reading. Neuron 50:191–204.

Gao X, Gentile F, Rossion B (2018) Fast periodic stimulation (FPS): a highly effective approach
in fMRI brain mapping. Brain Struct Funct 223:2433–2454.

Geissler A, Lanzenberger R, Barth M, Tahamtan AR, Milakara D, Gartus A, Beisteiner R (2005)
Influence of fMRI smoothing procedures on replicability of fine scale motor localization.
Neuroimage 24:323–331.

Ghuman AS, Brunet NM, Li Y, Konecky RO, Pyles JA, Walls SA, Destefino V, Wang W,

Richardson RM (2014) Dynamic encoding of face information in the human fusiform gyrus. Nat Commun 5:5672.

Ghuman AS, Martin A (2019) Dynamic Neural Representations: An Inferential Challenge for fMRI. Trends Cogn Sci 23:534–536.

Gomez J, Natu V, Jeska B, Barnett M, Grill-Spector K (2018) Development differentially sculpts receptive fields across early and high-level human visual cortex. Nat Commun 9:788.

Grill-Spector K, Weiner KS (2014) The functional architecture of the ventral temporal cortex and its role in categorization. Nat Rev Neurosci 15:536–548.

Halgren E, Baudena P, Heit G, Clarke M, Marinkovic K (1994) Spatio-temporal stages in face and word processing. 1. Depth recorded potentials in the human occipital and parietal lobes. J Physiol - Paris 88:1–50.

Harris RJ, Rice GE, Young AW, Andrews TJ (2016) Distinct but overlapping patterns of response to words and faces in the fusiform gyrus. Cereb Cortex 26:3161–3168.

Hasson U, Levy I, Behrmann M, Hendler T, Malach R (2002) Eccentricity Bias as an Organizing Principle for Human High-Order Object Areas. Neuron 34:479–490.

Haxby J V, Horwitz B, Ungerleider LG, Maisog JM, Pietrini P, Grady CL (1994) The functional organization of human extrastriate cortex: a PET-rCBF study of selective attention to faces and locations. J Neurosci 14:6336–6353.

Haxby J V, Ungerleidert LG, Horwitz B, Maisog JM, Rapoport SI, Grady CL (1996) Face encoding and recognition in the human brain. Proc Natl Acad Sci 93:922–927.

Hermes D, Miller KJ, Noordmans HJ, Vansteensel MJ, Ramsey NF (2010) Automated

electrocorticographic electrode localization on individually rendered brain surfaces. J Neurosci Methods 185:293–298.

Hermes D, Miller KJ, Vansteensel MJ, Aarnoutse EJ, Leijten FSS, Ramsey NF (2012) Neurophysiologic correlates of fMRI in human motor cortex. Hum Brain Mapp 33:1689–1699.

Hirshorn EA, Li Y, Ward MJ, Richardson RM, Fiez JA, Ghuman AS (2016) Decoding and disrupting left midfusiform gyrus activity during word reading. Proc Natl Acad Sci 113:8162–8167.

Jacques C, Witthoft N, Weiner KS, Foster BL, Rangarajan V, Hermes D, Miller KJ, Parvizi J, Grill-Spector K (2016) Corresponding ECoG and fMRI category-selective signals in human ventral temporal cortex. Neuropsychologia 83:14–28.

Jonas J, Jacques C, Liu-Shuang J, Brissart H, Colnat-Coulbois S, Maillard L, Rossion B (2016) A face-selective ventral occipito-temporal map of the human brain with intracerebral potentials. Proc Natl Acad Sci 113:E4088–E4097.

Kadipasaoglu CM, Conner CR, Whaley ML, Baboyan VG, Tandon N (2016) Category-Selectivity in Human Visual Cortex Follows Cortical Topology: A Grouped icEEG Study Ben Hamed S, ed. PLoS One 11:e0157109.

Kay KN, Yeatman JD (2017) Bottom-up and top-down computations in word- and face-selective cortex. Elife 6:e22341.

Konkle T, Caramazza A (2013) Tripartite organization of the ventral stream by animacy and object size. J Neurosci 33:10235–10242.

Krauss GL, Fisher R, Plate C, Hart J, Uematsu S, Gordon B, Lesser RP (1996) Cognitive effects

of resecting basal temporal language areas. Epilepsia 37:476–483.

Kravitz DJ, Saleem KS, Baker CI, Ungerleider LG, Mishkin M (2013) The ventral visual

pathway: an expanded neural framework for the processing of object quality. Trends Cogn

Sci 17:26–49.

Kutas M, Hillyard SA (1980) Reading senseless sentences: Brain potentials reflect semantic

incongruity. Science (80-) 207:203–205.

Lachaux JP, George N, Tallon-Baudry C, Martinerie J, Hugueville L, Minotti L, Kahane P,

Renault B (2005) The many faces of the gamma band response to complex visual stimuli.

Neuroimage 25:491–501.

Lerma-Usabiaga G, Carreiras M, Paz-Alonso PM (2018) Converging evidence for functional and

structural segregation within the left ventral occipitotemporal cortex in reading. Proc Natl

Acad Sci U S A 115:E9981–E9990.

Leszczyński M, Barczak A, Kajikawa Y, Ulbert I, Falchier AY, Tal I, Haegens S, Melloni L,

Knight RT, Schroeder CE (2020) Dissociation of broadband high-frequency activity and

neuronal firing in the neocortex. Sci Adv 6:977–989.

Li Y, Richardson RM, Ghuman AS (2018) Posterior Fusiform and Midfusiform Contribute to

Distinct Stages of Facial Expression Processing. Cereb Cortex 29:3209–3219.

Lochy A, Jacques C, Maillard L, Colnat-Coulbois S, Rossion B, Jonas J (2018) Selective visual

representation of letters and words in the left ventral occipito-temporal cortex with

intracerebral recordings. Proc Natl Acad Sci 115:E7595–E7604.

Mani J, Diehl B, Piao Z, Schuele SS, Lapresto E, Liu P, Nair DR, Dinner DS, Lüders HO (2008) Evidence for a basal temporal visual language center: cortical stimulation producing pure alexia. Neurology 71:1621–1627.

Maris E, Oostenveld R (2007) Nonparametric statistical testing of EEG- and MEG-data. J Neurosci Methods 164:177–190.

Matsuo T, Kawasaki K, Kawai K, Majima K, Masuda H, Murakami H, Kunii N, Kamitani Y, Kameyama S, Saito N, Hasegawa I (2015) Alternating Zones Selective to Faces and Written Words in the Human Ventral Occipitotemporal Cortex. Cereb Cortex 25:1265–1277.

Miller KJ (2010) Broadband spectral change: Evidence for a macroscale correlate of population firing rate? J Neurosci 30:6477–6479.

Miller KJ, Schalk G, Hermes D, Ojemann JG, Rao RPN (2016) Spontaneous Decoding of the Timing and Content of Human Object Perception from Cortical Surface Recordings Reveals Complementary Information in the Event-Related Potential and Broadband Spectral Change Sporns O, ed. PLOS Comput Biol 12:e1004660.

Mo C, He D, Fang F (2018) Attention priority map of face images in human early visual cortex. J Neurosci 38:149–157.

Op de Beeck HP, Pillet I, Ritchie JB (2019) Factors Determining Where Category-Selective Areas Emerge in Visual Cortex. Trends Cogn Sci 23:784–797.

Otten LJ, Donchin E (2000) Relationship between P300 amplitude and subsequent recall for distinctive events: Dependence on type of distinctiveness attribute. Psychophysiology 37:644–661.

Price CJ, Devlin JT (2003) The myth of the visual word form area. Neuroimage 19:473–481.

Puce A, Allison T, Asgari M, Gore JC, McCarthy G (1996) Differential Sensitivity of Human
Visual Cortex to Faces, Letterstrings, and Textures: A Functional Magnetic Resonance
Imaging Study. J Neurosci 16:5205–5215.

Quian Quiroga R, Mukamel R, Isham EA, Malach R, Fried I (2008) Human single-neuron
responses at the threshold of conscious recognition. Proc Natl Acad Sci U S A 105:3599–
3604.

Ray S, Maunsell JHR (2011) Different Origins of Gamma Rhythm and High-Gamma Activity in
Macaque Visual Cortex Ungerleider L, ed. PLoS Biol 9:e1000610.

Saygin ZM, Osher DE, Norton ES, Youssoufian DA, Beach SD, Feather J, Gaab N, Gabrieli
JDE, Kanwisher N (2016) Connectivity precedes function in the development of the visual
word form area. Nat Neurosci 19:1250–1255.

Srihasam K, Vincent JL, Livingstone MS (2014) Novel domain formation reveals proto-
architecture in inferotemporal cortex. Nat Neurosci 17:1776–1783.

Stevens WD, Kravitz DJ, Peng CS, Tessler MH, Martin A (2017) Privileged functional
connectivity between the visual word form area and the language system. J Neurosci
37:5288–5297.

Stevens WD, Tessler MH, Peng CS, Martin A (2015) Functional connectivity constrains the
category-related organization of human ventral occipitotemporal cortex. Hum Brain Mapp
36:2187–2206.

Susilo T, Duchaine B (2013) Dissociations between faces and words: Comment on Behrmann

and Plaut. Trends Cogn Sci 17:545.

Susilo T, Wright V, Tree JJ, Duchaine B (2015) Acquired prosopagnosia without word

recognition deficits. Cogn Neuropsychol 32:321–339.

Tadel F, Baillet S, Mosher JC, Pantazis D, Leahy RM (2011) Brainstorm: a user-friendly

application for MEG/EEG analysis. Comput Intell Neurosci 2011:879716.

Tang H, Buia C, Madhavan R, Crone NE, Madsen JR, Anderson WS, Kreiman G (2014)

Spatiotemporal Dynamics Underlying Object Completion in Human Ventral Visual Cortex.

Neuron 83:736–748.

Thorpe S, Fize D, Marlot C (1996) Speed of processing in the human visual system. Nature

381:520–522.

Tsao DY, Moeller S, Freiwald WA (2008) Comparing face patch systems in macaques and

humans. Proc Natl Acad Sci U S A 105:19514–19519.

Vinckier F, Dehaene S, Jobert A, Dubus JP, Sigman M, Cohen L (2007) Hierarchical Coding of

Letter Strings in the Ventral Stream: Dissecting the Inner Organization of the Visual Word-

Form System. Neuron 55:143–156.

Weiner KS, Grill-Spector K (2010) Sparsely-distributed organization of face and limb

activations in human ventral temporal cortex. Neuroimage 52:1559–1573.

White AL, Palmer J, Boynton GM, Yeatman JD (2019) Parallel spatial channels converge at a

bottleneck in anterior word-selective cortex. Proc Natl Acad Sci U S A 116:10087–10096.

Zhen Z, Yang Z, Huang L, Kong X zhen, Wang X, Dang X, Huang Y, Song Y, Liu J (2015)

Quantifying interindividual variability and asymmetry of face-selective regions: A

probabilistic functional atlas. Neuroimage 113:13–25.

1 Table 1. iEEG participant coverage

Number	Tasks completed	Electrodes in VTC	Face-selective	Word-selective	House-selective	Word medial to face-selective	Alternating word- and face-selective
1	CL	L: 6	0	0	0	N/A	N/A
2	CL	L: 11	0	0	0	N/A	N/A
3	CL	L: 34, R: 18	0	0	L: 2, R: 2	N/A	N/A
4	CL	L: 20, R: 14	0	0	R: 2	N/A	N/A
5	CL	R: 18	R: 2	0	0	N/A	N/A
6 (Fig. 4B)	CL	L: 11	L: 1	L: 1	0	Yes	N/A
7	CL, WID	L: 17	L: 2	L: 1	0	No	No
8	CL	R: 9	0	0	R: 2	N/A	N/A
9	CL, WID	R: 21	0	R: 1	0	N/A	N/A
10 (Fig. 4B)	CL, WID, FID	L: 25, R: 16	L: 2, R:1	L: 2	0	Yes	Yes
11	CL, FID	L: 4, R: 23	R: 5	L: 1, R: 1	R: 5	Yes	Yes
12	CL, FID	R: 42	R: 8	R: 4	R: 6	Yes	Yes
13	CL, FID	L: 38	0	L: 2	L: 2	N/A	N/A
14	CL, FID	L: 23, R: 24	L:2, R: 1	0	L: 2, R: 2	N/A	N/A
15	CL, FID	L: 30	L: 1	0	L: 2	N/A	N/A
16	CL, FID	L: 23, R: 11	0	L: 1	R: 3	N/A	N/A
17 (Fig. 4B)	CL, WID, FID	L: 48	L: 6	L: 4	L: 2	Yes	Yes
18	CL, FID	L: 23	0	0	L: 7	N/A	N/A
19	CL	L: 4	0	L: 2	L: 2	N/A	N/A
20	CL	L: 23	0	0	0	N/A	N/A
21	CL	R: 11	0	0	R: 1	N/A	N/A
22	CL, WID, FID	R: 41	0	R: 6	0	N/A	N/A
23	CL	L: 10	L: 1	L: 2	0	No	No
24	CL, FID	L: 26, R: 25	L: 3, R: 1	R: 2	R: 1	Yes	Yes
25	CL, WID, FID	L: 21, R: 19	0	L: 6, R: 1	0	N/A	N/A
26	CL	L: 21, R: 28	L: 2	L: 3	R: 3	No	No
27	CL, FID	L: 5, R: 18	0	L: 1, R: 5	R: 3	N/A	N/A
28 (Fig. 4A)	CL, WID, FID	L: 55	L: 6	L: 4	0	Yes	Yes
29	CL, FID	L: 42	L: 2	L: 2	0	Yes	No
30	CL, FID	L: 26, R: 28	L: 1, R: 2	R: 1	L: 2, R: 1	Yes	No
31	CL, WID, FID	L: 19, R: 36	L: 1, R: 6	0	R: 2	N/A	N/A

32	CL, WID	L: 10, R: 34	L: 1	0	L: 1, R: 3	N/A	N/A
33	CL, WID, FID	L 39, R 50	0	L: 4	L: 3, R:2	N/A	N/A
34	CL, WID	L 24, R: 27	R: 2	L: 5, R: 2	L: 1, R: 3	No	No
35	CL, FID	L: 116	L: 16	L: 8	L: 6	Yes	Yes
36	CL, WID, FID	L: 129	L: 33	L: 15	L: 12	Yes	Yes
Total: 36	CL: 32, WID: 12, FID: 20	L 883, R: 513	L: 80, R: 28	L: 64, R: 23	L: 44, R: 41	L: 7/10, R: 4/5	L: 5/9, R: 3/5

Number of word-, face-, and house-selective electrodes in the left (L) and right (R) hemisphere

of each iEEG participant out of the total number of implanted VTC electrodes. All participants

underwent a category localizer task (CL) from which word, face, and house-selectivity was

determined by comparing electrode responses to six categories of objects (see *Materials and*

Methods and Figure 1B). The table indicates whether any word-selective electrodes were medial

to any face-selective electrodes in participants that had at least one word- and one face-selective

electrode within the same hemisphere. The table also indicates whether there was alternation of

word- and face-selective electrodes along the medial to lateral axis within participants that had at

least two word-selective electrodes and one face-selective electrode within the same hemisphere

or vice-versa. Participants with insufficient coverage word or face-selective regions to determine

their relative anatomical location are listed as not available (N/A). A subset of participants also

participated in a face individuation task (FID) or word individuation task (WID) from which the

individuation capacity of word- and face-selective electrodes was tested. Participants illustrated

in figures are noted next to the corresponding participant number.

Figure 1. Spatial topography of word- and face-selective iEEG electrodes

A) Heat map of electrode coverage (both category-selective and non-selective) across 36 iEEG participants. Electrodes below the inferior temporal sulcus and anterior to the posterior edge of the fusiform gyrus on the participant's native space were considered VTC. There was a greater proportion of left hemisphere coverage relative to right hemisphere coverage. B) Electrodes that responded preferentially to words, faces, or houses and could significantly discriminate these stimuli from all others using six-way Naïve Bayes classification ($p < 0.05$, Bonferroni corrected within participant). House-selective electrodes are primarily medial to word- and face-selective electrodes. Multiple adjoining word- and face-selective patches are found along the medio-lateral axis of ventral temporal cortex, bilaterally. Depth stereotactic EEG electrodes have been brought to the ventral surface for clarity.

1   Figure 2. Distribution of face-selective and word-selective electrodes by participant

3   Distribution of highly face-selective (left) and word-selective (right) electrodes by participant.

4   Color represents individual participants and corresponds across figure panels. Each group-level

5   cluster of word- and face-selective electrodes is represented by data from multiple participants.

Figure 3. Independence of word- and face-processing networks

A) Average decoding time-course for word- (top, n = 39) and face- (bottom, n = 75) selective

electrodes that contained significant amounts of information about the other object category. 21

out of 28 (75 %) face-selective electrodes in the right hemisphere and 54 out of 80 (66 %) in the

left hemisphere could significantly discriminate words from the other object categories excluding

faces (e.g. d' sensitivity for words was above chance for 5-way classification of the non-face

object categories) at the $p < 0.05$ level (Bonferroni corrected for multiple comparisons in time

and electrodes within participant, see *Materials and Methods*). 9 out of 23 (39 %) word-selective

electrodes in the right hemisphere and 30 out of 64 (47 %) in the left hemisphere could

discriminate faces from the other object categories excluding words. Error bars indicate standard

error from the mean across electrodes. Colored bars under the curves indicate times where the

average selectivity is above chance ($p < 0.001$ corrected for temporal comparisons). B) Average

decoding time-course for word- (top, n = 48) and face- (bottom, n = 33) selective electrodes that

did not contain above chance information for the other object category. Although decoding

accuracy was above chance at later time points for the non-preferred category across the

population of electrodes, decoding accuracy was much smaller for the non-preferred compared to

preferred category. C) Example decoding time courses from three highly word-selective

electrodes that did not display face-selectivity. D) Decoding time courses of three highly face-

selective electrodes that did not display word-selectivity. The patient from which the middle

electrode recording was obtained was not presented with pictures of hammers.

Figure 4. Multiple adjoining word- and face-selective patches in individual participants

A) Representative single participant demonstrated alternating bands of word- and face-selectivity

along the left fusiform gyrus. Shaded electrodes are those selective to words (yellow) and faces

(blue). Non-filled circles represent ventral temporal electrodes that did not reach the selectivity

criterion for either of these categories. Raw event-related broadband activity is shown for each of

the numbered electrodes in the right panel. Moving from medial to lateral, electrodes

demonstrate a preferential response to words, mixed response to both words and faces,

preferential response to faces then preferential response to words. Abbreviations: collateral

sulcus (COS), midfusiform sulcus (MFS), occipitotemporal sulcus (OTS). B) Three additional

examples of patients with multiple adjoining word- and face-selective regions or word-selectivity

medial to face-selectivity in VTC. Major VTC sulci have been labeled for clarity.

Figure 5. Interdigitation of BOLD responses to words and faces across eight healthy participants

Eight healthy participants that underwent an identical category localizer task as the iEEG

participants demonstrated similar category selectivity. A) Individual maps demonstrate word

versus face-selectivity across left VTC. In six out of eight of these participants there was strong

word-selectivity medial to face-selectivity along the midfusiform sulcus. The bar graphs below

each participant indicates the selectivity of these word-selective regions when defining them

based on one half of the data and testing on the other half of the data. Word-selective responses

were less consistent in the right hemisphere across participants. B) Bar graph of word-selectivity

in left hemisphere medial word-selective regions across participants for the left-out half of the

data. Symbols: ** $p < 0.01$, *** $p < 0.001$.

Figure 6. Comparing category-selectivity in single-trial potentials and high-frequency broadband

Differing spatial distribution of electrodes that demonstrated selectivity in single-trial potentials

(stP) and single-trial high-frequency broadband activity (stHFBB). A) Electrodes that

demonstrated selectivity in both stP and stHFBB were clustered around the fusiform and

parahippocampal gyri. B) Electrodes selective in only stP were much more widely distributed

and included medial and anterior word-selective regions not typically seen in fMRI. C)

Electrodes that were only selective in stHFBB were less prevalent than those only selective in

stP, but also extended outside of the fusiform and parahippocampal gyri.

Figure 7. Differences in the dynamics of left versus right word- and face-selective regions

Latency of word (yellow) and face (blue) sensitivity onset, latency of peak sensitivity, and

magnitude of peak sensitivity across hemispheres. Latency of sensitivity onset is defined as the

first timepoint that reached a d' corresponding to $p < 0.001$ non-parametrically defined by the

pre-stimulus baseline period. All time points reference the leading edge of the classification

window. Box plots represent median, 25th and 75th percentiles. Summary statistics of each box

plot are presented in the table. Abbreviations: confidence interval (CI), degrees of freedom (d.f.),

left (L), right (R). Symbols: n.s. $p > 0.05$, * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.

Figure 8. Spatiotemporal clustering of word- and face-selective regions

A) Spatiotemporal clustering of word-selective VTC electrodes. The illustrated clustering

solution was robust to different weightings of spatial and temporal information. Left hemisphere

word-selective electrodes were clustered into four spatial clusters. A cluster was found on the

fusiform gyrus (gray, 21 electrodes from 5 participants), as well as medial (green, 20 electrodes

from 10 participants), anterior (blue, 11 electrodes from 5 participants), and lateral (red, 12

electrodes from 7 participants) to the fusiform gyrus. Right hemisphere word-selective regions

had later onsets and were more clearly separated along the posterior to anterior axis (posterior:

cyan; 8 electrodes from 4 participants, mid: yellow; 8 electrodes from 3 participants, anterior:

magenta; 7 electrodes from 6 participants). B) Average d' timecourse of each group of electrodes

in A when jointly classifying stP and stHFBB. Error bars represent standard error across

electrodes. C) Average d' time course of each group of electrodes when classifying only stP. D)

Average d' time course of each electrodes when classifying only stBB. Word-selective

electrodes on the fusiform demonstrate strong selectivity in both stP and stHFBB, whereas other

regions display distinct dynamics across these signal components. E) Spatiotemporal clustering

of face-selective VTC electrodes. Left hemisphere electrodes were clustered into three spatial

clusters roughly posterior to (green, 21 electrodes from 3 participants), on (gray, 46 electrodes

from 12 participants), and anterior to the fusiform gyrus (blue, 13 electrodes from 7 participants).

Right hemisphere, face-selective electrodes were primarily clustered along the posterior to

anterior VTC axis into four clusters (posterior: cyan; 9 electrodes from 5 participants, mid:

yellow; 13 electrodes from 6 participants and black; 3 electrodes from 2 participants, anterior:

magenta; n = 3 electrodes from 3 participants). F) Average d' time course of each group of

electrodes illustrated in E when jointly classifying stP and stHFBB. G) Average d' timecourse of

each group of electrodes when classifying only stP. H) Average d' time course of each group of

electrodes when classifying only stBB. Face-selective electrodes on the fusiform demonstrate

strong selectivity in both stP and stHFBB, whereas other regions display distinct dynamics

across these signal components.

A

Anterior

Right ⟷ Left

Posterior

total number of electrodes

B

Face
Word
House

face-selective electrodes

word-selective electrodes

A Word-selective electrodes with significant face information

Face-selective electrodes with significant word information

B Word-selective electrodes without significant face information

Face-selective electrodes without significant word information

Legend (panels A, B): Face, Word

Legend (panels C, D): face, word, hammer, house, scrambled

Axis labels: sensitivity (d′), Time (ms)

word-selective

A   B stP & stHFBB

C stP   D stHFBB

face-selective

E   F stP & stHFBB

G stP   H stHFBB