Human blood contains circulating cell-free mitochondria, but are they really

functional?

Antoine Stier [a,b]

[a] Department of Biology, University of Turku, Turku, Finland

[b] Institute of Biodiversity, Animal Health and Comparative Medicine, University of Glasgow,

Glasgow, UK

*Corresponding author: amstie@utu.fi / antoine.stier@gmail.com, Natura Building room 343

20500 Turku, Finland. +447778208723

Running title: Circulating cell-free mitochondrial bioenergetics

Abstract

Dache et al. (2020, *FASEB J.* **15**, e2002338–15) recently reported the presence of *respiratory-competent* cell-free mitochondria in human blood (up to 3.7 x 10^6 per mL of blood), providing exciting perspectives on the potential role of these extra-cellular mitochondria. While their evidence for the presence of cell-free mitochondria in human blood is compelling, their conclusion that these cell-free mitochondria are respiratory-competent or functional has to be re-evaluated. To this end, we evaluated the functionality of cell-free mitochondria in human blood using high-resolution respirometry and mitochondria extracted from platelets of the same blood samples as positive controls. While cell-free mitochondria were present in human plasma (*i.e.* significant complex IV activity), there was no evidence suggesting that their mitochondrial electron transport system (ETS) was functional (*i.e.* respiration rate not significantly different from 0; no significant responses to ADP, uncoupler or mitochondrial inhibitors oligomycin and antimycin A). Yet, *in vitro* complex IV activity was detectable and even slightly higher than levels found in mitochondria extracted from platelets, suggesting that cell-free mitochondria in human blood only retain a non-functional part of the electron transport system. Despite being unlikely to be fully functional in the narrow-sense (*i.e.* capable of oxidative phosphorylation), circulating cell-free mitochondria may have significant physiological roles that remain to be elucidated.

Keywords: mitochondrial function, blood cells, platelet, extra-cellular mitochondria

Introduction

Mitochondria are the powerhouse of eukaryotic cells using oxidative phosphorylation to convert nutrients into ATP. They also have key roles in cell signalling (1) and ageing (2). Vertebrates possess functional mitochondria in every cell type, with the notable exception of mammalian mature erythrocytes (3). Dache et al. (2020) recently reported the presence of cell-free mitochondria in human blood, providing exciting perspectives on the role of these extra-cellular mitochondria, and their potential to be used as prognosis tool for various diseases (4). While their evidence for the presence of cell-free mitochondria in human blood is compelling and has been supported by other studies (*e.g.* (5, 6)), their conclusion that these cell-free mitochondria are respiratory-competent or functional has to be re-evaluated (7) since this is critical to uncover the physiological role they may or may not play (8).

Indeed, the interpretation that circulating cell-free mitochondria are functional is exclusively based on the authors' graphical interpretation of *in vitro* O_2 consumption rates (Fig. 7 in (4)), and not on statistical testing considering their sample size was N = 1 (despite having n = 3 replicates of the same pooled sample; Dache et al. *pers. com.*). Based on the data being presented in (4), it seems impossible to conclude that circulating cell-free mitochondria are functional, as recently highlighted by (7). The *in vitro* respirometry assay used in (4) to assess functionality has several limitations. First, the addition of ascorbate and TMPD to test complex IV maximal capacity indeed elicits a rise in O_2 consumption, but no negative control is shown by the authors, while there is a known chemical O_2 consumption linked to the auto-oxidation of ascorbate and TMPD (9). Second, there is no visible inhibition of O_2 consumption by antimycin A (*i.e.* a complex III inhibitor), suggesting that the minor O_2 consumption being observed is unlikely to be of mitochondrial origin. Third, the role of mitochondria is not to consume O_2 *per se*, but to produce ATP, and no test of oxidative phosphorylation functionality

(*e.g.* stimulation of respiration by ADP addition, or inhibition of respiration by oligomycin

addition, an inhibitor of ATP synthase) is provided (7).

Two alternative explanations could explain the apparent lack of functionality of these

circulating cell-free mitochondria. It is possible that circulating cell-free mitochondria are not

functional because they are damaged, or that only a small fraction of them (*i.e.* potentially the

ones that arrived in the blood stream only very recently) are still somewhat functional.

Mitochondrial integrity might be evaluated to some extent by the addition of exogenous

cytochrome c during respirometry assay. Cytochrome c does not penetrate intact

mitochondria but penetrates within partially damaged ones eliciting a rise in O_2 consumption

(10). Alternatively, it is possible that circulating cell-free mitochondria are functional, but that

the *in vitro* respirometry assay of (4) was not well suited for these mitochondria (7). Indeed,

the mitochondrial uncoupler FCCP has been added directly to the respiration medium at a

concentration of 4μM. However, uncouplers like FCCP have a narrow optimal working range

and require a progressive titration (*e.g.* 0.5μM steps; (10)). Indeed, when added in excess,

uncouplers can partially to fully inhibit mitochondrial respiration, and partially damaged

mitochondria are likely to be more sensitive to such phenomenon than intact ones (10).

Consequently, it is possible that the cell-free circulating mitochondria observed in (4) were

functional, but that they were fully inhibited by an inappropriate amount of mitochondrial

uncoupler in the respiratory medium.

A proper assessment of the functionality of circulating cell-free mitochondria is

therefore required to establish if they are really functional (7), or potentially non-functional

remnants of mitochondria that have been extruded by cells in the blood stream. In this study,

we aim to provide elements to answer this question by characterizing mitochondrial

bioenergetics properties of circulating cell-free mitochondria using high-resolution

respirometry and mitochondria extracted from platelets of the same blood samples as positive

controls.

Methods

<u>Sample collection</u>

This study was performed in accordance with the WMA Declaration of Helsinki. Blood

samples were collected at the University of Turku by a trained and habilitated lab technician

from six healthy volunteers (2 males and 4 females) following signature of an informed

consent form. Approximately 80mL of blood was taken from each volunteer using BD

vacutainer K_2EDTA tubes, which were progressively cooled and stored on ice before

processing (max. 1 hour).

<u>Sample processing</u>

We used a protocol very similar to the one used by (6). Blood samples were first

centrifugated at 150g for 20min (4°C) to isolate platelet-rich plasma (PRP) from other blood

cells. PRP was subsequently centrifugated at 2500g for 30min (4°C) to pellet platelets and

obtain cell-free plasma. Mitochondria were isolated from platelets pellet following (11), with

4 cycles of 10 passes in glass/glass Potter-Elvehjem pestle homogenizer and 2500g

centrifugation (4°C, 1min) to re-form platelet pellet and facilitate cellular membrane

disruption by homogenization. Platelet samples were then centrifugated at 800g (4°C, 10min)

and the supernatant containing isolated mitochondria was subsequently processed along with

cell-free plasma. Both sample types were centrifugated at 10000g (4°C, 10min) to pellet

mitochondria. Mitochondrial pellets were resuspended in 2mL of Mir05 respiration buffer (0.5

mM EGTA, 3 mM $MgCl_2$, 60 mM K-lactobionate, 20 mM taurine, 10 mM KH_2PO_4, 20 mM Hepes,

110 mM sucrose, free fatty acid bovine serum albumin (1 g/L)), and subsequently washed by

centrifugation at 10000g (4°C, 10min). Mitochondrial pellets were finally resuspended in

100μL of Mir05. Mitochondrial proteins were quantified using the Biuret method, accounting

for the fact that Mir05 contains 1mg.mL^{-1} of BSA.

<u>High-resolution respirometry assay</u>

Mitochondrial respiration was assessed using the high-resolution respirometry system

O2k (Oroboros Instruments, Innsbruck, Austria) at 37°C using *ca.* 100μg (range: [75 ; 125]) of

mitochondrial protein per assay (for comparison, 64μg of mitochondrial protein was used in

(4)). Mitochondria were added to the respirometry chambers containing 2mL of Mir05 pre-

equilibrated at 37°C and chambers were closed after 2min of equilibration. A substrate-

uncoupler-inhibitor titration (SUIT) protocol was used to assess the functionality of different

components of the mitochondrial electron transport system. Substrates of complex I (P:

pyruvate 5mM and M: malate 2mM) were first added, followed by a saturating amount of ADP

(2mM). Substrate of complex II (S: succinate 10mM) was then added to stimulate

mitochondrial respiration fueled by both complexes I and II. Cytochrome c (Cyt c: 10μM) was

added to test potential issues linked to mitochondrial inner membrane integrity. ATP synthesis

was then inhibited with oligomycin (O: 2.5μM). A progressive titration (0.5μM steps) of the

uncoupler FCCP was subsequently conducted until no further stimulation could be observed

(maximum stimulation was reached at *ca.* 1.5μM), at which stage mitochondrial respiration

was fully inhibited using the complex III inhibitor antimycin A (AA: 2.5μM). Finally, the maximal

activity of complex IV was assessed using ascorbate (Asc: 2mM) and N,N,N',N'-tetramethyl-p-

phenylenediamine (TMPD: 0.5mM). Since there is a known chemical O_2 consumption linked

to the auto-oxidation of ascorbate and TMPD (9), 2 negative control samples (*i.e.* 50μL of

Mir05 instead of mitochondria) were analyzed following the full SUIT protocol, so chemical O_2

consumption could be subtracted from ascorbate+TMPD values of biological samples (see Fig.

1A).

Data analysis

Oxygen consumption rates ($pmol.s^{-1}.mL^{-1}$) were extracted using DatLab 7.0 (Oroboros

Instruments, Innsbruck, Austria) after correction for background noise. Responses to SUIT

protocol of both plasma and platelet mitochondria were analyzed using a linear mixed model

with sample type (cell-free *vs.* platelet), stage (P+M, ADP, S, Cyt C, O, FCCP, AA, Asc+TMPD)

and their interaction as fixed factor; while controlling for pseudo-replication by including

stage(ID) as a random factor to account for the fact that plasma and platelet samples came

from the same individual (ID), as well as that measurements were repeated over the same

samples in the different stages of the SUIT protocol. Since a significant interaction between

sample type and stage was detected (see results), data was subsequently analyzed for each

sample type individually to assess the differences between stages using Bonferroni-corrected

post-hoc tests.

Mitochondrial respirations rates were calculated by subtracting non-mitochondrial

oxygen consumption (after AA inhibition) to specific stages ($OXPHOS_{CI+II}$ = S - AA; $LEAK_{CI+II}$ = O

- AA; ETS_{CI+II} = FCCP - AA), and by subtracting chemical O_2 consumption (*i.e.* obtained from

negative controls, see Fig 1A) to Asc+TMPD values (COX = Asc+TMPD - AA - chemical O_2

consumption). Those values were corrected by the quantity of mitochondrial protein used in

each assay and therefore expressed as $pmol.s^{-1}.mg^{-1}$. Differences between cell-free and

platelet mitochondria for each parameter were analyzed using non-parametric Wilcoxon

paired tests, and differences to 0 of each parameter were tested using non-parametric one-

way Wilcoxon tests. All analyses were conducted using SPSS 25.0, p-values < 0.05 were

considered significant and values are expressed as mean ± 95% C.I.

Results

Responses to SUIT protocol of both cell-free and platelet mitochondria, as well as

negative controls, are presented in Fig. 1A. There was a significant interaction between sample

type and stage in the way mitochondria responded to the SUIT protocol ($F_{7,40}$ = 16.99, p <

0.001). Platelet mitochondria responded in the expected way to the SUIT protocol ($F_{7,8.1}$ =

84.4, p < 0.001; Fig. 1A), with an increase in O_2 consumption following ADP addition (P+M *vs.*

ADP: p = 0.012), stimulation by complex II substrate S (ADP *vs.* S: p = 0.005), non-significant

effect of Cyt c addition (S *vs.* Cyt c: p = 1.00), a decrease in O_2 consumption following ATP

synthase inhibition (Cyt c *vs.* O: p = 0.042), stimulation by FCCP titration (O *vs.* FCCP: p = 0.028),

full inhibition by complex III inhibitor AA (FCCP *vs.* AA: p = 0.007) and massive stimulation by

fueling directly complex IV with electrons (AA *vs.* Asc+TMPD: p < 0.001). On the contrary,

although cell-free mitochondrial O_2 consumption was affected by SUIT stage ($F_{7,7.9}$ = 159.1, p

< 0.001; Fig. 1A), they did not significantly respond to the SUIT protocol (P+M *vs.* ADP: p =

1.00; ADP *vs.* S: p = 1.00; S *vs.* Cyt c: p = 1.00; Cyt c *vs.* O: p = 1.00; O *vs.* FCCP: p = 1.00; FCCP

vs. AA: p = 1.00), except for the massive stimulation obtained by fueling directly complex IV

with electrons (AA *vs.* Asc+TMPD, p < 0.001).

While mitochondrial respiration rates of platelet mitochondria were all significantly

different from 0 (all p < 0.03; Fig. 1B), this was not the case for cell-free mitochondria

(*OXPHOS$_{CI+II}$*: p = 0.60; *LEAK $_{CI+II}$*: p = 0.60; *ETS $_{CI+II}$*: p = 0.25), except for complex IV respiration

(*COX*: p = 0.028). Mitochondrial respiration rates of platelet mitochondria were significantly

higher than cell-free ones (*OXPHOS$_{CI+II}$*, *LEAK $_{CI+II}$* and *ETS $_{CI+II}$*: all p < 0.03; Fig 1B), except for

complex IV respiration (*i.e. COX*), for which cell-free mitochondria actually had a slightly higher

respiration rate than platelet ones (*COX*: p = 0.046; Fig. 1B).

Fig. 1: *Mitochondrial bioenergetics properties of human circulating cell-free mitochondria and mitochondria isolated from platelets. (A) Responses to a typical substrate-uncoupler-inhibitor protocol evaluated by high-resolution respirometry assay (see methods for full details, and results for information on statistics). Negative control samples consisted of 50μL of respiration buffer instead of mitochondria. While platelets exhibited a typical response to SUIT protocol (all p < 0.05), cell-free mitochondria only responded to Asc+TMPD (p < 0.001, all other p = 1.00). (B) Mitochondrial respiration rates of platelet vs. cell-free mitochondria. Significant difference from 0 for each parameter is indicated by an *. All mean values are plotted ± 95% C.I., N = 6 in every case except for negative controls (N = 2).*

Discussion

As noted by (7), the evidence provided by (4) was insufficient to infer the functionality

or the respiratory competence of human circulating cell-free mitochondria. By conducting a

comprehensive analysis of mitochondrial bioenergetics of human circulating cell-free

mitochondria in direct comparison to mitochondria extracted from platelets of the same

blood samples, we found no significant support for the functionality or the respiratory

competence of those cell-free mitochondria. Indeed, there was no significant evidence that

cell-free mitochondria possess a functional electron transport system since they exhibited no

responses to mitochondrial substrates, ADP or inhibitors. Yet, when fueling directly complex

IV with exogenous electrons (*i.e.* Asc+TMPD), a significant mitochondrial O_2 consumption was

detected (similarly to (4), except that no negative control was provided in their study). This

shows that cell-free mitochondria retain a small fraction of the ETS, but that they are unlikely

to be functional or respiratory competent *in vivo* since electrons have to enter the ETS through

complex I and/or II. In contrast, mitochondria isolated from platelets show the typical

response to the SUIT protocol. Mitochondrial quantity per assay is unlikely to be a bias in the

data being presented here since cell-free mitochondria even had a slightly higher complex IV

activity (sometimes used as a proxy of mitochondrial quantity (12)) than mitochondria isolated

from platelets. The absence of narrow-sense functionality (*i.e.* capability of oxidative

phosphorylation) of circulating cell-free mitochondria suggested by our results is a key point

regarding the identification of their potential physiological role(s). For instance, (8)

hypothesized that extracellular mitochondria could help in restoring homeostasis by

accumulating at sites of energy deficits, which is unlikely to be true if those mitochondria have

no or an extremely low capacity for oxidative phosphorylation.

Yet, using flow-cytometry fluorescence assays, (6) recently reported that at least part

of the cell-free mitochondria of human and murine blood could remain functional. Indeed,

part of the cell-free mitochondria identified by flow cytometry and MitoTracker Green

fluorescence appear capable of maintaining a mitochondrial membrane potential since they

retained TMRE probe, and such retention was lowered by the addition of the mitochondrial

uncoupler FCCP (6). Considering that the isolation protocol for cell-free mitochondria is likely

to result in many microparticles not containing 'intact' mitochondria (6), one hypothesis could

be that mitochondrial quantity was too low in our study to enable the detection of significant

functionality by high-resolution respirometry. Yet, considering the estimated density of

functional cell-free mitochondria provided by (6) (*i.e.* 1.4×10^6/mL of blood), the samples used

in the present study should have contained *ca.* 110×10^6 intact extracellular mitochondria.

Considering that human platelets contain on average 4.9 mitochondria per cell (13),

mitochondrial respiration of our cell-free mitochondria samples should be approximately

similar to the respiration of 22×10^6 platelets. Mitochondrial respiration of human platelets

has been reported by others to be around 0.35 pmol.s^{-1}.10^6 cells^{-1} (*OXPHOS$_{CI+II}$* in (14)).

Thereby, if cell-free mitochondria from human blood had respiration rates in the same range

as platelet mitochondria, we would expect our cell-free mitochondria to exhibit a respiration

of *ca.* 7.6 pmol.s^{-1}.mL^{-1}, which is approximately 8 times above the detection limit of our high-

resolution respirometry assay (*e.g. LEAK$_{CI+II}$* of platelets was 7.3 ± 2.4 in this study).

Although fully rejecting the hypothesis that human circulating cell-free mitochondria

are functional would require a way larger sample size (*e.g.* N > 90 for reaching a statistical

power of 0.80 for *OXPHOS$_{CI+II}$* based on data presented in Fig. 1B), our results do not support

the functionality of circulating cell-free mitochondria. Further studies should be performed to

further ascertain the functionality or lack of functionality of circulating cell-free mitochondria

since this appears critical as to elucidate their physiological role(s) (*e.g.* (8)).

Acknowledgement

AS is grateful to the authors of (4, 6, 7) for helping him to find motivation to conduct this study

following the rejection without peer-review of his initial commentary on (4). AS is also grateful

to Tiina Hentinen and Hekki Minn for helping to organize the study, to Riikka Sjöroos for

collecting blood, to the six volunteers for giving their blood, as well as to the 'Turku Collegium

for Science and Medicine' for funding his research, and to the CafR team of La Piclière La Clusaz

for hosting him while writing the initial commentary.

Conflict of interest

The author declares having no conflict of interest.

Author contribution

AS designed the study and conducted all the laboratory work, statistical analysis and writing

of the manuscript.

References

1.   Bohovych, I. and Khalimonchuk, O. (2016) Sending Out an SOS: Mitochondria as a
Signaling Hub. *Front. Cell Dev. Biol.* **4**, 524–15

2.   Sun, N., Youle, R. J., and Finkel, T. (2016) The Mitochondrial Basis of Aging. *Molecular*
Cell **61**, 654–666

3.   Stier, A., Reichert, S., Criscuolo, F., and Bize, P. (2015) Red blood cells open promising
avenues for longitudinal studies of ageing in laboratory, non-model and wild animals.
Experimental Gerontology **71**, 118–134

4. Dache, Al, Z., Otandault, A., Tanos, R., Pastor, B., Meddeb, R., Sanchez, C., Arena, G.,
Lasorsa, L., Bennett, A., Grange, T., Messaoudi, El, S., Mazard, T., Prevostel, C., and
Thierry, A. R. (2020) Blood contains circulating cell-free respiratory competent
mitochondria. *FASEB J.* **15**, e2002338–15

5. Song, X., Hu, W., Yu, H., Wang, H., Zhao, Y., Korngold, R., and Zhao, Y. (2020) Existence
of Circulating Mitochondria in Human and Animal Peripheral Blood. *IJMS* **21**, 2122–12

6. Stephens, O. R., Grant, D., Frimel, M., Wanner, N., Yin, M., Willard, B., Erzurum, S. C.,
and Asosingh, K. (2020) Characterization and origins of cell-free mitochondria in
healthy murine and human blood. *Mitochondrion* **54**, 102–112

7. Bertero, E., O'Rourke, B., and Maack, C. (2020) Response by Bertero et al to Letter
Regarding Article, "Mitochondria Do Not Survive Calcium Overload". *Circulation*
Research **126**, 5069–2

8. Esch, T., Stefano, G. B., Ptacek, R., and Kream, R. M. (2020) Emerging Roles of Blood-
Borne Intact and Respiring Mitochondria as Bidirectional Mediators of Pro- and Anti-
Inflammatory Processes. *Med Sci Monit* **26**, e924337

9. Kuznetsov, A. V. and Gnaiger, E. (2010) Oxygraph assay of cytochrome c oxidase
activity: chemical background correction. *Mitochondr Physiol Netw* **6**, 1–4

10. Lanza, I. R. and Nair, K. S. (2009) Chapter 20 - Functional Assessment of Isolated
Mitochondria In Vitro. In *Mitochondrial Function, Part B: Mitochondrial Protein*
Kinases, Protein Phosphatases and Mitochondrial Diseases vol. 457, pp. 349–372,
Elsevier Inc.

11. Frezza, C., Cipolat, S., and Scorrano, L. (2007) Organelle isolation: functional
mitochondria from mouse liver, muscle and cultured filroblasts. *Nat Protoc* **2**, 287–
295

12. Larsen, S., Nielsen, J., Hansen, C. N., Nielsen, L. B., Wibrand, F., Stride, N., Schroder, H.
D., Boushel, R., Helge, J. W., Dela, F., and Hey-Mogensen, M. (2012) Biomarkers of
mitochondrial content in skeletal muscle of healthy young human subjects. *The*
Journal of Physiology **590**, 3349–3360

13. Shuster, R. C., Rubenstein, A. J., and Wallace, D. C. (1988) Mitochondrial DNA in
anucleate human blood cells. *Biochemical and biophysical research communications*
155, 1360–1365

14. Sjövall, F., Ehinger, J. K. H., Marelsson, S. E., Morota, S., Åsander Frostner, E., Uchino,
H., Lundgren, J., Arnbjörnsson, E., Hansson, M. J., Fellman, V., and Elmér, E. (2013)
Mitochondrial respiration in human viable platelets—Methodology and influence of
gender, age and storage. *Mitochondrion* **13**, 7–14

315