MSM and sesame seed oil improve hyperlipidemia and PUFA metabolism in the *db/db* mouse

Cameron V. Fili[1], Ling Lin[2,3], Jonathan Chapman[2], David Hamilton[1], Charles R. Yates[4*]

[1] Department of Comparative Medicine, University of Tennessee College of Medicine, Memphis,

TN, USA

[2] Department of Pharmaceutical Sciences, University of Tennessee College of Pharmacy,

Memphis, TN, USA

[3] Regenerative Medicine Research Center, West China Hospital, Sichuan University, Chengdu,

Sichuan 610041, P.R. China

[4] National Center for Natural Products Research, University of Mississippi, Oxford, MS, USA

*Corresponding author

Email: cryates2@olemiss.edu

Abstract

Background: An estimated 463 million people worldwide suffer from diabetes, and that number is projected to rise significantly to 700 million by 2045. One of the hallmarks of type 2 diabetes and metabolic syndrome is alterations in the lipid profile and polyunsaturated fatty acid metabolism.

Objective: The objective of this study was to identify lipid alterations in leptin receptor deficient BKS.Cg-*Dock7^m +/+ Lepr^{db}*/J (*db/db*) mice, a murine model of type 2 diabetes, for the purpose of establishing a baseline biological signature for the subsequent evaluation of natural products with purported lipid-altering activity.

Methods: Six-week old male *db/db* mice (n = 10/group) were randomized to the following groups: 1) diabetic control with no treatment, 2) methylsulfonylmethane (MSM) treatment (3.81 $\pm$ 0.33 g/kg), 3) sesame seed oil (SSO) treatment (23.54 $\pm$ 2.91 mg/kg), and 4) MSM and SSO combination treatment. Eight-week old male C57BL6/J mice (n=10) were used as a non-diabetic control group.

Results: Serum triglycerides and total cholesterol were significantly increased in the *db/*db model compared to nondiabetic control, mimicking the diabetic condition in people. HDL-cholesterol was significantly increased in all *db/db* treatment groups, with the most significant treatment effect in the MSM and SSO combination group, with a corresponding decrease in non-HDL cholesterol (LDL and VLDL). With regard to fatty acid metabolism, serum total polyunsaturated fatty acid (PUFA) levels were significantly reduced in diabetic mice compared to control mice. In contrast, feeding of only SSO reversed this effect such that fed mice exhibited serum PUFA levels comparable to control mice.

40   **Conclusions**: Treatment of *db/db* mice with MSM and SSO improved commonly measured

41   clinical parameters in serum lipid panels. The combination of MSM and SSO treatment's effects

42   on HDL and non-HDL cholesterol and fatty acid metabolism could lead to improved clinical

43   outcomes in people such as reduced incidence of atherosclerosis and hepatic steatosis.

Introduction

The International Diabetes Federation estimates there are at least 463 million people (1 in 11 adults aged 20-79 years) worldwide suffering from diabetes, and that number is projected to rise significantly to 700 million by 2045 [1]. One of the hallmarks of type 2 diabetes mellitus and related metabolic syndrome and obesity is dyslipidemia and impaired lysophospholipid metabolism [1, 2]. While most type 2 diabetic patients require pharmaceutical intervention such as insulin therapy and peroxisome proliferator-activated receptor (PPARγ) agonists, changes in the diet may also improve dyslipidemia and the diabetic condition [3]. An important contributor to the rise in the number of diabetic patients is the increasing rates of obesity due to the Modern Western Diet. In the past three decades, total fat and saturated fat intake as a percentage of total calories has continuously increased in Western diets, while the intake of omega-6 fatty acids increased and omega-3 fatty acids decreased. This shift has resulted in a large increase in the omega-6: omega-3 ratio from 1:1 during evolution to 20:1 today or even higher [4]. This high omega-6: omega-3 ratio is directly linked to chronic inflammatory diseases including obesity, metabolic syndrome, and type 2 diabetes mellitus [5].

Current anti-diabetic therapeutic strategies, e.g., insulin, metformin, and PPARγ agonists, are employed once an individual has developed type 2 diabetes. A more recent approach to controlling the diabetes epidemic is to elucidate the mechanisms of pre-diabetes as a means to understand factors that contribute to conversion of pre-diabetes to diabetes. Pre-diabetes, or impaired glucose tolerance and insulin resistance prior to onset of type 2 diabetes, has been historically diagnosed based on glycemic criteria including hemoglobin A1c of 5.7-6.4%, fasting blood glucose of 100-125 mg/dL, and blood glucose of 140-199 mg/dL two hours post oral glucose

tolerance test [1, 6]. Early detection of the pre-diabetic condition and insulin resistance (IR) can

lead to intervention and prevention of the development of type 2 diabetes [1]. The recent

development of tests such as the Quantose^{TM} Insulin Resistance test, which measures biomarkers

of insulin sensitivity other than blood glucose and A1c, can detect pre-diabetes even earlier in the

disease process than these standard glycemic markers [7]. One biomarker included in the

Quantose^{TM} IR is linoleoylglycerophosphocholine (LGPC), also known as

lysophosphatidylcholine-linoleic acid (LPC-LA, 18:2), which is a representative omega-6 fatty

acid that decreases in people with pre-diabetes [6].

The dietary ingredient sesamin, a phytonutrient of the lignans class found abundantly in

sesame seed oil, is known to decrease formation of the pro-inflammatory omega-6 metabolite

arachidonic acid in the liver by inhibiting the D-5 desaturase activity. This effect results in

enhanced liver formation of EPA and DHA relative to arachidonic acid [8]. Sesamin's impact on

the PUFA balance is limited by the fact that the liver inefficiently delivers EPA and DHA to the

systemic circulation and thus the tissues in the setting of diabetes. This occurs because a key

contributor to PUFA release from the liver is the presence of methyl donors such as B12 and folic

acid, which are significantly reduced in diabetes [9]. Methylsulfonylmethane, with the chemical

formula $(CH_3)_2SO_2$, also known as MSM, DMSO2, methyl sulfone, or dimethyl sulfone, is a

naturally-derived sulfur compound found in many plants and food sources and a putative methyl

donor [7, 8]. MSM provides remarkable defense against oxidative stress and inflammatory injuries

[9, 10]. The objective of this research was to investigate whether supplementation with sesame

seed oil, MSM, or the combination of sesame seed oil and MSM will improve the dyslipidemia

seen in the diabetic condition, and thus reverse the pro-inflammatory environment found in an

animal model of type 2 diabetes mellitus, the leptin receptor-deficient db/db mouse.

Materials and methods

Animals and housing

Six-week old male leptin receptor deficient BKS.Cg-$Dock7^m$ +/+ $Lepr^{db}$/J (hereafter referred to as *db/db*) mice were purchased from The Jackson Laboratory as a model for type 2 diabetes mellitus. This strain is leptin receptor deficient; therefore, leptin is unable to bind and produce satiety, resulting in development of obesity and type 2 diabetes mellitus. Eight-week old male C57BL6/J mice (n=10) were used as a non-diabetic control group. Nondiabetic mice (n=10) received no treatment, and diabetic mice were divided into four experimental groups (n=10 per group): 1) diabetic control with no treatment, 2) MSM treatment, 3) sesame seed oil treatment, and 4) MSM and sesame seed oil combination treatment. Upon arrival prior to treatments, mice were placed in individually ventilated caging at a housing density of 5 mice per cage on corncob bedding (Shepherd Specialty Papers, Watertown, TN) with a 12:12 light: dark cycle. Chlorinated water was delivered in polysulfone bottles (Alternative Design, Siloam Springs, AR) ad libitum. Mice were fed Tekland 7912 rodent diet ad libitum (Envigo, Madison, WI). Animal work was performed in an American Association for Accreditation of Laboratory Animal Care (AAALAC) accredited animal facility and approved by the University of Tennessee Health Science Center Institutional Animal Care and Use Committee (Protocol 17-095.0).

Diabetes confirmation and treatments

One week after arrival, non-fasted blood glucose was measured to confirm hyperglycemia in the *db/db* mice. Blood glucose was measured by handheld glucometer (FreeStyle Lite) on a drop of blood obtained from the tail vein with the mouse in a plastic mouse restraining device. Mice

with blood glucose readings greater than 250 mg/dL were considered diabetic. Records were

maintained for daily mouse weight, water intake, and food intake, and weekly blood glucose levels.

Criteria for early euthanasia was 20% weight loss from start of experiment. After confirmation of

diabetes, *db/db* mice then began treatment with either MSM alone, sesame seed oil alone (SSO),

or a combination of MSM and SSO for four weeks prior to euthanasia. MSM provided by

Bergstrom Nutrition was dissolved in Millipore filtered water in a polysulfone bottle (Alternative

Design, Siloam Springs, AR) at one percent v/v. Tekland 7912 feed was soaked in sesame seed

oil (Sigma Aldrich) for five minutes and then blotted dry with an absorbent paper towel.

Nondiabetic C57BL6/J mice and diabetic control *db/db* mice continued to receive unaltered 7912

diet and Millipore filtered water in polysulfone bottles for four weeks prior to euthanasia.

All drug solutions were sterile filtered using a 0.22-micron filter prior to administration.

Blood was collected via cardiocentesis under isoflurane anesthesia at euthanasia, and then

centrifuged at 2500 rpm for fifteen minutes for serum separation. Serum was aliquoted for Rodent

Lipid Profiles and fatty acid analysis via LC-MS/MS. Samples of 250 uL of serum per mouse were

mailed on ice overnight to IDEXX BioAnalytics for Rodent Lipid Profile measurement, including

cholesterol, high-density lipoprotein, low-density lipoprotein, and triglyceride levels. At

euthanasia, tissues (eyes, brain, and liver) were harvested for fatty acid analysis using LC-

MS/MS. Sections of liver were preserved in formalin, cryosectioned, and stained with Oil Red O

for lipid content. Histology slides were read and scored numerically from 1 to 5 for degree of

hepatocyte steatosis, or fatty change, by a blinded veterinary pathologist. The scores of 1 through

5 were none, mild, moderate, marked, and pathologic, respectively.

LC-MS/MS parameters

LC-MS/MS analysis was performed following the paper with minor modifications. The

LC-MS/MS system comprised a Sciex (Framingham, MA) 4500 triple quadrupole mass

spectrometer, equipped with a Turboionspray™ ionization interface in positive and negative ion

mode. The positive and negative spray voltages were 5,500 and 4,500 V, respectively, the heated

capillary temperature was 600°C, the sheath gas pressure was 50 psi, the auxiliary gas setting was

50 psi, and the heated vaporizer temperature was 350°C. The collision gas was argon at a pressure

of 1.5 mTorr. The analytical data were processed using the software program Analyst (Version

1.3). LysoGP analyses were performed in the multiple reaction monitoring (MRM) mode, in

positive ion mode for LPCs and in negative ion mode for LPAs and LPEs. The collision Energy

(CE), collision cell exit potential (CXP), entrance potential (EP), and declustering potential (DP)

were optimized for each compound to obtain optimum sensitivity.

A reverse-phase column (Thermo Scientific Hypersil Gold C4; 2.1mm ID; 10mm length;

5um particle size; 175A pore size) with a gradient elution of solvent A (5 mM ammonium formate

in water, pH 4.0) and solvent B (5 mM ammonium formate in 95% (v/v) acetonitrile, pH 4.0) was

used for chromatographic separation. Flowrate was at 0.5 ml/min. Solvent A was prepared by

mixing 95 ml of Milli-Q (Millipore) water and 5 ml of 100 mM ammonium formate, then pH was

adjusted to 4.0 with formic acid. Similarly, solvent B was prepared by mixing 95 ml acetonitrile

and 5 ml of 100 mM ammonium formate, adjusting pH 4.0 with formic acid. The gradient started

at 25% of mobile phase B, maintained 25% B for 1 min and then followed by a linear gradient to

100% B from 1 to 4 min, and held at 100% B for 6 min. Subsequently, the mobile phase was

immediately returned to the initial condition 25% B and maintained for 1 min until the end of the

run. Column effluents were introduced into the mass spectrometer between 2 and 7.5 min after

injections. The run time for a single sample was 11 min with most of phospholipids eluting from

3 mins to 5 mins.

Lipid extraction

Liver, brain, and eye tissues were weighed and placed into 1.5 ml siliconized sample tubes.

500 µl volumes of acidic methanol (pH 4.0) with 200 ng/ml 17:0-LPA and 17:0-LPC were added.

The obtained mixtures were homogenized for 1 min by Bio-Gen PRO200 Homogenizer (with a 5

mm probe, 3,000 rpm, under ice). Mixtures were continuously homogenized for 10 mins under ice

in an ultrasonic bath. After an initial centrifugation at 1,000 g for 10 min at 4°C, the supernatants

were further centrifuged at 13,500 g for 10 min at 4°C. The resulting supernatants were filtered

with a 0.2 µm pore size and 4 mm inner diameter filters. About 150 µl of filtrations were transferred

to a 96-well plate for LC-MS/MS.

Serum was incubated at 37°C for 1 h, allowed to stand at 4°C for 12 h, and centrifuged at

1,500 g for 10 min at 25°C. The serum samples (30 µl each) were placed in 1.5 ml siliconized

sample tubes, deproteinized by mixing with 150 µl acidic methanol (pH 4.0), and homogenized

for 10 min under ice in an ultrasonic bath. After an initial centrifugation at 1,000 g for 10 min, the

supernatants were further centrifuged at 13,500 g for 10 min at 4°C, as was done with the liver,

brain, and eye samples. The resulting supernatants were filtered and 150 µl of filtrations were

subjected to LC-MS/MS.

Statistics

181   Mouse weights, Rodent Lipid Panels, and results of the fatty acid analysis via LC-MS/MS

182   were analyzed via two-way ANOVA with Tukey posthoc analysis for statistical significance. All

183   data were analyzed in SPSS and considered significant if the p value was less than 0.05.

Results

Weight, water and food intake, and blood glucose

Mice gained weight appropriately according to their age throughout the study, and no mice

reached early euthanasia criteria. There were no significant differences in diabetic mouse weights

among the treatment groups at the end of treatment (Fig 1). Diabetic mice weighed significantly

more than nondiabetic control mice (p<0.05), which was to be expected due to the *db/db* obese

condition. The average daily intake of MSM in water was 4.49 ± 0.53 g/kg of body weight for the

MSM treatment group and 3.81 ± 0.33 g/kg of body weight for the MSM + SSO treatment group.

The average daily intake of the ingredient sesamin in sesame seed oil-soaked rodent feed was

23.54 ± 2.91 mg/kg of body weight for the SSO treatment group and 22.70 ± 2.57 mg/kg of body

weight for the MSM + SSO treatment group. Blood glucose readings on day 0 and day 28 of

treatment did not significantly differ within any treatment group, although the blood glucose of

db/db mice on day 28 of treatment with MSM (326 ± 21 mg/dL) was mildly decreased compared

with day 0 of treatment (375 ± 30 mg/dL).

Rodent lipid panels

Results of the Rodent Lipid Panels received from IDEXX demonstrated significant

differences between the nondiabetic control group and the diabetic groups. Serum cholesterol was

significantly increased in the type 2 diabetic control and all type 2 diabetic treatment groups

compared with the nondiabetic control group (Fig 2). Serum triglycerides were significantly

increased in the diabetic control, sesame seed oil treated, and MSM/sesame seed oil treated diabetic

groups compared with the nondiabetic control group. In the MSM treated diabetic mice, serum

triglycerides were not significantly different from the nondiabetic control group (Fig 3).

There was no statistical difference of serum HDL between the nondiabetic and the diabetic

control groups. Serum HDL was significantly increased in all diabetic treatment groups compared

with the nondiabetic control group. Additionally, the HDL levels of diabetic mice receiving

treatment with MSM/sesame seed oil combination were significantly increased compared with

treatment with either MSM alone or sesame seed oil alone (Fig 4). Serum non-HDL cholesterol

levels were significantly increased in all diabetic groups compared with the nondiabetic control.

There is a decreasing trend in all diabetic mice receiving treatment with either MSM, sesame seed

oil, or combination of the two, although the difference is not statistically significant (Fig 5).

Liver histology

In the diabetic mice, liver histology revealed variable microvesicular steatosis with

intrahepatic variability and increased red-staining lipid found in the centrilobular regions around

the central veins of the liver. There was intragroup variability and no significant differences among

treatment groups (Fig 6A-D). All liver histology slides were scored either a 1 for appearing normal

or a 2 for mild hepatic steatosis, but there was no trend among the groups, and therefore, no

statistics were performed on the liver histology scores.

Free Fatty Acids

Serum

In the serum, LA levels were significantly increased in *db/db* mice treated with sesamin

or MSM compared with nondiabetic and diabetic control mice. DGLA and AA levels were

significantly decreased in *db/db* mice compared with nondiabetic control mice, and this effect

was reversed in mice treated with either sesamin or MSM alone, where these levels were

significantly increased compared with *db/db* control mice and not significantly different from

nondiabetic control mice. In diabetic control and diabetic mice treated with combination therapy,

ALA levels were significantly increased compared with nondiabetic controls. In diabetic mice

treated with sesame seed oil alone, ALA levels were returned to levels that were not significantly

different from nondiabetic control and were significantly decreased compared with diabetic

control mice. EPA levels were significantly increased in diabetic mice treated with MSM alone

compared with both nondiabetic control and diabetic control mice. DHA levels were

significantly decreased in diabetic control mice and diabetic mice treated with combination

therapy, but this effect was reversed in diabetic mice treated with sesame seed oil alone or MSM

alone, whose DHA levels were significantly increased compared with diabetic control mice and

not significantly different from nondiabetic control mice (Fig 7A).

Liver

The livers of diabetic mice treated with sesame seed oil alone and combination therapy

had significantly increased levels of LA compared with nondiabetic controls. ALA levels were

significantly increased in diabetic control mice and diabetic mice treated with either sesame seed

oil alone or MSM alone. This effect was reversed in diabetic mice treated with MSM and sesame

seed oil, and the ALA levels were not significantly different from nondiabetic control mice.

EPA levels were significantly increased in diabetic control mice and diabetic mice treated with

MSM alone compared with nondiabetic control mice but were significantly decreased in those

treated with sesame seed oil alone compared with diabetic control. Mice receiving combination

therapy trended towards levels of EPA seen in nondiabetic control mice. There were no

significant differences regarding AA and DHA levels among groups (Fig 7B).

Brain

In the brain, LA levels were significantly decreased in diabetic mice treated with MSM

alone compared with diabetic control mice. Diabetic control mice and diabetic mice treated with

combination therapy had significantly increased DGLA levels compared with nondiabetic

controls. This effect was reversed in diabetic mice treated with MSM alone, whose DGLA levels

were significantly decreased compared with diabetic control mice and were not significantly

different from nondiabetic control mice. AA levels in diabetic mice treated with MSM alone

were significantly increased compared with both nondiabetic and diabetic control mice. There

was no significant difference in ALA levels among the groups. Diabetic control mice and

diabetic mice treated with MSM alone had significantly increased levels of EPA. This effect was

reversed in diabetic mice treated with sesame seed oil or combination therapy, with EPA levels

that were not significantly different from nondiabetic control mice and were significantly

decreased compared with diabetic control mice. DHA levels were significantly increased in

diabetic mice treated with MSM alone compared with nondiabetic control mice (Fig 7C).

Eye

In the eye, the levels of LA and ALA were significantly decreased in all diabetic mice

regardless of treatment compared with nondiabetic control mice. DGLA and AA were

significantly increased in diabetic mice receiving MSM, sesame seed oil, or combination therapy

compared with nondiabetic control mice. Diabetic mice receiving sesame seed oil or

combination treatment had significantly decreased EPA levels compared with diabetic control

mice and were not significantly different from nondiabetic control mice. DHA was significantly

increased in diabetic mice regardless of treatment compared with nondiabetic control mice (Fig

7D).

Discussion

Dyslipidemia and impaired lysophospholipid metabolism are hallmarks of type 2 diabetes

mellitus and related metabolic syndrome and obesity [1]. Diet and exercise, along with

administration of cholesterol-lowering pharmaceuticals such as statins, are recognized as standard

interventions to control hyperlipidemia. In recent years, however, there has been increasing interest

and compelling evidence to support the adjuvant use of natural products to help control

hyperlipidemia [11]. Our goal was to identify lipid alterations in *db/db* mice, a murine model of

type 2 diabetes, for the purpose of establishing a baseline biological signature for the subsequent

evaluation of natural products with purported lipid-altering activity. In this article, we utilized

mass spectrometry-based lipidomics to investigate diabetes-associated changes in fatty acid

metabolism and the extent to which the dietary ingredients sesame seed oil and/or

methylsulfonylmethane (MSM) impacted biomarkers of metabolic syndrome.

Sesamin, the primary constituent lignan found in SSO, exerts potent lipid-lowering effects

by targeting key enzymes fatty acid synthesis (e.g., delta-5 desaturase), fatty acid oxidation (e.g.,

PPARα), and cholesterol metabolism (e.g., 3-hydroxy-3-methyl-glutaryl-CoA reductase)

(reviewed in [12]). For example, low-density lipoprotein (LDL) receptor deficient mice fed a diet

of SSO were found to have reduced plasma total cholesterol, TG, very low-density lipoprotein

cholesterol, and LDL cholesterol while high-density lipoprotein (HDL) was significantly increased

compared with atherosclerotic diet-fed animals[13]. Similarly, we found that diabetic mice fed

SSO alone showed decreased non-HDL cholesterol (LDL + VLDL) accompanied by an increase

in HDL (**Figures 4 and 5**). Because the HDL-cholesterol level is increasing with treatment, the

ratio of LDL:HDL is in turn decreasing. This change in the LDL:HDL ratio due to combination

MSM and SSO treatment could lead to improved clinical outcomes in people such as reduced

incidence of atherosclerosis.

With regard to fatty acid metabolism, serum total polyunsaturated fatty acid (PUFA) levels

were significantly reduced in diabetic mice compared to control mice (**Table 1**). This finding is

consistent with literature data demonstrating that in diabetics there is a reduction in the activity of

D5D, the rate-limiting step in PUFA metabolism [14]. In contrast, feeding of only SSO reversed

this effect such that fed mice exhibited serum PUFA levels comparable to control mice. SSO

contains high amounts of omega-6 PUFAs such as linoleic acid [15], which partially explains the

effect of SSO on omega-6 PUFA levels. However, since SSO is not an appreciable source of

omega-3 PUFAs, dietary PUFAs in SSO cannot entirely account for the increase in total PUFAs,

viz., omega-6 plus omega-3 PUFAs. A clue to an alternative mechanism to explain SSO's effect

on total PUFA levels may be found in its effect on the $\Delta5$ desaturase index (AA/DGLA), which

represents a measure of PUFA metabolism where a higher index reflects greater $\Delta5$ desaturase

activity. The $\Delta5$ desaturase activity in diabetic mice (~500) was higher than either control (~100)

or SSO only fed mice (~100). Paradoxically, it appears that total serum PUFAs are reduced in

diabetic mice compared to SSO-only fed mice despite $\Delta5$ desaturase being 5X greater in diabetic

mice. Regardless of the mechanism, reduced PUFA availability can significantly impact

lysophospholipid metabolism.

Oral administration of methylsulfonylmethane to *db/db* mice results in a dose-dependent

decrease in liver cholesterol, triglycerides, lipogenesis-dependent gene expression (e.g., fatty acid

synthase), and fatty oxidation[8]. Since MSM contains two methyl groups, the authors speculated

that MSM may serve as a methyl donor to ameliorate hepatic steatosis. Building on this hypothesis,

we sought to demonstrate that MSM, acting as a methyl donor, could mitigate diabetes-related

hyperlipidemia and aberrant PUFA metabolism. Similar to SSO, administration of MSM alone

led to decreased non-HDL cholesterol (LDL + VLDL) and increased HDL (**Figures 4 and 5**).

319   The genes for delta-5 (*FADS1*) and delta-6 (*FADS2*) desaturase enzymes are situated head-

to-head (5'-5') in a cluster configuration on chromosome 11 (11q12.2). This arrangement affords

a common promoter for *FADS1* and *FADS2* that contains a potential "enhancer signature"

comprising methylation loci located within a 5' regulatory region [16]. Methylation probe studies

demonstrated an inverse relationship between methylation proportion and both $\Delta 5$ (AA/DGLA)

and $\Delta 6$ (GLA/LA) desaturase activity. These data could potentially explain the association

between methyl donor deficiency and reduced D5D activity in diabetics [14]. As mentioned above,

our data demonstrate that D5D activity is greatly enhanced in *db/db* mice compared to control or

MSM alone treated mice (Table 1). These data are consistent with the notion that reduced methyl

donor availability in diabetic mice is associated with increased D5D activity and that

administration of the putative methyl donor MSM significantly decreases D5D activity. Future

studies examining the serum and liver methyl donor status (e.g., choline, betaine, etc.) in *db/db*

mice are needed to test this hypothesis.

332   In conclusion, treatment of *db/db* mice with MSM and SSO improved commonly

measured clinical parameters in serum lipid panels. Serum triglycerides and total cholesterol

were significantly increased in the *db/db* model compared to nondiabetic control, mimicking the

diabetic condition in people. Most notably, HDL-cholesterol was significantly increased in all

db/db treatment groups, with the most significant treatment effect in the MSM + SSO

combination group. Because the HDL-cholesterol level is increasing with treatment, the ratio of

LDL:HDL is in turn decreasing. This change in the LDL:HDL ratio due to combination MSM

and SSO treatment could lead to improved clinical outcomes in people such as reduced incidence

of atherosclerosis.

343 Acknowledgements

Figure legends

Figure 1: Body weights of *db/db* mice over the 4 wk treatment.

Figure 2: Cholesterol levels of C57BL/6J mice (Control) and *db/db* mice after 4 wk treatment.

Figure 3: Triglyceride levels of C57BL/6J mice (Control) and *db/db* mice after 4 wk treatment.

Figure 4: HDL levels of C57BL/6J mice (Control) and *db/db* mice after 4 wk treatment.

Figure 5: Non-HDL cholesterol levels of C57BL/6J mice (Control) and *db/db* mice after 4 wk treatment.

Figure 6: Representative liver histology stained with Oil Red O from diabetic control group (A), diabetic mice treated with MSM (B), diabetic mice treated with sesame seed oil (C), and diabetic mice treated with both MSM and sesame seed oil (D).

Figure 7: Free fatty acids of diabetic *db/db* mouse serum (A), liver (B), brain (C), and eye (D).

359 **Figures**

361   **Figure 1. Weights**

362
363
364

Figure 2. Total Cholesterol

Figure 3. Triglycerides

Figure 4. HDL Cholesterol

375
376
377

Figure 5. Non-HDL Cholesterol

384 **Figure 6. Liver histology**

385 A

386 B

387 C

388 D

Figure 7. Serum Free Fatty Acids

394   LA   α-LA

396   DGLA   EPA

398   AA   DHA

399
400

402 **Figure 8. Liver Free Fatty Acids**

403 LA   α-LA

405 DGLA   EPA

406
407 AA   DHA

408
409

410 **Figure 9. Brain Free Fatty Acids**

411 LA   α-LA

413 DGLA   EPA

414
415 AA   DHA

416
417

418   **Figure 10. Eye Free Fatty Acids**

419   LA   α-LA

421   DGLA   EPA

422
423
424   AA   DHA

442 **Tables**

445 **Table 1. Serum Free Fatty Acids in db/db mice**

Free fatty acids	Non-DB Control (Mean ± SD)	T2D Control (Mean ± SD)	T2D Sesame oil (Mean ± SD)	T2D MSM (Mean ± SD)	T2D MSM + Sesame oil (Mean ± SD)
AA	102.35±8.37	52.14±17.07 [a]	117.02±7.46 [b]	113.87±14.62 [b]	78.53±18.57
DGLA	1.13±0.30	0.13±0.06 [a]	1.48±0.13 [b]	1.71±0.51 [b]	0.17±0.06
LA	4.15±0.55	3.80±1.09 [a]	6.33±0.98 [b]	5.93±1.15 [b]	5.39±1.09
DHA	157.67±12.73	100.48±29.68	147.51±13.16	156.42±18.36 [ab]	107.40±23.35
EPA	29.63±7.47	34.39±14.04	40.69±7.19 [ab]	97.29±12.78 [ab]	16.97±7.80
α-LA	0.59±0.16	1.69±0.47 [a]	0.89±0.21 [b]	0.96±0.32	1.77±0.37 [a]
NPD1	0.03±0.01	0.04±0.01	0.05±0.02	0.03±0.02	0.04±0.01
PGE2	0.10±0.01	0.16±0.07 [a]	0.13±0.01 [a]	0.12±0.03	0.12±0.03
total n-3	187.89±19.59	136.56±42.69	189.09±18.11	254.67±30.50 [b]	126.14±30.67 [a]
total n-6	107.64±8.56	56.07±17.89 [a]	124.83±7.94 [b]	121.52±15.85 [b]	84.09±19.36
Total	295.52±25.26	192.63±60.04	313.92±25.17 [a]	376.18±45.13 [a]	210.23±48.08
n-6/n-3	0.58±0.06	0.41±0.05 [a]	0.66±0.04 [b]	0.48±0.03	0.68±0.10 [b]
AA/DHA	0.65±0.05	0.52±0.06	0.80±0.05 [ab]	0.73±0.04 [b]	0.74±0.11 [b]
AA/EPA	3.68±1.07	1.58±0.30 [a]	2.94±0.42 [a]	1.17±0.09 [a]	5.18±1.50 [b]

Table 2. Liver Free Fatty Acids in db/db mice

Free fatty acids	Non-DB Control (Mean ± SD)	T2D Control (Mean ± SD)	T2D Sesame oil (Mean ± SD)	T2D MSM (Mean ± SD)	T2D MSM + Sesame oil (Mean ± SD)
AA	37.99±10.64	36.31±6.22	45.63±15.35	40.63±15.54	44.78±10.31
DGLA	0.07±0.03	0.07±0.02	0.08±0.03	0.09±0.04	0.09±0.02
LA	0.92±0.29	1.23±0.25	1.51±0.49 [a]	1.21±0.32	1.47±0.30 [a]
A-LA	0.09±0.03	0.18±0.06 [a]	0.17±0.05 [a]	0.21±0.09 [a]	0.11±0.03
DHA	50.17±13.54	40.33±5.12	45.09±13.85	48.31±14.24	45.67±7.37
EPA	6.78±2.28	31.98±4.97 [a]	13.20±1.56 [b]	42.04±11.71 [a]	15.78±4.34
NPD1	0.01±0.00	0.00±0.00	0.01±0.00	0.00±0.00	0.01±0.00
PGE2	0.05±0.01	0.04±0.01	0.05±0.02	0.06±0.02	0.05±0.02
total n-3	57.04±15.47	72.49±8.29	58.46±15.01	90.56±25.42 [a]	61.56±11.28
total n-6	38.99±10.91	37.62±6.39	47.22±15.83	41.94±15.87	46.33±10.49
Total PUFAs	96.03±26.06	110.11±14.15	105.68±30.01	132.50±39.38	107.90±21.60
n-6/n-3	0.69±0.06	0.52±0.05	0.80±0.12 [b]	0.46±0.10 [a]	0.75±0.05 [b]
AA/DHA	0.76±0.08	0.90±0.06	1.02±0.16 [a]	0.84±0.18	0.97±0.09 [a]
AA/EPA	5.85±1.43	1.15±0.18 [a]	3.40±0.85 [b]	0.96±0.25 [a]	2.89±0.39

Table 3. Brain Free Fatty Acids in db/db mice

Free fatty acids	Non-DB Control (Mean ± SD)	T2D Control (Mean ± SD)	T2D Sesame oil (Mean ± SD)	T2D MSM (Mean ± SD)	T2D MSM + Sesame oil (Mean ± SD)
AA	1,025.36±195.89	1,162.52±314.12	1,253.63±451.91	1,503.06±205.06 [ab]	1,358.28±323.89
LA	9.73±2.20	11.52±1.91	13.61±3.76	8.73±1.77	12.62±2.48
DGLA	7.19±2.89	14.85±3.97 [a]	10.67±3.06	10.44±1.40 [b]	11.52±2.41 [a]
DHA	1,038.56±223.54	1,180.76±354.49	1,149.79±340.71	1,489.79±226.12 [a]	1,364.83±312.89
EPA	98.07±20.99	150.31±29.30 [a]	81.67±18.03 [b]	198.73±33.64 [a]	104.68±28.58 [b]
α-LA	1.04±0.77	1.52±0.48	1.21±0.40	1.08±0.36 [b]	1.28±0.50
NPD1	0.51±0.77	0.38±0.28	0.23±0.15	0.16±0.08	0.28±0.15
PGE2	1.70±0.73	1.88±0.28	1.66±0.31	1.96±0.26	2.07±0.36
total n-3	1,137.67±222.40	1,332.59±369.65	1,232.67±355.44	1,689.60±243.27 [a]	1,470.79±330.90
total n-6	1,042.28±196.96	1,188.90±312.11	1,277.91±452.79	1,522.24±204.48 [a]	1,382.42±324.02
total PUFAs	2,179.95±402.22	2,521.48±675.66	2,510.58±775.13	3,211.84±446.24 [a]	2,853.21±645.70
n-6/n-3	0.92±0.11	0.90±0.07	1.03±0.16 [b]	0.90±0.02	0.94±0.08
AA/DHA	1.00±0.12	1.00±0.09	1.09±0.17	1.01±0.03	1.00±0.07
AA/EPA	11.01±4.06	7.84±1.99	15.27±4.00 [b]	7.67±1.04	13.59±3.82 [b]

Table 4. Eye Free Fatty Acids in db/db mice

Free fatty acids	Non-DB Control (Mean ± SD)	T2D Control (Mean ± SD)	T2D Sesame oil (Mean ± SD)	T2D MSM (Mean ± SD)	T2D MSM + Sesame oil (Mean ± SD)
AA	71.03±4.46	75.73±7.93	84.13±5.39 [a]	84.74±6.39 [a]	86.57±10.45 [ab]
LA	6.08±1.15	1.55±0.66 [a]	1.91±0.39 [a]	1.51±0.27 [a]	1.93±0.59 [a]
DGLA	0.64±0.06	0.76±0.11	0.80±0.10 [a]	0.84±0.11 [a]	0.87±0.14 [a]
DHA	149.18±14.20	174.65±12.86 [a]	167.85±8.54	177.71±9.97 [a]	184.75±12.78 [a]
EPA	9.99±1.45	12.16±2.02	8.38±0.99 [b]	14.58±1.66	9.87±1.61 [a]
α-LA	0.48±0.12	0.20±0.09 [a]	0.17±0.04 [a]	0.18±0.04 [a]	0.21±0.07 [a]
NPD1	0.05±0.01	0.01±0.00 [a]	0.01±0.00 [a]	0.01±0.00 [a]	0.01±0.00 [a]
PGE2	0.04±0.01	0.04±0.01	0.04±0.00	0.04±0.00	0.04±0.01
total n-3	159.65±14.97	187.01±13.54 [a]	176.40±8.56	192.48±10.12 [a]	194.82±13.63 [a]
total n-6	77.75±4.29	78.03±8.38	86.84±5.43	87.09±6.61	89.37±10.67 [ab]
total PUFAs	237.41±17.52	265.04±20.86 [a]	263.24±11.59	279.57±14.58 [a]	284.20±22.13 [a]
n-6/n-3	0.49±0.04	0.42±0.03 [a]	0.49±0.03 [b]	0.45±0.03	0.46±0.04 [b]
AA/DHA	0.48±0.04	0.43±0.03	0.50±0.03 [b]	0.48±0.04	0.47±0.05
AA/EPA	7.21±0.84	6.31±0.76	10.13±0.97 [ab]	5.85±0.54	8.85±0.75 [b]

491
492
493
494

References

1.	Saeedi P, Petersohn I, Salpea P, Malanda B, Karuranga S, Unwin N, et al. Global and regional diabetes prevalence estimates for 2019 and projections for 2030 and 2045: Results from the International Diabetes Federation Diabetes Atlas, 9(th) edition. Diabetes Res Clin Pract. 2019;157:107843. Epub 2019/09/14. doi: 10.1016/j.diabres.2019.107843. PubMed PMID: 31518657.

2.	Barber MN, Risis S, Yang C, Meikle PJ, Staples M, Febbraio MA, et al. Plasma lysophosphatidylcholine levels are reduced in obesity and type 2 diabetes. PLoS One. 2012;7(7):e41456. Epub 2012/08/01. doi: 10.1371/journal.pone.0041456. PubMed PMID: 22848500; PubMed Central PMCID: PMCPMC3405068.

3.	Del Bas JM, Caimari A, Rodriguez-Naranjo MI, Childs CE, Paras Chavez C, West AL, et al. Impairment of lysophospholipid metabolism in obesity: altered plasma profile and desensitization to the modulatory properties of n-3 polyunsaturated fatty acids in a randomized controlled trial. Am J Clin Nutr. 2016;104(2):266-79. Epub 2016/06/17. doi: 10.3945/ajcn.116.130872. PubMed PMID: 27305954.

4.	Simopoulos AP. An Increase in the Omega-6/Omega-3 Fatty Acid Ratio Increases the Risk for Obesity. Nutrients. 2016;8(3):128. Epub 2016/03/08. doi: 10.3390/nu8030128. PubMed PMID: 26950145; PubMed Central PMCID: PMCPMC4808858.

5.	Lazic M, Inzaugarat ME, Povero D, Zhao IC, Chen M, Nalbandian M, et al. Reduced dietary omega-6 to omega-3 fatty acid ratio and 12/15-lipoxygenase deficiency are protective against chronic high fat diet-induced steatohepatitis. PLoS One. 2014;9(9):e107658. Epub 2014/09/25. doi: 10.1371/journal.pone.0107658. PubMed PMID: 25251155; PubMed Central PMCID: PMCPMC4175074.

6.	Cobb J, Eckhart A, Perichon R, Wulff J, Mitchell M, Adam KP, et al. A novel test for IGT utilizing metabolite markers of glucose tolerance. J Diabetes Sci Technol. 2015;9(1):69-76. Epub 2014/09/28. doi: 10.1177/1932296814553622. PubMed PMID: 25261439; PubMed Central PMCID: PMCPMC4495543.

7.	Kim LS, Axelrod LJ, Howard P, Buratovich N, Waters RF. Efficacy of methylsulfonylmethane (MSM) in osteoarthritis pain of the knee: a pilot clinical trial. Osteoarthritis Cartilage. 2006;14(3):286-94. Epub 2005/11/29. doi: 10.1016/j.joca.2005.10.003. PubMed PMID: 16309928.

8.	Sousa-Lima I, Park SY, Chung M, Jung HJ, Kang MC, Gaspar JM, et al. Methylsulfonylmethane (MSM), an organosulfur compound, is effective against obesity-induced metabolic disorders in mice. Metabolism. 2016;65(10):1508-21. doi: 10.1016/j.metabol.2016.07.007. PubMed PMID: 27621186.

9.	Bournival J, Plouffe M, Renaud J, Provencher C, Martinoli MG. Quercetin and sesamin protect dopaminergic cells from MPP+-induced neuroinflammation in a microglial (N9)-neuronal (PC12) coculture system. Oxid Med Cell Longev. 2012;2012:921941. Epub 2012/08/25. doi: 10.1155/2012/921941. PubMed PMID: 22919443; PubMed Central PMCID: PMCPMC3418684.

10.	Amirshahrokhi K, Bohlooli S. Effect of methylsulfonylmethane on paraquat-induced acute lung and liver injury in mice. Inflammation. 2013;36(5):1111-21. Epub 2013/04/19. doi: 10.1007/s10753-013-9645-8. PubMed PMID: 23595869.

11.   El-Tantawy WH, Temraz A. Natural products for controlling hyperlipidemia: review.
Archives   of   physiology   and   biochemistry.   2019;125(2):128-35.   doi:
10.1080/13813455.2018.1441315.
12.   Majdalawieh AF, Dalibalta S, Yousef SM. Effects of sesamin on fatty acid and cholesterol
metabolism, macrophage cholesterol homeostasis and serum lipid profile: A comprehensive
review. European Journal of Pharmacology. 2020:173417.
13.   Bhaskaran S, Santanam N, Penumetcha M, Parthasarathy S. Inhibition of atherosclerosis
in low-density lipoprotein receptor-negative mice by sesame oil. J Med Food. 2006;9(4):487-90.
Epub 2007/01/05. doi: 10.1089/jmf.2006.9.487. PubMed PMID: 17201634.
14.   Kroger J, Schulze MB. Recent insights into the relation of Delta5 desaturase and Delta6
desaturase activity to the development of type 2 diabetes. Curr Opin Lipidol. 2012;23(1):4-10. doi:
10.1097/MOL.0b013e32834d2dc5. PubMed PMID: 22123669.
15.   Carvalho R, Galvão E, Barros J, Conceição M, Sousa E. Extraction, fatty acid profile and
antioxidant activity of sesame extract (Sesamum Indicum L.). Brazilian Journal of Chemical
Engineering. 2012;29(2):409-20.
16.   Howard TD, Mathias RA, Seeds MC, Herrington DM, Hixson JE, Shimmin LC, et al. DNA
Methylation in an Enhancer Region of the FADS Cluster Is Associated with FADS Activity in
Human Liver. PLOS ONE. 2014;9(5):e97510. doi: 10.1371/journal.pone.0097510.
560