1

A novel and effective Cre/*lox*-based genetic tool for repeated, targeted and markerless integration in *Yarrowia lipolytica*

Qinghua Zhou [†], Liangcheng Jiao[†], Wenjuan Li, Zhiming Hu, Yunchong Li, Houjin Zhang, Li Xu [*], Yunjun Yan[*]

Key Laboratory of Molecular Biophysics of the Ministry of Education, College of Life Science and Technology, Huazhong University of Science and Technology, Wuhan 430074, P. R. China

[†]These authors contributed equally to this work

*Corresponding author

E-mail: xuli@mail.hust.edu.cn (Li Xu), yanyunjun@hust.edu.cn (Yunjun Yan)

Running title: A novel and effective genetic tool for gene integration

Keywords: Genetic tool, Cre/*lox*, repeatedly targeted and markerless integration, *Yarrowia lipolytica*

Abstract

The unconventional yeast *Yarrowia lipolytica* is extensively applied in bioproduction fields owing to its excellent metabolite and protein production ability. Nonetheless, utilization of this promising host is still restricted by limited availability of precise and effective gene integration tools. In this study, a novel and efficient genetic tool was developed for targeted, repeated, and markerless gene integration based on Cre/*lox* site-specific recombination system. The developed tool required only a single selection marker and could completely excise all of the unnecessary sequences. A total of three plasmids were created and seven rounds of marker-free gene integration were examined in *Y. lipolytica*. All the integration efficiencies remained above 90%, and analysis of protein production and growth characteristics of the engineered strains confirmed that genome modification via the novel genetic tool was feasible. Further work also confirmed the genetic tool was effective for integration of other genes, loci, and strains. Thus, this study significantly promotes the application of Cre/*lox* system and presents a powerful tool for genome engineering in *Y. lipolytica*.

Introduction

With the development of synthetic biology, various molecular genetic tools are being increasingly designed for precise genome modification in yeasts. As a promising host, the generally regarded as safe (GRAS) yeast, *Yarrowia lipolytica,* has been employed in a wide range of fields such as protein production, metabolic engineering, degradation of *n*-alkanes, and dimorphism (Blazeck et al. 2014; Madzak 2015; Carly et al. 2017; Abdel-Mawgoud et al. 2018; Czajka et al. 2018; Madzak 2018). Nevertheless, only few effective genetic tools have been developed for repeatedly targeted gene integration in this host, which still restricts further exploitation and application of *Y. lipolytica*.

In traditional methods, heterologous and endogenous genes are transformed into *Y. lipolytica* genome via insertion of the whole plasmid at target loci or integration of tandem fragment of target gene and selection marker at zeta loci (Barth and Gaillardin 1996; Pignede et al. 2000; Larroude et al. 2018). However, introduction of non-target DNA fragments, especially antibiotics resistance genes, may harm the GRAS status of *Y. lipolytica* and hinder its application in food and medicine industries. Besides, the bioproduction of *Y. lipolytica* usually requires multi-round integrations of target gene or co-expression of various genes (Ledesma-Amaro and Nicaud 2016; Carly et al. 2017; Larroude et al. 2018; Markham et al. 2018; Yang et al. 2019), which can be hardly realized via these conventional methods owing to the limited availability of selection markers. Consequently, there is an urgent need to develop efficient genetic tool(s) for targeted, repeatable, and markerless gene integration in *Y. lipolytica*.

To date, several site-specific nuclease technologies have been established for genome modification in various species (Cong et al. 2013; Gaj et al. 2013; Hsu et al.

2014; Wu et al. 2014). Among them, CRISPR-Cas9 system can accomplish relatively

high-efficiency gene disruption and marker-free gene integration in *Y. lipolytica*

(Schwartz et al. 2017a; Holkenbrink et al. 2018; Bae et al. 2020). A previous study

examined 17 loci for targeted and markerless gene integration in *Y. lipolytica,* among

which only five loci exhibited gene integration, with highest integration efficiency of

68.9%±25.5% (Schwartz et al. 2017b). Another study accomplished gene integration into

*PEX*10 locus via homologous recombination (HR) and homology-mediated end-joining,

with integration efficiencies of 6.7%±3.6% and 37.5%±8.8%, respectively (Gao et al.

2018). These studies indicate the need for further improvement in achieving more

desirable gene integration efficiency (similar to gene disruption efficiency of about 90%)

via CRISPR-Cas9-based tools. Besides, gene insertion via HR-mediated double-strand

break (DSB) repair usually demands synchronous introduction of two plasmids harboring

different selection markers (Schwartz et al. 2017b; Gao et al. 2018), resulting in reduction

of transformation efficiency and additional procedure for plasmids recovery.

The Cre/*lox*-site-specific recombination system derived from bacteriophage P1 is

known to be appropriate for genome editing (Kuhn and Torres 2002; Kawano et al. 2016;

Richardson et al. 2017; Wu et al. 2018), and its action mechanism has been clearly

described (Albert et al. 1995; Leibig et al. 2008; Yan et al. 2008). Cre recombinase can

recognize and bind to *lox*P site, and can catalyze molecular recombination reactions

including insertion, translocation, inversion and deletion (Albert et al. 1995). However,

these reactions can retain redundant *lox*P sites in the genome, which may disturb further

recombination (Albert et al. 1995). To address this issue, some mutational *lox* sites have

been successfully designed (Leibig et al. 2008; Yan et al. 2008). For example, the

recombination between a left element-mutant *lox* site (LE) *lox*71 and a right element-

mutant *lox* site (RE) *lox*66 can generate a double element-mutant *lox* site (LE+RE) *lox*72

and a *lox*P site (Leibig et al. 2008; Yan et al. 2008). Meanwhile, *lox*P site can be deleted

and *lox*72 site can be retained in the genome through rational design. In particular, the

*lox*72 site, which can weakly bind to Cre recombinase, is less likely to participate in

subsequent recombination (Leibig et al. 2008; Yan et al. 2008). As a result, *lox* mutants

can exhibit enhanced controllability of recombination and accelerate the utilization of

Cre/*lox* system in yeasts (Gueldener et al. 2002; Fickers et al. 2003; Hentges et al. 2005;

Hochrein et al. 2018). Although this method can employ marker-free genome

modification, it often presents low efficiency or is time-consuming. In addition, a few

studies have attempted markerless gene integration via Cre/*lox* system in *Y. lipolytica*.

Among them, an inspired method combining Cre/*lox* system and 26S rDNA achieved

79.2% integration efficiency in *Y. lipolytica* (Lv et al. 2019); however, the integration

loci were random and the copy numbers of target genes were uncontrollable, requiring lot

of time for selection and evaluation of transformants. Moreover, to rescue selection

marker in the genome, an additional plasmid harboring *cre* gene must be introduced into

the recombinants by another transformation, and this plasmid should subsequently be

recovered. These tedious procedures require two different selection markers and further

extend the experimental period.

Therefore, in the present study, a novel and effective genetic tool was designed

based on Cre/*lox* system. Only a single selection marker was adequate to enable repeated,

targeted, and markerless gene integration, and all unnecessary fragments could be totally

excised. To demonstrate the efficacy of the developed tool, an important lipase gene from

Rhizomucor miehei (*rml*) was chosen as a qualitative reporter, and three new plasmids were constructed. A total of seven rounds of marker-recyclable integration were implemented, and the positive colony proportions reached 90.63%–100%. The experiment conditions, feasibility, and controllability of the genetic tool were evaluated and optimized. Moreover, to confirm the universality of the developed tool, other six different genes were further integrated into *leu2* locus, and a derived tool was successfully employed to integrate another target gene into the *axp1-2* locus. The results obtained revealed that Cre/*lox*-based genetic tool could resolve the limitations of the existing methods and bring significant advancement in gene editing in yeast species.

Results

Genetic tool design and integration process analysis

The novel Cre/*lox*-based genetic tool comprised three plasmids, namely, Cre-Y1, Cre-Y2, and Cre-Y3 (Figure 1). Plasmid Cre-Y1 was applied for generating the initial strain, and plasmids Cre-Y2 and Cre-Y3 were employed to iteratively introduce target genes into the host strain. Cre-Y1 consisted of a yeast selection marker, *cre* gene expression cassette, bacterial element (antibiotic resistance gene and origin of replication), homologous fragment of target locus in the genome (HF-A), and two convergently oriented *lox* mutants. The *cre* expression cassette should adopt an inducible promoter to control the production of Cre recombinase. Cre-Y2 was derived from Cre-Y1 by replacing HF-A and two convergently oriented *lox* mutants with another homologous fragment of target locus (HF-B), a *lox* mutant and its reverse complement (rc*lox*), and a target gene expression cassette. To avoid excising original segments from the genome, HF-B should be close to

HF-A. Cre-Y3 was similar to Cre-Y2, but with the corresponding *lox* mutants of Cre-Y2

(e.g., LE for Cre-Y2 and RE for Cre-Y3). All of the tool elements were flexible and

nonspecific; thus, the Cre/*lox*-based genetic tool was appropriate for different species.

Plasmid design   **Repeated, targeted and markerless integration**

Figure 1. Plasmid design and integration process analysis.

In the present study, the plasmid elements used for *Y. lipolytica* were as follows

(Figure 1): the auxotroph Ura3 (UniProtKB, Q12724) was adopted as selection marker,

cre expression cassette was assembled as pPOX2-*cre*-lip2t (pPOX2 promoter, *cre* gene,

and *lip2* terminator, GenBank Accession Nos. AJ001300.1, AB449974.1, and

AJ012632.1), bacterial element (Amp-ori) and target gene expression cassette (*rml*) were

subcloned from plasmid hp12d-*rml*, and HF-B and HF-A were respectively designed as

leu2 gene (GenBank Accession No. AF260230.1) and its upstream segment (Upleu) for

the convenience of plasmid construction and linearization. In addition, two *lox*71 sites, a

*lox*66 site and a reverse complement of *lox*66 site (rc*lox*66) and a reverse complement of

*lox*71 site (rc*lox*71) and a *lox*71 site were used in plasmids Cre-Y1, Cre-Y2, and Cre-Y3,

respectively.

To demonstrate repeated and markerless gene integration in *Y. lipolytica* via Cre/*lox*-

based genetic tool, the experimental process was as follows (Figure 1). First, Cre-Y1

containing two *lox*71 sites was introduced into the Upleu locus of Po1f to construct an

initial strain harboring a single *lox*71 site. Under the condition of suitable screening and

induction, recombination between two *lox*71 sites could remove unnecessary plasmid

elements and produce the initial strain. Then, Cre-Y2 was integrated into *leu2* locus of

the initial strain, forming a transitional strain harboring three *lox* sites (*lox*71, *lox*66, and

rc*lox*66) in the genome. Various recombination reactions among these three *lox* sites

could excise unnecessary DNA fragments and retain a *lox*72 site, an rc*lox*66 site, and a

target gene expression cassette in the genome. Meanwhile, two types of stable strains

were eventually generated. Similarly, integration of Cre-Y3 produced strains harboring

two target gene expression cassettes. Consequently, repeated, targeted, and markerless

gene integration could be efficiently realized following iterative integration of Cre-Y2

and Cre-Y3.

First-round integration to construct initial strain FY1

To construct strain harboring a single *lox*71 fragment, the plasmid Cre-Y1 was inserted

into auxotrophic host Po1f (Figure 2). The transformed cell pellet was screened twice in 5

mL of MD-Leu medium to enrich the Ura$^+$ recombinant strain Po1f/Cre-Y1 (Ura$^+$ and

Leu$^-$) and avoid interference from untransformed Po1f cells (Supplemental Fig. S10).

Subsequently, the culture solution containing Po1f/Cre-Y1 was transferred into MO-Leu-

Ura medium for 12 h with oleic acid as the sole carbon source, which could activate

pPOX2 to induce the expression of *cre* gene. In this case, the produced Cre recombinase

could mediate deletion reaction between two repeat *lox*71 sites in the Po1f/Cre-Y1

genome, causing excision of the bacterial element (Amp-ori), selection marker (Ura3),

and *cre* expression cassette, further generating Ura$^-$ recombinants. Meanwhile, the extra

added uracil allowed Ura$^-$ recombinants to grow normally. The induced cell suspension

was streaked on YPD plate for isolating single colony with Ura$^-$ genotype, and the

phenotype of the isolated colonies were determined on YPD and MD-Leu media. As

shown in Supplemental Fig. S1A, a total of 32 recombinants were randomly chosen and

all of them could not grow on MD-Leu plate, proving that the selection marker Ura3 was

recycled successfully. Moreover, 10 positive colonies were picked up randomly and

identified by genome PCR (Supplemental Fig. S2A), and the PCR product obtained via

primers Upleu-F3/Upleu-R3 was further sequenced. The results shown in Supplemental

Fig. S3A indicated that only a single *lox*71 site was retained in Upleu locus of the

eventually obtained strain (named FY1). This finding also revealed that all recombination

and integration events occurred as expected (Figure 2), and the adopted plasmid elements

of the novel genetic tool were well suitable for *Y. lipolytica*.

Figure 2. Schematic diagram of first-round integration to obtain initial strain FY1. The Cre-Y1 was linearized at homologous fragment Upleu and introduced into Po1f. The Po1f/Cre-Y1 was screened twice in MD-Leu medium and then inoculated into MO-Leu-Ura medium for inducible expression of *cre* gene. The bacterial element (Amp-ori), selection marker (Ura3) and *cre* expression cassette (pPOX2-*cre*-lip2t) were removed by deletion reaction of two *lox*71 sites, retaining a single *lox*71 site

in the FY1 genome. Primers used for PCR amplification are labeled in the figure and described in

Supplemental Table S3.

Second-round integration and analysis

To evaluate whether the designed plasmids could realize markerless gene integration,

Cre-Y2 was linearized within *leu2* and introduced into FY1, to obtain the strain FY1/Cre-

Y2. As shown in Figure 3, three *lox* sites (*lox*71, *lox*66, and rc*lox*66) were noted in the

FY1/Cre-Y2 genome, and three kinds of recombination reactions could occur when Cre

recombinase was expressed. In the first-type recombination (a), deletion reaction

occurred between the *lox*71 and *lox*66 sites in the FY1/Cre-Y2 genome, and the strain

FY2-*rml* carrying *rml* expression cassette was obtained. In the second-type

recombination (b), the *lox*71 and rc*lox*66 sites were combined with Cre proteins,

producing a *lox*72 site and reverse complement of *lox*P site (rc*lox*P). Here, the *lox*72 site

was hardly bound to Cre recombinase, and the rc*lox*P site could not stably coexist with

the unreacted rc*lox*66 site in the genome, which could recombine with Cre recombinase

again, generating FY2-rc*rml*, a derived strain carrying the reverse complement of *rml*

gene expression cassette (rc*rml*). In the third-type recombination (c), the *lox*66 and

rc*lox*66 sites were recombined for inversion reaction, forming a transitional strain that

also contained three *lox* sites similar to those of FY1/Cre-Y2. The recombination reaction

among the three *lox* sites continued until the first or second type of recombination

occurred, and the strain FY2-*rml* or FY2-rc*rml* was finally generated. Consequently, two

kinds of stable strains were obtained from three types of recombination reactions in

second-round integration. Here, all of the unnecessary and redundant fragments,

including Upleu, *leu2-270*, bacterial element, selection marker, and *cre* expression

cassette, were completely deleted, and only gene expression cassette and rc*lox*66 site

remained in the genome.

Y2. The FY1/Cre-Y2 was screened twice in MD medium and then induced in MO-Ura medium for

the production of Cre protein. Therein, three kinds of different recombination reactions for *lox* sites

were noted in the genome: deletion reaction between *lox*71 and *lox*66 sites (a), inversion reaction

between *lox*71 and rc*lox*66 sites (b), and inversion reaction between *lox*66 and rc*lox*66 sites (c).

Eventually, two forms of stable strains FY2-*rml* and FY2-rc*rml* were produced. Meanwhile, all of the

unnecessary and redundant sequences were excised, the *rml* expression cassette was integrated into

the genome, and the remaining rc*lox*66 site locus was employed in third-round integration. Primers

used for PCR amplification are labeled in the figure and described in Supplemental Table S3.

Induction time optimization in second-round integration

In the present study, oleic acid induced pPOX2 promoter controlled the production of Cre

recombinase in MO liquid medium. Therein, induction time affected Cre protein

concentration in the cell, and further influenced the recombination efficiency in the

engineered strains. In the first-round gene integration, there were two *lox* sites in the

yeast genome. The recombinants were induced for sufficiently long time to obtain high

proportion of positive colonies. However, in subsequent gene integrations, there were

three or more *lox* sites in the genome and the recombination reaction was complex and

diverse. As reported in previous studies (Albert et al. 1995; Suzuki et al. 2005), high

concentration Cre protein is more likely to mediate recombination in *lox*72 site, resulting

in the removal of the target gene expression cassette from the engineering strain genome.

Therefore, induction of *cre* gene expression should be controlled in an appropriate time

range. In the present study, to determine the optimal induction time, the recombinant

strain FY1/Cre-Y2 was incubated in MO-Ura liquid medium for 4–16 h and streaked

onto YPD medium every 2 h (approximately one cell generation cycle). Furthermore, 32

recombinants were randomly selected from each YPD medium and then transferred onto
MD and BMSY media. The proportions of positive colonies under different induction
times were examined, and the results are shown in Table 1 and Supplemental Fig. S1B.
All the selected recombinants could complete recombination reaction after 12 h of
induction. For more accurate confirmation, 64 colonies induced for 12 h were chosen and
none of them could normally grow on MD medium. Consequently, the induction time
was set as 12 h, which could be further increased if the proportion of positive colonies
decreased in multi-round integrations.

Table 1. Positive colony proportion, genotype rate, and *rml* copy number determined in this study.

Integration round	Induction time (h)	Positive colonies/ all colonies	Proportion (%)	Selected colonies		
				Genotype	Rate	*rml* copy number
1	12	32/32	100	FY1	-	0
2	4	2/32	6.25	FY2-*rml*, FY2-rc*rml*	5/5	0.93
	6	2/32	6.25			
	8	14/32	43.75			
	10	24/32	75			
	12	32/32	100			
	14	32/32	100			
3	12	32/32	100	FY3-2*rml*, FY3-*rml*-rc*rml*	17/13	1.99
4	12	31/32	96.88	FY4-3*rml*, FY4-2*rml*-rc*rml*	4/2	3.10
5	12	29/32	90.63	FY5-4*rml*, FY5-3*rml*-rc*rml*	3/3	4.08
6	12	17/32	53.12	FY6-5*rml*, FY6-4*rml*-rc*rml*	4/2	5.12
	14	28/32	87.5			
	16	32/32	100			
7	16	32/32	100	FY7-6*rml*, FY7-5*rml*-rc*rml*	3/3	5.94

A total of 10 positive colonies were randomly chosen for identification via PCR
sequencing. The *rml*/rc*rml* expression cassette and rc*lox*66 site in all the selected colonies

matched with the genome of FY2-*rml*/FY2-rc*rml* (Supplemental Figs. S3B, C),

demonstrating that the second-round gene integration was performed according to the

predictive analysis (Figure 3). In addition, for the convenience of subsequent protein

production evaluation, strains that do not carry rc*rml* expression cassette were employed

for further gene integration, because the existing plasmids and method (Zhou et al. 2019)

are more applicable to construct strains only harboring *rml* expression cassettes.

Iterative insertion of Cre-Y3 and Cre-Y2

To achieve third-round integration, Cre-Y3 was utilized to transform FY2, and the

schematic diagram is presented in Figure 4. Similar to the second-round integration, there

were three *lox* mutants (rc*lox*66, rc*lox*71, and *lox*71) and a *lox*72 site in FY2-*rml*/Cre-Y3

genome, resulting in three different recombination reactions and eventually generating

two genotypes of the recombinant strains (FY3-2*rml* and FY3-*rml*-rc*rml*). Subsequently,

FY3-2*rml* and FY3-*rml*-rc*rml* were distinguished by PCR amplification with primer pairs

KNC-F2/Apa-R6 and SXB-R2/Apa-R3 (Supplemental Fig. S2C). The obtained products

were further sequenced and the results (Supplemental Fig. S3D, E) were consistent with

the prediction presented in Fig. 4. Thus, another copy of *rml* gene was targeted on *leu2*

locus and all unnecessary DNA fragments were excised in the third-round integration.

Figure 4. Schematic diagram of third-round integration to construct FY3-2*rml*/FY3-*rml*-rc*rml*. The *Apa*I-linearized Cre-Y3 was used to transform FY2-*rml* 3#, generating the strain FY2-*rml*/Cre-Y3.

There were three kinds of different recombination reactions for *lox* sites in FY2-*rml*/Cre-Y3 genome.

Herein, the recombination reactions excised unnecessary DNA fragments (a *leu2*, Amp-ori, Ura3, and

pPOX2-*cre*-lip2t) and eventually generated two kinds of stable strains FY3-2*rml* and FY3-*rml*-rc*rml*.

The remaining *lox*71 site could be used for subsequent integration process. Primers used for PCR

amplification are labeled in the figure and described in Supplemental Table S3.

Similarly, Cre-Y2 was introduced into FY3-2*rml* to generate FY4-3*rml*/FY4-2*rml*-

rc*rml*, and Cre-Y3 was used to transform FY4-2*rml* for constructing FY5-4*rml*/FY5-

3*rml*-rc*rml* (Supplemental Fig. S4A). Subsequently, Cre-Y2 and Cre-Y3 were iteratively

integrated into the corresponding recombinant strains to obtain FY6-5*rml*/FY6-4*rml*-

rc*rml* and FY7-6*rml*/FY7-5*rml*-rc*rml*. The genome PCR identification results

(Supplemental Fig. S4B) proved that repeated, targeted, and markerless gene integration

was successfully implemented by the novel genetic tool.

Increasing induction time in sixth and seventh rounds of integration

As shown in Table 1, the positive colony proportions slightly decreased in multi-round

integration processes, which might be owing to the following possibility: an additional

*lox*72 site could have been inserted into the genome in each round of gene integration,

except in the first-round integration, and all of the *lox* sites can bind to Cre recombinase

(Albert et al. 1995; Leibig et al. 2008; Yan et al. 2008). Thus, *lox*72 site might disturb the

combination between LE/RE *lox* sites and Cre recombinase, resulting in increasingly less

recombination reactions during the induction process and a decrease in efficiency in

multi-round integrations. Nevertheless, an increase in induction time caused higher

accumulation of Cre protein in the cell, and more Cre recombinase could recognize

LE/RE mutants and mediate effective recombination.

Accordingly, the transformants were induced for longer time to obtain higher

recombination efficiency in the sixth-round integration. As a result, the proportion of

positive colonies was enhanced, and 16 h was adequate to complete recombination in the

engineered strains (Table 1 and Supplemental Fig. S1C). In the seventh-round integration,

16 h was employed as the induction time and all the chosen colonies could not grow on

MD medium (Supplemental Fig. S1C). Therefore, recombination efficiency might be

roughly inversely proportional to the integration round; however, this problem could be

alleviated by increasing the induction time in MO-Ura liquid medium. Finally, seven

rounds of gene integration were implemented in this study, and the positive colony

proportions reached 90.63%–100% with an induction time of 12–16 h (Table 1). In fact,

the realized integration rounds were adequate to meet the demands of engineering strains

construction, and the potential of the developed genetic tool could also be explored in

future research.

Protein expression level and growth characteristics evaluation

The recombinants harboring different *rml* gene copies were cultured in BMSY liquid

medium for protein production, and the results are presented in Figure 5A. The

recombinant strains FY5-4*rml* 31# showed the highest RML expression level, while the

lipase activity of FY6-5*rml*/FY6-4*rml*-rc*rml* and FY7-6*rml*/FY7-5*rml*-rc*rml* was slightly

lower. Thus, gene integration was terminated in the seventh round. Moreover, several

colonies were chosen as representatives for each type of the recombinant strains, and

their *rml* gene copies were determined via qPCR. As shown in Table 1, all the

transformants carried the same gene copy number as previously predicted, indicating that

no *rml* expression cassette was excised from the genome and also confirming the

excellent controllability and stability of the genetic tool.

To evaluate the protein expression level of the recombinant strains generated via the

developed genetic tool, the plasmids hp12d-5*rml* and hp12d-6*rml* were constructed

according to the previous method (Zhou et al. 2019). Subsequently, the plasmid hp12d-

n*rml* (n=1–6) were linearized and introduced into Po1f to form the recombinant strains

Po1f/hp12d-n*rml* (carrying n copies of *rml* expression cassette, n=1–6). Then,

Po1f/hp12d-n*rml* (n=1–6) and engineered strains from seven rounds of gene integration

were respectively inoculated into BMSY medium under the same culture conditions, and

their lipase activity was measured (Figure 5B), which revealed that the two types of

recombinant strains showed similar protein expression level. Therefore, it can be

concluded that genome editing via the developed genetic tool did not affect target protein

production of the engineering strains. When compared with the previous method (Zhou et

al. 2019), the developed genetic tool has many advantages such as utilization of reusable

marker, only requiring three plasmids, no gene redundancy, and flexible direction of gene

expression cassette. Moreover, the novel genetic tool could assemble engineering strains

harboring any copies of the target gene, only requiring increase/decrease in integration

round.

Furthermore, the strains Po1f, FY1, FY5-4*rml* 31#, and FY7-6*rml* 13# were

cultivated in YPD and BMSY media, respectively, and their growth characteristics were

tested by measuring the optical density at 600 nm (OD_{600}) at different incubation times

(Fig. 5C, D). The results revealed no observable differences in the growth curves of the

four types of strains, indicating that the genome modification in *leu2* locus via Cre/*lox-*

based genetic tool did not influence cell growth.

Figure 5. *(A)* Lipase activity of recombinant strains harboring different copies of *rml* expression cassettes. *(B)* Comparison of lipase activity of recombinant strains obtained via different methods. Data are the mean values of each kind of recombinant strain. The strains Po1f and FY1 showed low lipase activity owing to expression of autologous lipase genes. Weakly significant difference was observed between the strains FY7-6*rml*/FY7-5*rml*-rc*rml* and Po1f/hp12d-6*rml* ($P=0.044$). *(C)* Lipase activity of transformants from third-round integration. *(D)* Transcriptional level analysis of *rml* mRNA from FY3-2*rml* and FY3-*rml*-rc*rml*. The endogenous *act1* was adopted as reference gene, and the *rml* transcriptional level of FY3-2*rml* 4# was defined as 1. *(E)* Growth curves of Po1f, FY1, FY5-4*rml* 31#, and FY7-6*rml* 13# in YPD medium. *(F)* Growth curves of Po1f, FY1, FY5-4*rml* 31#, and FY7-6*rml* 13# in BMSY medium. NS, [*], [**], and [***] indicate non-significant, $P < 0.05$, $P < 0.01$, and $P < 0.001$, respectively.

Comparison of two-genotype recombinant strains

Two genotypes of the recombinant strains were generated in each round of integration (except for the first round), and their numbers were presented in Table 1. The rates of two forms of strains were approximately equal, suggesting that there was nearly equal possibility to choose a strain with certain genotype. The colonies with high lipase activities were selected for protein production evaluation. The results shown in Table 1 also implied that the protein expression levels of the recombinant strains in each round of integration were similar, which was confirmed in Figure 5A. In contrast, some previous studies have reported that the relative arrangement of multiple genes might affect corresponding transcription and protein production in host cells (Eszterhas et al. 2002; Lee et al. 2014). Thus, a total of 30 positive colonies from the third-round integration were identified via the primer pairs KNC-F2/Apa-R6 and SXB-R2/Apa-R3 (data not

shown), and their lipase activities were measured. The results presented in Figure 5E

showed that the RML activity of each recombinant randomly changed within a proper

range. Moreover, transcriptional level analysis further indicated that there was no

significant difference in the transcriptional quantity of *rml* mRNA isolated from FY3-

2*rml* and FY3-*rml*-rc*rml* cells (Figure 5F). Thus, the relative direction of *rml* expression

cassettes had no observable effect on protein production in this study, which might be

owing to the difference in the host type. Accordingly, both the genotypes of the strains

were concluded to be suitable for RML production in *Y. lipolytica*.

Successfully expanding to six different genes and *axp1-2* locus

To evaluate the universality of the novel genetic tool, six genes with varied lengths and

resources were additionally integrated into the engineered strain FY5-4*rml*. The genes

ire1, *kar2*, *pdi*, *sls1,* and *hac1* are the homologous genes of *Y. lipolytica*, and *vgb* is the

coding gene of *Vitreoscilla* hemoglobin. These genes may differentially affect RML

expression level, which determines their integration times. The corresponding plasmids

were derived from Cre-Y2 and Cre-Y3 by replacing the *rml* gene with different target

genes (Supplemental Fig. S5A), and then used to transform FY5-4*rml*. The subsequent

integration processes were similar to those of Cre-Y2 and Cre-Y3, and the proportions of

positive colonies were all above 90% (Table 2), confirming the availability of the

developed genetic tool for repeated, targeted, and markerless integration of different

genes.

Furthermore, a derived tool was established to verify whether the developed genetic

tool could be applied for other target genes, integration loci, and strains. For this purpose,

three plasmids containing *Rhizopus oryzae* lipase (*rol*) gene were respectively

constructed based on Cre-Y1, Cre-Y2, and Cre-Y3 (Supplemental Fig. S5B), and

integrated into defective *axp* gene locus (*axp1-2*) of *Y. lipolytica* Po1h. The

transformation, screening and induction, isolation of positive colonies, and identification

processes were similar to those employed for *rml* gene integration into *leu2* locus of Po1f.

First, the Cre-axp1 plasmid was linearized by *Nsi*I and inserted into the Upaxp locus to

generate an initial strain. Then, the Cre-axp2 and Cre-axp3 plasmids were digested with

*Cla*I and iteratively introduced into the *axp1-2* locus for repeated, targeted, and

markerless integration of *rol* in Po1h. As a result, three rounds of integration were

conducted and all of the integration rates were above 96%. These findings revealed that

such extension of the novel genetic tool is feasible and convenient.

Table 2. Integration rates of various genes, loci and strains in this study.

Gene	GenBank	Lenth (bp)	Integration round	Positive colonies/ all colonies	Proportion (%)
Integration locus: *leu2* gene of FY5-4*rml*, induction time: 16 h					
ire1	XM_002142952.1	3663	1	32/32	100
kar2	U63136.1	2013	1	32/32	100
			2	32/32	100
			3	32/32	100
pdi	XM_503481.1	1515	1	32/32	100
			2	32/32	100
			3	31/32	96.88
			4	31/32	96.88
sls1	Z50154.1	1281	1	32/32	100
			2	32/32	100
hac1	XM_500811.1	918	1	32/32	100
vgb	AF292694.1	441	1	32/32	100
			2	32/32	100
Integration locus: *axp1-2* of Po1h, induction time: 12 h					
rol	AF229435.1	1101	1	31/32	96.88
			2	32/32	100
			3	32/32	100

Discussion

In the present study, a Cre/*lox*-based genetic tool was precisely and skillfully designed.

The plasmids Cre-Y2 and Cre-Y3 adopted two inversely oriented *lox* sites, and the Ura3

marker cannot be removed from the genome if the strain does not undergo integration

process as expected. Thus, the Ura$^+$ strain could be distinguished by isolating positive

colonies on the MD medium. It must be noted that except for the *cre* gene and *lox* sites,

all of the genetic tool elements are nonspecific and replaceable. Any DNA sequence can

be chosen as the homologous fragment as long as no gene is disrupted, and the length of

the homologous fragment is very flexible because of the excision of unnecessary

sequences. The *rml* gene in plasmids Cre-Y2 and Cre-Y3 can be replaced by

heterologous and endogenous genes of *Y. lipolytica* to accomplish repeated markerless

integration, and the plasmid construction process is simple and convenient. In addition,

the Cre/*lox*-based genetic tool can also be applied to other genes and loci if the

corresponding plasmid elements are available.

Recently, several genetic tools have been developed for markerless gene integration

of *Y. lipolytica* based on CRISPR-Cas9 system, Cre/*lox* system and Ura/5-fluoroorotic

acid resistance method (Supplemental Fig. S6) (Schwartz et al. 2017b; Gao et al. 2018;

Holkenbrink et al. 2018; Lv et al. 2019). As shown in Supplemental Table S1, the

integration efficiency markedly fluctuates with different genome loci and single-guide

RNAs (sgRNAs) for CRISPR-Cas9-mediated gene integrations (Schwartz et al. 2017b;

Holkenbrink et al. 2018). According to most of the existing Cre/*lox*-based methods

(Holkenbrink et al. 2018; Lv et al. 2019), the DNA fragment can be integrated into the

host genome via gene insertion or gene replacement. However, insertion event may

introduce unnecessary DNA segments into the genome, whereas replacement events,

which occur occasionally, can avoid this issue, but requires substantial screening of target

transformants. Furthermore, while the Ura/5-fluoroorotic acid resistance method (Ura

pop-in/pop-out method) can also accomplish markerless gene integration, reverse

mutation is difficult to avoid in the marker recovery process. In addition, new plasmids

carrying different sgRNA or homologous fragments must be constructed to repeatedly

realize gene integration in a specific genome locus via the three above-mentioned genetic

tools. These limitations can be completely overcome by the novel genetic tool designed

in the present study. The target gene is integrated into the host genome via efficient,

targeted, and repeatable gene insertion event, and the introduced non-target sequences

can be deleted by recombination reaction of the *lox* sites. Moreover, despite the

occurrence of inefficient insertion event in the integration process, a few recombinants

can be enriched via twice auxotroph screening. The existing methods based on CRISPR-

Cas9 and Cre/*lox* systems require two different selection markers and additional plasmid

recovery procedure (Schwartz et al. 2017b; Gao et al. 2018; Holkenbrink et al. 2018; Lv

et al. 2019). In contrast, only a single marker is required for the novel Cre/*lox*-based

genetic tool, and the process of gene integration, markers reuse, and plasmid recovery can

be combined and accomplished simultaneously, which can significantly reduce the

experimental period and workload. Overall, the novel genetic tool developed in this study

can overcome many limitations such as low HR rate, limited selection marker, random

integration site, and additional plasmid recovery.

In conclusion, a powerful and efficient Cre/*lox*-based genetic tool for repeated,

targeted, and markerless gene integration was developed in this study. In practice, only a

single selection marker can enable iterative integration of target gene into the genome,

which can facilitate genetic modifications of the host strains. Furthermore, in each round

of integration, all of the unnecessary segments are removed, the relative direction of each

target gene expression cassette is controllable, and the gene recombination on genome is

foreseeable. These characteristics can significantly decrease gene redundancy in the

engineering strain, making it convenient for the use of host strain in pharmaceutical, food,

and other health-related areas. Besides, the developed genetic tool is also suitable for

various other genes and loci. The findings of this study significantly broaden the

application of Cre/*lox* system and strengthen the capabilities for genome modification in

Y. lipolytica.

Methods

Strains and media

Escherichia coli TOP10 and TOP10 F' (Shanghai Weidi Biotechnology Co., Ltd,

Shanghai, China) were applied for plasmid amplification. The plasmid hp12d-*rml* (Zhou

et al. 2019) and auxotrophic *Y. lipolytica* Po1f (Madzak et al. 2000) and Po1h (Madzak et

al. 2004) were respectively utilized for plasmid construction and transformation. The T4

DNA ligase, Phanta Max Master Mix polymerase, Rapid Taq Master Mix polymerase,

and ClonExpress II One Step Cloning Kit were obtained from Vazyme Biotech Co., Ltd

(Nanjing, China) and used for gene amplification and genome identification. The *E. coli*

and *Y. lipolytica* strains were respectively cultured at 37 °C and 28 °C. All strains and

plasmids used in this study are listed in Supplemental Table S2. The LB, YPD, MD, and

BMSY media have been previously described (Zhou et al. 2019). The MO medium (20

mL/L oleic acid, 0.5 mL/L Tween 80, 13.4 g/L yeast nitrogen base with no amino acids,

and 0.4 mg/L biotin) was derived from MD medium, and leucine (final concentration,

262 mg/L) or uracil (final concentration, 22.4 mg/L) was added for auxotrophic strains.

Plasmid construction

Figure S7 illustrates plasmid Cre-Y1 construction. The pPOX2 promoter, *cre* gene,

tandem fragment of lip2t and *lox*71, and fusion segment of Upleu and *lox*71 were

respectively cloned via corresponding primer pairs (Pox2-F1/Pox2-R1, Cre-F1/Cre-R1,

lip2t-F1/lip2t-R1, and Upleu-F1/Upleu-R1). The four amplified products were assembled

by overlap extension PCR and then double-digested with *Nde*I and *Apa*I. Subsequently,

the digested product was inserted into *Nde*I/*Apa*I-digested plasmid hp12d-*rml* to

construct plasmid 1296-*cre*. Furthermore, the *ura3* gene (amplification using Ura-

F1/Ura-R1) was subcloned into *Nde*I-linearized 1296-*cre* by HR, generating Cre-Y1. All

the plasmids were confirmed by DNA sequencing by TsingKe Biological Technology

Co., Ltd. (Wuhan, China), and all the primers and *lox* sites used in this study are

presented in Supplemental Table S3.

As shown in Supplemental Fig. S8, the plasmids Cre-Y1 and hp12d-*rml* were

respectively double-digested using *Apa*I/*Nhe*I and *Apa*I/*Xba*I, and the two digestion

products with same cohesive ends were ligated to generate the plasmid hp12d-*rml-cre*.

Subsequently, the gene segment lip2t-*lox*66 was cloned by lip2t-F1/lip2t-R2 and

recombined into *Avr*II/*Mlu*I-digested hp12d-*rml-cre*, forming hp12d-*rml-cre-lox*66.

Furthermore, the fusion fragment of rc*lox*66 and partial *leu2* was amplified by leu-F2/leu-

R2 and recombined with *Nde*I/*Apa*I-opened hp12d-*rml-cre-lox*66 to generate plasmid

Cre-Y2. Similarly, the gene segment lip2t-rc*lox*71 was amplified by lip2t-F1/lip2t-R3 and used for the construction of hp12d-*rml-cre*-rc*lox*71, and hp12d-*rml-cre*-rc*lox*71 was recombined with *lox*71-partial *leu2* to obtain plasmid Cre-Y3 (Supplemental Fig. S9).

The *rml* gene of Cre-Y2 and Cre-Y3 was replaced with six different genes (*ire1*, *kar2*, *pdi*, *sls1*, *hac1,* and *vgb*) to assemble the corresponding plasmids, and their structures are presented in Supplemental Fig. S5A. Moreover, the homologous fragment Upleu, *leu2,* and target gene *rml* were respectively substituted with Upaxp, *axp1-2,* and *rol* to generate the plasmids Cre-axp1, Cre-axp2, and Cre-axp3 (Supplemental Fig. S5B). The abovementioned procedures were achieved by conventional PCR amplification, enzyme digestion, and ligation.

Yeast transformation, screening and induction

The integration process was performed as follows (Supplemental Fig. S10). The *Sal*I-linearized plasmid Cre-Y1 was integrated into Po1f genome based on the previously described method (Chen et al. 1997). Then, the transformants were screened twice in 5 mL of MD liquid medium supplemented with leucine (MD-Leu). Subsequently, the derivatives were inoculated into MO liquid medium supplemented with leucine and uracil (MO-Leu-Ura) for inducible expression of *cre* gene. To isolate single colony, the cell suspension solution was streaked onto YPD plate, and the selected colonies were respectively transferred onto YPD and MD-Leu plates for phenotype verification, where the positive colonies could only grow on YPD medium. In addition, the genomic DNA of positive colonies was extracted for further identification via PCR amplification with appropriate primers and DNA sequencing by TsingKe Biological Technology Co., Ltd.

(Wuhan, China). Thus, the initial strain FY1 was obtained. The subsequent integration

was almost similar to the first-round integration; however, Cre-Y2 or Cre-Y3 was

linearized by *Apa*I, leucine was not required in MD or MO medium, and YPD medium

was replaced with BMSY medium containing 10 mL/L glyceryl tributyrate to select

colonies with high RML activity.

Shaking flask culture and lipase activity determination

Transformants harboring clear and large transparent halos were inoculated into 5 mL of

YPD for 24 h. Then, the cultures were transferred into 500-mL shake flasks containing 30

mL of BMSY medium and incubated for 120 h to produce RML. The RML activity in

cell culture supernatant was evaluated using previously described method (Zhou et al.

2019). All the experiments were performed in triplicate, and the results were analyzed by

ANOVA with Duncan's multiple range test at $P < 0.05$.

Determination of gene copy number and transcriptional analysis

Based on the protocol of ChanQ^{TM} Universal SYBR qPCR Master Mix (Vazyme Biotech

Co., Ltd, Nanjing, China), the *rml* gene copy numbers were determined by real-time

qPCR performed on StepOnePlus apparatus with StepOne software version 2.3 (Applied

Biosystems, Foster City, CA, USA). The endogenous *act1* was set as reference gene, and

the plasmids pMD19-*act1* and hp4d-*rml* (Zhou et al. 2019) were utilized for establishing

standard curves, respectively. The *rml* gene transcriptional level analysis was conducted

by TsingKe Biological Technology Co., Ltd (Wuhan, China). Each sample was collected

in triplicate.

Acknowledgments

We acknowledge financial support by the National Natural Science Foundation of China (Nos: 31070089, 31170078, 31971206, 41972319 and J1103514), the National High Technology Research and Development Program of China (Nos: 2013AA065805 and 2014AA093510), the Natural Science Foundation of Hubei Province (No. 2015CFA085), and the Fundamental Research Funds for HUST (Nos. 2017KFYXJJ212, 2017KFXKJC010 and 2017KFTSZZ001). We appreciate the Research Core Facilities for Life Science (HUST) for their valuable assistance in qPCR measurements. We thank International Science Editing (http://www.internationalscienceediting.com) for editing this manuscript.

Author contributions: Q.Z. and L.J. conducted the project design. Q.Z., L.J., W.L. and Y.L. carried out the experiments. Q.Z., L.J., Z.H. and H.Z. helped with the formal analysis. Q.Z., L.J., L.X. and Y.Y. conducted the manuscript preparation.

Competing interests: All authors declare that they have no competing interests.

References

Abdel-Mawgoud AM, Markham KA, Palmer CM, Liu N, Stephanopoulos G, Alper HS. 2018. Metabolic engineering in the host *Yarrowia lipolytica*. *Metab Eng* **50**:192-208. doi: 10.1016/j.ymben.2018.07.016

Albert H, Dale EC, Lee E, Ow DW. 1995. Site-specific integration of DNA into wild-type and mutant *lox* sites placed in the plant genome. *Plant J* **7**:649-659. doi: 10.1046/j.1365-313x.1995.7040649.x

Bae SJ, Park BG, Kim BG, Hahn JS. 2020. Multiplex gene disruption by targeted base editing of *Yarrowia lipolytica* genome using cytidine deaminase combined with the CRISPR/Cas9 system. *Biotechnol J* **15**:e1900238. doi: 10.1002/biot.201900238

Barth G, Gaillardin C. 1996. *Yarrowia lipolytica*. In: Nonconventional yeasts in biotechnology: a handbook. pp. 313-388. Springer, Berlin, Heidelberg.

Blazeck J, Hill A, Liu L, Knight R, Miller J, Pan A, Otoupal P, Alper HS. 2014. Harnessing *Yarrowia lipolytica* lipogenesis to create a platform for lipid and biofuel production. *Nat Commun* **5**:3131. doi: 10.1038/ncomms4131

Carly F, Vandermies M, Telek S, Steels S, Thomas S, Nicaud JM, Fickers P. 2017. Enhancing erythritol productivity in *Yarrowia lipolytica* using metabolic engineering. *Metab Eng* **42**:19-24. doi: 10.1016/j.ymben.2017.05.002

Chen DC, Beckerich JM, Gaillardin C. 1997. One-step transformation of the dimorphic yeast *Yarrowia lipolytica*. *Appl Microbiol Biotechnol* **48**:232-235. doi: 10.1007/s002530051043

Cong L, Ran FA, Cox D, Lin S, Barretto R, Habib N, Hsu PD, Wu X, Jiang W, Marraffini LA, et al. 2013. Multiplex genome engineering using CRISPR/Cas

systems. *Science* **339**:819-823. doi: 10.1126/science.1231143

Czajka JJ, Nathenson JA, Benites VT, Baidoo E, Cheng Q, Wang Y, Tang YJ. 2018.

Engineering the oleaginous yeast *Yarrowia lipolytica* to produce the aroma

compound beta-ionone. *Microb Cell Fact* **17**:136. doi: 10.1186/s12934-018-0984-x

Eszterhas SK, Bouhassira EE, Martin DI, Fiering S. 2002. Transcriptional interference by

independently regulated genes occurs in any relative arrangement of the genes and is

influenced by chromosomal integration position. *Mol Cell Biol* **22**:469-479. doi:

10.1128/mcb.22.2.469-479.2002

Fickers P, Le Dall MT, Gaillardin C, Thonart P, Nicaud JM. 2003. New disruption

cassettes for rapid gene disruption and marker rescue in the yeast *Yarrowia*

lipolytica. *J Microbiol Methods* **55**:727-737. doi: 10.1016/j.mimet.2003.07.003

Gaj T, Gersbach CA, Barbas CR. 2013. ZFN, TALEN, and CRISPR/Cas-based methods

for genome engineering. *Trends Biotechnol* **31**:397-405. doi:

10.1016/j.tibtech.2013.04.004

Gao D, Smith S, Spagnuolo M, Rodriguez G, Blenner M. 2018. Dual CRISPR-Cas9

cleavage mediated gene excision and targeted integration in *Yarrowia lipolytica*.

Biotechnol J **13**:e1700590. doi: 10.1002/biot.201700590

Gueldener U, Heinisch J, Koehler GJ, Voss D, Hegemann JH. 2002. A second set of *lox*P

marker cassettes for Cre-mediated multiple gene knockouts in budding yeast.

Nucleic Acids Res **30**:e23. doi: 10.1093/nar/30.6.e23

Hentges P, Van Driessche B, Tafforeau L, Vandenhaute J, Carr AM. 2005. Three novel

antibiotic marker cassettes for gene disruption and marker switching in

Schizosaccharomyces pombe. *Yeast* **22**:1013-1019. doi: 10.1002/yea.1291

Hochrein L, Mitchell LA, Schulz K, Messerschmidt K, Mueller-Roeber B. 2018. L-

SCRaMbLE as a tool for light-controlled Cre-mediated recombination in yeast. *Nat*

Commun **9**:1931. doi: 10.1038/s41467-017-02208-6

Holkenbrink C, Dam MI, Kildegaard KR, Beder J, Dahlin J, Domenech BD, Borodina I.

2018. EasyCloneYALI: CRISPR/Cas9-based synthetic toolbox for engineering of

the yeast *Yarrowia lipolytica*. *Biotechnol J* **13**:e1700543. doi:

10.1002/biot.201700543

Hsu PD, Lander ES, Zhang F. 2014. Development and applications of CRISPR-Cas9 for

genome engineering. *Cell* **157**:1262-1278. doi: 10.1016/j.cell.2014.05.010

Kawano F, Okazaki R, Yazawa M, Sato M. 2016. A photoactivatable Cre-*lox*P

recombination system for optogenetic genome engineering. *Nat Chem Biol* **12**:1059-

1064. doi: 10.1038/nchembio.2205

Kuhn R, Torres RM. 2002. Cre/*lox*P recombination system and gene targeting. *Methods*

Mol Biol **180**:175-204. doi: 10.1385/1-59259-178-7:175

Larroude M, Celinska E, Back A, Thomas S, Nicaud JM, Ledesma-Amaro R. 2018. A

synthetic biology approach to transform *Yarrowia lipolytica* into a competitive

biotechnological producer of beta-carotene. *Biotechnol Bioeng* **115**:464-472. doi:

10.1002/bit.26473

Larroude M, Rossignol T, Nicaud JM, Ledesma-Amaro R. 2018. Synthetic biology tools

for engineering *Yarrowia lipolytica*. *Biotechnol Adv* **36**:2150-2164. doi:

10.1016/j.biotechadv.2018.10.004

Ledesma-Amaro R, Nicaud JM. 2016. *Yarrowia lipolytica* as a biotechnological chassis

to produce usual and unusual fatty acids. *Prog Lipid Res* **61**:40-50. doi:

10.1016/j.plipres.2015.12.001

Lee TJ, Parikh RY, Weitz JS, Kim HD. 2014. Suppression of expression between

adjacent genes within heterologous modules in yeast. *G3 (Bethesda)* **4**:109-116. doi:

10.1534/g3.113.007922

Leibig M, Krismer B, Kolb M, Friede A, Gotz F, Bertram R. 2008. Marker removal in

staphylococci via Cre recombinase and different *lox* sites. *Appl Environ Microbiol*

74:1316-1323. doi: 10.1128/AEM.02424-07

Lv Y, Edwards H, Zhou J, Xu P. 2019. Combining 26s rDNA and the Cre-*lox*P system

for iterative gene integration and efficient marker curation in *Yarrowia lipolytica*.

ACS Synth Biol **8**:568-576. doi: 10.1021/acssynbio.8b00535

Madzak C. 2015. *Yarrowia lipolytica*: recent achievements in heterologous protein

expression and pathway engineering. *Appl Microbiol Biotechnol* **99**:4559-4577. doi:

10.1007/s00253-015-6624-z

Madzak C. 2018. Engineering *Yarrowia lipolytica* for use in biotechnological

applications: a review of major achievements and recent innovations. *Mol*

Biotechnol **60**:621-635. doi: 10.1007/s12033-018-0093-4

Madzak, C, Gaillardin C, Beckerich J M. 2004. Heterologous protein expression and

secretion in the non-conventional yeast *Yarrowia lipolytica*: a review. *J Biotechnol*

109(1-2): p. 63-81. doi: 10.1016/j.jbiotec.2003.10.027

Madzak C, Treton B, Blanchin-Roland S. 2000. Strong hybrid promoters and integrative

expression/secretion vectors for quasi-constitutive expression of heterologous

proteins in the yeast *Yarrowia lipolytica*. *J Mol Microbiol Biotechnol* **2**:207-216.

PMID: 10939246

Markham KA, Palmer CM, Chwatko M, Wagner JM, Murray C, Vazquez S,

Swaminathan A, Chakravarty I, Lynd NA, Alper HS. 2018. Rewiring *Yarrowia*

lipolytica toward triacetic acid lactone for materials generation. *Proc Natl Acad Sci*

USA **115**:2096-2101. doi: 10.1073/pnas.1721203115

Pignede G, Wang HJ, Fudalej F, Seman M, Gaillardin C, Nicaud JM. 2000. Autocloning

and amplification of LIP2 in *Yarrowia lipolytica*. *Appl Environ Microbiol* **66**:3283-

3289. doi: 10.1128/aem.66.8.3283-3289.2000

Richardson SM, Mitchell LA, Stracquadanio G, Yang K, Dymond JS, DiCarlo JE, Lee D,

Huang CL, Chandrasegaran S, Cai Y, et al. 2017. Design of a synthetic yeast

genome. *Science* **355**:1040-1044. doi: 10.1126/science.aaf4557

Schwartz C, Frogue K, Ramesh A, Misa J, Wheeldon I. 2017a. CRISPRi repression of

nonhomologous end-joining for enhanced genome engineering via homologous

recombination in *Yarrowia lipolytica*. *Biotechnol Bioeng* **114**:2896-2906. doi:

10.1002/bit.26404

Schwartz C, Shabbir-Hussain M, Frogue K, Blenner M, Wheeldon I. 2017b. Standardized

markerless gene integration for pathway engineering in *Yarrowia lipolytica*. *ACS*

Synth Biol **6**:402-409. doi: 10.1021/acssynbio.6b00285

Suzuki N, Nonaka H, Tsuge Y, Inui M, Yukawa H. 2005. New multiple-deletion method

for the *Corynebacterium glutamicum* genome, using a mutant *lox* sequence. *Appl*

Environ Microbiol **71**:8472-8480. doi: 10.1128/AEM.71.12.8472-8480.2005

Wu X, Scott DA, Kriz AJ, Chiu AC, Hsu PD, Dadon DB, Cheng AW, Trevino AE,

Konermann S, Chen S, et al. 2014. Genome-wide binding of the CRISPR

endonuclease Cas9 in mammalian cells. *Nat Biotechnol* **32**:670-676. doi:

10.1038/nbt.2889

Wu Y, Zhu RY, Mitchell LA, Ma L, Liu R, Zhao M, Jia B, Xu H, Li YX, Yang ZM, et al.

2018. In vitro DNA SCRaMbLE. *Nat Commun* **9**:1935. doi: 10.1038/s41467-018-

03743-6

Yan X, Yu HJ, Hong Q, Li SP. 2008. Cre/*lox* system and PCR-based genome engineering

in *Bacillus subtilis*. *Appl Environ Microbiol* **74**:5556-5562. doi:

10.1128/AEM.01156-08

Yang K, Qiao Y, Li F, Xu Y, Yan Y, Madzak C, Yan J. 2019. Subcellular engineering of

lipase dependent pathways directed towards lipid related organelles for highly

effectively compartmentalized biosynthesis of triacylglycerol derived products in

Yarrowia lipolytica. *Metab Eng* **55**:231-238. doi: 10.1016/j.ymben.2019.08.001

Zhou Q, Jiao L, Qiao Y, Wang Y, Xu L, Yan J, Yan Y. 2019. Overexpression of GRAS

Rhizomucor miehei lipase in *Yarrowia lipolytica* via optimizing promoter, gene

dosage and fermentation parameters. *J Biotechnol* **306**:16-23. doi:

10.1016/j.jbiotec.2019.09.004