1

A novel and effective Cre/*lox*-based genetic tool for repeated, targeted and markerless gene integration

Qinghua Zhou [†], Liangcheng Jiao[†], Wenjuan Li, Zhiming Hu, Yunchong Li, Houjin Zhang, Li Xu [*], Yunjun Yan[*]

Key Laboratory of Molecular Biophysics of the Ministry of Education, College of Life Science and Technology, Huazhong University of Science and Technology, Wuhan 430074, P. R. China

[†]These authors contributed equally to this work

*Corresponding author

E-mail: xuli@mail.hust.edu.cn (L. Xu), yanyunjun@hust.edu.cn (Y. Yan)

Highlights

● A powerful genetic tool was developed for targeted and markerless gene integration

● The genetic tool was effective for various genes, loci, and strains

● Only a single selection marker was adequate for repeated gene integration

● All of the useless and redundant sequences would be excised

● All integration efficiencies could remain above 90%

Abstract

The unconventional yeast *Yarrowia lipolytica* is extensively applied in bioproduction

fields owing to its excellent metabolite and protein production ability. Nonetheless,

utilization of this promising host is still restricted by limited availability of precise and

effective gene integration tools. In this study, a novel and efficient genetic tool was

developed for targeted, repeated, and markerless gene integration based on Cre/*lox* site-

specific recombination system. The developed tool required only a single selection

marker and could completely excise all of the unnecessary sequences. A total of three

plasmids were created and seven rounds of marker-free gene integration were examined

in *Y. lipolytica*. All the integration efficiencies remained above 90%, and analysis of

protein production and growth characteristics of the engineered strains confirmed that

genome modification via the novel genetic tool was feasible. Further work also

confirmed the genetic tool was effective for integration of other genes, loci, and strains.

Thus, this study significantly promotes the application of Cre/*lox* system and presents a

powerful tool for genome engineering in *Y. lipolytica*.

Keywords

Genetic tool, Cre/*lox*, repeatedly targeted and markerless integration, *Yarrowia lipolytica*

Graphical abstract

Plasmid design   **Repeated, targeted and markerless integration**

The novel genetic tool comprised three plasmids, namely, Cre-Y1, Cre-Y2, and Cre-Y3.

Cre-Y1 was introduced into the Upleu locus of Po1f, and then recombination between

two *lox*71 sites could remove unnecessary plasmid elements and produce an initial strain

harboring a *lox*71 site. Subsequently, Cre-Y2 was integrated onto the *leu2* locus, forming

a transitional strain with three *lox* sites (*lox*71, *lox*66, and rc*lox*66). Recombination

reactions among these three *lox* sites could excise unnecessary DNA fragments and retain

a *lox*72 site, an rc*lox*66 site, and a target gene expression cassette in the genome.

Similarly, integration of Cre-Y3 produced strains harboring two target gene expression

cassettes. Consequently, repeated, targeted, and markerless gene integration could be

efficiently realized following iterative integration of Cre-Y2 and Cre-Y3.

1. Introduction

With the development of synthetic biology, various molecular genetic tools are being increasingly designed for precise genome modification in yeasts. As a promising host, the generally regarded as safe (GRAS) yeast, *Yarrowia lipolytica*, has been employed in a wide range of fields such as protein production, metabolic engineering, degradation of *n*-alkanes, and dimorphism (Abdel-Mawgoud et al., 2018; Blazeck et al., 2014; Carly et al., 2017; Czajka et al., 2018; Madzak, 2015; Madzak, 2018). Nevertheless, only few effective genetic tools have been developed for repeatedly targeted gene integration in this host, which still restricts further exploitation and application of *Y. lipolytica*.

In traditional methods, heterologous and endogenous genes are transformed into *Y. lipolytica* genome via insertion of the whole plasmid at target loci or integration of tandem fragment of target gene and selection marker at zeta loci (Barth and Gaillardin, 1996; Larroude et al., 2018; Pignede et al., 2000). However, introduction of non-target DNA fragments, especially antibiotics resistance genes, may harm the GRAS status of *Y. lipolytica* and hinder its application in food and medicine industries. Besides, the bioproduction of *Y. lipolytica* usually requires multi-round integrations of target gene or co-expression of various genes (Carly et al., 2017; Larroude et al., 2018; Ledesma-Amaro and Nicaud, 2016; Markham et al., 2018; Yang et al., 2019), which can be hardly realized via these conventional methods owing to the limited availability of selection markers. Consequently, there is an urgent need to develop efficient genetic tool(s) for targeted, repeatable, and markerless gene integration in *Y. lipolytica*.

To date, several site-specific nuclease technologies have been established for genome modification in various species (Cong et al., 2013; Gaj et al., 2013; Hsu et al.,

2014; Wu et al., 2014). Among them, CRISPR-Cas9 system can accomplish relatively

high-efficiency gene disruption and marker-free gene integration in *Y. lipolytica* (Bae et

al., 2020; Holkenbrink et al., 2018; Schwartz et al., 2017a). A previous study examined

17 loci for targeted and markerless gene integration in *Y. lipolytica,* among which only

five loci exhibited gene integration, with highest integration efficiency of 68.9%±25.5%

(Schwartz et al., 2017b). Another study accomplished gene integration into *PEX*10 locus

via homologous recombination (HR) and homology-mediated end-joining, with

integration efficiencies of 6.7%±3.6% and 37.5%±8.8%, respectively (Gao et al., 2018).

These studies indicate the need for further improvement in achieving more desirable gene

integration efficiency (similar to gene disruption efficiency of about 90%) via CRISPR-

Cas9-based tools. Besides, gene insertion via HR-mediated double-strand break (DSB)

repair usually demands synchronous introduction of two plasmids harboring different

selection markers (Gao et al., 2018; Schwartz et al., 2017b), resulting in reduction of

transformation efficiency and additional procedure for plasmids recovery.

The Cre/*lox*-site-specific recombination system derived from bacteriophage P1 is

known to be appropriate for genome editing (Kawano et al., 2016; Kuhn and Torres,

2002; Richardson et al., 2017; Wu et al., 2018), and its action mechanism has been

clearly described (Albert et al., 1995; Leibig et al., 2008; Yan et al., 2008). Cre

recombinase can recognize and bind to *lox*P site, and can catalyze molecular

recombination reactions including insertion, translocation, inversion and deletion (Albert

et al., 1995). However, these reactions can retain redundant *lox*P sites in the genome,

which may disturb further recombination (Albert et al., 1995). To address this issue,

some mutational *lox* sites have been successfully designed (Leibig et al., 2008; Yan et al.,

2008). For example, the recombination between a left element-mutant *lox* site (LE) *lox*71

and a right element-mutant *lox* site (RE) *lox*66 can generate a double element-mutant *lox*

site (LE+RE) *lox*72 and a *lox*P site (Leibig et al., 2008; Yan et al., 2008). Meanwhile,

*lox*P site can be deleted and *lox*72 site can be retained in the genome through rational

design. In particular, the *lox*72 site, which can weakly bind to Cre recombinase, is less

likely to participate in subsequent recombination (Leibig et al., 2008; Yan et al., 2008).

As a result, *lox* mutants can exhibit enhanced controllability of recombination and

accelerate the utilization of Cre/*lox* system in yeasts (Fickers et al., 2003; Gueldener et al.,

2002; Hentges et al., 2005; Hochrein et al., 2018). Although this method can employ

marker-free genome modification, it often presents low efficiency or is time-consuming.

In addition, a few studies have attempted markerless gene integration via Cre/*lox* system

in *Y. lipolytica*. Among them, an inspired method combining Cre/*lox* system and 26S

rDNA achieved 79.2% integration efficiency in *Y. lipolytica* (Lv et al., 2019); however,

the integration loci were random and the copy numbers of target genes were

uncontrollable, requiring lot of time for selection and evaluation of transformants.

Moreover, to rescue selection marker in the genome, an additional plasmid harboring *cre*

gene must be introduced into the recombinants by another transformation, and this

plasmid should subsequently be recovered. These tedious procedures require two

different selection markers and further extend the experimental period.

Therefore, in the present study, a novel and effective genetic tool was designed

based on Cre/*lox* system. Only a single selection marker was adequate to enable repeated,

targeted, and markerless gene integration, and all unnecessary fragments could be totally

excised. To demonstrate the efficacy of the developed tool, an important lipase gene from

Rhizomucor miehei (*rml*) was chosen as a qualitative reporter, and three new plasmids

were constructed. A total of seven rounds of marker-recyclable integration were

implemented, and the positive colony proportions reached 90.63%–100%. The

experiment conditions, feasibility, and controllability of the genetic tool were evaluated

and optimized. Moreover, to confirm the universality of the developed tool, other six

different genes were further integrated into *leu2* locus, and a derived tool was

successfully employed to integrate another target gene into the *axp1-2* locus. The results

obtained revealed that Cre/*lox*-based genetic tool could resolve the limitations of the

existing methods and bring significant advancement in gene editing in yeast species.

2. Material and methods

2.1 Strains and media

Escherichia coli TOP10 and TOP10 F' (Shanghai Weidi Biotechnology Co., Ltd,

Shanghai, China) were applied for plasmid amplification. The plasmid hp12d-*rml* (Zhou

et al., 2019) and auxotrophic *Y. lipolytica* Po1f (Madzak et al., 2000) and Po1h (Madzak

et al., 2004) were respectively utilized for plasmid construction and transformation. The

T4 DNA ligase, Phanta Max Master Mix polymerase, Rapid Taq Master Mix polymerase,

and ClonExpress II One Step Cloning Kit were obtained from Vazyme Biotech Co., Ltd

(Nanjing, China) and used for gene amplification and genome identification. The *E. coli*

and *Y. lipolytica* strains were respectively cultured at 37 °C and 28 °C. All strains and

plasmids used in this study are listed in Supplemental Table S1. The LB, YPD, MD, and

BMSY media have been previously described (Zhou et al., 2019). The MO medium (20

mL/L oleic acid, 0.5 mL/L Tween 80, 13.4 g/L yeast nitrogen base with no amino acids,

and 0.4 mg/L biotin) was derived from MD medium, and leucine (final concentration,

262 mg/L) or uracil (final concentration, 22.4 mg/L) was added for auxotrophic strains.

2.2 Plasmid construction

Supplemental Fig. S1 illustrates plasmid Cre-Y1 construction. The pPOX2 promoter, *cre*

gene, tandem fragment of lip2t and *lox*71, and fusion segment of Upleu and *lox*71 were

respectively cloned via corresponding primer pairs (Pox2-F1/Pox2-R1, Cre-F1/Cre-R1,

lip2t-F1/lip2t-R1, and Upleu-F1/Upleu-R1). The four amplified products were assembled

by overlap extension PCR and then double-digested with *Nde*I/*Apa*I. Subsequently, the

digested product was inserted into *Nde*I/*Apa*I-digested plasmid hp12d-*rml* to construct

plasmid 1296-*cre*. Furthermore, the *ura3* gene (amplification using Ura-F1/Ura-R1) was

subcloned into *Nde*I-linearized 1296-*cre* by HR, generating Cre-Y1. All the plasmids

were confirmed by DNA sequencing by TsingKe Biological Technology Co., Ltd.

(Wuhan, China), and all the primers and *lox* sites used in this study are presented in

Supplemental Table S2.

As shown in Supplemental Fig. S2, the plasmids Cre-Y1 and hp12d-*rml* were

respectively double-digested using *Apa*I/*Nhe*I and *Apa*I/*Xba*I, and the two digestion

products with same cohesive ends were ligated to generate the plasmid hp12d-*rml-cre*.

Subsequently, the gene segment lip2t-*lox*66 was cloned by lip2t-F1/lip2t-R2 and

recombined into *Avr*II/*Mlu*I-digested hp12d-*rml-cre*, forming hp12d-*rml-cre-lox*66.

Furthermore, the fusion fragment of rc*lox*66 and partial *leu2* was amplified by leu-F2/leu-

R2 and recombined with *Nde*I/*Apa*I-opened hp12d-*rml-cre-lox*66 to generate plasmid

Cre-Y2. Similarly, the gene segment lip2t-rc*lox*71 was amplified by lip2t-F1/lip2t-R3 and

used for the construction of hp12d-*rml-cre*-rc*lox*71, and hp12d-*rml-cre*-rc*lox*71 was

recombined with *lox*71-partial *leu2* to obtain plasmid Cre-Y3 (Supplemental Fig. S3).

The *rml* gene of Cre-Y2 and Cre-Y3 was replaced with six different genes (*ire1*,

kar2, *pdi*, *sls1*, *hac1,* and *vgb*) to assemble the corresponding plasmids, and their

structures are presented in Supplemental Fig. S4A. Moreover, the homologous fragment

Upleu, *leu2,* and target gene *rml* were respectively substituted with Upaxp, *axp1-2,* and

rol to generate the plasmids Cre-axp1, Cre-axp2, and Cre-axp3 (Supplemental Fig. S4B).

The abovementioned procedures were achieved by conventional PCR amplification,

enzyme digestion, and ligation.

2.3 Yeast transformation, screening and induction

The integration process was performed as follows (Supplemental Fig. S5). The *Sal*I-

linearized plasmid Cre-Y1 was integrated into Po1f genome based on the previously

described method (Chen et al., 1997). Then, the transformants were screened twice in 5

mL of MD liquid medium supplemented with leucine (MD-Leu). Subsequently, the

derivatives were inoculated into MO liquid medium supplemented with leucine and uracil

(MO-Leu-Ura) for inducible expression of *cre* gene. To isolate single colony, the cell

suspension solution was streaked onto YPD plate, and the selected colonies were

respectively transferred onto YPD and MD-Leu plates for phenotype verification, where

the positive colonies could only grow on YPD medium. In addition, the genomic DNA of

positive colonies was extracted for further identification via PCR amplification with

appropriate primers and DNA sequencing by TsingKe Biological Technology Co., Ltd.

(Wuhan, China). Thus, the initial strain FY1 was obtained. The subsequent integration

was almost similar to the first-round integration; however, Cre-Y2 or Cre-Y3 was

linearized by *Apa*I, leucine was not required in MD or MO medium, and YPD medium

was replaced with BMSY medium containing 10 mL/L glyceryl tributyrate to select

colonies with high RML activity.

2.4 Shaking flask culture and lipase activity determination

Transformants harboring clear and large transparent halos were inoculated into 5 mL of

YPD for 24 h. Then, the cultures were transferred into 500-mL shake flasks containing 30

mL of BMSY medium and incubated for 120 h to produce RML. The RML activity in

cell culture supernatant was evaluated using previously described method (Zhou et al.,

2019). All the experiments were performed in triplicate, and the results were analyzed by

ANOVA with Duncan's multiple range test at $P < 0.05$.

2.5 Determination of gene copy number and transcriptional analysis

Based on the protocol of ChanQ^{TM} Universal SYBR qPCR Master Mix (Vazyme Biotech

Co., Ltd, Nanjing, China), the *rml* gene copy numbers were determined by real-time

qPCR performed on StepOnePlus apparatus with StepOne software version 2.3 (Applied

Biosystems, Foster City, CA, USA). The endogenous *act1* was set as reference gene, and

the plasmids pMD19-*act1* and hp4d-*rml* (Zhou et al., 2019) were utilized for establishing

standard curves, respectively. The *rml* gene transcriptional level analysis was conducted

by TsingKe Biological Technology Co., Ltd (Wuhan, China). Each sample was collected

in triplicate.

3. Results

3.1 Genetic tool design and integration process analysis

The novel Cre/*lox*-based genetic tool comprised three plasmids, namely, Cre-Y1, Cre-Y2,

and Cre-Y3. Plasmid Cre-Y1 was applied for generating the initial strain, and plasmids

Cre-Y2 and Cre-Y3 were employed to iteratively introduce target genes into the host

strain. Cre-Y1 consisted of a yeast selection marker, *cre* gene expression cassette,

bacterial element (antibiotic resistance gene and origin of replication), homologous

fragment of target locus in the genome (HF-A), and two convergently oriented *lox*

mutants. The *cre* expression cassette should adopt an inducible promoter to control the

production of Cre recombinase. Cre-Y2 was derived from Cre-Y1 by replacing HF-A and

two convergently oriented *lox* mutants with another homologous fragment of target locus

(HF-B), a *lox* mutant and its reverse complement (rc*lox*), and a target gene expression

cassette. To avoid excising original segments from the genome, HF-B should be close to

HF-A. Cre-Y3 was similar to Cre-Y2, but with the corresponding *lox* mutants of Cre-Y2

(e.g., LE for Cre-Y2 and RE for Cre-Y3). All of the tool elements were flexible and

nonspecific; thus, the Cre/*lox*-based genetic tool was appropriate for different species.

In the present study, the plasmid elements used for *Y. lipolytica* were as follows: the

auxotroph Ura3 (UniProtKB, Q12724) was adopted as selection marker, *cre* expression

cassette was assembled as pPOX2-*cre*-lip2t (pPOX2 promoter, *cre* gene, and *lip2*

terminator, GenBank Accession Nos. AJ001300.1, AB449974.1, and AJ012632.1),

bacterial element (Amp-ori) and target gene expression cassette (*rml*) were subcloned

from plasmid hp12d-*rml*, and HF-B and HF-A were respectively designed as *leu2* gene

(GenBank Accession No. AF260230.1) and its upstream segment (Upleu) for the

convenience of plasmid construction and linearization. In addition, two *lox*71 sites, a

*lox*66 site and a reverse complement of *lox*66 site (rc*lox*66) and a reverse complement of

*lox*71 site (rc*lox*71) and a *lox*71 site were used in plasmids Cre-Y1, Cre-Y2, and Cre-Y3,

respectively.

To demonstrate repeated and markerless gene integration in *Y. lipolytica* via Cre/*lox*-

based genetic tool, the experimental process was as follows. First, Cre-Y1 containing two

*lox*71 sites was introduced into the Upleu locus of Po1f to construct an initial strain

harboring a single *lox*71 site. Under the condition of suitable screening and induction,

recombination between two *lox*71 sites could remove unnecessary plasmid elements and

produce the initial strain. Then, Cre-Y2 was integrated into *leu2* locus of the initial strain,

forming a transitional strain harboring three *lox* sites (*lox*71, *lox*66, and rc*lox*66) in the

genome. Various recombination reactions among these three *lox* sites could excise

unnecessary DNA fragments and retain a *lox*72 site, an rc*lox*66 site, and a target gene

expression cassette in the genome. Meanwhile, two types of stable strains were

eventually generated. Similarly, integration of Cre-Y3 produced strains harboring two

target gene expression cassettes. Consequently, repeated, targeted, and markerless gene

integration could be efficiently realized following iterative integration of Cre-Y2 and

Cre-Y3.

3.2 First-round integration to construct initial strain FY1

To construct strain harboring a single *lox*71 fragment, the plasmid Cre-Y1 was inserted

into auxotrophic host Po1f (Fig. 1). The transformed cell pellet was screened twice in 5

mL of MD-Leu medium to enrich the Ura$^+$ recombinant strain Po1f/Cre-Y1 (Ura$^+$ and

Leu⁻) and avoid interference from untransformed Po1f cells (Supplemental Fig. S5).

Subsequently, the culture solution containing Po1f/Cre-Y1 was transferred into MO-Leu-

Ura medium for 12 h with oleic acid as the sole carbon source, which could activate

pPOX2 to induce the expression of *cre* gene. In this case, the produced Cre recombinase

could mediate deletion reaction between two repeat *lox*71 sites in the Po1f/Cre-Y1

genome, causing excision of the bacterial element (Amp-ori), selection marker (Ura3),

and *cre* expression cassette, further generating Ura⁻ recombinants. Meanwhile, the extra

added uracil allowed Ura⁻ recombinants to grow normally. The induced cell suspension

was streaked on YPD plate for isolating single colony with Ura⁻ genotype, and the

phenotype of the isolated colonies were determined on YPD and MD-Leu media. As

shown in Supplemental Fig. S6A, a total of 32 recombinants were randomly chosen and

all of them could not grow on MD-Leu plate, proving that the selection marker Ura3 was

recycled successfully. Moreover, 10 positive colonies were picked up randomly and

identified by genome PCR (Supplemental Fig. S7A), and the PCR product obtained via

primers Upleu-F3/Upleu-R3 was further sequenced. The results shown in Supplemental

Fig. S8A indicated that only a single *lox*71 site was retained in Upleu locus of the

eventually obtained strain (named FY1). This finding also revealed that all recombination

and integration events occurred as expected (Fig. 1), and the adopted plasmid elements of

the novel genetic tool were well suitable for *Y. lipolytica*.

Fig. 1. Schematic diagram of first-round integration to obtain initial strain FY1. The Cre-Y1 was linearized at homologous fragment Upleu and introduced into Po1f. The Po1f/Cre-Y1 was screened twice in MD-Leu medium and then inoculated into MO-Leu-Ura medium for inducible expression of *cre* gene. The bacterial element (Amp-ori), selection marker (Ura3) and *cre* expression cassette (pPOX2-*cre*-lip2t) were removed by deletion reaction of two *lox*71 sites, retaining a single *lox*71 site in the FY1 genome. Primers used for PCR amplification are labeled in this figure and described in Supplemental Table S2.

3.3 Second-round integration and analysis

To evaluate whether the designed plasmids could realize markerless gene integration,

Cre-Y2 was linearized within *leu2* and introduced into FY1, to obtain the strain FY1/Cre-

Y2. As shown in Fig. 2, three *lox* sites (*lox*71, *lox*66, and rc*lox*66) were noted in the

FY1/Cre-Y2 genome, and three kinds of recombination reactions could occur when Cre

recombinase was expressed. In the first-type recombination (a), deletion reaction

occurred between the *lox*71 and *lox*66 sites in the FY1/Cre-Y2 genome, and the strain

FY2-*rml* carrying *rml* expression cassette was obtained. In the second-type

recombination (b), the *lox*71 and rc*lox*66 sites were combined with Cre proteins,

producing a *lox*72 site and reverse complement of *lox*P site (rc*lox*P). Here, the *lox*72 site

was hardly bound to Cre recombinase, and the rc*lox*P site could not stably coexist with

the unreacted rc*lox*66 site in the genome, which could recombine with Cre recombinase

again, generating FY2-rc*rml*, a derived strain carrying the reverse complement of *rml*

gene expression cassette (rc*rml*). In the third-type recombination (c), the *lox*66 and

rc*lox*66 sites were recombined for inversion reaction, forming a transitional strain that

also contained three *lox* sites similar to those of FY1/Cre-Y2. The recombination reaction

among the three *lox* sites continued until the first or second type of recombination

occurred, and the strain FY2-*rml* or FY2-rc*rml* was finally generated. Consequently, two

kinds of stable strains were obtained from three types of recombination reactions in

second-round integration. Here, all of the unnecessary and redundant fragments,

including Upleu, *leu2-270*, bacterial element, selection marker, and *cre* expression

cassette, were completely deleted, and only gene expression cassette and rc*lox*66 site

remained in the genome.

Fig. 2. Schematic diagram of second-round integration to generate FY2-*rml*/FY2-rc*rml*. The Cre-Y2

was linearized on *leu2* and integrated into the FY1 genome to obtain recombinant strain FY1/Cre-Y2.

The FY1/Cre-Y2 was screened twice in MD medium and then induced in MO-Ura medium for the

production of Cre protein. Therein, three kinds of different recombination reactions for *lox* sites were

noted in the genome: deletion reaction between *lox*71 and *lox*66 sites (a), inversion reaction between

*lox*71 and rc*lox*66 sites (b), and inversion reaction between *lox*66 and rc*lox*66 sites (c). Eventually,

two forms of stable strains FY2-*rml* and FY2-rc*rml* were produced. Meanwhile, all of the unnecessary

and redundant sequences were excised, the *rml* expression cassette was integrated into the genome,

and the remaining rc*lox*66 site locus was employed in third-round integration. Primers used for PCR

amplification are labeled in this figure and described in Supplemental Table S2.

324 **3.4 Induction time optimization in second-round integration**

In the present study, oleic acid induced pPOX2 promoter controlled the production of Cre

recombinase in MO liquid medium. Therein, induction time affected Cre protein

concentration in the cell, and further influenced the recombination efficiency in the

engineered strains. In the first-round gene integration, there were two *lox* sites in the

yeast genome. The recombinants were induced for sufficiently long time to obtain high

proportion of positive colonies. However, in subsequent gene integrations, there were

three or more *lox* sites in the genome and the recombination reaction was complex and

diverse. As reported in previous studies (Albert et al., 1995; Suzuki et al., 2005), high

concentration Cre protein is more likely to mediate recombination in *lox*72 site, resulting

in the removal of the target gene expression cassette from the engineering strain genome.

Therefore, induction of *cre* gene expression should be controlled in an appropriate time

range. In the present study, to determine the optimal induction time, the recombinant

strain FY1/Cre-Y2 was incubated in MO-Ura liquid medium for 4–16 h and streaked

onto YPD medium every 2 h (approximately one cell generation cycle). Furthermore, 32

recombinants were randomly selected from each YPD medium and then transferred onto

MD and BMSY media. The proportions of positive colonies under different induction

times were examined, and the results are shown in Table 1 and Supplemental Fig. S6B.

All the selected recombinants could complete recombination reaction after 12 h of

induction. For more accurate confirmation, 64 colonies induced for 12 h were chosen and

none of them could normally grow on MD medium. Consequently, the induction time

was set as 12 h, which could be further increased if the proportion of positive colonies

decreased in multi-round integrations.

Table 1. Positive colony proportion, genotype rate, and *rml* copy number determined in this study.

Integration round	Induction time (h)	Positive colonies/ all colonies	Proportion (%)	Selected colonies		
				Genotype	Rate	*rml* copy number
1	12	32/32	100	FY1	-	0
2	4	2/32	6.25	FY2-*rml*, FY2-rc*rml*	5/5	0.93
	6	2/32	6.25			
	8	14/32	43.75			
	10	24/32	75			
	12	32/32	100			
	14	32/32	100			
3	12	32/32	100	FY3-2*rml*, FY3-*rml*-rc*rml*	17/13	1.99
4	12	31/32	96.88	FY4-3*rml*, FY4-2*rml*-rc*rml*	4/2	3.10
5	12	29/32	90.63	FY5-4*rml*, FY5-3*rml*-rc*rml*	3/3	4.08
6	12	17/32	53.12	FY6-5*rml*, FY6-4*rml*-rc*rml*	4/2	5.12
	14	28/32	87.5			
	16	32/32	100			
7	16	32/32	100	FY7-6*rml*, FY7-5*rml*-rc*rml*	3/3	5.94

A total of 10 positive colonies were randomly chosen for identification via PCR

sequencing (Supplemental Fig. S7B). The *rml*/rc*rml* expression cassette and rc*lox*66 site

in all the selected colonies matched with the genome of FY2-*rml*/FY2-rc*rml*

(Supplemental Figs. S8B, C), demonstrating that the second-round gene integration was

performed according to the predictive analysis (Fig. 2). In addition, for the convenience

of subsequent protein production evaluation, strains that do not carry rc*rml* expression

cassette were employed for further gene integration, because the existing plasmids and

method (Zhou et al., 2019) are more applicable to construct strains only harboring *rml*

expression cassettes.

361 **3.5 Iterative insertion of Cre-Y3 and Cre-Y2**

To achieve third-round integration, Cre-Y3 was utilized to transform FY2, and the

schematic diagram is presented in Fig. 3. Similar to the second-round integration, there

were three *lox* mutants (rc*lox*66, rc*lox*71, and *lox*71) and a *lox*72 site in FY2-*rml*/Cre-Y3

genome, resulting in three different recombination reactions and eventually generating

two genotypes of the recombinant strains (FY3-2*rml* and FY3-*rml*-rc*rml*). Subsequently,

FY3-2*rml* and FY3-*rml*-rc*rml* were distinguished by PCR amplification with primer pairs

KNC-F2/Apa-R6 and SXB-R2/Apa-R3 (Supplemental Fig. S7C). The obtained products

were further sequenced and the results (Supplemental Figs. S8D, E) were consistent with

the prediction presented in Fig. 3. Thus, another copy of *rml* gene was targeted on *leu2*

locus and all unnecessary DNA fragments were excised in the third-round integration.

Fig. 3. Schematic diagram of third-round integration to construct FY3-2*rml*/FY3-*rml*-rcr*ml*. The ApaI-linearized Cre-Y3 was used to transform FY2-*rml* 3#, generating the strain FY2-*rml*/Cre-Y3. There

were three kinds of different recombination reactions for *lox* sites in FY2-*rml*/Cre-Y3 genome. Herein,

the recombination reactions excised unnecessary DNA fragments (a *leu2*, Amp-ori, Ura3, and pPOX2-

cre-lip2t) and eventually generated two kinds of stable strains FY3-2*rml* and FY3-*rml*-rc*rml*. The

remaining *lox*71 site could be used for subsequent integration process. Primers used for PCR

amplification are labeled in this figure and described in Supplemental Table S2.

Similarly, Cre-Y2 was introduced into FY3-2*rml* to generate FY4-3*rml*/FY4-2*rml*-

rc*rml*, and Cre-Y3 was used to transform FY4-2*rml* for constructing FY5-4*rml*/FY5-

3*rml*-rc*rml* (Supplemental Fig. S9A). Subsequently, Cre-Y2 and Cre-Y3 were iteratively

integrated into the corresponding recombinant strains to obtain FY6-5*rml*/FY6-4*rml*-

rc*rml* and FY7-6*rml*/FY7-5*rml*-rc*rml*. The genome PCR identification results

(Supplemental Fig. S9B) proved that repeated, targeted, and markerless gene integration

was successfully implemented by the novel genetic tool.

3.6 Increasing induction time in sixth and seventh rounds of integration

As shown in Table 1, the positive colony proportions slightly decreased in multi-round

integration processes, which might be owing to the following possibility: an additional

*lox*72 site could have been inserted into the genome in each round of gene integration,

except in the first-round integration, and all of the *lox* sites can bind to Cre recombinase

(Albert et al., 1995; Leibig et al., 2008; Yan et al., 2008). Thus, *lox*72 site might disturb

the combination between LE/RE *lox* sites and Cre recombinase, resulting in increasingly

less recombination reactions during the induction process and a decrease in efficiency in

multi-round integrations. Nevertheless, an increase in induction time caused higher

accumulation of Cre protein in the cell, and more Cre recombinase could recognize

LE/RE mutants and mediate effective recombination.

Accordingly, the transformants were induced for longer time to obtain higher

recombination efficiency in the sixth-round integration. As a result, the proportion of

positive colonies was enhanced, and 16 h was adequate to complete recombination in the

engineered strains (Table 1 and Supplemental Fig. S6C). In the seventh-round integration,

16 h was employed as the induction time and all the chosen colonies could not grow on

MD medium (Supplemental Fig. S6C). Therefore, recombination efficiency might be

roughly inversely proportional to the integration round; however, this problem could be

alleviated by increasing the induction time in MO-Ura liquid medium. Finally, seven

rounds of gene integration were implemented in this study, and the positive colony

proportions reached 90.63%–100% with an induction time of 12–16 h (Table 1). In fact,

the realized integration rounds were adequate to meet the demands of engineering strains

construction, and the potential of the developed genetic tool could also be explored in

future research.

3.7 Protein expression level and growth characteristics evaluation

The recombinants harboring different *rml* gene copies were cultured in BMSY liquid

medium for protein production, and the results are presented in Fig. 5A. The recombinant

strains FY5-4*rml* 31# showed the highest RML expression level, while the lipase activity

of FY6-5*rml*/FY6-4*rml*-rc*rml* and FY7-6*rml*/FY7-5*rml*-rc*rml* was slightly lower. Thus,

gene integration was terminated in the seventh round. Moreover, several colonies were

chosen as representatives for each type of the recombinant strains, and their *rml* gene

copies were determined via qPCR. As shown in Table 1, all the transformants carried the

same gene copy number as previously predicted, indicating that no *rml* expression

cassette was excised from the genome and also confirming the excellent controllability

and stability of the genetic tool.

To evaluate the protein expression level of the recombinant strains generated via the

developed genetic tool, the plasmids hp12d-5*rml* and hp12d-6*rml* were constructed

according to the previous method (Zhou et al., 2019). Subsequently, the plasmid hp12d-

n*rml* (n=1–6) were linearized and introduced into Po1f to form the recombinant strains

Po1f/hp12d-n*rml* (carrying n copies of *rml* expression cassette, n=1–6). Then,

Po1f/hp12d-n*rml* (n=1–6) and engineered strains from seven rounds of gene integration

were respectively inoculated into BMSY medium under the same culture conditions, and

their lipase activity was measured (Fig. 5B), which revealed that the two types of

recombinant strains showed similar protein expression level. Therefore, it can be

concluded that genome editing via the developed genetic tool did not affect target protein

production of the engineering strains. When compared with the previous method (Zhou et

al., 2019), the developed genetic tool has many advantages such as utilization of reusable

marker, only requiring three plasmids, no gene redundancy, and flexible direction of gene

expression cassette. Moreover, the novel genetic tool could assemble engineering strains

harboring any copies of the target gene, only requiring increase/decrease in integration

round.

Furthermore, the strains Po1f, FY1, FY5-4*rml* 31#, and FY7-6*rml* 13# were

cultivated in YPD and BMSY media, respectively, and their growth characteristics were

tested by measuring the optical density at 600 nm (OD_{600}) at different incubation times

(Figs. 5C, D). The results revealed no observable differences in the growth curves of the

four types of strains, indicating that the genome modification in *leu2* locus via Cre/*lox*-

based genetic tool did not influence cell growth.

Fig. 4. (A) Lipase activity of recombinant strains harboring different copies of *rml* expression

cassettes. (B) Comparison of lipase activity of recombinant strains obtained via different methods.

Data are the mean values of each kind of recombinant strain. The strains Po1f and FY1 showed low

lipase activity owing to expression of autologous lipase genes. Weakly significant difference was

observed between the strains FY7-6*rml*/FY7-5*rml*-rc*rml* and Po1f/hp12d-6*rml* ($P=0.044$). (C) Lipase

activity of transformants from third-round integration. (D) Transcriptional level analysis of *rml*

mRNA from FY3-2*rml* and FY3-*rml-rcrml*. The endogenous *act1* was adopted as reference gene, and

the *rml* transcriptional level of FY3-2*rml* 4# was defined as 1. (E) Growth curves of Po1f, FY1, FY5-

4*rml* 31#, and FY7-6*rml* 13# in YPD medium. (F) Growth curves of Po1f, FY1, FY5-4*rml* 31#, and

FY7-6*rml* 13# in BMSY medium. NS, [*], [**], and [***] indicate non-significant, $P < 0.05$, $P < 0.01$, and P

< 0.001, respectively.

3.8 Comparison of two-genotype recombinant strains

Two genotypes of the recombinant strains were generated in each round of integration

(except for the first round), and their numbers were presented in Table 1. The rates of two

forms of strains were approximately equal, suggesting that there was nearly equal

possibility to choose a strain with certain genotype. The colonies with high lipase

activities were selected for protein production evaluation. The results shown in Table 1

also implied that the protein expression levels of the recombinant strains in each round of

integration were similar, which was confirmed in Fig. 5A. In contrast, some previous

studies have reported that the relative arrangement of multiple genes might affect

corresponding transcription and protein production in host cells (Eszterhas et al., 2002;

Lee et al., 2014). Thus, a total of 30 positive colonies from the third-round integration

were identified via the primer pairs KNC-F2/Apa-R6 and SXB-R2/Apa-R3 (data not

shown), and their lipase activities were measured. The results presented in Fig. 5E

showed that the RML activity of each recombinant randomly changed within a proper

range. Moreover, transcriptional level analysis further indicated that there was no

significant difference in the transcriptional quantity of *rml* mRNA isolated from FY3-

2*rml* and FY3-*rml*-rc*rml* cells (Fig. 5F). Thus, the relative direction of *rml* expression

cassettes had no observable effect on protein production in this study, which might be

owing to the difference in the host type. Accordingly, both the genotypes of the strains

were concluded to be suitable for RML production in *Y. lipolytica*.

3.9 Successfully expanding to six different genes and *axp1-2* locus

To evaluate the universality of the novel genetic tool, six genes with varied lengths and

resources were additionally integrated into the engineered strain FY5-4*rml*. The genes

ire1, *kar2*, *pdi*, *sls1,* and *hac1* are the homologous genes of *Y. lipolytica*, and *vgb* is the

coding gene of *Vitreoscilla* hemoglobin. These genes may differentially affect RML

expression level, which determines their integration times. The corresponding plasmids

were derived from Cre-Y2 and Cre-Y3 by replacing the *rml* gene with different target

genes (Supplemental Fig. S4A), and then used to transform FY5-4*rml*. The subsequent

integration processes were similar to those of Cre-Y2 and Cre-Y3, and the proportions of

positive colonies were all above 90% (Table 2), confirming the availability of the

developed genetic tool for repeated, targeted, and markerless integration of different

genes.

Furthermore, a derived tool was established to verify whether the developed genetic

tool could be applied for other target genes, integration loci, and strains. For this purpose,

three plasmids containing *Rhizopus oryzae* lipase (*rol*) gene were respectively

constructed based on Cre-Y1, Cre-Y2, and Cre-Y3 (Supplemental Fig. S4B), and

integrated into defective *axp* gene locus (*axp1-2*) of *Y. lipolytica* Po1h. The

transformation, screening and induction, isolation of positive colonies, and identification

processes were similar to those employed for *rml* gene integration into *leu2* locus of Po1f.

First, the Cre-axp1 plasmid was linearized by *Nsi*I and inserted into the Upaxp locus to

generate an initial strain. Then, the Cre-axp2 and Cre-axp3 plasmids were digested with

*Cla*I and iteratively introduced into the *axp1-2* locus for repeated, targeted, and

markerless integration of *rol* in Po1h. As a result, three rounds of integration were

conducted and all of the integration rates were above 96%. These findings revealed that

such extension of the novel genetic tool is feasible and convenient.

Table 2. Integration rates of various genes, loci and strains in this study.

Gene	GenBank	Lenth (bp)	Integration round	Positive colonies/ all colonies	Proportion (%)
Integration locus: *leu2* gene of FY5-4*rml*, induction time: 16 h					
ire1	XM_002142952.1	3663	1	32/32	100
kar2	U63136.1	2013	1	32/32	100
			2	32/32	100
			3	32/32	100
pdi	XM_503481.1	1515	1	32/32	100
			2	32/32	100
			3	31/32	96.88
			4	31/32	96.88
sls1	Z50154.1	1281	1	32/32	100
			2	32/32	100
hac1	XM_500811.1	918	1	32/32	100
vgb	AF292694.1	441	1	32/32	100
			2	32/32	100
Integration locus: *axp1-2* of Po1h, induction time: 12 h					
rol	AF229435.1	1101	1	31/32	96.88
			2	32/32	100
			3	32/32	100

4. Discussion

In the present study, a Cre/*lox*-based genetic tool was precisely and skillfully designed. The plasmids Cre-Y2 and Cre-Y3 adopted two inversely oriented *lox* sites, and the Ura3 marker cannot be removed from the genome if the strain does not undergo integration process as expected. Thus, the Ura$^+$ strain could be distinguished by isolating positive colonies on the MD medium. It must be noted that except for the *cre* gene and *lox* sites, all of the genetic tool elements are nonspecific and replaceable. Any DNA sequence can be chosen as the homologous fragment as long as no gene is disrupted, and the length of the homologous fragment is very flexible because of the excision of unnecessary sequences. The *rml* gene in plasmids Cre-Y2 and Cre-Y3 can be replaced by heterologous and endogenous genes of *Y. lipolytica* to accomplish repeated markerless integration, and the plasmid construction process is simple and convenient. In addition, the Cre/*lox*-based genetic tool can also be applied to other genes and loci if the corresponding plasmid elements are available.

Recently, several genetic tools have been developed for markerless gene integration of *Y. lipolytica* based on CRISPR-Cas9 system, Cre/*lox* system and Ura/5-fluoroorotic acid resistance method (Supplemental Fig. S10) (Gao et al., 2018; Holkenbrink et al., 2018; Lv et al., 2019; Schwartz et al., 2017b). As shown in Supplemental Table S3, the integration efficiency markedly fluctuates with different genome loci and single-guide RNAs (sgRNAs) for CRISPR-Cas9-mediated gene integrations (Holkenbrink et al., 2018; Schwartz et al., 2017b). According to most of the existing Cre/*lox*-based methods (Holkenbrink et al., 2018; Lv et al., 2019), the DNA fragment can be integrated into the host genome via gene insertion or gene replacement. However, insertion event may

introduce unnecessary DNA segments into the genome, whereas replacement events,

which occur occasionally, can avoid this issue, but requires substantial screening of target

transformants. Furthermore, while the Ura/5-fluoroorotic acid resistance method (Ura

pop-in/pop-out method) can also accomplish markerless gene integration, reverse

mutation is difficult to avoid in the marker recovery process. In addition, new plasmids

carrying different sgRNA or homologous fragments must be constructed to repeatedly

realize gene integration in a specific genome locus via the three above-mentioned genetic

tools. These limitations can be completely overcome by the novel genetic tool designed

in the present study. The target gene is integrated into the host genome via efficient,

targeted, and repeatable gene insertion event, and the introduced non-target sequences

can be deleted by recombination reaction of the *lox* sites. Moreover, despite the

occurrence of inefficient insertion event in the integration process, a few recombinants

can be enriched via twice auxotroph screening. The existing methods based on CRISPR-

Cas9 and Cre/*lox* systems require two different selection markers and additional plasmid

recovery procedure (Gao et al., 2018; Holkenbrink et al., 2018; Lv et al., 2019; Schwartz

et al., 2017b). In contrast, only a single marker is required for the novel Cre/*lox*-based

genetic tool, and the process of gene integration, markers reuse, and plasmid recovery can

be combined and accomplished simultaneously, which can significantly reduce the

experimental period and workload. Overall, the novel genetic tool developed in this study

can overcome many limitations such as low HR rate, limited selection marker, random

integration site, and additional plasmid recovery.

In conclusion, a powerful and efficient Cre/*lox*-based genetic tool for repeated,

targeted, and markerless gene integration was developed in this study. In practice, only a

single selection marker can enable iterative integration of target gene into the genome,

which can facilitate genetic modifications of the host strains. Furthermore, in each round

of integration, all of the unnecessary segments are removed, the relative direction of each

target gene expression cassette is controllable, and the gene recombination on genome is

foreseeable. These characteristics can significantly decrease gene redundancy in the

engineering strain, making it convenient for the use of host strain in pharmaceutical, food,

and other health-related areas. Besides, the developed genetic tool is also suitable for

various other genes and loci. The findings of this study significantly broaden the

application of Cre/*lox* system and strengthen the capabilities for genome modification in

Y. lipolytica.

Conflict of interest

The authors declare that they have no conflict of interest.

Acknowledgments

We acknowledge financial support by the National Natural Science Foundation of China

(No: 41972319), the National High Technology Research and Development Program of

China (Nos: 2013AA065805 and 2014AA093510), and the Fundamental Research Funds

for HUST (Nos. 2017KFXKJC010 and 2017KFTSZZ001). We appreciate the Research

Core Facilities for Life Science (Huazhong University of Science and Technology) for

their valuable assistance in qPCR measurements. We thank International Science Editing

(http://www.internationalscienceediting.com) for editing this manuscript.

Author contributions

Q.Z. and L.J. conducted the project design. Q.Z., L.J., W.L. and Y.L. carried out the

experiments. Q.Z., L.J., Z.H. and H.Z. helped with the formal analysis. Q.Z., L.J., L.X.

and Y.Y. conducted the manuscript preparation.

References

Abdel-Mawgoud, A. M., et al., 2018. Metabolic engineering in the host *Yarrowia*

lipolytica. Metab. Eng. 50, 192-208. https://doi.org/10.1016/j.ymben.2018.07.016

Albert, H., Dale, E. C., Lee, E., Ow, D. W., 1995. Site-specific integration of DNA into

wild-type and mutant *lox* sites placed in the plant genome. Plant. J. 7, 649-59.

https://doi.org/10.1046/j.1365-313x.1995.7040649.x

Bae, S. J., Park, B. G., Kim, B. G., Hahn, J. S., 2020. Multiplex gene disruption by

targeted base editing of *Yarrowia lipolytica* genome using cytidine deaminase

combined with the CRISPR/Cas9 system. Biotechnol. J. 15, e1900238.

https://doi.org/10.1002/biot.201900238

Barth, G., Gaillardin, C., 1996. *Yarrowia lipolytica*., Nonconventional Yeasts in

Biotechnology: A Handbook. Springer Berlin Heidelberg, Berlin, Heidelberg, pp.

313-388.

Blazeck, J., et al., 2014. Harnessing *Yarrowia lipolytica* lipogenesis to create a platform

for lipid and biofuel production. Nat. Commun. 5, 3131.

https://doi.org/10.1038/ncomms4131

Carly, F., et al., 2017. Enhancing erythritol productivity in *Yarrowia lipolytica* using

metabolic engineering. Metab. Eng. 42, 19-24.

https://doi.org/10.1016/j.ymben.2017.05.002

Chen, D. C., Beckerich, J. M., Gaillardin, C., 1997. One-step transformation of the

dimorphic yeast *Yarrowia lipolytica*. Appl. Microbiol. Biotechnol. 48, 232-5.

https://doi.org/10.1007/s002530051043

Cong, L., et al., 2013. Multiplex genome engineering using CRISPR/Cas systems.

605   Science. 339, 819-23. https://doi.org/10.1126/science.1231143

606 Czajka, J. J., et al., 2018. Engineering the oleaginous yeast *Yarrowia lipolytica* to

607   produce the aroma compound beta-ionone. Microb. Cell. Fact. 17, 136.

608   https://doi.org/10.1186/s12934-018-0984-x

609 Eszterhas, S. K., Bouhassira, E. E., Martin, D. I., Fiering, S., 2002. Transcriptional

610   interference by independently regulated genes occurs in any relative arrangement of

611   the genes and is influenced by chromosomal integration position. Mol. Cell. Biol. 22,

612   469-79. https://doi.org/10.1128/mcb.22.2.469-479.2002

613 Fickers, P., et al., 2003. New disruption cassettes for rapid gene disruption and marker

614   rescue in the yeast *Yarrowia lipolytica*. J. Microbiol. Methods. 55, 727-37.

615   https://doi.org/10.1016/j.mimet.2003.07.003

616 Gaj, T., Gersbach, C. A., Barbas, C. R., 2013. ZFN, TALEN, and CRISPR/Cas-based

617   methods for genome engineering. Trends. Biotechnol. 31, 397-405.

618   https://doi.org/10.1016/j.tibtech.2013.04.004

619 Gao, D., et al., 2018. Dual CRISPR-Cas9 cleavage mediated gene excision and targeted

620   integration in *Yarrowia lipolytica*. Biotechnol. J. 13, e1700590.

621   https://doi.org/10.1002/biot.201700590

622 Gueldener, U., et al., 2002. A second set of *lox*P marker cassettes for Cre-mediated

623   multiple gene knockouts in budding yeast. Nucleic. Acids. Res. 30, e23.

624   https://doi.org/10.1093/nar/30.6.e23

625 Hentges, P., et al., 2005. Three novel antibiotic marker cassettes for gene disruption and

626   marker switching in *Schizosaccharomyces pombe*. Yeast. 22, 1013-9.

627   https://doi.org/10.1002/yea.1291

Hochrein, L., et al., 2018. L-SCRaMbLE as a tool for light-controlled Cre-mediated

recombination in yeast. Nat. Commun. 9, 1931. https://doi.org/10.1038/s41467-017-

02208-6

Holkenbrink, C., et al., 2018. EasyCloneYALI: CRISPR/Cas9-Based synthetic toolbox

for engineering of the yeast *Yarrowia lipolytica*. Biotechnol. J. 13, e1700543.

https://doi.org/10.1002/biot.201700543

Hsu, P. D., Lander, E. S., Zhang, F., 2014. Development and applications of CRISPR-

Cas9   for   genome   engineering.   Cell.   157,   1262-78.

https://doi.org/10.1016/j.cell.2014.05.010

Kawano, F., Okazaki, R., Yazawa, M., Sato, M., 2016. A photoactivatable Cre-*lox*P

recombination system for optogenetic genome engineering. Nat. Chem. Biol. 12,

1059-1064. https://doi.org/10.1038/nchembio.2205

Kuhn, R., Torres, R. M., 2002. Cre/*lox*P recombination system and gene targeting.

Methods. Mol. Biol. 180, 175-204. https://doi.org/10.1385/1-59259-178-7:175

Larroude, M., et al., 2018. A synthetic biology approach to transform *Yarrowia lipolytica*

into a competitive biotechnological producer of beta-carotene. Biotechnol. Bioeng.

115, 464-472. https://doi.org/10.1002/bit.26473

Larroude, M., Rossignol, T., Nicaud, J. M., Ledesma-Amaro, R., 2018. Synthetic biology

tools for engineering *Yarrowia lipolytica*. Biotechnol. Adv. 36, 2150-2164.

https://doi.org/10.1016/j.biotechadv.2018.10.004

Ledesma-Amaro, R., Nicaud, J. M., 2016. *Yarrowia lipolytica* as a biotechnological

chassis to produce usual and unusual fatty acids. Prog. Lipid. Res. 61, 40-50.

https://doi.org/10.1016/j.plipres.2015.12.001

Lee, T. J., Parikh, R. Y., Weitz, J. S., Kim, H. D., 2014. Suppression of expression

between adjacent genes within heterologous modules in yeast. G3 (Bethesda). 4,

109-16. https://doi.org/10.1534/g3.113.007922

Leibig, M., et al., 2008. Marker removal in staphylococci via Cre recombinase and

different *lox* sites. Appl. Environ. Microbiol. 74, 1316-23.

https://doi.org/10.1128/AEM.02424-07

Lv, Y., Edwards, H., Zhou, J., Xu, P., 2019. Combining 26s rDNA and the Cre-*lox*P

system for iterative gene integration and efficient marker curation in *Yarrowia*

lipolytica. ACS. Synth. Biol. 8, 568-576. https://doi.org/10.1021/acssynbio.8b00535

Madzak, C., 2015. *Yarrowia lipolytica*: recent achievements in heterologous protein

expression and pathway engineering. Appl. Microbiol. Biotechnol. 99, 4559-77.

https://doi.org/10.1007/s00253-015-6624-z

Madzak, C., 2018. Engineering *Yarrowia lipolytica* for Use in Biotechnological

Applications: A review of major achievements and recent innovations. Mol.

Biotechnol. 60, 621-635. https://doi.org/10.1007/s12033-018-0093-4

Madzak, C., Gaillardin, C., Beckerich, J. M., 2004. Heterologous protein expression and

secretion in the non-conventional yeast *Yarrowia lipolytica*: a review. J. Biotechnol.

109, 63-81. https://doi.org/10.1016/j.jbiotec.2003.10.027

Madzak, C., Treton, B., Blanchin-Roland, S., 2000. Strong hybrid promoters and

integrative expression/secretion vectors for quasi-constitutive expression of

heterologous proteins in the yeast *Yarrowia lipolytica*. J. Mol. Microbiol. Biotechnol.

2, 207-16. PMID: 10939246

Markham, K. A., et al., 2018. Rewiring *Yarrowia lipolytica* toward triacetic acid lactone

for materials generation. Proc. Natl. Acad. Sci. USA. 115, 2096-2101.

https://doi.org/10.1073/pnas.1721203115

Pignede, G., et al., 2000. Autocloning and amplification of LIP2 in *Yarrowia lipolytica*.

Appl. Environ. Microbiol. 66, 3283-9. https://doi.org/10.1128/aem.66.8.3283-

3289.2000

Richardson, S. M., et al., 2017. Design of a synthetic yeast genome. Science. 355, 1040-

1044. https://doi.org/10.1126/science.aaf4557

Schwartz, C., et al., 2017a. CRISPRi repression of nonhomologous end-joining for

enhanced genome engineering via homologous recombination in *Yarrowia lipolytica*.

Biotechnol. Bioeng. 114, 2896-2906. https://doi.org/10.1002/bit.26404

Schwartz, C., et al., 2017b. Standardized markerless gene integration for pathway

engineering in *Yarrowia lipolytica*. ACS. Synth. Biol. 6, 402-409.

https://doi.org/10.1021/acssynbio.6b00285

Suzuki, N., et al., 2005. New multiple-deletion method for the *Corynebacterium*

glutamicum genome, using a mutant *lox* sequence. Appl. Environ. Microbiol. 71,

8472-80. https://doi.org/10.1128/AEM.71.12.8472-8480.2005

Wu, X., et al., 2014. Genome-wide binding of the CRISPR endonuclease Cas9 in

mammalian cells. Nat. Biotechnol. 32, 670-6. https://doi.org/10.1038/nbt.2889

Wu, Y., et al., 2018. In vitro DNA SCRaMbLE. Nat. Commun. 9, 1935.

https://doi.org/10.1038/s41467-018-03743-6

Yan, X., Yu, H. J., Hong, Q., Li, S. P., 2008. Cre/*lox* system and PCR-based genome

engineering in *Bacillus subtilis*. Appl. Environ. Microbiol. 74, 5556-62.

https://doi.org/10.1128/AEM.01156-08

Yang, K., et al., 2019. Subcellular engineering of lipase dependent pathways directed

towards lipid related organelles for highly effectively compartmentalized

biosynthesis of triacylglycerol derived products in *Yarrowia lipolytica*. Metab. Eng.

55, 231-238. https://doi.org/10.1016/j.ymben.2019.08.001

Zhou, Q., et al., 2019. Overexpression of GRAS *Rhizomucor miehei* lipase in *Yarrowia*

lipolytica via optimizing promoter, gene dosage and fermentation parameters. J.

Biotechnol. 306, 16-23. https://doi.org/10.1016/j.jbiotec.2019.09.004