Rif2 protects Rap1-depleted telomeres from MRX-mediated degradation in *Saccharomyces cerevisiae*

Fernando R. Rosas Bringas[1], Sonia Stinus[1], Pien de Zoeten[1], Marita Cohn[2], and Michael Chang[1,*]

[1]European Research Institute for the Biology of Ageing, University of Groningen, University Medical Center Groningen, Groningen, The Netherlands

[2]Department of Biology, Lund University, Lund, Sweden

*For correspondence: m.chang@umcg.nl

Abstract

Rap1 is the main protein that binds double-stranded telomeric DNA in *Saccharomyces cerevisiae*. Examination of the telomere functions of Rap1 is complicated by the fact that it also acts as a transcriptional regulator of hundreds of genes and is encoded by an essential gene. In this study, we disrupt Rap1 telomere association by expressing a mutant telomerase RNA subunit (tlc1-tm) that introduces mutant telomeric repeats. *tlc1-tm* cells grow similar to wild-type cells, although depletion of Rap1 at telomeres causes defects in telomere length regulation and telomere capping. Rif2 is a protein normally recruited to telomeres by Rap1, but we show that Rif2 can still associate with Rap1-depleted *tlc1-tm* telomeres, and that this association is required to inhibit telomere degradation by the MRX complex. We find that Rap1, Rif2, and the Ku complex work in parallel to prevent telomere degradation, and absence of all three at telomeres causes lethality. The partially redundant mechanisms may explain the rapid evolution of telomere components in budding yeast species.

Keywords: Rap1, Rif2, Ku complex, telomere, telomerase, tlc1-tm, yeast

Introduction

Telomeres, nucleoprotein complexes located at the ends of eukaryotic chromosomes, protect chromosome ends from degradation, from telomere-telomere fusion events, and from being recognized as double-stranded DNA breaks (Wellinger & Zakian, 2012; de Lange, 2018). In most eukaryotic species, telomeric DNA consists of tandem G/C-rich repeats of double-stranded DNA with the G-rich strand extending to form a 3′ single-stranded overhang. These repeats are bound by specialized proteins—some to the double-stranded region and others to the 3′ overhang—which are important for proper telomere function. Telomere length is maintained by a dynamic process of shortening and lengthening. Telomeres shorten due to incomplete DNA replication and nucleolytic degradation, and are lengthened by the action of a specialized reverse transcriptase called telomerase (Wellinger, 2014). At its core, telomerase consists of a catalytic protein subunit and an RNA subunit, and extends telomeres by iterative reverse transcription of a short G-rich sequence to the 3′ overhang, using the RNA subunit as a template.

Rap1 is the main double-stranded telomeric DNA binding protein in the budding yeast *Saccharomyces cerevisiae* (Buchman et al., 1988; Conrad et al., 1990), with important roles in regulating telomere length (Lustig et al., 1990; Conrad et al., 1990; Marcand et al., 1997), transcriptional silencing of subtelomeric genes (Kyrion et al., 1993), and preventing telomere-telomere fusions (Pardo & Marcand, 2005). Rap1 mediates these functions in part by recruiting additional proteins (i.e. Rif1, Rif2, Sir3, and Sir4) via its C-terminal domain (Hardy et al., 1992; Wotton & Shore, 1997; Moretti et al., 1994). Expression of a mutant Rap1 lacking the C-terminal domain, which retains the ability to bind telomeric sequences, mimics the deletion of the *RIF* and *SIR* genes in terms of telomere length regulation and subtelomeric gene silencing (Wotton & Shore, 1997; Kyrion et al., 1993). However, the telomeric functions of Rap1 are not limited to the recruitment of Rif and Sir proteins. For

example, although Rif2 and Sir4 are important for the inhibition of telomere end-to-end

fusions, Rap1 also inhibits fusions independently of Rif2 and Sir4 (Marcand et al., 2008). In

addition, binding of Rap1 to telomeric repeats is thought to promote replication fork pausing

and breakage of dicentric chromosomes at telomere fusions independently of the Rif and Sir

proteins (Makovets et al., 2004; Guérin et al., 2019; Douglas & Diffley, 2021). Despite the

central role of Rap1 in yeast telomere biology, its study is complicated by the fact that Rap1

is also a transcription factor that regulates the expression of a few hundred genes (Lieb et al.,

2001). Moreover, *RAP1* is essential for viability (Shore & Nasmyth, 1987), preventing

analysis of a gene deletion mutant.

In this study, we examined the function of Rap1 at *S. cerevisiae* telomeres by

expressing a mutant telomerase RNA subunit (tlc1-tm) that introduces $[(TG)_{0-}$

$_4TGG]_nATTTGG$ mutant telomeric repeats instead of wild-type $(TG)_{0-6}TGGGTGTG(G)_{0-1}$

repeats (Chang et al., 2007; Förstemann & Lingner, 2001). We find that Rap1 binds very

poorly to mutant telomeric sequences, yet *tlc1-tm* cells are viable and grow similar to wild-

type cells. The depletion of Rap1, not surprisingly, causes telomere length homeostasis

defects. Unexpectedly, the overall levels of Rif2 at telomeres are unaffected. Rif2 recruitment

to *tlc1-tm* telomeres is dependent on the MRX complex, and Rif2 is crucial for preventing

tlc1-tm telomeres from degradation by the MRX complex. The yeast Ku complex functions

in parallel to protect *tlc1-tm* telomeres, and absence of both Rif2 and the Ku complex renders

tlc1-tm cells inviable. Our findings reveal multiple redundant mechanisms that may have

been important for the rapid evolution of telomere components in budding yeasts.

Results

Rap1 binds poorly to *tlc1-tm* sequences

To examine the function of Rap1 specifically at telomeres, we made use of a telomere RNA

mutant, *tlc1-tm*. Rap1 has a consensus DNA binding sequence of 5′-CACCCAYACMYM-3′

(where Y is C or T, and M is A or C) containing an invariable CCC core (Graham &

Chambers, 1994; Lieb et al., 2001). The template region of wild-type TLC1 is 3′-

CACACACCCACACCAC-5′, resulting in the addition of $(TG)_{0-6}TGGGTGTG(G)_{0-1}$

telomeric repeats (Förstemann & Lingner, 2001). The tlc1-tm template region is 3′-

CACCUAAACCACACAC-5′, resulting in $[(TG)_{0-4}TGG]_nATTTGG$ mutant telomeric

repeats (Chang et al., 2007). The lack of the GGG motif in the mutant repeat sequence should

disrupt Rap1 binding, yet the *tlc1-tm* mutant grows similar to a wild-type strain (Fig. 1A).

Telomeres in the *tlc1-tm* mutant are on average longer and more heterogeneous in length than

in wild-type strains (Fig. 1B), but the telomere profile of *tlc1-tm* is much less dramatically

altered compared to most other *TLC1* mutants with altered template sequences (Förstemann

et al., 2003; Lin et al., 2004). Telomeres in *tlc1-tm* cells were previously reported to be

slightly shorter than in wild-type cells, as measured by telomere PCR (Chang et al., 2007).

This discrepancy is likely due to the tendency of PCR to amplify shorter fragments more

efficiently that longer ones.

Because telomerase only adds repeats to the distal end of telomeres, the proximal

region of *tlc1-tm* telomeres still contains some wild-type repeats with the capacity to be

bound by Rap1. Thus, we engineered strains in which the left arm of chromosome VII (VII-

L) ends with a telomere that consists entirely of either wild-type or *tlc1-tm* telomeric

sequence (Fig. 1C). These strains were constructed by replacing the native VII-L telomere

with the *URA3* gene followed by 81 bp of wild-type or 84 bp of *tlc1-tm* telomeric seed

sequence, which was lengthened in vivo by either wild-type or *tlc1-tm* mutant telomerase,

respectively. As expected, the VII-L telomere containing only mutant sequence (VII-L-MUT)

was longer and more heterogeneous than the wild-type control telomere (VII-L-WT) (Fig.

Figure 1. Rap1 binds poorly to *tlc1-tm* telomere sequences

(A) Exponentially growing cells of the indicated genotypes were diluted to an OD_{600} of 0.05 and monitored for 8 h. Three clones of each genotype were followed, and the average of the measurements at each time point is plotted. Error bars are too small to be visualized.

(B) Telomere Southern blot analysis of *TLC1* and *tlc1-tm* cells (three clonal isolates per genotype). Black arrowhead indicates a 1.8 kb DNA fragment generated from the BsmAI-digestion of plasmid pYt103 (Shampay et al., 1984). The major terminal restriction fragment is below the 1.8 kb control band.

(C) Southern blot analysis of the artificial VII-L telomere, with either wild-type or *tlc1-tm* mutant sequence, using a probe to the adjacent *URA3* gene. Multiple clones were examined, with each clone propagated for 1 to 4 passages (each passage corresponds to approximately 25 generations). A wild-type strain (lacking the artificial VII-L telomere) was used as a control.

(D, F, G) ChIP-qPCR analysis of the association of (D) Protein A-tagged Rap1, (F) Myc-tagged Cdc13, and (G) Myc-tagged Tbf1 to six Y′ telomeres, the VII-L telomere, or to the non-telomeric *ARO1* locus. Untagged wild-type and *tlc1-tm* strains were used as controls. The mean percentage of input +/- SEM is shown (n=3, ** p<0.01).

(E) EMSA of Rap1 protein incubated with radiolabeled oligonucleotides with either wild-type or tlc1-tm mutant telomeric sequence.

1C). We assessed the association of Rap1 to the VII-L-WT or VII-L-MUT telomeres, as well

as to native Y′ telomeres, by chromatin immunoprecipitation coupled with quantitative PCR

(ChIP-qPCR). Binding of Rap1 is substantially reduced at the VII-L-MUT telomere (Fig.

1D). Rap1 is more modestly reduced at Y′ telomeres due to the retention of some wild-type

repeats at native telomeres. We confirmed the Rap1 ChIP-qPCR result by performing an

electrophoretic mobility shift assay (EMSA) using recombinant full-length Rap1 protein. We

find that Rap1 shows a very poor binding capacity to the *tlc1-tm* oligonucleotides compared

to the wild-type telomeric sequence (Fig. 1E).

Cdc13 is the main sequence-specific single-stranded telomeric DNA binding protein

in *S. cerevisiae* (Nugent et al., 1996). Cdc13 requires a minimum of 11 nucleotides of GT-

rich sequence for full binding affinity, with only three bases—forming a GNGT motif—

recognized with high specificity (Eldridge et al., 2006). The GNGT motif is present in the

tlc1-tm mutant sequence, so Cdc13 binding should not be altered. Consistent with this

hypothesis, we find that Cdc13 associates equally well with both wild-type and mutant

telomeres (Fig. 1F).

The ability of *tlc1-tm* strains to survive with greatly reduced Rap1 telomere

association was surprising given that Rap1 is the main telomere binding protein in *S.*

cerevisiae, important for recruiting a number of other proteins to telomeres, and is encoded

by an essential gene. However, the essential function of Rap1 is likely linked to its ability to

bind nontelomeric sites in the genome where it acts as a transcriptional regulator of several

hundred genes (Azad & Tomar, 2016). In addition, *S. cerevisiae* strains expressing a mutant

telomerase that adds vertebrate TTAGGG telomeric repeats are devoid of Rap1, indicating

that Rap1 is not essential for telomere capping (Alexander & Zakian, 2003; Brevet et al.,

2003). In these strains, Tbf1 binds to the vertebrate repeats and is able to regulate telomere

length homeostasis in a Rap1-independent manner (Alexander & Zakian, 2003; Berthiau et

al., 2006). Thus, we assessed binding of Tbf1 to *tlc1-tm* telomeres. We find no change in the

levels of Tbf1 at *tlc1-tm* telomeres compared to wild-type telomeres by ChIP-qPCR (Fig.

1G), consistent with neither wild-type nor mutant telomere sequences containing the RCCCT

Tbf1 consensus binding sequence (Preti et al., 2010), indicating that the loss of Rap1 is not

compensated by recruitment of Tbf1. The Tbf1 association observed at the native Y′

telomeres is due to the presence of TTAGGG repeats at subtelomeric regions (Brigati et al.,

1993), while the association observed at the artificial VII-L telomere is likely due to the

presence of several RCCCT motifs in the adjacent sequence (Supplementary Fig. 1).

Loss of Rap1-mediated telomere length regulation at *tlc1-tm* telomeres

The decreased binding of Rap1 to *tlc1-tm* sequences likely explains the long, heterogeneous-

sized telomeres in *tlc1-tm* strains, because Rap1 negatively regulates telomerase through

what is called the "protein counting" model. This model posits that Rap1, through its

recruitment of Rif1 and Rif2, inhibits telomerase; the longer a telomere is, the more Rap1,

Rif1, and Rif2 will be present, and the stronger the inhibition of telomerase will be (Marcand

et al., 1997; Levy & Blackburn, 2004). Reduced binding of Rap1 would cause *tlc1-tm*

telomeric sequences to not be recognized as telomeric in terms of Rap1-mediated telomere

length regulation. To test this hypothesis, we again modified telomere VII-L to generate a

telomere that would be seeded with either 84 bp or 300 bp of *tlc1-tm* telomeric sequence, but

in a *TLC1* strain expressing wild-type telomerase so that the tip of the telomere would contain

wild-type telomeric sequences. In both cases, the size of the telomere VII-L terminal

restriction fragment increased in size, with the magnitude of the increase roughly equivalent

to the size of the *tlc1-tm* telomeric seed sequence, indicating that this sequence is not being

sensed by the Rap1 protein counting mechanism (Fig. 2A).

Figure 2. Telomere length regulation is disrupted in *tlc1-tm* cells

(A) Southern blot analysis of the artificial VII-L telomere using a probe to the adjacent *URA3* gene. The telomere was seeded with either wild-type or *tlc1-tm* mutant sequence of the indicated lengths in a strain expressing wild-type *TLC1*. Multiple clones of each strain were examined. A wild-type strain (lacking the artificial VII-L telomere) was used as a control.

(B) In vivo extension of *tlc1-tm* telomeres was examined using the iSTEX assay. Telomere VI-R was amplified and sequenced after the induction of wild-type telomerase. Each bar represents an individual telomere. The black and red portions of each bar represent wild-type and *tlc1-tm* sequence, respectively, that is identical in sequence and thus present before telomerase induction. The length of the black/wild-type sequence is 48 bp. Sequence that is divergent from the black and red sequence is shown in grey and green. Grey represents newly added wild-type sequence after the induction of telomerase. Green represents divergent *tlc1-tm* sequence, most likely a result of homologous recombination. Telomeres are sorted based on the length of the undiverged (black plus red) sequence.

(C) Telomere VI-R sequences obtained from the iSTEX analysis in B were binned into groups of 10 nt in size according to telomere length before telomerase induction. iSTEX data for the extension of wild-type telomeres were taken from previous studies (Strecker et al., 2017; Stinus et al., 2017) and included for comparison. Groups containing less than four telomeres were excluded from this analysis. Frequency of extension and average telomere length before telomerase induction were calculated and plotted for each group. Logarithmic regression curves for each data set were also included in the plot. Telomeres shorter than 40 nt before telomerase induction, which are not efficiently recognized and extended by telomerase (Strecker et al., 2017), were removed from the regression analysis.

To examine telomere length regulation in *tlc1-tm* strains further, we monitored

telomerase-mediated telomere extension events at nucleotide resolution after a single cell

cycle using the iSTEX (inducible Single Telomere EXtension) assay (Strecker et al., 2017).

At wild-type telomeres, telomerase extends only a subset of telomeres in each cell cycle, with

a strong preference for the extension of short telomeres (Teixeira et al., 2004; Strecker et al.,

2017). At *tlc1-tm* telomeres, we find that telomere extension frequency is dramatically

increased, with nearly all (92%) of telomeres being extended during a single cell cycle (Fig.

2B and 2C). This observation is consistent with the protein counting model; the lack of Rap1

at *tlc1-tm* telomeres causes them to all be recognized as short telomeres in need of

elongation.

***tlc1-tm* telomeres are rapidly degraded in the absence of telomerase**

The increase in telomere extension frequency of *tlc1-tm* telomeres is much more dramatic

than the increase seen in *rif1Δ*, *rif2Δ*, and *pif1-m2* cells (Teixeira et al., 2004; Stinus et al.,

2017), even though these mutants have similar or longer telomeres than the *tlc1-tm* mutant

(e.g. see Fig. 5 for comparison to *rif1Δ* and *rif2Δ*). Ablation of Rap1 leads to readily

detectable telomere degradation within a few hours (Vodenicharov et al., 2010), so we

hypothesized that while decreased Rap1 at *tlc1-tm* telomeres may favor their extension by

telomerase, increased degradation may limit the length of *tlc1-tm* telomeres. If true, removal

of telomerase should trigger rapid entry into senescence. To test this idea, we sporulated

diploid strains that were heterozygous for *EST2*, which encodes the protein catalytic subunit

of telomerase (Lingner et al., 1997), or *TLC1*, with either wild-type or mutant telomeres (i.e.

est2Δ/EST2 versus *est2Δ/EST2 tlc1-tm/tlc1-tm* and *tlc1Δ/TLC1* versus *tlc1Δ/tlc1-tm*), and

performed senescence assays with the haploid meiotic progeny. In the presence of wild-type

telomeres but absence of telomerase (*est2Δ* and *tlc1Δ*), telomeres shortened until the cells

senesced after 60 to 70 population doublings, as expected (Fig. 3A). A small subset of the

senescent population was then able to lengthen the telomeres by recombination-mediated

mechanisms, forming so-called survivors (Lundblad & Blackburn, 1993). We found that cells

containing mutant telomeres senesced rapidly, only ~40 population doublings after

telomerase loss (Fig. 3A, *est2Δ tlc1-tm** and *tlc1Δ**). We examined the telomere length of

cells that had undergone ~30 population doublings after isolation of the haploid spores and

found that, upon loss of telomerase, mutant telomeres became extremely heterogeneous and

degraded (i.e. a continuous smear of telomeric DNA-hybridizing signal extending from the

wells to the bottom; Fig. 3B).

Rif1 and Sir4, but not Rif2 nor Sir3, association to *tlc1-tm* telomeres is decreased

Rap1, through its C-terminal domain, recruits Rif1, Rif2, Sir3, and Sir4 to telomeres (Hardy

et al., 1992; Wotton & Shore, 1997; Moretti et al., 1994). We reasoned that the reduction of

Rap1 at the *tlc1-tm* telomeres should also decrease recruitment of Rif1, Rif2, Sir3, and Sir4,

which we tested by ChIP-qPCR (Fig. 4A). Consistent with this hypothesis, we find that Rif1

and Sir4 recruitment to the VII-L-MUT telomere is reduced compared to the VII-L-WT

telomere (to 16% for Rif1 and 31% for Sir4). Surprisingly, the recruitment of neither Rif2 nor

Sir3 is affected. Rif2 can interact with the Xrs2 and Rad50 subunits of the MRX complex

(Hirano et al., 2009; Hailemariam et al., 2019; Roisné-Hamelin et al., 2021), which binds to

DNA ends and telomeres (Oh & Symington, 2018); Rif2 can also interact directly with

double-strand DNA (Cassani et al., 2016; Hailemariam et al., 2019). Sir3 possesses multiple

domains that can interact with histones (Gartenberg & Smith, 2016). Thus, Rif2 and Sir3 may

associate to *tlc1-tm* telomeres via Rap1-independent mechanisms.

Transcription is not affected at *tlc1-tm* telomeres

Figure 3. Fast degradation of *tlc1-tm* telomeres in the absence of telomerase

(A) Senescence rates were measured by serial passaging of strains of the indicated genotypes, derived from the sporulation of *est2Δ/EST2* and *est2Δ/EST2 tlc1-tm/tlc1-tm* diploids (top panel) or *tlc1Δ/TLC1* and *tlc1Δ/tlc1-tm* diploids (bottom panel). Cell density was measured every 24 h, followed by dilution to 1x10⁵ cell/ml. Mean +/- SEM of four independent isolates per genotype is plotted.

(B) Telomere Southern blot analysis of samples obtained at the first time point of the senescence assays in A. Black arrowhead indicates the 1.8 kb telomere sequence-containing fragment loaded as control, as in Figure 1B.

Figure 4. Recruitment of Rif1 and Sir4, but not Rif2 nor Sir3, is significantly reduced at *tlc1-tm* telomeres

(A) ChIP-qPCR analysis of the association of Myc-tagged Rif1, Rif2, Sir4, and Protein A-tagged Sir3 to six Y' telomeres, the VII-L telomere, or to the non-telomeric *ARO1* locus. Untagged wild-type and tlc1-tm strains were used as controls.

(B) Total RNA was reverse transcribed and TERRA from specific telomeres (I-L and XV-L) was analyzed by qPCR. TERRA values were normalized to ACT1 levels, and to the respective wild type (*TLC1*).

(C) Tenfold serial dilutions of strains with the indicated genotypes were spotted on SC plates without or with 5-FOA.

(D) The expression of the subtelomerically integrated *URA3* gene was measured in the indicated yeast strains by RT-qPCR.

All data (except C) are shown as mean +/- SEM (n=3, *** p<10^{-3}, **** p<10^{-4}).

Rap1, through its recruitment of the Rif and Sir proteins, represses the transcription of non-

coding telomeric repeat-containing RNA (TERRA) (Iglesias et al., 2011). Reduced levels of

Rap1, Rif1, and Sir4 at *tlc1-tm* telomeres may alter TERRA transcription. However, we find

no change in TERRA abundance in *tlc1-tm* cells (Fig. 4B). Rap1, through its recruitment of

the Sir complex, also mediates the transcriptional silencing of subtelomeric genes (Moretti et

al., 1994). The VII-L-WT and VII-L-MUT telomeres possess a *URA3* gene immediately

adjacent to the telomeric repeats. Silencing of the *URA3* gene was monitored by assaying for

growth in the presence of 5-fluoroorotic acid (5-FOA), which kills cells expressing *URA3*

(Fig. 4C). Strains deleted for *SIR2*, which causes a defect in silencing (Aparicio et al., 1991),

were used as controls. Because hypersensitivity to 5-FOA does not correlate with loss of

silencing in some genetic backgrounds (Rossmann et al., 2011), we also measured URA3

transcript levels by reverse-transcriptase quantitative PCR (Fig. 4D). In both assays, we find

no difference between VII-L-WT and VII-L-MUT, indicating that the reduced levels of Rap1

and Sir4 at *tlc1-tm* telomeres still allow functional silencing.

Rif2 prevents degradation of *tlc1-tm* telomeres

Rap1 regulates telomere length homeostasis through its recruitment of Rif1, Rif2, and Sir4.

Rif1 and Rif2 negatively regulate telomerase (Hardy et al., 1992; Wotton & Shore, 1997;

Levy & Blackburn, 2004). In contrast, Sir4 functions in a pathway to recruit telomerase to

telomeres (Hass & Zappulla, 2015). We tested what effect deleting *RIF1*, *RIF2*, and *SIR4*

would have on telomere length in *tlc1-tm* cells. Deletion of *SIR4* results in a small decrease in

telomere length in *TLC1* cells, as expected (Hass & Zappulla, 2015), but no noticeable

change in *tlc1-tm* cells (Fig. 5A). The lack of an effect could be due to Sir4 already being

reduced at *tlc1-tm* telomeres (Fig. 4A). Deletion of *RIF1* increases telomere length in both

TLC1 and *tlc1-tm* cells (Fig. 5B and Supplementary Fig. 2), despite Rif1 also being reduced

at *tlc1-tm* telomeres (Fig. 4A). Rif1 may still impact *tlc1-tm* telomeres because it plays a

much bigger role in telomere length homeostasis than Sir4.

Deletion of *RIF2* causes dramatic degradation of *tlc1-tm* telomeres, with a continuous

smear of telomeric DNA-hybridizing signal observed (Fig. 5B and Supplementary Fig. 2),

similar to *tlc1-tm* cells upon loss of telomerase (Fig. 3B). This finding, along with the

observation that Rif2 is still present at *tlc1-tm* telomeres (Fig. 4A), indicates that Rif2 has an

important role in protecting *tlc1-tm* telomeres. Since uncapped telomeres expose

chromosome ends to non-homologous end joining (NHEJ) and homologous recombination

(HR), we tested whether the *rif2Δ tlc1-tm* telomere profile would change by deleting *DNL4*

or *RAD52*, which are required for NHEJ and HR, respectively (Wilson et al., 1997;

Symington et al., 2014). The extensive smear was still observed in *dnl4Δ rif2Δ tlc1-tm* and

rad52Δ rif2Δ tlc1-tm triple mutants, but was reduced in intensity, especially at low molecular

weight, in *rad52Δ rif2Δ tlc1-tm* (Fig. 5C). *rif2Δ tlc1-tm* cells grow poorly, and this growth

defect is exacerbated by additional deletion of *RAD52*; *rad52Δ rif2Δ tlc1-tm* spores mostly

fail to germinate, and those that do grow even more poorly than *rif2Δ tlc1-tm* (Fig. 5D-E),

suggesting that Rad52-mediated HR helps cope with *rif2Δ tlc1-tm* uncapped telomeres.

Interestingly, *rif1Δ rif2Δ* and *rif1Δ rif2Δ tlc1-tm* cells have very similar telomere

profiles: very long telomeres but without the extensive degradation seen in *rif2Δ tlc1-tm* cells

(Fig. 5B). This observation is reminiscent of the elevated levels of telomere-telomere fusion

seen in *rif2Δ tel1Δ* cells, and not in *rif1Δ rif2Δ tel1Δ* cells (Marcand et al., 2008), but it is

currently unclear whether there is a common underlying mechanism that explains both

observations.

Rif2 protects *tlc1-tm* telomeres by inhibiting the MRX complex

Figure 5. Rif2 prevents degradation of *tlc1-tm* telomeres

(A-C) Telomere Southern blot analysis of strains of the indicated genotypes. Black arrowhead indicates the 1.8 kb telomere sequence-containing fragment loaded as control, as in Figure 1B.

(D) Colony sizes of haploid meiotic progeny derived from the sporulation of *rad52Δ/RAD52 rif2Δ/RIF2* and *rad52Δ/RAD52 rif2Δ/RIF2 tlc1-tm/tlc1-tm* diploid strains were measured and normalized to wild-type (top panel) or *tlc1-tm* (bottom panel). The number of colonies analyzed is indicated in parenthesis. Error bars show SEM.

(E) Tenfold serial dilutions of strains with the indicated genotypes were spotted on YPD plates and grown at 30°C.

TEL1 is epistatic to *RIF2* with respect to telomere length regulation; while *rif2Δ* cells have

long telomeres, *tel1Δ* and *rif2Δ tel1Δ* cells have very short telomeres (Craven & Petes, 1999).

Thus, we tested whether the same would be true in terms of the telomere degradation seen in

rif2Δ tlc1-tm cells. Deletion of *TEL1* shortens the telomeres of *tlc1-tm* cells, but does not

affect the telomere profile of *rif2Δ tlc1-tm* cells (Fig. 5B), indicating that Rif2 does not

inhibit Tel1 to prevent the degradation of *tlc1-tm* telomeres. Recent studies have shown that

Rif2 attenuates Tel1 activity at telomeres by inhibiting the MRX complex (consisting of

Mre11, Rad50, and Xrs2), which is responsible for recruiting Tel1 to telomeres (Hailemariam

et al., 2019; Sabourin et al., 2007). Rif2 discharges the ATP-bound state of Rad50, thereby

making the MRX complex incapable of activating Tel1 (Hailemariam et al., 2019). Similarly,

Rif2 enhancement of Rad50 ATPase activity limits MRX-mediated tethering of DNA ends

during double-strand break repair (Cassani et al., 2016). Rif2 also inhibits MRX-mediated

resection of telomeric DNA ends (Bonetti et al., 2010a; Bonetti et al., 2010b). Rif2 inhibition

of the MRX complex involves a direct interaction between the BAT/MIN motif of Rif2 with

the ATPase domain of Rad50 (Roisné-Hamelin et al., 2021; Khayat et al., 2021). Therefore,

we hypothesized that Rif2 could be inhibiting MRX-mediated degradation of *tlc1-tm*

telomeres. Consistent with this idea, we find that Mre11 is about fivefold more associated

with the VII-L-MUT telomere in comparison with the VII-L-WT telomere (Fig. 6A), and that

the telomere degradation observed in *rif2Δ tlc1-tm* cells is absent in *rad50Δ rif2Δ tlc1-tm*

cells (Fig. 6B).

The *rad50Δ tlc1-tm* and *rad50Δ rif2Δ tlc1-tm* cells (also *cdc13-1 tlc1-tm* cells,

discussed below) show a telomere profile reminiscent of type II survivors, which maintain

their telomeres in a telomerase-independent manner that relies on recombination-mediated

amplification of telomeric sequence (Lundblad & Blackburn, 1993; Teng & Zakian, 1999).

Independently generated isogenic isolates of type II survivors will have similar but slightly

Figure 6. Rif2 protects *tlc1-tm* telomeres by inhibiting the MRX complex and acts in parallel with the Yku complex

(A&C) ChIP-qPCR analysis of the association of Myc-tagged Mre11 and Yku70 to six Y' telomeres, the VII-L telomere, or to the non-telomeric *ARO1* locus. Untagged wild-type and *tlc1-tm* strains were used as controls. The mean percentage of input +/- SEM is shown (n=3, ** p<0.01).

(B&D) Telomere Southern blot analysis of strains of the indicated genotypes. Black arrowhead indicates the 1.8 kb telomere sequence-containing fragment loaded as control, as in Figure 1B. (B) Parental diploid 1 is *rad50Δ/RAD50 rif2Δ/RIF2*; parental diploid 2 is *rad50Δ/RAD50 rif2Δ/RIF2 tlc1-tm/tlc1-tm*. The last six lanes are derived from cells grown at 22°C. (D) Parental diploid 1 is *exo1Δ/EXO1 yku70Δ/YKU70*; parental diploid 2 is *exo1Δ/EXO1 yku70Δ/YKU70 tlc1-tm/tlc1-tm*.

(E) Colony sizes of haploid meiotic progeny derived from the sporulation of *rif2Δ/RIF2 yku70Δ/YKU70* and *rif2Δ/RIF2 yku70Δ/YKU70 tlc1-tm/tlc1-tm* diploid strains were measured and normalized to wild-type (top panel) or *tlc1-tm* (bottom panel). The number of colonies analyzed is indicated in parenthesis. Error bars show SEM. Representative images of dissected tetrads are shown on the right. Each column of colonies arose from a single tetrad.

different telomere profiles, much like that observed in different isolates of isogenic *tlc1-tm*

strains (as seen in *rad50Δ tlc1-tm* and *rad50Δ rif2Δ tlc1-tm* cells, but also in *tlc1-tm* strains in

general). However, this effect is still observed in *rad50Δ tlc1-tm* and *rad50Δ rif2Δ tlc1-tm*

strains with an additional deletion of *RAD52*, which is required for homologous

recombination and survivor formation (Lundblad & Blackburn, 1993; Claussin & Chang,

2015) (Supplementary Fig. 3A). Interestingly, the severe growth defect of *rad52Δ rif2Δ tlc1-*

tm strains (Fig. 5D) is suppressed by an additional deletion of *RAD50* (Supplementary Fig.

3B). At present, we do not know the reason for the slight telomere profile differences seen

between different isolates of isogenic *tlc1-tm* strains, but it may be related to a previous

observation that the very tip of the telomere has unique sequence features that affect telomere

capping and length regulation (Grossi et al., 2001).

Rif2 and the Yku complex act in parallel to protect *tlc1-tm* telomeres

The CST (Cdc13-Stn1-Ten1) and Yku (Yku70-Yku80) complexes are also important for

inhibiting nucleolytic degradation at telomeres (Garvik et al., 1995; Grandin et al., 1997;

Grandin et al., 2001; Gravel et al., 1998), and their importance might be increased at *tlc1-tm*

telomeres. Cdc13 and Yku70 are present at similar levels at VII-L-WT and VII-L-MUT

telomeres (Fig. 1F; Fig. 6C). The *cdc13-1* temperature-sensitive mutant causes an

accumulation of single-stranded telomeric DNA (Garvik et al., 1995), and is defective even at

the permissive temperature of 23°C (Paschini et al., 2012). *cdc13-1 tlc1-tm* cells do not

exhibit the extensive telomere degradation seen in *rif2Δ tlc1-tm* cells (Fig. 6B). In contrast,

we observe extensive telomere degradation in *yku70Δ tlc1-tm* cells (Fig. 6D). Telomere

degradation is not seen in *exo1Δ yku70Δ tlc1-tm* cells, suggesting that the Yku complex

inhibits the Exo1 exonuclease to protect *tlc1-tm* telomeres. *rif2Δ yku70Δ tlc1-tm* cells are not

viable (Fig. 6E), indicating that Rif2 and the Yku complex function in parallel to protect *tlc1-*

tm telomeres.

Rif2 recruitment to *tlc1-tm* telomeres requires a DNA end and the MRX complex

The telomere degradation seen in *tlc1-tm rif2Δ* cells has also been observed in other *tlc1*

template mutants, such as the *tlc1-476A* mutant (Fig. 7A), even without deletion of *RIF2*

(Chan et al., 2001; Lin et al., 2004). The *tlc1-476A* mutant changes the invariable CCC core

of the template region to CAC, causing the addition of long stretches of TG dinucleotide

repeats, occasionally interrupted by a TGG trinucleotide, that are predicted to disrupt Rap1

binding even more so than *tlc1-tm* repeats (Chan et al., 2001; Graham & Chambers, 1994).

We find that *tlc1-476A* mutants are completely dependent on Rif2 for survival, as *tlc1-476A*

rif2Δ double mutants are inviable (Fig. 7B).

To examine how Rif2 is able to associate with the Rap1-depleted telomeres of *tlc1-tm*

cells, we first asked whether a DNA end is required. We inserted 300 bp of either wild-type

or *tlc1-tm* telomeric sequence at an internal genomic locus on chromosome III (III-ITS;

chromosome III interstitial telomeric sequence) and assessed Rif2 association by ChIP-qPCR.

We find that Rif2 associates with the wild-type ITS, but not the *tlc1-tm* ITS (Fig. 7C),

indicating that Rap1-independent recruitment of Rif2 to *tlc1-tm* sequence requires a DNA

end. Since the MRX complex binds to DNA ends and telomeres (Oh & Symington, 2018),

and Rif2 is known to interact with both Rad50 and Xrs2 (Hirano et al., 2009; Hailemariam et

al., 2019; Roisné-Hamelin et al., 2021), Rif2 recruitment to *tlc1-tm* telomeres might require

the MRX complex. Consistent with this hypothesis, we find that Rif2 association to the *tlc1-*

tm telomeres is lost in *rad50Δ* strains (Fig. 7D).

Discussion

Figure 7. Rif2 association to *tlc1-tm* telomeres requires the MRX complex

(A) Telomere Southern blot analysis of strains of the indicated genotypes. Black arrowhead indicates the 1.8 kb telomere sequence-containing fragment loaded as control, as in Figure 1B.

(B) Colony sizes of haploid meiotic progeny derived from the sporulation of a *rif2Δ/RIF2 yku70Δ/YKU70 tlc1-476A/TLC1* diploid strain were measured and normalized to wild-type. The number of colonies analyzed is indicated in parenthesis. Error bars show SEM. Representative images of dissected tetrads are shown on the right. Each column of colonies arose from a single tetrad.

(C) ChIP-qPCR analysis of the association of Myc-tagged Rif2 to a 300-bp ITS located on chromosome III containing either wild-type or *tlc1-tm* sequence. The mean percentage of input +/- SEM is shown (n=3, *** p<10^{-3}).

(D) ChIP-qPCR analysis of the association of Myc-tagged Rif2 to six Y' telomeres, the VII-L telomere, or to the non-telomeric *ARO1* locus in a *rad50Δ* background. Untagged strains were used as controls. The mean percentage of input +/- SEM is shown (n=3, **** p<10^{-4}).

In this study, we investigated the consequences of depleting Rap1 from *S. cerevisiae*

telomeres. Our findings suggest that impairing Rap1 binding does not significantly affect cell

proliferation. The depletion of telomeric Rap1 causes defects in telomere length regulation

and telomere capping. Surprisingly, Rif2 is still recruited to Rap1-depleted telomeres, and its

recruitment is dependent on the MRX complex. *tlc1-tm* telomeres become extensively

degraded in the absence of Rif2 or the Yku complex, and *rif2Δ yku70Δ tlc1-tm* triple mutants

are inviable. Thus, Rap1, Rif2, and the Yku complex perform separate tasks at telomeres that

are together essential for telomere capping and cell viability.

Telomere capping is essential for cell viability; yeast cells cannot survive the loss of a

single telomere (Sandell & Zakian, 1993). Telomere capping is executed by telomere-binding

proteins, so it was surprising to find that yeast cells can survive when Rap1, the main protein

that binds double-stranded telomeric DNA, is significantly reduced at telomeres. Previous

studies have reported that telomeres of *S. cerevisiae* expressing mutant telomerase that adds

TTAGGG repeats are also devoid of Rap1; however, the absence of Rap1 in this situation

appears to be compensated by the binding of Tbf1 to the TTAGGG sequence (Alexander &

Zakian, 2003; Brevet et al., 2003; Berthiau et al., 2006). In contrast, there is no change in

Tbf1 binding at *tlc1-tm* telomeres (Fig. 1G). The difference between yeast telomeres with

TTAGGG repeats and *tlc1-tm* telomeres is further highlighted by the fact that Rif2 is absent

at the former (Alexander & Zakian, 2003), but is crucial for telomere protection of the latter.

In addition, multiple viable *tlc1* template mutants, including *tlc1-tm*, result in the addition of

mutant telomeric sequence that does not conform to the consensus binding sites of both Rap1

and Tbf1 (Prescott & Blackburn, 2000; Förstemann et al., 2003; Lin et al., 2004), supporting

the notion that telomeres can remain sufficiently capped in the absence of both proteins.

Previously studied *tlc1* template mutations that abolish Rap1 binding to the resulting

mutant telomere sequences generally fall into two categories; they either cause telomere

shortening or they cause rapid telomere elongation, which in some cases is accompanied by

extensive telomere degradation (Prescott & Blackburn, 2000; Lin et al., 2004), similar to that

seen in *rif2Δ tlc1-tm* mutants (Fig. 5B). The first category is likely due to a loss of telomerase

enzymatic activity, because most *tlc1* template mutations result in a reduction in the

nucleotide addition processivity of telomerase (Förstemann et al., 2003). The extent of

telomere elongation and degradation in the second category appears to be correlated with the

decrease in Rap1 binding (Prescott & Blackburn, 2000). Rap1 association to fully mutant

tlc1-tm telomeres is reduced to approximately 13% compared to wild-type telomeres (Fig.

1D). We suspect that this level of Rap1 prevents the more extensive elongation and

degradation seen in several other *tlc1* template mutants (e.g. *tlc1-476A*; Fig. 7A) that likely

have even less Rap1 telomere association (Prescott & Blackburn, 2000; Lin et al., 2004).

Consistent with this hypothesis, *tlc1Δ* cells derived from the sporulation of *tlc1-tm/tlc1Δ*

diploids with mutant telomeres also exhibit extensive degradation (Fig. 3B), which is likely

the result of further reducing Rap1 telomere association due to telomere shortening.

Our findings build upon previous work to show that Rap1 and Rif2 inhibit the MRX

complex to prevent telomere degradation. First, decreased telomere association of Rap1 in

tlc1-tm cells (Fig. 1D) is accompanied by an increase in Mre11 telomere association (Fig.

6A), which is consistent with a previous report showing that Rap1 binding inhibits Mre11

recruitment (Negrini et al., 2007). Second, the telomere degradation observed in *rif2Δ tlc1-tm*

cells is eliminated by deletion of *RAD50* (Fig. 6B), which is consistent with previous studies

showing that Rif2 inhibits the MRX complex by discharging the ATP-bound form of Rad50

(Cassani et al., 2016; Hailemariam et al., 2019). Rif2 and the Yku complex have previously

been shown to separately inhibit the resection of telomeric ends in *TLC1* cells, but *rif2Δ ykuΔ*

double mutants remain viable (Bonetti et al., 2010a; Bonetti et al., 2010b). In contrast, we

find that *rif2Δ yku70Δ tlc1-tm* cells are inviable (Fig. 6E), indicating that Rap1 has an

important role alongside Rif2 and the Yku complex to inhibit telomere degradation.

Furthermore, the inviability of *tlc1-476A rif2Δ* double mutants (Fig. 7B) indicates that Rap1

and Rif2 are more important than the Yku complex in this regard.

The redundant capping mechanisms offer an explanation for the rapid evolution of the

telomere sequence and telomere binding proteins in budding yeast species of the

Saccharomycotina subdivision, of which *S. cerevisiae* is a member (Steinberg-Neifach &

Lue, 2015). Alterations in the template region of telomerase RNA or in the DNA binding

domain of telomere-binding proteins are more easily tolerated if there are redundant

mechanisms to ensure sufficient telomere capping and, therefore, cell viability. Indeed, the

budding yeast *Kluyveromyces lactis* can also tolerate changes to the template region of its

telomerase RNA that disrupts binding to Rap1 (Krauskopf & Blackburn, 1996). The more

distantly related fission yeast *Schizosaccharomyces pombe* can also survive without its main

telomere-binding protein, Taz1 (Cooper et al., 1997), suggesting that redundancy of telomere

capping mechanisms has also evolved independently outside of the Saccharomycotina

subdivision.

The association of Rif2 to Rap1-depleted *tlc1-tm* telomeres was unexpected given the

well-characterized role of the Rap1 C-terminal domain in recruiting Rif2 to telomeres

(Wotton & Shore, 1997; Shi et al., 2013). We find that this Rap1-independent recruitment of

Rif2 is dependent on the MRX complex (Fig. 7D), which itself is increased at *tlc1-tm*

telomeres (as determined by ChIP-qPCR analysis of Mre11; Fig. 6A). The MRX complex

plays a central role in the repair of double-strand breaks (DSBs) (Oh & Symington, 2018), so

why does Rif2 play such a negligible role? It has recently been reported that Rif2 and Sae2

have opposing roles in regulating the MRX complex, and the role of Rif2 in DSB repair is

more readily detected in the absence of Sae2 (Marsella et al., 2021). While Sae2 has a

prominent role in DSB repair, it has a minor role at telomeres (Bonetti et al., 2009); the

inverse is true of Rif2 at DSBs and telomeres. Thus, Sae2 appears to limit Rif2 function at

DSBs, and Rif2 may limit the role of Sae2 at telomeres. In wild-type cells, the prominent role

of Rif2 at telomeres can be explained by its recruitment by Rap1. However, it is unclear how

Rif2 maintains its prominent role at telomeres without Rap1-dependent recruitment. One

possibility is that there is preferential recruitment of Sae2 to DSBs compared to telomeres.

Further studies are needed to examine this hypothesis.

Materials and Methods

Yeast strains and plasmids

Standard yeast media and growth conditions were used (Sherman, 2002; Lundblad & Struhl,

2010). Yeast strains used in this study are listed in Supplementary Table S1. The artificial

VII-L telomere was created essentially as previously described (Gottschling et al., 1990), by

transforming restriction-digested pVII-L URA-TEL plasmid into yeast. To obtain different

versions of the VII-L telomere, this plasmid was modified by replacing the original 81 bp of

telomeric repeats with oligonucleotides containing wild-type or *tlc1-tm* telomeric sequence of

different length (Supplementary Table S2). Myc and protA C-terminal epitope-tagged

proteins were created by integrative transformation of 13 copies of the human c-myc (Myc)

epitope or the Staphylococcal protein A IgG binding domain.

Southern blotting

Southern blots to detect native telomeres were performed essentially as previously described

(van Mourik et al., 2018). Yeast genomic DNA was isolated using a Wizard Genomic DNA

Purification Kit (Promega) and digested with XhoI. For each sample, 4 µg of digested

genomic DNA, along with 1.25 ng of BsmAI-digested pYt103 (Askree et al., 2004), was

separated on a 0.8% (w/v) agarose gel, and transferred to a Hybond-N+ membrane (GE

Healthcare). The membrane was hybridized to telomere-specific digoxygenin-labeled probe

(wild type probe: 5′-CACCACACCCACACACCACACCCACA-3′; *tlc1-tm* mutant probe:

5′-ACCACACCACACCACACACACCACAC-3′). For detection of the artificial VII-L

telomere, a similar procedure was performed, except that 6 μg yeast genomic DNA was

digested with EcoRV and the membrane was hybridized at 42ºC with a digoxigenin-labeled

probe complementary to *URA3* sequence. Unless otherwise mentioned (i.e. Fig. 1C and 3B),

all strains were propagated for at least 100 generations before Southern blot analysis.

Chromatin immunoprecipitation and quantitative PCR (ChIP-qPCR)

ChIP-qPCR was performed essentially as previously described (Graf et al., 2017). For protein

A ChIP, IgG Sepharose 6 Fast Flow beads (GE Healthcare) were used. For Myc ChIP,

nProtein A Sepharose 4 Fast Flow beads (GE Healthcare) were used; after preclearing, 9 μl

anti-c-Myc Monoclonal antibody (Clontech/Takara) were added to each sample. qPCR was

performed using a LightCycler 480 II (Roche) and SYBR-Green (Thermo Scientific)

detection with an annealing temperature of 60ºC (40 cycles). Primers are listed in

Supplementary Table S3. Measured Cq values were corrected to input and graphs were

created using R. All strains were propagated for ~50 generations before ChIP-qPCR analysis.

Electrophoretic mobility shift assay (EMSA)

The recombinant *S. cerevisiae* Rap1 full-length protein was expressed in *E. coli* BL21 cells

as previously described (Wahlin & Cohn, 2000). The ability of Rap1 to bind the *tlc1-tm*

telomeric sequences was assessed by EMSA with two double-stranded (ds) oligonucleotides

containing representative sequences; Mut-1 5'-

*GTCATACGTCACAC*TGTGGTGTGTGGTGTGGTGTGTGGTGGTGGTGTGTGT-3' (37

bp) and Mut-2 5'-

*GTCATACGTCACAC*TGTGGTGTGTGTGGTGTGTGGATTTGGTGTGTGG-3' (34 bp).

The respective G-rich forward strands and C-rich reverse strands were annealed in 1 mM

Tris-HCl, pH 8.0, 0.1 mM $MgCl_2$ by boiling for 2 min and slowly cooling down to room

temperature. The correct annealing was guided by the 14 nt non-telomeric region indicated in

italics. A ds-oligonucleotide with wild-type telomeric sequence was used as the positive

binding control: 5'-TGTGGTGTGTGGGTGTGTG-3' (19 bp). The 5' ends of ds-probes were

radioactively labelled with $[\gamma\text{-}^{32}P]$-ATP using T4 polynucleotide kinase (New England

Biolabs), purified on Illustra Microspin G-25 columns (GE), and diluted in 10 mM Tris-HCl,

pH 7.5. In binding assays, 10 fmol of labelled probe was mixed with 1.5 µg competitor mix

(0.5 µg each of sheared *E.coli* DNA (~250 bp), salmon sperm DNA, and yeast t-RNA) in 1x

Binding buffer (10 mM Tris-HCl, pH 7.5, 7 mM $MgCl_2$, 8% glycerol), and varying amounts

(~0.1-0.4 µg) of Rap1 protein extract, in a total of 15 µl. The binding reaction was incubated

at 25°C for 15 min and then loaded onto 4% polyacrylamide gels (29:1 acrylamide:bis-

acrylamide) and run in 1x TBE (89 mM Tris-borate, 2 mM EDTA, pH 8.0), 150V at 4°C.

Radioactive signals from dried gels were visualized with a Typhoon FLA 9500 biomolecular

imager (GE Life Sciences).

Inducible STEX (iSTEX) assay

iSTEX was performed essentially as previously described (Strecker et al., 2017), except that

the starting P_{GALL}-*EST1* strain contained the *tlc1-tm* allele. This strain was then transformed

with a PCR product containing the wild-type *TLC1* allele and immediately grown on media

containing glucose, which shut off expression of *EST1*. Successfully transformed cells were

cultured in YPD media and eventually arrested in late G1 phase by the addition of alpha

factor (Sigma-Aldrich). Cells were washed and resuspended in YPGal to induce the

expression of wild-type telomerase, allowing the extension of *tlc1-tm* mutant telomeres

during a single cell division.

Liquid culture senescence assay

Senescence assays in liquid culture were performed essentially as previously described (van

Mourik et al., 2016).

TERRA and URA3 level analysis by quantitative reverse transcription and PCR

TERRA levels were measured essentially as previously described (Graf et al., 2017), while

URA3 and ACT1 levels were measured by one-step RT-PCR with specific primers

(Supplementary Table S3) following the standard protocol of the RNeasy® Mini kit

(Qiagen). All strains were propagated for at least 100 generations before analysis.

Data availability

488 All data generated or analyzed during this study are included in this published article (and its

489 supplementary information files).

Acknowledgments

We thank K. Paeschke and L. Wanders for critical reading of the manuscript; S. Pandey and

O. Rosas Bringas for help with data analysis; and H.G. Kazemier for technical assistance.

F.R.R.B. was supported by a CONACYT scholarship. S.S. was supported by an EMBO

Short-Term Fellowship to visit the lab of B. Luke to perform TERRA measurements and to

gain experience with ChIP-qPCR experiments, and we thank B. Luke and M. Graf for their

technical assistance. M. Cohn was supported by grants from the Carl Trygger Foundation, the

Erik Philip-Sörensen Foundation, and the Royal Physiographic Society in Lund. Work in the

laboratory of M. Chang was supported by a Vidi grant from the Netherlands Organization for

Scientific Research.

Author Contributions

503   Conceptualization, F.R.R.B., S.S., M. Chang; Investigation, F.R.R.B., S.S., P.Z., M. Cohn;

504   Writing – Original Draft, F.R.R.B., S.S., M. Chang; Writing – Reviewing & Editing,

505   F.R.R.B., S.S., P.Z., M. Cohn, M. Chang; Supervision, M. Chang.

Competing Interests

508   The authors declare no competing interests.

References

Alexander, M. K., & Zakian, V. A. (2003, Apr 1). Rap1p telomere association is not required for mitotic stability of a C$_3$TA$_2$ telomere in yeast. *EMBO J, 22*(7), 1688-1696. https://doi.org/10.1093/emboj/cdg154

Aparicio, O. M., Billington, B. L., & Gottschling, D. E. (1991, Sep 20). Modifiers of position effect are shared between telomeric and silent mating-type loci in *S. cerevisiae* [Research Support, Non-U.S. Gov't
Research Support, U.S. Gov't, Non-P.H.S.
Research Support, U.S. Gov't, P.H.S.]. *Cell, 66*(6), 1279-1287. http://www.ncbi.nlm.nih.gov/pubmed/1913809

Askree, S. H., Yehuda, T., Smolikov, S., Gurevich, R., Hawk, J., Coker, C., Krauskopf, A., Kupiec, M., & McEachern, M. J. (2004, Jun 08). A genome-wide screen for *Saccharomyces cerevisiae* deletion mutants that affect telomere length. *Proc Natl Acad Sci U S A, 101*(23), 8658-8663. https://doi.org/10.1073/pnas.0401263101

Azad, G. K., & Tomar, R. S. (2016, Jun 1). The multifunctional transcription factor Rap1: a regulator of yeast physiology. *Front Biosci (Landmark Ed), 21*, 918-930. https://doi.org/10.2741/4429

Berthiau, A. S., Yankulov, K., Bah, A., Revardel, E., Luciano, P., Wellinger, R. J., Geli, V., & Gilson, E. (2006, Feb 22). Subtelomeric proteins negatively regulate telomere elongation in budding yeast. *EMBO J, 25*(4), 846-856. https://doi.org/10.1038/sj.emboj.7600975

Bonetti, D., Clerici, M., Anbalagan, S., Martina, M., Lucchini, G., & Longhese, M. P. (2010a). Shelterin-like proteins and Yku inhibit nucleolytic processing of *Saccharomyces cerevisiae* telomeres. *PLoS Genet, 6*(5), e1000966. https://doi.org/10.1371/journal.pgen.1000966

Bonetti, D., Clerici, M., Manfrini, N., Lucchini, G., & Longhese, M. P. (2010b, Nov 30). The MRX complex plays multiple functions in resection of Yku- and Rif2-protected DNA ends. *PLoS One, 5*(11), e14142. https://doi.org/10.1371/journal.pone.0014142

Bonetti, D., Martina, M., Clerici, M., Lucchini, G., & Longhese, M. P. (2009, Jul 10). Multiple pathways regulate 3' overhang generation at *S. cerevisiae* telomeres. *Mol Cell, 35*(1), 70-81. https://doi.org/10.1016/j.molcel.2009.05.015

Brevet, V., Berthiau, A. S., Civitelli, L., Donini, P., Schramke, V., Geli, V., Ascenzioni, F., & Gilson, E. (2003, Apr 1). The number of vertebrate repeats can be regulated at yeast

551   telomeres by Rap1-independent mechanisms. *EMBO J, 22*(7), 1697-1706.
552   https://doi.org/10.1093/emboj/cdg155
553

554 Brigati, C., Kurtz, S., Balderes, D., Vidali, G., & Shore, D. (1993, Feb). An essential yeast
555   gene encoding a TTAGGG repeat-binding protein. *Mol Cell Biol, 13*(2), 1306-1314.
556   https://doi.org/10.1128/mcb.13.2.1306
557

558 Buchman, A. R., Kimmerly, W. J., Rine, J., & Kornberg, R. D. (1988, Jan). Two DNA-
559   binding factors recognize specific sequences at silencers, upstream activating
560   sequences, autonomously replicating sequences, and telomeres in *Saccharomyces*
561   *cerevisiae. Mol Cell Biol, 8*(1), 210-225. https://doi.org/10.1128/mcb.8.1.210
562

563 Cassani, C., Gobbini, E., Wang, W., Niu, H., Clerici, M., Sung, P., & Longhese, M. P. (2016,
564   Feb). Tel1 and Rif2 Regulate MRX Functions in End-Tethering and Repair of DNA
565   Double-Strand Breaks. *PLoS Biol, 14*(2), e1002387.
566   https://doi.org/10.1371/journal.pbio.1002387
567

568 Chan, S. W., Chang, J., Prescott, J., & Blackburn, E. H. (2001, Aug 21). Altering telomere
569   structure allows telomerase to act in yeast lacking ATM kinases. *Curr Biol, 11*(16),
570   1240-1250. https://doi.org/10.1016/s0960-9822(01)00391-8
571

572 Chang, M., Arneric, M., & Lingner, J. (2007, Oct 1). Telomerase repeat addition processivity
573   is increased at critically short telomeres in a Tel1-dependent manner in
574   *Saccharomyces cerevisiae. Genes & Development, 21*(19), 2485-2494.
575   https://doi.org/10.1101/gad.1588807
576

577 Claussin, C., & Chang, M. (2015). The many facets of homologous recombination at
578   telomeres. *Microbial Cell, 2*(9), 308-321. https://doi.org/10.15698/mic2015.09.224
579

580 Conrad, M. N., Wright, J. H., Wolf, A. J., & Zakian, V. A. (1990, Nov 16). RAP1 protein
581   interacts with yeast telomeres in vivo: overproduction alters telomere structure and
582   decreases chromosome stability. *Cell, 63*(4), 739-750. https://doi.org/10.1016/0092-
583   8674(90)90140-a
584

585 Cooper, J. P., Nimmo, E. R., Allshire, R. C., & Cech, T. R. (1997, Feb 20). Regulation of
586   telomere length and function by a Myb-domain protein in fission yeast. *Nature,*
587   *385*(6618), 744-747.
588   http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=C
589   itation&list_uids=9034194
590

Craven, R. J., & Petes, T. D. (1999, Aug). Dependence of the regulation of telomere length
on the type of subtelomeric repeat in the yeast *Saccharomyces cerevisiae. Genetics,*

152(4), 1531-1541.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=C
itation&list_uids=10430581

de Lange, T. (2018, Nov 23). Shelterin-Mediated Telomere Protection. *Annual Review of*
Genetics, 52, 223-247. https://doi.org/10.1146/annurev-genet-032918-021921

Douglas, M. E., & Diffley, J. F. X. (2021, Jun 4). Budding yeast Rap1, but not telomeric
DNA, is inhibitory for multiple stages of DNA replication in vitro. *Nucleic Acids Res,*
49(10), 5671-5683. https://doi.org/10.1093/nar/gkab416

Eldridge, A. M., Halsey, W. A., & Wuttke, D. S. (2006, Jan 24). Identification of the
determinants for the specific recognition of single-strand telomeric DNA by Cdc13.
Biochemistry, 45(3), 871-879. https://doi.org/10.1021/bi0512703

Förstemann, K., & Lingner, J. (2001, Nov). Molecular basis for telomere repeat divergence in
budding yeast. *Mol Cell Biol, 21*(21), 7277-7286.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=C
itation&list_uids=11585910

Förstemann, K., Zaug, A. J., Cech, T. R., & Lingner, J. (2003, Mar 15). Yeast telomerase is
specialized for C/A-rich RNA templates. *Nucleic Acids Res, 31*(6), 1646-1655.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=C
itation&list_uids=12626706

Gartenberg, M. R., & Smith, J. S. (2016, Aug). The Nuts and Bolts of Transcriptionally
Silent Chromatin in *Saccharomyces cerevisiae*. *Genetics, 203*(4), 1563-1599.
https://doi.org/10.1534/genetics.112.145243

Garvik, B., Carson, M., & Hartwell, L. (1995, Nov). Single-stranded DNA arising at
telomeres in *cdc13* mutants may constitute a specific signal for the *RAD9* checkpoint.
Mol Cell Biol, 15(11), 6128-6138.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=C
itation&list_uids=7565765

Gottschling, D. E., Aparicio, O. M., Billington, B. L., & Zakian, V. A. (1990). Position effect
at *S. cerevisiae* telomeres: Reversible repression of Pol II transcription. *Cell, 63*, 751-
762. https://doi.org/10.1016/0092-8674(90)90141-Z

Graf, M., Bonetti, D., Lockhart, A., Serhal, K., Kellner, V., Maicher, A., Jolivet, P., Teixeira,
M. T., & Luke, B. (2017, Jun 29). Telomere Length Determines TERRA and R-Loop

Regulation through the Cell Cycle. *Cell, 170*(1), 72-85 e14.
https://doi.org/10.1016/j.cell.2017.06.006

Graham, I. R., & Chambers, A. (1994). Use of a selection technique to identify the diversity of binding sites for the yeast RAP1 transcription factor. *Nucleic Acids Research, 22*, 124-130.

Grandin, N., Damon, C., & Charbonneau, M. (2001, Mar 1). Ten1 functions in telomere end protection and length regulation in association with Stn1 and Cdc13. *EMBO J, 20*(5), 1173-1183. https://doi.org/10.1093/emboj/20.5.1173

Grandin, N., Reed, S. I., & Charbonneau, M. (1997, Feb 15). Stn1, a new *Saccharomyces cerevisiae protein*, is implicated in telomere size regulation in association with Cdc13. *Genes Dev, 11*(4), 512-527.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=9042864

Gravel, S., Larrivee, M., Labrecque, P., & Wellinger, R. J. (1998, May 1). Yeast Ku as a regulator of chromosomal DNA end structure. *Science, 280*(5364), 741-744.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=9563951

Grossi, S., Bianchi, A., Damay, P., & Shore, D. (2001, Dec). Telomere formation by Rap1p binding site arrays reveals end-specific length regulation requirements and active telomeric recombination. *Mol Cell Biol, 21*(23), 8117-8128.
https://doi.org/10.1128/MCB.21.23.8117-8128.2001

Guérin, T. M., Béneut, C., Barinova, N., López, V., Lazar-Stefanita, L., Deshayes, A., Thierry, A., Koszul, R., Dubrana, K., & Marcand, S. (2019, Jul 11). Condensin-Mediated Chromosome Folding and Internal Telomeres Drive Dicentric Severing by Cytokinesis. *Mol Cell, 75*(1), 131-144 e133.
https://doi.org/10.1016/j.molcel.2019.05.021

Hailemariam, S., De Bona, P., Galletto, R., Hohl, M., Petrini, J. H., & Burgers, P. M. (2019, Dec 6). The telomere-binding protein Rif2 and ATP-bound Rad50 have opposing roles in the activation of yeast Tel1(ATM) kinase. *J Biol Chem, 294*(49), 18846-18852. https://doi.org/10.1074/jbc.RA119.011077

Hardy, C. F., Sussel, L., & Shore, D. (1992, May). A RAP1-interacting protein involved in transcriptional silencing and telomere length regulation. *Genes Dev, 6*(5), 801-814.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=1577274

Hass, E. P., & Zappulla, D. C. (2015, Jul 28). The Ku subunit of telomerase binds Sir4 to recruit telomerase to lengthen telomeres in *S. cerevisiae*. *eLife, 4*. https://doi.org/10.7554/eLife.07750

Hirano, Y., Fukunaga, K., & Sugimoto, K. (2009, Feb 13). Rif1 and Rif2 inhibit localization of Tel1 to DNA ends. *Mol Cell, 33*(3), 312-322. https://doi.org/S1097-2765(09)00024-0 [pii]
10.1016/j.molcel.2008.12.027

Iglesias, N., Redon, S., Pfeiffer, V., Dees, M., Lingner, J., & Luke, B. (2011, Jun). Subtelomeric repetitive elements determine TERRA regulation by Rap1/Rif and Rap1/Sir complexes in yeast. *EMBO Rep, 12*(6), 587-593. https://doi.org/10.1038/embor.2011.73

Khayat, F., Cannavo, E., Alshmery, M., Foster, W. R., Chahwan, C., Maddalena, M., Smith, C., Oliver, A. W., Watson, A. T., Carr, A. M., Cejka, P., & Bianchi, A. (2021, Jun 22). Inhibition of MRN activity by a telomere protein motif. *Nat Commun, 12*(1), 3856. https://doi.org/10.1038/s41467-021-24047-2

Krauskopf, A., & Blackburn, E. H. (1996, Sep 26). Control of telomere growth by interactions of RAP1 with the most distal telomeric repeats. *Nature, 383*(6598), 354-357. https://doi.org/10.1038/383354a0

Kyrion, G., Liu, K., Liu, C., & Lustig, A. J. (1993, Jul). RAP1 and telomere structure regulate telomere position effects in *Saccharomyces cerevisiae*. *Genes Dev, 7*(7A), 1146-1159. https://doi.org/10.1101/gad.7.7a.1146

Levy, D. L., & Blackburn, E. H. (2004, Dec). Counting of Rif1p and Rif2p on *Saccharomyces cerevisiae* telomeres regulates telomere length. *Mol Cell Biol, 24*(24), 10857-10867. https://doi.org/24/24/10857 [pii]
10.1128/MCB.24.24.10857-10867.2004

Lieb, J. D., Liu, X., Botstein, D., & Brown, P. O. (2001, Aug). Promoter-specific binding of Rap1 revealed by genome-wide maps of protein-DNA association. *Nat Genet, 28*(4), 327-334. https://doi.org/10.1038/ng569

Lin, J., Smith, D. L., & Blackburn, E. H. (2004, Apr). Mutant telomere sequences lead to impaired chromosome separation and a unique checkpoint response. *Mol Biol Cell, 15*(4), 1623-1634. http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=14742705

719 Lingner, J., Hughes, T. R., Shevchenko, A., Mann, M., Lundblad, V., & Cech, T. R. (1997,
720   Apr 25). Reverse transcriptase motifs in the catalytic subunit of telomerase. *Science,*
721   *276*(5312), 561-567.
722   http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=C
723   itation&list_uids=9110970
724

725 Lundblad, V., & Blackburn, E. H. (1993, Apr 23). An alternative pathway for yeast telomere
726   maintenance rescues *est1-* senescence. *Cell, 73*(2), 347-360. https://doi.org/0092-
727   8674(93)90234-H [pii]
728

729 Lundblad, V., & Struhl, K. (2010). Yeast. *Current Protocols in Molecular Biology, 92*(1).
730   https://doi.org/10.1002/0471142727.mb1300s92
731

732 Lustig, A. J., Kurtz, S., & Shore, D. (1990, Oct 26). Involvement of the silencer and UAS
733   binding protein RAP1 in regulation of telomere length. *Science, 250*(4980), 549-553.
734

735 Makovets, S., Herskowitz, I., & Blackburn, E. H. (2004, May). Anatomy and dynamics of
736   DNA replication fork movement in yeast telomeric regions. *Mol Cell Biol, 24*(9),
737   4019-4031.
738   http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=C
739   itation&list_uids=15082794
740

741 Marcand, S., Gilson, E., & Shore, D. (1997, Feb 14). A protein-counting mechanism for
742   telomere length regulation in yeast [Research Support, Non-U.S. Gov't
743 Research Support, U.S. Gov't, P.H.S.]. *Science, 275*(5302), 986-990.
744   http://www.ncbi.nlm.nih.gov/pubmed/9020083
745

746 Marcand, S., Pardo, B., Gratias, A., Cahun, S., & Callebaut, I. (2008, May 1). Multiple
747   pathways inhibit NHEJ at telomeres. *Genes Dev, 22*(9), 1153-1158.
748   https://doi.org/10.1101/gad.455108
749

750 Marsella, A., Gobbini, E., Cassani, C., Tisi, R., Cannavo, E., Reginato, G., Cejka, P., &
751   Longhese, M. P. (2021, Mar 30). Sae2 and Rif2 regulate MRX endonuclease activity
752   at DNA double-strand breaks in opposite manners. *Cell Rep, 34*(13), 108906.
753   https://doi.org/10.1016/j.celrep.2021.108906
754

755 Moretti, P., Freeman, K., Coodly, L., & Shore, D. (1994, Oct 1). Evidence that a complex of
756   SIR proteins interacts with the silencer and telomere-binding protein RAP1. *Genes*
757   *Dev, 8*(19), 2257-2269. https://www.ncbi.nlm.nih.gov/pubmed/7958893
758

Negrini, S., Ribaud, V., Bianchi, A., & Shore, D. (2007). DNA breaks are masked by multiple Rap1 binding in yeast: Implications for telomere capping and telomerase regulation. *Genes & Development, 21*, 292-302. https://doi.org/10.1101/gad.400907

Nugent, C. I., Hughes, T. R., Lue, N. F., & Lundblad, V. (1996, Oct 11). Cdc13p: a single-strand telomeric DNA-binding protein with a dual role in yeast telomere maintenance. *Science, 274*(5285), 249-252. http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=8824190

Oh, J., & Symington, L. S. (2018, Nov 29). Role of the Mre11 Complex in Preserving Genome Integrity. *Genes (Basel), 9*(12). https://doi.org/10.3390/genes9120589

Pardo, B., & Marcand, S. (2005, Sep 07). Rap1 prevents telomere fusions by nonhomologous end joining. *EMBO J, 24*(17), 3117-3127. https://doi.org/10.1038/sj.emboj.7600778

Paschini, M., Toro, T. B., Lubin, J. W., Braunstein-Ballew, B., Morris, D. K., & Lundblad, V. (2012, May). A naturally thermolabile activity compromises genetic analysis of telomere function in *Saccharomyces cerevisiae*. *Genetics, 191*(1), 79-93. https://doi.org/10.1534/genetics.111.137869

Prescott, J. C., & Blackburn, E. H. (2000, Apr). Telomerase RNA template mutations reveal sequence-specific requirements for the activation and repression of telomerase action at telomeres. *Mol Cell Biol, 20*(8), 2941-2948. https://doi.org/10.1128/mcb.20.8.2941-2948.2000

Preti, M., Ribeyre, C., Pascali, C., Bosio, M. C., Cortelazzi, B., Rougemont, J., Guarnera, E., Naef, F., Shore, D., & Dieci, G. (2010, May 28). The telomere-binding protein Tbf1 demarcates snoRNA gene promoters in *Saccharomyces cerevisiae*. *Mol Cell, 38*(4), 614-620. https://doi.org/10.1016/j.molcel.2010.04.016

Roisné-Hamelin, F., Pobiega, S., Jézéquel, K., Miron, S., Dépagne, J., Veaute, X., Busso, D., Du, M. L., Callebaut, I., Charbonnier, J. B., Cuniasse, P., Zinn-Justin, S., & Marcand, S. (2021, May 12). Mechanism of MRX inhibition by Rif2 at telomeres. *Nat Commun, 12*(1), 2763. https://doi.org/10.1038/s41467-021-23035-w

Rossmann, M. P., Luo, W., Tsaponina, O., Chabes, A., & Stillman, B. (2011, Apr 8). A common telomeric gene silencing assay is affected by nucleotide metabolism. *Mol Cell, 42*(1), 127-136. https://doi.org/10.1016/j.molcel.2011.03.007

Sabourin, M., Tuzon, C. T., & Zakian, V. A. (2007, Jul 24). Telomerase and Tel1p Preferentially Associate with Short Telomeres in *S. cerevisiae*. *Mol Cell*.

801   http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=C
802   itation&list_uids=17656141
803

804   Sandell, L. L., & Zakian, V. A. (1993, Nov 19). Loss of a yeast telomere: arrest, recovery,
805   and chromosome loss. *Cell, 75*(4), 729-739.
806   https://www.ncbi.nlm.nih.gov/pubmed/8242745
807

808   Shampay, J., Szostak, J. W., & Blackburn, E. H. (1984). DNA sequences of telomeres
809   maintained in yeast. *Nature, 310*, 154-157. https://doi.org/10.1038/310154a0
810

811   Sherman, F. (2002). Getting started with yeast. *Methods Enzymol, 350*, 3-41.
812   https://www.ncbi.nlm.nih.gov/pubmed/12073320
813

814   Shi, T., Bunker, R. D., Mattarocci, S., Ribeyre, C., Faty, M., Gut, H., Scrima, A., Rass, U.,
815   Rubin, S. M., Shore, D., & Thoma, N. H. (2013, Jun 6). Rif1 and Rif2 shape telomere
816   function and architecture through multivalent Rap1 interactions. *Cell, 153*(6), 1340-
817   1353. https://doi.org/10.1016/j.cell.2013.05.007
818

819   Shore, D., & Nasmyth, K. (1987, Dec 4). Purification and cloning of a DNA binding protein
820   from yeast that binds to both silencer and activator elements [Research Support, Non-
821   U.S. Gov't]. *Cell, 51*(5), 721-732. http://www.ncbi.nlm.nih.gov/pubmed/3315231
822

823   Steinberg-Neifach, O., & Lue, N. F. (2015). Telomere DNA recognition in Saccharomycotina
824   yeast: potential lessons for the co-evolution of ssDNA and dsDNA-binding proteins
825   and their target sites. *Front Genet, 6*, 162. https://doi.org/10.3389/fgene.2015.00162
826

827   Stinus, S., Paeschke, K., & Chang, M. (2017, Oct 20). Telomerase regulation by the Pif1
828   helicase: a length-dependent effect? *Curr Genet*. https://doi.org/10.1007/s00294-017-
829   0768-6
830

831   Strecker, J., Stinus, S., Caballero, M. P., Szilard, R. K., Chang, M., & Durocher, D. (2017,
832   Aug 03). A sharp Pif1-dependent threshold separates DNA double-strand breaks from
833   critically short telomeres. *eLife, 6*, e23783. https://doi.org/10.7554/eLife.23783
834

835   Symington, L. S., Rothstein, R., & Lisby, M. (2014, Nov). Mechanisms and Regulation of
836   Mitotic Recombination in *Saccharomyces cerevisiae*. *Genetics, 198*(3), 795-835.
837   https://doi.org/10.1534/genetics.114.166140
838

Teixeira, M. T., Arneric, M., Sperisen, P., & Lingner, J. (2004, Apr 30). Telomere length
homeostasis is achieved via a switch between telomerase- extendible and -
nonextendible states. *Cell, 117*(3), 323-335.

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=C
itation&list_uids=15109493

Teng, S. C., & Zakian, V. A. (1999, Dec). Telomere-telomere recombination is an efficient
bypass pathway for telomere maintenance in *Saccharomyces cerevisiae*. *Mol Cell*
Biol, 19(12), 8083-8093.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=C
itation&list_uids=10567534

van Mourik, P. M., de Jong, J., Agpalo, D., Claussin, C., Rothstein, R., & Chang, M. (2016).
Recombination-Mediated Telomere Maintenance in *Saccharomyces cerevisiae* Is Not
Dependent on the Shu Complex. *PLoS One, 11*(3), e0151314.
https://doi.org/10.1371/journal.pone.0151314

van Mourik, P. M., de Jong, J., Sharma, S., Kavsek, A., Chabes, A., & Chang, M. (2018, Jul
31). Upregulation of dNTP Levels After Telomerase Inactivation Influences
Telomerase-Independent Telomere Maintenance Pathway Choice in *Saccharomyces*
cerevisiae. *G3 (Bethesda), 8*(8), 2551-2558. https://doi.org/10.1534/g3.118.200280

Vodenicharov, M. D., Laterreur, N., & Wellinger, R. J. (2010, Sep 1). Telomere capping in
non-dividing yeast cells requires Yku and Rap1. *EMBO J, 29*(17), 3007-3019.
https://doi.org/10.1038/emboj.2010.155

Wahlin, J., & Cohn, M. (2000). *Saccharomyces cerevisiae* RAP1 binds to telomeric
sequences with spatial flexibility. *Nucleic Acids Research, 28*, 2292-2301.
https://doi.org/10.1093/nar/28.12.2292

Wellinger, R. J. (2014, Mar 20). In the end, what's the problem? *Mol Cell, 53*(6), 855-856.
https://doi.org/10.1016/j.molcel.2014.03.008

Wellinger, R. J., & Zakian, V. A. (2012, Aug). Everything you ever wanted to know about
Saccharomyces cerevisiae telomeres: beginning to end [Research Support, N.I.H.,
Extramural
Research Support, Non-U.S. Gov't
Review]. *Genetics, 191*(4), 1073-1105. https://doi.org/10.1534/genetics.111.137851

Wilson, T. E., Grawunder, U., & Lieber, M. R. (1997, Jul 31). Yeast DNA ligase IV mediates
non-homologous DNA end joining. *Nature, 388*(6641), 495-498.
https://doi.org/10.1038/41365

Wotton, D., & Shore, D. (1997, Mar 15). A novel Rap1p-interacting factor, Rif2p, cooperates
with Rif1p to regulate telomere length in *Saccharomyces cerevisiae*. *Genes Dev,*

11(6), 748-760.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=C
itation&list_uids=9087429
887
888

    Figure Legends

Figure 1. Rap1 binds poorly to *tlc1-tm* telomere sequences

(A) Exponentially growing cells of the indicated genotypes were diluted to an OD_{600} of 0.05

and monitored for 8 h. Three clones of each genotype were followed, and the average of the

measurements at each time point is plotted. Error bars are too small to be visualized.

(B) Telomere Southern blot analysis of *TLC1* and *tlc1-tm* cells (three clonal isolates per

genotype). Black arrowhead indicates a 1.8 kb DNA fragment generated from the BsmAI-

digestion of plasmid pYt103 (Shampay et al., 1984). The major terminal restriction fragment

is below the 1.8 kb control band.

(C) Southern blot analysis of the artificial VII-L telomere, with either wild-type or *tlc1-tm*

mutant sequence, using a probe to the adjacent *URA3* gene. Multiple clones were examined,

with each clone propagated for 1 to 4 passages (each passage corresponds to approximately

25 generations). A wild-type strain (lacking the artificial VII-L telomere) was used as a

control.

(D, F, G) ChIP-qPCR analysis of the association of (D) Protein A-tagged Rap1, (F) Myc-

tagged Cdc13, and (G) Myc-tagged Tbf1 to six Y′ telomeres, the VII-L telomere, or to the

non-telomeric *ARO1* locus. Untagged wild-type and *tlc1-tm* strains were used as controls.

The mean percentage of input ±SEM is shown (n=3, ** p<0.01).

(E) EMSA of Rap1 protein incubated with radiolabeled oligonucleotides with either wild-

type or *tlc1-tm* mutant telomeric sequence.

Figure 2. Telomere length regulation is disrupted in *tlc1-tm* cells

(A) Southern blot analysis of the artificial VII-L telomere using a probe to the adjacent *URA3*

gene. The telomere was seeded with either wild-type or *tlc1-tm* mutant sequence of the

indicated lengths in a strain expressing wild-type TLC1. Multiple clones of each strain were

examined. A wild-type strain (lacking the artificial VII-L telomere) was used as a control.

(B) In vivo extension of *tlc1-tm* telomeres was examined using the iSTEX assay. Telomere

VI-R was amplified and sequenced after the induction of wild-type telomerase. Each bar

represents an individual telomere. The black and red portions of each bar represent wild-type

and *tlc1-tm* sequence, respectively, that is identical in sequence and thus present before

telomerase induction. The length of the black/wild-type sequence is 48 bp. Sequence that is

divergent from the black and red sequence is shown in grey and green. Grey represents newly

added wild-type sequence after the induction of telomerase. Green represents divergent *tlc1-*

tm sequence, most likely a result of homologous recombination. Telomeres are sorted based

on the length of the undiverged (black plus red) sequence.

(C) Telomere VI-R sequences obtained from the iSTEX analysis in B were binned into

groups of 10 nt in size according to telomere length before telomerase induction. iSTEX data

for the extension of wild-type telomeres were taken from previous studies (Strecker et al.,

2017; Stinus et al., 2017) and included for comparison. Groups containing less than four

telomeres were excluded from this analysis. Frequency of extension and average telomere

length before telomerase induction were calculated and plotted for each group. Logarithmic

regression curves for each data set were also included in the plot. Telomeres shorter than 40

nt before telomerase induction, which are not efficiently recognized and extended by

telomerase (Strecker et al., 2017), were removed from the regression analysis.

Figure 3. Fast degradation of *tlc1-tm* telomeres in the absence of telomerase

(A) Senescence rates were measured by serial passaging of strains of the indicated genotypes,

derived from the sporulation of *est2Δ/EST2* and *est2Δ/EST2 tlc1-tm/tlc1-tm* diploids (top

panel) or *tlc1Δ/TLC1* and *tlc1Δ/tlc1-tm* diploids (bottom panel). Cell density was measured

every 24 h, followed by dilution to 1×10^5 cell/ml. Mean ±SEM of four independent isolates

per genotype is plotted.

(B) Telomere Southern blot analysis of samples obtained at the first time point of the

senescence assays in A. Black arrowhead indicates the 1.8 kb telomere sequence-containing

fragment loaded as control, as in Figure 1B.

Figure 4. Recruitment of Rif1 and Sir4, but not Rif2 nor Sir3, is significantly reduced at

tlc1-tm **telomeres**

(A) ChIP-qPCR analysis of the association of Myc-tagged Rif1, Rif2, Sir4, and Protein A-

tagged Sir3 to six Y′ telomeres, the VII-L telomere, or to the non-telomeric *ARO1* locus.

Untagged wild-type and *tlc1-tm* strains were used as controls.

(B) Total RNA was reverse transcribed and TERRA from specific telomeres (I-L and XV-L)

was analyzed by qPCR. TERRA values were normalized to ACT1 levels, and to the

respective wild type (*TLC1*).

(C) Tenfold serial dilutions of strains with the indicated genotypes were spotted on SC plates

without or with 5-FOA.

(D) The expression of the subtelomerically integrated *URA3* gene was measured in the

indicated yeast strains by RT-qPCR.

All data (except C) are shown as mean ±SEM (n=3, *** $p<10^{-3}$, **** $p<10^{-4}$).

Figure 5. Rif2 prevents degradation of *tlc1-tm* **telomeres**

(A-C) Telomere Southern blot analysis of strains of the indicated genotypes. Black

arrowhead indicates the 1.8 kb telomere sequence-containing fragment loaded as control, as

in Figure 1B.

(D) Colony sizes of haploid meiotic progeny derived from the sporulation of *rad52Δ/RAD52*

rif2Δ/RIF2 and *rad52Δ/RAD52 rif2Δ/RIF2 tlc1-tm/tlc1-tm* diploid strains were measured and

normalized to wild-type (top panel) or *tlc1-tm* (bottom panel). The number of colonies

analyzed is indicated in parenthesis. Error bars show SEM.

(E) Tenfold serial dilutions of strains with the indicated genotypes were spotted on YPD

plates and grown at 30ºC.

Figure 6. Rif2 protects *tlc1-tm* telomeres by inhibiting the MRX complex and acts in

parallel with the Yku complex

(A&C) ChIP-qPCR analysis of the association of Myc-tagged Mre11 and Yku70 to six Y′

telomeres, the VII-L telomere, or to the non-telomeric *ARO1* locus. Untagged wild-type and

tlc1-tm strains were used as controls. The mean percentage of input ±SEM is shown (n=3, **

$p < 0.01$).

(B&D) Telomere Southern blot analysis of strains of the indicated genotypes. Black

arrowhead indicates the 1.8 kb telomere sequence-containing fragment loaded as control, as

in Figure 1B. (B) Parental diploid 1 is *rad50Δ/RAD50 rif2Δ/RIF2*; parental diploid 2 is

rad50Δ/RAD50 rif2Δ/RIF2 tlc1-tm/tlc1-tm. The last six lanes are derived from cells grown at

22°C. (D) Parental diploid 1 is *exo1Δ/EXO1 yku70Δ/YKU70*; parental diploid 2 is

exo1Δ/EXO1 yku70Δ/YKU70 tlc1-tm/tlc1-tm.

(E) Colony sizes of haploid meiotic progeny derived from the sporulation of *rif2Δ/RIF2*

yku70Δ/YKU70 and *rif2Δ/RIF2 yku70Δ/YKU70 tlc1-tm/tlc1-tm* diploid strains were measured

and normalized to wild-type (top panel) or *tlc1-tm* (bottom panel). The number of colonies

analyzed is indicated in parenthesis. Error bars show SEM. Representative images of

dissected tetrads are shown on the right. Each column of colonies arose from a single tetrad.

Figure 7. Rif2 association to *tlc1-tm* telomeres requires the MRX complex

(A) Telomere Southern blot analysis of strains of the indicated genotypes. Black arrowhead indicates the 1.8 kb telomere sequence-containing fragment loaded as control, as in Figure 1B.

(B) Colony sizes of haploid meiotic progeny derived from the sporulation of a *rif2Δ/RIF2 yku70Δ/YKU70 tlc1-476A/TLC1* diploid strain were measured and normalized to wild-type. The number of colonies analyzed is indicated in parenthesis. Error bars show SEM. Representative images of dissected tetrads are shown on the right. Each column of colonies arose from a single tetrad.

(C) ChIP-qPCR analysis of the association of Myc-tagged Rif2 to a 300-bp ITS located on chromosome III containing either wild-type or *tlc1-tm* sequence. The mean percentage of input ±SEM is shown (n=3, *** $p<10^{-3}$).

(D) ChIP-qPCR analysis of the association of Myc-tagged Rif2 to six Y′ telomeres, the VII-L telomere, or to the non-telomeric *ARO1* locus in a *rad50Δ* background. Untagged strains were used as controls. The mean percentage of input ±SEM is shown (n=3, **** $p<10^{-4}$).

1004   **Supplemental Information**

1005

Supplementary Table S1. Yeast strains used in this study.

Supplementary Table S2. Wild-type and mutant sequences used to replace the VII-L

telomere.

Supplementary Table S3. qPCR primers.

Supplementary Figure 1. Tbf1 binding motifs adjacent to the artificial VII-L telomere.

Supplementary Figure 2. Rif2 protects *tlc1-tm* telomeres from degradation.

Supplementary Figure 3. Slightly differences in the telomere profile of isogenic *tlc1-tm*

strains are not due to Rad52-dependent homologous recombination.

1014

```
>adh4::URA3-VII-L-telomere
TTAATATTCATAGGCTTTCTTGATAATGGCAACCACTTGTTCTTTGGTGAATTGAACTGGGTTAGTCAAATGG
CAGGCATCATGCATGGCGTGTTCAGCCAAAATGTCAAAATCTTCGGTTTTAACACCTAAGTCTTTCAAGTTTC
TTGGAATGTTCATGGTTCTGTTTAAAACGTGCAAAGCCTTGATGGTTTCTTCTGGATCTTCTTGAGAAGCACC
GCAATGCAAGGCAATTTCACCTAATCTCTTCTTGGCCTTTGGACATTGCATGTTGGCCTCTTGAACATGAGGC
AACAAGACAGCGTTACAAACACCATGAGGCAAGTGGTAGAAACCACCAAGTTGATGAGCAAGGGCATGAACAT
AACCTAGAGAAGCATTGTTGAAAGCCATACCTGCCAAGTATTCTGCGTAACACATATCAGTTCTGGCCTTCTT
GTCTTTACCGTCTTTGTATGCGGCGACCAAGCTTTTCAATTCATCATTTTTTTTTTATTCTTTTTTTTGATTT
CGGTTTCCTTGAAATTTTTTTGATTCGGTAATCTCCGAACAGAAGGAAGAACGAAGGAAGGAGCACAGACTTA
GATTGGTATATATACGCATATGTAGTGTTGAAGAAACATGAAATTGCCCAGTATTCTTAACCCAACTGCACAG
AACAAAAACCTGCAGGAAACGAAGATAAATCATGTCGAAAGCTACATATAAGGAACGTGCTGCTACTCATCCT
AGTCCTGTTGCTGCCAAGCTATTTAATATCATGCACGAAAAGCAAACAAACTTGTGTGCTTCATTGGATGTTC
GTACCACCAAGGAATTACTGGAGTTAGTTGAAGCATTAGGTCCCAAAATTTGTTTACTAAAAACACATGTGGA
TATCTTGACTGATTTTTCCATGGAGGGCACAGTTAAGCCGCTAAAGGCATTATCCGCCAAGTACAATTTTTTA
CTCTTCGAAGACAGAAAATTTGCTGACATTGGTAATACAGTCAAATTGCAGTACTCTGCGGGTGTATACAGAA
TAGCAGAATGGGCAGACATTACGAATGCACACGGTGTGGTGGGCCCAGGTATTGTTAGCGGTTTGAAGCAGGC
GGCAGAAGAAGTAACAAAGGAACCTAGAGGCCTTTTGATGTTAGCAGAATTGTCATGCAAGGGCTCCCTATCT
ACTGGAGAATATACTAAGGGTACTGTTGACATTGCGAAGAGCGACAAAGATTTTGTTATCGGCTTTATTGCTC
AAAGAGACATGGGTGGAAGAGATGAAGGTTACGATTGGTTGATTATGACACCCGGTGTGGGTTTAGATGACAA
GGGAGACGCATTGGGTCAACAGTATAGAACCGTGGATGATGTGGTCTCTACAGGATCTGACATTATTATTGTT
GGAAGAGGACTATTTGCAAAGGGAAGGGATGCTAAGGTAGAGGGTGAACGTTACAGAAAAGCAGGCTGGGAAG
CATATTTGAGAAGATGCGGCCAGCAAAACTAAAAAACTGTATTATAAGTAAATGCATGTATACTAAACTCACA
AATTAGAGCTTCAATTTAATTATATCAGTTATTACCCTCGACGGATCCGGTGGTGTGGTG...
```

ADH4
URA3
Telomere
Tbf1 binding motif

Supplementary Fig. 1. Tbf1 binding motifs adjacent to the artificial VII-L telomere.
The sequence immediately adjacent to the artificial VII-L telomeric repeats is shown. The
truncated *ADH4* and full-length *URA3* sequences are highlighted, as well as sequences
conforming to the RCCCT Tbf1 consensus binding motif.

Supplementary Fig. 2. Rif2 protects *tlc1-tm* telomeres from degradation.
Southern blot analysis of the artificial VII-L telomere using a probe to the adjacent *URA3* gene in strains of the indicated genotypes.

A

B

Supplementary Fig. 3. Slightly differences in the telomere profile of isogenic *tlc1-tm* strains are not due to Rad52-dependent homologous recombination.

(A) Telomere Southern blot analysis of strains of the indicated genotypes (duplicate isolates per strain). Black arrowhead indicates the 1.8 kb telomere sequence-containing fragment loaded as control, as in Figure 1B.

(B) Colony sizes of haploid meiotic progeny derived from the sporulation of a *rad50Δ/RAD50 rad52Δ/RAD52 rif2Δ/RIF2 tlc1-tm/tlc1-tm* diploid strain were measured and normalized to *tlc1-tm*. The number of colonies analyzed is indicated in parenthesis. Error bars show SEM.