Mirror Neurons are Modulated by Grip Force and Reward Expectation in the Sensorimotor

Cortices (S1, M1, PMd, PMv)

Md Moin Uddin Atique[1] and Joseph Thachil Francis[1-2*]

Number of Pages: 42

Number of Figures: 10

Number of Tables: 0

Number of words for Abstract: 204

Number of words for Introduction: 817

Number of words for Discussion: 1296

Conflict of interest statement: None

Acknowledgements: We thank Matthew J. Perry for helping with the manuscript. Research was

supported by NIH 1R01NS092894-01, NSF IIS-17 1527558, DARPA REPAIR Project N66001-10-

C-2008.

[1]Department of Biomedical Engineering, Cullen College of Engineering, The University of Houston,

Houston, TX 77204, United States;

[2] Department of Electrical and Computer Engineering, Cullen College of Engineering, The

University of Houston, Houston, TX 77204, United States;

* Corresponding author Contact- jtfrancis@uh.edu, joey199us@gmail.com

Abstract

Mirror Neurons (MNs) respond similarly when primates make, or observe, grasping movements. Recent work indicates that reward expectation influences rostral M1 (rM1) during manual, observational, and Brain Machine Interface (BMI) reaching movements. Previous work showed MNs are modulated by subjective value. Here we expand on the above work utilizing two non-human primates (NHPs), one male *Macaca* Radiata (NHP S) and one female *Macaca* Mulatta (NHP P), that were trained to perform a cued reward level isometric grip-force task, where the NHPs had to apply visually cued grip-force to move and transport a virtual object. We found a population of (S1 area 1-2, rM1, PMd, PMv) units that significantly represented grip-force during manual and observational trials. We found the neural representation of visually cued force was similar during observational trials and manual trials for the same units, however, the representation was weaker during observational trials. Comparing changes in neural time lags between manual and observational tasks indicated that a subpopulation fit the standard MN definition of observational neural activity lagging the visual information. Neural activity in (S1 areas 1-2, rM1, PMd, PMv) significantly represented force and reward expectation. In summary, we present results indicating that sensorimotor cortices have MNs for visually cued force and value.

Introduction

Mirror neuron (MN) activity has been observed while studying kinematic behaviors, [1–4], and through indirect measures in human motor cortex [5]. In order to ask questions about force information at the single unit level, we utilized an isometric grip-force paradigm (Fig.1), and cued reward level [6,7], while modulating both cued grip-force and reward level in the current work. We utilized this grip-force paradigm initially for brain machine interfacing research [8], and subsequently included observation only trials to ask questions about mirror neuron activity within the sensorimotor regions when kinematics are not actively being controlled, but rather force output is the controlled variable.

A better understanding of how reward expectation influences the primary- and pre- sensorimotor

cortices has practical implications towards the production of stable and autonomously updating

brain machine interfaces (BMIs) [6,7,9–12]. Neural modulation related to reward has been well

characterized within many brain structures [13–15], and is known to occur at many levels, such as

single units and local field potentials (LFP) within rM1 [6]. The neural response to both reward and

conditioned stimuli (CS) that predict reward have been demonstrated in a multitude of brain regions

[7,16–22]. It has recently been shown that reward expectation changes directional and force related

tuning functions within the primary sensorimotor cortices during reaching movements and BMI

reaching movements [8,23]. The above studies indicate the need to conduct more research on the

influence reward has on these brain regions, and how such variables may modulate MNs. The

influence of reward on the caudal primary somatosensory cortex (cS1, area1 and 2) was reported

in conference proceedings [24–26]. While these experiments demonstrated our ability to classify

reward expectation from S1 they did not provide insight on its influence of grip-force during action

and action observation as we do here. A similar reward signal has been demonstrated in primary

somatosensory cortex utilizing fMRI, where a stronger BOLD response was recorded for higher

reward delivery [27], the investigation of reward-correlated signals within S1 at the single unit level

has not been fully developed.

Our previous work focused on identifying significant units related to reward and/or force within

single blocks of data [8] during manual trials and BMI control. To expand on these findings, we

thought it important to investigate the cause behind each unit's modulated activity across multiple

blocks, during both manual [8], and observational tasks. We hypothesized modulation responses

during observation might be explained by the presence of mirror neural activity when NHPs

observed the task, received reward, or possibly both. To this end, our goal was to detect units that

encoded the following: 1) Motor actions that are the physical application of force; 2) Cued "force"

during visual observation; 3) units that encoded both 1, and 2. 4) Determine which of these (1-3)

were further modulated by reward level, cued or uncued. We focused on "grasping" movements

with cued isometric grip-force control, to expand beyond our previous work on reaching

movements [6,28]. Below we demonstrate our ability to decode actual and observed grip-force neural

responses, during manual and observational trials, as well as reward's influence on the neural

population during reward cued and uncued trials. Others have described activity of MNs in rM1

during observation of movement [3,28–31], but little has been reported on the activity in S1 during

observational tasks, or any of the 4 regions as it pertains to isometric grip-force control while

modulating reward level as we do here. There are some studies on human and NHP that showed

evidence of MN activation [32–34] and grip force related modulation [35,36] is present in somatosensory

areas which implies the possibility of MN related activity due to grip force in cS1 cortex. Generally,

our paper presents findings of MN responses related to varying levels of grip-force (Fig.6) and

reward's influence on grip-force MN activity (Fig.8-10) within cS1, rM1, PMd and PMv.

Methods

All NHP manipulations described in this work were approved by the Institutional Animal Care and

Use Committee of the State University of New York at Downstate Medical Center and conformed

to National Institutes of Health (NIH) and United States Department of Agriculture (USDA) animal

care and use guidelines. In addition, this work complies with the ARRIVE guidelines. Two non-

human primates (NHPs), one 9.0kg male *Macaca* Radiata (NHP S) and one 5.0kg female *Macaca*

Mulatta (NHP P), were trained to perform a behavioral grip-force task using their right hand and

subsequently implanted with 96-channel electrode arrays in cS1, rM1, PMd and PMv.

<div align="center">**Behavioral Task**</div>

Figure 1: Flow diagram illustrating a single trial. The reward cue (green square, R1) was shown

for R1 trials in cued blocks only. Absence of the green square indicated a non-rewarding (R0) trial

during cued blocks of the task. The robotic arm started in a resting position for all trials (left-most

image). During uncued blocks, there was no cue provided during the "Reward cue" period for any

trials. Please note, under experimental conditions with NHPs the background of the grid and free

space around the robotic arm and object were black. For the reader's benefit and visual clarity,

both areas have been changed to white here. Blue rectangles are grip-force targets and were seen

for the duration of a trial. Red rectangle shows visual force feedback of the NHPs grip-force output

initially but is no longer shown once the NHP has reached the acceptable force range (blue

rectangles).

We describe the sequence for one complete trial of the manual reward-cued version of the task

seen in Fig.1. First, the virtual robot moves to the start position, which is always to the far left of

the task space. Second, cue scene, a green square moved across the top of the screen for 0.5s

from the left to indicate a rewarding (R1) trial, absence of this green square indicated a non-

rewarding (R0) trial. Third, the virtual robot autonomously moved its arm toward the cylindrical

object in the reach scene. Fourth, during the grasp scene of manual trials, the NHPs had to apply

isometric grip-force using a stationary handle that contained a force transducer. The amount of

force required was indicated by two blue rectangles, where the inner vertical edge of each blue

rectangle indicated minimum force, and the outer edges indicated maximum force. The minimum

threshold of force for NHP S was randomly chosen from either 150 or 200 and for NHP P it was

100 or 150. If the force output went below that minimum value the trial was considered a failure.

The upper threshold was determined by randomly adding either 300 or 500 to the minimum for

NHP S and adding randomly either 250 or 350 to the minimum for NHP P. Therefore, the ranges

used for NHP S were randomly selected from the following, 150-450, 150-650, 200-500, 200-700,

and for NHP P the ranges were, 100-350, 100-450, 150-400, 150-500. If the output force of the

NHP was above these max levels the trial was considered a failure. The peak grip force values

across the manual trials, R1 and R0, in cued and uncued blocks are provided in supplementary

figure S7. When the NHP's applied force was within the acceptable tolerance range, the red force

rectangle's edges would also be within the blue rectangles. When this occurred, the visual force

feedback (red rectangle) was removed, and the robot hand would grasp the object. We utilized this

method so that the NHPs could not simply use visual feedback to perform the task, but rather had

to learn to control their force output based on the visual force targets and somatosensory feedback

for manual trials. Thus, they had to learn to produce a given grip-force, and this may have led to

the positive results during observational tasks, as they most likely had built an internal

representation of the force cues and their production of such force as well as the expected

somatosensory feedback. During the transport scene, the robot arm autonomously moved the

object to a predetermined target location, indicated by a pink circle. If the NHPs applied the

appropriate grip-force during object grasp and transport, and then released when the object was

touching the virtual ground at the pink target location, the trial was successful. Feedback was

provided to NHPs for successful placement when the blue force squares turned green, and the

robotic arm released the cylinder during the release scene. If they did not apply proper force during

initial grasp, failed to maintain an acceptable force range at any time during the grasp or transport

scenes, or released their grip early or late, the trial was considered a failure.

There were four types of task blocks, which were experienced by each NHP in the following order:

first, manually performed tasks with the presence of a reward level conditioned stimulus (CS)

(manual cued); second, manually performed tasks without a reward level CS (manual uncued);

third, observational task with the presence of a reward level CS (observational cued); fourth,

observational task without a CS (observational uncued). All trials (R0 and R1) during uncued

blocks lacked a visual reward cue so the NHP had no indication of trial value until the post result

period, and thus, no explicitly cued expectation for reward outcome as the trial sequence was

randomized with no clear autocorrelation (see supplementary Fig.S1). The time limit to complete

a successful trial was within 10s. The NHPs had to repeat failed trials under the same reward

conditions until successful, thus giving incentive to perform non-rewarding trials. Without this

added stipulation the NHPs would choose to fail R0 trials to move onto a possible R1 trial,

indicating the NHPs clearly understood the cue reward values. Juice rewards were delivered via a

system-controlled solenoid driven by task logic (Crist instruments). All elements of the grip-force

task were developed in Linux using robot operating system (ROS) [37]. ROS and Python controlled

the task logic, outputs to the reward delivery system, and provided timestamp synchronization with

external systems to simultaneously and accurately record task state, reward delivery, and neural

data.

During observational blocks NHPs were visually monitored by researchers in real-time via cameras

during all sessions to make sure NHPs remained attentive and focused on the projection screen,

especially during observational trials. During observational trials NHPs did not have access to the

force transducer handle, and their arms were blocked behind a plexiglass box meant to keep the

NHPs hands away from the trainers (BKIN's Arms-Free restraint chair). Additionally, NHPs were

trained, and performed all experiments in a dark, quiet, and distraction-free isolation chamber to

encourage their attention remained on the large projection screen. Each NHPs' rear-mounted

cranial head post was affixed to the BKIN primate chair to restrict head movement for neural

recordings. The virtual environment was projected onto a vertical screen in the animal's visual

field. The visual projection of the virtual robotic system was approximately the same size as the

real WAM robotic arm ~1m reach (WAM Barrett). Recording sessions were broken into blocks of

trials that averaged ten minutes of randomized R1 and R0 trials.

Chronic Implantation

We performed implantation procedures as described in-depth in a previous methods paper [38], but

give a summary here. Following training, when both NHPs achieved greater than 80% success

rate on all trials in a block, the animals were implanted with chronic electrode arrays consisting of

a 10 by 10 array of 1.5mm electrodes of which 96 were active for all regions (rM1, PMd, PMv)

except cS1, which was implanted with electrodes of 1mm length. Spacing between individual

electrodes within the arrays were 400μm (Utah array, Blackrock Microsystems) (Fig.2). All surgical

procedures were conducted using aseptic technique. NHPs were initially anesthetized with

Ketamine, followed by isoflurane and a continuous infusion of fentanyl. The NHP was then placed

into a stereotactic frame before the surgical site was shaved and cleaned. An incision was made

along the skull to expose desired implant locations. The craniotomy window was large enough to

accommodate implant locations while leaving enough margin between the dural flap and skull.

The dural flap was kept under tension using stay sutures until electrode arrays were implanted and

the site was ready to close. We performed intraoperative probing of cS1 to ensure implantation

within the hand region, by using a four shank, 32-channel silicon microelectrode array [39] within the

post-central gyrus as determined by stereotactic coordinates. The NHP's contralateral hand was

continuously stimulated by touch to assess cS1 hand region boundaries. Neural responses were

amplified and sent to an audio speaker to verify stimulation areas. Utah arrays were then

chronically implanted in cS1's hand region, in rM1 directly reflected across the central sulcus rostral

from cS1 for NHP S. However, NHP P had large blood vessels that interfered, and we had to

implant rM1 more lateral than in NHP S as seen in Fig.2. In addition, as seen in Fig.2, NHP P's

PMd implant was slightly more medial than NHP P, and NHP

195 S's PMv was slightly more lateral than NHP P. In both NHPs our PMv arrays had fewer clear single

196 units than the other arrays. We cannot be certain but believe this was due to the PMv wire bundles

197 exiting the craniotomy closer to the array and at a stiffer region of the wire than the other arrays.

198 The craniotomy was replaced according to previously described methods [38].

200 **Figure 2**: Position of four Utah arrays in relation to the central sulcus for NHP S (left) and P (right).

201 The four arrays were implanted in cS1, rM1, PMd and PMv cortices. The yellow line indicates the

202 Central Sulcus. Note in NHP P we had to implant the rM1 array more lateral than in NHP S due to

203 a large set of blood vessel running through that region. Likewise, PMd was implanted more medial

204 in NHP P for this same reason as compared to NHP S. The color map on each array is for the

205 number of MN (mirror neuron) units recorded in each location, we did not have the electrode map

206 for NHP P PMv. The white arrow inside the color map represents the mean direction of MN position

207 with respect to the center of the array. A Rayleigh test (p<0.01) showed the distribution of MNs

208 around the center of the array is non-uniform only for M1 in both NHP.

209 **Neural Recording**

210 Neural recordings were performed using three synchronized multi-channel acquisition processors

211 (Plexon, Dallas, TX), each having 128-spike waveform recording channels and 32 analogue

212 channels used for simultaneous recording of local field potentials [6,12]. Single unit recordings were

amplified and retained using waveform voltage thresholds. Thresholds were set using an auto-

scale (Plexon recording software) feature, followed by manual adjustments that eliminated noise

on channels prior to recording. Robot operating system [37] and Python programs controlled task

logic and embedded timestamps into neural recordings using a common clock. The common clock

was maintained by a microprocessor that delivered a 2kHz pulse to keep task logic and neural

data synchronized. Initially, unit waveforms were automatically clustered using a k-means

algorithm in principal component space [40,41]. Afterward, we used Plexon's offline sorter software to

adjust clusters to remove noise and artifacts.

Significance Related to Reward

We only considered successfully completed trials for all analyses and results presented in this

work. We analyzed single unit activity from four types of task blocks for each NHP, cued or uncued,

and these were either manual or observational. For trials that contained a visual reward cue, the

post-cue (0-500ms) analysis window began immediately after the green visual cue came to rest.

In uncued trials, the "post-cue" period (0-500ms) began when the robotic arm returned to its rest

position from the previous trial. We defined the "post-result" period (0-500ms) as the time after the

cylindrical object was successfully placed at the target location. We analyzed spike activity in the

post-cue and post-result periods separately for cued and uncued blocks to identify units significant

for reward modulation. We binned the spike activity into non-overlapping bins of 100ms covering

500ms for both the post-cue and post-result periods. We collected the p-values from t-test for

Spearman Rank correlation and the test was done between the binned spike rate and

corresponding reward levels (0 for R0 and 1 for R1). For reward levels, 0 is used for R0 and 1 is

for R1. We then extracted units that were significant (t-test, $p < 0.05$) for reward expectation (post-

cue activity, R1 vs. R0 trials) and reward result (post-result activity, R1 (after reward delivery) vs.

R0 (after successful trial with no reward)). The p-values from the test are adjusted for multiple

comparisons using the false discovery rate (FDR) procedure by Benjamini and Hochberg (BH

Method) (Benjamini et al. 1995) for number of times t-test was applied which is equal to the number

of units for each case, which is given in supplementary (Fig.S2). Finally, we compared spike rate

activity between cued and uncued blocks to gain better insight as to how reward feedback

information was affected by the presence or absence of a reward cue that is reward expectation

based on explicit environmental cues. Our interest in reward expectation stems from our work

towards autonomously updating BMIs, and thus our use of this paradigm.

Grip-force Trajectory Prediction

We applied linear regression in two steps to identify units that showed significant prediction of grip-

force from cS1, rM1, PMd and PMv. In these two steps, the first step identified all units that were

significantly related to grip-force, which was used below in the sections labeled "*Grip-force Tuning*

Curve Analysis" and "*Identifying Observation Modulated Neurons*". The second step was applied

to sort out a selective number of units, from the units collected in the first step, for force prediction

to reduce the number of predictors and avoid computational complexity. Unit activity from the

aforementioned cortices were analyzed along with force profiles for both NHPs during manual and

observation task blocks, see supplementary material for data not shown in the main text. During

some trials, initially the NHP's applied more grip-force than necessary and in an effort to meet the

target value, reduced their grip-force drastically. This resulted in an overcorrection where the NHP

had to apply more force to meet the requirement and resulted in force profiles with multiple peaks.

We manually inspected and pruned all force profiles that contained multiple peaks due to highly

variable or corrective grip-forces applied by NHPs. We defined force onset as the point where force

values increased from zero to positive value of 50 au and force offset when values returned below

50 au, as the NHPs only grasped the force transducer during the grasp scene, and a trial was

considered successful when their output force < 50 au, which generally meant they let go of the

handle. Force data was collected from 0.5s prior to force onset through 0.5s after force offset for

all trials. The collected force data was smoothed using a Gaussian kernel (100ms wide). During

observational trials the visually cued "force" targets, and "force" output profiles were trapezoidal

and were inferred from the trapezoidal velocity grasping motion of the hand. Therefore,

smoothening gave the "force" profile a gaussian shape like the force data on manual blocks. From

the force data collected in each trial, we evaluated neural spike activity 0.5s prior to, and following

each force sample value. This neural activity was further placed into ten non-overlapping time bins

centered on each force value (100ms bins, covering 0.5s pre- and 0.5s post-force sample value)

to determine a unit's significance to grip-force, so for each unit there were 10-time bins used as

predictors. We performed this separately on each unit from all four cortices. As we smoothed the

unit raw data using a Gaussian kernel (100ms wide), we wanted to find a generalized kernel width

that would be used for both cued and uncued data blocks of the same kind, manual vs.

observational. The force prediction accuracy for different kernel bin widths from 10ms to 250ms

were determined for all data. We manually selected the above value (100ms) for the kernel width

so that the predictions were close to the peak prediction values for all blocks. See Fig.S3 for

differences between analysis on the smoothed data, vs. unsmoothed. The collected binned data

was square root transformed to achieve a more normalized distribution. To make sure this square

root transformation is not affecting the prediction results significantly a comparison of grip force

prediction accuracy is given before and after the transformation is applied in supplementary Fig.S4.

Data recorded from each NHP was split into a fit (90%) and test (10%) set to check goodness of

fit and verify predictive performance of the final linear regression model (equation 2). However,

before testing the final linear regression model, we extracted significant units related to force in a

two-step process. To do this, we subdivided the 90% fit set into 80% (fit) and 10% (validate) data

subsets, see the supplementary section for a cartoon of this procedure Fig.S5.

$$y_f = a\mathbf{X}_{fr}^{n_i} + b \quad (eq.1)$$

Equation 1 shows the linear model used, where $\mathbf{y}_f$ is the vector of force values, $\mathbf{X}_{fr}^{n_i}$ is the

binned spike rate matrix for neuron n_i, while $\mathbf{a}$ and b are coefficients fit to the data. In the first

step, we applied linear regression using equation 1 with the 80% subset data for the fit and

collected the F-statistics from the analysis of variance (ANOVA) for the model. Units were sorted

according to their p-value from the above F-test and only those with a significant fit ($p < 0.05$) were

considered in the second step of the process. For the second step, we took significant force units,

starting with the most significant (lowest p-value), and used it to test prediction of force values in

the remaining 10% validation subset. We calculated the R-squared value after each unit was added

to the model and if the R-squared value increased (improved prediction results), the unit was kept.

Remaining units that did not improve the model's force prediction were pruned. After the two-step

process, the subset of significant force units was utilized in the final linear regression model (eq.

2). This subset of units was used with updated coefficients fit to the original 90% validation dataset.

Finally, we used the held out 10% test set to determine prediction and validate accuracy of the

linear regression model (eq. 2) by comparing them against actual recorded force values.

$$\mathbf{y}_f = \boldsymbol{\alpha}\mathbf{X}_{fr} + \boldsymbol{\beta} \quad (2)$$

The regression model used for final grip-force prediction from neural activity is shown in equation

2. Here $\mathbf{y}_f$ denotes grip-force and $\mathbf{X}_{fr}$ represents binned firing rates for the population of units

being used, $\boldsymbol{\alpha}$ and $\boldsymbol{\beta}$ are model coefficients fit to the data.

Grip-force Tuning Curve Analysis

Our previous research concerning force tuning curves in rM1 showed a significant difference

between R1 and R0 trials [8]. We applied the same analysis to our cS1, rM1, PMd and PMv data to

determine if neural activity led to different force tuning curves when taken from R1 vs. R0 trials. As

in Zhao et al. 2018 we utilized analysis of covariance (ANCOVA). We analyzed each unit previously

identified as significant for force to measure whether the slopes of the tuning curves between R1

and R0 were significantly different (F-test, $p < 0.01$). The p-values from the ANCOVA test are

adjusted for multiple comparisons using the false discovery rate (FDR) procedure by Benjamini

and Hochberg (BH Method) (Benjamini et al. 1995) for number of population on which the test was

applied. Significant units identified by ANCOVA were considered to have a force representation

that was modulated by reward expectation, and units that also passed the BH method are stated

explicitly.

Identifying Observation Modulated Neurons

We started by identifying and tracking single unit activity across multiple blocks of recorded data.

We compared single unit activity between reward-cued manual and observational blocks, and

again between reward-uncued manual and observational blocks for each NHP and cortices (CS1,

rM1, PMd and PMv). We tracked single unit activity from the two manual and two observation

blocks performed in a single day for each NHP. If a unit on a channel retained the same waveform

over all 4 recorded blocks, we considered it the same unit. In addition, we verified this single unit

activity by checking the correlation coefficient between the waveform shapes across all blocks,

where a high correlation (> 0.98) indicated the same unit. We also confirmed the consistency of

single unit activity across blocks by checking the first two principal components using principal

component analysis (PCA). Performing these checks allowed us to track single unit activity with

significant correlation to reward, force, or both during manual and observation task during cued

and uncued blocks towards identifying putative mirror neural activity responding to force and

reward modulation.

Results

The results section is structured as follows: 1) We start by presenting raw data from manual and

observational versions of our isometric grip-force task showing peri-event-time-histograms

(PETHs) and rasters of sample units, indicating the modulation of single units by both cued grip-

force and cued reward level during manual and observational trials in the caudal S1 areas 1-2

(cS1), rostral M1 (rM1), PMd and PMv, for NHP S and P, Fig.3 and Fig.4 respectively. 2) We

present population results from linear regression focusing on cued grip-force, showing that

subpopulations of these brain regions encode grip-force trajectories (Fig.5) even during

observational trials. 3) In Figure 6 we test for temporal shifts between the neural representation of

force, between manual and observational versions of the tasks, to determine if the same units are

acting in response to the visual input, classical MNs, or predicting such input, as expected by

mental simulation of the predicted visual stimulus, which could in turn be either mental simulation

of the expected movement [2], or MNs responding in an anticipatory way due to the familiarity with

the task [43]. 4) Subsequently, we give information on units that are modulated by both cued grip-

force and cued reward level during the force output period of the manual and observational tasks

for the same single units that is putative MNs for force modulated by reward. To be clear, the term

force, when used for the observational trials, indicates the force that would be expected from the

NHP if the NHP were performing the task manually based on the visual force target cues still being

presented during the observational task.

In Fig.3 and 4 (b-c) we present raster plots for example units with all trials aligned to the onset or

offset of the grip-force. The mean force profile (indicated with green lines on the raster plots) shows

force onset and offset, plotted with arbitrary units for R0 (blue raster) and R1 (red raster) trials. The

mean duration that NHP S applied grip-force was 1.4 ± 0.4s (cued), 1.42 ±0.42s (uncued) during

manual blocks. During observational blocks the "force" times where 1.12 ±0.26s (cued), 1.16

±0.27s (uncued). For NHP P, the mean force duration was 1.2 ±0.28s (cued), 1.45 ±0.29s (uncued)

during manual blocks, and 1.12 ±0.18s (cued), 1.21 ±0.23s (uncued) during observational blocks,

see Fig.S6 – S8 for more on force level, duration and reaction times. NHP S achieved success

rates of 77% (cued manual) and 82% (uncued manual) during the two manual-task blocks, while

NHP P achieved rates of 58% (cued manual) and 72% (uncued manual). The number of R0 and

R1 trials recorded on each individual block type is given in supplementary Fig.S9. The R0 and R1

mean force profiles are plotted with the same scale for comparison. All units shown in both Fig.3

(4 units) and 4 (4 units), where significant (F-test, BH corrected for p=0.05 for the number of units

within the given brain region) for grip-force. For each NHP and cortices the number of units

recorded is given in supplementary (Fig. S2), min ~50 units and max ~170 units. In Fig.3 ax (x=1,

2, 3 and, 4) show the mean spike waveform with standard error (shaded) for a single unit (top) and

its PCA space (bottom) for all 4 blocks, manual cued (MC), manual uncued (MU), observational

uncued (OU) and observational cued (OC), which were all recorded on the same day and in the

aforementioned order for each NHP. Each raster subplot (b-e) shows activity for force onset (left)

and force offset (right) periods.

Units in Figs.3 - 4 were chosen as they represented the variety of responses we saw in the

population. We utilized 100ms bins during the post-force onset spike activity from the point when

force sensor values crossed above 50 (a.u.) and pre-force offset activity until the reading went

below 50 (a.u.). Fifty was the same value used for the task logic and was chosen based on

experience with these NHPs performing this task in order to minimize false starts without missing

movements. We collected p-values from t-tests on Spearman Rank correlation between the binned

(100ms) spike rate and corresponding reward levels (1 for R0 and 2 for R1) indicating significance

with an asterisk (*). For example, in Fig.3.b1-c1 we see an cS1 unit that decreases its activity at

or just after the onset of "force" and is suppressed around the force offset period. Our hypothesis

is this unit along with other example units that are showing suppression of activity is due to these

units possibly being connected to extensor muscle groups of forearm and thus showing inhibitory

spike activity during gripping action, see Fig.S17 for support of this. In addition, the unit in Fig.3.b1-

c1 shows modulation by reward. Similarly, in Fig.3.b2-c2, we see an rM1 unit with a similar

response to the cS1 unit, where the rM1 unit is suppressed at force onset and activated at force

offset with some reward modulation as well. In Fig.3.b3-c3 a PMd unit is shown that is suppressed

before force onset and has its peak activation pre-offset. Monkey P data was not as responsive to

reward activity utilizing the spearman rank test as Monkey S. The Fig.3 responses shown for cS1, rM1 and PMd resemble extensor patters of activation while the PMv response shown resembles a more flexor typical response [44,45]. In Fig.4 we have chosen units that show some other responses seen in the population for both NHPs. The observational response to "force" was weaker compared to the actual force during manual trials and did not always align with the force onset and offset times as precisely. For additional raster plots on example units during grip force observation please see supplementary Fig.S10. All units seen in Figs.3-4 had significant fits to grip-force (F-test, p <0.05). This F-test was performed on the regression model (equation 1) to test whether it was a better fit than a degenerate model, which consisted of only a constant term. Figs. 3-4 show a total of 8 example units; 1 from each cortical region (rM1, cS1, PMd and PMv) for each NHP, P and S, from reward cued, and uncued blocks.

Mirror neuron activity in cS1, rM1, PMd and PMv (NHP S)

Figure 3: Raster plots of example units from NHP S during the manual and observational blocks. The plot numbers, 1-4, indicate the brain regions (1-4 for cS1, rM1, PMd and PMv respectively). The letters, a-c, after the numbers, 1-4, are associated with (a) single unit waveform for the example unit, (b) force onset plots and (c) for force offset plots. For (b) and (c) there are four plots

for manual cued task (MC), observational cued task (OC), manual uncued task (MU), and

observational uncued task (OU) sequentially. The post-force onset and pre-force offset time is set

as 60% of the mean force length of all trials for that task. On each subplot (b, c), the x-axis

represents time in seconds, for raster plots the y-axis represents trial number, for spike rate plots

(bottom of subplots) y-axis is spike rate in 'Hz'. The dashed horizontal black line on each plot

divides the R1 trials from R0 trials. Below the raster plots, solid red and blue lines indicate mean

spike rate (Hz) for R1 and R0, respectively. An asterisk (*) indicates post force onset or pre-force

offset spike activity is significantly (t-test, $p<0.05$, $p<0.01$ and $p<0.001$ denoted by *, ** and ***

respectively) different between R0 and R1 trials.

Figure 4: Raster plots of example units from NHP P during the manual and observational blocks.

The plot numbers, 1-4, indicate the brain regions (1-4 for cS1, rM1, PMd and PMv respectively).

The letters, a-c, after the numbers, 1-4, are associated with the (a) single unit waveform for an

example unit, (b) force onset plots and (c) force offset plots. For (b) and (c) there are four plots for,

manual cued task (MC), observational cued task (OC), manual uncued task (MU), and

observational uncued task (OU) sequentially. The post-force onset and pre-force offset time is set

as 60% of the mean force length of all trials for that task. On each subplot (b, c), the x-axis

represents time in seconds, for raster plots the y-axis represents trial number, for spike rate plots

(bottom of subplots) y-axis is spike rate in 'Hz'. The dashed horizontal black line on each plot

divides the R1 trials from R0 trials. Below the raster plots, solid red and blue lines indicate mean

spike rate (Hz) for R1 and R0, respectively. An asterisk (*) indicates post force onset or pre-force

offset spike activity is significantly (t-test, $p<0.05$, $p<0.01$ denoted by * and ** respectively) different

between R0 and R1 trials.

Grip-force Decoding During Manual and Observational Trials

As described in the methods section and further in the supplementary section (Fig.S5), data blocks

were analyzed using a two-step process. We identified significant units related to force using the

simple linear model eq.1, $\boldsymbol{y_f} = \boldsymbol{a}\mathbf{X}_{fr}^{n_i} + b$, where $\boldsymbol{y_f}$ is the vector of grip-force trajectory, $\boldsymbol{a}$ is the

vector of regression parameters that multiply unit $n_i's$ firing rate, where we used the associated F-

statistic with a ($p < 0.05$) for significance determination. Note, we utilized neural data around a

given grip-force timepoint, from -0.5s to 0.5s in 100ms bins for this regression, thus 10 bins. We

extracted single units with activity that improved force decoding. Model 1 variables $\boldsymbol{a}$ (10 per unit)

and b were fit using least-squares estimates from 90% of the recorded data (see Fig.S5 for data

split cartoon). The remaining 10% of the data was used as a test set to verify the performance of

force decoding. The R-squared values for decoding on the test set of force are shown below. We

found high prediction accuracies for both NHPs. The results shown in Fig.5 are force decoding

prediction from cS1, rM1, PMd and PMv cortices for cued manual blocks, top two rows, with the

corresponding observational blocks seen in the bottom two rows. Note intertrial intervals have

been clipped out for presentation purposes.

483   **Neural Decoding of Force (Manual and Observational)**

Figure 5: Force decoding (Red lines) from subpopulations of cS1, rM1, PMd, and PMv cortical

recordings taken from manually performed blocks top two rows, and observational blocks bottom

two rows. Linear regression (eq. 2) was used to predict the force from a subpopulation (see

methods). The black dots on each plot indicate that force profile was taken from an R0 trial, while

all others were R1 trials. Plots *a, b, c* and d show results for NHP S, for cS1, rM1, PMd and PMv

cortices respectively. Plots e*, f, g,* and h show similar results for NHP P. The top of each figure

shows R-squared values (R^2) using the 10% test set and the number (n) of significant units used

for regression to predict force. See table S2 for full stats on the regression models.

It is clear the manual tasks generally had higher levels of prediction as compared to the

observational versions. We detected the units that were consistently present in all four data blocks

(two manual and two observational) and those significant (F-test, BH adjusted for p=0.05 and

number of units in the population) for grip-force fit in all data blocks. For NHP S we found 31 units

in cS1, 42 in rM1, 46 in PMd and 29 in PMv that were consistently present in all four data blocks.

For NHP P the numbers were 107 in cS1, 95 in rM1, 79 in PMd and 49 in PMv. Below in Fig. 6 we

show the total percentage of these units from each region that showed significant linear regression

model fit under both the manual and observational versions of the task that are the putative MNs.

We used the peak value from the absolute correlation coefficients between grip force and each of

the 10 bins of spike rate (100ms from -0.5ms pre grip force spike rate to 0.5s post, described "Grip

force trajectory prediction" section inside method) to detect inhibitory or excitatory activity. We

considered a unit's representation excitatory if the actual value of that peak absolute correlation

was positive otherwise it was considered inhibitory. We detected congruent units among the

significant units that showed similar activity in all four data blocks of either inhibitory (blue) or

excitatory (red) spike rate in relation to grip-force. Our hypothesis is the units that are showing

inhibitory activity during grip force application are connected to the extensor muscle groups of the

forearm which relaxes during gripping activity and show evidence for this in Fig.S17. Incongruent

units showed opposing behavior between manual and observational tasks, such as excitatory

during manual and inhibitory during observational, purple, or inhibitory during manual and

excitatory during observational (green) in Fig.6. Other (gray) units didn't follow any simple pattern

during all four data blocks as seen in Figs.S11-S12. The number of units for each possible

combination of inhibitory or excitatory spike activity during grip force for four data blocks are given

in supplementary (Fig.S11 for NHP S and Fig.S12 for NHP P) section. Monkey S showed what

one might expect PMv>PMd>rM1>cS1, whereas NHP P's data was not in this clear expected order

and could be due to the placement of the electrode arrays due to the vasculature-imposed

restrictions as seen in Fig.2. However, in both NHPs the largest single group of MNs appear to be

congruent and inhibitory for our task (blue bar sections).

% Units with Force-Mirror-Neuron Properties

Figure 6: Percentage of well isolated MN single units from cS1, rM1, PMd and PMv that showed

significant (F-test, BH post-hoc adjusted for number of units in the population) fit to grip-force

during all four blocks of data (two manual and two observational) for each NHP. The different color

bars represent congruent inhibitory (blue), excitatory (red), incongruent inhibitory to excitatory

(green), incongruent excitatory to inhibitory (purple), and other (gray) unit responses. The

percentage (y-axis) shows how many units (# at top) are significant for force from units present in

all four data blocks. See Fig.S11-S12 for a breakdown of units by trial type.

Distribution of Reactive vs. Predictive Mirror Neurons

The analysis in this section was conducted in order to determine if there was an obvious shift in

time lag between the neural correlates of force between manual trials and observational trials. One

would expect the neural data responsible for force production, or imagining force production, to

lead the force output, whereas classical MN activity would lag the viewed "force" output. However,

this may not be the case for predictable movements, such as used in our task, as described by

others [43], where MNs can still lead the observed task. In Fig.7 units that were significant for force

(F-test, $p < 0.05$, eq.1) on all block types were plotted in the time bin where their correlation

coefficient with force was maximum in absolute value, either negative or positive. We asked if

these distributions significantly (signed Rank test, $p < 0.01$) deviated from zero between the manual

and the observational versions of both the cued and uncued tasks. There were no significant shifts

in the histograms for NHP P. NHP S showed one task with significant shifts for PMd during the

cued tasks. For each plot the zero-time shift shows the probability of a unit not shifting in time

between manual and observational. The positive differences are the classical MN response where

observational neural activity lags the visual information, whereas the negative time shifts represent

the probability of the observational neural responses being earlier, or more "predictive" of moment,

than during the manual trials. For more on correlational analysis see Fig.S13 and Fig.S14.

Changes in Neural Time Lag (Observational vs. Manual)

Figure 7: Changes in neural time lags where neural data best correlate with peak grip-force. Bar plots showing the shift of peak correlation between similar (cued/Uncued) Manual and Observational blocks for cS1, rM1, PMd and PMv cortices (cortex is labelled on the title of each subplot). The shift from a block type to another block type (manual-to-observational) is at the top right of each subplot for each NHP (Mc= Manual Cued, Mu= Manual Uncued, Oc= Observation Cued, Ou= Observation Uncued). The number of units used is indicated in the upper right corner of each subplot. The left column for each NHP shows the shift between cued blocks and the right column shows the shift between uncued block. An Asterisks (*) symbol before the number of units represents a significant (p<0.01) shift in histogram from manual to observational tasks. Probability of unit shifts was calculated by subtracting the position of the peak correlation time bin for manual from the observational tasks for the same unit. For more on this correlational analysis see Fig.S13 and Fig.S14.

Force and Reward Modulation of Single Mirror Neurons

Figure 8 shows that a good percentage of units show MN activity encoding force during both manual and observational trials, Fig.8.a. We also show the percentage and number of units that were significantly correlated with force or reward separately, as well as units that were significant for both, but not simultaneously, which would be comodulation and is shown in Figs.9 - 10. Units in Fig.8 were significant under both manual and observational trials. Single unit activities were tracked across manual and observational cued and uncued blocks to determine the significance of their correlation with force under these different conditions. To clarify, a unit that was present during both cued manual and cued observational blocks was checked for significant correlation with force and/or reward and included in Fig.8 only if it was significant for both manual and observational trials towards possible discovery of MNs. The same procedure was performed for uncued manual and observational blocks separately. The plots in Fig.8 show the % of units (left y-axis) and total number of units (top of each bar) for each category that were identified from cS1, rM1, PMd and PMv recordings. The green bars are for reward level cued trials while the purple bars are for reward level uncued trials. The analysis windows used for reward were, for post-cue from 0s to 0.5s, and for post-result from 0s to 0.5s while for force significance the window was 0.5s starting at pre-force-onset to 0.5s after post-force-offset. Note these time windows were not overlapping, thus Fig. 8 is not describing results of force tuning modulation by reward, which is described in Figs.9 - 10.

Percentage of Force-Mirror-Neurons That Multiplex Force and Reward

Figure 8. Mirror Neurons in cS1, rM1, PMd and PMv Multiplex Reward and Grip-force. The

percentage of units with significant modulation via either grip-force (F test, BH method for p=0.05),

reward (post-cue or post-reward, t-test, BH method for p=0.05), or both, during manual and

observational blocks of a similar type (e.g. cued or uncued). Number of hypothesis tests for the

post hoc BH method was the number of MNs in that brain region under study, seen above each

bar. Green represents units for cued blocks and Purple for uncued blocks. Each subplot shows the

percentage of significant units between cued/uncued for units showing such activity during both

manual and observational blocks for Force (plot a and b), Reward (plot c and d) and significant for

both force and reward (plot e and f) during separate times in the task. For subplots c, d, e and f

the left plot shows units for post reward cue and right plot shows units for post reward activity after

reward delivery (150ms) was completed.

Comodulation of Force Tuning Curves by Cued Reward Level

In Figs.9-10 we are asking questions about the modulation of the force tuning curves of MNs by

cued reward level. Force tuning curves for R1 and R0 trials as well as significant differences

between their slopes were calculated as described in the methods using MATLAB's ANCOVA

function. F-statistics were conducted on the reward group*force interaction which expresses the

difference in slopes and the p-values for those interactions were collected. Fig.9 shows the force

tuning curves for example units recorded from reward cued observational blocks for each brain

region (see Fig.S15 for the manual version). The relationship between R1 and R0 trials as spike

rate varied with force can be seen in Fig.9. Note this is in comparison to the previous sections

when force decoding meant the neural activity could be used to determine the force level, whereas

here we are looking at the change in firing rate as a change in force level and its modulation by

reward (encoding). The smoothed spike rate against force is shown in the left subplots while the

line plots to the right show force tuning curves obtained from the ANCOVA. The two example units

from each NHP, S and P, had significant differences between R0 and R1, in agreement with our

previous rM1 results for manual trials [8], again, results in Fig.9 are for observational trials only and

manual versions can be seen in Fig.S15.

 Spike Rate vs Force (Observational)

Figure 9: Plots of spike rate vs force (left subplots) and their linear tuning curves (right subplots) for example units from cS1 (plot *a, e*), rM1 (plot *b, f*), PMd (plot *c, g*) and PMv (plot *d, h*) cortices of both NHPs (for NHP S plots *a, b, c* and *d* and for NHP P plots *e, f, g* and *h*). The units presented had significant differences between R0 and R1 groups (ANCOVA, F-test, $p < 0.05$) force tuning curves during observational blocks. Red lines indicate rewarding trials (R1) and blue indicate non-rewarding trials (R0). See Fig.10 for population results and Fig.S15 for the manual version of this figure.

In Fig. 10 we show results indicating the degree to which cS1, rM1, PMd and PMv not only multiplex

information on both force and reward expectation during both manual and observational tasks, but

how reward expectation modulates the force tuning functions that is the degree of comodulation.

Fig. 10 green bar plots indicate the percentage of single units that have significant grip-force tuning

curves (ANVOCA, $p<0.01$) for the NHP and task type indicated at the subplot title. Purple bars

show the subpopulation of these units (green bars) that pass a rather stringent post-hoc false

discovery rate correction (BH, $p<0.01$, # of hypothesis corrected for is the number of units being

considered for tuning curve testing in that brain region). Red bars indicate the subpopulation of the

grip-force units that are modulated for both task types (manual and observational) that is MNs.

Likewise, yellow bars indicate the MN subpopulation after the post-hoc test (BH, $p<0.01$, # of units).

Fig. 10 indicates that Comodulation of force by reward is most likely significant in manual tasks but

support for the MN activity during observation of this comodulation of force by reward is rather

weak (1-2% max), which could in part be due to the lower number of trials recorded during

observation compounded by the lower firing rate. See tables S3-S4 for more information on the

ANCOVA results.

Figure 10: The percentage of single units with force tuning curves that are significantly modulated by reward level. The titles represent which NHP and block type are plotted, p-values were adjusted using the Benjamini-Hochberg method (BH method) correcting for false discovery rate where the correction was applied using p=0.01 and the number of hypothesis tests applied was the number of units in the population under study. The y-axis is the percentage of units and x-axis represents the cortical region. The green bar is showing the number of significant units (p<0.01, ANCOVA) and red is showing the MN population among them. The purple bar is showing the significant units after BH method was applied. The yellow bar inside the purple is showing the percentage of putative MNs after BH method. The number after the '#' sign on top of each bar is showing the number of significant units for that given case and the number presented with 'n' is the total units recorded for that case.

Discussion

The main outcome of this work is the clear support showing that MN's can encode grip force (Fig.5-6 and Tables.S1 and S2), a non-kinematic variable in cS1 that is areas 1 and 2 as well as in rM1, PMd and PMv (see Fig.2). Secondly, that the MN activity during observational trials, as compared

to manual trials, can shift their temporal relationship predicting/responding to visually cued force

about equally in either the predictive or responsive time lag directions seen in Fig.7. Thirdly, we

have shown that the firing rate in each of these regions is also modulated by reward level

expectation during the post-reward-cue period for reward-level-cued trials, and during the post-

feedback time period for both cued and uncued reward level trials seen in Fig.8. Fourthly, that MNs

activity of visually cued grip-force can coexist (multiplex) within single units that also code for

reward level seen in Fig.8. Finally, we have shown evidence, albeit weak (~3-4% of MNs for

manual and ~1-2% of MNs for observational tasks), that the neural grip-force tuning functions can

be modulated by reward expectation (comodulation) in the brain regions under study (cS1, rM1,

PMd and PMv) during both manual and observational trials Figs.9-10.

To the best of our knowledge, we are reporting the first evidence of mirror neural responses to

expected, or visually cued effort (grip-force), within these sensorimotor regions at the single unit

level. As the NHPs had a great deal of experience with the cued grip-force task we expect the

mirror responses were due to the visual cueing of the force targets that the NHPs had come to

understand based on their manual trials. In Fig.7 we showed that the shift in peak correlation

between cued force and neural time lag could be in either the direction expected for motor

production, or that expected from sensory stimulation/gating, which would be more in line with the

initial MN work [1], and the former more in line with movement rehearsal [2]. We predict that such

force MN activity could be seen in more natural settings if the context was understood by the NHPs,

such as lifting heavy vs. light objects that they were familiar with, or perhaps when the NHP

squeezes a deformable object, or observes such a grasp, again with knowledge of the object's

physical properties. However, further work is needed to show such activity exists under more

natural conditions within each of these regions. It should be noted that this type of prior knowledge

being necessary for MN activity patterns is not new. Mirror neurons that encode subjective value

in addition to grasping information clearly need the NHP to have familiarity with the object in some

way to have formed a subjective value [46].

We have shown that reward modulation can occur post-cue, when the cue indicates the reward

value of the current trial, as well as post-feedback when the reward level that was cued is delivered.

When there is no reward cue given in uncued blocks there is no "post-cue" reward modulation as

expected, but there is still post-feedback modulation in each of the 4 brain regions, and again

during both manual and observational trial types. Based on the evidence we have presented, we

see that cS1, rM1, PMd and PMv contain units that encode reward expectation and reward itself,

as has been shown previously for the task used in this work as well as other tasks for rM1 [6,8,23,28],

indicating this reward modulation may be a generalized broadcast signal to regions involved in the

ongoing sensorimotor planning and production of movement, as well as during observation of such

movements, in a manner that can be predictive, as expected during mental simulation [2], and

responsive to the visual stimuli, as expected for classical MNs (Rizzolatti et al. 1996; Gallese et al.

1996). In addition, we have now shown such reward modulation simultaneously for rM1, cS1, PMd

and PMv during both manual and observational trial types for the same single units tracked

between tasks.

There are several limitations to the work presented here that should be addressed in future work.

EMGs were not available for these particular datasets that allowed us to track single units between

manual and observational trials, and therefore we can't state with certainty that the NHPs were not

activating their arm muscles covertly, however, they were not making obvious movements, and did

not have access to the force transducing handle, in addition, our previous studies found no

correlation between EMGs and curser motion in observational trials of a reaching task [28]. In

addition, we show in Fig.S16 for both NHPs there was no significant EMG activity during

observational trials on different days when the EMG signals were usable. The grip-force output in

the current task was generally phasic in nature with a bell-shaped profile, even if the cued

amplitudes were different, the force profiles were still fairly stereotyped, and so it is possible that

some other phasic neural response was allowing the regression models to predict force output,

which was bell-shaped, by some non-force related phasic response, such as that related to the

cued reward information. However, the regression models did not fall apart in the uncued reward

level task, which indicates at least cued reward does not explain our force decoding results. In

addition, when we conducted analysis on the time series of just the peak force amplitude of each

trial, we still obtained positive results showing MNs for force, although clearly not as strong as

when using the fuller dataset including the force trajectories (see Fig.S19). Additionally, in Fig.S20

we show significance holds for several measures on the manual data between R0 and R1 trials

post data pruning to keep the trials between these two categories statistically indistinguishable.

Previous work has suggested that MN activity is not seen in S1 [47], however they did see non-

frequency discriminative activation of S1 due to auditory stimuli. Others have suggested that S1 is

in fact modulated in a MN manner, however, they utilized fMRI [48]. Histological studies where MN

brain regions were injected with tracers did show that IPL injections led to staining of S1 [49], and

these studies by Meyer et al. and Bruni et al. would imply our results are perhaps expected. The

data we present for MNs encoding grip force are very clear and significant as are the results

showing that these MNs multiplex cued reward level, or reward expectation and force, however,

the results for comodulation of force tuning curves by reward were less convincing, but still

significant.

The work presented here has a practical application past basic neuroscience. We believe it is

important to continue gaining a better understanding of encoded information, such as reward,

within the sensorimotor cortices toward the development of a close-loop brain machine interface

(BMI) for the restoration of motor control and beyond, such as towards a better understanding and

tracking of psychological state of the individual. BMI neural signals are often recorded and decoded

from a subset of cS1, rM1, PMd, and PMv while sensory feedback is obtained either by natural

vision, or stimuli routed to cS1, such as via the thalamus [50] for somatosensory feedback. It has

recently been shown that reward expectation can change directional and force based tuning curves

in rM1 and cS1 [8,23], and here we have shown that all of these brain regions are influenced by

reward expectation, effort, or sensory feedback on expected effort, and therefore research into

these relationship is warranted. Differentiating between mirror activity in these regions and

intentional activity for movement is key towards making BMIs more stable with respect to the users

intended movements compared to observed movements, and our future work will address these

issues.

References

1. Gallese, V., Fadiga, L., Fogassi, L. & Rizzolatti, G. Action recognition in the premotor cortex.

Brain **119 (Pt 2)**, 593–609 (1996).

2. Cisek, P. & Kalaska, J. F. Neural correlates of mental rehearsal in dorsal premotor cortex.

Nature **431**, 993–6 (2004).

3. Dushanova, J. & Donoghue, J. Neurons in primary motor cortex engaged during action

observation. *European Journal of Neuroscience* **31**, 386–398 (2010).

4. Vigneswaran, G., Philipp, R., Lemon, R. N. & Kraskov, A. M1 corticospinal mirror neurons and

their role in movement suppression during action observation. *Curr Biol* **23**, 236–43 (2013).

5. Alaerts, K., de Beukelaar, T. T., Swinnen, S. P. & Wenderoth, N. Observing how others lift light

or heavy objects: time-dependent encoding of grip force in the primary motor cortex.

Psychological Research **76**, 503–513 (2012).

6. An, J., Yadav, T., Hessburg, J. P. & Francis, J. T. Reward Expectation Modulates Local Field

Potentials, Spiking Activity and Spike-Field Coherence in the Primary Motor Cortex. *eNeuro* **6**,

(2019).

7. Marsh, B. T., Tarigoppula, V. S., Chen, C. & Francis, J. T. Toward an autonomous brain

machine interface: integrating sensorimotor reward modulation and reinforcement learning. *J*

Neurosci **35**, 7374–87 (2015).

8. Zhao, Y., Hessburg, J. P., Asok Kumar, J. N. & Francis, J. T. Paradigm Shift in Sensorimotor

Control Research and Brain Machine Interface Control: The Influence of Context on

Sensorimotor Representations. *Front. Neurosci.* **12**, (2018).

9. Sanchez, J. C. *et al.* Control of a center-out reaching task using a reinforcement learning Brain-

Machine Interface. in *2011 5th International IEEE/EMBS Conference on Neural Engineering*

525–528 (2011). doi:10.1109/NER.2011.5910601.

10. Pohlmeyer, E. A., Mahmoudi, B., Geng, S., Prins, N. & Sanchez, J. C. Brain-machine interface

control of a robot arm using actor-critic reinforcement learning. in *2012 Annual International*

Conference of the IEEE Engineering in Medicine and Biology Society 4108–4111 (2012).

doi:10.1109/EMBC.2012.6346870.

11. Pohlmeyer, E. A., Mahmoudi, B., Geng, S., Prins, N. W. & Sanchez, J. C. Using Reinforcement

Learning to Provide Stable Brain-Machine Interface Control Despite Neural Input

Reorganization. *PLoS One* **9**, (2014).

12. An, J., Yadav, T., Ahmadi, M. B., Tarigoppula, V. S. A. & Francis, J. T. Near Perfect Neural

Critic from Motor Cortical Activity Toward an Autonomously Updating Brain Machine Interface.

in *2018 40th Annual International Conference of the IEEE Engineering in Medicine and Biology*

Society (EMBC) 73–76 (2018). doi:10.1109/EMBC.2018.8512274.

13. Rolls, E. T. *Emotion and decision-making explained*. (Oxford University Press, 2014).

14. *Neuroeconomics: Decision Making and the Brain*. (Academic Press, 2008).

15. Schultz, W., Dayan, P. & Montague, P. R. A Neural Substrate of Prediction and Reward.

Science **275**, 1593–1599 (1997).

16. Zaghloul, K. A. *et al.* Human Substantia Nigra Neurons Encode Unexpected Financial

Rewards. *Science* **323**, 1496–1499 (2009).

17. Sato, M. & Hikosaka, O. Role of Primate Substantia Nigra Pars Reticulata in Reward-Oriented

Saccadic Eye Movement. *J. Neurosci.* **22**, 2363–2373 (2002).

18. Cohen, J. Y., Haesler, S., Vong, L., Lowell, B. B. & Uchida, N. Neuron-type-specific signals for

reward and punishment in the ventral tegmental area. *Nature* **482**, 85–88 (2012).

19. Murray, G. K. *et al.* Substantia nigra/ventral tegmental reward prediction error disruption in

psychosis. *Molecular Psychiatry* **13**, 267–276 (2008).

20. Tremblay, L. & Schultz, W. Relative reward preference in primate orbitofrontal cortex. *Nature*

398, 704–708 (1999).

21. Roesch, M. R. & Olson, C. R. Impact of Expected Reward on Neuronal Activity in Prefrontal

Cortex, Frontal and Supplementary Eye Fields and Premotor Cortex. *Journal of*

Neurophysiology **90**, 1766–1789 (2003).

22. Ramkumar, P., Dekleva, B., Cooler, S., Miller, L. & Kording, K. Premotor and Motor Cortices

Encode Reward. *PLoS ONE* **11**, e0160851 (2016).

23. Ramakrishnan, A. *et al.* Cortical neurons multiplex reward-related signals along with sensory

and motor information. *PNAS* **114**, E4841–E4850 (2017).

24. McNiel, D., Bataineh, M., Choi, J., Hessburg, J. & Francis, J. Classifier Performance in Primary

Somatosensory Cortex Towards Implementation of a Reinforcement Learning Based Brain

Machine Interface. in *2016 32nd Southern Biomedical Engineering Conference (SBEC)* 17–

18 (2016). doi:10.1109/SBEC.2016.19.

25. McNiel, D. B., Choi, J. S., Hessburg, J. P. & Francis, J. T. Reward value is encoded in primary

somatosensory cortex and can be decoded from neural activity during performance of a

psychophysical task. in 3064–3067 (IEEE, 2016).

26. Atique, M. M. & Francis, J. T. Reward and force modulation of neurons in the primate primary

somatosensory cortex (S1). https://www.abstractsonline.com/pp8/#!/7883/presentation/59642

(2019).

27. Pleger, B., Blankenburg, F., Ruff, C. C., Driver, J. & Dolan, R. J. Reward Facilitates Tactile

Judgments and Modulates Hemodynamic Responses in Human Primary Somatosensory

Cortex. *J. Neurosci.* **28**, 8161–8168 (2008).

28. Marsh, B. T., Tarigoppula, V. S. A., Chen, C. & Francis, J. T. Toward an Autonomous Brain

Machine Interface: Integrating Sensorimotor Reward Modulation and Reinforcement Learning.

J. Neurosci. **35**, 7374–7387 (2015).

29. Tkach, D., Reimer, J. & Hatsopoulos, N. G. *J. Neurosci.* **27**, 13241–13250 (2007).

30. Vigneswaran, G., Philipp, R., Lemon, R. N. & Kraskov, A. M1 Corticospinal Mirror Neurons and Their Role in Movement Suppression during Action Observation. *Current Biology* **23**, 236–243 (2013).

31. Mazurek, K. A., Rouse, A. G. & Schieber, M. H. Mirror Neuron Populations Represent Sequences of Behavioral Epochs During Both Execution and Observation. *J. Neurosci.* **38**, 4441–4455 (2018).

32. Rizzolatti, G. & Craighero, L. The Mirror-Neuron System. *Annual Review of Neuroscience* **27**, 169–192 (2004).

33. Gazzola, V. & Keysers, C. The Observation and Execution of Actions Share Motor and Somatosensory Voxels in all Tested Subjects: Single-Subject Analyses of Unsmoothed fMRI Data. *Cereb Cortex* **19**, 1239–1255 (2009).

34. Keysers, C., Kaas, J. H. & Gazzola, V. Somatosensation in social perception. *Nature Reviews Neuroscience* **11**, 417–428 (2010).

35. Rossi, S. *et al.* Somatosensory processing during movement observation in humans. *Clinical Neurophysiology* **113**, 16–24 (2002).

36. Keisker, B., Hepp-Reymond, M.-C., Blickenstorfer, A., Meyer, M. & Kollias, S. S. Differential force scaling of fine-graded power grip force in the sensorimotor network. *Human Brain Mapping* **30**, 2453–2465 (2009).

37. Quigley, M. *et al.* ROS: an open-source Robot Operating System. in *ICRA workshop on open source software* vol. 3 5 (Kobe, Japan, 2009).

38. Chhatbar, P. Y., von Kraus, L. M., Semework, M. & Francis, J. T. A bio-friendly and economical technique for chronic implantation of multiple microelectrode arrays. *Journal of Neuroscience Methods* **188**, 187–194 (2010).

39. Kipke, D. R., Vetter, R. J., Williams, J. C. & Hetke, J. F. Silicon-substrate intracortical microelectrode arrays for long-term recording of neuronal spike activity in cerebral cortex. *IEEE Transactions on Neural Systems and Rehabilitation Engineering* **11**, 151–155 (2003).

40. Kretzberg, J., Coors, T. & Furche, J. Comparison of valley seeking and T-distributed EM
algorithm for spike sorting. *BMC Neuroscience* **10**, P47 (2009).

41. Yuan, Y., Yang, C. & Si, J. The M-Sorter: An automatic and robust spike detection and
classification system. *Journal of Neuroscience Methods* **210**, 281–290 (2012).

42. Benjamini, Y. & Hochberg, Y. Controlling the False Discovery Rate: A Practical and Powerful
Approach to Multiple Testing. *Journal of the Royal Statistical Society: Series B*
(Methodological) **57**, 289–300 (1995).

43. Maranesi, M., Livi, A., Fogassi, L., Rizzolatti, G. & Bonini, L. Mirror neuron activation prior to
action observation in a predictable context. *J Neurosci* **34**, 14827–32 (2014).

44. Hendrix, C. M., Mason, C. R. & Ebner, T. J. Signaling of Grasp Dimension and Grasp Force in
Dorsal Premotor Cortex and Primary Motor Cortex Neurons During Reach to Grasp in the
Monkey. *Journal of Neurophysiology* **102**, 132–145 (2009).

45. Ro, J. Y., Debowy, D., Ghosh, S. & Gardner, E. P. Depression of neuronal firing rates in
somatosensory and posterior parietal cortex during object acquisition in a prehension task.
Experimental Brain Research **135**, 1–11 (2000).

46. Caggiano, V. *et al.* Mirror neurons encode the subjective value of an observed action. *Proc*
Natl Acad Sci U S A **109**, 11848–53 (2012).

47. Lemus, L., Hernández, A., Luna, R., Zainos, A. & Romo, R. Do Sensory Cortices Process More
than One Sensory Modality during Perceptual Judgments? *Neuron* **67**, 335–348 (2010).

48. Meyer, K., Kaplan, J. T., Essex, R., Damasio, H. & Damasio, A. Seeing Touch Is Correlated
with Content-Specific Activity in Primary Somatosensory Cortex. *Cerebral Cortex* **21**, 2113–
2121 (2011).

49. Bruni, S. *et al.* Cortical and subcortical connections of parietal and premotor nodes of the
monkey hand mirror neuron network. *Brain Struct Funct* **223**, 1713–1729 (2018).

50. Choi, J. S. *et al.* Eliciting naturalistic cortical responses with a sensory prosthesis via optimized
microstimulation. *J. Neural Eng.* **13**, 056007 (2016).

SUPPLEMENTARY INFORMATION

Mirror Neurons are Modulated by Grip Force and Reward Expectation in the Sensorimotor

Cortices (S1, M1, PMd, PMv)

Abbreviated title:

Md Moin Uddin Atique[1] and Joseph Thachil Francis[1-2] [*]

Correspondence: Joey199us@gmail.com

Acknowledgements: Research was supported by NIH 1R01NS092894-01, NSF IIS-
17 1527558, DARPA REPAIR Project N66001-10-C-2008.

[1]Department of Biomedical Engineering, Cullen College of Engineering, The University of

Houston, Houston, TX 77204, United States.

[2] Department of Electrical and Computer Engineering, Cullen College of Engineering, The

University of Houston, Houston, TX 77204, United States.

[*] Corresponding author - jtfracis@uh.edu

Supplementary Information

Autocorrelation on the reward level target sequence:

Figure S1: Autocorrelation on the reward level target sequence (R0 and R1) of trials for all data

blocks. Plot a, c, e, and g are for NHP S and b, d, f, and h is for NHP P. The block type is given

on the right side of each row. The x-axis represents lag in trial numbers and y axis represents

corresponding autocorrelation.

The number of single units recorded for each cortex:

40   The figure below shows the total number of single units recorded in all the cortices used (S1, M1,

41   PMd, and PMv) and for each of the block types (MC, MU, OU, and OC). The block types are

42   color coded as seen in the legend in the right subplot. The left subplot shows the number of units

43   from NHP S and the right shows from NHP P. The x-axis is the cortical region and the y-axis is

44   the number of units.

Figure S2: The total number of units recorded in all four cortices (S1, M1, PMd, and PMv) for

47   both NHP S (left) and NHP P (right). For each cortex there is four columns for four data block

48   types used (MC, MU, OU, and OC) that are color cued as in the legend on the right subplot. The

49   y-axis represents the number of units.

Decoded Force R-square value with and without smoothening (Fig.S3):

We applied MLR (Multiple linear regression) to decoded grip force from smoothened spike rates

(shown in main text figure 5). In order to show that the smoothening was not playing a role in our

decoding results we also tested the square root transformed spike rate without smoothing to

decode the force following the same procedure described in the method section. The following

figure shows the R-square values of force decoding with (red) or without (blue) smoothening.

Figure S3: The R-square of the decoded grip force with (blue) or without (red) smoothening was

applied on the spike rate data. The rows of subplot are showing data from S1, M1, PMd and PMv

cortices of NHP S and bottom row is same for NHP P.

Grip Force prediction without Square root transformation (Fig.S4):

In the following figure S4, we show a comparison between the grip decoding accuracy with R-

square values between decoded grip force on the test set of data and the actual grip force

applied (manual) or observed (observational). The bar plots showed that the R-squared doesn't

deviate much when a square root transform is applied (blue) on the spike rate from the case

when the transformation was not applied (Red). The figure 8 in the main text shows that NHP S

had a higher percentage of significant units, especially in M1, PMd and PMv cortex due to

reward cue compared to NHP P.

Figure S4: The R-square of the decoded grip force with (blue) or without (red) square root

transformation was applied on the spike rate data. The rows of subplot are showing data from

S1, M1, PMd and PMv cortices of NHP S and bottom row is same for NHP P.

85   **Algorithm for Linear Regression:** The following algorithm shows the two-step process of

86   collecting a subset of units significant for force to achieve a comparatively better prediction of

87   force using linear regression. **Fig.S5**

89   **Figure S5:** Algorithm for linear Regression

The Force duration for each data block:

The durations of applied grip force in R0 (blue) and R1 (red) trials are given on the following box

plots for each of the block types (MC= Manual Cued, MU= Manual Uncued, OC= Observation

Cued and OU=Observation Uncued). The box plots are showing the highest (upper line), 75[th]

percentile, median, 25[th] percentile and the lowest values for force duration of all the trials. A

paired t-test (p-value <0.05) was performed to check if the R0 peak force was significantly

different from R1 trials. The R0 and R1 force durations are significantly different only for the case

of MC block of Monkey S.

Figure S6: The box plot is showing the median, maximum and minimum values with 75th and

25th percentile of the grip force duration of all the trials on each block individually. The x-axis is

showing the data block type (also shown on the legend which represents same information).

The y-axis is representing the force duration in seconds. The black line and asterisk (*) on the

top are showing the significant difference between R0 (blue) and R1 (red) trials.

Peak grip force for each data block:

The applied/observed peak grip force on each manual/observational blocks are plotted on the

following figures. The Red is representing the R1 trials and blue is R0 trials. The block type (MC,

MU, OC, OU) is mentioned in the x-axis. A paired t-test (p-value <0.05) was performed to check

if the R0 peak force was significantly different from R1 trials. The significance test showed no p-

value less than 0.05 and hence no asterisk (*) sign was plotted in any of the pairs. The minimum

threshold of force for monkey S was 150 and P was 100 for the trial that had lowest force peak

threshold. If the value goes below that value, the trial will be failed. The value mentioned is the

minimum value for different trails a random value was added with the to increase the minimum

threshold from 100 (monkey P) and 150 (monkey S).

Figure S7: The peak force from each block type is plotted on the following box plot and for each

block type the R0 (blue) and R1 (red) trials are plotted separately for NHP S (left) and NHP P

(right). The x-axis is showing the block type (MC, MU, OC and OU). The y-axis is representing

the value of the grip force applied with arbitrary units. The grip force values from R0 and R1 trials

were significantly different for any of the case and hence no asterisk (*) is present.

Reaction time to apply grip force:

The reaction time to apply grip force for R0 and R1 trials is plotted on the following subplots. The

manual blocks for Monkey S (left) and Monkey P (right) were plotted. The blue and red color

represents R0 (blue) and R1 (red) trials from the same data block (MC, MU). The asterisk (*) on

the top of the black line indicates the reaction time is significantly different between R0 and R1

trials (paired t-test, p-value <0.05). The reaction time in this case is the difference between the

times, NHP was allowed to apply grip force (start of the grip force scene) to the time when they

applied 10% of the peak grip force. From the following plot the reaction time is significantly

different only for the manual cued block for Monkey S.

Figure S8: The reaction time for R0 (blue) and R1 (red) trials for all the block type is given on the

following plots. The x-axis is showing the block types MC, MU for the case of manual blocks. The

y-axis is showing the reaction time in seconds. The asterisk (*) on the top of each pair

represents they reaction time is significantly different between R0 and R1 trials. The left subplot

is for Monkey S and right subplot is for Monkey P.

The number of trials in each data block:

The following figure shows the number of trials for each data block that were used for analysis.

Figure S9: The number of trials used for the data analysis is shown on the figure above for

monkey S (left) and monkey P (right). Each bar represents the total number of trials and the blue

147   part of the bar is showing R0 trials where the red part is for R1 trials. The x-axis is representing

148   the block types (MC, MU, OC and OU) and y-axis is showing the number of trials.

161   **Observation Results on force (Fig.S10):**

162   This figure shows raster plots of some more example units from each of the 4 cortices during grip

163   force onset and offset taken from observation blocks showing the full average force time period

164   with both increases and decreases to force onset.

Figure S10: Raster plots for units from S1 (plots a, b, c and d), M1 region (plots e, f, g and h),

PMd (plots i, j, k and l) and PMv (plots m, n, o and p) are shown for NHPs S and P. These plots

show the grip force values expected given the cued force targets shown to the NHPs during

observational blocks. The black bold line on each raster and spike rate plot shows the onset of

"force" and the dotted horizontal line divides R1 (top, red) and R0 (bottom, blue) trials. On each

neural data subplot, the x-axis represents time in seconds and for raster plots y-axis represents

trial number (upper part of the subplots) and for the spike rate plot y-axis is spike rate in 'Hz'

(bottom part of the subplots). The Asterisk (*) symbol on the right top corner of the mean spike

rate plots represents, that unit was significant (F-test, $p<0.05$) for force fit using linear regression

model in eq 1.

Responses of Mirror neuron activity due to grip force for different data blocks:

The following figures are showing the number of MN units in terms of their activity during

different blocks. These units are already shown in figure 6 of the main manuscript but the other

category was not elaborated there, which is given in detail in figure S11 and S12. For analysis,

four data blocks for each NHP (NHP S and NHP P) were used. The abbreviations for the block

types are, MC=Manual Cued, MU=Manual Uncued, OU=Observational Uncued and

OC=Observational Cued. For figure 6, we used the peak value from the absolute correlation

coefficients between grip force and each of the 10 bins of spike rate (100ms from -0.5ms pre grip

force spike rate to 0.5s post, described "Grip force trajectory prediction" section inside method) to

detect inhibitory or excitatory activity. We considered a unit's representation excitatory if the

actual value of that peak absolute correlation was positive otherwise it was considered

inhibitory. We detected congruent units among the significant units that showed similar activity in

all four data blocks of either inhibitory (blue) or excitatory (red) spike rate in relation to grip-force.

Incongruent units showed opposing behavior between manual and observational tasks, such as

excitatory during manual and inhibitory during observational, purple, or inhibitory during manual

and excitatory during observational (green). Other (gray) units didn't follow any simple pattern

during all four data blocks.

In the following two figures (S11 and S12) we are showing the "other" units in detail for each of

the combination pattern. On the x axis label is showing four numbers for four data blocks in the

order of (MC-MU-OU-OC) and a number '0' represents inhibitory activity for that block and '1'

represents excitatory activity. So, for example the bar with the first label '0000', shows the

number of units that showed inhibitory spike rate during grip force activity during all four data

blocks (MC, Mu, OU and OC). The bar colors are kept as like figure 6 and given as the legend.

The bars that contain the other unit category from figure 6 are kept gray colored here.

Figure S11: The Number of MN units from NHP S, for each possible activity combination during

blocks in order of MC, MU, OU and OC. The y-axis represents the number of units and x-axis

label shows the combination of inhibitory (0) or excitatory activity for the given four data blocks.

The title on each subplot shows the corresponding cortex and NHP name.

Figure S12: The Number of MN units from NHP P, for each possible activity combination during

blocks in order of MC, MU, OU and OC. The y-axis represents the number of units and x-axis

label shows the combination of inhibitory (0) or excitatory activity for the given four data blocks.

The title on each subplot shows the corresponding cortex and NHP name.

 Time Dependent Correlation Analysis (Fig.S13):

The time dependent peak correlation plots for all the blocks are given below. Only units

significant for force encoding that were tracked through all manual and observational blocks are

shown below.

Figure S13: The number of units that had their highest correlation between force and spike rate

for each time bin are shown leading (0.5s before) or lagging (0.5s after) each force value (100ms

non-overlapping time bins). Units significant (F-test, $p < 0.05$) on all block types were plotted in

the time bin where their correlation coefficient with force was maximum in absolute value, either

negative or positive. The y-axis represents the number of units (positively correlated units on the

upper side and negatively correlated units on the lower). The x-axis is representing the time

lag/lead for which the maximum correlation was found. Units from S1 (a, b), M1 (c, d), PMd (e, f)

and PMv (g, h) cortex are given on each plot and the legend is showing the color diagram for

each cortex.

Distribution of Significant Correlation Time Lags Between Neural Activity and Force

(Fig.S14):

The following figure is showing the time shift of significant peak correlation from manual cued

blocks to manual uncued blocks and from observation cued blocks to observation uncued blocks.

See Fig.7 for more of such plots between manual and observational.

Figure S14: The shift of peak force correlation from manual to manual and observational to

observational block. Bar plots showing the shift of peak correlation between Manual blocks and

Observational blocks for S1, M1, PMd and PMv cortex for individual MNs. The shift from a block

type to another block type is given on the left to of each figure and short form of the block type

(Mc= Manual Cued, Mu= Manual Uncued, Oc= Observation Cued, Ou= Observation Uncued)

was used. The number of units used is included on the upper right corner of each subplot. An

asterisk (*) symbol before the number of units represents a significant shift in histogram between

the blocks used for that subplot. Shift was calculated by subtracting the manual histogram for

cued from the Manual histogram for uncued and similarly for the observational cases.

Spike Rate vs Force Fig.S15 (Manual)

Figure. S15: Plots of spike rate vs force (left subplots) and their linear tuning curve (right

subplots) for Grip force manual trials as compared to Fig.9 for observational. The subplots show

example units from S1 (plot *a, e*), M1 (plot *b, f*), PMd (plot *c, g*) and PMv (plot *d, h*) cortices of

both NHPs (for NHP S plots *a, b, c* and *d* and for NHP P plots *e, f, g* and *h*). The units presented

here showed a significant difference between R0 and R1 groups (ANCOVA, F-test, $p<0.05$) force

tuning curves during observation of robotic manipulator applying grip force on a force sensor.

Red lines indicate rewarding trials (R1) and blue indicate non-rewarding trials (R0).

Regression on additional observational blocks:

Table S1 shows the regression data analysis for NHP S and P. The table contains the R-

square value for model fit given in equation 1 and the corresponding p-value and F-statistic. The

R-squared value for prediction and its p-value is also shown on the table. This table contains two

additional observational blocks that were recorded on a different day from the data shown in the

main text.

Table S1: Regression Model Output for NHP S & P for the two additional observational blocks

recorded on a different day with units we could not track from the previous data sets.

NHP S - S1

Block	R-fit	p-value	F-stat	R-predict	p-value
Observation Cued	0.28	3e-91	3.81	0.19	8.3e-21
Observation Uncued	0.34	4.3e-154	4.79	0.04	9.3e-6

NHP P – S1

Block	R-fit	p-value	F-stat	R-predict	p-value
Observation Cued	0.46	1.3e-229	8.34	0.27	1.9e-26
Observation Uncued	0.53	0	10.09	0.13	1.8e-14

NHP S – M1

Block	R-fit	p-value	F-stat	R-predict	p-value
Observation Cued	0.65	0	11.59	0.6	2.6e-87
Observation Uncued	0.76	0	21.19	0.66	4.3e-113

NHP P – M1

Block	R-fit	p-value	F-stat	R-predict	p-value
Observation Cued	0.61	0	13.19	0.55	2.7e-64
Observation Uncued	0.69	0	18.07	0.55	3.1e-75

NHP S – PMd

Block	R-fit	p-value	F-stat	R-predict	p-value
Observation Cued	0.53	1.9e-273	6.63	0.36	2.7e-43

NHP P – PMd

Block	R-fit	p-value	F-stat	R-predict	p-value
Observation Cued	0.5	6.4e-274	10	0.52	6.3e-59

Observation Uncued	0.67	0	11.57	0.39	0.55		Observation Uncued	0.55	0	13.14	0.37	2e-44

NHP S – PMv							NHP P – PMv					
Block	R-fit	p-value	F-stat	R-predict	p-value		Block	R-fit	p-value	F-stat	R-predict	p-value
Observation Cued	0.6	0	11.84	0.72	5.7e-121		Observation Cued	0.35	3.5e-197	11.24	0.28	4.9e-27
Observation Uncued	0.8	0	34.11	0.72	6.1e-134		Observation Uncued	0.46	0	13.18	0.36	1.4e-42

294 **Table S2:** Regression model statistics for Fig. 5 in the main text.

295 **Table S2: Regression Model Output for NHPs S & P**

NHP S - S1							NHP P – S1					
Block	R^2-fit	p-value	F-stat	R^2-predict	p-value		Block	R^2-fit	p-value	F-stat	R^2-predict	p-value
Manual Cued	0.91	0	$F_{550,2173}$ 39.83	0.78	1.9e-101		Manual Cued	0.78	0	$F_{320,2142}$ 24.19	0.84	2.26e-109
Manual Uncued	0.89	0	$F_{540,3597}$ 51.68	0.82	3.5e-173		Manual Uncued	0.9	0	$F_{470,2528}$ 46.77	0.87	4.6e-148
Obs Cued	0.34	7.4e-40	$F_{250,1330}$ 3.32	**0.11**	9.5e-6		**Obs Cued**	0.68	5e-88	$F_{220,707}$ 7.12	**0.66**	1.4e-25
Obs Uncued	0.29	9.4e-46	$F_{300,1679}$ 3.47	**0.14**	7.8e-9		**Obs Uncued**	0.47	2.3e-42	$F_{220,652}$ 4.6	**0.15**	9.5e-5

NHP S – M1							NHP P – M1					
Block	R^2-fit	p-value	F-stat	R^2-predict	p-value		Block	R^2-fit	p-value	F-stat	R^2-predict	p-value
Manual Cued	0.92	0	$F_{590,2133}$ 44.24	0.80	7.8e-109		Manual-Cued	0.80	0	$F_{360,2102}$ =23.71	0.87	5.58e-123
Manual Uncued	0.93	0	$F_{530,3607}$ 85.7	0.83	1.2e-180		Manual-Uncued	0.89	0	$F_{440,2558}$ 46.22	0.87	1.6e-151
Obs Cued	0.69	1.7e-147	$F_{240,1340}$ 6.91	**0.51**	1.6e-28		**Obs Cued**	0.76	6.3e-96	$F_{300,627}$ 7.29	**0.56**	7e-20
Obs Uncued	0.65	1.2e-197	$F_{350,1629}$ 8.19	**0.45**	1.8e-30		**Obs Uncued**	0.57	4.5e-52	$F_{180,692}$ 4.81	**0.32**	1.5e-09

NHP S –PMd	NHP P – PMd

Block	R^2-fit	p-value	F-stat	R^2-predict	p-value
Manual Cued	0.92	0	$F_{550,2173}$ 44.25	0.84	1.3e-122
Manual Uncued	0.92	0	$F_{640,3497}$ 58.86	0.83	9.1e-181
Obs Cued	0.66	6.3e-172	$F_{270,1310}$ 8.99	**0.54**	1.5e-31
Obs Uncued	0.73	2.5e-282	$F_{330,1649}$ 11.95	**0.61**	6.4e-47

Block	R^2-fit	p-value	F-stat	R^2-predict	p-value
Manual Cued	0.85	0	$F_{260,2202}$ 47.17	0.94	4.9e-164
Manual Uncued	0.93	0	$F_{540,2458}$ 63.09	0.91	6.4e-175
Obs Cued	0.73	1.3e-117	$F_{240,627}$ 9.79	**0.66**	1.4e-25
Obs Uncued	0.72	4.7e-82	$F_{200,672}$ 6.71	**0.36**	7.3e-11

NHP S –PMv

NHP P – PMv

Block	R^2-fit	p-value	F-stat	R^2-predict	p-value
Manual Cued	0.83	0	$F_{430,2293}$ 26.72	0.77	2.3e-97
Manual Uncued	0.87	0	$F_{610,3527}$ 40.29	0.67	4.4e-112
Obs Cued	0.81	4e-265	$F_{320,1260}$ 13.65	**0.69**	4.9e-46
Obs Uncued	0.86	0	$F_{430,1549}$ 16.25	**0.77**	7.1e-71

Block	R^2-fit	p-value	F-stat	R^2-predict	p-value
Manual Cued	0.61	5.3e-308	$F_{24,2222}$ 14.26	0.53	9.3e-47
Manual Uncued	0.6	0	$F_{160,2838}$ 26.31	0.57	5.6e-63
Obs Cued	0.43	1.2e-60	$F_{80,847}$ 7.89	**0.29**	2.7e-09
Obs Uncued	0.33	7.9e-45	$F_{80,792}$ 8.15	**0.40**	2.9e-12

Table S2: Results from the regression analysis for NHPs S and P between neural activity and force trajectories. The table contains the R-square value for the model (eq. 2) fit and prediction including their corresponding p-values and F-statistics. We have used bold text for the prediction during observation tasks. Note all single units were followed through each of the 4 task types on a given day. From table 1 it appears that these brain regions are representing force during both the manual trails and the observational versions of these trials. In Fig.6 we show the number of single units that significantly predict force during both manual and observational trails indicating that these same units are representing this information to some extent during both manual and observed movements, which may represent a new form of mirror neuron activity.

Table S3: Units with significantly different slopes for the force tuning curves between R0 and R1. The values in parenthesis show the adjusted significant units after Benjamini and Hochberg's

309   method accounting for the false discovery rate was applied. The BH method applied for number

310   of units on which the hypothesis testing was applied in the population is described in the method

311   section titled "grip force tuning curve analysis".

	S1 #	S1 %	M1 #	M1 %	PMd #	PMd %	PMv #	PMv %
NHP S Manual	11 (3)	8 (2)	20 (4)	14 (3)	28 (24)	20 (17)	13 (4)	8 (2)
NHP P Manual	9 (1)	7 (1)	9 (4)	8 (3)	16 (6)	14 (5)	4 (0)	5 (0)
NHP S Obs.	2 (2)	2 (2)	2 (1)	2 (1)	2 (0)	2 (0)	0 (0)	0 (0)
NHP P Obs.	0 (0)	0 (0)	1 (0)	1 (0)	2 (0)	2 (0)	1 (1)	2 (2)

313   **Table S4**: Units with significantly different y-intercept for the force tuning curves for R0 and R1.

314   The values in parenthesis show the adjusted significant units after Benjamini and Hochberg's

315   method accounting for the false discovery rate was applied. The BH method applied for number

316   of population is described in the method section titled "grip force tuning curve analysis".

	S1 #	S1 %	M1 #	M1 %	PMd #	PMd %	PMv #	PMv %
NHP S Manual	14 (2)	10 (1)	29 (21)	21 (15)	32 (24)	23 (17)	27 (17)	16 (10)
NHP P Manual	8 (0)	6 (0)	7 (1)	6 (1)	14 (4)	12 (3)	2 (0)	2 (0)
NHP S Obs.	3 (3)	3 (3)	4 (4)	3 (3)	1 (1)	1 (1)	4 (4)	3 (3)
NHP P Obs.	0 (0)	1 (0)	1 (0)	1 (0)	1 (0)	1 (0)	2 (0)	3 (0)

EMG Analysis

Thee data used in the main text had corrupted EMG and we were unable to confirm whether the

NHPs were still using their arm muscles during the observation task, however, from observation

notes of the experiments, and the clear decrease in activity seen in Figs 3-4 of the main text, and

the fact that the NHPs could not reach for, or touch the force transducing handle, we do not

believe the NHPs were physically rehearsing grip movements during observation. In addition, as

seen below in Fig.S16, EMG on other days from these same NHPs was available during

observation trials and did show any obvious activation patterns in the mean activity, however, it

appears there was some small and insignificant increase in biceps for NHP P. The neural results

of these below data sets held to that shown in the main text and figs. This does not rule out the

NHPs covertly imagining movement, which was addressed in the main text.

Figure S16: EMG collected from the Biceps (section a and b) and Forearm (section c and d) of

NHP S (section a and c) and P (section b and d) recorded on different days. The blue color

represents manual blocks (execution) and red is for observation. On each of four sections a, b, c

and d, left plots show the mean EMG (0.5s pre-force onset to 1s post force onset activity) for all

the trials with standard deviation as shaded error bar. The right plots are showing the mean

power spectral density (PSD) with standard deviation for same data. A band pass filter was

applied on the EMG for the power line noise and its harmonics (30Hz, 60Hz, 90Hz… etc) and the

effect of that can be seen in the PSD plots (b). The p-value for the significance test between the

mean manual and observational is given on the title of each subplot.

Units significant for Extensor Muscle EMG:

Figure S13 shows, most recorded units are showing negative correlation with grip force and we

speculated that it could be a reason of they are connected to the extensor muscle group in the

forearm. To investigate further we wanted to observe units that are showing negative correlation

with force has any positive correlation with the EMG recorded from extensor muscle group. We

used manual data block recorded on another day for monkey S which had EMG recorded for the

extensor muscle from forearm. Unfortunately monkey P extensor EMG was not recorded on the

other days. For correlation EMG and spike data was taken from all trials and concatenated. We

used binned (100ms) spike data taken from 500ms pre force onset to 1000ms post and

measured the spearman rank correlation with corresponding binned envelope of the EMG data

from extensor muscle of forearm. The units that showed significance for force fit and a negative

correlation with the grip force were further investigated to observe their correlation with extensor

muscle EMG with the procedure described above. The percentage of such units that has positive

correlation with extensor muscle EMG and negative correlation with grip force for NHP S is

shown in figure S17. Figure S18 is showing the mean EMG and spike rate for an example unit

with the raster plot.

Figure S17: Percentage of with positive correlation between spike activity and extensor EMG

envelope from the units that showed negative correlation with grip force.

Figure S18: Example unit taken from NHP S, M1 cortex that shows positive correlation with the

Extensor EMG activity. The left plots show the mean EMG envelope (top) and the mean spike

activity (bottom) for all trials. The right raster plot shows spike information for all trials. The title

on the raster pot is showing the spearman rank correlation (Re) with the significance (Pe)

between binned (100ms) spike rate and extensor EMG done on concatenated data from all trials.

Significant units for Peak Grip force:

As force profiles were stereotypical smooth Gaussian-like profiles, we wanted to make sure our

regression model fits and predictions were meaningful, and not solely due to a unit having a

phasic response that could be used to fit the stereotypical waveform. Therefore, we determined

how neural activity correlated with the peak values of the Gaussian-shaped force profiles during

each trial. We considered force values around the peak force applied on each trial (three values

around the peak including the peak value) and we placed neural data, centered around each of

them, into ten 100ms non-overlapping bins from 500ms pre peak force to 500ms post peak force

neural activity. Each MN (figure 6 on the main manuscript) was tested for peak force significance

(F-test, BH corrected for number of population that is the number of corresponding units) by

fitting a linear regression model with peak force values and spike rate for each of the ten bins.

The blue bars are showing the units with Force-Mirror-neuron properties (Figure 6 inside main

manuscript). The red bars on figure S11 are showing significant units for peak grip force activity

among the force-mirror-neuron units.

Figure S19: Significant mirror neurons (MN) for grip force (blue) and peak grip force (red) for

Cued and uncued Manual (left subplots) and Observational (right subplots) block from NHP P

(bottom four subplots) and S (top four subplots). The blue bars are representing units significant

for grip force (F-test, BH corrected for number of population) and red bars are representing units

significant for peak grip force values (F-test, BH corrected for number of population). The

abbreviations on the subplot titles have the following meanings where Mc = Manual cued block,

Oc = Observation cued block, Mu = Manual uncued block, and Ou = Observation uncued block.

This plot is showing the mirror neuron units information from the Figure 6 of the main manuscript.

 Behavioral Summary:

 We used two NHP for our experiment, one male *Macaca* Radiata (NHP S) and one female

 Macaca Mulatta (NHP P). Both NHP performed manual and observational task. During the

 observational version of the task a plexiglass box restrained their hand from reaching the grip

 force sensor. For the manual version of the task they were able to reach the force sensor. During

 manual trials when they were cued to apply grip force, they reached to the grip force sensor and

 at the end of a trial they usually rested their arm on the base of the grip force sensor. NHP S

 achieved success rates of 77% (cued manual) and 82% (uncued manual) during the two manual-

 task blocks, while NHP P achieved rates of 58% (cued manual) and 72% (uncued manual). This

 comparatively low success rate for the manual block when they were cued for reward at the

 beginning shows that both NHP had an idea about the reward information. Between the two

 monkeys, performance of S during the trial was affected more by reward cue information

 compared to P. Figure S6 shows the force duration is significantly different between R0 and R1

 trials for NHP S which is not same for NHP P. Also, from figure S7 we can see the reaction time

 is significantly longer when NHP S completes R0 trials compared to R1 trials. Both results are

 indication that NHP S task performance was more dependent on reward compared to NHP P.

 One thing that should be mentioned that, for NHP S, although reaction time and grip force

 duration was significantly different the peak force applied on each trial was not significantly

 different (figure S11) since the applied grip force was supposed to be within a boundary value

 cued during task.

 Mirror neuron distribution from the array center:

 The position of the mirror neurons with respected to the electrode they were recorded from is

 shown on the Fig.2 on the main manuscript. To observe any non-uniform distribution of the

 mirror neurons with respect to the center of the array we applied Rayleigh test which used the

 angular position of all the mirror neurons from the center of the array and test the hypothesis if

the angles of the mirror neurons are uniformly distributed or not. For each cortex we applied the

test for both NHP individually and also we combined data from both NHP for each cortex and

applied the statistics again. The table S5 shows the p-value recorded from the statistical tests.

Only M1 cortex showed significant for all cases and the mean direction for M1 is toward (Fig. 2,

Main text) pre-motor cortex.

Table S5: the p-value from the Rayleigh test.

	S1	M1	PMd	PMv
NHP S	0.234	0.0036	0.7366	0.0595
NHP P	0.4617	0.0045	0.4104	
Combined	0.5714	1.48E-05	0.3726	

Similar motor behavior data analysis for NHP S:

Figure S6 and S8 show significant differences between force duration and reaction time for R0

and R1 trials for NHP S, MC data blocks. Note no such differences were noticed for NHP P. We

wanted to make sure that even if we only used the trials that have similar motor behavior during

R0 and R1 trials from MC block, the neural activity would still show significant reward related

differences for individual units. We pruned trials from MC data block (NHP S) that had higher

force duration (>2s) and reaction times (>1.5s) and the remining trials showed the difference

between these pruned R0 and R1 trials were not significantly different for both duration and

reaction times. See Fig.S20 figure legend for the results of this comparison.

Figure S20: Statistical results from MC block of NHP S before and after pruning trials for the

intention to use trials with similar motor behavior is given. The force duration and reaction time

for R0 and R1 trials are given in left column respectively. The number of significant units for

reward cue, reward grip force prediction and significant for reward outcome are indicated above

the respective plots.