Genomic insights of high-risk clones of ESBL-producing *Escherichia coli* isolated from community infections and commercial meat in Southern Brazil.

João Gabriel Material Soncini[1], Louise Cerdeira[2,3], Vanessa Lumi Koga[4], Ariane Tiemy Tizura[1], Gerson Nakazato[4], Renata Katsuko Takayama Kobayashi[4], Caio Augusto Martins Aires[5], Nilton Lincopan[6], Eliana Carolina Vespero[1*]

[1]Laboratory of Clinical Microbiology, Department of Pathology, Clinical and Toxicological Analysis, Health Sciences Center, State University of Londrina, Londrina, Brazil

[2]Department of Infectious Diseases, Central Clinical School, Monash University, Melbourne, Australia

[3]Department of Vector Biology, Liverpool School of Tropical Medicine, Liverpool, United Kingdom.

[4]Departament of Microbiology, Biological Science Center, State University of Londrina, Londrina, Brazil

[5]Department of Health Science, Health and Biological Science Center, Federal Rural University of Semi-Arid, Mossoró, Brazil.

[6]Department of Microbiology, Institute of Biomedical Sciences, University of São Paulo, Brazil.

*eliana.vespero@gmail.com

Keywords: Escherichia coli, EPEC, one health, WGS, food-producing, ST131, ST648, resistome.

ABSTRACT

During a microbiological and genomic surveillance study to investigate the molecular epidemiology of extended-spectrum beta-lactamase (ESBL)-producing *Escherichia coli* from community-acquired urinary tract infections (UTI) and commercial meat samples, in a Brazilian city with a high occurrence of infections by ESBL-producing bacteria, we have identified the presence of CTX-M (-55, -27, -24, -15, -14 and -2)-producing *E. coli* belonging to the international clones ST354, ST131, ST117, and ST38. The ST131 was more prevalent in human samples, and worryingly the high-risk ST131-C1-M27 was identified in human infections for the first time. We also detected CTX-M-55-producing *E. coli* ST117 isolates from meat samples (i.e., chicken and pork) and human infections. Moreover, we have identified the important clone CTX-M-24-positive *E. coli* ST354 from human samples in Brazil for the first time. In brief, our results suggest a potential of commercialized meat as a

reservoir of high-priority *E. coli* lineages in the community. In contrast, the identification of *E. coli* ST131-C1-M27 indicates that novel pandemic clones have emerged in Brazil, constituting a public health issue.

INTRODUCTION

Escherichia coli is a commensal of the human intestinal tract and most warm-blooded mammals and figures as an important pathogen for humans and animals[1,8]. In humans, urinary tract infection (UTI) is the second most common bacterial infection managed in primary care, and uropathogenic *E. coli* (UPEC) is responsible for 75% to 95% of the cases[1]. The increasing antimicrobial resistance (AMR) detected in clinical UPEC isolates has been of concern[1] and infections caused by antimicrobial-resistant bacteria as extended-spectrum β-lactamase (ESBL)-producing *E. coli* represent significant healthcare issues[8] since it compromises the effective treatment, being responsible for a large number of morbidity and mortality[4].

Since *E. coli* can act as a large reservoir of resistance genes that directly impact treatment in human and veterinary medicine, the debate over the transmission of multiresistant *E. coli* strains between animals and humans through numerous pathways has become increasingly important. However, the interaction between food-producing animals, humans, and the environment regarding the transmission of these resistant pathogens is not yet fully understood[8,9].

The isolation of ESBL-producing *E. coli* from food-production animals is increased worldwide, mostly from chicken meat[8,9]. The excessive use of antimicrobials in livestock is one of the practices that help in the emergence of pathogens resistant to humans. The consumption of meat, direct contact with colonized animals, or manure spread in the environment are sources for the transmission of livestock AMR to humans[5,6]. Besides that, AMR gene transfer may occur between different bacterial species in the gut of animals and humans[7].

The CTX-M type is one of the largest groups of ESBL, and recent studies that addressed the epidemiology of these enzymes in Brazil, show that CTX-M-2, CTX-M-8, CTX-M-9, and CTX-M-15 are the predominant variants in the country[10–12]. Many types of CTX-M-producing *E. coli* have been recognized as belonging to specific clones commonly isolated from UTI cases originating in a particular locale, country, or even globally. Some studies

show that isolates from foods CTX-M genotypes sometimes correspond with the locally

dominant human types[13,14].

Considering the emerging AMR in Brazil, both in human medicine as in livestock, and the

need for understanding this panorama, we conducted next-generation sequencing (NGS)-

based analysis adopting a One Health approach to assess national transmission of CTX-M-

producing *E. coli* isolated from meat products and human patients.

RESULTS

The results for each of 91 *E. coli* isolates included in this study can be seen in **Figure 1**. It is

notorious high rates of resistance to ampicillin (100%), ceftriaxone (87.91%), nalidixic acid

(87.91%), cefepime (83.52%), trimethoprim-sulfamethoxazole (82.42 %), nitrofurantoin

(76.92%), norfloxacin (75.82%) and ciprofloxacin (72.53%). Less than half showed

resistance to gentamicin (36.26%) and amoxicillin/clavulanate (21.98%). Only three (3.30%)

isolates were resistant to piperacillin-tazobactam and two (2.20%) to amikacin. Some isolates

also showed intermediate resistance levels: 28.57% to amoxicillin/clavulanate; 4.40% to

piperacillin-tazobactam and gentamicin; 1.10% to ciprofloxacin and norfloxacin.

Genomic analysis revealed 57 genes associated with resistance to aminoglycosides (n = 15),

β-lactams (n = 12), trimethoprim (n = 8), phenicols (n = 5), tetracyclines (n = 4), macrolides

(n = 4), sulfonamides (n = 3), quinolones (n = 3), lincosamides (n = 2) and fosfomycin (n =

1). Regarding aminoglycosides, the most prevalent genes were *strA* and *strB*, both with

39.56%, followed by the *aadA1* gene (36.26%). The *dfrA17* and *dfrA1* genes, associated with

resistance to trimethoprim, were detected in 31 (34.07%) and 13 (14.29%) isolates,

respectively. Genes related to phenicols resistance had similar prevalence, being *catB3*

(8.79%), *floR* (5.49%), *catA1* (4.40%) and *cmlA1* (4.40%). Concerning to tetracyclines

resistance, we detected the *tet(A)* (38.46%) and *tet(B)* (27.47%) genes. About macrolides the

mph(A) gene (29.67%) was found and the detected genes related to sulfonamides were *sul1*

(56.04%) and *sul2* (53.85%). Few isolates had lincosamide resistance genes, two of them had

Inu(F) (2.20%) and one *Inu(A)* (1.10%). The *fosA* gene found in three isolates was the only

one associated with fosfomycin resistance.

The genes associated with resistance to β-lactams were bla_{TEM-1B} (48.35%), bla_{OXA-1} (7.69%),

bla_{CMY-2} (6.59%), and bla_{TEM-1A} (2.20%), in addition to eight variants of the bla_{CTX-M} gene

that encode CTX-M-type ESBL enzymes. Among the ESBL coding genes, $bla_{CTX-M-55}$ was

the most detected (21.98%), mainly from chicken meats (n = 10), followed by humans (n = 6)

and porks (n = 4). The $bla_{CTX-M-15}$ was found predominantly in human isolates (n = 14) and

only in one pork isolate. On the other hand, $bla_{CTX-M-2}$ was also detected in 15 isolates

(16.48%), being them chicken meat (n = 7), human (n = 6) and pork (n = 2). The CTX-M-8

and CTX-M-14 coding genes were present in eight and five human isolates, and two and one

chicken meat isolates, respectively. The CTX-M-24 (n = 4), CTX-M27 (n = 3) and CTX-M-3

(n = 1) coding genes were present only in human isolates.

In this work, 52 plasmid incompatibility groups belong to the p0111, IncF, IncI1, and IncN

families. In human isolates, the most frequent pMLST were IncI1[ST-113] (n=9), IncF[F-: A-

: B-] (n=7), IncF[F1: A2: B20] (n=5), IncF[F48: A1: B49] (n=5) and p0111 (n=5). In chicken

meat isolates, IncF[F18: A-: B1] (n=8), p0111 (n=7) and IncN [Unknown ST] (n=5) were the

most frequent pMLST. In isolates of pork, the most frequent incompatibility groups were

IncN[Unknown ST] (n=4) and IncF[F33: A-: B1] (n=3).

In total, 40 sequence types (STs) were found, the most observed were the ST131 (n = 12),

ST38 (n = 8), ST648 (n = 7), and ST354 (n = 6). Some STs were detected in more than one

source, demonstrating a genetic relationship between these isolates, mainly between humans

and chicken meat. The ST38, ST131, ST354, and ST1196 were found in both urine and

chicken meat strains in the respective quantities of 5 and 3, 12 and 1, 4 and 3 and 1 and 1.

The ST410 was the only observed in urine (n = 1) and pork (n = 1) strains. The ST117 was

present in the three sources studied, with two strains from urine, one from chicken meat and

pork. The clonal relationship between the isolates in this study and the dissemination

distribution in Brazil can be seen in **Figure 2A-C**. Additionally, it is possible observed that

UK sample (ST131) clustered with other samples isolates in Brazil (ST131), all results could

be view in Microreact link

(https://microreact.org/project/2mKg54AHdWj5xdJ5VFejY8).

DISCUSSION

This study presents the first reports of *E. coli* ST131-C1-M27 in human infection and CTX-

M-24-positive *E. coli* ST354 from ITU, in Brazil. In Latin America CTX-M-producing *E.*

coli are endemic. Our data show a wide distribution of these isolates belonging to the

international clones in livestock and the community. The extensive presence of CTX-M

enzyme-producing strains in several sources raises the hypothesis that the spread occurs with

greater frequency and efficiency, especially among enterobacteria[10].

E. coli ST131 globally known and is related to the spread of resistance genes, including
specific CTX-M coding genes[15]. Recent studies have shown that ST131 is rare among animal
isolates, becoming almost exclusively a human pathogen, as demonstrated by our results,
where ST131 is predominantly found in strains of human urine[16]. The subclade C2 is
associated with $bla_{CTX-M-15}$ that can be carried by different groups of plasmids[17]. Here we also
observe that all $bla_{CTX-M-15}$ are involved with the incompatibility group IncF. In a study by
Peirano et al. (2020), it was shown that clade C was related to the highest rates of UTI, with
subclade C2 being the most common and associated with incompatibility group IncFII[18].
Besides, CTX-M-15-producing *E. coli* ST131 has already been shown to be involved in
outbreaks in health institutions and is the most prevalent ESBL-producing *E. coli*
worldwide[19].

The CTX-M-27-producing ST131-C1 has been considered a new epidemic clone, and there
have been no reports of human infections so far, in Brazil. Clade C1-M27 is associated with
CTX-M-27 and was first observed as colonizing children in France in 2012. Recent studies
suggest that the subclade C1-M27 was recently selected since SNPs have a smaller difference
between isolates of this same subclade than SNPs of isolates of subclade C2 and A. In
addition, the plasmid predominantly involved with the dissemination of $bla_{CTX-M-27}$ is
IncF[F1:A2:B20], as found in our study. Resistance to fluoroquinolones, macrolides,
tetracyclines, aminoglycosides, and sulfonamides appears to be part of the profile of C1-M27
isolates[20,21].

The CTX-M-14 and CTX-M-24 enzymes belong to the CTX-M-9 group. Although the first
one is widely distributed worldwide, especially in China, South-East Asia, Japan, South
Korea, and Spain, microorganisms producing CTX-M-24 remain relatively rare, reported
with greater incidence in countries such as Peru and Bolivia[18,22,23]. This study found an
important association between CTX-M-24 and *E. coli* ST354 detected in two human isolates,
never before reported in UTI in Brazil. In a study by Dagher et al. (2018), ST354 isolates
were positive to $bla_{CTX-M-24}$ and resistant to ciprofloxacin, associated with extra-intestinal
infections, animals and humans, reinforcing the zooanthroponotic hypothesis of these
clones[24].

ESBL type CTX-M-2 and CTX-M-55 are frequently found, and their coding genes are spread
in several ways. Some studies suggest that the plasmid IncF[F33: A-: B-] is involved in
disseminating these genes, which may explain the coexistence of these two genes in two
strains belonging to ST1725 isolated from urine samples[25]. Although another strain of *E. coli*

ST6448 isolated from chicken meat also showed the coexistence of $bla_{CTX-M-2}$ and $bla_{CTX-M-5}$,

the plasmids that carried them belonged to IncF [F24: A-: B73] and IncI1 [ST -131],

respectively. In the last ten years, IncI1-type plasmids have had a high spread, mainly in

animal reservoirs. There are reports of $bla_{CTX-M-2}$, $bla_{CTX-M-8}$ and $bla_{CTX-M-55}$ genes frequently

found on IncI plasmids from *E. coli* isolated from chickens and pigs several countries, such

as China, France, the United States of America, and the United Kingdom[26–28].

The international clone ST117, found in the three different sources of this study, is often

found in chicken meats and pork, and it is also associated with human infections. Studies

have already reported the multiple resistance profile of ST117 and associated it with CTX-M-

55 expression, consistent with our results[29,30]. Likewise, ST38 is also widely found in

chickens and humans, worldwide, and is related to several ESBL genes, such as $bla_{CTX-M-14}$,

$bla_{CTX-M-27}$, and $bla_{CTX-M-55}$. One of the hypotheses for the successful dissemination of these

genes among the *E. coli* clones is that the families of plasmids IncI1 and IncF are important

vectors for disseminating bla_{CTX-M}. In China, South Korea, and Japan, studies suggest an

epidemic of bla_{CTX-M} genes carried by plasmids IncI1, IncF[F33: A-: B-], IncF[F46: A-: B20]

and IncF[F18: A-: B1], found in cattle, pigs, chickens, pets and humans[31–33]. The second

hypothesis suggests that *E. coli* ST131 isolated in UK in 2001, could be the origin clone of *E.*

coli ST131 disseminated in Brazil, and may be after arrived in Brazil this clone acquired a

plasmid carrying $bla_{CTX-M-55}$ gene.

In conclusion, *E. coli* carrying bla_{CTX-M} genes from different sources seem to be related to the

spread of internationally known clones (ST354, ST131, ST117, ST38). Some clones

associated with some CTX-M variants are more prevalent in some sources than others do not

exclude the possibility that new clones are entering and establishing themselves in different

niches, as shown in this study. Thus, novel studies should continue to be carried out with

more samples and sources to understand further the dynamics of dissemination, shift, and

establishment of ESBL-producing *E. coli* clones at the interface between animal sources and

human health.

MATERIAL AND METHODS

Study population

During June 2016 to May 2019, 195.080 urine cultures were performed in a by public health

services, in a city in south of Brazil. A total of 34.293 (17,6%) were positive for gram-

positive or gram-negative microorganisms; of these 22.698 (66,2%) were E. coli strains and a

total of 2.033 (6,2%) ESBL producing bacteria, being 1.389 (51,2%) ESBL production *E.*

coli. Concomitantly, a surveillance study from January to May 2019 was carried out, to

research ESBL-producing *E. coli*, in chicken and pork meat, bought at markets and butcher

shop near public health services. Fluoroquinolone-resistant and ESBL-producing *E. coli* were

investigated in chicken meat (n = 50), and pork (n = 50) samples. A total of 102 *E. coli* was

isolated from chicken meat marketed, with 52 ESBL positive. And 67 resistant *E. coli* were

isolated in pigs, 31 ESBL positive. This study included for sequencing 91 *E. coli* strains of

102 total isolates: 59 isolated from urine culture(n=59), chicken meat (n=24) and pork (n=8).

These strains were selected by the similarity profile established by ERIC-PCR analysis of

1.389 ESBL-producing isolates. The study was approved by the Ethics and Research

Committee of the State University of Londrina CAAE 56869816.0.0000.5231.

Microbiological methods

Urine collected from women patients was inoculated on CHROMagar (Becton Dickinson,

Heidelberg, Germany) and MacConkey (Merck, Darmstadt, Germany) plates using a

calibrated inoculating loop with a capacity of 10 μl and incubated at 37°C for 24h.

The samples of chicken meat and pork were dipped in Brain Heart Infusion broth (Oxoid)

with cefotaxime (4μg / mL), ciprofloxacin (4μg / mL), and both (Sigma-Aldrich, Munich,

Germany) to selected resistant *E. coli* strains. After incubation, the solution was inoculated in

the same way used for urine samples. All the isolates were stored in Tryptic Soy Broth (TSB)

with 15% glycerol (−20◦C).

The identification and bacterial susceptibility were performed by the automated VITEK® 2

system, using the VITEK® 2 AST 239 card and the VITEK® 2 GN ID card (BioMérieux,

USA). The bacterial susceptibility was tested for 14 antibiotics: ampicillin,

amoxicillin/clavulanate, ceftriaxone, cefepime, ertapenem, meropenem, nalidixic acid,

ciprofloxacin, norfloxacin, gentamicin, amikacin nitrofurantoin, trimethoprim-

sulfamethoxazole, and piperacillin-tazobactam. The CLSI 2020 (Clinical and Laboratory

Standards Institute) criteria were used for interpretation. *E. coli* ATCC®25922 strain was

used as quality control.

ERIC-PCR

1.389 ESBL-producing isolates were subjected to Enterobacterial Repetitive Intergenic

Consensus (ERIC-PCR), by Versalovic et al. (1991)[34]. Analysis of genomic fingerprinting

was performed using GelJ v.2.0 software by the Dice similarity method (HERAS et al.,
2015)[35]. Strains were considered genetically related if the similarity index was ≥85 %.

DNA isolation and whole-genome sequencing

For DNA extraction, strains were grown on Mueller-Hinton Agar overnight at 37 °C.
Subsequently, a single colony was inoculated in 2 mL of Luria-Bertani broth for 12 hours at
37 °C. The suspension was used to continue extraction and purification by the DNA
extraction kit (Invitrogen, Carlsbad, CA). The extracted DNA was quantified by Qubit
dsDNA (double-stranded DNA) BR assay kit (Invitrogen, Carlsbad, CA). After
quantification, the DNA was used to construct a paired-end library (150 bp), sequenced using
the NextSeq platform (Illumina). The instructions of each manufacturer were followed in all
steps.

Bioinformatic analysis

Genome quality filter and assemblies were performed by the CLC Genomics Workbench
version 7.0 (Aarhus, Denmark). Multilocus sequence type (MLST), resistome, and virulome
were identified using MLST v2.0 (Larsen et al., 2012), ResFinder v3.1(Bortolaia et al.,
2020), VirulenceFinder v2.0, (Joensen et al., 2014), PlasmidFinder v2.1 (Carattoli et al.,
2014), FimTyper v1.0 (Roer et al., 2017) and SerotypeFinder v.2.0 (Joensen et al., 2015),
respectively. The BacMet database (Pal et al., 2013) was used to identify biocides and heavy
metal (HM)[31,36–42]. The EnteroBase (https://enterobase.warwick.ac.uk/) was used to create a
single nucleotide polymorphisms (SNPs) project to strains that showed the same STs
genomes were aligned against genomes of other Brazilian studies.

REFERENCES

1. Subashchandrabose, S. & Mobley, H. L. T. Virulence and Fitness Determinants of
Uropathogenic Escherichia coli. *Microbiol. Spectr.* **3**, 235–261. doi:
10.1128/microbiolspec.UTI-0015-2012 (2015).

4. Allcock, S. *et al.* Antimicrobial resistance in human populations: Challenges and
opportunities. *Glob. Heal. Epidemiol. Genomics* **2**. doi: 10.1017/gheg.2017.4 (2017).

5. Clifford, K. *et al.* Antimicrobial resistance in livestock and poor quality veterinary
medicines. *Bull. World Health Organ.* **96**, 662–664. doi: 10.2471/BLT.18.209585,
(2018).

6. Gozi, K. S. *et al.* Dissemination of multidrug-resistant commensal escherichia coli in
feedlot lambs in southeastern Brazil. *Front. Microbiol.* **10**. doi:

262    10.3389/fmicb.2019.01394 (2019).

7. Paphitou, N. I. Antimicrobial resistance: Action to combat the rising microbial challenges. *Int. J. Antimicrob. Agents* **42**, S25–S28. doi: 10.1016/j.ijantimicag.2013.04.007 (2013).

8. Poirel, L. *et al.* Antimicrobial Resistance in Escherichia coli. *Antimicrob. Resist. Bact. from Livest. Companion Anim.* 289–316. doi: 10.1128/microbiolspec.arba-0026-2017 (2018).

9. Reid, C. J., McKinnon, J. & Djordjevic, S. P. Clonal st131-h22 Escherichia coli strains from a healthy pig and a human urinary tract infection carry highly similar resistance and virulence plasmids. *Microb. Genomics* **5**. doi: 10.1099/mgen.0.000295 (2019).

10. Cyoia, P. S. *et al.* Distribution of ExPEC virulence factors, blaCTX-M, fosA3, and mcr-1 in Escherichia coli isolated from commercialized chicken carcasses. *Front. Microbiol.* **10**, 1–9. 10.3389/fmicb.2018.03254 (2019).

11. Daga, A. P. *et al.* Escherichia coli Bloodstream Infections in Patients at a University Hospital: Virulence factors and clinical characteristics. *Front. Cell. Infect. Microbiol.* **9**. doi: 10.3389/fcimb.2019.00191 (2019).

12. Irrgang, A. *et al.* CTX-M-15-producing E. coli isolates from food products in Germany are mainly associated with an IncF-type plasmid and belong to two predominant clonal E. coli lineages. *Front. Microbiol.* **8**. doi: 10.3389/fmicb.2017.02318 (2017).

13. Gonçalves, L. F. *et al.* Multidrug resistance dissemination by extended-spectrum β-lactamase-producing Escherichia coli causing community-acquired urinary tract infection in the Central-Western Region, Brazil. *J. Glob. Antimicrob. Resist.* **6**, 1–4. doi: 10.1016/j.jgar.2016.02.003 (2016).

14. Silva, N. *et al.* Genetic characterization of extended-spectrum β-lactamases in Escherichia coli isolated from retail chicken products including CTX-M-9 containing isolates: A food safety risk factor. *Br. Poult. Sci.* **53**, 747–755. doi: 10.1080/00071668.2012.740554 (2012).

15. Kondratyeva, K., Salmon-Divon, M. & Navon-Venezia, S. Meta-analysis of Pandemic Escherichia coli ST131 Plasmidome Proves Restricted Plasmid-clade Associations. *Sci. Rep.* **10**, 1–11. doi: 10.1038/s41598-019-56763-7 (2020).

16. Pitout, J. D. D. & DeVinney, R. Escherichia coli ST131: A multidrug-resistant clone primed for global domination. *F1000Research* **6**, 1–7. doi: 10.12688/f1000research.10609.1 (2017).

17. Coque, T. M. *et al.* Dissemination of clonally related E.coli strains expressing ESBL CTX-M-15. *Emerg. Infect. Dis.* **14**, 195–200. doi: 10.3201/eid1402.070350 (2008).

18. Peirano, G. & Pitout, J. D. D. Extended-Spectrum β-Lactamase-Producing Enterobacteriaceae: Update on Molecular Epidemiology and Treatment Options. *Drugs* **79**, 1529–1541. doi: 10.1007/s40265-019-01180-3 (2019).

19. Nicolas-Chanoine, M. H. *et al.* Intercontinental emergence of Escherichia coli clone O25:H4-ST131 producing CTX-M-15. *J. Antimicrob. Chemother.* **61**, 273–281. doi: 10.1093/jac/dkm464 (2008).

20. Birgy, A. *et al.* Acrossm of the Emergence and Dominance of Escherichia coli

304   Sequence. 1–10. doi: https://doi.org/10.1128/AAC .00382-19 (2019).

21.   Fernandes, M. R. *et al.* Emergence of CTX-M-27-producing Escherichia coli of ST131
and clade C1-M27 in an impacted ecosystem with international maritime traffic in
South America. *J. Antimicrob. Chemother.* **75**, 1647–1649. doi: 10.1093/jac/dkaa069
(2020).

22.   Song, J., Oh, S. S., Kim, J. & Shin, J. Extended-spectrum β-lactamase-producing
Escherichia coli isolated from raw vegetables in South Korea. *Sci. Rep.* **10**, 1–7. doi:
10.1038/s41598-020-76890-w (2020).

23.   Delgado, D. Y. C., Barrigas, Z. P. T., Astutillo, S. G. O., Jaramillo, A. P. A. & Ausili,
A. Detection and molecular characterization of β-lactamase genes in clinical isolates of
Gram-negative bacteria in Southern Ecuador. *Brazilian J. Infect. Dis.* **20**, 627–630.
doi: 10.1016/j.bjid.2016.07.001 (2016).

24.   Dagher, C. *et al.* Molecular characterization of carbapenem resistant *escherichia coli*
recovered from a tertiary hospital in Lebanon. *PLoS One* **13**, 1–13. doi:
10.1371/journal.pone.0203323 (2018).

25.   C., Cunha, M. P. V, Lincopan, N., Cerdeira, L. & Esposito, F. Coexistence of CTX-M-
2, CTX-M- 55, CMY-2, FosA3, and QnrB19 in extra-intestinal pathogenic Escherichia
coli from poultry in Brazil. *Antimicrob Agents Chemother.* 1–3 (2017).

26.   Song, J., Oh, S. S., Kim, J., Park, S. & Shin, J. Clinically Relevant Extended-Spectrum
β-Lactamase–Producing Escherichia coli Isolates From Food Animals in South Korea.
Front. Microbiol. **11**, 1–9. doi: 10.3389/fmicb.2020.00604 (2020).

27.   Castellanos, L. R. *et al.* High heterogeneity of Escherichia coli sequence types
harbouring ESBL/AmpC genes on IncI1 plasmids in the Colombian poultry chain.
PLoS One **12**, 1–15. doi: 10.1371/journal.pone.0170777 (2017).

28.   Ferreira, J. C., Penha Filho, R. A. C., Andrade, L. N., Berchieri Junior, A. & Darini, A.
L. C. Evaluation and characterization of plasmids carrying CTX-M genes in a non-
clonal population of multidrug-resistant Enterobacteriaceae isolated from poultry in
Brazil. *Diagn. Microbiol. Infect. Dis.* **85**, 444–448. doi:
10.1016/j.diagmicrobio.2016.05.011 (2016).

29.   Cormier, A. *et al.* Diversity of CTX-M-positive Escherichia coli recovered from
animals in Canada. *Vet. Microbiol.* **231**, 71–75. doi: 10.1016/j.vetmic.2019.02.031
(2019).

30.   Norizuki, C. *et al.* Detection of Escherichia coli producing CTX-M-1-group extended-
spectrum β -lactamases from pigs in Aichi Prefecture, Japan, between 2015 and 2016.
Jpn. J. Infect. Dis. **71**, 33–38. doi: 10.7883/yoken.JJID.2017.206 (2018).

31.   Lupo, A., Saras, E., Madec, J. Y. & Haenni, M. Emergence of bla CTX-M-55
associated with fosA, rmtB and mcr gene variants in Escherichia coli from various
animal species in France. *J. Antimicrob. Chemother.* **73**, 867–872. doi:
10.1093/jac/dkx489 (2018).

32.   Fernandes, M. R., Sellera, F. P., Moura, Q., Souza, T. A. & Lincopan, N. Draft
genome sequence of a CTX-M-8, CTX-M-55 and FosA3 co-producing Escherichia
coli ST117/B2 isolated from an asymptomatic carrier. *J. Glob. Antimicrob. Resist.* **12**,
183–184. doi: 10.1016/j.jgar.2018.01.015 (2018).

33. Xia, L. *et al.* Prevalence of ST1193 clone and IncI1/ST16 plasmid in E-coli isolates
carrying blaCTX-M-55 gene from urinary tract infections patients in China. *Sci. Rep.*
7, 1–8. doi: 10.1038/srep44866 (2017).

34. Versalovic, J., Koeuth, T., Lupski, J. R. & Plaza, O. B. Institute for Molecular
Genetics and department of Pediatrics , Baylor College of Medicine ,. *Cell* **19**, 6823–
6831 (1991).

35. Heras, J. *et al.* GelJ - a tool for analyzing DNA fingerprint gel images. *BMC*
Bioinformatics **16**, 1–8. doi: 10.1186/s12859-015-0703-0 (2015).

36. Larsen, M. V. *et al.* Multilocus sequence typing of total-genome-sequenced bacteria. *J.*
Clin. Microbiol. **50**, 1355–1361. doi: 10.1128/JCM.06094-11 (2012).

37. Bortolaia, V. *et al.* ResFinder 4.0 for predictions of phenotypes from genotypes. *J.*
Antimicrob. Chemother. **75**, 3491–3500. doi: 10.1093/jac/dkaa345 (2020).

38. Kleinheinz, K. A., Joensen, K. G. & Larsen, M. V. Applying the ResFinder and
VirulenceFinder web-services for easy identification of acquired antibiotic resistance
and E. coli virulence genes in bacteriophage and prophage nucleotide sequences.
Bacteriophage **4**, e27943. doi: 10.4161/bact.27943 (2014).

39. Carattoli, A. *et al.* In Silico detection and typing of plasmids using plasmidfinder and
plasmid multilocus sequence typing. *Antimicrob. Agents Chemother.* **58**, 3895–3903.
doi: 10.1128/AAC.02412-14 (2014).

40. Roer, L. *et al.* Development of a web tool for Escherichia coli subtyping based on fimh
alleles. *J. Clin. Microbiol.* **55**, 2538–2543. doi: 10.1128/JCM.00737-17 (2017).

41. Joensen, K. G., Tetzschner, A. M. M., Iguchi, A., Aarestrup, F. M. & Scheutz, F.
Rapid and easy in silico serotyping of Escherichia coli isolates by use of whole-
genome sequencing data. *J. Clin. Microbiol.* **53**, 2410–2426. doi: 10.1128/JCM.00008-
15 (2015).

42. Pal, C., Bengtsson-Palme, J., Rensing, C., Kristiansson, E. & Larsson, D. G. J.
BacMet: Antibacterial biocide and metal resistance genes database. *Nucleic Acids Res.*
42, 737–743 doi: 10.1093/nar/gkt1252 (2014).

ACKNOWLEDGEMENTS
The authors would like to thank the financial support of the Bill and Melinda Gates
Foundation's Grand Challenges Explorations Brazil – New Approaches to characterize the
global burden of antimicrobial resistance (OPP1193112), the Research Support Facilities
Center of the State University of São Paulo (CEFAP-USP), Institute of Biomedical Sciences
of State University of São Paulo (ICB-USP) and Master's program Clinical and Laboratory
Pathophysiology of the State University of Londrina.

AUTHOR INFORMATION

Affiliations

Laboratory of Clinical Microbiology, Department of Pathology, Clinical and Toxicological Analysis, Health Sciences Center, State University of Londrina, Londrina, Brazil

João Gabriel Material Soncini

Department of Infectious Diseases, Central Clinical School, Monash University, Melbourne, Australia

Department of Vector Biology, Liverpool School of Tropical Medicine, Liverpool, United Kingdom.

Louise Cerdeira

Department of Microbiology, Biological Science Center, State University of Londrina, Londrina, Brazil

Vanessa Lumi Koga

Laboratory of Clinical Microbiology, Department of Pathology, Clinical and Toxicological Analysis, Health Sciences Center, State University of Londrina, Londrina, Brazil

Ariane Tiemy Tizura

Department of Microbiology, Biological Science Center, State University of Londrina, Londrina, Brazil

Gerson Nakazato

Department of Microbiology, Biological Science Center, State University of Londrina, Londrina, Brazil

Renata Katsuko Takayama Kobayashi

Department of Health Science, Health and Biological Science Center, Federal Rural University of Semi-Arid, Mossoró, Brazil.

Caio Augusto Martins Aires

Department of Microbiology, Institute of Biomedical Sciences, University of São Paulo, Brazil

Nilton Lincopan

Laboratory of Clinical Microbiology, Department of Pathology, Clinical and Toxicological Analysis, Health Sciences Center, State University of Londrina, Londrina, Brazil

Eliana Carolina Vespero

Data availability

Draft whole-genome assembly was deposited in GenBank under the bioproject PRJNA578368. The data of the figures can be accessed in Figshare (https://doi.org/10.6084/m9.figshare.12808439.v1).

ETHICS DECLARATIONS
Competing interests
The authors declare no competing interests.

Ethical approval
The study was approved by the Ethics and Research Committee of the State University of Londrina CAAE 56869816.0.000.5231.

Figure 01 – Heatmap show AMR profile, ST and Resistome.

Figure 02 – (A) *E. coli* Phylogenomic SNP tree with circular heatmap shows source type (first inner circle), sequence type (second inner circle) and ESBL-gene (third inner circle. (B) *E. coli* Phylogenomic SNP tree with columns shows ESBL-gene (column 1) and source type (column 2); light pink highlights the clade where UK-ST131 samples clustered with Brazilian ST131 samples. (C) dissemination distribution map in Brazil.