1    Structural insights into Cullin4-RING ubiquitin ligase remodelling by Vpr from

2                           simian immunodeficiency viruses

Sofia Banchenko[1¶], Ferdinand Krupp[1¶], Christine Gotthold[1], Jörg Bürger[1,2], Andrea Graziadei[3], Francis

O'Reilly[3], Ludwig Sinn[3], Olga Ruda[1], Juri Rappsilber[3,4], Christian M. T. Spahn[1], Thorsten Mielke[2], Ian

A. Taylor[5], David Schwefel[1*]

[1] Institute of Medical Physics and Biophysics, Charité – Universitätsmedizin Berlin, corporate member

of Freie Universität Berlin, Humboldt-Universität zu Berlin, and Berlin Institute of Health, Berlin,

Germany

[2] Microscopy and Cryo-Electron Microscopy Service Group, Max-Planck-Institute for Molecular

Genetics, Berlin, Germany

[3] Bioanalytics Unit, Institute of Biotechnology, Technische Universität Berlin, Berlin, Germany

[4] Wellcome Centre for Cell Biology, University of Edinburgh, Edinburgh, United Kingdom

[5] Macromolecular Structure Laboratory, The Francis Crick Institute, London, United Kingdom

*Corresponding author

E-mail: david.schwefel@charite.de (DS)

¶These authors contributed equally to this work

## Abstract

Viruses have evolved means to manipulate the host's ubiquitin-proteasome system, in order to down-regulate antiviral host factors. The Vpx/Vpr family of lentiviral accessory proteins usurp the substrate receptor DCAF1 of host Cullin4-RING ligases (CRL4), a family of modular ubiquitin ligases involved in DNA replication, DNA repair and cell cycle regulation. CRL4$^{DCAF1}$ specificity modulation by Vpx and Vpr from certain simian immunodeficiency viruses (SIV) leads to recruitment, poly-ubiquitylation and subsequent proteasomal degradation of the host restriction factor SAMHD1, resulting in enhanced virus replication in differentiated cells. To unravel the mechanism of SIV Vpr-induced SAMHD1 ubiquitylation, we conducted integrative biochemical and structural analyses of the Vpr protein from SIVs infecting *Cercopithecus cephus* (SIV$_{mus}$). X-ray crystallography reveals commonalities between SIV$_{mus}$ Vpr and other members of the Vpx/Vpr family with regard to DCAF1 interaction, while cryo-electron microscopy and cross-linking mass spectrometry highlight a divergent molecular mechanism of SAMHD1 recruitment. In addition, these studies demonstrate how SIV$_{mus}$ Vpr exploits the dynamic architecture of the multi-subunit CRL4$^{DCAF1}$ assembly to optimise SAMHD1 ubiquitylation. Together, the present work provides detailed molecular insight into variability and species-specificity of the evolutionary arms race between host SAMHD1 restriction and lentiviral counteraction through Vpx/Vpr proteins.

## Author summary

Due to the limited size of virus genomes, virus replication critically relies on host cell components. In addition to the host cell's energy metabolism and its DNA replication and protein synthesis apparatus, the protein degradation machinery is an attractive target for viral re-appropriation. Certain viral factors divert the specificity of host ubiquitin ligases to antiviral host factors, in order to mark them for destruction by the proteasome, to lift intracellular barriers to virus replication. Here, we present molecular details of how the simian immunodeficiency virus accessory protein Vpr interacts with a substrate receptor of host Cullin4-RING ubiquitin ligases, and how this interaction redirects the specificity of Cullin4-RING to the antiviral host factor SAMHD1. The studies uncover the mechanism of Vpr-induced SAMHD1 recruitment and subsequent ubiquitylation. Moreover, by comparison to related accessory proteins from other immunodeficiency virus species, we illustrate the surprising variability in the molecular strategies of SAMHD1 counteraction, which these viruses adopted during evolutionary adaptation to their hosts. Lastly, our work also provides deeper insight into the inner workings of the host's Cullin4-RING ubiquitylation machinery.

## Introduction

A large proportion of viruses have evolved means to co-opt their host's ubiquitylation machinery, in order to improve replication conditions, either by introducing viral ubiquitin ligases and deubiquitinases, or by modification of host proteins involved in ubiquitylation [1-3]. In particular, host ubiquitin ligases are a prominent target for viral usurpation, to redirect specificity towards antiviral host restriction factors. This results in recruitment of restriction factors as non-endogenous *neo*-substrates, inducing their poly-ubiquitylation and subsequent proteasomal degradation [4-8]. This counteraction of the host's antiviral repertoire is essential for virus infectivity and spread [9-12], and mechanistic insights into these specificity changes extend our understanding of viral pathogenesis and might pave the way for novel treatments.

Frequently, virally encoded modifying proteins associate with, and adapt the Cullin4-RING ubiquitin ligases (CRL4) [5]. CRL4 consists of a Cullin4 (CUL4) scaffold that bridges the catalytic RING-domain subunit ROC1 to the adaptor protein DDB1, which in turn binds to exchangeable substrate receptors (DCAFs, DDB1- and CUL4-associated factors) [13-17]. In some instances, the DDB1 adaptor serves as an anchor for virus proteins, which then act as "viral DCAFs" to recruit the antiviral substrate. Examples are the simian virus 5 V protein and mouse cytomegalovirus M27, which bind to DDB1 and recruit STAT1/2 proteins for ubiquitylation, in order to interfere with the host's interferon response [18-20]. Similarly, CUL4-dependent downregulation of STAT signalling is important for West Nile Virus replication [21]. In addition, the hepatitis B virus X protein hijacks DDB1 to induce proteasomal destruction of the structural maintenance of chromosome (SMC) complex to promote virus replication [22, 23].

Viral factors also bind to and modify DCAF receptors in order to redirect them to antiviral substrates. Prime examples are the lentiviral accessory proteins Vpr and Vpx. All contemporary human and simian immunodeficiency viruses (HIV/SIV) encode Vpr, while only two lineages, represented by HIV-2 and SIV infecting mandrills, carry Vpx [24]. Vpr and Vpx proteins are packaged into progeny virions and released into the host cell upon infection, where they bind to DCAF1 in the nucleus [25]. In this work, corresponding simian immunodeficiency virus Vpx/Vpr proteins will be indicated with their host species as subscript, with the following abbreviations used: mus – moustached monkey (*Cercopithecus*

*cephus*), mnd – mandrill (*Mandrillus sphinx*), rcm – red-capped mangabey (*Cercocebus torquatus*), sm

– sooty mangabey (*Cercocebus atys*), deb – De Brazza's monkey (*Cercopithecus neglectus*), syk –

Syke's monkey (*Cercopithecus albogularis*), agm – african green monkey (*Chlorocebus* spec).

$Vpr_{HIV-1}$ is important for virus replication *in vivo* and in macrophage infection models [26]. Recent

proteomic analyses revealed that DCAF1 specificity modulation by $Vpr_{HIV-1}$ proteins results in down-

regulation of hundreds of host proteins in a DCAF1- and proteasome-dependent manner [27], including

the previously reported $Vpr_{HIV-1}$ degradation targets UNG2 [28], HLTF [29], MUS81 [30, 31], MCM10

[32] and TET2 [33]. This surprising promiscuity in degradation targets is also partially conserved in

more distant clades exemplified by $Vpr_{agm}$ and $Vpr_{mus}$ [27]. However, Vpr pleiotropy, and the lack of

easily accessible experimental models, have prevented a characterisation of how these degradation

events precisely promote replication [26].

By contrast, Vpx, exhibits a much narrower substrate range. It has recently been reported to target

stimulator of interferon genes (STING) and components of the human silencing hub (HUSH) complex

for degradation, leading to inhibition of antiviral cGAS-STING-mediated signalling and reactivation of

latent proviruses, respectively [34-36]. Importantly, Vpx also recruits the SAMHD1 restriction factor to

DCAF1, in order to mark it for proteasomal destruction [37, 38]. SAMHD1 is a deoxynucleotide

triphosphate (dNTP) triphosphohydrolase that restricts retroviral replication in non-dividing cells by

lowering the dNTP pool to levels that cannot sustain viral reverse transcription [39-46]. Retroviruses

that express Vpx are able to alleviate SAMHD1 restriction, allowing for replication in differentiated

myeloid lineage cells, resting T cells and memory T cells [38, 47, 48]. As a result of the constant

evolutionary arms race between the host's SAMHD1 restriction and its viral antagonist Vpx, the

mechanism of Vpx-mediated SAMHD1 recruitment is highly virus species- and strain-specific: The

Vpx clade represented by $Vpx_{HIV-2}$ recognises the SAMHD1 C-terminal domain (CtD), while $Vpx_{mnd2/rcm}$

binds the SAMHD1 N-terminal domain (NtD) in a fundamentally different way [24, 49-52].

In the course of evolutionary adaptation to their primate hosts, and due to selective pressure to evade

SAMHD1 restriction, two groups of SIVs that do not have Vpx, $SIV_{agm}$, and $SIV_{deb/mus/syk}$, *neo-*

functionalised their Vpr to bind SAMHD1 and induce its degradation [24, 49, 53]. Consequently, these

species evolved "hybrid" Vpr proteins that retain targeting of some host factors depleted by HIV-1-type

Vpr [27], and additionally induce SAMHD1 degradation.

To uncover the molecular mechanisms of DCAF1- and SAMHD1-interaction of such a "hybrid" Vpr,

we initiated integrative biochemical and structural analyses of the Vpr protein from an SIV infecting

*Cercopithecus cephus*, $Vpr_{mus}$. These studies reveal similarities and differences to Vpx and Vpr proteins

from other lentivirus species and pinpoint the divergent molecular mechanism of $Vpr_{mus}$-dependent

SAMHD1 recruitment to CUL4/ROC1/DDB1/DCAF1 ($CRL4^{DCAF1}$). Furthermore, cryo-electron

microscopic (cryo-EM) reconstructions of a $Vpr_{mus}$-modified $CRL4^{DCAF1}$ protein complex allow for

insights into the structural plasticity of the entire CRL4 ubiquitin ligase assembly, with implications for

the ubiquitin transfer mechanism.

**Results**

**SAMHD1-CtD is necessary and sufficient for $Vpr_{mus}$-binding and ubiquitylation in vitro**

To investigate the molecular interactions between $Vpr_{mus}$, the *neo*-substrate SAMHD1 from rhesus

macaque and CRL4 subunits DDB1/DCAF1 C-terminal domain (DCAF1-CtD), protein complexes were

reconstituted *in vitro* from purified components and analysed by gel filtration (GF) chromatography.

The different protein constructs that were employed are shown schematically in S1A Fig. In the absence

of additional binding partners, $Vpr_{mus}$ is insoluble after removal of the GST affinity purification tag (S1B

Fig) and accordingly could not be applied to the GF column. No interaction of SAMHD1 with

DDB1/DCAF1-CtD could be detected in the absence of $Vpr_{mus}$ (S1C Fig). Analysis of binary protein

combinations ($Vpr_{mus}$ and DDB1/DCAF1-CtD; $Vpr_{mus}$ and SAMHD1) shows that $Vpr_{mus}$ elutes together

with DDB1/DCAF1-CtD (S1D Fig) or with SAMHD1 (S1E Fig). Incubation of $Vpr_{mus}$ with

DDB1/DCAF1B and SAMHD1 followed by GF resulted in co-elution of all three components (Fig 1A).

Together, these results show that $Vpr_{mus}$ forms stable binary and ternary protein complexes with

DDB1/DCAF1-CtD and/or SAMHD1 *in vitro*. Furthermore, incubation with any of these interaction

partners apparently stabilises $Vpr_{mus}$ by alleviating its tendency for aggregation/insolubility.

Previous cell-based assays indicated that residues 583-626 of rhesus macaque SAMHD1 (SAMHD1-

CtD) are necessary for $Vpr_{mus}$-induced proteasomal degradation [49]. To test this finding in our *in vitro*

system, constructs containing SAMHD1-CtD fused to T4 lysozyme (T4L-SAMHD1-CtD), or

containing only the N-terminal domains of SAMHD1, and lacking SAMHD1-CtD (SAMHD1-ΔCtD),

were incubated with $Vpr_{mus}$ and DDB1/DCAF1-CtD, and complex formation was assessed by GF

chromatography. Analysis of the resulting chromatograms by SDS-PAGE shows that SAMHD1-ΔCtD

did not co-elute with DDB1/DCAF1-CtD/$Vpr_{mus}$ (Fig 1B). By contrast, T4L-SAMHD1-CtD

accumulated in the same elution peak as DDB1/DCAF1-CtD and $Vpr_{mus}$ (Fig 1C). These results confirm

that SAMHD1-CtD is necessary for stable association with DDB1/DCAF1-CtD/$Vpr_{mus}$ *in vitro*, and

demonstrate that SAMHD1-CtD is sufficient for $Vpr_{mus}$-mediated recruitment of the T4L-SAMHD1-

CtD fusion construct to DDB1/DCAF1-CtD.

To correlate these data with enzymatic activity, *in vitro* ubiquitylation assays were conducted by

incubating SAMHD1, SAMHD1-ΔCtD or T4L-SAMHD1-CtD with purified CRL4$^{DCAF1-CtD}$, E1

(UBA1), E2 (UBCH5C), ubiquitin and ATP. Input proteins are shown in S2A Fig, and control reactions

in S2B, C Fig. In the absence of $Vpr_{mus}$, no SAMHD1 ubiquitylation was observed (Figs 1D and S2D),

while addition of $Vpr_{mus}$ resulted in robust SAMHD1 ubiquitylation, as demonstrated by an upward shift

of SAMHD1 in the SDS PAGE analysis, induced by covalent modification with increasingly more

ubiquitin molecules, leading to almost complete loss of the band corresponding to unmodified SAMHD1

after 15 min incubation (Figs 1E and S2E). In agreement with the analytical GF data, SAMHD1-ΔCtD

was not ubiquitylated in the presence of $Vpr_{mus}$ (Figs 1F and S2F). By contrast, T4L-SAMHD1-CtD was

efficiently ubiquitylated, resulting in >90% loss of the band corresponding to unmodified T4L-

SAMHD1-CtD after 15 min (Figs 1G and S2F). Again, these data substantiate the functional importance

of SAMHD1-CtD for $Vpr_{mus}$-mediated recruitment to the CRL4$^{DCAF1}$ ubiquitin ligase.

**Crystal Structure analysis of apo- and $Vpr_{mus}$-bound DDB1/DCAF1-CtD protein**

**complexes**

To obtain structural information regarding $Vpr_{mus}$ and its mode of binding to the CRL4 substrate receptor

DCAF1, the X-ray crystal structures of a DDB1/DCAF1-CtD complex, and DDB1/DCAF1-CtD/T4L-

$Vpr_{mus}$ (residues 1-92) fusion protein ternary complex were determined. The structures were solved

using molecular replacement and refined to resolutions of 3.1 Å and 2.5 Å respectively (S1 Table).

Vpr$_{mus}$ adopts a three-helix bundle fold, stabilised by coordination of a zinc ion by His and Cys residues

on Helix-1 and at the C-terminus (Fig 2A). Superposition of Vpr$_{mus}$ with previously determined Vpx$_{sm}$

[50], Vpx$_{mnd2}$ [51, 52], and Vpr$_{HIV-1}$ [54] structures reveals a conserved three-helix bundle fold, and

similar position of the helix bundles on DCAF1-CtD (S3A Fig). In addition, the majority of side chains

involved in DCAF1-interaction are type-conserved in all Vpx and Vpr proteins (Figs S3B-G and S6A),

strongly suggesting a common molecular mechanism of host CRL4-DCAF1 hijacking by the Vpx/Vpr

family of accessory proteins. However, there are also significant differences in helix length and register

as well as conformational variation in the loop region N-terminal of Helix-1, at the start of Helix-1 and

in the loop between Helices-2 and -3 (S3A Fig).

Vpr$_{mus}$ binds to the side and on top of the disk-shaped 7-bladed β-propeller (BP) DCAF1-CtD domain

with a total contact surface area of ~1600 Å$^2$ comprising three major regions of interaction. The extended

Vpr$_{mus}$ N-terminus attaches to the cleft between DCAF1 BP blades 1 and 2 through several hydrogen

bonds, electrostatic and hydrophobic interactions (S3B-D Fig). A second, smaller contact area is formed

by hydrophobic interaction between Vpr$_{mus}$ residues L31 and E34 from Helix-1, and DCAF1 W1156,

located in a loop on top of BP blade 2 (S3E Fig). The third interaction surface comprises the C-terminal

half of Vpr$_{mus}$ Helix-3, which inserts into a ridge on top of DCAF1 (S3F, G Fig).

Superposition of the apo-DDB1/DCAF1-CtD and Vpr$_{mus}$-bound crystal structures reveals

conformational changes in DCAF1 upon Vpr$_{mus}$ association. Binding of the N-terminal arm of Vpr$_{mus}$

induces only a minor rearrangement of a loop in BP blade 2 (S3C Fig). By contrast, significant structural

changes occur on the upper surface of the BP domain: polar and hydrophobic interactions of DCAF1

residues P1329, F1330, F1355, N1371, L1378, M1380 and T1382 with Vpr$_{mus}$ side chains of T79, R83,

R86 and E87 in Helix-3 result in the stabilisation of the sequence stretch that connect BP blades 6 and

7 ("C-terminal loop", Figs 2B and S3F). Moreover, side chain electrostatic interactions of Vpr$_{mus}$

residues R15, R75 and R76 with DCAF1 E1088, E1091 and E1093 lock the conformation of an "acidic

loop" upstream of BP blade 1, which is also unstructured and flexible in the absence of Vpr$_{mus}$ (Figs 2B,

C and S3D, F).

Notably, in previously determined structures of Vpx/DCAF1/SAMHD1 complexes the "acidic loop" is

a central point of ternary contact, providing a binding platform for positively charged amino acid side

chains in either the SAMHD1 N- or C-terminus [50-52]. For example, $Vpx_{sm}$ positions SAMHD1-CtD

in such a way, that SAMHD1 K622 engages in electrostatic interaction with the DCAF1 "acidic loop"

residue D1092 (Fig 2C, left panel). However, in the $Vpr_{mus}$ crystal structure the bound $Vpr_{mus}$ now blocks

access to the corresponding SAMHD1-CtD binding pocket, in particular by the positioning of an

extended N-terminal loop that precedes Helix-1. Additionally, $Vpr_{mus}$ side chains R15, R75 and R76

neutralise the DCAF1 "acidic loop", precluding the formation of further salt bridges to basic residues in

SAMHD1-CtD (Fig 2C, right panel).

To validate the importance of $Vpr_{mus}$ residues R15 and R75 for DCAF1-CtD-binding, charge reversal

mutations to glutamates were generated by site-directed mutagenesis. The circular dichroism (CD)

spectrum of the $Vpr_{mus}$ R15E R75E double mutant GST-fusion protein was identical to the wild type,

indicating similar secondary structure content and thus no major structural disturbances caused by the

amino acid substitutions (S3H Fig). The effect of the $Vpr_{mus}$ R15E R75E double mutant on complex

assembly was then analysed by GF chromatography. SDS-PAGE analysis of the resulting

chromatographic profile shows an almost complete loss of the DDB1/DCAF1-CtD/$Vpr_{mus}$/SAMHD1

complex peak (Fig 2D, fraction 6), when compared to the wild type, concomitant with enrichment of (i)

some proportion of $Vpr_{mus}$ R15E R75E-bound DDB1/DCAF1-CtD (Fig 2D, fraction 7), (ii) free

DDB1/DCAF1-CtD (fraction 7-8), and of (iii) $Vpr_{mus}$ R15E R75E/SAMHD1 binary complex (Fig 2D,

fraction 8-9). This suggests that charge reversal of $Vpr_{mus}$ side chains R15 and R75 weakens the strong

association with DCAF1 observed in wild type $Vpr_{mus}$, due to loss of electrostatic interaction with the

"acidic loop", in accordance with the crystal structure. Consequently, some proportion of Vpr-bound

SAMHD1 dissociates. This notion is further supported by GF analysis of binary combinations of the

$Vpr_{mus}$ R15E R75E double mutant with either SAMHD1 or DDB1/DCAF1-CtD. Incubation of $Vpr_{mus}$

R15E R75E with SAMHD1 followed by GF leads to co-elution of both proteins, concomitant with a

shift of the elution peak towards higher apparent molecular weight, compared to SAMHD1 alone (Fig.

2E, fractions 8-9). By contrast, incubation of the $Vpr_{mus}$ double mutant with DDB1/DCAF1-CtD does

not change the elution volume of the DDB/DCAF1-CtD species, and no band corresponding to Vpr can

be detected in the SDS-PAGE analysis of the corresponding fractions (Fig. 2F, fractions 7-8). These

data clearly demonstrate loss of interaction with DDB1/DCAF1-CtD upon charge reversal of $Vpr_{mus}$

residues 15 and 75, while the SAMHD1-binding activity is retained.

**Cryo-EM analysis of CRL4-NEDD8$^{DCAF1-CtD}$/$Vpr_{mus}$/SAMHD1 conformational states and**

**dynamics**

To obtain structural insight into $Vpr_{mus}$ in the context of a complete CRL4 assembly, and to understand

the SAMHD1 recruitment mechanism, we initiated cryo-EM analyses of the

CRL4$^{DCAF1-CtD}$/$Vpr_{mus}$/SAMHD1 assembly. In these studies, the small ubiquitin-like protein NEDD8

was enzymatically attached to the CUL4 subunit, in order to obtain its active form (S4A Fig) [55]. A

CRL4-NEDD8$^{DCAF1-CtD}$/$Vpr_{mus}$/SAMHD1 complex was reconstituted *in vitro* and purified by GF

chromatography (S4B Fig). 2D classification of the resulting particle images revealed considerable

compositional and conformational heterogeneity, especially regarding the presence and position of the

CUL4-NEDD8/ROC1 sub-complex (stalk) relative to DDB1/DCAF1/$Vpr_{mus}$ (core) (S4C Fig).

Two consecutive rounds of 3D classification yielded three particle populations, resulting in 3D

reconstructions at 8-10 Å resolution, which contained both the $Vpr_{mus}$-bound CRL4 core and the stalk

(conformational states-1, -2 and -3, Figs 3A and S4D-F). The quality of the 3D volumes was sufficient

to fit crystallographic models of core (Fig 2) and the stalk (PDB 2hye) [15] as rigid bodies (Figs 3B and

S4G). For the catalytic RING-domain subunit ROC1, only fragmented electron density was present near

the position it occupies in the crystallographic model (S4G Fig). In all three states, electron density was

selectively absent for the C-terminal CUL4 winged helix B (WHB) domain (residues 674-759), which

contains the NEDD8 modification site (K705), and for the preceding α-helix, which connects the CUL4

N-terminal domain to the WHB domain (S4G Fig). In accordance with this observation, the positions

of CRL5-attached NEDD8 and of the CRL4 ROC1 RING domain are sterically incompatible upon

superposition of their respective crystal structures (S4H Fig) [56].

Alignment of 3D volumes from states-1, -2 and -3 shows that core densities representing DDB1 BPA,

BPC, DCAF1-CtD and $Vpr_{mus}$ superimpose well, indicating that these components do not undergo major

conformational fluctuations and thus form a rigid platform for substrate binding and attachment of the

CRL4 stalk (Fig 3). However, rotation of DDB1 BPB around a hinge connecting it to BPC results in

three different orientations of state-1, -2 and -3 stalk regions relative to the core. BPB rotation angles

were measured as 69° between state-1 and -2, and 50° between state-2 and -3.

These data are in line with previous prediction based on extensive comparative crystal structure

analyses, which postulated an approx. 150° rotation of the CRL4 stalk around the core [13, 15, 16, 19,

57]. However, the left- and rightmost CUL4 orientations observed here, states-1 and -3 from our cryo-

EM analysis, indicate a slightly narrower stalk rotation range (119°), when compared to the outermost

stalk conformations modelled from previously determined crystal structures (143°) (Fig 3C). A possible

explanation for this discrepancy arises from inspection of the cryo-EM densities and fitted models,

revealing that along with the main interaction interface on DDB1 BPB there are additional molecular

contacts between CUL4 and DDB1. Specifically, in state-1, there is a contact between the loop

connecting helices D and E of CUL4 cullin repeat (CR)1 (residues 161-169) and a loop protruding from

BP blade 3 of DDB1 BPC (residues 795-801, S4I Fig). In state-3, the loop between CUL4 CR2 helices

D and E (residues 275-282) abuts a region in the C-terminal helical domain of DDB1 (residues 1110-

1127, S4J Fig). These auxiliary interactions might be required to lock the outermost stalk positions

observed here in order to confine the rotation range of CUL4.

**Molecular mechanism of SAMHD1-targeting**

A reanalysis of the cryo-EM data involving template-based particle picking and extensive 3D

classification allowed for separation of an additional homogeneous particle population (S5A, B Fig).

This subset of particle images yielded a 3D reconstruction at a nominal resolution of 7.3 Å that only

contained electron density corresponding to the CRL4 core (S5A-D Fig). Molecular models of DDB1

BP domains A and C (BPA, BPC), DCAF1-CtD and Vpr$_{mus}$, derived from our crystal structure (Fig 2),

could be fitted as rigid bodies into this cryo-EM volume (Fig 4A). No obvious electron density was

visible for the bulk of SAMHD1. However, close inspection revealed an additional tubular, slightly

arcing density feature, approx. 35 Å in length, located on the upper surface of the Vpr$_{mus}$ helix bundle,

approximately 17 Å away from and opposite of the Vpr$_{mus}$/DCAF1-CtD binding interface (Fig 4A, red

arrows). One end of the tubular volume contacts the middle of Vpr$_{mus}$ Helix-1, and the other end forms

additional contacts to the C-terminus of Helix-2 and the N-terminus of Helix-3 (Fig 4B). A local

resolution of 7.5-8 Å (S5C Fig) precluded the fitting of an atomic model. Considering the biochemical

data, showing that SAMHD1-CtD is sufficient for recruitment to DDB1/DCAF1/Vpr$_{mus}$, we hypothesise

that this observed electron density feature corresponds to a region of SAMHD1-CtD which physically

interacts with Vpr$_{mus}$. Given its dimensions, the putative SAMHD1-CtD density could accommodate

approx. 10 amino acid residues in a fully extended conformation or up to 23 residues in a kinked helical

arrangement. All previous crystal structure analyses [46], as well as secondary structure predictions

indicate that SAMHD1 residues C-terminal to the catalytic HD domain and C-terminal lobe (amino

acids 599-626) are disordered in the absence of additional binding partners. Accordingly, the N-terminal

globular domains of the SAMHD1 molecule might be flexibly linked to the C-terminal tether identified

here. In that case, the bulk of SAMHD1 samples a multitude of positions relative to the DDB1/DCAF1-

CtD/Vpr$_{mus}$ core, and consequently is averaged out in the process of cryo-EM reconstruction.

To test this hypothesis, Vpr$_{mus}$ amino acid residues in close proximity to the putative SAMHD1-CtD

density were substituted by site-directed mutagenesis. Specifically, Vpr$_{mus}$ W29 was changed to alanine

to block a hydrophobic contact with SAMHD1-CtD involving the aromatic side chain, and Vpr$_{mus}$ A66

was changed to a bulky tryptophan, in order to introduce a steric clash with SAMHD1-CtD (Fig 4B).

The structural integrity of the Vpr$_{mus}$ W29A A66W double mutant was confirmed by CD spectroscopy

(S3H Fig), and it was then assessed for complex formation with DDB1/DCAF1-CtD and SAMHD1 by

analytical GF. In comparison to wild type Vpr$_{mus}$, the W29A A66W mutant showed reduced

DDB1/DCAF1-CtD/Vpr$_{mus}$/SAMHD1 complex peak intensity (Fig 4C, fraction 6), concomitant with (i)

enrichment of DDB1/DCAF1-CtD/Vpr$_{mus}$ ternary complex, sub-stoichiometrically bound to SAMHD1

(Fig 4C, fraction 7), (ii) excess DDB1/DCAF1-CtD complex (Fig 4C, fraction 8), and (iii) excess

SAMHD1 (Fig 4C, fractions 9-10). In addition, binary combinations of the Vpr$_{mus}$ W29A A66W double

mutant with DDB1/DCAF1-CtD or SAMHD1 were also analysed by GF. These data show loss of

SAMHD1 interaction (Fig 4D), while the ability to bind DDB1/DCAF1-CtD is retained (Fig 4E).

Together, these biochemical analyses support a location of the SAMHD1-CtD binding site on the upper

surface of the Vpr$_{mus}$ helix bundle, as suggested by medium-resolution cryo-EM reconstruction.

To obtain additional experimental evidence, the CRL4$^{DCAF1-CtD}$/Vpr$_{mus}$/SAMHD1 assembly was further

examined by cross-linking mass spectrometry (CLMS), using the photo-reactive cross-linker sulfo-

SDA. This bi-functional compound contains an NHS-ester functional group on one end that reacts with

primary amines and hydroxyl groups, while the other end covalently links to any amino acid sidechain

within reach upon UV-activation via a carbene intermediate [58]. Accordingly, incubation of proteins

or protein complexes with sulfo-SDA, followed by UV-illumination, allows for high-density cross-

linking of lysine, and to a lesser extent serine, threonine and tyrosine side chains to amino acids within

reach of the SDA spacer group, with faster kinetics than pure NHS ester-based cross-linkers, due to the

short half-life of the UV-activated intermediate. Cross-linked peptides are subsequently identified by

mass spectrometry, and provide insights into the topology and residue-residue distances of proteins and

complexes [59]. In the case of the CRL4$^{DCAF1-CtD}$/Vpr$_{mus}$/SAMHD1 assembly, the majority of identified

cross-links that can be mapped onto the structure (468/519, 90.2%) are within the 25 Å violation

threshold imposed by the geometry of the SDA spacer. Interestingly, 8 of the 11 cross-links between

DDB1 and CUL4 are satisfied by the state-1 model, but increasingly violated in states-2 and -3 (S5E

Fig), supporting in solution the rotational flexibility of the CRL4 stalk with respect to DDB1/DCAF1-

CtD, as observed in cryo-EM (Fig 3).

An additional 300 cross-links involved SAMHD1, extending to the C-terminal half of CUL4, to a DDB1

sequence stretch comprising amino acid residues 900-1000, to parts of DCAF1-CtD and to Vpr$_{mus}$ (Figs

5A, S5E). The CRL4$^{DCAF1-CtD}$/Vpr$_{mus}$ residues exhibiting cross-links to SAMHD1 were mapped onto the

state-2 model, and showed the presence of a large, yet defined, interaction surface (Fig 5B). Importantly,

cross-links were apparent between the C-terminus of SAMHD1-CtD (residues K622, K626) and a region

in Vpr$_{mus}$ Helix-1 (residues 27-36), which forms a part of the putative SAMHD1-CtD binding interface

observed in cryo-EM, and which contains Vpr$_{mus}$ W29, one of the residues substituted in the mutagenesis

and biochemical analysis presented above (Fig 5B, purple spheres). In addition, amino acid residues

from the N-terminal portion of SAMHD1-CtD (residues K595, K596, T602-S606) cross-linked close to

the DCAF1-CtD "acidic loop" (residues 1092-1096), which is immobilised by Vpr near the proposed

SAMHD1-CtD binding site, and to the very C-terminus of CUL4 (residues Y744, A759), which is also

adjacent to the predicted SAMHD1-CtD binding position (Fig. 5B, pink spheres). Lastly, cross-links in

the SAMHD1 N-terminal SAM and catalytic HD domains almost exclusively involved patches of the

CRL4$^{DCAF1-CtD}$/Vpr$_{mus}$ surface surrounding and facing towards the putative SAMHD1-CtD attachment

point (Fig. 5B, light brown spheres). These observations are compatible with recruitment of SAMHD-CtD on the upper surface of the Vpr helix bundle, as indicated by cryo-EM. In addition, the spatial distribution of cross-links involving the SAMHD1 N-terminal domains suggest that these are flexibly connected to SAMHD1-CtD, leading to highly variable positioning relative to $\text{CRL4}^{\text{DCAF1-CtD}}/\text{Vpr}_{\text{mus}}$ and thus offering a multitude of cross-linking opportunities to nearby CRL4 components, again in line with the cryo-EM reconstruction results, especially upon consideration of the positional heterogeneity of the CUL4 stalk (Fig. 3).

In order to evaluate the distance information inherent in SAMHD1-CtD cross-links in a more quantitative way, the volume accessible to SAMHD1-CtD for interaction with $\text{CRL4}^{\text{DCAF1-CtD}}/\text{Vpr}_{\text{mus}}$, consistent with the CLMS distance restraints, was simulated using the DisVis software tool [60, 61]. For this analysis, SAMHD1-CtD was modelled as peptide in extended conformation. During the simulation, the state-2 $\text{CRL4}^{\text{DCAF1-CtD}}/\text{Vpr}_{\text{mus}}$ molecular model was kept fixed, and a six-dimensional search of all possible degrees of freedom of rotation and translation for the SAMHD1-CtD model in molecular contact with $\text{CRL4}^{\text{DCAF1-CtD}}/\text{Vpr}_{\text{mus}}$ was computed and ranked according to agreement with CLMS distance restraints. To visualise the output, all possible spatial positions of the centre of mass of SAMHD1-CtD, which satisfy >50% of the CLMS restraints, were plotted as density map on the structure of DCAF1-CtD/$\text{Vpr}_{\text{mus}}$ (Fig 5C). In accordance with the cryo-EM reconstruction, this independent computational analysis also locates SAMHD1-CtD on top of the $\text{Vpr}_{\text{mus}}$ helix bundle.

Taken together, the structural, biochemical and CLMS data are consistent with a model where the very C-terminus of SAMHD1 is recruited by $\text{Vpr}_{\text{mus}}$, to place the remaining SAMHD1 domains appropriately for access to the catalytic machinery at the distal end of the CRL4 stalk.

These data allow for structural comparison with *neo*-substrate binding modes of Vpx and Vpr proteins from different retrovirus lineages (Fig 6A-D). $\text{Vpx}_{\text{HIV-2}}$ and $\text{Vpx}_{\text{sm}}$ position SAMHD1-CtD at the side of the DCAF1 BP domain through interactions with the N-termini of Vpx Helices-1 and -3 (Fig 6B) [50]. $\text{Vpx}_{\text{mnd2}}$ and $\text{Vpx}_{\text{rcm}}$ bind SAMHD1-NtD using a bipartite interface comprising the side of the DCAF1 BP and the upper surface of the Vpx helix bundle (Fig 6C) [51, 52]. $\text{Vpr}_{\text{HIV-1}}$ engages its ubiquitylation substrate UNG2 using both the top and the upper edge of the $\text{Vpr}_{\text{HIV-1}}$ helix bundle (Fig 6D) [54]. Of note, these upper-surface interaction interfaces only partially overlap with the $\text{Vpr}_{\text{mus}}$/SAMHD1-CtD

binding interface identified here and employ fundamentally different sets of interacting amino acid residues. Thus, it appears that the molecular interaction interfaces driving Vpx/Vpr-mediated *neo-substrate* recognition and degradation are not conserved between related SIV and HIV Vpx/Vpr accessory proteins, even in cases where identical SAMHD1-CtD regions are targeted for recruitment.

## Discussion

Our X-ray crystallographic studies of the DDB1/DCAF1-CtD/Vpr$_{mus}$ assembly provide the first structural insight into a class of "hybrid" SIV Vpr proteins. These are present in the SIV$_{agm}$ and SIV$_{mus/deb/syk}$ lineages of lentiviruses and combine characteristics of related Vpr$_{HIV-1}$ and SIV Vpx accessory proteins.

Like SIV Vpx, "hybrid" Vpr proteins down-regulate the host restriction factor SAMHD1 by recruiting it to CRL4$^{DCAF1}$ for ubiquitylation and subsequent proteasomal degradation. However, using a combination of X-ray, cryo-EM and CLMS analyses, we show that the molecular strategy, which Vpr$_{mus}$ evolved to target SAMHD1, is strikingly different from Vpx-containing SIV strains. In the two clades of Vpx proteins, divergent amino acid sequence stretches just upstream of Helix-1 (variable region (VR)1, S6A Fig), together with polymorphisms in the SAMHD1-N-terminus of the respective host species, determine if HIV-2-type or SIV$_{mnd}$-type Vpx recognise SAMHD1-CtD or SAMHD1-NtD, respectively. These recognition mechanisms result in positioning of SAMHD1-CtD or -NtD on the side of the DCAF1 BP domain in a way that allows for additional contacts between SAMHD1 and DCAF1, thus forming ternary Vpx/SAMHD1/DCAF1 assemblies with very low dissociation rates [50-52, 62]. In Vpr$_{mus}$, different principles determine the specificity for SAMHD1-CtD. Here, VR1 is not involved in SAMHD1-CtD-binding at all, but forms additional interactions with DCAF1, which are not observed in Vpx/DCAF1 protein complexes (S6A Fig). Molecular contacts between Vpr$_{mus}$ and SAMHD1 are dispersed on Helices-1 and -3, facing away from the DCAF1 interaction site and immobilising SAMHD1-CtD on the top side of the Vpr$_{mus}$ helix bundle (S6A Fig). Placement of SAMHD1-CtD in such a position precludes stabilising ternary interaction with DCAF1-CtD, but still results in robust SAMHD1 ubiquitylation *in vitro* and SAMHD1 degradation in cell-based assays [24].

The catalytic dNTPase activity of SAMHD1 depends on nucleotide-dependent oligomerisation, mediated by two allosteric nucleotide-binding sites, where guanine-based nucleotides in the first site induce dimer formation, and dNTP-binding to the second site leads to assembly of the catalytically active tetramer. SAMHD1-CtD is essential for tetramer formation by contributing critical molecular contacts to neighbouring protomers. Furthermore, tetramer destabilisation by CDK1/2-cyclinA-dependent phosphorylation of T592 in SAMHD1-CtD endogenously attenuates SAMHD1 activity in cycling cells [46, 63-65]. Accordingly, it is conceivable that by sequestering SAMHD1-CtD, $Vpr_{mus}$ destabilises the SAMHD1 tetramer, and in this way abrogates SAMHD1 activity, prior to inducing its proteasomal degradation, in accordance with previous observation of $Vpx_{HIV-2}$-mediated SAMHD1 tetramer disassembly and inhibition of dNTPase activity [62].

Predictions regarding the molecular mechanism of SAMHD1-binding by other "hybrid" Vpr orthologues are difficult due to sequence divergence. Even in $Vpr_{deb}$, the closest relative to $Vpr_{mus}$, only approximately 50% of amino acid side chains lining the putative SAMHD1-CtD binding pocket are conserved (S6A Fig). Previous *in vitro* ubiquitylation and cell-based degradation experiments did not show a clear preference of $Vpr_{deb}$ for recruitment of either SAMHD1-NtD or –CtD [24, 49]. Furthermore, it is disputed if $Vpr_{deb}$ actually binds DCAF1 [66], which might possibly be explained by amino acid variations in the very N-terminus and/or in Helix-3 (S6A Fig). $Vpr_{syk}$ is specific for SAMHD1-CtD [49], but the majority of residues forming the binding platform for SAMHD1-CtD observed in the present study are not conserved. The $SIV_{agm}$ lineage of Vpr proteins is even more divergent, with significant differences not only in possible SAMHD1-contacting residues, but also in the sequence stretches preceding Helix-1, and connecting Helices-2 and -3, as well as in the N-terminal half of Helix-3 (S6A Fig). Furthermore, there are indications that recruitment of SAMHD1 by the $Vpr_{agm.GRI}$ sub-type involves molecular recognition of both SAMHD1-NtD and –CtD [49, 53]. In conclusion, recurring rounds of evolutionary lentiviral adaptation to the host SAMHD1 restriction factor, followed by host re-adaptation, resulted in highly species-specific, diverse molecular modes of Vpr-SAMHD1 interaction. Similar molecular arms races between cell-intrinsic antiviral host factors and viral antagonists shaped the species-specific lentiviral antagonism of e.g. host restriction factors of the APOBEC3 family and tetherin, through induction of their degradation by the respective viral antagonists

Vif or Nef/Vpu [67-69]. Furthermore, viral re-adaptation to certain simian and human variants of these

restriction factors, following cross-species transmission, took part in the emergence of pandemic HIV

strains, thus highlighting the importance of structural insight into these processes [9]. In addition to the

instance presented here, further structural characterisation of SAMHD1-Vpr complexes will be

necessary to fully define outcomes of this particular virus-host molecular arms race.

Previous structural investigation of DDB1/DCAF1/Vpr$_{HIV-1}$ in complex with the *neo*-substrate UNG2

demonstrated that Vpr$_{HIV-1}$ engages UNG2 by mimicking the DNA phosphate backbone. More precisely,

UNG2 residues, which project into the major groove of its endogenous DNA substrate, insert into a

hydrophobic cleft formed by Vpr$_{HIV-1}$ Helices-1, -2 and the N-terminal half of Helix-3 [54]. This

mechanism might rationalise Vpr$_{HIV-1}$'s extraordinary binding promiscuity, since the list of potential

Vpr$_{HIV-1}$ degradation substrates is significantly enriched in DNA- and RNA-binding proteins [27].

Moreover, promiscuous Vpr$_{HIV-1}$-induced degradation of host factors with DNA- or RNA-binding

activity has been proposed to induce cell cycle arrest at the G2/M phase border, which is the most

thoroughly described phenotype of Vpr proteins so far [26, 27, 70]. In Vpr$_{mus}$, the N-terminal half of

Helix-1 as well as the bulky amino acid residue W48, which is also conserved in Vpr$_{agm}$ and Vpx,

constrict the hydrophobic cleft (S6A, B Fig). Furthermore, the extended N-terminus of Vpr$_{mus}$ Helix-3

is not compatible with UNG2-binding due to steric exclusion (S6C Fig). In accordance with these

observations, Vpr$_{mus}$ does not down-regulate UNG2 in a human T cell line [27]. However, Vpr$_{mus}$, Vpr$_{syk}$

and Vpr$_{agm}$ also cause G2/M cell cycle arrest in their respective host cells [66, 71, 72]. This strongly

hints at the existence of further structural determinants in Vpr$_{mus}$, Vpr$_{syk}$, Vpr$_{agm}$ and potentially Vpr$_{HIV-1}$,

which regulate recruitment and ubiquitylation of DNA/RNA-binding host factors, in addition to the

hydrophobic, DNA-mimicking cleft on top of the three-helix bundle. Future efforts to structurally

characterise these determinants will further extend our understanding of how the Vpx/Vpr helical

scaffold binds, and in this way adapts to a multitude of *neo*-substrate epitopes.

Our cryo-EM reconstructions of CRL4$^{DCAF1-CtD}$/Vpr$_{mus}$/SAMHD1, complemented by CLMS, also

provide insights into the structural dynamics of CRL4 assemblies prior to ubiquitin transfer. The data

confirm previously described rotational movement of the CRL4 stalk, in the absence of constraints

imposed by a crystal lattice, creating a ubiquitylation zone around the Vpr$_{mus}$-modified substrate receptor

(Figs 3 and 7A) [13, 15, 16, 19, 57]. Missing density for the neddylated CUL4 WHB domain and for

the catalytic ROC1 RING domain indicates that these distal stalk elements are highly mobile and likely

sample a multitude of orientations relative to the CUL4 scaffold (Fig 7B). These observations are in line

with structure analyses of CRL1 and CRL5, where CUL1/5 neddylation leads to re-orientation of the

cullin WHB domain, and to release of the ROC1 RING domain from the cullin scaffold, concomitant

with stimulation of ubiquitylation activity [56]. Moreover, recent cryo-EM structure analysis of

CRL1$^{\beta\text{-TRCP}}$/IκBα demonstrated substantial mobility of pre-catalytic CUL1-NEDD8 WHB and ROC1

RING domains [73]. Such flexibility seems necessary to structurally organise multiple CRL1-dependent

processes, in particular the nucleation of a catalytic assembly, involving intricate protein-protein

interactions between NEDD8, CUL1, ubiquitin-charged E2 and substrate receptor. This synergistic

assembly then steers the ubiquitin C-terminus towards a substrate lysine for priming with ubiquitin [73].

Accordingly, our cryo-EM studies might indicate that similar principles apply for CRL4-catalysed

ubiquitylation. However, to unravel the catalytic architecture of CRL4, sophisticated cross-linking

procedures as in reference (73) will have to be pursued.

Intrinsic mobility of CRL4 stalk elements might assist the accommodation of a variety of sizes and

shapes of substrates in the CRL4 ubiquitylation zone and might rationalise the wide substrate range

accessible to CRL4 ubiquitylation through multiple DCAF receptors. Owing to selective pressure to

counteract the host's SAMHD1 restriction, HIV-2 and certain SIVs, amongst other viruses, have taken

advantage of this dynamic CRL4 architecture by modification of the DCAF1 substrate receptor with

Vpx/Vpr-family accessory proteins. By tethering either SAMHD1-CtD or -NtD to DCAF1, and in this

way flexibly recruiting the bulk of SAMHD1, the accessibility of lysine side chains both tether-proximal

and on the SAMHD1 globular domains to the CRL4 catalytic assembly might be further improved (Fig

7C, D). This ensures efficient Vpx/Vpr-mediated SAMHD1 priming, poly-ubiquitylation and

proteasomal degradation to stimulate virus replication. In addition, structural insight into this

evolutionary optimised, highly specific protein degradation machinery might inform the positioning of

novel CRL4$^{DCAF1}$-based synthetic modalities for targeted protein degradation, e.g. in the form of

proteolysis-targeting chimera-(PROTAC-) or molecular glue-type compounds [74, 75].

Lastly, while current highly active antiretroviral therapy (HAART) regimens are able to control HIV-1

replication in infected patients [76], they cannot eradicate the virus due to viral rebound after treatment

cessation, and they lead to emergence of resistant virus variants [77]. Accordingly, identification and

inhibition of novel targets, in addition to those already covered by HAART, are of high interest. In this

context, HIV accessory proteins have for a long time been regarded as promising drug targets [78, 79].

For example, preclinical research aiming at inhibition of Vif-mediated APOBEC3 degradation

culminated in the isolation of several compounds which impede HIV-1 replication [80-82]. Due to its

promiscuous host interactions, $Vpr_{HIV-1}$ is also considered an attractive antiretroviral target [83]. The

comparative structural analyses of Vpr/DCAF1-CtD interaction presented here might inform future

efforts to disrupt this interaction interface with small molecules, in order to abolish all Vpr-induced host

protein degradation processes at once, to maximise the inhibitory effect on virus replication.

**Methods**

**Protein expression and purification**

Constructs were PCR-amplified from cDNA templates and inserted into the indicated expression

plasmids using standard restriction enzyme methods (S2 Table). pAcGHLT-B-DDB1 (plasmid #48638)

and pET28-UBA1 (plasmid #32534) were obtained from Addgene. The pOPC-UBA3-GST-APPBP1

co-expression plasmid, and the pGex6P2-UBC12 plasmid were obtained from MRC-PPU Reagents and

Services (clones 32498, 3879). Bovine erythrocyte ubiquitin and recombinant hsNEDD8 were

purchased from Sigma-Aldrich (U6253) and BostonBiochem (UL-812) respectively. Point mutations

were introduced by site-directed mutagenesis using KOD polymerase (Novagen). All constructs and

variants are summarised S3 Table.

Proteins expressed from vectors pAcGHLT-B, pGex6P1/2, pOPC and pET49b contained an N-terminal

GST-His-tag; pHisSUMO – N-terminal His-SUMO-tag; pET28, pRSF-Duet-1 – N-terminal His-tag;

pTri-Ex-6 – C-terminal His-tag. Constructs in vectors pAcGHLT-B and pTri-Ex-6 were expressed in

Sf9 cells, and constructs in vectors pET28, pET49b, pGex6P1/2, pRSF-Duet-1, and pHisSUMO in *E.*

*coli* Rosetta 2(DE3).

Recombinant baculoviruses (*Autographa californica nucleopolyhedrovirus* clone C6) were generated

as described previously [84]. Sf9 cells were cultured in Insect-XPRESS medium (Lonza) at 28°C in an

Innova 42R incubator shaker (New Brunswick) at a shaking speed of 180 rpm. In a typical preparation,

1 L of Sf9 cells at $3 \times 10^6$ cells/mL were co-infected with 4 mL of high titre DDB1 virus and 4 mL of

high titre DCAF1-CtD virus for 72 h.

For a typical *E. coli* Rosetta 2 (DE3) expression, 2 L of LB medium was inoculated with 20 mL of an

overnight culture and grown in a Multitron HT incubator shaker (Infors) at 37°C, 150 rpm until $OD_{600}$

reached 0.7. At that point, temperature was reduced to 18°C, protein expression was induced by addition

of 0.2 mM IPTG, and cultures were grown for further 20 h. During co-expression of CUL4 and ROC1

from pRSF-Duet, 50 µM zinc sulfate was added to the growth medium before induction.

Sf9 cells were pelleted by centrifugation at 1000 rpm, 4°C for 30 min using a JLA 9.1000 centrifuge

rotor (Beckman). *E. coli* cells were pelleted by centrifugation at 4000 rpm, 4°C for 15 min using the

same rotor. Cell pellets were resuspended in buffer containing 50 mM Tris, pH 7.8, 500 mM NaCl, 4

mM MgCl2, 0.5 mM tris-(2-carboxyethyl)-phosphine (TCEP), mini-complete protease inhibitors (1

tablet per 50 mL) and 20 mM imidazole (for His-tagged proteins only). 100 mL of lysis buffer was used

for resuspension of a pellet from 1 L Sf9 culture, and 35 mL lysis buffer per pellet from 1 L *E. coli*

culture. Before resuspension of CUL4/ROC1 co-expression pellets, the buffer pH was adjusted to 8.5.

5 µL Benzonase (Merck) was added and the cells lysed by passing the suspension at least twice through

a Microfluidiser (Microfluidics). Lysates were clarified by centrifugation at 48000xg for 45 min at 4°C.

Protein purification was performed at 4°C on an Äkta pure FPLC (GE) using XK 16/20 chromatography

columns (GE) containing 10 mL of the appropriate affinity resin. GST-tagged proteins were captured

on glutathione-Sepharose (GSH-Sepharose FF, GE), washed with 250 mL of wash buffer (50 mM Tris-

HCl pH 7.8, 500 mM NaCl, 4 mM $MgCl_2$, 0.5 mM TCEP), and eluted with the same buffer

supplemented with 20 mM reduced glutathione. His-tagged proteins were immobilised on Ni-Sepharose

HP (GE), washed with 250 mL of wash buffer supplemented with 20 mM imidazole, and eluted with

wash buffer containing 0.3 M imidazole. Eluent fractions were analysed by SDS-PAGE, and appropriate

fractions were pooled and reduced to 5 mL using centrifugal filter devices (Vivaspin). If applicable, 100

µg GST-3C protease, or 50 µg thrombin, per mg total protein, was added and the sample was incubated

for 12 h on ice to cleave off affinity tags. As second purification step, gel filtration chromatography

(GF) was performed on an Äkta prime plus FPLC (GE), with Superdex 200 16/600 columns (GE),

equilibrated in 10 mM Tris-HCl pH 7.8, 150 mM NaCl, 4 mM $MgCl_2$, 0.5 mM TCEP buffer, at a flow

rate of 1 mL/min. For purification of the CUL4/ROC1 complex, the pH of all purification buffers was

adjusted to 8.5. Peak fractions were analysed by SDS-PAGE, appropriate fractions were pooled and

concentrated to approx. 20 mg/mL, flash-frozen in liquid nitrogen in small aliquots and stored at -80°C.

Protein concentrations were determined with a NanoDrop spectrophotometer (ND 1000, Peqlab), using

theoretical absorption coefficients calculated based upon the amino acid sequence by ProtParam on the

ExPASy webserver [85].

**Analytical gel filtration analysis**

Prior to gel filtration analysis affinity tags were removed by incubation of 30 µg GST-3C protease with

6 µM of each protein component in a volume of 150 µL wash buffer, followed by incubation on ice for

12 h. In order to remove the cleaved GST-tag and GST-3C protease, 20 µL GSH-Sepharose FF beads

(GE) were added and the sample was rotated at 4 °C for one hour. GSH-Sepharose beads were removed

by centrifugation at 4°C, 3500 rpm for 5 min, and 120 µL of the supernatant was loaded on an analytical

GF column (Superdex 200 10/300 GL, GE), equilibrated in 10 mM Tris-HCl pH 7.8, 150 mM NaCl,

4 mM $MgCl_2$, 0.5 mM TCEP, at a flow rate of 0.5 mL/min. 1 mL fractions were collected and analysed

by SDS-PAGE. For the following control samples, 120 µl of purified protein was applied directly to the

GF column, because no purification tag had to be removed by cleavage: SAMHD1, T4L-SAMHD1-

CtD, SAMHD1-ΔCtD. For these samples, the concentration was adjusted to 18 µM, 37 µM and 30 µM,

respectively, to account for the lower extinction coefficient of these isolated protein components, in

order to allow for better visualisation of the elution peak. The GF column was calibrated using the high

molecular weight range (HMW) Gel Filtration Calibration Kit (GE) according to the manufacturer's

instructions.

***In vitro* ubiquitylation assays**

160 µL reactions were prepared, containing 0.5 µM substrate (indicated SAMHD1 constructs, S2 Fig),

0.125 µM DDB1/DCAF1-CtD, 0.125 µM CUL4/ROC1, 0.125 µM HisSUMO-T4L-Vpr$_{mus}$ (residues 1-

92), 0.25 µM UBCH5C, 15 µM ubiquitin in 20 mM Tris-HCl pH 7.8, 150 mM NaCl, 2.5 mM MgCl$_2$,

2.5 mM ATP. In control reactions, certain components were left out as indicated in S2 Fig. A 30 µl

sample for SDS-PAGE analysis was taken (t=0). Reactions were initiated by addition of 0.05 µM UBA1,

incubated at 37°C, and 30 µl SDS-PAGE samples were taken after 1 min, 2 min, 5 min and 15 min,

immediately mixed with 10 µl 4x SDS sample buffer and boiled at 95°C for 5 min. Samples were

analysed by SDS-PAGE.

***In vitro* neddylation of CUL4/ROC1**

For initial neddylation tests, a 200 µL reaction was prepared, containing 8 µM CUL4/ROC1, 1.8 µM

UBC12, 30 µM NEDD8 in 50 mM Tris-HCl pH 7.8, 150 mM NaCl, 2.5 mM MgCl$_2$, 2.5 mM ATP. 2x

30 µL samples were taken for SDS-PAGE, one was immediately mixed with 10 µL 4x SDS sample

buffer, the other one incubated for 60 min at 25°C. The reaction was initiated by addition of 0.7 µM

APPBP1/UBA3, incubated at 25°C, and 30 µL SDS-PAGE samples were taken after 1 min, 5 min,

10 min, 30 min and 60 min, immediately mixed with 10 µL 4x SDS sample buffer and boiled at 95°C

for 5 min. Samples were analysed by SDS-PAGE. Based on this test, the reaction was scaled up to 1 mL

and incubated for 5 min at 25°C. Reaction was quenched by addition of 5 mM TCEP and immediately

loaded onto a Superdex 200 16/600 GF column (GE), equilibrated in 10 mM Tris-HCl pH 7.8, 150 mM

NaCl, 4 mM MgCl$_2$, 0.5 mM TCEP at a flow rate of 1 mL/min. Peak fractions were analysed by SDS-

PAGE, appropriate fractions were pooled and concentrated to ~20 mg/mL, flash-frozen in liquid

nitrogen in small aliquots and stored at -80°C.

**X-ray crystallography sample preparation, crystallisation, data collection and structure**

**solution**

**DDB1/DCAF1-CtD complex**. DDB1/DCAF1-CtD crystals were grown by the hanging drop vapour

diffusion method, by mixing equal volumes (1 µL) of DDB1/DCAF1-CtD solution at 10 mg/mL with

reservoir solution containing 100 mM Tri-Na citrate pH 5.5, 18% PEG 1000 and suspending over a 500

µl reservoir. Crystals grew over night at 18°C. Crystals were cryo-protected in reservoir solution

supplemented with 20% glycerol and cryo-cooled in liquid nitrogen. A data set from a single crystal was

collected at Diamond Light Source (Didcot, UK) at a wavelength of 0.92819 Å. Data were processed

using XDS [86] (S1 Table), and the structure was solved using molecular replacement with the program

MOLREP [87] and available structures of DDB1 (PDB 3e0c) and DCAF1-CtD (PDB 4cc9) [50] as

search models. Iterative cycles of model adjustment with the program Coot [88], followed by refinement

using the program PHENIX [89] yielded final $R/R_{free}$ factors of 22.0%/27.9% (S1 Table). In the model,

94.5 % of residues have backbone dihedral angles in the favoured region of the Ramachandran plot, the

remainder fall in the allowed regions, and none are outliers. Details of data collection and refinement

statistics are presented in S1 Table. Coordinates and structure factors have been deposited in the PDB,

accession number 6zue.

**DDB1/DCAF1-CtD/T4L-Vpr$_{mus}$ (1-92) complex**. The DDB1/DCAF1-CtD/Vpr$_{mus}$ complex was

assembled by incubation of purified DDB1/DCAF1-CtD and HisSUMO-T4L-Vpr$_{mus}$ (residues 1-92), at

a 1:1 molar ratio, in a buffer containing 50 mM Bis-tris propane pH 8.5, 0.5 M NaCl, 4 mM $MgCl_2$, 0.5

mM TCEP, containing 1 mg of HRV-3C protease for HisSUMO-tag removal. After incubation on ice

for 12 h, the sample was loaded onto a Superdex 200 16/600 GF column (GE), with a 1 mL GSH-

Sepharose FF column (GE) connected in line. The column was equilibrated with 10 mM Bis-tris propane

pH 8.5, 150 mM NaCl, 4 mM MgCl2, and 0.5 mM TCEP. The column flow rate was 1 mL/min. GF

fractions were analysed by SDS-PAGE, appropriate fractions were pooled and concentrated to 4.5

mg/mL.

Crystals were prepared by the sitting drop vapour diffusion method, by mixing equal volumes (200 nL)

of the protein complex at 4.5 mg/mL and reservoir solution containing 8-10% PEG 4000 (w/v), 200 mM

$MgCl_2$, 100 mM HEPES-NaOH, pH 7.0-8.2. The reservoir volume was 75 µL. Crystals grew after at

least 4 weeks of incubation at 4°C. Crystals were cryo-protected in reservoir solution supplemented with

20% glycerol and cryo-cooled in liquid nitrogen. Data sets from two single crystals were collected,

initially at BESSY II (Helmholtz-Zentrum Berlin, HZB) at a wavelength of 0.91841 Å, and later at

ESRF (Grenoble) at a wavelength of 1 Å. Data sets were processed separately using XDS [86] and

XDSAPP [90]. The structure was solved by molecular replacement, using the initial BESSY data set,

with the program PHASER [91], and the following structures as search models: DDB1/DCAF1-CtD (this work) and T4L variant E11H (PDB 1qt6) [92]. After optimisation of the initial model and refinement against the higher-resolution ESRF data set, $Vpr_{mus}$ was placed manually into the density, using an NMR model of $Vpr_{HIV-1}$ (PDB 1m8l) [93] as guidance. Iterative cycles of model adjustment with the program Coot [88], followed by refinement using the program PHENIX [89] yielded final $R/R_{free}$ factors of 21.61%/26.05%. In the model, 95.1 % of residues have backbone dihedral angles in the favoured region of the Ramachandran plot, the remainder fall in the allowed regions, and none are outliers. Details of data collection and refinement statistics are presented in S1 Table. Coordinates and structure factors have been deposited in the PDB, accession number 6zx9.

**Cryo-EM sample preparation and data collection**

**Complex assembly**. Purified CUL4-NEDD8/ROC1, DDB1/DCAF1-CtD, GST-$Vpr_{mus}$ and rhesus macaque SAMHD1, 1 µM each, were incubated in a final volume of 1 mL of 10 mM Tris-HCl pH 7.8, 150 mM NaCl, 4 mM $MgCl_2$, 0.5 mM TCEP, supplemented with 1 mg of GST-3C protease. After incubation on ice for 12 h, the sample was loaded onto a Superdex 200 16/600 GF column (GE), equilibrated with the same buffer at 1 mL/min, with a 1 mL GSH-Sepharose FF column (GE) connected in line. GF fractions were analysed by SDS-PAGE, appropriate fractions were pooled and concentrated to 2.8 mg/mL.

**Grid preparation.** 3.5 µl protein solution containing 0.05 µM CUL4-NEDD8/ROC1/DDB1/DCAF1-CtD/$Vpr_{mus}$/SAMHD1 complex and 0.25 µM UBCH5C-ubiquitin conjugate (S4 A, B Fig) were applied to a 300 mesh Quantifoil R2/4 Cu/Rh holey carbon grid (Quantifoil Micro Tools GmbH) coated with an additional thin carbon film as sample support and stained with 2% uranyl acetate for initial characterisation. For cryo-EM, a fresh 400 mesh Quantifoil R1.2/1.3 Cu holey carbon grid (Quantifoil Micro Tools GmbH) was glow-discharged for 30 s using a Harrick plasma cleaner with technical air at 0.3 mbar and 7 W. 3.5 µl protein solution containing 0.4 µM CUL4-NEDD8/ROC1/DDB1/DCAF1-CtD/$Vpr_{mus}$/SAMHD1 complex and 2 µM UBCH5C-ubiquitin conjugate were applied to the grid, incubated for 45 s, blotted with a Vitrobot Mark II device (FEI, Thermo Fisher Scientific) for 1-2 s at 8°C and 80% humidity, and plunged in liquid ethane. Grids were stored in liquid nitrogen until imaging.

**Cryo-EM data collection.** Initial negative stain and cryo-EM datasets were collected automatically

for sample quality control and low-resolution reconstructions on a 120 kV Tecnai Spirit cryo-EM (FEI,

Thermo Fisher Scientific) equipped with a F416 CMOS camera (TVIPS) using Leginon [94, 95].

Particle images were then analysed by 2D classification and initial model reconstruction using SPHIRE

[96], cisTEM [97] and Relion 3.07 [98]. These data revealed the presence of the complexes containing

both DDB1/DCAF1-CtD/Vpr$_{mus}$ (core) and CUL4/ROC1 (stalk). High-resolution data was collected on

a 300 kV Tecnai Polara cryo-EM (FEI, Thermo Fisher Scientific) equipped with a K2summit direct

electron detector (Gatan) at a nominal magnification of 31000x, with a pixel size of 0.625 Å/px on the

object scale. In total, 3644 movie stacks were collected in super-resolution mode using Leginon [94, 95]

with the following parameters: defocus range of 0.5-3.0 µm, 40 frames per movie, 10 s exposure time,

electron dose of 1.25 e/Å$^2$/s and a cumulative dose of 50 e/Å$^2$ per movie.

**Cryo-EM computational analysis**

Movies were aligned and dose-weighted using MotionCor2 [99] and initial estimation of the contrast

transfer function (CTF) was performed with the CTFFind4 package [100]. Resulting micrographs were

manually inspected to exclude images with substantial contaminants (typically large protein aggregates

or ice contaminations) or grid artefacts. Power spectra were manually inspected to exclude images with

astigmatic, weak, or poorly defined spectra. After these quality control steps the dataset included 2322

micrographs (63% of total). At this stage, the data set was picked twice and processed separately, to

yield reconstructions of states-1, -2 and -3 (analysis 1) and of the core (analysis 2).

For **analysis 1**, particle positions were determined using cisTEMs Gaussian picking routine, yielding

959,155 particle images in total. After two rounds of 2D-classification, 227,529 particle images were

selected for further processing (S4C, D Fig). Using this data, an initial model was created using Relion

3.07. The resulting map yielded strong signal for the core but only fragmented stalk density, indicating

a large heterogeneity in the stalk-region within the data set. This large degree of compositional (+/-

stalk) and conformational heterogeneity (movement of the stalk relative to the core) made the

classification challenging. Accordingly, alignment and classification were carried out simultaneously.

The first objective was to separate the data set into three categories: "junk", "core" and "core+stalk".

Therefore, the stalk was deleted from the initial model using the "Eraser"-tool in Chimera [101]. This

core-map was used as an initial model for the Tier 1 3D-classification with Relion 3.07 at a decimated

pixel size of 2.5 Å/px. The following parameters were used: number of classes K=6, T=10, global step

search=7.5°, number of iterations= 25. The classification yielded two classes containing the stalk

(classes 3 and 5 containing 23% and 22% of the particle images, respectively) (S4D Fig). These particles

were pooled and directed into Tier 2 3D-classification using the following parameters: number of classes

K=6, T=10, global step search=7.5°, number of iterations=25. Three of these classes yielded medium-

resolution maps with interpretable features (states-1, -2 and -3, S4D Fig). These three classes were

refined individually using 3D autorefinement in Relion 3.07, resulting in maps with resolution ranging

from 7.8 Å – 8.9 Å (S4E-G Fig).

For **analysis 2**, particle positions were determined using template matching with a filtered map

comprising core and stalk using the software Gautomatch (https://www2.mrc-

lmb.cam.ac.uk/research/locally-developed-software/zhang-software/). 712,485 particle images were

found, extracted with Relion 3.07 and subsequently 2D-classified using cryoSPARC [102], resulting in

505,342 particle images after selection (S5A, B Fig). These particle images were separated into two

equally sized subsets and Tier 1 3D-classification was performed using Relion 3.07 on both of them to

reduce computational burden (S5B Fig). The following parameters were used: initial model="core",

number of classes K=4, T=10, global step search=7.5°, number of iterations=25, pixel size 3.75 Å/px.

From these, the ones possessing both core and stalk were selected. Classes depicting a similar stalk

orientation relative to the core were pooled and directed into Tier 2 as three different subpopulations

containing 143,172, 193,059 and 167,666 particle images, respectively (S5B Fig).

For Tier 2, each subpopulation was classified separately into 4 classes each. From these 12 classes, all

particle images exhibiting well-defined densities for core and stalk were pooled and labelled

"core+stalk", resulting in 310,801 particle images in total. 193,096 particle images representing classes

containing only the core were pooled and labelled "core" (S5B Fig)

For Tier 3, the "core" particle subset was separated into 4 classes which yielded uninterpretable

reconstructions lacking medium- or high-resolution features. The "core+stalk" subset was separated into

6 classes, with 5 classes containing both stalk and core (S5B Fig) and one class consisting only of the

core with Vpr$_{mus}$ bound. The 5 classes with stalk showed similar stalk orientations as the ones obtained

from analysis 1 (see above, S4 Fig), but refined individually to lower resolution as in analysis 1 and

were discarded. However, individual refinement of the core-only tier 3 class yielded a 7.3 Å

reconstruction (S5C, D Fig).

**Molecular visualisation, rigid body fitting, 3D structural alignments, rotation and**

**interface analysis**

Density maps and atomic models were visualised using Coot [88], PyMOL (Schrödinger) and UCSF

Chimera [101]. Rigid body fits and structural alignments were performed using the program UCSF

Chimera [101]. Rotation angles between extreme DDB1 BPB domain positions were measured using

the DynDom server [103] (http://dyndom.cmp.uea.ac.uk/dyndom/runDyndom.jsp). Molecular

interfaces were analysed using the EBI PDBePISA server [104] (https://www.ebi.ac.uk/msd-

srv/prot_int/cgi-bin/piserver).

**Multiple sequence alignment**

705 A multiple sequence alignment was calculated using the EBI ClustalOmega server [105]

706 (https://www.ebi.ac.uk/Tools/msa/clustalo/), and adjusted manually using the program GeneDoc [106].

**Cross-linking mass spectrometry (CLMS)**

**Complex assembly.** Purified CUL4/ROC1, DDB1/DCAF1-CtD, GST-Vpr$_{mus}$ and rhesus macaque

SAMHD1, 1 μM each, were incubated in a volume of 3 mL buffer containing 10 mM HEPES pH 7.8,

150 mM NaCl, 4 mM MgCl$_2$, 0.5 mM TCEP, supplemented with 1 mg GST-3C protease. After

incubation on ice for 12 h, the sample was loaded onto a Superdex 200 16/600 GF column (GE),

equilibrated with the same buffer, at a flow rate of 1 mL/min with a 1 mL GSH-Sepharose FF column

(GE) connected in line. GF fractions were analysed by SDS-PAGE, appropriate fractions were pooled

and concentrated to 6 mg/mL.

**Photo-Crosslinking.** The cross-linker sulfo-SDA (sulfosuccinimidyl 4,4′-azipentanoate) (Thermo

Scientific) was dissolved in cross-linking buffer (10 mM HEPES pH 7.8, 150 mM NaCl, 4 mM

MgCl$_2$, 0.5 mM TCEP) to 100 mM before use. The labelling step was performed by incubating

18 µg aliquots of the complex at 1 mg/mL with 2, 1, 0.5, 0.25, 0.125 mM sulfo-SDA, added,

respectively, for an hour. The samples were then irradiated with UV light at 365 nm, to form cross-

links, for 20 min and quenched with 50 mM NH$_4$HCO$_3$ for 20 min. All steps were performed on

ice. Reaction products were separated on a Novex Bis-Tris 4–12% SDS−PAGE gel (Life

Technologies). The gel band corresponding to the cross-linked complex was excised and digested

with trypsin (Thermo Scientific Pierce) [107] and the resulting tryptic peptides were extracted and

desalted using C18 StageTips [108]. Eluted peptides were fractionated on a Superdex Peptide

3.2/300 increase column (GE Healthcare) at a flow rate of 10 µL/min using 30% (v/v) acetonitrile

and 0.1 % (v/v) trifluoroacetic acid as mobile phase. 50 µL fractions were collected and vacuum-

dried.

**CLMS acquisition.** Samples for analysis were resuspended in 0.1% (v/v) formic acid, 3.2% (v/v)

acetonitrile. LC-MS/MS analysis was performed on an Orbitrap Fusion Lumos Tribrid mass

spectrometer (Thermo Fisher) coupled on-line with an Ultimate 3000 RSLCnano HPLC system

(Dionex, Thermo Fisher). Samples were separated on a 50 cm EASY-Spray column (Thermo Fisher).

Mobile phase A consisted of 0.1% (v/v) formic acid and mobile phase B of 80% (v/v) acetonitrile with

0.1% (v/v) formic acid. Flow rates were 0.3 µL/min using gradients optimized for each chromatographic

fraction from offline fractionation, ranging from 2% mobile phase B to 55% mobile phase B over

90 min. MS data were acquired in data-dependent mode using the top-speed setting with a 3 s cycle

time. For every cycle, the full scan mass spectrum was recorded using the Orbitrap at a resolution of

120,000 in the range of 400 to 1,500 m/z. Ions with a precursor charge state between 3+ and 7+ were

isolated and fragmented. Analyte fragmentation was achieved by Higher-Energy Collisional

Dissociation (HCD) [109] and fragmentation spectra were then recorded in the Orbitrap with a resolution

of 50,000. Dynamic exclusion was enabled with single repeat count and 60 s exclusion duration.

**CLMS processing.** A recalibration of the precursor m/z was conducted based on high-confidence

(<1% false discovery rate (FDR)) linear peptide identifications. The re-calibrated peak lists were

searched against the sequences and the reversed sequences (as decoys) of cross-linked peptides using

the Xi software suite (v.1.7.5.1) for identification [110]. Final crosslink lists were compiled using the

identified candidates filtered to <1% FDR on link level with xiFDR v.2.0 [111] imposing a minimum

of 20% sequence coverage and 4 observed fragments per peptide.

**CLMS analysis.** In order to sample the accessible interaction volume of the SAMHD1-CtD consistent

with CLMS data, a model for SAMHD1 was generated using I-TASSER [112]. The SAMHD1-CtD,

which adopted a random coil configuration, was extracted from the model. In order to map all crosslinks,

missing loops in the complex structure were generated using MODELLER [113]. An interaction volume

search was then submitted to the DisVis webserver [61] with an allowed distance between 1.5 Å and 22

Å for each restraint using the "complete scanning" option. The rotational sampling interval was set to

9.72° and the grid voxel spacing to 1Å. The accessible interaction volume was visualised using UCSF

Chimera [101].

**Circular dichroism (CD) spectroscopy**

In order to evaluate the secondary structure content of GST-tagged $Vpr_{mus}$, and the double mutants

R15E/R75E and W29A/A66W, CD spectroscopy was performed. Protein samples were dialysed

overnight at 4°C against CD buffer (100 mM NaF, 10 mM $K_2HPO_4/KH_2PO_4$ mixture, pH 8.5), and

protein concentration then adjusted to 0.2 mg/ml. For each sample, three replicate CD spectra were

recorded in 1.0 nm steps over the range of 190-260 nm. Measurements were performed in 0.05 cm path-

length quartz cuvettes (Hellma, Mühlheim, Germany) at 20°C on a Chirascan spectropolarimeter

(Applied Photophysics, London, UK). A reference spectrum of CD buffer was subtracted from the

averaged protein sample spectrum. Measured ellipticity in millidegrees (m°) was converted to molar

ellipticity [Θ] according to equation $\Theta = m° * M/(10 \times L \times C)$, where M is the average protein molecular

weight, L is path length of the measurement cell, and C is the molar protein concentration. Molar

ellipticity was plotted against wavelength using SigmaPlot version 14.0 (Systat Software Inc., San Jose,

California, USA).

**Acknowledgments**

We thank the MPI-MG for granting access to the TEM instruments of the microscopy and cryo-EM

service group. We thank Dr. Manfred Weiss and the scientific staff of the BESSY-MX (Macromolecular

X-ray Crystallography)/Helmholtz Zentrum Berlin für Materialien und Energie at beamlines BL14.1,

BL14.2, and BL14.3 operated by the Joint Berlin MX-Laboratory at the BESSY II electron storage ring

(Berlin-Adlershof, Germany) as well as the scientific staff of the ESRF (Grenoble, France) at beamlines

ID30A-3, ID30B, ID23-1, ID23-2, and ID29 for continuous support. We acknowledge Diamond Light

Source (Didcot, UK) for access and support of the synchrotron beamline I04 and cryo-EM facilities at

the UK's national Electron Bio-imaging Centre (eBIC). Furthermore, the authors acknowledge the

North-German Supercomputing Alliance (HLRN) and the HPC for Research cluster of the Berlin

Institute of Health for providing HPC resources. We are grateful to Prof. Udo Heinemann and Jianhui

Wang (Max-Delbrück-Centrum, Berlin, Germany) for access to and assistance during CD spectroscopy.

The pHisSUMO plasmid was a generous gift from Dr. Evangelos Christodoulou (The Francis Crick

Institute, London, UK). The rhesus macaque SAMHD1 cDNA template was a generous gift from Prof.

Michael Emerman (Fred Hutchinson Cancer Research Center, Seattle, USA). Recombinant

BAC10:1629KO bacmid was a generous gift from Prof. Ian Jones (University of Reading, UK).

pAcGHLT-B-DDB1 was a gift from Prof. Ning Zheng (Addgene plasmid 48638). pET28-mE1 was a

gift from Prof. Jorge Eduardo Azevedo (Addgene plasmid 32534).

## 790    Data availability

The coordinates and structure factors for the crystal structures have been deposited at the Protein Data

Bank (PDB) with the accession codes 6ZUE (DDB1/DCAF1-CtD) and 6ZX9 (DDB1/DCAF1-

CtD/T4L-Vpr$_{mus}$ 1-92). Cryo-EM reconstructions have been deposited at the Electron Microscopy Data

Bank (EMDB) with the accession codes EMD-10611 (core), EMD-10612 (conformational state-1),

EMD-10613 (state-2) and EMD-10614 (state-3). CLMS data have been deposited at the PRIDE database

[114] with the accession code PXD020453, reviewer password fCrQG2u8.

# References

1. Randow F, Lehner PJ. Viral avoidance and exploitation of the ubiquitin system. Nat Cell Biol. 2009;11(5):527-34. doi: 10.1038/ncb0509-527.

2. Isaacson MK, Ploegh HL. Ubiquitination, ubiquitin-like modifiers, and deubiquitination in viral infection. Cell host & microbe. 2009;5(6):559-70. doi: 10.1016/j.chom.2009.05.012.

3. Gustin JK, Moses AV, Fruh K, Douglas JL. Viral takeover of the host ubiquitin system. Front Microbiol. 2011;2:161. doi: 10.3389/fmicb.2011.00161.

4. Barry M, Fruh K. Viral modulators of cullin RING ubiquitin ligases: culling the host defense. Science's STKE : signal transduction knowledge environment. 2006;2006(335):pe21. Epub 2006/05/18. doi: 10.1126/stke.3352006pe21.

5. Mahon C, Krogan NJ, Craik CS, Pick E. Cullin E3 ligases and their rewiring by viral factors. Biomolecules. 2014;4(4):897-930. Epub 2014/10/15. doi: 10.3390/biom4040897.

6. Becker T, Le-Trilling VTK, Trilling M. Cellular Cullin RING Ubiquitin Ligases: Druggable Host Dependency Factors of Cytomegaloviruses. Int J Mol Sci. 2019;20(7). doi: 10.3390/ijms20071636.

7. Seissler T, Marquet R, Paillart JC. Hijacking of the Ubiquitin/Proteasome Pathway by the HIV Auxiliary Proteins. Viruses. 2017;9(11). doi: 10.3390/v9110322.

8. Zheng N, Shabek N. Ubiquitin Ligases: Structure, Function, and Regulation. Annu Rev Biochem. 2017;86:14.1-29.

9. Sauter D, Kirchhoff F. Key Viral Adaptations Preceding the AIDS Pandemic. Cell host & microbe. 2019;25(1):27-38. doi: 10.1016/j.chom.2018.12.002.

10. Sharp PM, Hahn BH. Origins of HIV and the AIDS pandemic. Cold Spring Harbor perspectives in medicine. 2011;1(1):a006841. Epub 2012/01/10. doi: 10.1101/cshperspect.a006841.

11. Hatziioannou T, Del Prete GQ, Keele BF, Estes JD, McNatt MW, Bitzegeio J, et al. HIV-1-induced AIDS in monkeys. Science. 2014;344(6190):1401-5. Epub 2014/06/21. doi: 10.1126/science.1250761.

12. Malim MH, Bieniasz PD. HIV Restriction Factors and Mechanisms of Evasion. Cold Spring Harbor perspectives in medicine. 2012;2(5):a006940. Epub 2012/05/04. doi: 10.1101/cshperspect.a006940.

13. Fischer ES, Scrima A, Bohm K, Matsumoto S, Lingaraju GM, Faty M, et al. The molecular basis of CRL4DDB2/CSA ubiquitin ligase architecture, targeting, and activation. Cell. 2011;147(5):1024-39. Epub 2011/11/29. doi: 10.1016/j.cell.2011.10.035.

14.    Lee J, Zhou P. DCAFs, the missing link of the CUL4-DDB1 ubiquitin ligase. Molecular cell.

2007;26(6):775-80. Epub 2007/06/26. doi: 10.1016/j.molcel.2007.06.001.

15.    Angers S, Li T, Yi X, MacCoss MJ, Moon RT, Zheng N. Molecular architecture and assembly of the

DDB1-CUL4A ubiquitin ligase machinery. Nature. 2006;443(7111):590-3. Epub 2006/09/12. doi:

10.1038/nature05175.

16.    Scrima A, Konickova R, Czyzewski BK, Kawasaki Y, Jeffrey PD, Groisman R, et al. Structural basis of

UV DNA-damage recognition by the DDB1-DDB2 complex. Cell. 2008;135(7):1213-23. Epub 2008/12/27. doi:

10.1016/j.cell.2008.10.045.

17.    Zimmerman ES, Schulman BA, Zheng N. Structural assembly of cullin-RING ubiquitin ligase

complexes. Current opinion in structural biology. 2010;20(6):714-21. Epub 2010/10/01. doi:

10.1016/j.sbi.2010.08.010.

18.    Andrejeva J, Young DF, Goodbourn S, Randall RE. Degradation of STAT1 and STAT2 by the V

proteins of simian virus 5 and human parainfluenza virus type 2, respectively: consequences for virus replication

in the presence of alpha/beta and gamma interferons. Journal of virology. 2002;76(5):2159-67. doi:

10.1128/jvi.76.5.2159-2167.2002.

19.    Li T, Chen X, Garbutt KC, Zhou P, Zheng N. Structure of DDB1 in complex with a paramyxovirus V

protein: viral hijack of a propeller cluster in ubiquitin ligase. Cell. 2006;124(1):105-17. Epub 2006/01/18. doi:

10.1016/j.cell.2005.10.033.

20.    Trilling M, Le VT, Fiedler M, Zimmermann A, Bleifuss E, Hengel H. Identification of DNA-damage

DNA-binding protein 1 as a conditional essential factor for cytomegalovirus replication in interferon-gamma-

stimulated cells. PLoS pathogens. 2011;7(6):e1002069. doi: 10.1371/journal.ppat.1002069.

21.    Paradkar PN, Duchemin JB, Rodriguez-Andres J, Trinidad L, Walker PJ. Cullin4 Is Pro-Viral during

West Nile Virus Infection of Culex Mosquitoes. PLoS pathogens. 2015;11(9):e1005143. doi:

10.1371/journal.ppat.1005143.

22.    Decorsiere A, Mueller H, van Breugel PC, Abdul F, Gerossier L, Beran RK, et al. Hepatitis B virus X

protein identifies the Smc5/6 complex as a host restriction factor. Nature. 2016;531(7594):386-9. doi:

10.1038/nature17170.

23.    Murphy CM, Xu Y, Li F, Nio K, Reszka-Blanco N, Li X, et al. Hepatitis B Virus X Protein Promotes

Degradation of SMC5/6 to Enhance HBV Replication. Cell reports. 2016;16(11):2846-54. doi:

10.1016/j.celrep.2016.08.026.

24.    Lim ES, Fregoso OI, McCoy CO, Matsen FA, Malik HS, Emerman M. The ability of primate

lentiviruses to degrade the monocyte restriction factor SAMHD1 preceded the birth of the viral accessory protein

Vpx. Cell host & microbe. 2012;11(2):194-204. Epub 2012/01/31. doi: 10.1016/j.chom.2012.01.004.

25.    Romani B, Cohen EA. Lentivirus Vpr and Vpx accessory proteins usurp the cullin4-DDB1 (DCAF1)

E3 ubiquitin ligase. Current opinion in virology. 2012;2(6):755-63. Epub 2012/10/16. doi:

10.1016/j.coviro.2012.09.010.

26.    Fabryova H, Strebel K. Vpr and Its Cellular Interaction Partners: R We There Yet? Cells. 2019;8(11).

doi: 10.3390/cells8111310.

27.    Greenwood EJD, Williamson JC, Sienkiewicz A, Naamati A, Matheson NJ, Lehner PJ. Promiscuous

Targeting of Cellular Proteins by Vpr Drives Systems-Level Proteomic Remodeling in HIV-1 Infection. Cell

reports. 2019;27(5):1579-96 e7. doi: 10.1016/j.celrep.2019.04.025.

28.    Schrofelbauer B, Yu Q, Zeitlin SG, Landau NR. Human immunodeficiency virus type 1 Vpr induces the

degradation of the UNG and SMUG uracil-DNA glycosylases. Journal of virology. 2005;79(17):10978-87. doi:

10.1128/JVI.79.17.10978-10987.2005.

29.    Lahouassa H, Blondot ML, Chauveau L, Chougui G, Morel M, Leduc M, et al. HIV-1 Vpr degrades the

HLTF DNA translocase in T cells and macrophages. Proceedings of the National Academy of Sciences of the

United States of America. 2016;113(19):5311-6. doi: 10.1073/pnas.1600485113.

30.    Laguette N, Bregnard C, Hue P, Basbous J, Yatim A, Larroque M, et al. Premature activation of the

SLX4 complex by Vpr promotes G2/M arrest and escape from innate immune sensing. Cell. 2014;156(1-2):134-

45. Epub 2014/01/15. doi: 10.1016/j.cell.2013.12.011.

31.    Zhou X, DeLucia M, Ahn J. SLX4-SLX1 Protein-independent Down-regulation of MUS81-EME1

Protein by HIV-1 Viral Protein R (Vpr). The Journal of biological chemistry. 2016;291(33):16936-47. doi:

10.1074/jbc.M116.721183.

32.    Romani B, Shaykh Baygloo N, Aghasadeghi MR, Allahbakhshi E. HIV-1 Vpr Protein Enhances

Proteasomal Degradation of MCM10 DNA Replication Factor through the Cul4-DDB1[VprBP] E3 Ubiquitin

Ligase to Induce G2/M Cell Cycle Arrest. The Journal of biological chemistry. 2015;290(28):17380-9. doi:

10.1074/jbc.M115.641522.

33.    Lv L, Wang Q, Xu Y, Tsao LC, Nakagawa T, Guo H, et al. Vpr Targets TET2 for Degradation by

CRL4(VprBP) E3 Ligase to Sustain IL-6 Expression and Enhance HIV-1 Replication. Molecular cell.

2018;70(5):961-70 e5. doi: 10.1016/j.molcel.2018.05.007.

34.    Su J, Rui Y, Lou M, Yin L, Xiong H, Zhou Z, et al. HIV-2/SIV Vpx targets a novel functional domain

of STING to selectively inhibit cGAS-STING-mediated NF-kappaB signalling. Nat Microbiol. 2019;4(12):2552-

64. doi: 10.1038/s41564-019-0585-4.

35.    Chougui G, Munir-Matloob S, Matkovic R, Martin MM, Morel M, Lahouassa H, et al. HIV-2/SIV viral

protein X counteracts HUSH repressor complex. Nat Microbiol. 2018;3(8):891-7. doi: 10.1038/s41564-018-

0179-6.

36.    Yurkovetskiy L, Guney MH, Kim K, Goh SL, McCauley S, Dauphin A, et al. Primate

immunodeficiency virus proteins Vpx and Vpr counteract transcriptional repression of proviruses by the HUSH

complex. Nat Microbiol. 2018;3(12):1354-61. doi: 10.1038/s41564-018-0256-x.

37.    Hrecka K, Hao C, Gierszewska M, Swanson SK, Kesik-Brodacka M, Srivastava S, et al. Vpx relieves

inhibition of HIV-1 infection of macrophages mediated by the SAMHD1 protein. Nature. 2011;474(7353):658-

61. Epub 2011/07/02. doi: 10.1038/nature10195.

38.    Laguette N, Sobhian B, Casartelli N, Ringeard M, Chable-Bessia C, Segeral E, et al. SAMHD1 is the

dendritic- and myeloid-cell-specific HIV-1 restriction factor counteracted by Vpx. Nature. 2011;474(7353):654-

7. Epub 2011/05/27. doi: 10.1038/nature10117.

39.    Powell RD, Holland PJ, Hollis T, Perrino FW. Aicardi-Goutieres syndrome gene and HIV-1 restriction

factor SAMHD1 is a dGTP-regulated deoxynucleotide triphosphohydrolase. The Journal of biological chemistry.

2011;286(51):43596-600. Epub 2011/11/10. doi: 10.1074/jbc.C111.317628.

40.    Goldstone DC, Ennis-Adeniran V, Hedden JJ, Groom HC, Rice GI, Christodoulou E, et al. HIV-1

restriction factor SAMHD1 is a deoxynucleoside triphosphate triphosphohydrolase. Nature.

2011;480(7377):379-82. Epub 2011/11/08. doi: 10.1038/nature10623.

41.    Zhu C, Gao W, Zhao K, Qin X, Zhang Y, Peng X, et al. Structural insight into dGTP-dependent

activation of tetrameric SAMHD1 deoxynucleoside triphosphate triphosphohydrolase. Nature communications.

2013;4:2722. Epub 2013/11/13. doi: 10.1038/ncomms3722.

42.    Kim B, Nguyen LA, Daddacha W, Hollenbaugh JA. Tight interplay among SAMHD1 protein level,

cellular dNTP levels, and HIV-1 proviral DNA synthesis kinetics in human primary monocyte-derived

macrophages. The Journal of biological chemistry. 2012;287(26):21570-4. Epub 2012/05/17. doi:

10.1074/jbc.C112.374843.

43.    Lahouassa H, Daddacha W, Hofmann H, Ayinde D, Logue EC, Dragin L, et al. SAMHD1 restricts the

replication of human immunodeficiency virus type 1 by depleting the intracellular pool of deoxynucleoside

triphosphates. Nature immunology. 2012;13(3):223-8. Epub 2012/02/14. doi: 10.1038/ni.2236.

44. St Gelais C, de Silva S, Amie SM, Coleman CM, Hoy H, Hollenbaugh JA, et al. SAMHD1 restricts

HIV-1 infection in dendritic cells (DCs) by dNTP depletion, but its expression in DCs and primary CD4+ T-

lymphocytes cannot be upregulated by interferons. Retrovirology. 2012;9:105. Epub 2012/12/13. doi:

10.1186/1742-4690-9-105.

45. Rehwinkel J, Maelfait J, Bridgeman A, Rigby R, Hayward B, Liberatore RA, et al. SAMHD1-

dependent retroviral control and escape in mice. The EMBO journal. 2013;32(18):2454-62. Epub 2013/07/23.

doi: 10.1038/emboj.2013.163.

46. Morris ER, Taylor IA. The missing link: allostery and catalysis in the anti-viral protein SAMHD1.

Biochem Soc Trans. 2019;47(4):1013-27. doi: 10.1042/BST20180348.

47. Baldauf HM, Pan X, Erikson E, Schmidt S, Daddacha W, Burggraf M, et al. SAMHD1 restricts HIV-1

infection in resting CD4(+) T cells. Nature medicine. 2012;18(11):1682-7. Epub 2012/09/14. doi:

10.1038/nm.2964.

48. Shingai M, Welbourn S, Brenchley JM, Acharya P, Miyagi E, Plishka RJ, et al. The Expression of

Functional Vpx during Pathogenic SIVmac Infections of Rhesus Macaques Suppresses SAMHD1 in CD4+

Memory T Cells. PLoS pathogens. 2015;11(5):e1004928. doi: 10.1371/journal.ppat.1004928.

49. Fregoso OI, Ahn J, Wang C, Mehrens J, Skowronski J, Emerman M. Evolutionary toggling of Vpx/Vpr

specificity results in divergent recognition of the restriction factor SAMHD1. PLoS pathogens.

2013;9(7):e1003496. Epub 2013/07/23. doi: 10.1371/journal.ppat.1003496.

50. Schwefel D, Groom HC, Boucherit VC, Christodoulou E, Walker PA, Stoye JP, et al. Structural basis of

lentiviral subversion of a cellular protein degradation pathway. Nature. 2014;505(7482):234-8. Epub 2013/12/18.

doi: 10.1038/nature12815.

51. Schwefel D, Boucherit VC, Christodoulou E, Walker PA, Stoye JP, Bishop KN, et al. Molecular

Determinants for Recognition of Divergent SAMHD1 Proteins by the Lentiviral Accessory Protein Vpx. Cell

host & microbe. 2015;17(4):489-99. Epub 2015/04/10. doi: 10.1016/j.chom.2015.03.004.

52. Wu Y, Koharudin LM, Mehrens J, DeLucia M, Byeon CH, Byeon IJ, et al. Structural Basis of Clade-

specific Engagement of SAMHD1 (Sterile alpha Motif and Histidine/Aspartate-containing Protein 1) Restriction

Factors by Lentiviral Viral Protein X (Vpx) Virulence Factors. The Journal of biological chemistry.

2015;290(29):17935-45. doi: 10.1074/jbc.M115.665513.

53. Spragg CJ, Emerman M. Antagonism of SAMHD1 is actively maintained in natural infections of simian

immunodeficiency virus. Proceedings of the National Academy of Sciences of the United States of America.

2013;110(52):21136-41. Epub 2013/12/11. doi: 10.1073/pnas.1316839110.

54.      Wu Y, Zhou X, Barnes CO, DeLucia M, Cohen AE, Gronenborn AM, et al. The DDB1-DCAF1-Vpr-

UNG2 crystal structure reveals how HIV-1 Vpr steers human UNG2 toward destruction. Nature structural &

molecular biology. 2016;23(10):933-40. doi: 10.1038/nsmb.3284.

55.      Enchev RI, Schulman BA, Peter M. Protein neddylation: beyond cullin-RING ligases. Nature reviews

Molecular cell biology. 2015;16(1):30-44. doi: 10.1038/nrm3919.

56.      Duda DM, Borg LA, Scott DC, Hunt HW, Hammel M, Schulman BA. Structural insights into NEDD8

activation of cullin-RING ligases: conformational control of conjugation. Cell. 2008;134(6):995-1006. doi:

10.1016/j.cell.2008.07.022.

57.      Fischer ES, Bohm K, Lydeard JR, Yang H, Stadler MB, Cavadini S, et al. Structure of the DDB1-

CRBN E3 ubiquitin ligase in complex with thalidomide. Nature. 2014;512(7512):49-53. doi:

10.1038/nature13527.

58.      Schneider M, Belsom A, Rappsilber J. Protein Tertiary Structure by Crosslinking/Mass Spectrometry.

Trends in biochemical sciences. 2018;43(3):157-69. Epub 2018/02/06. doi: 10.1016/j.tibs.2017.12.006.

59.      O'Reilly FJ, Rappsilber J. Cross-linking mass spectrometry: methods and applications in structural,

molecular and systems biology. Nature structural & molecular biology. 2018;25(11):1000-8. Epub 2018/10/31.

doi: 10.1038/s41594-018-0147-0.

60.      van Zundert GC, Bonvin AM. DisVis: quantifying and visualizing accessible interaction space of

distance-restrained biomolecular complexes. Bioinformatics. 2015;31(19):3222-4. Epub 2015/05/31. doi:

10.1093/bioinformatics/btv333.

61.      van Zundert GC, Trellet M, Schaarschmidt J, Kurkcuoglu Z, David M, Verlato M, et al. The DisVis and

PowerFit Web Servers: Explorative and Integrative Modeling of Biomolecular Complexes. Journal of molecular

biology. 2017;429(3):399-407. Epub 2016/12/13. doi: 10.1016/j.jmb.2016.11.032.

62.      DeLucia M, Mehrens J, Wu Y, Ahn J. HIV-2 and SIVmac accessory virulence factor Vpx down-

regulates SAMHD1 enzyme catalysis prior to proteasome-dependent degradation. The Journal of biological

chemistry. 2013;288(26):19116-26. doi: 10.1074/jbc.M113.469007.

63.      Yan J, Kaur S, DeLucia M, Hao C, Mehrens J, Wang C, et al. Tetramerization of SAMHD1 is required

for biological activity and inhibition of HIV infection. The Journal of biological chemistry. 2013;288(15):10406-

17. Epub 2013/02/22. doi: 10.1074/jbc.M112.443796.

64.      Arnold LH, Groom HC, Kunzelmann S, Schwefel D, Caswell SJ, Ordonez P, et al. Phospho-dependent

Regulation of SAMHD1 Oligomerisation Couples Catalysis and Restriction. PLoS pathogens.

2015;11(10):e1005194. Epub 2015/10/03. doi: 10.1371/journal.ppat.1005194.

65.    Yan J, Hao C, DeLucia M, Swanson S, Florens L, Washburn MP, et al. CyclinA2-Cyclin-dependent

Kinase Regulates SAMHD1 Protein Phosphohydrolase Domain. The Journal of biological chemistry.

2015;290(21):13279-92. Epub 2015/04/08. doi: 10.1074/jbc.M115.646588.

66.    Berger G, Lawrence M, Hue S, Neil SJ. G2/M cell cycle arrest correlates with primate lentiviral Vpr

interaction with the SLX4 complex. Journal of virology. 2014. Epub 2014/10/17. doi: 10.1128/JVI.02307-14.

67.    Daugherty MD, Malik HS. Rules of engagement: molecular insights from host-virus arms races. Annual

review of genetics. 2012;46:677-700. Epub 2012/11/14. doi: 10.1146/annurev-genet-110711-155522.

68.    Duggal NK, Emerman M. Evolutionary conflicts between viruses and restriction factors shape

immunity. Nat Rev Immunol. 2012;12(10):687-95. Epub 2012/09/15. doi: 10.1038/nri3295.

69.    Uriu K, Kosugi Y, Ito J, Sato K. The Battle between Retroviruses and APOBEC3 Genes: Its Past and

Present. Viruses. 2021;13(1). Epub 2021/01/23. doi: 10.3390/v13010124.

70.    Guenzel CA, Herate C, Benichou S. HIV-1 Vpr-a still "enigmatic multitasker". Front Microbiol.

2014;5:127. doi: 10.3389/fmicb.2014.00127.

71.    Stivahtis GL, Soares MA, Vodicka MA, Hahn BH, Emerman M. Conservation and host specificity of

Vpr-mediated cell cycle arrest suggest a fundamental role in primate lentivirus evolution and biology. Journal of

virology. 1997;71(6):4331-8.

72.    Planelles V, Jowett JB, Li QX, Xie Y, Hahn B, Chen IS. Vpr-induced cell cycle arrest is conserved

among primate lentiviruses. Journal of virology. 1996;70(4):2516-24.

73.    Baek K, Krist DT, Prabu JR, Hill S, Klugel M, Neumaier LM, et al. NEDD8 nucleates a multivalent

cullin-RING-UBE2D ubiquitin ligation assembly. Nature. 2020;578(7795):461-6. doi: 10.1038/s41586-020-

2000-y.

74.    Schapira M, Calabrese MF, Bullock AN, Crews CM. Targeted protein degradation: expanding the

toolbox. Nat Rev Drug Discov. 2019;18(12):949-63. doi: 10.1038/s41573-019-0047-y.

75.    Hanzl A, Winter GE. Targeted protein degradation: current and future challenges. Curr Opin Chem

Biol. 2020;56:35-41. doi: 10.1016/j.cbpa.2019.11.012.

76.    Antiretroviral Therapy Cohort C. Life expectancy of individuals on combination antiretroviral therapy

in high-income countries: a collaborative analysis of 14 cohort studies. Lancet. 2008;372(9635):293-9. Epub

2008/07/29. doi: 10.1016/S0140-6736(08)61113-7.

77.    Gunthard HF, Calvez V, Paredes R, Pillay D, Shafer RW, Wensing AM, et al. Human

Immunodeficiency Virus Drug Resistance: 2018 Recommendations of the International Antiviral Society-USA

Panel. Clin Infect Dis. 2019;68(2):177-87. Epub 2018/07/28. doi: 10.1093/cid/ciy463.

78.    Miller RH, Sarver N. HIV accessory proteins as therapeutic targets. Nature medicine. 1997;3(4):389-94.

Epub 1997/04/01. doi: 10.1038/nm0497-389.

79.    Puhl AC, Garzino Demo A, Makarov VA, Ekins S. New targets for HIV drug discovery. Drug Discov

Today. 2019;24(5):1139-47. Epub 2019/03/20. doi: 10.1016/j.drudis.2019.03.013.

80.    Nathans R, Cao H, Sharova N, Ali A, Sharkey M, Stranska R, et al. Small-molecule inhibition of HIV-1

Vif. Nat Biotechnol. 2008;26(10):1187-92. Epub 2008/09/23. doi: 10.1038/nbt.1496.

81.    Zuo T, Liu D, Lv W, Wang X, Wang J, Lv M, et al. Small-molecule inhibition of human

immunodeficiency virus type 1 replication by targeting the interaction between Vif and ElonginC. Journal of

virology. 2012;86(10):5497-507. Epub 2012/03/02. doi: 10.1128/JVI.06957-11.

82.    Ma L, Zhang Z, Liu Z, Pan Q, Wang J, Li X, et al. Identification of small molecule compounds

targeting the interaction of HIV-1 Vif and human APOBEC3G by virtual screening and biological evaluation.

Scientific reports. 2018;8(1):8067. Epub 2018/05/26. doi: 10.1038/s41598-018-26318-3.

83.    Gonzalez ME. The HIV-1 Vpr Protein: A Multifaceted Target for Therapeutic Intervention. Int J Mol

Sci. 2017;18(1). Epub 2017/01/12. doi: 10.3390/ijms18010126.

84.    Zhao Y, Chapman DA, Jones IM. Improving baculovirus recombination. Nucleic acids research.

2003;31(2):E6-. Epub 2003/01/16.

85.    Wilkins MR, Gasteiger E, Bairoch A, Sanchez JC, Williams KL, Appel RD, et al. Protein identification

and analysis tools in the ExPASy server. Methods Mol Biol. 1999;112:531-52. Epub 1999/02/23. doi: 10.1385/1-

59259-584-7:531.

86.    Kabsch W. Xds. Acta crystallographica Section D, Biological crystallography. 2010;66(Pt 2):125-32.

Epub 2010/02/04. doi: 10.1107/S0907444909047337.

87.    Vagin A, Teplyakov A. Molecular replacement with MOLREP. Acta crystallographica Section D,

Biological crystallography. 2010;66(Pt 1):22-5. Epub 2010/01/09. doi: 10.1107/S0907444909042589.

88.    Emsley P, Cowtan K. Coot: model-building tools for molecular graphics. Acta crystallographica Section

D, Biological crystallography. 2004;60(Pt 12 Pt 1):2126-32. Epub 2004/12/02. doi:

10.1107/S0907444904019158.

89.    Liebschner D, Afonine PV, Baker ML, Bunkoczi G, Chen VB, Croll TI, et al. Macromolecular structure

determination using X-rays, neutrons and electrons: recent developments in Phenix. Acta Crystallogr D Struct

Biol. 2019;75(Pt 10):861-77. Epub 2019/10/08. doi: 10.1107/S2059798319011471.

90.    Sparta KM, Krug M, Heinemann U, Mueller U, Weiss MS. XDSAPP2.0. Journal of Applied

Crystallography. 2016;49(3):1085-92. doi: doi:10.1107/S1600576716004416.

91. McCoy AJ, Grosse-Kunstleve RW, Adams PD, Winn MD, Storoni LC, Read RJ. Phaser

crystallographic software. Journal of Applied Crystallography. 2007;40(4):658-74. doi:

doi:10.1107/S0021889807021206.

92. Kuroki R, Weaver LH, Matthews BW. Structural basis of the conversion of T4 lysozyme into a

transglycosidase by reengineering the active site. Proceedings of the National Academy of Sciences of the

United States of America. 1999;96(16):8949-54. Epub 1999/08/04. doi: 10.1073/pnas.96.16.8949.

93. Morellet N, Bouaziz S, Petitjean P, Roques BP. NMR structure of the HIV-1 regulatory protein VPR.

Journal of molecular biology. 2003;327(1):215-27. Epub 2003/03/05.

94. Carragher B, Kisseberth N, Kriegman D, Milligan RA, Potter CS, Pulokas J, et al. Leginon: an

automated system for acquisition of images from vitreous ice specimens. J Struct Biol. 2000;132(1):33-45. Epub

2000/12/21. doi: 10.1006/jsbi.2000.4314.

95. Suloway C, Pulokas J, Fellmann D, Cheng A, Guerra F, Quispe J, et al. Automated molecular

microscopy: the new Leginon system. J Struct Biol. 2005;151(1):41-60. Epub 2005/05/14. doi:

10.1016/j.jsb.2005.03.010.

96. Moriya T, Saur M, Stabrin M, Merino F, Voicu H, Huang Z, et al. High-resolution Single Particle

Analysis from Electron Cryo-microscopy Images Using SPHIRE. J Vis Exp. 2017;(123). Epub 2017/06/02. doi:

10.3791/55448.

97. Grant T, Rohou A, Grigorieff N. cisTEM, user-friendly software for single-particle image processing.

Elife. 2018;7. Epub 2018/03/08. doi: 10.7554/eLife.35383.

98. Zivanov J, Nakane T, Forsberg BO, Kimanius D, Hagen WJ, Lindahl E, et al. New tools for automated

high-resolution cryo-EM structure determination in RELION-3. Elife. 2018;7. Epub 2018/11/10. doi:

10.7554/eLife.42166.

99. Zheng SQ, Palovcak E, Armache JP, Verba KA, Cheng Y, Agard DA. MotionCor2: anisotropic

correction of beam-induced motion for improved cryo-electron microscopy. Nat Methods. 2017;14(4):331-2.

Epub 2017/03/03. doi: 10.1038/nmeth.4193.

100. Mindell JA, Grigorieff N. Accurate determination of local defocus and specimen tilt in electron

microscopy. J Struct Biol. 2003;142(3):334-47. Epub 2003/06/05. doi: 10.1016/s1047-8477(03)00069-8.

101. Pettersen EF, Goddard TD, Huang CC, Couch GS, Greenblatt DM, Meng EC, et al. UCSF Chimera--a

visualization system for exploratory research and analysis. J Comput Chem. 2004;25(13):1605-12. Epub

2004/07/21. doi: 10.1002/jcc.20084.

102. Punjani A, Rubinstein JL, Fleet DJ, Brubaker MA. cryoSPARC: algorithms for rapid unsupervised cryo-EM structure determination. Nat Methods. 2017;14(3):290-6. Epub 2017/02/07. doi: 10.1038/nmeth.4169.

103. Hayward S, Lee RA. Improvements in the analysis of domain motions in proteins from conformational change: DynDom version 1.50. J Mol Graph Model. 2002;21(3):181-3. Epub 2002/12/05. doi: 10.1016/s1093-3263(02)00140-7.

104. Krissinel E, Henrick K. Inference of macromolecular assemblies from crystalline state. Journal of molecular biology. 2007;372(3):774-97. Epub 2007/08/08. doi: 10.1016/j.jmb.2007.05.022.

105. Madeira F, Park YM, Lee J, Buso N, Gur T, Madhusoodanan N, et al. The EMBL-EBI search and sequence analysis tools APIs in 2019. Nucleic acids research. 2019;47(W1):W636-W41. Epub 2019/04/13. doi: 10.1093/nar/gkz268.

106. Nicholas KB, Nicholas Jr., H. B., Deerfield II., D. W. GeneDoc: Analysis and Visualization of Genetic Variation. embnetnews. 1997;4(2):1-4.

107. Shevchenko A, Tomas H, Havlis J, Olsen JV, Mann M. In-gel digestion for mass spectrometric characterization of proteins and proteomes. Nature protocols. 2006;1(6):2856-60. Epub 2007/04/05. doi: 10.1038/nprot.2006.468.

108. Rappsilber J, Ishihama Y, Mann M. Stop and go extraction tips for matrix-assisted laser desorption/ionization, nanoelectrospray, and LC/MS sample pretreatment in proteomics. Anal Chem. 2003;75(3):663-70. Epub 2003/02/15. doi: 10.1021/ac026117i.

109. Kolbowski L, Mendes ML, Rappsilber J. Optimizing the Parameters Governing the Fragmentation of Cross-Linked Peptides in a Tribrid Mass Spectrometer. Anal Chem. 2017;89(10):5311-8. Epub 2017/04/14. doi: 10.1021/acs.analchem.6b04935.

110. Mendes ML, Fischer L, Chen ZA, Barbon M, O'Reilly FJ, Giese SH, et al. An integrated workflow for crosslinking mass spectrometry. Mol Syst Biol. 2019;15(9):e8994. Epub 2019/09/27. doi: 10.15252/msb.20198994.

111. Fischer L, Rappsilber J. Quirks of Error Estimation in Cross-Linking/Mass Spectrometry. Anal Chem. 2017;89(7):3829-33. Epub 2017/03/08. doi: 10.1021/acs.analchem.6b03745.

112. Yang J, Zhang Y. Protein Structure and Function Prediction Using I-TASSER. Curr Protoc Bioinformatics. 2015;52:5 8 1-5 8 15. Epub 2015/12/19. doi: 10.1002/0471250953.bi0508s52.

113. Webb B, Sali A. Comparative Protein Structure Modeling Using MODELLER. Curr Protoc Bioinformatics. 2016;54:5 6 1-5 6 37. Epub 2016/06/21. doi: 10.1002/cpbi.3.

114.    Perez-Riverol Y, Csordas A, Bai J, Bernal-Llinares M, Hewapathirana S, Kundu DJ, et al. The PRIDE

database and related tools and resources in 2019: improving support for quantification data. Nucleic acids

research. 2019;47(D1):D442-D50. Epub 2018/11/06. doi: 10.1093/nar/gky1106.

115.    Shabek N, Ruble J, Waston CJ, Garbutt KC, Hinds TR, Li T, et al. Structural insights into DDA1

function as a core component of the CRL4-DDB1 ubiquitin ligase. Cell Discov. 2018;4:67. Epub 2018/12/20.

doi: 10.1038/s41421-018-0064-8.

1107

 **Figures**

**Fig 1. Biochemical analysis of Vpr_mus-induced CRL4^DCAF1 specificity redirection.**

(**A-C**) GF analysis of *in vitro* reconstitution of protein complexes containing DDB1/DCAF1-CtD, Vpr_mus and SAMHD1 (**A**), SAMHD1-ΔCtD (**B**) or T4L-SAMHD1-CtD (**C**). Elution volumes of protein molecular weight standards are indicated above the chromatogram in **A**. Coomassie blue-stained SDS-PAGE analyses of fractions collected during the GF runs are shown below the chromatograms, with boxes colour-coded with respect to the chromatograms. SAM – sterile α-motif domain, HD – histidine-aspartate domain, T4L – T4 Lysozyme. The asterisk and double asterisk indicate slight contaminations with remaining GST-3C protease and the GST purification tag, respectively. (**D-G**) *In vitro* ubiquitylation reactions with purified protein components in the absence (**D**) or presence (**E-G**) of Vpr_mus, with the indicated SAMHD1 constructs as substrate. Reactions were stopped after the indicated times, separated on SDS-PAGE and visualised by Coomassie blue staining.

**Fig 2. Crystal structure of the DDB1/DCAF1-CtD/Vpr_mus complex.**

(**A**) Overall structure of the DDB1/DCAF1-CtD/Vpr_mus complex in two views. DCAF1-CtD is shown

as grey cartoon and semi-transparent surface. Vpr_mus is shown as a dark green cartoon with the co-

ordinated zinc ion shown as grey sphere. T4L and DDB1 have been omitted for clarity. (**B**)

Superposition of apo-DCAF1-CtD (light blue cartoon) with Vpr_mus-bound DCAF1-CtD (grey/green

cartoon). Only DCAF1-CtD regions with significant structural differences between apo- and Vpr$_{mus}$-

bound forms are shown. Disordered loops are indicated as dashed lines. (**C**) Comparison of the binary

Vpr$_{mus}$/DCAF1-CtD and ternary Vpx$_{sm}$/DCAF1-CtD/SAMHD1-CtD complexes. For DCAF1-CtD, only

the N-terminal "acidic loop" region is shown. Vpr$_{mus}$, DCAF1-CtD and bound zinc are coloured as in

**A**; Vpx$_{sm}$ is represented as orange cartoon and SAMHD1-CtD as pink cartoon. Selected

Vpr/Vpx/DCAF1-CtD side chains are shown as sticks, and electrostatic interactions between these side

chains are indicated as dotted lines. (**D**) GF analysis of *in vitro* reconstitution of protein complexes

containing DDB1/DCAF1-CtD/Vpr$_{mus}$ or the Vpr$_{mus}$ R15E/R75E mutant, and SAMHD1. SDS-PAGE

analyses of corresponding GF fractions are shown below the chromatogram, with boxes colour-coded

with respect to the chromatogram. (**E-F**) *In vitro* reconstitution of protein complexes containing

SAMHD1 and Vpr$_{mus}$ R15E/R75E (**E**) or DDB1/DCAF1-CtD and Vpr$_{mus}$ R15E/R75E (**F**). SDS-PAGE

analyses of corresponding GF fractions are shown below the chromatogram, with boxes colour-coded

with respect to the chromatogram. The asterisk and double asterisk indicate slight contaminations with

remaining GST-3C protease and the GST purification tag, respectively.

**Fig 3. Cryo-EM analysis of CRL4-NEDD8$^{DCAF1-CtD}$/Vpr$_{mus}$/SAMHD1 conformational states.**

(**A**) Two views of an overlay of CRL4-NEDD8$^{DCAF1-CtD}$/Vpr$_{mus}$/SAMHD1 cryo-EM reconstructions (conformational state-1 – light green, state-2 – salmon, state-3 – purple). The portions of the densities corresponding to DDB1 BPA/BPC, DCAF1-CtD and Vpr$_{mus}$ have been superimposed. (**B**) Two views

of a superposition of DDB1/DCAF1-CtD/Vpr$_{mus}$ and CUL4/ROC1 (PDB 2hye) [15] molecular models,

which have been fitted as rigid bodies to the corresponding cryo-EM densities; the models are oriented

as in **A**. DDB1/DCAF1-CtD/Vpr$_{mus}$ is shown as in Fig 2A, CUL4 is shown as cartoon, coloured as in **A**

and ROC1 is shown as cyan cartoon. (**C**) Comparison of outermost CUL4 stalk orientations observed

in the cryo-EM analysis presented here (states-1 and -3, coloured as in **B**, show 119.5° rotation of DDB1

BPB) to the two most extreme stalk positions present in previous crystal structures (PDB 4a0l [13], PDB

6dsz [115], coloured grey, show 143.4° DDB1 BPB rotation).

**Fig 4. Mechanism of SAMHD1-CtD recruitment by Vpr$_{mus}$.**

(**A**) Two views of the cryo-EM reconstruction of the CRL4-NEDD8$^{DCAF1-CtD}$/Vpr$_{mus}$/SAMHD1 core. The crystal structure of the DDB1/DCAF1-CtD/Vpr$_{mus}$ complex was fitted as a rigid body into the cryo-EM density and is shown in the same colours as in Fig 2A. The DDB1 BPB model and density was removed for clarity. The red arrows mark additional density on the upper surface of the Vpr$_{mus}$ helix bundle. (**B**) Detailed view of the SAMHD1-CtD electron density. The model is in the same orientation as in **A**, left panel. Selected Vpr$_{mus}$ residues W29 and A66, which are in close contact to the additional density, are shown as red space-fill representation. (**C**) *In vitro* reconstitution of protein complexes containing DDB1/DCAF1-CtD, Vpr$_{mus}$ or the Vpr$_{mus}$ W29A/A66W mutant, and SAMHD1, assessed by analytical GF. SDS-PAGE analyses of corresponding GF fractions are shown below the chromatogram,

with boxes colour-coded with respect to the chromatogram. (**D-E**) *In vitro* reconstitution of protein

complexes containing SAMHD1 and Vpr$_{mus}$ W29A/A66W (**D**) or DDB1/DCAF1-CtD and Vpr$_{mus}$

W29A/A66W (**E**). SDS-PAGE analyses of corresponding GF fractions are shown below the

chromatogram, with boxes colour-coded with respect to the chromatogram. The asterisk and double

asterisk indicate slight contaminations with remaining GST-3C protease and the GST purification tag,

respectively.

**Fig 5. Cross-linking mass spectrometry (CLMS) analysis of CRL4$^{DCAF1-CtD}$/ Vpr$_{mus}$/SAMHD1**

(**A**) Schematic representation of sulfo-SDA cross-links between CRL4$^{DCAF1}$/Vpr$_{mus}$ and SAMHD1, identified by CLMS. Proteins are colour-coded as in Figs 3, 4, and SAMHD1 black/white. SAMHD1-CtD is highlighted in yellow. Crosslinks to the N-terminal SAMHD1 globular SAM and HD domains are coloured light brown, while cross-links to the N-terminal half of SAMHD1-CtD are highlighted in pink and cross-links to the C-terminal end of SAMHD1-CtD are coloured purple. (**B**) Sulfo-SDA cross-links from **A**, in the same colour scheme, mapped on the molecular model of CRL4-NEDD8$^{DCAF1-CtD}$/Vpr$_{mus}$/SAMHD1 (state-2), obtained from cryo-EM analysis (Fig 3). SAMHD1-CtD density from the CRL4-NEDD8$^{DCAF1-CtD}$/Vpr$_{mus}$/SAMHD1 (core) cryo-EM analysis (Fig. 4) is shown as yellow mesh. (**C**) The accessible interaction space of SAMHD1-CtD, calculated by the DisVis server [61], consistent with at least 14 of 26 observed cross-links, is visualised as grey mesh. DCAF1-CtD and Vpr$_{mus}$ are oriented and coloured as in Fig 4A.

**Fig 6. Variability of *neo*-substrate recognition in Vpx/Vpr proteins.**

Comparison of *neo*-substrate recognition modes of $Vpr_{mus}$ (**A**), $Vpx_{sm}$ (**B**), $Vpx_{mnd2}$ (**C**) and $Vpr_{HIV-1}$ (**D**).

DCAF1-CtD is shown as grey cartoon and semi-transparent surface, $Vpr_{mus}$ – green, $Vpx_{sm}$ – orange,

$Vpx_{mnd2}$ – blue and $Vpr_{HIV-1}$ – light brown are shown as cartoon. Models of the recruited ubiquitylation

substrates are shown as strongly filtered, semi-transparent calculated electron density maps with the

following colouring scheme: SAMHD1-CtD bound to $Vpr_{mus}$ – yellow, SAMHD1-CtD (bound to $Vpx_{sm}$,

PDB 4cc9) [50] – mint green, SAMHD1-NtD ($Vpx_{mnd2}$, PDB 5aja) [51] – magenta, UNG2 ($Vpr_{HIV-1}$,

PDB 5jk7) [54] – light violet.

**Fig 7. Schematic illustration of structural plasticity in Vpr_{mus}-modified CRL4^{DCAF1-CtD}, and implications for ubiquitin transfer**.

(**A**) Rotation of the CRL4 stalk increases the space accessible to catalytic elements at the distal tip of the stalk, forming a ubiquitylation zone around the core. (**B**) Modification of CUL4-WHB with NEDD8 leads to increased mobility of these distal stalk elements (CUL4-WHB, ROC1 RING domain) [56], further extending the ubiquitylation zone and activating the formation of a catalytic assembly for ubiquitin transfer (see also **D**) [73]. (**C**) Flexible tethering of SAMHD1 to the core by Vpr_{mus} places the bulk of SAMHD1 in the ubiquitylation zone and optimises surface accessibility. (**D**) Dynamic processes **A-C** together create numerous possibilities for assembly of the catalytic machinery (CUL4-NEDD8 WHB, ROC1, ubiquitin-(ubi-)charged E2) on surface-exposed SAMHD1 lysine side chains. Here, three of these possibilities are exemplified schematically. In this way, ubiquitin coverage on SAMHD1 is maximised.