1
Time-dependent pleiotropic phenotypes are caused by progressive DNA polymerase α deficiency in zebrafish

Alex Y. Lin[1,2], Georgia K. Thomas[1,2], Khai Chung Ang[1,2], Damian B. van Rossum[1,2], Victor A. Canfield[1,2], and Keith

C. Cheng[1,2]*

[1]The Jake Gittlen Laboratories for Cancer Research, Penn State College of Medicine, Hershey, PA, USA;

[2]Division of Experimental Pathology, Department of Pathology, Penn State College of Medicine, Hershey, PA, USA

*kcheng76@gmail.com (KCC)

27 Short title: Pol α deficiency mediates cell-specific atypia and cell death

28 Keywords: Pol α, *pola2*, nuclear atypia, cell death, zebrafish

Abstract

The mechanisms underlying variation in cell- and tissue-specific phenotypes caused by mutations are poorly understood. For example, why some cancers show cellular atypia more than others is unclear. As an illustration of phylogenetic variation in phenotype, mutations in a subunit of DNA polymerase α (Pol α), one of the primary eukaryotic replicative DNA polymerases, result in immediate cell cycle arrest in yeast and *Arabidopsis*; a null mutation in the same subunit results in lethality in 5 to 7 days in zebrafish, in association with striking tissue-specific, time-dependent pleiotropic cellular phenotypes. Here, we describe a zebrafish null mutant in the gene encoding the B subunit of Pol α, *huli hutu (hht)*, in which cell death with nuclear fragmentation is found in the central nervous system where the largest number of cells arise, while cells present in smaller numbers, such as the gastrointestinal epithelial cells, instead show cellular atypia that is associated with hepatic dysgenesis and pancreatic and air bladder agenesis. Wild-type maternal mRNA in genotypically mutant eggs becomes undetectable by 24 hours post-fertilization (hpf). Furthermore, the treatment of wild-type larvae with DNA synthesis inhibitors aphidicolin or hydroxyurea from 24 hpf phenocopies *hht*. Taken together, the data are consistent with a model in which reduced capacity for cell proliferation caused by either genetic or chemical mechanisms of DNA synthesis beginning at about 24 hpf results in cell death in tissues with large cell number, and abnormal nuclear morphology in tissues with fewer cells, including the gastrointestinal tract.

Introduction

Changes in nuclear morphology occur during normal cellular processes such as mitosis, differentiation, and cell migration (Jevtić *et al.* 2014, Calero-Cuenca *et al.* 2018), but histologically-defined cellular atypia, characterized by darker hematoxylin staining and large, irregular nuclear shape, is common in both neoplasia and in settings that involve physical or chemical damage to DNA, such as UV-irradiation and chemotherapy (Blum 1978, Stenbäck 1978, Carr and LiVolsi 1989, Kim *et al.* 2016). Nuclear atypia is also used as a diagnostic signature in many human cancers and pre-cancers (Rosai 2004, Billings and Goldblum 2010, Kumar *et al.* 2010, Lanzkowsky *et al.* 2016, Pizzorno *et al.* 2016). The presence and severity of nuclear atypia generally correlates with higher tumor grade and poor prognosis (Rosai 2004, Kodota *et al.* 2014, Manimaran *et al.* 2014, Yamaguchi *et al.* 2015, Poropatich *et al.* 2016, Zhou 2018). Despite the clinical diagnostic relevance of nuclear atypia, its genetic and mechanistic origins are poorly understood.

The small size of zebrafish larvae allowed us to generate histological arrays for pursuit of a histology-based forward genetic screen to identify genes that can cause nuclear atypia (Mohideen et al. 2003). Our laboratory identified a recessive larval-lethal mutant, *huli hutu* (*hht*), that develops optical opacity of the brain, small eyes, an upward curvature of the body, and dies after 5 to 7 days of development. Histology revealed that virtually all organs and tissues exhibited severe cytological abnormalities (Mohideen *et al.* 2003). Highly proliferative tissues, particularly of the retina and brain, and to a lesser extent, the gastrointestinal tract organs, were affected most dramatically.

Positional cloning shown here revealed a frameshift mutation in *pola2* in *hht*, which encodes the

B subunit of the DNA polymerase α-primase complex (Pol α). Immediate growth arrest

phenotypes associated with *pola2* mutation in two other organisms, *Saccharomyces cerevisiae*,

(Foiani *et al* 1994) and *Arabidopsis thaliana* (Yang *et al.* 2009) begged the question of the reason

underlying the longer survival of zebrafish *pola2* mutants. In *Caenorhabditis elegans*, mutation

of the *pola2* paralog *div-1* substantially increased the duration of interphase and produced embryos

that failed to hatch or to generate intestinal or pharyngeal cells (Encalada *et al.* 2000). We detected

wild-type maternal *pola2* mRNA in homozygous mutant embryos, providing an explanation for

the extended lifespan of *pola2* mutant zebrafish. Consistent with the disappearance of detectable

pola2 mRNA at 24 hpf, two independent chemical means of inhibiting DNA synthesis beginning

at 24 hpf phenocopied cell-specific *hht* phenotypes. Here, we show correlations between

replicative demand and DNA damage that appear to explain the disparity in mutant phenotype

between cell types.

Results

A zebrafish mutant with abnormal nuclear morphology

The gross phenotype of *hht* mutants is evident at 3 days post-fertilization (dpf) under a dissecting microscope and includes reduced head and eye size, a curved body, and a pronounced yolk (Mohideen et al. 2003, **Fig. 1A**). Further characterization by 3-dimensional x-ray histotomography using micro-CT confirmed that these mutants lack the pneumatic duct and swim bladder (Ding *et al*. 2019). This gross phenotype persists until death, which occurs between 5 and 7 dpf (**Fig. 1B**). A detailed examination by histology at 5 dpf showed that while all organs and tissues were distinguishable, virtually all exhibited severe cytological changes (**Fig. 1C**). Highly proliferative tissues, including retina, brain, and gastrointestinal tract organs, were affected most dramatically. Enterocytes in the *hht* intestinal lumen were irregular in polarity, size and shape, cellular and nuclear boundaries were difficult to define, and cell nuclei were often hyperchromatic and contained prominent nucleoli. Other endodermally-derived organs such as pancreas and liver also exhibited severe cytologic atypia, often presenting with enlarged nuclei and high variability in cellular and nuclear size and shape. Dysmorphologies were also detected in cartilage, where the normal aligned arrangement of cartilage cells in neat rows (in wild-type siblings) is replaced by cells with variable shapes, sizes, and polarity, and irregular arrangements. The eyes of *hht* mutants have misshapen lenses and retinas with a substantial reduction in cell number, a loss of stratification of retinal layers, and nuclear fragmentation.

Pola2 mutation is responsible for the hht phenotype

<u>*Positional cloning indicates the causative mutation is in one of five candidate genes*</u>

Microsatellite-based positional cloning and gene knockdown by morpholino oligonucleotides

revealed that the causative gene in *hht* mutants was *pola2*. The identification of zero recombinants

in 1948 meioses with markers *z10868* and *z15236* revealed that the gene responsible for the *hht*

phenotype was less than 0.05 centimorgans (cM) away, which corresponds to about 35.5 kb in

zebrafish (Shimoda *et al*. 1999). These two markers were flanked on either side by microsatellite

markers *z26580* and *z13225*, each of which had one recombinant in 1948 meioses, indicating that

the causative mutation resides between these markers (**Fig. 2A**). Five candidate genes, *cenph*,

dimt1l, *mier3b*, *mrps36*, and *pola2*, are annotated in this region.

<u>*Morpholino antisense knockdown of pola2 results in hht phenotype*</u>

To identify the gene responsible for the *hht* phenotype, one translational blocking and two splice-

junction blocking morpholino oligonucleotides (MOs) were designed per candidate gene and

injected into wild-type zebrafish embryos at the one-cell stage to inhibit translation or RNA

splicing, respectively. Only *pola2* and *dimt1l* morphants exhibited a combination of reduced head

and eye size, a pronounced yolk, and a dorsal curvature that were grossly similar to *hht*

(**Supplemental S1**). Histological examination of cellular phenotype at 3 dpf revealed striking

similarities between *pola2* morphants and *hht*, both demonstrating reduced retinal volume and cell

number, loss of retinal layers, fragmented cell nuclei, and a thickened uneven corneal epithelium

(**Fig. 2B**). The eyes of *dimt1l* morphants, while distinctly abnormal compared with wild-type

zebrafish, retained the overall size, ovoid shape, some retinal layering, and cell types of wild-type

eyes, and lacked the profound deficiencies found in *hht* eyes. Taken together, mapping data and

the near identity in phenotypes of *hht* and *pola2* morphants indicate that the affected gene in *hht*

is *pola2*.

Sequencing of gDNA and cDNA of wild-type and mutant *pola2* revealed a 2-nucleotide AC

insertion at the double AC repeat at the 68^{th} nucleotide from the translation start site (**Fig. 2C**).

The frameshift mutation results in a premature stop codon at the 38^{th} amino acid position (**Fig.**

2D), suggesting that the final protein product is truncated and non-functional. The AC insertion,

located at the 2^{nd} exon of the 19-exon zebrafish *pola2* gene, is therefore most likely the causative

mutation. To verify this insertion, genomic DNA from 20 each of *hht* larvae and wild-type larvae

from wild-type siblings of their parents was pooled to sequence PCR amplicons of the 2^{nd} exon of

pola2; the 2-nucleotide insertion was present in the *hht* pool, but not the wild-type pool.

Partial rescue of the huli hutu mutant phenotype with wild-type pola2 mRNA

One prediction of *pola2* being the affected gene in *hht*, is that injection of wild-type mRNA would

rescue at least some of the several phenotypes of *hht* fish. Wild-type *pola2* mRNA was transcribed

from a cDNA clone from the wild-type Connors background strain and injected into embryos from

a $\frac{hht}{+}$ x $\frac{hh}{+}$ cross at the 1-cell stage with the expectation of a 1:2:1 ratio of $\frac{+}{+}$, $\frac{hht}{+}$, and $\frac{hht}{hh}$

genotypes (**Supplemental S2**). At 3 dpf, 90% of injected $\frac{hht}{hht}$ larvae, identified by genotyping,

were partially rescued, exhibiting an intermediate gross phenotype – a straight body and yolk

resembling that of a wild-type, but with eyes and brain that are smaller than wild-type but larger

than *hht* (**Fig. 3, Table 1**). This intermediate phenotype has not been seen in hundreds of crosses

between *hht* heterozygotes. All larvae with an intermediate phenotype were genotypically

homozygous mutant $(\frac{hh}{hht})$. Retinal cell layering of the rescued fish was reminiscent of those seen

in homozygous wild-type larvae. The *hht* intestinal epithelial phenotype, in which cell nuclei show

loss of polarity and prominent nucleoli, was absent in mRNA-rescued *hht* larvae. Fragmented

nuclei in the brain and eyes of *hht* larvae were also absent in both wild-type and rescued fish. The

incomplete rescue phenotype can be explained by an expected degradation and/or dilution of the

injected wild-type *pola2* mRNA over time, leading to a slower rate of cell division than in wild-

type fish.

Another prediction of *pola2* mutation being responsible for the *hht* phenotype is that knockout of

pola2 on both chromosomes by genome editing would show a *hht* phenotype. Wild-type larvae

were rendered mutant for *pola2* using the CRISPR/Cas9 genome editing system. Homozygous

knockout fish harboring a 16-nucleotide insertion between nucleotides 927-928 (**Supplemental**

S3A) exhibited a gross phenotype that was strikingly similar to *hht* (**Supplemental S3B**). The

frameshift in this CRISPR insertion allele disrupts the protein sequence after amino acid 308.

Under the assumption that human and zebrafish POLA2 proteins share similar structures, these

pola2 knockout zebrafish would produce a mutant protein without a functional phosphodiesterase

(PDE) domain, which has been reported to be important for binding between human POLA2 and

the catalytic A subunit of Pol α, predicting a null phenotype (Klinge *et al*. 2009, Suwa *et al*. 2015).

The gross phenotypes of *hht* and *pola2* knockout larvae were indistinguishable, further confirming that the

affected gene in *hht* is *pola2*.

We have established that *hht* is mutant in the *pola2* gene by positional cloning, morpholino

knockdown to generate a similar histological phenotype, CRISPR knockout, and partial rescue

with *pola2* mRNA. Human POLA2 protein, the B subunit of DNA polymerase alpha, is known to

mediate binding of the A catalytic subunit and the chromosome (Suwa *et al.* 2015). As predicted

by this model, insertional mutant of the A subunit, encoded by *pola1*, resulted in gross and

histological brain and eye phenotypes indistinguishable from *hht* mutants (**Supplemental S4**).

pola2-deficient zebrafish mutants exhibit reduced DNA synthesis

Since *pola2* plays a critical role in DNA replication, we investigated how DNA synthesis was

affected in the *hht* mutants. Wild-type and *hht* larvae between 2 to 5 dpf were assessed for DNA

replication by incorporation of 5-ethynyl-2′-deoxyuridine (EdU), a synthetic thymidine analog.

After 30 min of EdU incubation, wild-type larvae exhibited strongly positive EdU staining at all

ages examined. In contrast, *hht* larvae were EdU-negative from 2 to 5 dpf, indicating a reduction

of DNA synthesis below the limits of detection (**Fig. 4**).

The observed global reduction of DNA synthesis in *hht* mutants suggests a model in which

inhibition of DNA synthesis by any mechanism can potentially reproduce the *hht* phenotype,

which includes largely normal development up to 24 hpf followed by progressive dysmorphology

and cell death. Based on this timing, we hypothesized that chemical inhibition of DNA synthesis

by multiple mechanisms may be able to phenocopy *hht*, if started at about 24 hpf. To test this idea,

wild-type larvae were exposed to hydroxyurea, an inhibitor of deoxyribonucleotide production, or

aphidicolin, an inhibitor of replicative DNA polymerases α, δ, and ε (Zhang *et al.* 2008), beginning

at different times. Wild-type embryos were treated with empirically determined concentrations of

50 μM aphidicolin or 150 mM hydroxyurea to induce the *hht* phenotype. Starting inhibition at 2

hpf resulted in death by 24 hpf (**Fig. 5A**). Inhibiting DNA synthesis beginning at 24 hpf (but not

before) phenocopied *hht* by 72 hpf (**Fig. 5B; Supplemental S5, S6**). Cellular phenotypes in the

eyes, brain, and intestine were confirmed by histology (**Supplemental S7**).

Since inhibition of DNA synthesis predicts cell cycle arrest in S phase, higher proportions of cells

are expected to be in S phase. We measured the proportion of cells in S phase by flow cytometric

measurements of DNA per cell in propidium iodide-labeled cells dissociated from wild-type and

hht larvae (**Fig. 6**). At 3 dpf, the proportion of cells in S phase was approximately 3-fold higher in

hht (45.9 ± 5.8 %) than in wild-type (15.5 ± 3.2 %; $p < 0.05$, t-test). S phase accumulation in *hht*

cells persisted to 5 dpf ($p < 0.05$, t-test). In this analysis, all cell types were examined and the

proportion of cells in each phase of the cell cycle represented an average across heterogeneous cell

populations.

Tissue specificity of cell death and DNA damage phenotypes in* hht *zebrafish

Nuclear fragmentation as seen in the eyes of *hht* larvae indicates cell death. To assess the extent

and timing of cell death, live wild-type and *hht* larvae at 24, 36, 48, 72, 96, and 120 hpf were

incubated with acridine orange (AO), a fluorescent vital dye that labels dying cells. There was no

significant difference between the number of AO-positive foci in wild-type and *hht* larvae at 24

hpf but AO-positive foci in *hht* larvae increased dramatically between 24-36 hpf (**Fig. 7A**). After

peaking at 36 hpf, the number of detectable foci in *hht* continued to decline until 120 hpf. In

contrast, the number of AO-positive foci in wild-type larvae remained consistently low and was

significantly less than in *hht* larvae - at the 36 hpf peak, the number of AO-positive foci in *hht* was

approximately 4-fold higher than that in wild-type. AO-positive cells in *hht* larvae accumulated

predominantly in the most numerous cell types – those in the brain, eyes, and spinal cord (**Fig. 7B-**

C).

To assess the induction of double strand DNA breaks, wild-type and *hht* larvae at 24, 36, 48, 72,

96, and 120 hpf were probed with an antibody specific for γ-H2AX, the phosphorylated form of

H2AX, a member of the H2A histone family. H2AX becomes phosphorylated on serine139 in

response to double-stranded DNA breaks (DSB, Kuo and Yang 2008). At all ages examined, less

than 30 positive foci were detected per wild-type larva. In contrast, the number of positive γ-H2AX

foci in *hht* was indistinguishable from wild-type at 24 hpf, but increased significantly, from 8- to

13-fold, between 36-120 hpf. (**Fig. 8A**). γ-H2AX staining predominated in the brain, eyes, and

spinal cord of *hht,* following the pattern of cell death as assessed by AO (**Fig. 8B-C**).

Wild-type maternal* pola2 *mRNA sustains survival of* pola2*-deficient mutants

To explain the extended survival of *pola2* null mutants of zebrafish compared with the *pola2-*

deficient phenotypes of other model organisms, we hypothesized the presence of wild-type

maternal *pola2* mRNA contributed from the wild-type chromosome of the heterozygous mother's

primary oocytes. Wild-type Pola2 protein originating from translation of wild-type maternal *pola2*

mRNA would then support normal DNA replication during early embryogenesis and could be

expected to be progressively diluted as cells divide in the absence of the wild-type gene in *hht*

mutants. This scenario explains the relatively normal appearance of *hht* fish at 24 hpf, followed

by the increasing disruption of development and eventual death at days 5-7 that characterize *hht*

fish. The similarity in *hht* and chemically-inhibited fish phenotypes only after 24 hpf suggested

are consistent with active DNA replication before 24 hpf in *hht* homozygotes.

241 To determine whether maternal wild-type *pola2* mRNA facilitates the survival of *hht* larvae,

242 cDNA from 3 homozygous wild-type and 3 *hht* embryos at 2.5, 6, 24, 48, and 72 hpf were analyzed

243 by quantitative allele-specific PCR to detect the presence of wild-type maternal *pola2* mRNA (**Fig.**

244 **9**). At 2.5 hpf, wild-type *pola2* transcripts were detected at similar levels in wild-type and *hht*

245 embryos ($n = 3$; $p > 0.5$, t-test). By 6 hpf, the quantity of wild-type *pola2* transcripts in *hht* embryos

246 was significantly reduced compared to wild-type embryos ($n = 3$; $p < 0.05$, t-test). Wild-type *pola2*

247 mRNA was not detected in *hht* larvae at or beyond 24 hpf. Taken together, our results are

248 consistent with a model in which the presence of wild-type *pola2* mRNA and protein is responsible

249 for the sustained survival of the *hht* mutants, and that the degree of depletion of *pola2* mRNA and

250 protein would be dependent upon the number of cell divisions in a given lineage.

Discussion

253 Nuclear atypia, despite its clinical importance in the diagnosis and prognosis of human cancers,

254 has not been clearly linked to the genetic origins of cancer. Since the amount of hematoxylin stain

255 in histology is roughly proportional to DNA content, aneuploidy is associated with darker staining

256 nuclei. Nuclei showing increased hematoxylin staining in nuclei, referred to as hyperchromatic

257 nuclei, are a component of nuclear atypia, and are often associated with increased prominence of

258 nucleoli and irregular nuclear shapes. Cells of the brain and eyes have dense nuclear morphology,

259 making it impossible to assess prominence of nucleoli as a component of nuclear atypia. In

260 contrast, the nuclei of gut cells are far larger. Since larger euploid cells contain the same amount

261 of DNA per cell as cells with small nuclei, their nuclei are normally largely pale, with darker-

262 staining nuclear regions. The atypical nuclei of *hht* gut epithelium are consistent with hyperploidy,

which can be caused by arrest of DNA replication in S phase. Cells trapped in S phase in *hht*

mutant cells that are still alive would, by definition, contain more than 2n DNA content, and

therefore be more darkly stained.

The tissue-dependent cellular phenotypes caused by *pola2* deficiency in *hht* mutants are consistent

with differences in cell proliferation in different cell types. Since homozygous *hht* mutants cannot

generate wild-type *pola2* mRNA, the quantity of wild-type *pola2* mRNA, and presumably wild-

type Pola2 protein, is diluted after each round of cell division (**Fig. 10**). As shown in our RT-PCR

experiment, wild-type *pola2* mRNA is undetectable in *hht* fish by 24 hpf, leaving only the wild-

type Pola2 protein produced before the maternal-to-zygotic transition, to support growth thereafter.

Assuming the quantity of wild-type Pola2 protein in the progenitor cells for every tissue was the

same, and is diluted with each cell division, tissues with a larger number of cells must have

undergone more cell divisions and would thus contain less wild-type protein per cell. The

difference in cell number in different tissues may result from differences in proliferation between

organs.

The *pola2* gene encodes the B subunit of Pol α and does not itself exhibit any known enzymatic

activity. Molecular studies in mouse revealed that the B subunit is tightly associated with the zinc

finger motifs in the carboxy-terminal domain of the catalytic subunit A (Mizuno *et al.* 1999). It

functions as a "molecular tether" that recruits the catalytic polymerase subunit to the origin

recognition complex (ORC) for the initiation of DNA replication (Collins *et al* 1993; Uchiyama

and Wang 2004). Mutagenesis of the *POL12* gene, encoding the B subunit of Pol α in yeast,

showed that mutations in the conserved C-terminus of the protein caused either lethality or

temperature sensitivity while mutations in the non-conserved regions in the N-terminus had no

effect (Foiani *et al*. 1995). The B subunit has been shown to be phosphorylated and

dephosphorylated in a cell cycle-dependent manner and this phosphorylation is dependent on its

association with the catalytic subunit (Foiani *et al* 1995; Ferrari *et al* 1996). The ability of Pol α to

initiate DNA replication is maximized when the B subunit is phosphorylated while the N-terminus

of the catalytic subunit A remains unphosphorylated (Schub *et al*. 2001). Other studies have shown

that the B subunit pays a role in telomere maintenance in yeast (Grossi *et al*. 2004) and

reprogramming the regenerative potential of the tail fin of adult zebrafish (Want *et al.* 2019).

In non-vertebrate models, inactivating *pola2* mutations can result in immediate growth arrest (*S.*

cerevisiae: Collins *et al*. 1993, Foiani *et al.* 1994, Foiani *et al.* 1997; *Arabidopsis*: Yang *et al.*

2009). Despite the high likelihood that the *hht* zebrafish are null mutants, they exhibit an extended

lifespan of 5-7 dpf. As predicted by our replicative deficiency model, chemical inhibition of DNA

replication in wild-type embryos starting at 1 dpf phenocopied *hht* at 3 dpf. Treatment during early

embryogenesis caused death in wild-type larvae by 1 dpf, suggesting that normal DNA replication

is indeed required during early embryogenesis for the survival of *hht* mutants, just as it is required

in yeast and *Arabidopsis*. The detection of wild-type *pola2* mRNA in homozygous mutant embryos

supports our hypothesis that DNA replication during embryonic stages of the mutants requires

wild-type *pola2* mRNA. Since the homozygous mutant embryos have no genomic source for wild-

type *pola2* transcripts, wild-type mRNA must have come from the heterozygous mother (fathers

are also heterozygous but do not contribute mRNA to the oocyte). The maternal-to-zygotic

transition (MZT) designates the time at which the zygotic genome is activated, and maternal

transcripts become destabilized and rapidly degraded. In zebrafish, the transcription of the zygotic

genome begins around 3 hpf, and most of the maternal transcripts are degraded by 6 hpf (Kane

and Kimmel 1993, Schier 2007, Tadros and Lipshitz 2009). The time interval between MZT and

onset of the *hht* mutant phenotype suggests that persistence of the wild-type protein is responsible

for the survival of the mutant. That the pattern of subsequent somatic depletion of wild-type Pola2

protein in *hht* follows the pattern of DNA damage is consistent with our model for explaining the

cell-specific pattern of *hht*'s pleiotropic mutant phenotype.

The extended lifespan of our *pola2* mutant, *hht,* and its development of time- and tissue-specific

cellular abnormalities created an opportunity to explore the potential role of the progressive loss

of DNA replication in the causation of a diagnostically important tissue-specific phenotype

associated with cancer, nuclear atypia. We showed that the progressive pattern of phenotypic

change in *hht* can be explained by the initial presence of wild-type mRNA, followed by its

progressive loss in genotypically homozygous mutant zygotes. In the course of our work,

transcriptomic studies of wild-type, unfertilized zebrafish eggs showed that over 10,000 maternal

transcripts are present that include essential genes such as *pola2* (Rauwerda *et al*. 2016). The

present studies suggest that zebrafish mutants for maternally expressed essential genes may

provide an opportunity to assess, without bias, both the tissue-specific and potentially pleiotropic

effects of the progressive loss of gene function in the context of the whole animal. It is worth

noting that the tissue-specific phenotypes of *hht* were discovered here using histology, but that a

3-dimensional, pan-cellular, cell-resolution imaging method such as x-ray histotomography, will

be better-suited for the quantitative study of volumetric phenotypes such as cell and tissue volume

and shape across cell types and tissues (Ding *et al.* 2019). The incorporation of such approaches

in the systematic study of pleiotropy may help us to understand patterns of cell and tissue-specific

phenotypes that emerge in the course of disease.

Materials and Methods

Fish lines, mating, and embryo collection. Generation of ENU-mutagenized mutant *hht* was previously described (Mohideen *et al*. 2003). The background strain was a wild-type fish line acquired from the Connors fish farm, which differs from the Tu and AB laboratory strains that are in common use. The *hht* line is maintained as heterozygotes due to the larval-lethal nature of the mutation. Mating was carried out by placing male and female heterozygotes in Aquatic habitat tanks with dividers the afternoon prior to egg collection. Collected eggs were disinfected in 10% Ovadine (Syndel) for 1 min at room temperature then washed 3 times in charcoal-filtered water. Larvae were incubated at 28.5° C to maintain consistent speed of development.

Histology. Procedure for zebrafish histology has been previously described (Copper *et al*. 2018) Larvae were dechorionated prior to submersion in 10% Neutral Buffered Formalin (Fisher) at 4° C. Samples were fixed overnight at room temperature with gentle agitation and subsequently embedded in agarose. For long-term storage, fixative was replaced with 70% ethanol. Agarose-embedded larvae are subsequently embedded in paraffin, sectioned, and mounted in glass slides.

Sequencing. mRNA was extracted from 72 hpf wildtype and *hht* larvae using the RNeasy Mini Kit (Qiagen) and reverse transcribed to cDNA. PCR was performed on cDNA with 3 sets of primers:

-23~596:

Forward (POLA2F1: TTGAACATCAGAGGACAATA)

Reverse (POLA2R1: TCTCCTCCGTCCAGCATCTC)

503~1274:

Forward (POLA2F2: AGAGGTGGTTTCCACATTTG)

Reverse (POLA2R2: ACAATACCAACTTGCATGC)

1180~1823:

Forward (POLA2F3: ACCAGGTGACAGAAACATTT)

Reverse (POLA2R3: TAAAGTTCAAACATTGTATG).

PCR was performed on gDNA with 1 set of primer for exon 2 of *pola2*:

Forward (POLA2F1: TTGAACATCAGAGGACAATA) anneals to the 5' end of the 2nd exon of

pola2.

Reverse (POLA2gR: TGACTCCAAACAATGTTGTACTTTGATAGTCATTTG) anneals to the

2nd intron of *pola2*.

PCR products were purified with the QIAquick PCR purification kit (Qiagen) and submitted to

Genewiz for sequencing.

Genotyping. Allele-specific primers were designed based on the SNAP primer method described

by Drenkard *et al*. Forward primers span the AC repeats. Wild-type specific primers detect alleles

with only two repeats and mutant specific primers detect alleles with three repeats.

Wild-type forward primer (POLA2wtF: CACTTTCAACATAGCCTTCAACAATGACAGC);

Mutant forward primer (POLA2mtF: GACACTTTCAACATAGCCTTCAACAATGAGACA);

gDNA reverse primer

(gPOLA2ASR: TGACTCCAAACAATGTTGTACTTTGATAGTCATTTG).

cDNA Reverse primer: (cPOLA2ASR: GCTCCAACATTTTGTCGATGGTGCC)

RNA rescue. Wild-type and mutant *pola2* mRNA was extracted from homozygous wild-type or

homozygous mutant embryos, respectively, from a $\frac{hht}{+}$ x $\frac{hht}{+}$ cross with the RNeasy Mini Kit

(Qiagen) and reverse transcribed into cDNAs using the M-MLV Reverse Transcriptase kit

(ThermoFisher). cDNA from wild-type embryos were subjected to allele-specific PCR to eliminate

heterozygotes. cDNAs were amplified by PCR with the following primers:

Forward primer (POLA2-BglIIF): <u>AGATCT</u>TTGAACATCAGAGGACAATA

Reverse Primer (POLA20MluIR): <u>ACGCGT</u>TAAAGTTCAAACATTGTATG

and cloned into PCR®II-TOPO® vectors for sequencing. After sequences are verified, the inserts

are subcloned into PT3TS(4) vectors. Plasmid DNA was extracted from competent cells with the

QIAprep Spin Miniprep Kit (Qiagen) and linearized with Xmal. mRNA was generated from the

linearized plasmids with the T3 RNA polymerase (NEB). 200 pg of wild-type or mutant *pola2*

mRNA were injected into embryos from a $\frac{hht}{+}$ x $\frac{hht}{+}$ cross at the 1-cell stage. All injected embryos

in the first round of rescue experiments were genotyped to ensure that no phenotypic wild-type

larvae were genotypically *hht*.

CRISPR/Cas9 knockout. Gene-specific primer was designed for *pola2* and annealed to a universal

primer containing the sgRNA scaffold as described in protocol provided by Dr. Wenbiao Chen's

lab at Vanderbilt University. 1 μL of 10 μM gene-specific primer was mixed with 1 μL of 10 μM

sgRNA scaffold primer, 2.5 μL 2X NEB buffer 2 with BSA and annealed at 98° C for 1 min then

cooled to 37° C with a ramp speed of 0.1° C/sec. 0.5 μL of 500 μM dNTP and 0.5 μL of T4 DNA

polymerase were added to the mixture and incubated for 20 min. T4 DNA polymerase was then

inactivated by incubation at 75° C for 20 min. sgRNA was synthesized using the MaxiScrip T7 kit

(ThermoFisher Scientific). 1 μL of Turbo Dnase I was added and incubated at 37° C for 15 min to

remove residual DNA primers. sgRNA was purified using the mirVANA kit (ThermoFisher

Scientific). 600 ng/μL Cas9 protein was mixed with either 100 ng/μL *pola2* sgRNA and injected

into wild-type zebrafish embryo at 1-cell stage.

pola2KO: ATTAATACGACTCACTATAGGGGTCCGCGCTGGATGGGGAAgttttagagctagaaatagc;

sgRNA scaffold: ttttgcaccgactcggtgccactttttcaagtTgataaCggactagccttattttaacttgctatttctagctctaaaac

Cell cycle analysis. 20 wildtype and *hht* larvae were collected at 3 and 5 dpf and dissociated into single cells as previously described. Cells were resuspended in 2 mL 70% EtOH at room temperature for 10 min to fix then stored at 4° C before use. Cells were centrifuged at 500 x g for 5 min to remove supernatant. 1 mL propidium iodide solution from DNA QC Particles Kit (BD) was added to each cell mixture. Cells were sorted with FACSCalibur (BD) and cell cycle was analyzed using ModFit LT V3.3.11.

EdU staining. For labeling of proliferating cells, 2, 3, 4, and 5 dpf wildtype and *hht* larvae were incubated for 30 min at 28.5° C in 400 µM EdU solution (Life Technologies). Larvae were incubated in charcoal-filtered water for 30 min at 28.5° C, then fixed immediately in cold 10% neutral buffered formalin. Detection of EdU incorporation was performed according to manufacturer's instructions. Samples were visualized with Axio Zoom.V16 fluorescence stereo microscope (Zeiss).

Chemical inhibition. Wild-type embryos were submerged in charcoal-filtered water, hydroxyurea (Sigma-Aldrich), or aphidicolin (Sigma-Aldrich) at 0, 2, 6 hpf or 24 hpf. For the 24 hpf treatment, fresh solutions were administered at 48 hpf. Hydroxyurea was dissolved in deionized water to prepare a 500 mM stock and aphidicolin was dissolved in DMSO to prepare a 10mM stock. Stock solutions were diluted to appropriate concentrations for each experimental condition so that equal volume was added.

Acridine orange staining. 24, 36, 48, 72, 96, and 120 hpf wildtype and *hht* larvae were incubated in 2 µg/mL acridine orange for 30 min at 28.5° C, then washed for 5 min in charcoal-filtered water for 5 min at 28.5° C. Larvae were visualized with Axio Zoom.V16 fluorescence stereo microscope (Zeiss).

γ-H2AX staining. 24, 36, 48, 72, 96, and 120 hpf wildtype and *hht* larvae were fixed in cold 10%

neutral buffered formalin. Samples were permeablized in acetone at -20 °C for 7 min then washed

with distilled water for 5 min followed by two washes in PBS for 10 min. Samples were then

blocked in 5% goat serum for 1 hr at room temperature and incubated in 1:100 γ-H2AX primary

antibody (Genetex) overnight at 4 °C. After 3 washes in PBST for 15 min, samples were incubated

in 1:1000 Alexa Fluor 488 goat anti-rabbit secondary antibody (Life Technologies) for 3 hrs at

room temperature. After 3 washes in PBST for 15 min, samples were visualized with Axio

Zoom.V16 fluorescence stereo microscope (Zeiss).

Wildtype *pola2* transcript detection. DNA and RNA were extracted from 2.5, 6, 24, 48, and 72

hpf embryos with AllPrep DNA/RNA Mini Kit (Qiagen). Embryos were genotyped with allele-

specific PCR. RNA of genotyped homozygous wild-type and *hht* embryos were reverse transcribed

with oligo-dT. Quantitative PCR was performed on cDNA with allele-specific primers for wild-

type *pola2* and *actinb1*:

Forward (Actinb1F: CATCCGTAAGGACCTGTATGCCAAC)

Reverse (Actinb1R: AGGTTGGTCGTTCGTTTGAATCTC)

as loading control.

PCR products from three homozygous wild-type and mutant embryos were subjected to

electrophoresis. Relative quantity was calculated by densitometry as a ratio of *pola2*

product:*actinb1* product.

Reference

1. Aanes, H., Winata, C. L., Lin, C. H., Chen, J. P., Srinivasan, K. G., Lee, S. G. P., Lim, A. Y. M., Hajan, H. S., Collas, P., Bourque, G., Gong, Z., Korzh, V., Aleström, P., and S. Mathavan, 2011. Zebrafish mRNA sequencing deciphers novelties in transcriptome dynamics during maternal to zygotic transition. *Genome Res*. Aug; 21(8): 1328–1338.

2. Baranovskiy, A. G., Lada, A. G., Siebler, H. M., Zhang, Y., and Y. I. Pavlov, 2012. DNA polymerase δ and ζ switch by sharing accessory subunits of DNA polymerase δ. *J Biol Chem*. 287(21):17281-7.

3. Blum, H. F., 1978. Ultraviolet radiation and skin cancer in mice and men: accumulation of effect and uncertainty of prediction. *Natl Cancer Inst Monogr.* Dec; 50:11-2

4. Calero-Cuenca, F. J., Janota, C. S., and E. R. Gomes, 2018. Dealing with the nucleus during cell migration. *Curr Opin Cell Biol*. **50**: 35-41.

5. Carr, R. F. and V. A. Livolsi, 1989. Morphologic changes in the thyroid after irradiation for Hodgkin's and non-Hodgkin's lymphoma. *Cancer*. Aug 15; 64(4): 825-9

6. Collins, K. L., A. A. Russo, B. Y. Tseng, and T. J. Kelly, 1993. The role of the 70 kDa subunit of DNA polymerase alpha in DNA replication. *EMBO J*. 12: 4555-4566

7. Cooper, G. M, 2000. *The Cell: A Molecular Approach. 2nd Edition*. Sinauer Associates; Sunderland (MA). ISBN-10: 0-87893-106-6.

8. Date, H., Onodera, O., Tanaka, H., Iwabuchi, K., Uekawa, K., Igarashi, S., Koike, R., Hiroi, T., Yuaasa, T., Awaya, Y., Sakai, T., Takahashi, T., Nagatomo, H., Sekijima, Y., Kawachi, I., Takiyama, Y., Nishizawa, M., Fukuhara, N., Saito, K., Sugano, S., and S. Tsuji, 2001. Early-onset ataxia with ocular motor apraxia and hypoalbuminemia is caused by mutations in a new HIT superfamily gene. *Nat Genet*. 29 (2): 184-188

9. Ding, Y., Vanselow, D. J., Yakovlev, M. A., Katz, S. R., Lin, A. Y., Clark, D. P., Vargas,
P., Xin, X., Copper, J. E., Canfiend. V. A., Ang, K. C., Wang, Y., Xiao, X., De Carlo, F.,
van Rossum, D. B., La Riviere, P., and K. C. Cheng, 2019. Computational 3D histological
phenotyping of whole zebrafish by X-ray histotomography. *eLife*. 8:e44898. doi:
10.7554/eLife.44898

10. Dion, V., 2014**.** Tissue specificity in DNA repair: lessons from trinucleotide repeat
instability. *Trends Genet*. 30(6): 220-9

11. Drenkard, E., Richter, B. G., Rozen, S., Stutius, L. M., Angell, N. A., Mindrinos, M., Cho,
R. J., Oefner, P. J., Davis, R. W., and F. M. Ausubel, 2000. A simple procedure for the
analysis of single nucleotide polymorphisms facilitates map-based cloning in *Arabidopsis*.
Plant Physiol. Dec; 124(4): 1483-92. doi: 10.1104/pp.124.4.1483.

12. Encalada, S. E., P. R. Martin, J. B. Phillips, R. Lyczak, D. R. Hamill *et al*, 2000. DNA
replication defects delay cell division and disrupt cell polarity in *Caenorhabditis elegans*
embryos. *Dev. Biol*. 15: 225-238

13. Foiani, M., F. Marini, D. Gamba, G. Lucchini, and P. Plevani, 1994. The B subunit of the
DNA polymerase alpha-primase complex in *Saccharomyces cerevisiae* executes an
essential function at the initial stage of DNA replication. *Mol. Cell Biol*. 13: 923-933

14. Foiani, M., G. Liberi, G. Lucchini, and P. Plevani, 1995. Cell cycle-dependent
phosphorylation and dephosphorylation of the yeast DNA polymerase alpha-primase B
subunit. *Mol. Cell Biol*. 15: 883-891

15. Ferrari, M., G. Lucchini, P. Plevani, and M. Foiani, 1996. Phosphorylation of the DNA
polymerase alpha-primase B subunit is dependent on its association with the p180
polypeptide. *J. Biol. Chem*. 271: 8661-8666

16. Gillooly, J. F., Hein, A., and R. Damiani, 2015. Nuclear DNA content varies with cell size
across human cell types. *Cold Spring Harb Perspect Biol*. **7** (7): a019091.

17. Grossi, S., Puglisi, A., Dmitriev, P.V., Lopes, M., Shore, D. 2004. Pol12, the B subunit of
DNA polymerase α, functions in both telomere capping and length regulation. *Genes Dev*.
18(9):992-1006

18. He, J., Zhang, G., Almeida, A. D., Cayouette, M., Simons, B. D., and W. A. Harris, 2012.
How variable clones build an invariant retina. *Neuron*. Sep 6;75(5):786-98

19. Hu, M. and S. S. Easter, 1999. Retinal neurogenesis: the formation of the initial central
patch of postmitotic cells. *Dev Biol*. Mar 15;207(2):309-21.

20. Hübscher, U., Maga, G., and Spadari, S. 2002. Eukaryotic DNA polymerases. *Annu. Rev.*
Biochem. 71:133-63

21. Iyama T. and D. M. Wilson 3[rd], 2013. DNA repair mechanisms in dividing and non-
dividing cells. *DNA Repair (Amst)*. Aug; 12(8):620-36. doi: 10.1016/j.dnarep.2013.04.015.
Epub 2013 May 16.

22. Jevtić, P., Edens, L. J., Vuković, L. D., and D. L. Levy, 2014. Sizing and shaping the
nucleus: mechanisms and significance. *Curr Opin Cell Biol*. 28: 16-27.
504
     23. Johnson, R. E., Prakash, L., and S. Prakash, 2012. Pol31 and Pol32 subunits of yeast DNA
polymerase δ are also essential subunits of DNA polymerase ζ. *Proc Natl Acad Sci U S A*.
109(31): 12455–12460.

24. Kaguni, L.S. and Lehman, I.R. 1988. Eukaryotic DNA polymerase-primase: structure,
mechanism, and function. *Biochim Biophys Acta*. 950:87-101

25. Kane, D.A. and C. B. Kimmel, 1993. The zebrafish midblastula transition. *Development*.
119:447-56.

26. Kim, E. K., Yoon, G., and H. S. Kim, 2016. Chemotherapy-induced endometrial pathology:

mimicry of malignancy and viral endometritis. *Am J Transl Res*. 8(5): 2459-2467

27. Klinge, S., Núñez-Ramírez, R., Llorca, O., and L. Pellegrini, 2009. 3D architecture of DNA

Pol α reveals the functional core of multi-subunit replicative polymerases. *EMBO J*. **28**:

1978-1987.

28. Kodota, K., Nitadori, J., Woo, K. M., Sima, C. S., Finley, D. J., Rusch, V. W., Adusumilli,

P. S., and W. D. Travis, 2014. Comprehensive pathological analyses in lung squamous cell

carcinoma: single cell invasion, nuclear diameter and tumor budding are independent

prognostic factors for worse outcomes. *J Thorac Oncol*. **9** (8): 1126-1139

29. Li, Z., Hu, M., Ochocinska, M. J., Joseph, N. M., and S. S. Easter Jr., 2000. Modulation of

cell proliferation in the embryonic retina of zebrafish (*Danio rerio*). *Dev*

Dyn. Nov;219(3):391-401.

30. Lynch, H. T. and A. de la Chapelle, 1999. Genetic susceptibility to non-polyposis

colorectal cancer. *J Med Genet*. 36 (11): 801-818.

31. Manimaran, D., Karthikeyan, T. M., Khan, D. M., and R. T. Raman, 2014. Follicular

variant of papillary thyroid carcinoma: cytological indicators of diagnostic value. *J Clin*

Diagn Res. **8** (3): 46-48

32. Mizuno, T., Yamagishi, K., Miyazawa, H., and F. Hanaoka, 1999 Molecular architecture

of the mouse DNA polymerase α-primase complex. *Mol Cell Biol*. 19(11):7886-96

33. Mohideen, M. P. K., Beckwith, L. G., Tsao-Wu, G. S., Moore, J. L., Wong, A. C. C.,

Chinoy, M. R., and K. C. Cheng, 2003 Histology-based screen for zebrafish mutants with

abnormal cell differentiation. *Developmental Dynamics*. 228:414-423

34. Nouspikel, T. and P. C. Hanawalt, 2000. Terminally differentiated human neurons repair

transcribed genes but display attenuated global DNA repair and modulation of repair gene

expression. *Mol Cell Biol*. Mar; 20(5): 1562-1570. doi: 10.1128/mcb.20.5.1562-1570.2000

35. Poropatich, K., Yang, J. C., Goyal, R., Parini, V., and X. J. Yang, 2016. Nuclear size

measurement for distinguishing urothelial carcinomas from reactive urothelium on tissue

sections. *Diagn Pathol*. 11(1): 57.

36. Rauwerda, H., Wackers, P., Pagano, J. F., de Jong, M., Ensink, W., Dekker, R., Nehrdich,

U., Spaink, H. P., Jonker, M., and T. M. Breit, 2016. Mother-Specific Signature in the

Maternal Transcriptome Composition of Mature, Unfertilized Zebrafish Eggs. *PLoS*

One. Jan 22;11(1):e0147151

37. Rosai, J., 2004. *Rosai and Ackerman's Surgical Pathology*. Maryland Heights: MO; Mosby,

an affiliate of Elsevier Inc.

38. Schier, A.F. 2007. The maternal-zygotic transition: death and birth of RNAs. *Science*.

316:406-7

39. Schub, O., Rohaly, G., Smith, R.W.P., Schneider, A., Dehde, S., Dornreiter, I., Nasheuer,

H.P. 2001. Multiple phosphorylation sites of DNA polymerase α-primase cooperate to

regulate the initiation of DNA replication *in vitro*. *J Biol Chem*. 276(41):38076-83.

40. Shen, J., Gilmore, E. C., Marshall, C. A., Haddadin, M., Reynolds, J. J., Eyaid, W., Bodell,

A., Barry, B., Gleason, D., Allen, K., Ganesh, V. S., Chang, B. S., Grix, A., Hill, R. S.,

Topcu, M., Caldecott, K. W., Barkovich, A. J., and C. A. Walsh, 2000. Mutations in PNKP

cause microcephaly, seizures and defects in DNA repair. *Nat Genet*. 42(3): 245-249

41. Shimoda, N., Knapik, E. W., Ziniti, J., Sim, C., Yamada, E., Kaplan, S., Jackson, D., de
Sauvage, F., Jacob, H., and M. C. Fishman, 1999. Zebrafish genetic map with 2000
microsatellite markers. *Genomics*. Jun 15;58(3): 219-32. doi: 10.1006/geno.1999.5824.

42. Stenbäck, F., 1978. Life history and histopathology of ultraviolet light-induced skin
tumors. *Natl Cancer Inst Monogr.* Dec; 50: 57-70.

43. Suwa, Y., Gu, J., Baranovskiy, A. G., Babayeva, N. D., Pavlov, Y. I., and T. H. Tahirov,
2015. Crystal structure of the human Pol α B subunit in complex with the C-terminal
domain of the catalytic subunit. *J Biol Chem*. 290(23): 14328-14337.

44. Tadros, W. and H. D. Lipshitz, 2009. The maternal-to- zygotic transition: a play in two
acts. *Development*. 136:3033-42

45. Uchiyama, M. and T. S. Wang, 2004 The B-subunit of DNA polymerase alpha-primase
associates with the origin recognition complex for initiation of DNA replication. *Mol. Cell*
Biol. 24: 7419-7434

46. Waga, S., Bauer, G., Stillman, B. 1994. Reconstitution of complete SV40 DNA replication
with purified replication factors. *J Biol Chem*. 269(14):10923-34

47. Wallace, K. N., Akhter, S., Smith, E. M., Lorent, K., and M. Pack, 2005. Intestinal growth
and differentiation in zebrafish. *Mech Dev*. Feb;122(2):157-73.

48. Wan, Y., Almeida, A. D., Rulands, S., Chalour, N., Muresan, L., Wu, Y., Simons, B. D.,
He, J., and W. A. Harris, 2016. The ciliary marginal zone of the zebrafish retina: clonal
and time-lapse analysis of a continuously growing tissue. *Development*. Apr
1;143(7):1099-107

49. Wang, Y. T., Tseng, T. L., Kuo, Y. C., Yu, J. K., Su, Y. H., Poss, K. D., and C. H. Chen,

2019. Genetic reprogramming of positional memory in a regenerating appendage. *Curr*

Biol. Dec 16;29(24):4193-4207.e4. doi: 10.1016/j.cub.2019.10.038. Epub 2019 Nov 27.

50. Yamaguchi, T., Kawahara, A., Hattori, S., Taira, T., Abe, H., Sanada, S., Akiba, J., Nishio,

S., Ushijima, K., Kamura, T., and M. Kage, 2015. Cytological nuclear atypia classification

can predict prognosis in patients with endometrial cancer. *Cytopathology*. 26 (3): 157-66.

51. Yang, H., D. Xiang, S. P. Venglat, Y. Cao, E. Wang *et al*, 2009 *PolA2* is required for

embryo development in *Arabidopsis*. *Botany.* 87: 626-634

52. Zhang, L., Kendrick, C., Jülich, D. and S. A. Holley, 2008. Cell cycle progression is

required for zebrafish somite morphogenesis but not segmentation clock function.

Development. 135 (12): 2065-2070.

53. Kuo, L. J. and L. X. Yang, 2008. Gamma-H2AX – a novel biomarker for DNA double-

strand beaks. *In Vivo*. 22 (3): 305-9.

589   **Acknowledgements**

General: We thank Margaret Hubley, Gail Broda, and Kathryn Early for help in maintaining wild-

type and mutant zebrafish lines and generating embryos; Jean Copper and Lynn Budgeon for help

in development of zebrafish histology. **Funding:** This work was supported by NIH: 5R01

AR052535 (PI: KCC) and the Jake Gittlen Laboratories for Cancer Research. **Author**

contributions: The project was conceived by KCC; research design was by AYL and KCC;

experiments were done by AYL and GKT; analysis was by AYL, GKT, and KCC; AYL and KCC

wrote the paper; KCA, DBVR, and VAC contributed to experimental design and editing.

Competing interest: The authors declare no conflict of interest.

Figures

Figure 1. The pleiotropic *hht* mutant survives up to 7 days. A) The *hht* mutant gross phenotype includes reduced head and eye size, an enlarged yolk, and dorsal curvature of the tail. **B)** The *hht* mutants show increasing larval lethality over 7 days; none survive past 7 days. In the same time period, only 1 wild-type sibling died. **C)** 5 dpf *hht* mutants exhibit disruption in tissue organization and a range of cellular dysmorphologies across most cell types.

Figure 2. Microsatellite-based positional cloning of *pola2*. A) In positional cloning, the frequency of co-segregation of a phenotype of interest and polymorphic markers is used to locate the causative mutation. The map distance between the causative mutation and the markers is calculated from the frequency of recombinants as indicated above each marker. For markers *z26580* and *z13225*, 1 recombinant out of 1948 meiosis corresponds to ~ 0.05 centimorgans (cM), or 32.5 kb, in zebrafish. The lack of recombinants for markers *z10868* and *z15236* indicates that the causative mutation is physically closer to these two markers and resides in the interval between *z26580* and *z13225*. Five candidate genes, *mier3b* (mesoderm induction early response factor 1, family 3 b), *mrps36* (28S mitochondrial ribosomal protein S36), *cenph* (centromere protein H), *pola2* (DNA polymerase alpha, subunit B), and *dimtl1* (dimethyladenosine transferase), were annotated in the zebrafish genomic database in the identified GRCz11 region. **B)** Gross and histological examination of the eyes of wild-type, *hht*, *pola2* morphant, and *dimt1l* morphant revealed a striking similarity in reduced cell number, loss of stratification, thickened corneal epithelium, and disruption in the pattern of retinal pigmented epithelial pigment between *hht* and *pola2* morphant, suggesting that *pola2* was the affected gene in *hht*. **C)** a 2-nucleotitde AC insertion in the double AC repeat starting at the 68^{th} nucleotide was detected in *hht*. **D)** This frameshift mutation results in changes in amino acid sequence starting at the 25^{th} amino acid, ending with a premature stop codon at the 38^{th} amino acid position. (red box = AC insertion; yellow highlight = amino acid change; asterisk = stop codon).

602
603

Figure 3. Wild-type mRNA partially rescues the *hht* phenotype. Rescued *hht* mutants exhibit a straight body and a normal yolk. Sizes of the eyes and head are intermediate between wild-type and *hht* mutants. Rescued *hht* fish show histologically normal intestinal epithelium and partial preservation of the retinal layering found in wild-type eyes. The eyes and brain of partially rescued *hht* fish injected with wt mRNA show no evidence of nuclear fragmentation.

605
606

607
608

Figure 4. DNA synthesis is reduced in *hht* larvae. Representative images of wild-type and *hht* larvae stained with EdU. EdU-positive staining was observed in wild-type larvae, but absent in *hht* larvae at 48, 72, 96, and 120 hpf. 3 larvae were examined for each condition. White arrow = a positive focus.

609
610

Figure 5. Chemical inhibition of DNA synthesis in *wt* larvae after 24 hours phenocopies *hht*. A) Treatment wild-type embryos with 100 mM hydroxyurea or 50 μM aphidicolin from 2 hpf on caused death by 24 hpf. **B)** Treatment of wild-type larvae (the time maternal wt *pola2* mRNA becomes undetectable) with 100 mM hydroxyurea or 50 μM starting at 24 hpf yields *hht* phenocopies at 72 hpf.

Figure 6. Accumulation of cells in S phase in *hht*. Wild-type and *hht* larvae were dissociated into single cells at 3 and 5 dpf, stained with propidium iodide, and analyzed by flow cytometry. Compared to wild-type, the proportion of cells in S phase increased significantly in *hht* at 3 dpf ($p < 0.05$) and persisted to 5 dpf ($p < 0.05$). Error bars = standard error of the mean.

Figure 7. Acridine orange staining reveals increased cell death in the brain, eyes, and spinal cord of *hht* larvae. Cell death was assessed by counting the number of acridine orange-positive foci in 3 wild-type and 3 *hht* larvae. **A)** Cell death in *hht* was higher than in wild-type, peaking at 36 hpf. **B)** Cell death localized to the eyes and brain of 48 hpf *hht*. **C)** Cell death in the spinal cord in 48 hpf *hht*. Error bars = standard error of the mean. White arrow: brain, white asterisk: eye.

Figure 8. DNA damage is increased in *hht*. Double-stranded breaks were assessed by fluorescent antibody staining of γ-H2AX. **A)** DNA damage was minimal in wild-type, but striking in *hht* fish from 36-120 hpf. In 48 hpf *hht* fish, DNA damage was localized to the brain, eyes, and **B)** spinal cord **C**). Error bars = standard error of the mean. White arrow: brain, white asterisk: eye.

Figure 9. Wild-type *pola2* mRNA is detected in *hht* embryos. A) Wild-type *pola2* transcripts were detected in *hht* by allele-specific primers at 2.5 and 6 hpf. Wild-type transcripts were not detectable in *hht* after 24 hpf. **B)** Wild-type *pola2* transcripts were normalized to *actin1b*. Wild-type *pola2* transcripts in *hht* embryos were present at comparable levels to wild-type embryos at 2.5 hpf but were significantly reduced compared to wild-type embryos by 6 hpf (p < 0.05)

Figure 10. Differential dilution of *pola2* accounts for tissue-specific differences in cell death. Homozygous *hht* mutant embryos cannot generate wild-type *pola2* mRNA. The quantity of wild-type *pola2* mRNA and protein, is diluted after each round of cell division. In this model, highly proliferative cell types undergo more cell divisions per unit time, resulting in lower concentrations of wild-type *pola2* mRNA or protein per cell compared to cell types that proliferate slowly. Nuclear atypia is apparent in cells with large nuclei but inapparent in cells with small, condensed nuclei.

622 **Tables**

623 *Table 1: wild-type pola2 mRNA rescues hht mutants*

Injected mRNA	Total	$\frac{+}{+}$, $\frac{hht}{+}$	$\frac{hht}{hht}$	
		Phenotypically wild-type / (expected)	Partially rescued / total $\frac{hht}{hht}$	Phenotypically *hht*
None	182	136 / (137)	0 / 46	46
wt *pola2* RNA	302	234 / (227)	61 / 68	7
mutant *pola2* RNA	75	60 / (56)	0 / 15	15

A mixture of wild-type and mutant embryos from a $\frac{hht}{+}$ x $\frac{hht}{+}$ cross were injected at the 1-cell stage with phenol red, 200 pg wild-type *pola2* mRNA, or 200 pg mutant *pola2* mRNA. Number of embryos injected with no mRNA or wild-type *pola2* mRNA was aggregated from three separate experiments.

Supplemental S1. Morphants of the five candidate genes in the region identified by positional cloning, *mier3b*, *mrps36*, *cenph*, *pola2*, and *dimtl1*. Only *pola2* and *dimt1l* morphants exhibit a combination of small eyes, enlarged yolk, and dorsal curvature characteristic of *hht* mutants.

Supplemental S2. Wild-type *pola2* mRNA rescue schematic. Wild-type *pola2* mRNA was injected into 1-cell stage embryos from a $\frac{hht}{+}$ X $\frac{hht}{+}$ cross. Larvae were genotyped to confirm that rescued larvae were homozygous for the *hht* mutation.

```
wt  GTTGTTGTCATGGAAGGTATGAATCCCTC----------------GGGGGAGAAGCTTGTTGC
    |||||||||||||||||||||||||||||#|#              ||||||||||||||||||
hht GTTGTTGTCATGGAAGGTATGAATCCATGACAACAGGGGGGAGAAGGGGGAGAAGCTTGTTGC
```

B

Supplemental S3. Shared phenotype of CRISPR/Cas9 *pola2* knockouts and *hht*

A) The CRISPR/Cas9 knockout zebrafish carried a 16-nucleotide insertion between nucleotides 927-928, causing a frameshift that changes the protein sequence after amino acid 308 in the 600-amino acid zebrafish Pola2 protein. Differences in severity in phenotype are consistent with variation in genetic background. **B**) Representative images of wild-type, *hht*, and *pola2* CRISPR/Cas9 knockout zebrafish larvae at 5 dpf. *hht* and *pola2* knockout fish shared small eyes, small head, enlarged and rounded yolk, and dorsally curved body.

Supplemental S4. Mutants of *pola1* and *pola2* exhibit virtually indistinguishable gross and histological phenotypes. At 3 dpf, *hht*, our *pola2* null mutant, and a *pola2* insertional mutant both exhibit the combinatorial gross phenotype of small eyes, small head, rounded yolk, and curved body. Histologically, both mutants show a drastic reduction of cell number and loss of retinal layers in the eyes, reduced volume and disorganization of the white and gray matter of the brain. Scale bar = 100 μm.

Aphidicolin

Supplemental S5. Inhibition of replicative DNA polymerases by aphidicolin beginning in wild-type zebrafish larvae at 24 hours, but not earlier, phenocopies *hht* at 72 hpf. Wild-type larvae were continuously exposed to 0, 20, 50, or 100 µM aphidicolin in DMSO. Treatment began at 0, 2, 6, or 24 hpf. 10 larvae were treated for each condition. Representative images are shown. Black line = no treatment; red line = treatment.

Hydroxyurea

Supplemental S6. Inhibition of deoxyribonucleotide production by hydroxyurea in wild-type zebrafish larvae beginning at 24 hours, but not earlier, phenocopies *hht* at 72 hpf. Wild-type larvae were continuously exposed to 0, 40, 60, or 150 mM aqueous hydroxyurea. Chemical treatment began at 0, 2, 6, or 24 hpf. 10 larvae were treated for each condition. Representative images are shown. Black line = no treatment; red line = treatment.

Supplemental S7. Chemical inhibition of DNA synthesis starting at 24 hpf phenocopies *hht* **cellular phenotypes.** Continuous exposure of wild-type larvae to 50 μm aphidicolin or 150 mM hydroxyurea starting at 24 hpf results in nuclear fragmentation in the prefrontal cortex and eyes. Eyes also suffer a drastic reduction in cell number and organization of retinal layers. Cellular and nuclear atypia are observed in the intestine – cell and nuclear sizes are irregular, cell boundaries are obscure, and prominent nucleoli are present.

644