1 **Cyclic AMP signalling and glucose metabolism mediate pH taxis by African trypanosomes**
2

Sebastian Shaw[1,2,3], Sebastian Knüsel[1], Daniel Abbühl[1,4], Arunasalam Naguleswaran[1] and Isabel
Roditi[1,*]

[1] Institute of Cell Biology, University of Bern, Bern, Switzerland

[2] Graduate School of Cellular and Biomedical Sciences, University of Bern, Bern, Switzerland

[3] Present address: Department of Pathobiology, School of Veterinary Medicine, University of
Pennsylvania, 380 South University Avenue, Philadelphia, PA 19104, USA

[4] Present address: Trypanosome Cell Biology Unit, INSERM U1201, Institute Pasteur, 25 Rue du
Docteur Roux, 75015 Paris, France
11
* Corresponding author: isabel.roditi@izb.unibe.ch

Abstract

More than a decade ago it was proposed that the collective migration of African trypanosomes
on semi-solid surfaces could be explained by a combination of migration factors and repellents
released by the parasites, but the identity of these molecules was unknown. Here we show that
procyclic (insect midgut) forms acidify their environment as a consequence of glucose
metabolism, generating pH gradients by diffusion. Early and late procyclic forms exhibit self-
organising properties on surfaces. Both forms are attracted to alkali, but while early procyclic
forms are repelled by acid and migrate outwards, late procyclic forms remain at the inoculation
site. pH taxis relies on cyclic AMP signalling. Acid sensing requires a flagellar adenylate cyclase,
ACP5, and a cyclic AMP response protein, CARP3, that interacts with ACP5. Deletion of the
flagellar phosphodiesterase PDEB1 abolishes pH taxis completely. Trypanosomes can also
respond to exogenously formed gradients. pH sensing is likely to be biologically relevant as
trypanosomes experience large differences in pH as they progress through their tsetse fly host.
In addition, self-generated gradients may help reinforce directionality. Moreover, since
trypanosomes encode a large family of adenylate cyclases, these may govern other chemotactic
responses and tissue tropisms in both the mammal and the fly.

Introduction

Environmental sensing, chemotaxis and quorum sensing present possibilities for pathogens to orient themselves within their hosts. All three are widespread in bacteria, and the sensors and response mechanisms have been characterised in detail [1,2]. Genes involved in bacterial chemotaxis and quorum sensing have long been linked to biofilm formation and swarming behaviour[3,4]. By contrast, very little is known about chemotaxis in unicellular eukaryotes, with the exception of *Dictyostelium discoideum*, which generates and reacts to cyclic AMP (cAMP) gradients, promoting aggregation and differentiation[5–8].

Intracellular pathogens frequently exploit receptor-ligand interactions to recognise and invade the appropriate cell types [9–12]. For extracellular parasites such as *Trypanosoma brucei spp.*, causative agents of human and animal trypanosomiasis, it is not known what drives them to move from one host tissue to another. In their mammalian hosts, quorum sensing regulates the titre of *T. brucei* by triggering the differentiation of proliferative slender forms to quiescent stumpy forms[13,14]. When these are taken up by tsetse flies during a blood meal[15] they differentiate to early procyclic (midgut) forms and then to late procyclic forms[16]. In addition to the surface marker GPEET procyclin, which is only expressed by early procyclic forms, several metabolic enzymes, nutrient transporters and signal transducers are differentially expressed between the two forms[17,18]. Colonisation of the midgut by procyclic forms is followed by additional rounds of differentiation as parasites invade the proventriculus, then move to the salivary glands before being transmitted to a new mammalian host.

Early procyclic forms engage in a form of group migration, known as social motility (SoMo), when cultured on semi-solid surfaces such as agarose plates[18,19]. The parasites initially remain at the inoculation site, dividing approximately once every 24 h, then the communities start to form regularly spaced projections that extend by ~1 cm (500 body lengths) per day. Communities on the same plate are able to sense each other and reorient to avoid contact (Supplementary Video 1). Late procyclic forms do not exhibit SoMo, but they do produce a repellent that is sensed by early procyclic forms[18]. Communities on plates also react to other signals. Trypanosomes depleted of the spliced leader core protein SmD1 shed exosomes that act as repellents[20], while bacterial colonies can act as attractants[21]. Various species of *Leishmania*,

parasites related to trypanosomes, respond chemotactically to a range of compounds including

amino acids, carbohydrates and sera, in liquid culture[22–24]. In none of these cases is it known

which genes mediate the responses.

Few genes have been implicated in SoMo. Concomitant depletion of adenylate cyclases

ACP1&2, or depletion of ACP6 on its own, resulted in a hyper-social phenotype (hyper-SoMo)

where the communities formed more projections and migrated more rapidly, suggesting that

these are negative regulators[25]. The flagellar phosphodiesterase B1 (PDEB1)[19] and the

endoplasmic reticulum protein Rft1[26,27] are positive regulators of SoMo. Importantly, null

mutants of PDEB1 and Rft1 were not only impaired in SoMo, but also defective in colonising the

tsetse midgut[26,28]. We postulated that SoMo is largely a chemotactic response to factors

produced by the trypanosomes themselves. Here we show that procyclic forms can generate pH

gradients through glucose metabolism and that the response of early and late procyclic forms to

these gradients can explain the self-organising properties of communities on plates. In addition,

we identified two cAMP signalling components involved in the response to acid, ACP5 and the

cAMP response protein CARP3. Notably, pH sensing is completely abolished in PDEB1-KO, where

cAMP levels are perturbed. pH taxis could be biologically relevant as there is a pH gradient in the

tissues that are colonised by the parasites during their transmission through the insect host [29].

Results

Parasite-secreted factors change properties of the medium and repel migrating projections

What are the factors causing collective migration in SoMo? The fact that communities sense and

avoid each other over considerable distances means that they react to attractants or repellents

that are secreted, or otherwise released, by the parasites. We started by investigating if these

factors were produced under normal growth conditions in liquid culture. We concentrated the

supernatant of a dense culture (2×10^7 cells ml^{-1}) 10-fold and spotted this "conditioned medium"

(CM) next to migrating projections on SoMo plates (Fig. 1a, upper panel). Projections migrating

towards the site where CM was spotted reoriented away from it, indicating that it was perceived

as a repellent. In a second experiment, CM was either sterile filtered or dialysed (with a 12 kDa

cut-off) prior to concentration and spotted next to migrating projections. As seen with untreated

CM, filtered CM still repelled migrating projections (Fig. 1a, lower left panel), but dialysed CM

lost its repellent properties (Fig. 1a, lower right panel). On the contrary, migrating parasites were

attracted to dialysed CM in the same way that they were to concentrated fresh medium (DTM)

(Fig. 1a). These results suggested that repellents might be metabolites that were lost upon

dialysis.

Trypanosomes acidify their environment in liquid culture and on agarose plates

Trypanosomes secrete a variety of molecules that influence the pH of their environment[30,31].

Indicators for pH, such as phenol red, are commonly used for culture media and it is apparent

that dense cultures are more acidic than fresh medium or log phase cultures, as illustrated for

early procyclic forms growing in liquid culture (Fig. 1b). We therefore used a small tip micro glass

pH electrode to measure the pH in liquid culture and on agarose plates (Fig. 2). Fresh medium

has a pH of 7.46 ($\pm$ 0.02) and the supernatant of a log phase culture has a pH of 7.45 ($\pm$ 0.01) (Fig.

2a). The pH decreased to 6.95 ($\pm$ 0.00) when parasites were allowed to overgrow; this

corresponds to ~3-fold increase in the concentration of H^+ ions (Fig. 2a).

To measure the pH on plates, 2 x 10^5 parasites were inoculated either in the centre of a

plate (Fig. 2b, left row), or in pairs at a defined distance from each other (Fig. 2b, right row), and

the pH was measured after 2, 3 ,4 and 7 days post inoculation. We measured the pH in the centre

of a community, at the periphery of a plate and between two communities. The pH at the centres

of the communities decreased from 7.50 to 7.31 ($\pm$ 0.05) during the first three days and

decreased further to 6.96 ($\pm$ 0.05) between days 3 and 7 (Fig. 2b, c). The pH at the periphery also

started to decrease between day 3 and 7, possibly because of diffusion of H^+ ions from the

parasite communities. When communities were plated in pairs, the pH at the midpoint between

them also decreased over time (Fig. 2b, right row, arrow). Thus, communities are able to acidify

their vicinity resulting in pH gradients on the plates.

We showed previously that trypanosomes needed to reach a threshold number before

they started migrating[18]. We now postulate that it is not the cell number that is critical, but rather

the need to establish a discernible pH gradient. When the same number of parasites were

inoculated onto plates containing different volumes of semi-solid medium, the time point at

which projections started to form was inversely proportional to the volume (Fig. 2d). This would

be consistent with larger volumes buffering the pH more effectively and requiring more cells, and

thus more time, before a gradient was established.

Reaction of early and late procyclic forms to exposure to acid and alkali

As a next step, we tested if manipulating the pH gradient had an influence on migration by

exposing early procyclic forms to external sources of acid or alkali (Fig. 3a, b). Early procyclic

forms were inoculated onto plates and incubated until the projections reached a length of about

2 cm. At this point, solutions of HCl or NaOH were pipetted onto the surface at a defined distance

(Fig. 3a) and the plates were incubated for 16-20 h. The migrating cells responded very strongly

to both solutions (Fig. 3b, Supplementary Videos 2 & 3). Exogenous NaOH acted as an attractant,

causing projections to reorient towards it (OH^- treatment), while HCl caused the projections to

stop moving outwards, or to move away from the acid (H^+ treatment). Using the same assay,

another 22 chemicals were tested (Supplementary Fig. 1). In most cases, when a solution was

acidic, the projections were repelled, and when a solution was basic, the cells were attracted to

it. These experiments also showed that the projections reacted to the pH rather than to specific

ions in the solutions.

We next tested late procyclic forms since these do not perform SoMo. Late procyclic

forms did not respond to HCl (Fig. 3c, Supplementary Video 4), but they formed a limited number

of projections that migrated towards NaOH. Community lifts (Fig. 3c) confirmed that the

parasites were still GPEET-negative and had not dedifferentiated to early forms. In summary,

these results demonstrate that procyclic forms exhibit pH taxis, with alkali acting as an attractant

for both early and late procyclic forms. In contrast, there are stage-specific differences in their

reaction to acid.

RNA-seq analyses suggest that SoMo is a response to self-generated gradients

To better understand the mechanisms behind self-organisation of early and late procyclic forms,

and their pH reponses, we performed RNA-seq under different conditions. A culture of procyclic

forms (>60 % GPEET-positive) was plated and allowed to form projections (Fig. 4a, far right). A

community lift with antibodies against procyclins showed that the tips were strongly positive for

GPEET, while the centre of the community stained more strongly for EP. Cells were isolated from

the tips and roots of projections (Fig. 4a far right) and analysed by RNA-Seq. Data confirmed that

markers of early procyclic forms were more highly expressed in the tips than the roots (Fig. 4a

left; Supplementary Table 1a). Other transcripts that were more highly expressed in the tip were

not characteristic of early procyclic forms in liquid culture, such as cysteine-rich acidic integral

membrane protein (CRAM)[32], a microtubule-associated repetitive protein (MARP2)[33] and a

hypothetical protein (hypo; Tb927.9.10400). In addition, several adenylate cyclases were

differentially expressed on plates; ACP3 and ACP5 were ≥2-fold higher in the tip, while ACP6 and

another adenylate cyclase (GRESAG 4; Tb927.11.1480) were ≥2-fold higher in the root. The up-

regulation of high affinity glucose transporters (THT2 and 2A), glycolytic enzymes and

components of the glycosome (Fig. 4a middle and Supplementary Table 1a) is suggestive of

higher glucose metabolism by the tip than the root. Increased expression of these transcripts is

also characteristic of early procyclic forms in liquid culture[17,18].

RNA-seq was also performed on the tips of communities exposed to acid or alkali (Fig. 4b

and Supplementary Table 1b). There were no significant differences (≥2-fold, p=0.05) between

non-exposed communities and those exposed to alkali. A single cation transporter (CatT,

Tb927.11.9000) was more highly expressed in cells exposed to acid, but the read counts were

extremely low in both cases. CatT and a copy of a contig (Tb11.v5.0514) were the only significant

differences in a comparison of acid- versus alkali-treated communities. Taken together, these

results suggest that self-generated pH gradients might be the normal driver of migration on SoMo

plates.

Glucose contributes to the formation of pH gradients

Since the RNA-seq data indicated that cells at the tip might metabolise glucose more actively

than cells at the root, we investigated the role of glucose in SoMo. For these experiments SDM80

medium was prepared with 1 mM glucose, 6 mM glucose or 6mM N-acetylglucosamine (GlcNAc),

which inhibits glucose uptake by binding to hexose transporters[34]. Early procyclic forms of Lister

427 Bern were inoculated onto plates and incubated for 6 days (Fig. 5a). The parasites in SDM80

with 1 mM glucose started migrating later than cells in SDM80 containing 6 mM glucose, and the

SoMo defect was even more pronounced when glucose uptake was blocked by GlcNAc. Under

these conditions the cells only formed rudimentary "knobs" at the edge of the community and

failed to form projections, even at later time points.

The reduced ability to migrate on plates can have a variety of causes, especially in the

context of glucose restriction. It might be due to longer population doubling times, motility

defects or inefficient energy metabolism. Procyclic cultures of Lister 427 Bern are stably GPEET-

positive[26,35]. Nevertheless, we assessed the proportion of early procyclic forms by flow cytometry

(Fig. 5b). Regardless of the medium in which the cells were grown, all cultures were ≥97% GPEET-

positive. Furthermore, the glucose concentration had no impact on the population doubling time

(Fig. 5c) or intracellular ATP levels (Fig. 5d), both assessed in liquid culture. Previous work showed

that directional motility is important for SoMo and migration through the fly[28]. We therefore

assessed cell motility (Fig. 5e); no differences were observed between cells grown in the presence

or absence of glucose.

To investigate whether different glucose concentrations influence the culture pH, we

compared medium alone with liquid cultures of trypanosomes in log phase or high density (Fig.

5f). While the pH of log-phase cultures did not differ from that of medium alone, dense cultures

of parasites in SDM79 or SDM80 with 6 mM glucose strongly acidified the medium. The

concentration of H^+ in dense cultures, compared to log phase cultures, increased 2.53-fold

(±0.03) for SDM79 and 2.51-fold (±0.06) for SDM80 plus glucose. By contrast, cells grown in

SDM80 plus GlcNAc showed a smaller change in H^+ ion concentration (1.22-fold ±0.03). We next

measured pH changes for communities growing on plates (Fig. 5g). The ΔpH between the centre

and the periphery reached -0.31($\pm$ 0.02) for parasites on SDM79 and -0.37($\pm$ 0.02) for SDM80 +

6mM glucose, corresponding to a ≥2-fold difference in H^+ ions at the two locations, whereas

plates containing SDM80 + GlcNAc did not show a difference in pH. Taken together, these data

show that glucose is required for parasites to generate a pH gradient, and that this correlates

with their ability to perform SoMo.

Cyclic AMP signalling is involved in the response to pH

198   RNA-seq identified a number of differentially expressed transcripts, including that of the flagellar

199   pocket protein CRAM, which was more highly expressed in the tips than in the roots (Fig. 4a).

200   CRAM is not essential in procyclic culture forms[36]. When we generated knockouts, these grew at

the same rate as their parental line, but two out of three independent clones produced smaller projections. Nevertheless, all three reacted to acid and alkali in the same way as wild-type cells (Supplementary Fig. 2) so we did not proceed with additional analyses.

Several adenylate cyclases were also differentially expressed between the tip and the root (Fig. 4a). Before we started analysing these and other members of the cAMP signalling pathway, we tested whether PDEB1, which hydrolyses cAMP, had an influence on the pH response. When the PDEB1 null mutant was exposed to alkali or acid, it showed no pH response (Fig. 6a). Notably, the PDEB1 null mutant differs from both early procyclic forms, which react to acid and alkali, and late procyclic forms, which form projections in response to alkali but do not respond to acid.

Four cAMP response proteins (CARP1-CARP4) were previously identified in an RNAi screen as mediating resistance to an inhibitor of PDEB1[37], but they have not been studied in the context of SoMo. For CARP 1 and 2, we only obtained clones after single rounds of transfection. We did not pursue these further, however, as the resistance genes were not integrated at the correct loci. However, we were able to obtain knockouts of CARP3 and 4 (Supplementary Fig. 3a). CARP4 null mutants did not show any difference to the parental line when they were subjected to a standard SoMo assay or tested for pH responses (data not shown). Deletion of CARP3, while not affecting growth in liquid culture, motility or expression of the early marker GPEET (Supplementary Fig. 3b-3e) resulted in a major SoMo defect, reminiscent of the PDEB1 null mutant (Fig 6b, upper panel). CARP3 knockouts still reacted to alkali, but did not react to acid (Fig 6b, lower panel). An HA-tagged addback of CARP3, expressed from a procyclin locus, partially rescued the SoMo phenotype (Fig 6b, upper panel), although the projections were shorter than in the parental line. Importantly, the addback was now able to sense and react to both acid and alkali (Fig 6b, lower panel).

CARP3 is distributed between the cytoplasm and the flagellum (Supplementary Fig. 3f; www.tryptag.org). Since no interaction partners have been reported, a pulldown was performed with HA-tagged CARP3 (Fig. 6e) and analysed by mass spectrometry. This led to the identification of adenylate cyclases ACP3 and ACP5 in three biological replicates (Supplementary Table 2). These interactions were confirmed by *in situ* tagging and co-immunoprecipitation of myc-tagged versions of ACP3 and ACP5 in the CARP3-HA addback line (Fig. 6e). The reciprocal co-

immunoprecipitations with anti-myc antibody confirmed the interaction for both ACP3 and ACP5.

These results, combined with their differential expression in the tips and roots of projections (Fig.

4a) stimulated us to revisit some of the flagellar ACP previously characterised by Lopez and

coworkers[25]. In their study they showed that knockdown of ACP1/2 or ACP6 by RNAi resulted in

a hyper-SoMo phenotype. We obtained null mutants of ACP2 and ACP6, but it was only possible

to generate single knockouts of ACP1 and ACP5 (Supplementary Fig. 4a) implying that they are

essential at this stage of the life cycle. Although we attempted to knock out ACP3, and isolated

resistant cell lines, the antibiotic resistance genes were not integrated at the correct locus (data

not shown). The full knockout of ACP6 and the semi-knockout of ACP1 both exhibited hyper-

SoMo phenotypes, while the full knockout of ACP2 showed no phenotype (Supplementary Fig.

4b). These results are consistent with the findings with the RNAi lines; they also indicate that the

hyper-SoMo phenotype in the previously published ACP1/2 RNAi line[25] was due to depletion of

ACP1. Unexpectedly, however, and in contrast to the results with RNAi[25], single knockouts of

ACP5 were unable to form projections on plates (Fig. 6c, upper panel). In terms of their pH

response, ACP5 single knockouts resembled CARP3 double knockouts - they were able to form

projections migrating towards an alkaline source, but were refractory to acid (Fig. 6c, lower

panel). Further experiments showed that >80% of these cells were GPEET-positive and that the

phenotype of the ACP5 mutants was not due to defects in growth or motility (Supplementary Fig.

4c-4f). These experiments also revealed that ACP5-myc was localised exclusively at the flagellar

tip (Fig 6d). In summary, we have extended the catalogue of cAMP signalling components that

are involved in regulating SoMo. In addition, the finding that PDEB1, ACP5 and CARP3 play a role

in the perception of environmental acidity/alkalinity is consistent with SoMo being a response to

self-generated pH gradients.

Discussion

It was proposed more than a decade ago that the collective migration of trypanosomes on plates

could be explained by a combination of migration factors and repellents released by the

parasites[19], but the identity of these molecules was unknown. Here we show that trypanosomes

acidify their environment as a consequence of glucose metabolism, generating pH gradients by

diffusion. While it is commonly assumed that free glucose levels in tsetse are low between blood

meals, it is difficult to identify a primary publication on this topic. It was shown, however, that

the fly gut contains trehalose, a disaccharide composed of two glucose subunits[38]. Moreover, *T.*

b. brucei encodes a putative α-trehalose phosphorylase (Tb927.11.10670) which could generate

glucose and glucose-1-phosphate. Early and late procyclic forms exhibit self-organising

properties. It is the perception of these self-generated pH gradients that governs the outward

migration of early procyclic forms towards a more alkaline environment, while late procyclic

forms remain at the inoculation site. Local acidification of the environment, and the perception

thereof, can also explain the spacing between projections emanating from the same parasite

colony, as well as the cessation of migration or the reorientation of projections from two separate

communities on the same plate. In addition, it explains why the start of migration is not solely

dependent on a threshold cell number, as was previously reported[18], but rather on the external

pH. Early procyclic forms are also able to perceive exogenous sources of acid and alkali as

repellents and attractants, respectively. Late procyclic forms are also attracted to a more basic

environment, but unlike early procyclic forms they are not repelled by acid, explaining why they

remain at the inoculation site on plates and do not perform SoMo.

Analyses of the transcriptomes of cells spontaneously performing SoMo, or reacting to

exogenous alkali or acid, are consistent with them being the same response. Comparison of the

transcriptomes of trypanosomes at the tips and roots of projections showed that migrating cells

at the tip expressed high levels of markers for early procyclic forms. This mirrors what is seen in

liquid culture[17,18] as well as at early time points after flies are infected with stumpy forms (AN

and IR, manuscript in preparation). Among the adenylate cyclases differentially expressed

between the tip and the root, ACP6 is more highly expressed at the root and is also up-regulated

in late procyclic forms. By contrast, ACP3 and ACP5, which are more highly expressed in the tip,

are not stage-regulated between early and late procyclic forms in liquid culture[17]. This is also true

of a few other transcripts, such as CRAM, and may reflect a response to culture on plates.

cAMP signalling is required for the perception of pH gradients (summarised in Fig 6f). A

PDEB1 knockout was not only defective in SoMo[28], but was also impervious to acid and alkali. In

contrast to an RNAi mutant of ACP5 which still performed SoMo normally[25], an ACP5 single

knockout did not form projections and was insensitive to exogenous acid, but was still attracted

to alkali. In this respect, its behaviour resembles that of late procyclic forms. Down-regulation or

deletion of ACP6 ([25] and [Supplementary Fig. 4b](#)) or deletion of one copy of ACP1 cause hyper-

SoMo, while ACP2 is not required for SoMo or pH taxis. ACP1, ACP5 and ACP6 are localised at the

flagellar tip ([39] and Fig. 6d), highlighting its importance as a sensor.

For the first time, we show that there is a physical link between a cyclic AMP response

protein, CARP3, and adenylate cyclases ACP3 and ACP5. Moreover, the CARP3 null mutant had

the same defects in SoMo and acid sensing as the ACP5 single knockout. Interestingly, there is

strong evidence that CARP3 is N-myristoylated[40], which could enable it to associate with the inner

leaflet of the plasma membrane. N-myristoylation has been shown to regulate signalling in

mammals[41] and it is conceivable that acylated and non-acylated forms of CARP3 enable signals

to be transmitted from the membrane to the interior of the cell. *T. brucei*, which has the most

complex life cycle of all the African trypanosomes, also encodes the largest number of adenylate

cyclases. Many of these are stage-regulated, and it is plausible that they have functions in

reacting to their local environment.

pH taxis by trypanosomes is likely to be biologically relevant during fly transmission as

there are large differences between the pH of the midgut (pH ~8) and proventriculus (pH ~10.5)[29].

This would enable parasites to follow cues provided by the fly as they migrate to different

compartments. It is established that procyclic forms respond chemotactically to other effectors

[20,21], but their identities are unknown. Chemotaxis may also explain how parasites in the

mammalian host disperse from the bloodstream into other tissues [42–44]. Finally, the concept that

self-generated gradients help drive directionality and speed expansion into new territories,

recently described for bacteria[45], may also apply to trypanosomes.

Methods

Trypanosome stocks and culture conditions

Procyclic forms of EATRO1125 were derived from bloodstream forms of *T. b. brucei* AnTat1.1[46].

Early procyclic forms were cultivated at 27 °C and 2.5 % CO_2 in DTM supplemented with 15 %

foetal bovine serum (FBS) [47]. Late procyclic forms were cultured in SDM79[48] containing 10 % FBS

at 27 °C. Parasites were maintained at a density between 10^6 and 10^7 cells ml^{-1}. Procyclic forms

of *T. b. brucei* Lister 427 and derivatives thereof were cultured in SDM79 or SDM80 containing

10 % FBS at 27 °C. SDM80 components were as described[49], except that BME vitamin solution

100x (Bioconcept 5-20K00-H) was used instead of vitamin mix 100x (Invitrogen 010144) and 1%

Pen/Strep 10'000 IU/µg (Bioconcept 4-01F00-H) was used instead of 0.1 mM kanamycin. Glucose-

free SDM80 medium was supplemented with 10% FBS, which results in a final concentration of

~1 mM glucose. If indicated, SDM80 + 10 % FBS was supplemented with 6 mM D-glucose or 6 mM

N-Acetyl-D-glucosamine (GlcNAc) from a 1.2 M stock solution in water. Population doubling

times were determined over a period of at least 5 days in which the cell density was determined

daily.

SoMo assays and community lifts

The protocol to produce plates was adapted from Imhof and coworkers[18]. Plates were always

used on the same day that they were poured. Culture medium (10 ml) supplemented with 0.4 %

agarose (Promega V3125) was poured into petri dishes (85 mm diameter). Open plates were then

air-dried for 1 hour in a laminar flow hood. Cells from an exponentially growing culture were

centrifuged briefly and resuspended in the residual medium at a density of 4 x 10^7 cells ml^{-1}. Two

hundred thousand cells were spotted per inoculum; plates were sealed with parafilm and

incubated in a humidified environment at 27 °C and 2.5 % CO_2. Photographs of plates were taken

using a Nikon E8400 camera.

Community lifts for the detection of GPEET and EP procyclins were performed as

described[18], using nitrocellulose filters (Whatman Protran BA85), K1 rabbit anti-GPEET[35] at a

dilution of 1:1000 and TBRP1/247 mouse anti-EP (Cat. no. CLP001A, Cedarlane Laboratories,

Burlington, Canada) at a dilution of 1:2500 as primary antibodies (1 h incubation time at room

temperature). The secondary antibodies goat anti-mouse IRDye 800CW (LI-COR Biosciences, Bad

Homburg, Germany) and goat anti-rabbit IRDye 680LT (LI-COR Biosciences) were used at dilutions

of 1:10000 (1 h incubation time at room temperature). The filters were rinsed 3 times in TBS +

0.05 % Tween (Sigma), followed by one final rinse in PBS before being dried overnight at room

temperature and scanned on a LI-COR Odyssey Infrared Imager model 9120, using Odyssey

Application Software, Version 3.0.21.

pH measurements on SoMo plates and in liquid cultures

To monitor pH changes in liquid culture and on agarose plates, trypanosomes were cultured in a humidified incubator at 27 °C and 2.5 % CO_2. SoMo plates were kept in the same incubator, without sealing the lids with parafilm. For analysis of growth rate and pH measurements, cells were maintained between $1 – 10 \times 10^6$ cells ml^{-1} by daily monitoring and dilution. "Log-phase" samples had densities of $3 – 9 \times 10^6$ cells ml^{-1}. If not stated otherwise, "dense" samples were obtained from cultures with a concentration of $1 - 1.5 \times 10^7$ cells ml^{-1} without further dilution for 2 days. "Medium only" refers to complete medium incubated in a culture flask in a humidified incubator at 27 °C and 2.5 % CO_2 for at least 12 h. The pH of culture supernatants or SoMo plates was measured using a micro electrode (Thermo Scientific™ Orion™ 9810BN Micro pH Electrode; fisher scientific).

Preparation of concentrated supernatant

Early procyclic forms were grown until they reached a concentration of 3×10^7 cells ml^{-1}. The cells were centrifuged for 5 minutes at 2500 g and the conditioned medium was transferred to a new tube. Conditioned medium was either concentrated directly or first sterile filtered (Jet Biofil syringe filter, 0.22 µm, REF: FVP203030K) or dialysed. 2 ml of conditioned medium were dialysed with dialysis tubing cellulose membranes (Sigma-Aldrich, cut-off 12 kDa, REF: D9777-100FT) against 50 ml DTM supplemented with 15 % FBS at 4 °C in two rounds (16 hours, then 12 hours). To concentrate the conditioned medium, 1 ml was transferred into a 1.5 ml Eppendorf tube and vacuum centrifuged for several hours until 100 µl remained in the tube. As a control, DTM only was concentrated in parallel.

pH taxis assay

Procyclic forms were inoculated onto plates as described for SoMo assays. Following incubation at 27 °C for four days, concentrates of 30 µl conditioned medium or medium only, or different chemical solutions were added to the plates at a distance of 1.5 cm from migrating projections (Fig. 3a, D = distance). The plates were sealed with parafilm and incubated at 27 °C for 16-20 hours. Stock solutions were prepared at a concentration of 1M, if not stated otherwise.

Hot phenol extraction of RNA from migrating tips

Plates containing semi-solid DTM medium were inoculated with 2×10^5 cells from a culture of procyclic form EATRO1125 (60 – 80 % GPEET-positive). Following incubation for 6 – 7 days, tip and root samples were collected from projections that were ≥2.5 cm long. For collecting tips reacting to acid or alkali, 3×10^5 cells were seeded, incubated for five days and then 30 μl 1M NaOH or 1M HCl were added on two sides of migrating cells at a distance of 1.5 cm from the tips (Fig. 3a, D = distance). In total, between 70-100 tips of migrating projections, that were exposed to acidic or alkaline solutions for 16 hours, were harvested. Only tips that clearly showed a reaction were collected (usually 5-7 tips per community).

Cells were harvested by punching out a piece of agarose using an inverted P2 micro tip (outer and inner diameters 6 and 4.5 mm, respectively). The tip subpopulation refers to the outermost 3 mm of protrusions. The root refers to the region at the base of the projection as shown in Figure 4a. Between 20-25 tips or 3-5 root samples were transferred into 1 ml DTM in a 24-well plate well. Cells were washed off the agarose pieces by careful rinsing with a P1000 pipette and the cell suspension was transferred to a 1.5 ml Eppendorf tube. The agarose pieces were rinsed once more with 0.5 ml DTM, added to the cell suspension and centrifuged for 2 minutes at 3300 x g. On average, approximately 1×10^5 cells were retrieved from each tip and 6×10^5 cells from each root sample. For tips reacting to acid or alkali, the wells of a 6-well plate contained a larger number of samples (up to 100 tips). In this case, the supernatant was carefully aspirated from the well and collected in a 15 ml Falcon tube and the agarose pieces were washed twice with 4 ml medium. The washes were added to the cells in the 15 ml Falcon tube and centrifuged for 5 minutes at 2500 g. Cell pellets were resuspended in 50 μl residual medium and RNA was extracted with hot phenol as described[50].

RNA-seq analysis

Total RNA (maximum 10 μg) was processed and used for Illumina sequencing as described previously[17]. Poly(A)-selected RNAs were used for the preparation of cDNA libraries using an Illumina TruSeq kit. Sequencing of cDNA libraries was performed at Fasteris, Geneva, using Illumina HiSeq sequencing systems with 150 bp read lengths and sequencing depths of approximately 20 million reads per sample. Reads were mapped to the *T. b. brucei* 927 reference

genome version 5.1, using the bowtie tool available in the Galaxy Interface (usegalaxy.org) with

default parameters that allow a maximum of 2 mismatches per 28 bp seed (Galaxy version

1.1.2). Read counts were extracted using "Feature-count" tool available in Galaxy using

TritrypDB-41 GFF annotation file. DESeq2 analysis were performed to identify the differentially

regulated genes and RPM values were calculated to estimate the transcripts abundance. Raw

read files are deposited at the European Nucleotide Archives (ENA) http://www.ebi.ac.uk/ena as

study PRJEB41935.

Flow cytometry

Cells were fixed in 2 % (w/v) PFA/PBS at 4°C overnight at a concentration of 1×10^7 cells ml^{-1} and

processed for flow cytometric analysis as previously described[51,52]. Rabbit antiserum α-GPEET

K1[35] and Alexa Fluor 488-conjugated secondary antibody (Invitrogen, Thermo Fisher Scientific)

were used at 1:1000 dilution in 2 % BSA/PBS and incubated for 1.5 h at 4°C with rotation. To

remove particles of subcellular size, a cut-off of 7.5×10^5 was applied to the forward scatter. A

total of ten thousand events were recorded using a ACEA NovoCyte flow cytometer and analysed

without gating using FlowJo software version 10.6.1 (BD Life Sciences).

Determination of steady-state ATP levels

ATP concentrations were measured based on a previously described ATP production assay[53]. To

measure steady-state ATP levels in whole cells, cells were washed twice in ice-cold PBS and

resuspended in PBS at 2.5×10^7 cells ml^{-1}. Denaturation with perchloric acid, precipitation and

neutralisation with KOH was performed as described[53]. The cleared lysate was serially diluted 4-

fold in H_2O. Routinely, cells were harvested in triplicate and the second dilution of the cleared

lysate (16-fold dilution) was used for the bioluminescence assay. For each replicate, ATP

concentrations were measured in triplicate. ATP concentrations were determined using the ATP

Bioluminescence Assay Kit CLS II (Roche 11699695001, from Sigma-Aldrich) according to

manufacturer's instructions. For each dilution, 10 µl was added to 40 µl of 0.5 M Tris-acetate, pH

7.75 and supplemented with luciferase reagent to a final volume of 100 µl. Titrations of the ATP

standard supplied were analysed in parallel. Luminescence was measured in 96-well microtitre

plates (opaque, non-binding) using a Turner BioSystems Modulus Microplate Luminometer,

Model 9300-001. Signals were integrated over 5 sec in steady-glo mode. Three readings per plate

were acquired, with a period of 10 min. Values of the second reading were used for analysis,

since only the second and third readings yielded stable signals.

Motility assays

Cultures were allowed to grow to a density between 4 - 7 x 10^6 cells ml^{-1} and adjusted to a final

concentration of 5 x 10^6 prior to the experiment. These cells (in medium) were placed into

motility chambers[54]. To prevent cell adherence, the chamber was precoated with 1 % poly L-

glutamic acid (Fisher Scientific, P4886-25MG) for one hour and washed twice with 70 % EtOH.

The chambers were then washed twice with 100 µl medium and once with cell suspension before

loading 100 µl of cells at a concentration of 5 x 10^6 cells ml^{-1}. The edges of the coverslip were

sealed to avoid evaporation and capillary flow of liquid. Time lapse movies were recorded at 200

ms, with 250 cycles and 10 x magnification with a Leica DM 5500B microscope. The dataset was

analysed with a tracking macro provided by Richard Wheeler, University of Oxford[55]. Velocity,

speed and individual cell tracks were plotted using Fiji version 1.0 [56] and Prism Version 6.

Immunoprecipitation and proteomic analysis

Isolation of HA-tagged protein complexes was performed as previously described[57] using the

addback cell line expressing CARP3-HA, or derivatives expressing CARP3-HA and either ACP3-myc

or ACP5-myc. Myc-tagged proteins were immunoprecipitated with 5 µg monoclonal antibody

9E10 (Cat No: MA1-980) and Dynabeads™ Pan Mouse IgG (Cat No: 11041), both from

ThermoFisher Scientific (Invitrogen). Isolated protein complexes were subjected to Western blot

analysis, or protein bands were cut from Coomassie-stained polyacrylamide gels, subjected to

trypsin digestion and proteins were identified by LC-tandem mass spectometry[58] at the

Proteomics and Mass Spectrometry Core Facility of the University of Bern.

Generation of knockouts and addbacks

Single knockouts, double knockouts and tagged addbacks were generated in the parental 427

Cas9 cell line[28]. Targeting cassettes were amplified from pPOTv7-hygromycin and pPOTv7-G418[59]

plasmids. The primers used for PCR amplification of the targeting fragments, the sgRNA

templates and the primers used for genotyping PCRs can be found in Supplementary Table 3. The

knockouts were verified by PCR ([Supplementary Table 3](#)). Transfection was performed as
described previously[60] using TbBSF transfection buffer [61]. Stable transformants were selected
using 15 µg ml^{-1} Geneticin G418 sulphate and 25 µg ml^{-1} hygromycin.

Immunofluorescence analysis (IFA) of HA- and or Myc-tagged cells

Cells were harvested at a density of 2 – 5 x 10^6 cells and then washed once with PBS. They were
left to settle on a coverslip for 20 minutes and then fixed with 4 % PFA in PBS for 4 minutes. After
removing the supernatant, the cells were permeabilized with Triton-x 100 diluted to 0.2 % in PBS
for 5 minutes and the liquid was aspirated. The samples were blocked in PBS + 4 % BSA for 1 hour
at RT. The primary antibodies (monoclonal α-HA rat 3F10 and monoclonal α-myc mouse 9E10,
from Roche, distributed by Sigma Aldrich) were diluted 1:1000 and 1:500 respectively in PBS +
0.4 % BSA and the samples incubated for 1 hour at RT. The cells were then washed 3 times with
PBS and the supernatant aspirated after every washing step. They were then incubated with the
secondary antibodies AlexaFluor 488 conjugated donkey α-rat (Life Technologies) and AlexaFluor
488 conjugated goat α-mouse (Invitrogen/Thermo Fisher) both diluted 1:2000 in PBS + 0.4 % BSA
for 1 hour at RT. The samples were then washed 3 times as described previously and incubated
with Hoechst 33342 (20µg/ml) diluted 1:500 in PBS for 2 minutes. After three washing steps, the
slides were mounted on microscope slides
with Mowiol and imaged with a Leica DM 5500B microscope with a 100x objective. The
acquired images were processed with Fiji[56].

References

1. Aframian, N. & Eldar, A. A Bacterial Tower of Babel: Quorum-Sensing Signaling Diversity
and Its Evolution. *Annu. Rev. Microbiol.* **74**, 587–606 (2020).
2. Yang, W. & Briegel, A. Diversity of Bacterial Chemosensory Arrays. *Trends Microbiol.* **28**,
68–80 (2020).
3. Daniels, R., Vanderleyden, J. & Michiels, J. Quorum sensing and swarming migration in
bacteria. *FEMS Microbiol. Rev.* **28**, 261–289 (2004).
4. Burkart, M., Toguchi, A. & Harshey, R. M. The chemotaxis system, but not chemotaxis, is
essential for swarming motility in *Escherichia coli*. *Proc. Natl. Acad. Sci. U. S. A.* **95**, 2568–

2573 (1998).

5. Tyson, J. J. & Murray, J. D. Cyclic AMP waves during aggregation of *Dictyostelium*

amoebae. *Development* **106**, 421–426 (1989).

6. Tomchik, K. J. & Devreotes, P. N. Adenosine 3',5'-monophosphate waves in *Dictyostelium*

discoideum: a demonstration by isotope dilution--fluorography. *Science* **212**, 443–446

(1981).

7. Maeda, M. *et al.* Periodic signaling controlled by an oscillatory circuit that includes

protein kinases ERK2 and PKA. *Science* **304**, 875–878 (2004).

8. Wang, M., Aerts, R. J., Spek, W. & Schaap, P. Cell cycle phase in *Dictyostelium discoideum*

is correlated with the expression of cyclic AMP production, detection, and degradation.

Involvement of cyclic AMP signaling in cell sorting. *Dev. Biol.* **125**, 410–416 (1988).

9. Kahn, S. *et al. Trypanosoma cruzi* amastigote adhesion to macrophages is facilitated by

the mannose receptor. *J. Exp. Med.* **182**, 1243–1258 (1995).

10. Kanjee, U. *et al. Plasmodium vivax* strains use alternative pathways for invasion. *J. Infect.*

Dis. (2020) doi:10.1093/infdis/jiaa592.

11. Kedzierski, L. *et al.* A leucine-rich repeat motif of *Leishmania* parasite surface antigen 2

binds to macrophages through the complement receptor 3. *J. Immunol.* **172**, 4902–4906

(2004).

12. Xing, M. *et al.* A Sialic Acid-Binding Protein SABP1 of *Toxoplasma gondii* Mediates Host

Cell Attachment and Invasion. *J. Infect. Dis.* **222**, 126–135 (2020).

13. Reuner, B., Vassella, E., Yutzy, B. & Boshart, M. Cell density triggers slender to stumpy

differentiation of *Trypanosoma brucei* bloodstream forms in culture. *Mol. Biochem.*

Parasitol. **90**, 269–280 (1997).

14. Vassella, E., Reuner, B., Yutzy, B. & Boshart, M. Differentiation of African trypanosomes is

controlled by a density sensing mechanism which signals cell cycle arrest via the cAMP

pathway. *J. Cell Sci.* **110 (Pt 2**, 2661–2671 (1997).

15. Rico, E. *et al.* Bloodstream form pre-adaptation to the tsetse fly in *Trypanosoma brucei*.

Front. Cell. Infect. Microbiol. **3**, 78 (2013).

16. Vassella, E. *et al.* A major surface glycoprotein of *trypanosoma brucei* is expressed

transiently during development and can be regulated post-transcriptionally by glycerol or

hypoxia. *Genes Dev.* **14**, 615–626 (2000).

17. Naguleswaran, A., Doiron, N. & Roditi, I. RNA-Seq analysis validates the use of culture-

derived *Trypanosoma brucei* and provides new markers for mammalian and insect life-

cycle stages. *BMC Genomics* **19**, 227 (2018).

18. Imhof, S., Knüsel, S., Gunasekera, K., Vu, X. L. & Roditi, I. Social motility of African

trypanosomes is a property of a distinct life-cycle stage that occurs early in tsetse fly

transmission. *PLoS Pathog.* **10**, e1004493 (2014).

19. Oberholzer, M., Lopez, M. A., McLelland, B. T. & Hill, K. L. Social motility in african

trypanosomes. *PLoS Pathog.* **6**, e1000739 (2010).

20. Eliaz, D. *et al.* Exosome secretion affects social motility in *Trypanosoma brucei*. *PLoS*

Pathog. **13**, e1006245 (2017).

21. DeMarco, S. F., Saada, E. A., Lopez, M. A. & Hill, K. L. Identification of Positive Chemotaxis

in the Protozoan Pathogen *Trypanosoma brucei*. *mSphere* **5**, (2020).

22. Pozzo, L. Y. *et al.* Studying taxis in real time using optical tweezers: applications for

Leishmania amazonensis parasites. *Micron* **40**, 617–620 (2009).

23. Diaz, E. *et al.* Effect of aliphatic, monocarboxylic, dicarboxylic, heterocyclic and sulphur-

containing amino acids on *Leishmania spp.* chemotaxis. *Parasitology* **142**, 1621–1630

(2015).

24. Barros, V. C., Oliveira, J. S., Melo, M. N. & Gontijo, N. F. *Leishmania amazonensis*:

chemotaxic and osmotaxic responses in promastigotes and their probable role in

development in the phlebotomine gut. *Exp. Parasitol.* **112**, 152–157 (2006).

25. Lopez, M. A., Saada, E. A. & Hill, K. L. Insect stage-specific adenylate cyclases regulate

social motility in African trypanosomes. *Eukaryot. Cell* **14**, 104–112 (2015).

26. Imhof, S., Vu, X. L., Bütikofer, P. & Roditi, I. A Glycosylation Mutant of *Trypanosoma*

brucei Links Social Motility Defects In Vitro to Impaired Colonization of Tsetse Flies In

Vivo. *Eukaryot. Cell* **14**, 588–592 (2015).

27. Jelk, J. *et al.* Glycoprotein biosynthesis in a eukaryote lacking the membrane protein Rft1.

J. Biol. Chem. **288**, 20616–20623 (2013).

28. Shaw, S. *et al.* Flagellar cAMP signaling controls trypanosome progression through host
tissues. *Nat. Commun.* **10**, (2019).

29. Liniger, M. *et al.* Cleavage of trypanosome surface glycoproteins by alkaline trypsin-like
enzyme(s) in the midgut of *Glossina morsitans*. *Int. J. Parasitol.* **33**, 1319–1328 (2003).

30. Bringaud, F., Ebikeme, C. & Boshart, M. Acetate and succinate production in amoebae,
helminths, diplomonads, trichomonads and trypanosomatids: common and diverse
metabolic strategies used by parasitic lower eukaryotes. *Parasitology* **137**, 1315–1331
(2010).

31. Besteiro, S. *et al.* Succinate secreted by *Trypanosoma brucei* is produced by a novel and
unique glycosomal enzyme, NADH-dependent fumarate reductase. *J. Biol. Chem.* **277**,
38001–38012 (2002).

32. Lee, M. G., Bihain, B. E., Russell, D. G., Deckelbaum, R. J. & Van der Ploeg, L. H.
Characterization of a cDNA encoding a cysteine-rich cell surface protein located in the
flagellar pocket of the protozoan *Trypanosoma brucei*. *Mol. Cell. Biol.* **10**, 4506–4517
(1990).

33. Affolter, M., Hemphill, A., Roditi, I., Müller, N. & Seebeck, T. The repetitive microtubule-
associated proteins MARP-1 and MARP-2 of *Trypanosoma brucei*. *J. Struct. Biol.* **112**,
241–251 (1994).

34. Ebikeme, C. E. *et al.* N-acetyl D-glucosamine stimulates growth in procyclic forms of
Trypanosoma brucei by inducing a metabolic shift. *Parasitology* **135**, 585–594 (2008).

35. Ruepp, S. *et al.* Survival of *Trypanosoma brucei* in the tsetse fly is enhanced by the
expression of specific forms of procyclin. *J. Cell Biol.* **137**, 1369–1379 (1997).

36. Zheng, B., Yao, H. & Lee, G. S. Inactivation of the gene encoding the flagellar pocket
protein, CRAM, in African trypanosomes. *Mol. Biochem. Parasitol.* **100**, 235–242 (1999).

37. Gould, M. K. *et al.* Cyclic AMP effectors in African trypanosomes revealed by genome-
scale RNA interference library screening for resistance to the phosphodiesterase
inhibitor CpdA. *Antimicrob. Agents Chemother.* **57**, 4882–4893 (2013).

38. GEIGY, R., HUBER, M., WEINMAN, D. & WYATT, G. R. Demonstration of trehalose in the
vector of African trypanosomiasis: the tsetse fly. *Acta Trop.* **16**, 255–262 (1959).

39. Saada, E. A. *et al.* Insect stage-specific receptor adenylate cyclases are localized to

distinct subdomains of the Trypanosoma brucei Flagellar membrane. *Eukaryot. Cell* **13**,

1064–1076 (2014).

40. Wright, M. H., Paape, D., Price, H. P., Smith, D. F. & Tate, E. W. Global Profiling and

Inhibition of Protein Lipidation in Vector and Host Stages of the Sleeping Sickness

Parasite *Trypanosoma brucei*. *ACS Infect. Dis.* **2**, 427–441 (2016).

41. Kwon, H., Choi, M., Ahn, Y. & Pak, Y. N-myristoylation regulates insulin-induced

phosphorylation and ubiquitination of Caveolin-2 for insulin signaling. *Biochem. Biophys.*

Res. Commun. (2020) doi:10.1016/j.bbrc.2020.08.072.

42. Caljon, G. *et al.* The Dermis as a Delivery Site of *Trypanosoma brucei* for Tsetse Flies. *PLoS*

Pathog. **12**, e1005744 (2016).

43. Capewell, P. *et al.* The skin is a significant but overlooked anatomical reservoir for vector-

borne African trypanosomes. *Elife* **5**, (2016).

44. Trindade, S. *et al. Trypanosoma brucei* Parasites Occupy and Functionally Adapt to the

Adipose Tissue in Mice. *Cell Host Microbe* **19**, 837–848 (2016).

45. Cremer, J. *et al.* Chemotaxis as a navigation strategy to boost range expansion. *Nature*

575, 658–663 (2019).

46. Le Ray, D., Barry, J. D., Easton, C. & Vickerman, K. First tsetse fly transmission of the

'AnTat' serodeme of *Trypanosoma brucei*. *Ann. Soc. Belg. Med. Trop. (1920).* **57**, 369–381

(1977).

47. Overath, P., Czichos, J. & Haas, C. The effect of citrate/cis-aconitate on oxidative

metabolism during transformation of *Trypanosoma brucei*. *Eur. J. Biochem.* **160**, 175–182

(1986).

48. Brun, R. & Schönenberger. Cultivation and in vitro cloning or procyclic culture forms of

Trypanosoma brucei in a semi-defined medium. Short communication. *Acta Trop.* **36**,

289–292 (1979).

49. Lamour, N. *et al.* Proline metabolism in procyclic *Trypanosoma brucei* is down-regulated

in the presence of glucose. *J. Biol. Chem.* **280**, 11902–11910 (2005).

50. Roditi, I., Carrington, M. & Turner, M. Expression of a polypeptide containing a dipeptide

600   repeat is confined to the insect stage of *Trypanosoma brucei*. *Nature* **325**, 272–274

601   (1987).

602 51. Roditi, I. *et al.* Procyclin gene expression and loss of the variant surface glycoprotein

603   during differentiation of *Trypanosoma brucei*. *J. Cell Biol.* **108**, 737–746 (1989).

604 52. Knüsel, S. & Roditi, I. Insights into the regulation of GPEET procyclin during differentiation

605   from early to late procyclic forms of *Trypanosoma brucei*. *Mol. Biochem. Parasitol.* **191**,

606   66–74 (2013).

607 53. Allemann, N. & Schneider, A. ATP production in isolated mitochondria of procyclic

608   *Trypanosoma brucei*. *Mol. Biochem. Parasitol.* **111**, 87–94 (2000).

609 54. Gadelha, C., Wickstead, B., de Souza, W., Gull, K. & Cunha-e-Silva, N. Cryptic paraflagellar

610   rod in endosymbiont-containing kinetoplastid protozoa. *Eukaryot. Cell* **4**, 516–525 (2005).

611 55. Wheeler, R. J. Use of chiral cell shape to ensure highly directional swimming in

612   trypanosomes. *PLoS Comput. Biol.* **13**, e1005353 (2017).

613 56. Schindelin, J. *et al.* Fiji: An open-source platform for biological-image analysis. *Nat.*

614   *Methods* **9**, 676–682 (2012).

615 57. Florini, F., Naguleswaran, A., Gharib, W. H., Bringaud, F. & Roditi, I. Unexpected diversity

616   in eukaryotic transcription revealed by the retrotransposon hotspot family of

617   *Trypanosoma brucei*. *Nucleic Acids Res.* **47**, 1725–1739 (2019).

618 58. Al Kaabi, A., Traupe, T., Stutz, M., Buchs, N. & Heller, M. Cause or effect of arteriogenesis:

619   compositional alterations of microparticles from CAD patients undergoing external

620   counterpulsation therapy. *PLoS One* **7**, e46822 (2012).

621 59. Beneke, T. *et al.* A CRISPR Cas9 high-throughput genome editing toolkit for

622   kinetoplastids. *R. Soc. open Sci.* **4**, 170095 (2017).

623 60. Burkard, G., Fragoso, C. M. & Roditi, I. Highly efficient stable transformation of

624   bloodstream forms of *Trypanosoma brucei*. *Mol. Biochem. Parasitol.* **153**, 220–223

625   (2007).

626 61. Schumann Burkard, G., Jutzi, P. & Roditi, I. Genome-wide RNAi screens in bloodstream

627   form trypanosomes identify drug transporters. *Mol. Biochem. Parasitol.* **175**, 91–94

628   (2011).

Acknowledgements

Ruth Etzensperger, Berta Pozzi, Stephanie DeMarco and Boris Striepen are thanked for

stimulating discussions and constructive comments on the manuscript. This research was

supported by the Swiss National Science Foundation (Grant numbers 310030_184669 and

31003A_166427) and the Canton of Bern.

Author contributions

S.S., S.K., A.N., D.A. and I.R. conceived, designed and conducted experiments. All authors

analysed the data. S.S. and I.R. wrote the paper.

Fig. 1 Response of migrating cells to fresh medium (DTM) and conditioned medium (CM) after different treatments. **a** Upper panel: communities exposed to DTM and CM; lower left panel: communities exposed to filtered CM on both sides; lower right panel: communities exposed to dialysed CM (12 kDa cut-off) or undialysed CM. White circles with dots indicate where solutions were pipetted onto the plates. Community lifts stained for EP (green) and GPEET (red) are shown. **b** Continuous growth in liquid culture acidifies medium. Representative pictures of culture supernatants from early procyclic forms (EATRO1125) grown in DTM for 7 days without dilution. Left, DTM only. Samples were taken on days 0, 4 and 7. The colour change from red to yellow indicates that the culture becomes more acidic with time. Cell titres are indicated.

Fig. 2 Trypanosomes acidify their environment in liquid culture and on semi-solid agarose plates. Early procyclic forms (Lister 427 Bern) grown in SDM79 were analysed. **a** pH (left y axis) of medium only, a log-phase culture (log, 3.9×10^6 cells ml^{-1}) and a dense culture (dense, 5.6×10^7 cells ml^{-1}). Three technical replicates are shown. The H^+ concentration is indicated on the right y axis. Medium only was incubated for the same length of time as the cultures. **b** and **c** pH measurements on SoMo plates. **b** Trypanosomes (2×10^5 cells) were either spotted onto the centre of a plate as a single inoculum (left row) or in pairs (right row). White numbers indicate pH measurements on days 0, 2, 3, 4 and 7 in the centre of a community, at the periphery and in the space between two communities (indicated with a white arrow). **c** Graphs for pH measurement shown in **b**. Each data point represents the mean pH measured from replicate plates, error bars depict the range (n=2). Top: plates with single communities. pH at the plate periphery (○ dashed line) or in the centre of the community (● solid line). Bottom: paired communities. pH at the plate periphery (○ dashed line), in the centres of the two communities (▼ or ▲ solid line), or between them (◆ dotted line). Y axes are the same as in **a**.

d The time-point of migration depends on the volume of the substrate. Equal numbers of early procyclic forms were inoculated onto wells that contained different volumes of medium with agarose. Plates were photographed on days 3, 4 and 5 post inoculation.

a

b

c

Fig. 3 Response of early and late procyclic forms to acidic or basic solutions. **a** Schematic representation of the pH taxis assay. "x" represents the site where a solution was spotted. "D" is the distance between the closest projection and the spot. In these experiments D= 1.3 cm. **b** Early procyclic forms (EATRO1125) on DTM plates were exposed to NaOH (OH- treatment) or HCl (H+ treatment) on both sides of the population. **c** Late procyclic forms were exposed to NaOH on the right side and HCl on the left side of the community. Scale bar: 1 cm. For **b** and **c** community lifts stained for EP (green) and GPEET (red) are shown.

Fig. 4 Volcano plots comparing transcriptomes of cells from different regions of a community or following exposure to acid or alkali. DTM plates were inoculated with cells from liquid cultures of early procyclic forms (EATRO1125) in DTM medium. **a** Comparison of the roots and tips of projections. For clarity the same volcano plot is displayed twice with different transcripts annotated. Left: early procyclic form markers, such as GPEET, calflagin, hexokinase 1 and prostaglandin F synthetase are upregulated in tips. Adenylate cyclases such as ACP3, ACP5 and ACP6 are differentially expressed between tip and root. Right: glycolytic enzymes and genes involved in glucose metabolism are upregulated in tips. Far-right panel: community lift of a representative SoMo plate 7 days after inoculation stained for EP (green) and GPEET (red). Tips were isolated when projections were ≥ 2.5 cm in length. RNA-seq was performed in duplicate. **b** Pairwise comparisons of untreated communities and communities exposed to acid or alkali. Tips were isolated from projections showing a clear response to the treatment (indicated with red circles). RNA-seq was performed in triplicate. The illlustrations in the left and right bottom corners of the volcano plots show SoMo communities (in black) and the samples that were collected (red circles).

 Fig. 5 Glucose availability impacts SoMo by acidification of the culture medium. Procyclic forms (Lister 427

Bern) grown in SDM79 or SDM80 plus supplements were analysed: SDM79 (blue / ○), SDM80+G6 (+ 6 mM

glucose, green / □), SDM80+N6 (+ 6 mM GlcNAc, red / △). If not specified, cells were adapted from

SDM79 to the medium indicated for one to three weeks. **a** Social motility on semi-solid plates was

assessed in SDM80 with varying glucose availability. Left: SoMo plates on day 6 after inoculation, prior to
performing community lifts. Cells were adapted from SDM80+G6 to the corresponding liquid medium for
5 days before inoculation. Right: merge of the corresponding community lifts probed for GPEET (red) and
EP (green). **b** Flow cytometric analysis. The percentage of GPEET-positive cells (shaded area) is indicated.
Cells grown in SDM79 were used as a control (2° antibody only, grey). **c** Cumulative growth in different
media. Growth was monitored after transferring cells from SDM79 to the corresponding medium 3 days
earlier. Numbers indicate population doubling time averaged over 7 days (mean ± SD). **d** Steady-state ATP
levels in different media. Cell lysates were harvested in two independent experiments (replicate 1 or 2,
respectively). Cell densities at harvest were between 6.2 - 8.0 or 7.3 - 8.9 x 10^6 cells ml^{-1}, respectively.
Appropriate dilutions of cell lysates were assayed; the graph depicts means and error bars (SD) of technical
replicates shown as circles (n=12 or 9, respectively). Data are shown relative to SDM80+G6. **e** Directional
motility in different media. Cell densities at harvest were between 5.9 - 6.5 x 10^6 cells ml^{-1}. Centred motility
tracks of individual cells are shown at identical scale. > 6000 tracks in each medium were pooled from
nine movies acquired from three technical replicates.
f pH measurement of liquid cultures. Culture medium (medium only) and supernatants from log-phase
(log) or dense culture (dense). Cell titres per ml in SDM79, SDM80+G6 or SDM80+N6, respectively: 3.2 x
10^6, 3.8 x 10^6 or 3.3 x 10^6 (log) and 5.1x 10^7, 4 x 10^7or 3.6 x 10^7 (dense). Three technical replicates are
shown for each condition. The corresponding H$^+$ concentration is indicated on right y axis. **g** pH
measurements on SoMo plates inoculated with cells pre-adapted to the corresponding liquid medium.
ΔpH (centre – periphery) refers to pH in centre of the community minus the pH at the plate periphery.
Each data point represents the mean ΔpH from replicate plates, error bars depict the range (n=3). The
fold change in H$^+$ concentration is indicated on the right y axis.

Fig. 6 pH responses by SoMo-defective mutants. **a-c** Community lifts labelled with anti-GPEET (red) and anti-EP (green). The images in **a**, **b**, and **c** show merged EP and GPEET signals. **a** Lister 427 wt and PDEB1 knockout (KO). **b** 427 Cas9 parental line, CARP3-KO and CARP3-HA addback. Upper panel: untreated. Lower panel: exposed to acid and alkali. **c** 427 Cas9 parental line and 3 independent ACP5 single allele knockouts; panels as in **b**.

d Myc-tagged ACP5 is localised to the flagellar tip in a cell line co-expressing CARP3-HA. Immunofluorescence was performed with an anti-myc antibody (red). DNA was stained with Hoechst dye. Scale bar = 10 μm. **e** Western blot analysis of immunoprecipitations with anti-HA and anti-myc antibodies. Myc-tagged ACP3 and ACP5 were expressed independently in the CARP3-HA background. The parental line 427 Cas9 was used as a control. Input samples have 40 times fewer cell equivalents than the

immunoprecipitated samples. IP: immunoprecipitation with anti-HA or anti-myc antibodies. **f** Summary of mutants, SoMo phenotypes and responses to pH. -/-: double knockouts; +/-: single knockouts.

Supplementary data

All supplementary figures, tables and movies are available via the following link:

https://www.dropbox.com/sh/r7od7e0s5j04g3m/AAD3wUFIbHjKGeFavE27ndAxa?dl=0

Supplementary Fig. 1 Different chemicals were tested for their impact on migrating cells on semi-solid surfaces. Procyclic forms of EATRO1125 were incubated on DTM plates. **a** All chemicals were dissolved or diluted in water to give 1 M solutions. Chemicals highlighted in red could not be dissolved completely, but were used anyway. **b** Effect on migrating cells and pH of 1M solutions. **c** SoMo assays. Chemicals were spotted on day 4 after inoculation and photographs were taken 16-20 h later. Photographs were processed to improve contrast and converted to black and white.

Supplementary Fig. 2 Independent CRAM knockout clones exposed to acid and alkali. **a** Upper panel: community lifts of CRAM knockout clones and their 427 Cas9 parental cell line. Lower panel: community lifts of knockouts and the parental line exposed to acid (H^+) and alkali (OH^-). The images show merged EP and GPEET signals. **b** Growth comparison of three individual CRAM knockouts and parental 427 Cas9.

Supplementary Fig. 3 Characterisation of CARP3 and CARP4 knockouts. **a** Genotyping of CARP3 and CARP4 knockouts by PCR. Genomic DNA was isolated from individual clones and amplified with the primers listed in Supplementary Table 3. **b** Growth comparison of CARP3 and CARP4 knockouts and parental 427 Cas9. **c** Motility assay of CARP3 knockouts. Centred motility tracks of individual cells are shown at identical scale. > 8000 tracks for each cell line (427 Cas9, CARP3 KO Cl.1, CARP3 KO Cl.2 and Trypanin KO, respectively) were pooled from nine movies acquired from three technical replicates. The trypanin knockout was used as a motility-deficient control[28]. **d** Velocity and speed of each cell line. The colour code corresponds to the cell lines in d. Data points are displayed as box plot with whiskers representing 2.5 - 97.5 percentile, filled circles represent points outside that range. InterQuartileRange IQR (25 - 75 percentile), with Median (50 percentile). **e** Percentage of GPEET-positive cells (shaded area) is indicated. Cells without prior incubation with anti-GPEET antiserum were used as a negative control (2° antibody only, grey). **f** Localisation of CARP3-HA (green) by immunofluorescence. DNA was stained with Hoechst dye. Scale bar = 10 µm.

Supplementary Fig. 4 Characterisation of ACP knockouts. **a** Genotyping of ACP knockouts by PCR. Genomic DNA was isolated and PCRs were performed with the primers listed in Supplementary

Table 3. **b** SoMo and pH taxis assays. Plates were inoculated with the cell lines indicated and 4 days later HCl and NaOH were applied to the plates (indicated by dots). Pictures were taken 16-20 h after spotting of chemicals. **c** Growth of ACP5 single knockout clones compared to the parental 427 Cas9 cell line. **d** Motility assays of ACP5 single knockouts. Centred motility tracks of individual cells are shown at identical scale. > 7000 tracks for each clone were pooled from nine movies acquired from three technical replicates. The parental line 427 Cas9 and Trypanin KO were used as controls. **e** Velocity and speed of individual cell lines. The colour code corresponds to the cell lines in d. Data points are displayed as box plot with whiskers representing 2.5 - 97.5 percentile, filled circles represent points outside that range. InterQuartileRange IQR (25 - 75 percentile), with Median (50 percentile). **f** Flow cytometric analysis of percentages of GPEET-positive cells.

Supplementary Table 1 RNAseq data. **a** Comparison of the roots and tips of projections. Transcripts from the tips either upregulated >1 Log2FC or downregulated <1 Log2FC in comparison to the root. FC: fold change. **b** Pairwise comparisons of untreated communities and communities exposed to acid or alkali. Transcripts from tips that were either upregulated >1 Log2FC or downregulated <1 Log2FC. FC: fold change.
Raw read files are deposited at the European Nucleotide Archives (ENA)
http://www.ebi.ac.uk/ena as study PRJEB41935.

Supplementary Table 2 ACP3 and ACP5 are immunoprecipitated with CARP3-HA.

Supplementary Table 3 List of primers used for knockouts and genotyping.

Supplementary Video 1 Early procyclic form parasites exhibit social motility.

Supplementary Video 2 Response of early procyclic forms to hydrochloric acid (HCl). Black arrows indicate where solutions were spotted. The video is shown as a rewind loop.

Supplementary Video 3 Response of early procyclic forms to sodium hydroxide (NaOH). Black arrows indicate where solutions were spotted. The video is shown as a rewind loop.

Supplementary Video 4 Response of late procyclic forms to sodium hydroxide (NaOH). Black arrows indicate where solutions were spotted. The video is shown as a rewind loop.