Cyclic AMP signalling and glucose metabolism mediate pH taxis by African trypanosomes

Sebastian Shaw[1,2, #a], Sebastian Knüsel[1,♣], Daniel Abbühl[1, ♣, #b], Arunasalam Naguleswaran[1,♣],
Ruth Etzensperger[1], Mattias Benninger[1] and Isabel Roditi[1,*]

[1] Institute of Cell Biology, University of Bern, Bern, Switzerland

[2] Graduate School of Cellular and Biomedical Sciences, University of Bern, Bern, Switzerland

♣ These authors contributed equally

[#a] Current Address: Department of Pathobiology, School of Veterinary Medicine, University of
Pennsylvania, Philadelphia, PA 19104, USA

[#b] Current Address: Trypanosome Cell Biology Unit, INSERM U1201, Institute Pasteur, 75015
Paris, France

* Corresponding author

Email: isabel.roditi@izb.unibe.ch

Abstract

The collective movement of African trypanosomes on semi-solid surfaces, known as social motility, is presumed to be due to migration factors and repellents released by the parasites. Here we show that procyclic (insect midgut) forms acidify their environment as a consequence of glucose metabolism, generating pH gradients by diffusion. Early and late procyclic forms exhibit self-organising properties on agarose plates. While early procyclic forms are repelled by acid and migrate outwards, late procyclic forms remain at the inoculation site. Furthermore, trypanosomes respond to exogenously formed pH gradients, with both early and late procyclic forms being attracted to alkali. pH taxis is mediated by multiple cyclic AMP effectors: deletion of one copy of adenylate cyclase ACP5, or both copies of the cyclic AMP response protein CARP3, abrogates the response to acid, while deletion of phosphodiesterase PDEB1 completely abolishes pH taxis. The ability to sense pH is biologically relevant as trypanosomes experience large changes as they migrate through their tsetse host. Supporting this, a CARP3 null mutant is severely compromised in its ability to establish infections in flies. Based on these findings, we propose that the expanded family of adenylate cyclases in trypanosomes might govern other chemotactic responses in their two hosts.

Introduction

Environmental sensing, chemotaxis and quorum sensing present possibilities for pathogens to orient themselves within their hosts. All three are widespread in bacteria, and the sensors and response mechanisms have been characterised in detail[1,2]. Genes involved in bacterial chemotaxis and quorum sensing have long been linked to biofilm formation and swarming behaviour[3,4]. By contrast, very little is known about chemotaxis in unicellular eukaryotes, with the exception of *Dictyostelium discoideum*, which generates and reacts to cyclic AMP (cAMP) gradients, promoting aggregation and differentiation[5–8].

Intracellular pathogens frequently exploit receptor-ligand interactions to recognise and invade the appropriate cell types[9–12]. For extracellular parasites such as *Trypanosoma brucei spp.*, causative agents of human and animal trypanosomiasis, it is not known what drives them to move from one host tissue to another. In their mammalian hosts, quorum sensing regulates the titre of *T. brucei* by triggering the differentiation of proliferative slender forms to quiescent stumpy forms[13,14]. When these are taken up by tsetse flies during a blood meal[15] they differentiate to early procyclic (midgut) forms and then to late procyclic forms[16]. In addition to the surface marker GPEET procyclin, which is only expressed by early procyclic forms, several metabolic enzymes, nutrient transporters and signal transducers are differentially expressed between the two forms[17,18]. Colonisation of the midgut by procyclic forms is followed by additional rounds of differentiation as parasites invade the proventriculus, and then move to the salivary glands before being transmitted to a new mammalian host. At present, however, the cues allowing trypanosomes to orient themselves within their hosts are unknown.

Early procyclic forms engage in a form of group migration, known as social motility (SoMo), when cultured on semi-solid surfaces such as agarose plates[18,19]. The parasites initially remain at the inoculation site, dividing approximately once every 24 h, then the communities start to form regularly spaced projections that extend by ~1 cm (500 body lengths) per day. Communities on the same plate are able to sense each other and reorient to avoid contact (Supplementary Video 1). Late procyclic forms do not exhibit SoMo, but they do produce a repellent that is sensed by early procyclic forms[18]. Communities on plates also react to other signals. Trypanosomes depleted of the spliced leader core protein SmD1 shed exosomes that act

as repellents[20], while bacterial colonies can act as attractants[21]. Various species of *Leishmania*,

parasites related to trypanosomes, respond chemotactically to a range of compounds including

amino acids, carbohydrates and sera, in liquid culture[22–25]. In none of these cases is it known

which genes mediate the responses.

Few genes have been implicated in SoMo. Concomitant depletion of adenylate cyclases

ACP1&2, or depletion of ACP6 on its own, resulted in a hyper-social phenotype (hyper-SoMo)

where the communities formed more projections and migrated more rapidly, suggesting that

these are negative regulators[26]. Interestingly, ACP1 and ACP6 are upregulated in late procyclic

forms[17], which are SoMo-negative[18]. The flagellar phosphodiesterase B1 (PDEB1)[19] and the

endoplasmic reticulum protein Rft1[27,28] are positive regulators of SoMo. Importantly, null

mutants of PDEB1 and Rft1 were not only impaired in SoMo, but also defective in colonising the

tsetse midgut[27,29]. We postulated that SoMo is largely a chemotactic response to factors

produced by the trypanosomes themselves. Here we show that procyclic forms can generate pH

gradients through glucose metabolism and that the response of early and late procyclic forms to

these gradients can explain the self-organising properties of communities on plates. In addition,

we identified two cAMP signalling components involved in the response to acid, ACP5 and the

cAMP response protein CARP3. Notably, pH sensing is completely abolished in PDEB1 knockout,

where cAMP levels are perturbed. pH taxis could be biologically relevant as there is a pH gradient

in the tissues that are colonised by the parasites during their transmission through the insect

host[30]. In support of this, we show that CARP3 null mutant parasites are severely compromised

in their ability to establish fly infections.

Results

Parasite-secreted factors change properties of the medium and repel migrating projections

What are the factors causing collective migration in SoMo? The fact that communities sense and

avoid each other over considerable distances means that they react to attractants or repellents

that are secreted, or otherwise released, by the parasites. We started by investigating if these

factors were produced under normal growth conditions in liquid culture. We concentrated the

supernatant of a dense culture (2×10^7 cells ml^{-1}) 10-fold and spotted this "conditioned medium"

(CM) next to migrating projections on SoMo plates (Fig. 1a, upper panel). Projections migrating

towards the site where CM was spotted reoriented away from it, indicating that it was perceived

as a repellent. In a second experiment, CM was either sterile filtered or dialysed (with a 12 kDa

cut-off) prior to concentration and spotted next to migrating projections. As seen with untreated

CM, filtered CM still repelled migrating projections (Fig. 1a, lower left panel), but dialysed CM

lost its repellent properties (Fig. 1a, lower right panel). On the contrary, migrating parasites were

attracted to dialysed CM in the same way that they were to concentrated fresh medium (DTM)

(Fig. 1a). These results suggested that repellents might be metabolites that were lost upon

dialysis.

Trypanosomes acidify their environment in liquid culture and on agarose plates

Trypanosomes secrete a variety of molecules that influence the pH of their environment[31,32].

Indicators for pH, such as phenol red, are commonly used for culture media and it is apparent

that dense cultures are more acidic than fresh medium or log phase cultures, as illustrated for

early procyclic forms growing in liquid culture (Fig. 1b). We therefore used a small tip micro glass

pH electrode to measure the pH in liquid culture and on agarose plates (Fig. 2). Fresh medium

has a pH of 7.46 ($\pm$ 0.02) and the supernatant of a log phase culture has a pH of 7.45 ($\pm$ 0.01) (Fig.

2a). The pH decreased to 6.95 ($\pm$ 0.00) when parasites were allowed to overgrow; this

corresponds to ~3-fold increase in the concentration of H^+ ions (Fig. 2a).

To measure the pH on plates, 2 x 10^5 parasites were inoculated either in the centre of a

plate (Fig. 2b, left row), or in pairs at a defined distance from each other (Fig. 2b, right row), and

the pH was measured after 2, 3 ,4 and 7 days post inoculation. We measured the pH in the centre

of a community, at the periphery of a plate and between two communities. The pH at the centres

of the communities decreased from 7.50 ($\pm$ 0.01) to 7.31 ($\pm$ 0.05) during the first three days and

decreased further to 6.96 ($\pm$ 0.05) between days 3 and 7 (Fig. 2b, c). The pH at the periphery also

started to decrease between day 3 and 7, possibly because of diffusion of H^+ ions from the

parasite communities. When communities were plated in pairs, the pH at the midpoint between

them also decreased over time (Fig. 2b, right row, arrow). Thus, communities acidify their vicinity

resulting in pH gradients on the plates.

We showed previously that trypanosomes needed to reach a threshold number before

they started migrating[18]. We now demonstrate that it is not the cell number *per se* that is critical,

but rather the need to establish a discernible pH gradient. When the same number of parasites

were inoculated onto plates containing different volumes of semi-solid medium, the time point

at which projections started to form was inversely proportional to the volume (Fig. 2d). This

would be consistent with larger volumes buffering the pH more effectively and requiring more

cells, and thus more time, before a gradient was established. We cannot exclude, however, that

other factors are also affected by the volume of the medium.

Reaction of early and late procyclic forms to exposure to acid and alkali

As a next step, we tested if manipulating the pH gradient had an influence on migration by

exposing early procyclic forms to external sources of acid or alkali (Fig. 3a, b). Early procyclic

forms were inoculated onto plates and incubated until the projections reached a length of about

2 cm. At this point, solutions of HCl or NaOH were pipetted onto the surface at a defined distance

(Fig. 3a) and the plates were incubated for 16-20 h. The migrating cells responded very strongly

to both solutions (Fig. 3b, Supplementary Videos 2 & 3). Exogenous NaOH acted as an attractant,

causing projections to reorient towards it (OH$^-$ treatment), while HCl caused the projections to

stop moving outwards, or to move away from the acid (H$^+$ treatment). Using the same assay,

another 22 chemicals were tested (Supplementary Fig. 1). In most cases, when a solution was

acidic, the projections were repelled, and when a solution was basic, the cells were attracted to

it. These experiments also showed that the projections reacted to the pH rather than to specific

cations or anions in the solutions.

We next tested late procyclic forms since these do not perform SoMo. Late procyclic

forms did not respond to HCl (Fig. 3c, Supplementary Video 4), but they formed a limited number

of projections that migrated towards NaOH. Community lifts (Fig. 3c) confirmed that the

parasites were still GPEET-negative and had not dedifferentiated to early forms. In summary,

these results demonstrate that procyclic forms exhibit pH taxis, with alkali acting as an attractant

for both early and late procyclic forms. In contrast, there are stage-specific differences in their

reaction to acid.

RNA-seq analyses suggest that SoMo is a response to self-generated gradients

To better understand the mechanisms behind self-organisation of early and late procyclic forms, and their pH responses, we performed RNA-seq under different conditions. A culture of procyclic forms (>60 % GPEET-positive) was plated and allowed to form projections (Fig. 4a, far right). A community lift with antibodies against procyclins showed that the tips were strongly positive for GPEET, while the centre of the community stained more strongly for EP. Cells were isolated from the tips and roots of projections (Fig. 4a, far right) and analysed by RNA-seq. Data confirmed that markers of early procyclic forms were more highly expressed in the tips than the roots (Fig. 4a, left; Supplementary Table 1a). Other transcripts that were more highly expressed in the tip were not characteristic of early procyclic forms in liquid culture, such as cysteine-rich acidic integral membrane protein (CRAM)[33], a microtubule-associated repetitive protein (MARP2)[34] and a hypothetical protein (hypo; Tb927.9.10400). In addition, several adenylate cyclases were differentially expressed on plates; ACP3 and ACP5 were ≥2-fold higher in the tip, while ACP6 and another adenylate cyclase (GRESAG 4; Tb927.11.1480) were ≥2-fold higher in the root. The up-regulation of high affinity glucose transporters (THT2 and 2A), glycolytic enzymes and components of the glycosome (Fig. 4a, middle and Supplementary Table 1a) is suggestive of higher glucose metabolism by the tip than the root. Increased expression of these transcripts is also characteristic of early procyclic forms in liquid culture[17,18].

RNA-seq was also performed on the tips of communities exposed to acid or alkali (Fig. 4b and Supplementary Table 1b). There were no significant differences (≥2-fold, p=0.05) between non-exposed communities and those exposed to alkali. A single cation transporter (CatT, Tb927.11.9000) was more highly expressed in cells exposed to acid, but the read counts were extremely low in both cases. CatT and a copy of a contig (Tb11.v5.0514) were the only significant differences in a comparison of acid- versus alkali-treated communities. Taken together, these results suggest that self-generated pH gradients might be the normal driver of migration on SoMo plates.

Glucose contributes to the formation of pH gradients

Since the RNA-seq data indicated that cells at the tip might metabolise glucose more actively than cells at the root, we investigated the role of glucose in SoMo. For these experiments SDM80 medium was prepared with 1 mM glucose, 6 mM glucose or 6mM N-acetylglucosamine (GlcNAc), which inhibits glucose uptake by binding to hexose transporters[35]. Early procyclic forms of Lister 427 Bern were inoculated onto plates and incubated for 6 days (Fig. 5a). The parasites in SDM80 with 1 mM glucose started migrating later than cells in SDM80 containing 6 mM glucose, and the SoMo defect was even more pronounced when glucose uptake was blocked by GlcNAc. Under these conditions the cells only formed rudimentary "knobs" at the edge of the community and failed to form projections, even at later time points.

The reduced ability to migrate on plates can have a variety of causes, especially in the context of glucose restriction. It might be due to longer population doubling times, motility defects or inefficient energy metabolism. Procyclic cultures of Lister 427 Bern are stably GPEET-positive[27,36]. Nevertheless, we assessed the proportion of early procyclic forms by flow cytometry (Fig. 5b). Regardless of the medium in which the cells were grown, all cultures were ≥97 % GPEET-positive. Furthermore, the glucose concentration had no impact on the population doubling time (Fig. 5c) or intracellular ATP levels (Fig. 5d), both assessed in liquid culture. Previous work showed that directional motility is important for SoMo and migration through the fly[29]. We therefore assessed cell motility (Fig. 5e); no differences were observed between cells grown in the presence or absence of glucose.

To investigate whether different glucose concentrations influence the culture pH, we compared medium alone with liquid cultures of trypanosomes in log phase or high density (Fig. 5f). While the pH of log-phase cultures did not differ from that of medium alone, dense cultures of parasites in SDM79 or SDM80 with 6 mM glucose strongly acidified the medium. The concentration of H^+ in dense cultures, compared to log phase cultures, increased 2.53-fold (±0.03) for SDM79 and 2.51-fold (±0.06) for SDM80 plus glucose. By contrast, cells grown in SDM80 plus GlcNAc showed a smaller change in H^+ ion concentration (1.22-fold ±0.03). We next measured pH changes for communities growing on plates (Fig. 5g). The ΔpH between the centre and the periphery reached -0.31($\pm$ 0.02) for parasites on SDM79 and -0.37($\pm$ 0.02) for SDM80 +

205 6mM glucose, corresponding to a ≥2-fold difference in H^+ ions at the two locations, whereas

206 plates containing SDM80 + GlcNAc did not show a difference in pH. Taken together, these data

207 show that glucose is required for parasites to generate a pH gradient, and that this correlates

208 with their ability to perform SoMo.

Cyclic AMP signalling is involved in the response to pH

RNA-seq identified a number of differentially expressed transcripts, including that of the flagellar

pocket protein CRAM, which was more highly expressed in the tips than in the roots (Fig. 4a).

CRAM is not essential in procyclic culture forms[37]. When we generated knockouts, these grew at

the same rate as their parental line, but two out of three independent clones produced smaller

projections. Nevertheless, all three reacted to acid and alkali in the same way as wild-type cells

(Supplementary Fig. 2) so we did not proceed with additional analyses.

Several adenylate cyclases were also differentially expressed between the tip and the root

(Fig. 4a). Before we started analysing these and other members of the cAMP signalling pathway,

we tested whether PDEB1, which hydrolyses cAMP, had an influence on the pH response. When

the PDEB1 null mutant was exposed to alkali or acid, it showed no pH response (Fig. 6a). Notably,

the PDEB1 null mutant differs from both early procyclic forms, which react to acid and alkali, and

late procyclic forms, which form projections in response to alkali but do not respond to acid

(Supplementary Table 2).

Four cAMP response proteins (CARP1-CARP4) were previously identified in an RNAi

screen as mediating resistance to an inhibitor of PDEB1[38], but they have not been studied in the

context of SoMo. In our attempts to generate knockout cell lines for CARP 1 and 2, we only

obtained clones after single rounds of transfection. We did not pursue these further, however,

as the resistance genes were not integrated at the correct loci. We were able to obtain knockouts

of CARP3 and 4, however (Supplementary Fig. 3a). CARP4 null mutants did not show any

difference to the parental line when they were subjected to a standard SoMo assay

(Supplementary Fig. 3b) and were not analysed further. Deletion of CARP3, while not affecting

growth in liquid culture (Supplementary Fig. 3c) or on plates (Supplementary Fig. 4c and 4d), or

expression of the early marker GPEET (Supplementary Fig. 3d) resulted in a major SoMo defect

(Fig 6b, upper panel) reminiscent of the PDEB1 null mutant. The CARP3 knockout did not display

a motility defect in liquid culture (Supplementary Fig. 4a and b) and still reacted to alkali, but did

not react to acid on plates (Fig 6b, lower panel). An HA-tagged addback of CARP3, expressed

from a procyclin locus, partially rescued the SoMo phenotype (Fig 6b, upper panel), although the

projections were shorter than in the parental line. Importantly, the addback was now able to

sense and react to both acid and alkali (Fig 6b, lower panel).

We next assessed the requirement for CARP3 *in vivo*. For these experiments we generated

a null mutant in wild-type *T. b. brucei* Lister 427 using a recently described CRISPR/Cas9 transient

expression system[39]. In addition, we engineered an addback ectopically expressing an untagged

version of CARP3 (Supplementary Fig. 5). Flies were infected with either the wild type, the

knockout or the addback and evaluated for the prevalence and intensity of midgut infections

after 14 days. The knockout showed a significantly lower midgut infection rate ($p < 0.00001$,

Fisher's exact test, two-sided) when compared to the wild type. In addition, infection intensities

were reduced (Fig 6c). Both the prevalence and intensities of infection with the addback were

comparable to the wild type. Thus, the defect in establishing a midgut infection can be attributed

specifically to the loss of CARP3. In addition, whereas 55-60% of flies heavily infected with the

wild type, and 100% heavily infected with the addback, had trypanosomes in the proventriculus,

the knockout gave no heavy midgut infections nor did it colonise the proventriculus

(Supplementary Table 3).

CARP3 is present in both the cytoplasm and the flagellum (Supplementary Fig. 4e;

www.tryptag.org). Since no interaction partners have been reported, a pulldown was performed

with HA-tagged CARP3 (Fig. 6d) and analysed by mass spectrometry. This led to the identification

of adenylate cyclases ACP3 and ACP5 in two biological replicates (Supplementary Table 4). These

interactions were confirmed by *in situ* tagging and co-immunoprecipitation of myc-tagged

versions of ACP3 and ACP5 in the CARP3-HA addback line (Fig. 6d and Supplementary Table 4).

The reciprocal co-immunoprecipitations with anti-myc antibody confirmed the interaction for

both ACP3 and ACP5 (Fig. 6d and Supplementary Table 2). These results, combined with their

differential expression in the tips and roots of projections (Fig. 4a) stimulated us to revisit some

of the flagellar ACP previously characterised by Lopez and coworkers[26]. In their study they

showed that knockdown of ACP1/2 or ACP6 by RNAi resulted in a hyper-SoMo phenotype. We

obtained null mutants of ACP2 and ACP6, but it was only possible to generate single knockouts

of ACP1 and ACP5 (Supplementary Fig. 6a) implying that they are essential at this stage of the life

cycle. Although we attempted to knock out ACP3, and isolated resistant cell lines, the antibiotic

resistance genes were not integrated at the correct locus (data not shown). The full knockout of

ACP6 and the semi-knockout of ACP1 both exhibited hyper-SoMo phenotypes, while the full

knockout of ACP2 showed no phenotype (Supplementary Fig. 6b). These results are consistent

with the findings with the RNAi lines; they also indicate that the hyper-SoMo phenotype in the

previously published ACP1/2 RNAi line[26] was due to depletion of ACP1. Unexpectedly, however,

and in contrast to the results with RNAi[26], single knockouts of ACP5 were unable to form

projections on plates (Fig. 6e, upper panel). In terms of their pH response, ACP5 single knockouts

resembled CARP3 double knockouts - they were able to form projections migrating towards an

alkaline source, but were refractory to acid (Fig. 6e, lower panel). Further experiments showed

that >80 % of these cells were GPEET-positive and that the phenotype of the ACP5 mutants was

not due to defects in growth or motility (Supplementary Fig. 6c-6f). These experiments also

revealed that ACP5-myc was localised exclusively at the flagellar tip (Fig 6f). In summary, we have

extended the catalogue of cAMP signalling components that are involved in regulating SoMo. In

addition, the finding that PDEB1, ACP5 and CARP3 play a role in the perception of environmental

acidity/alkalinity is consistent with SoMo being a response to self-generated pH gradients.

Discussion

It was proposed more than a decade ago that the collective migration of trypanosomes on plates

could be explained by a combination of migration factors and repellents released by the

parasites[19], but the identity of these molecules was unknown. Here we show for the first time

that trypanosomes acidify their environment as a consequence of glucose metabolism,

generating pH gradients by diffusion. While it is commonly assumed that free glucose levels in

tsetse are low between blood meals, and this is often stated in reviews, there are no primary

publications on this topic. The fact that transcripts encoding high affinity glucose transporters

and glycolytic enzymes are not only upregulated in SoMo tips (Figure 4) and in early procyclic

forms[17], but also in trypanosomes at the beginning of a fly infection[40], is compatible with procyclic

forms utilising carbon sources other than proline in tsetse.

Early and late procyclic forms exhibit self-organising properties. It is the perception of

these self-generated pH gradients that governs the outward migration of early procyclic forms

towards a more alkaline environment, while late procyclic forms remain at the inoculation site.

Local acidification of the environment, and the response to it, can also explain the spacing

between projections emanating from the same parasite colony, as well as the cessation of

migration or the reorientation of projections from two separate communities on the same plate.

In addition, it explains why the start of migration is not solely dependent on a threshold cell

number, as was previously reported[18], but rather on the external pH. Early procyclic forms are

also able to perceive exogenous sources of acid and alkali as repellents and attractants,

respectively. Late procyclic forms are also attracted to a more basic environment, but unlike early

procyclic forms they are not repelled by acid, explaining why they remain at the inoculation site

on plates and do not perform SoMo.

Analyses of the transcriptomes of cells spontaneously performing SoMo, or reacting to

exogenous alkali or acid, are consistent with them being the same response. Comparison of the

transcriptomes of trypanosomes at the tips and roots of projections showed that migrating cells

at the tip expressed high levels of markers for early procyclic forms. This mirrors what is seen in

liquid culture[17,18] as well as at early time points after flies are infected with stumpy forms[40].

Among the adenylate cyclases differentially expressed between the tip and the root, ACP6 is

more highly expressed at the root and is also up-regulated in late procyclic forms. By contrast,

ACP3 and ACP5, which are more highly expressed in the tip, are not stage-regulated between

early and late procyclic forms in liquid culture[17]. This is also true of a few other transcripts, such

as CRAM, and may reflect a response to culture on plates.

cAMP signalling is required for the perception of pH gradients (summarised in Fig. 6g). A

PDEB1 knockout was not only defective in SoMo[29], but was also impervious to acid and alkali. In

contrast to an RNAi mutant of ACP5 which still performed SoMo normally[26], an ACP5 single

knockout did not form projections and was insensitive to exogenous acid, but was still attracted

to alkali. In this respect, its behaviour resembles that of late procyclic forms. Down-regulation or

deletion of ACP6 ([26] and Supplementary Fig. 6b) or deletion of one copy of ACP1 cause hyper-

SoMo, while ACP2 is not required for SoMo or pH taxis. ACP1, ACP5 and ACP6 are localised at the

flagellar tip ([41] and Fig. 6f), highlighting its importance as a sensor.

For the first time, we show that there is a physical link between a cyclic AMP response

protein, CARP3, and adenylate cyclases ACP3 and ACP5. Moreover, the CARP3 null mutant had

the same defects in SoMo and acid sensing as the ACP5 single knockout. Interestingly, there is

strong evidence that CARP3 is N-myristoylated[42], which could enable it to associate with the inner

leaflet of the plasma membrane. N-myristoylation has been shown to regulate signalling in

mammals[43] and it is conceivable that acylated and non-acylated forms of CARP3 enable signals

to be transmitted from the membrane to the interior of the cell. *T. brucei*, which has the most

complex life cycle of all the African trypanosomes, also encodes the largest number of adenylate

cyclases. Many of these are stage-regulated, and it is plausible that they have functions in

reacting to their local environment. It is also worth noting that CARP3 interacts with other

adenylate cyclases, although ACP3 and ACP5 were the top hits (Supplementary Table 4), and that

it was recently identified by proximity labelling as interacting with ACP1[44].

pH taxis by trypanosomes is likely to be biologically relevant during fly transmission as

there are large differences between the pH of the midgut (pH ~8) and proventriculus (pH ~10.5)[30].

This would enable parasites to follow cues provided by the fly as they migrate to different

compartments. The hindgut is also more basic than the posterior midgut[30], and trypanosomes

have been detected there, but these have abnormal morphologies[45]. At present we do not know

if there are additional chemotactic cues attracting trypanosomes towards to the proventriculus

or repelling them from the hindgut. Regardless of whether they migrate in one or both directions,

only parasites migrating to the proventriculus can survive and complete the life cycle. We have

shown that CARP3 is required for efficient colonisation of tsetse. Interestingly, the knockout has

a more severe phenotype than the PDEB1 knockout described previously[29]. The finding that

CARP3 interacts with several adenylate cyclases in addition to ACP3 and 5 suggests that it may

act as a node for signal perception via cAMP signalling. It is established that procyclic forms

respond chemotactically to other effectors[20,21], but their identities are unknown. Chemotaxis

may also explain how parasites in the mammalian host disperse from the bloodstream into other

tissues [46–48]. Finally, the concept that self-generated gradients help drive directionality and speed

expansion into new territories, recently described for bacteria[49], may also apply to

trypanosomes.

Methods

Trypanosome stocks and culture conditions

Procyclic forms of EATRO1125 were derived from bloodstream forms of *T. b. brucei* AnTat1.1[50].

Early procyclic forms were cultivated at 27 °C and 2.5 % CO_2 in DTM supplemented with 15 %

foetal bovine serum (FBS) [51]. Late procyclic forms were cultured in SDM79[52] containing 10 % FBS

at 27 °C. Parasites were maintained at a density between 10^6 and 10^7 cells ml^{-1}. Procyclic forms

of *T. b. brucei* Lister 427 and derivatives thereof were cultured in SDM79 or SDM80 containing

10 % FBS at 27 °C. SDM80 components were as described[53], except that BME vitamin solution

100x (Bioconcept 5-20K00-H) was used instead of vitamin mix 100x (Invitrogen 010144) and 1%

Pen/Strep stock solution (Bioconcept 4-01F00-H) was used instead of 0.1 mM kanamycin.

Glucose-free SDM80 medium was supplemented with 10 % FBS, which results in a final

concentration of ~1 mM glucose. If indicated, SDM80 + 10 % FBS was supplemented with 6 mM

D-glucose or 6 mM N-Acetyl-D-glucosamine (GlcNAc) from a 1.2 M stock solution in water.

Population doubling times were determined over a period of at least 5 days in which the cell

density was determined daily.

SoMo assays and community lifts

The protocol to produce plates was adapted from Imhof and coworkers[18]. Plates were always

used on the same day that they were poured. Culture medium (10 ml) supplemented with 0.4 %

agarose (Promega V3125) was poured into petri dishes (85 mm diameter). Open plates were then

air-dried for 1 hour in a laminar flow hood. Cells from an exponentially growing culture were

centrifuged briefly and resuspended in the residual medium at a density of 4 x 10^7 cells ml^{-1}. Two

hundred thousand cells were spotted per inoculum; plates were sealed with parafilm and

incubated in a humidified environment at 27 °C and 2.5 % CO_2. Photographs of plates were taken

using a Nikon E8400 camera.

Community lifts for the detection of GPEET and EP procyclins were performed as

described[18], using nitrocellulose filters (Whatman Protran BA85), K1 rabbit anti-GPEET[36] at a

dilution of 1:1000 and TBRP1/247 mouse anti-EP (Cat. no. CLP001A, Cedarlane Laboratories,

Burlington, Canada) at a dilution of 1:2500 as primary antibodies (1 h incubation time at room

temperature). The secondary antibodies goat anti-mouse IRDye 800CW (LI-COR Biosciences, Bad

Homburg, Germany) and goat anti-rabbit IRDye 680LT (LI-COR Biosciences) were used at dilutions

of 1:10000 (1 h incubation time at room temperature). The filters were rinsed 3 times in TBS +

0.05 % Tween (Sigma), followed by one final rinse in PBS before being dried overnight at room

temperature and scanned on a LI-COR Odyssey Infrared Imager model 9120, using Odyssey

Application Software, Version 3.0.21.

pH measurements on SoMo plates and in liquid cultures

To monitor pH changes in liquid culture and on agarose plates, trypanosomes were cultured in a

humidified incubator at 27 °C and 2.5 % CO_2. SoMo plates were kept in the same incubator,

without sealing the lids with parafilm. For analysis of growth rate and pH measurements, cells

were maintained between $1 - 10 \times 10^6$ cells ml^{-1} by daily monitoring and dilution. "Log-phase"

samples had densities of $3 - 9 \times 10^6$ cells ml^{-1}. If not stated otherwise, "dense" samples were

obtained from cultures with a concentration of $1 - 1.5 \times 10^7$ cells ml^{-1} without further dilution for

2 days. "Medium only" refers to complete medium incubated in a culture flask in a humidified

incubator at 27 °C and 2.5 % CO_2 for at least 12 h. The pH of culture supernatants or SoMo plates

was measured using a micro electrode (Thermo Scientific™ Orion™ 9810BN Micro pH Electrode;

fisher scientific).

Preparation of concentrated supernatant

Early procyclic forms were grown until they reached a concentration of 3×10^7 cells ml^{-1}. The cells

were centrifuged for 5 minutes at 2500 g and the conditioned medium was transferred to a new

tube. Conditioned medium was either concentrated directly or first sterile filtered (Jet Biofil

syringe filter, 0.22 µm, REF: FVP203030K) or dialysed. 2 ml of conditioned medium were dialysed

with dialysis tubing cellulose membranes (Sigma-Aldrich, cut-off 12 kDa, REF: D9777-100FT)

against 50 ml DTM supplemented with 15 % FBS at 4 °C in two rounds (16 hours, then 12 hours).

To concentrate the conditioned medium, 1 ml was transferred into a 1.5 ml Eppendorf tube and

vacuum centrifuged for several hours until 100 µl remained in the tube. As a control, DTM only

was concentrated in parallel.

pH taxis assay

Procyclic forms were inoculated onto plates as described for SoMo assays. Following incubation

at 27 °C for four days, concentrates of 30 µl conditioned medium or medium only, or different

chemical solutions were added to the plates at a distance of 1.5 cm from migrating projections

(Fig. 3a, D = distance). The plates were sealed with parafilm and incubated at 27 °C for 16-20

hours. Stock solutions were prepared at a concentration of 1M, if not stated otherwise.

Hot phenol extraction of RNA from migrating tips

Plates containing semi-solid DTM medium were inoculated with 2 x 10^5 cells from a culture of

procyclic form EATRO1125 (60 – 80 % GPEET-positive). Following incubation for 6 – 7 days, tip

and root samples were collected from projections that were ≥2.5 cm long. For collecting tips

reacting to acid or alkali, 3 x 10^5 cells were seeded, incubated for five days and then 30 µl 1M

NaOH or 1M HCl were added on two sides of migrating cells at a distance of 1.5 cm from the tips

(Fig. 3a, D = distance). In total, between 70-100 tips of migrating projections, that were exposed

to acidic or alkaline solutions for 16 hours, were harvested. Only tips that clearly showed a

reaction were collected (usually 5-7 tips per community).

Cells were harvested by punching out a piece of agarose using an inverted P2 micro tip

(outer and inner diameters 6 and 4.5 mm, respectively). The tip subpopulation refers to the

outermost 3 mm of protrusions. The root refers to the region at the base of the projection as

shown in Figure 4a. Between 20-25 tips or 3-5 root samples were transferred into 1 ml DTM in a

24-well plate well. Cells were washed off the agarose pieces by careful rinsing with a P1000

pipette and the cell suspension was transferred to a 1.5 ml Eppendorf tube. The agarose pieces

were rinsed once more with 0.5 ml DTM, added to the cell suspension and centrifuged for 2

minutes at 3300 x g. On average, approximately 1 x 10^5 cells were retrieved from each tip and 6

x 10^5 cells from each root sample. For tips reacting to acid or alkali, the wells of a 6-well plate

contained a larger number of samples (up to 100 tips). In this case, the supernatant was carefully

aspirated from the well and collected in a 15 ml Falcon tube and the agarose pieces were washed

twice with 4 ml medium. The washes were added to the cells in the 15 ml Falcon tube and

centrifuged for 5 minutes at 2500 g. Cell pellets were resuspended in 50 µl residual medium and

RNA was extracted with hot phenol as described[54].

RNA-seq analysis

Total RNA (maximum 10 µg) was processed and used for Illumina sequencing as described

previously[17]. Poly(A)-selected RNAs were used for the preparation of cDNA libraries using an

Illumina TruSeq kit. Sequencing of cDNA libraries was performed at Fasteris, Geneva, using

Illumina HiSeq sequencing systems with 150 bp read lengths and sequencing depths of

approximately 20 million reads per sample. Reads were mapped to the *T. b. brucei* 927 reference

genome version 5.1, using the bowtie2 tool available in the Galaxy Interface (usegalaxy.org) with

default parameters that allow a maximum of 2 mismatches per 28 bp seed (Galaxy version

1.1.2). Read counts were extracted using "featureCounts" tool available in Galaxy using

TritrypDB-41 GFF annotation file. DESeq2 analysis were performed to identify the differentially

regulated genes and RPM values were calculated to estimate the transcripts abundance. Raw

read files are deposited at the European Nucleotide Archives (ENA) http://www.ebi.ac.uk/ena as

study PRJEB41935.

Flow cytometry

Cells were fixed in 2 % (w/v) PFA/PBS at 4°C overnight at a concentration of 1 x 10^7 cells ml^{-1} and

processed for flow cytometric analysis as previously described[55,56]. Rabbit antiserum α-GPEET

K1[36] and Alexa Fluor 488-conjugated secondary antibody (Invitrogen, Thermo Fisher Scientific)

were used at 1:1000 dilution in 2 % BSA/PBS and incubated for 1.5 h at 4 °C with rotation. To

remove particles of subcellular size, a cut-off of 7.5 x 10^5 was applied to the forward scatter. A

total of ten thousand events were recorded using a ACEA NovoCyte flow cytometer and analysed

without gating using FlowJo software version 10.6.1 (BD Life Sciences).

Determination of steady-state ATP levels

ATP concentrations were measured based on a previously described ATP production assay[57]. To measure steady-state ATP levels in whole cells, cells were washed twice in ice-cold PBS and resuspended in PBS at 2.5×10^7 cells ml^{-1}. Denaturation with perchloric acid, precipitation and neutralisation with KOH was performed as described[57]. The cleared lysate was serially diluted 4-fold in H_2O. Routinely, cells were harvested in triplicate and the second dilution of the cleared lysate (16-fold dilution) was used for the bioluminescence assay. For each replicate, ATP concentrations were measured in triplicate. ATP concentrations were determined using the ATP Bioluminescence Assay Kit CLS II (Roche 11699695001, from Sigma-Aldrich) according to manufacturer's instructions. For each dilution, 10 µl was added to 40 µl of 0.5 M Tris-acetate, pH 7.75 and supplemented with luciferase reagent to a final volume of 100 µl. Titrations of the ATP standard supplied were analysed in parallel. Luminescence was measured in 96-well microtitre plates (opaque, non-binding) using a Turner BioSystems Modulus Microplate Luminometer, Model 9300-001. Signals were integrated over 5 sec in steady-glo mode. Three readings per plate were acquired, with a period of 10 min. Values of the second reading were used for analysis, since only the second and third readings yielded stable signals.

Motility assays

Cultures were allowed to grow to a density between 4 - 7×10^6 cells ml^{-1} and adjusted to a final concentration of 5×10^6 prior to the experiment. These cells (in medium) were placed into motility chambers[58]. To prevent cell adherence, the chamber was precoated with 1 % poly L-glutamic acid (Fisher Scientific, P4886-25MG) for one hour and washed twice with 70 % EtOH. The chambers were then washed twice with 100 µl medium and once with cell suspension before loading 100 µl of cells at a concentration of 5×10^6 cells ml^{-1}. The edges of the coverslip were sealed to avoid evaporation and capillary flow of liquid. Time lapse movies were recorded at 200 ms, with 250 cycles and 10 x magnification with a Leica DM 5500B microscope. The dataset was analysed with a tracking macro provided by Richard Wheeler, University of Oxford[59]. Velocity, speed and individual cell tracks were plotted using Fiji version 1.0[60] and Prism Version 6.

Immunoprecipitation and proteomic analysis

Isolation of HA-tagged protein complexes was performed as previously described[61] using the addback cell line expressing CARP3-HA, or derivatives expressing CARP3-HA and either ACP3-myc or ACP5-myc. Myc-tagged proteins were immunoprecipitated with 5 µg monoclonal antibody 9E10 (Cat No: MA1-980) and Dynabeads™ Pan Mouse IgG (Cat No: 11041), both from ThermoFisher Scientific (Invitrogen). Isolated protein complexes were subjected to Western blot analysis, or protein bands were cut from Coomassie-stained polyacrylamide gels, subjected to trypsin digestion and proteins were identified by LC-tandem mass spectometry[62] at the Proteomics and Mass Spectrometry Core Facility of the University of Bern.

Generation of knockouts and addbacks

Single knockouts, double knockouts and tagged addbacks used for all *in vitro* experiments were generated in the parental 427 Cas9 cell line[29]. The CARP3 double knockout used for fly experiments was generated in *T. b. brucei* Lister 427[36] using transiently expressed Cas9 and T7 RNA polymerase[39]. An addback ectopically expressing CARP3 was derived from the null mutant. Targeting cassettes were amplified from pPOTv7-hygromycin and pPOTv7-G418[63] plasmids. The primers used for PCR amplification of the targeting fragments and the sgRNA templates were designed using LeishGEdit. These primers and the primers used for genotyping PCRs can be found in Supplementary Table 5. The knockouts and the addback were verified by PCR (Supplementary Table 5). To generate addbacks, HA-tagged and untagged versions of CARP3 were cloned into the pGAPRONE-ΔLII[64] derivate pG-mcs-puro[65], linearised with Spe I and integrated upstream of a procyclin locus on chromosome 6. Transfection was performed as described previously[66] using TbBSF transfection buffer[67]. Stable transformants were selected using 15 µg ml^{-1} Geneticin G418 sulphate, 25 µg ml^{-1} hygromycin and 1 µg ml^{-1} puromycin.

Immunofluorescence analysis (IFA) of HA- and or Myc-tagged cells

Cells were harvested at a density of $2 - 5 \times 10^6$ cells and then washed once with PBS. They were left to settle on a coverslip for 20 minutes and then fixed with 4 % PFA in PBS for 4 minutes. After removing the supernatant, the cells were permeabilized with Triton-x 100 diluted to 0.2 % in PBS

for 5 minutes and the liquid was aspirated. The samples were blocked in PBS + 4 % BSA for 1 hour

at RT. The primary antibodies (monoclonal α-HA rat 3F10 and monoclonal α-myc mouse 9E10,

from Roche, distributed by Sigma Aldrich) were diluted 1:1000 and 1:500 respectively in PBS +

0.4 % BSA and the samples incubated for 1 hour at RT. The cells were then washed 3 times with

PBS and the supernatant aspirated after every washing step. They were then incubated with the

secondary antibodies AlexaFluor 488 conjugated donkey α-rat (Life Technologies) and AlexaFluor

488 conjugated goat α-mouse (Invitrogen/Thermo Fisher) both diluted 1:2000 in PBS + 0.4 % BSA

for 1 hour at RT. The samples were then washed 3 times as described for the primary antibodies

and incubated with Hoechst 33342 (20µg/ml) diluted 1:500 in PBS for 2 minutes. After three

washing steps, the cover slips were mounted on microscope slides with Mowiol and imaged with

a Leica DM 5500B microscope with a 100x objective. The acquired images were processed with

Fiji[60].

Fly infections

Glossina morsitans morsitans pupae were purchased from the Department of Entomology,

Slovak Academy of Science, Bratislava, Slovakia or were a kind gift from Jan van den Abbeele,

Institute of Tropical Medicine, Antwerp, Belgium. Teneral flies were infected by membrane

feeding with 2.5×10^6 parasites ml^{-1} and maintained on horse blood as described[36], except that

pupae and flies were kept at 24°C, 65 -70 % humidity, rather than 27 °C in the original protocol.

Flies were dissected and midgut infections were analysed on days 13 and 14 post infection. The

intensities of infections were graded as described[36].

References

1.   Aframian, N. & Eldar, A. A Bacterial Tower of Babel: Quorum-Sensing Signaling Diversity

and Its Evolution. *Annu. Rev. Microbiol.* **74**, 587–606 (2020).

2.   Yang, W. & Briegel, A. Diversity of Bacterial Chemosensory Arrays. *Trends Microbiol.* **28**,

68–80 (2020).

3.   Daniels, R., Vanderleyden, J. & Michiels, J. Quorum sensing and swarming migration in

bacteria. *FEMS Microbiol. Rev.* **28**, 261–289 (2004).

4.   Burkart, M., Toguchi, A. & Harshey, R. M. The chemotaxis system, but not chemotaxis, is

essential for swarming motility in *Escherichia coli*. *Proc. Natl. Acad. Sci. U. S. A.* **95**, 2568–

2573 (1998).

5.   Tyson, J. J. & Murray, J. D. Cyclic AMP waves during aggregation of *Dictyostelium*

amoebae. *Development* **106**, 421–426 (1989).

6.   Tomchik, K. J. & Devreotes, P. N. Adenosine 3',5'-monophosphate waves in *Dictyostelium*

discoideum: a demonstration by isotope dilution--fluorography. *Science* **212**, 443–446

(1981).

7.   Maeda, M. *et al.* Periodic signaling controlled by an oscillatory circuit that includes

protein kinases ERK2 and PKA. *Science* **304**, 875–878 (2004).

8.   Wang, M., Aerts, R. J., Spek, W. & Schaap, P. Cell cycle phase in *Dictyostelium discoideum*

is correlated with the expression of cyclic AMP production, detection, and degradation.

Involvement of cyclic AMP signaling in cell sorting. *Dev. Biol.* **125**, 410–416 (1988).

9.   Kahn, S. *et al. Trypanosoma cruzi* amastigote adhesion to macrophages is facilitated by

the mannose receptor. *J. Exp. Med.* **182**, 1243–1258 (1995).

10.   Kanjee, U. *et al. Plasmodium vivax* strains use alternative pathways for invasion. *J. Infect.*

Dis. (2020) doi:10.1093/infdis/jiaa592.

11.   Kedzierski, L. *et al.* A leucine-rich repeat motif of *Leishmania* parasite surface antigen 2

binds to macrophages through the complement receptor 3. *J. Immunol.* **172**, 4902–4906

(2004).

12.   Xing, M. *et al.* A Sialic Acid-Binding Protein SABP1 of *Toxoplasma gondii* Mediates Host

Cell Attachment and Invasion. *J. Infect. Dis.* **222**, 126–135 (2020).

13. Reuner, B., Vassella, E., Yutzy, B. & Boshart, M. Cell density triggers slender to stumpy

differentiation of *Trypanosoma brucei* bloodstream forms in culture. *Mol. Biochem.*

Parasitol. **90**, 269–280 (1997).

14. Vassella, E., Reuner, B., Yutzy, B. & Boshart, M. Differentiation of African trypanosomes is

controlled by a density sensing mechanism which signals cell cycle arrest via the cAMP

pathway. *J. Cell Sci.* **110 (Pt 2**, 2661–2671 (1997).

15. Rico, E. *et al.* Bloodstream form pre-adaptation to the tsetse fly in *Trypanosoma brucei*.

Front. Cell. Infect. Microbiol. **3**, 78 (2013).

16. Vassella, E. *et al.* A major surface glycoprotein of *trypanosoma brucei* is expressed

transiently during development and can be regulated post-transcriptionally by glycerol or

hypoxia. *Genes Dev.* **14**, 615–626 (2000).

17. Naguleswaran, A., Doiron, N. & Roditi, I. RNA-Seq analysis validates the use of culture-

derived *Trypanosoma brucei* and provides new markers for mammalian and insect life-

cycle stages. *BMC Genomics* **19**, 227 (2018).

18. Imhof, S., Knüsel, S., Gunasekera, K., Vu, X. L. & Roditi, I. Social motility of African

trypanosomes is a property of a distinct life-cycle stage that occurs early in tsetse fly

transmission. *PLoS Pathog.* **10**, e1004493 (2014).

19. Oberholzer, M., Lopez, M. A., McLelland, B. T. & Hill, K. L. Social motility in african

trypanosomes. *PLoS Pathog.* **6**, e1000739 (2010).

20. Eliaz, D. *et al.* Exosome secretion affects social motility in *Trypanosoma brucei*. *PLoS*

Pathog. **13**, e1006245 (2017).

21. DeMarco, S. F., Saada, E. A., Lopez, M. A. & Hill, K. L. Identification of Positive Chemotaxis

in the Protozoan Pathogen *Trypanosoma brucei*. *mSphere* **5**, (2020).

22. Diaz, E. *et al.* Effect of aliphatic, monocarboxylic, dicarboxylic, heterocyclic and sulphur-

containing amino acids on *Leishmania spp.* chemotaxis. *Parasitology* **142**, 1621–1630

(2015).

23. Pozzo, L. Y. *et al.* Studying taxis in real time using optical tweezers: applications for

Leishmania amazonensis parasites. *Micron* **40**, 617–620 (2009).

24. Barros, V. C., Oliveira, J. S., Melo, M. N. & Gontijo, N. F. *Leishmania amazonensis*:

chemotaxic and osmotaxic responses in promastigotes and their probable role in

development in the phlebotomine gut. *Exp. Parasitol.* **112**, 152–157 (2006).

25. Findlay, R. C. *et al.* High-speed, three-dimensional imaging reveals chemotactic behaviour

specific to human-infective Leishmania parasites. *Elife* **10**, (2021).

26. Lopez, M. A., Saada, E. A. & Hill, K. L. Insect stage-specific adenylate cyclases regulate

social motility in African trypanosomes. *Eukaryot. Cell* **14**, 104–112 (2015).

27. Imhof, S., Vu, X. L., Bütikofer, P. & Roditi, I. A Glycosylation Mutant of *Trypanosoma*

brucei Links Social Motility Defects In Vitro to Impaired Colonization of Tsetse Flies In

Vivo. *Eukaryot. Cell* **14**, 588–592 (2015).

28. Jelk, J. *et al.* Glycoprotein biosynthesis in a eukaryote lacking the membrane protein Rft1.

J. Biol. Chem. **288**, 20616–20623 (2013).

29. Shaw, S. *et al.* Flagellar cAMP signaling controls trypanosome progression through host

tissues. *Nat. Commun.* **10**, (2019).

30. Liniger, M. *et al.* Cleavage of trypanosome surface glycoproteins by alkaline trypsin-like

enzyme(s) in the midgut of *Glossina morsitans*. *Int. J. Parasitol.* **33**, 1319–1328 (2003).

31. Bringaud, F., Ebikeme, C. & Boshart, M. Acetate and succinate production in amoebae,

helminths, diplomonads, trichomonads and trypanosomatids: common and diverse

metabolic strategies used by parasitic lower eukaryotes. *Parasitology* **137**, 1315–1331

(2010).

32. Besteiro, S. *et al.* Succinate secreted by *Trypanosoma brucei* is produced by a novel and

unique glycosomal enzyme, NADH-dependent fumarate reductase. *J. Biol. Chem.* **277**,

38001–38012 (2002).

33. Lee, M. G., Bihain, B. E., Russell, D. G., Deckelbaum, R. J. & Van der Ploeg, L. H.

Characterization of a cDNA encoding a cysteine-rich cell surface protein located in the

flagellar pocket of the protozoan *Trypanosoma brucei*. *Mol. Cell. Biol.* **10**, 4506–4517

(1990).

34. Affolter, M., Hemphill, A., Roditi, I., Müller, N. & Seebeck, T. The repetitive microtubule-

associated proteins MARP-1 and MARP-2 of *Trypanosoma brucei*. *J. Struct. Biol.* **112**,

241–251 (1994).

35. Ebikeme, C. E. *et al.* N-acetyl D-glucosamine stimulates growth in procyclic forms of
Trypanosoma brucei by inducing a metabolic shift. *Parasitology* **135**, 585–594 (2008).
36. Ruepp, S. *et al.* Survival of *Trypanosoma brucei* in the tsetse fly is enhanced by the
expression of specific forms of procyclin. *J. Cell Biol.* **137**, 1369–1379 (1997).
37. Zheng, B., Yao, H. & Lee, G. S. Inactivation of the gene encoding the flagellar pocket
protein, CRAM, in African trypanosomes. *Mol. Biochem. Parasitol.* **100**, 235–242 (1999).
38. Gould, M. K. *et al.* Cyclic AMP effectors in African trypanosomes revealed by genome-
scale RNA interference library screening for resistance to the phosphodiesterase
inhibitor CpdA. *Antimicrob. Agents Chemother.* **57**, 4882–4893 (2013).
39. Shaw, S., Knüsel, S., Hoenner, S. & Roditi, I. A transient CRISPR/Cas9 expression system
for genome editing in Trypanosoma brucei. *BMC Res. Notes* **13**, 268 (2020).
40. Naguleswaran, A. *et al.* Developmental changes and metabolic reprogramming during
establishment of infection and progression of Trypanosoma brucei brucei through its
insect host. *PLoS Negl. Trop. Dis.* **15**, e0009504 (2021).
41. Saada, E. A. *et al.* Insect stage-specific receptor adenylate cyclases are localized to
distinct subdomains of the Trypanosoma brucei Flagellar membrane. *Eukaryot. Cell* **13**,
1064–1076 (2014).
42. Wright, M. H., Paape, D., Price, H. P., Smith, D. F. & Tate, E. W. Global Profiling and
Inhibition of Protein Lipidation in Vector and Host Stages of the Sleeping Sickness
Parasite *Trypanosoma brucei*. *ACS Infect. Dis.* **2**, 427–441 (2016).
43. Kwon, H., Choi, M., Ahn, Y. & Pak, Y. N-myristoylation regulates insulin-induced
phosphorylation and ubiquitination of Caveolin-2 for insulin signaling. *Biochem. Biophys.*
Res. Commun. (2020) doi:10.1016/j.bbrc.2020.08.072.
44. Vélez-Ramírez, D. E. *et al.* APEX2 Proximity Proteomics Resolves Flagellum Subdomains
and Identifies Flagellum Tip-Specific Proteins in Trypanosoma brucei. *mSphere* **6**, (2021).
45. Rose, C. *et al. Trypanosoma brucei* colonizes the tsetse gut via an immature peritrophic
matrix in the proventriculus. *Nat. Microbiol.* **5**, 909–916 (2020).
46. Caljon, G. *et al.* The Dermis as a Delivery Site of *Trypanosoma brucei* for Tsetse Flies. *PLoS*
Pathog. **12**, e1005744 (2016).

47.   Capewell, P. *et al.* The skin is a significant but overlooked anatomical reservoir for vector-
651   borne African trypanosomes. *Elife* **5**, (2016).

652   48.   Trindade, S. *et al. Trypanosoma brucei* Parasites Occupy and Functionally Adapt to the
653   Adipose Tissue in Mice. *Cell Host Microbe* **19**, 837–848 (2016).

654   49.   Cremer, J. *et al.* Chemotaxis as a navigation strategy to boost range expansion. *Nature*
655   **575**, 658–663 (2019).

656   50.   Le Ray, D., Barry, J. D., Easton, C. & Vickerman, K. First tsetse fly transmission of the
657   'AnTat' serodeme of *Trypanosoma brucei. Ann. Soc. Belg. Med. Trop. (1920).* **57**, 369–381
658   (1977).

659   51.   Overath, P., Czichos, J. & Haas, C. The effect of citrate/cis-aconitate on oxidative
660   metabolism during transformation of *Trypanosoma brucei. Eur. J. Biochem.* **160**, 175–182
661   (1986).

662   52.   Brun, R. & Schönenberger. Cultivation and in vitro cloning or procyclic culture forms of
663   *Trypanosoma brucei* in a semi-defined medium. Short communication. *Acta Trop.* **36**,
664   289–292 (1979).

665   53.   Lamour, N. *et al.* Proline metabolism in procyclic *Trypanosoma brucei* is down-regulated
666   in the presence of glucose. *J. Biol. Chem.* **280**, 11902–11910 (2005).

667   54.   Roditi, I., Carrington, M. & Turner, M. Expression of a polypeptide containing a dipeptide
668   repeat is confined to the insect stage of *Trypanosoma brucei. Nature* **325**, 272–274
669   (1987).

670   55.   Roditi, I. *et al.* Procyclin gene expression and loss of the variant surface glycoprotein
671   during differentiation of *Trypanosoma brucei. J. Cell Biol.* **108**, 737–746 (1989).

672   56.   Knüsel, S. & Roditi, I. Insights into the regulation of GPEET procyclin during differentiation
673   from early to late procyclic forms of *Trypanosoma brucei. Mol. Biochem. Parasitol.* **191**,
674   66–74 (2013).

675   57.   Allemann, N. & Schneider, A. ATP production in isolated mitochondria of procyclic
676   *Trypanosoma brucei. Mol. Biochem. Parasitol.* **111**, 87–94 (2000).

677   58.   Gadelha, C., Wickstead, B., de Souza, W., Gull, K. & Cunha-e-Silva, N. Cryptic paraflagellar
678   rod in endosymbiont-containing kinetoplastid protozoa. *Eukaryot. Cell* **4**, 516–525 (2005).

59. Wheeler, R. J. Use of chiral cell shape to ensure highly directional swimming in

trypanosomes. *PLoS Comput. Biol.* **13**, e1005353 (2017).

60. Schindelin, J. *et al.* Fiji: An open-source platform for biological-image analysis. *Nat.*

Methods **9**, 676–682 (2012).

61. Florini, F., Naguleswaran, A., Gharib, W. H., Bringaud, F. & Roditi, I. Unexpected diversity

in eukaryotic transcription revealed by the retrotransposon hotspot family of

Trypanosoma brucei. *Nucleic Acids Res.* **47**, 1725–1739 (2019).

62. Al Kaabi, A., Traupe, T., Stutz, M., Buchs, N. & Heller, M. Cause or effect of arteriogenesis:

compositional alterations of microparticles from CAD patients undergoing external

counterpulsation therapy. *PLoS One* **7**, e46822 (2012).

63. Beneke, T. *et al.* A CRISPR Cas9 high-throughput genome editing toolkit for

kinetoplastids. *R. Soc. open Sci.* **4**, 170095 (2017).

64. Furger, A., Schürch, N., Kurath, U. & Roditi, I. Elements in the 3' untranslated region of

procyclin mRNA regulate expression in insect forms of Trypanosoma brucei by

modulating RNA stability and translation. *Mol. Cell. Biol.* **17**, 4372–4380 (1997).

65. Furger, A., Jungi, T. W., Salomone, J. Y., Weynants, V. & Roditi, I. Stable expression of

biologically active recombinant bovine interleukin-4 in Trypanosoma brucei. *FEBS Lett.*

508, 90–94 (2001).

66. Burkard, G., Fragoso, C. M. & Roditi, I. Highly efficient stable transformation of

bloodstream forms of *Trypanosoma brucei*. *Mol. Biochem. Parasitol.* **153**, 220–223

(2007).

67. Schumann Burkard, G., Jutzi, P. & Roditi, I. Genome-wide RNAi screens in bloodstream

form trypanosomes identify drug transporters. *Mol. Biochem. Parasitol.* **175**, 91–94

(2011).

Acknowledgements

Berta Pozzi, Stephanie DeMarco and Boris Striepen are thanked for stimulating discussions and

constructive comments on the manuscript. We are indebted to Jan van den Abbeele, Institute

of Tropical Medicine, Antwerp for a gift of tsetse pupae.

This research was supported by the Swiss National Science Foundation (Grant numbers
310030_184669 and 31003A_166427) and the Canton of Bern.

Author contributions

S.S., S.K., A.N., D.A. and I.R. conceived, designed and conducted experiments. R.E. and M.B.

conducted experiments. S.S., S.K., A.N., D.A., R.E. and I.R. analysed the data. S.S. and I.R. wrote

the paper.

Figure Legends

Fig. 1 Response of migrating cells to fresh medium (DTM) and conditioned medium (CM) after

different treatments. **a** Upper panel: communities exposed to DTM and CM; lower left panel:

communities exposed to filtered CM on both sides; lower right panel: communities exposed to

dialysed CM (12 kDa cut-off) or undialysed CM. White circles with dots indicate where solutions

were pipetted onto the plates. Community lifts stained for EP (green) and GPEET (red) are shown.

b Continuous growth in liquid culture acidifies medium. Representative pictures of culture

supernatants from early procyclic forms (EATRO1125) grown in DTM for 7 days without dilution.

Left, DTM only. Samples were taken on days 0, 4 and 7. The colour change from red to yellow

indicates that the culture becomes more acidic with time. Cell titres are indicated.

Fig. 2 Trypanosomes acidify their environment in liquid culture and on semi-solid agarose plates**.**

Early procyclic forms (Lister 427 Bern) grown in SDM79 were analysed. **a** pH (left y axis) of

medium only, a log-phase culture (log, 3.9 x 10^6 cells ml^{-1}) and a dense culture (dense, 5.6 x 10^7

cells ml^{-1}). Three technical replicates are shown. The H$^+$ concentration is indicated on the right y

axis. Medium only was incubated for the same length of time as the cultures. **b** and **c** pH

measurements on SoMo plates. **b** Trypanosomes (2 x 10^5 cells) were either spotted onto the

centre of a plate as a single inoculum (left row) or in pairs (right row). White numbers indicate

pH measurements on days 0, 2, 3, 4 and 7 in the centre of a community, at the periphery and in

the space between two communities (indicated with a white arrow). **c** Graphs for pH

measurement shown in **b**. Each data point represents the mean pH measured from replicate

plates, error bars depict the range (n=2). Top: plates with single communities. pH at the plate

periphery (○ dashed line) or in the centre of the community (● solid line). Bottom: paired

communities. pH at the plate periphery (○ dashed line), in the centres of the two communities

(▼ or ▲ solid line), or between them (◆ dotted line). Y axes are the same as in **a.**

d The time-point of migration depends on the volume of the substrate. Equal numbers of early

procyclic forms were inoculated onto wells that contained different volumes of medium with

agarose. Plates were photographed on days 3, 4 and 5 post inoculation.

Fig. 3 Response of early and late procyclic forms to acidic or basic solutions. **a** Schematic

representation of the pH taxis assay. "x" represents the site where a solution was spotted. "D" is

the distance between the closest projection and the spot. In these experiments D= 1.3 cm. **b** Early

procyclic forms (EATRO1125) on DTM plates were exposed to NaOH (OH- treatment) or HCl (H+

treatment) on both sides of the population. **c** Late procyclic forms were exposed to NaOH on the

right side and HCl on the left side of the community. Scale bar: 1 cm. For **b** and **c** community lifts

stained for EP (green) and GPEET (red) are shown.

Fig. 4 Volcano plots comparing transcriptomes of cells from different regions of a community or

following exposure to acid or alkali. DTM plates were inoculated with cells from liquid cultures of

early procyclic forms (EATRO1125) in DTM medium. **a** Comparison of the roots and tips of

projections. For clarity the same volcano plot is displayed twice with different transcripts

annotated. Left: early procyclic form markers, such as GPEET, calflagin, hexokinase 1 and

prostaglandin F synthetase are upregulated in tips. Adenylate cyclases such as ACP3, ACP5 and

ACP6 are differentially expressed between tip and root. Right: glycolytic enzymes and genes

involved in glucose metabolism are upregulated in tips. Far-right panel: community lift of a

representative SoMo plate 7 days after inoculation stained for EP (green) and GPEET (red). Tips

were isolated when projections were ≥ 2.5 cm in length. RNA-seq was performed in duplicate. **b**

Pairwise comparisons of untreated communities and communities exposed to acid or alkali. Tips

were isolated from projections showing a clear response to the treatment (indicated with red

circles). RNA-seq was performed in triplicate. The illustrations in the left and right bottom corners

of the volcano plots show SoMo communities (in black) and the samples that were collected (red

circles).

Fig. 5 Glucose availability impacts SoMo by acidification of the culture medium. Procyclic forms (Lister 427 Bern) grown in SDM79 or SDM80 plus supplements were analysed: SDM79 (blue / ○), SDM80+G6 (+ 6 mM glucose, green / □), SDM80+N6 (+ 6 mM GlcNAc, red / △). If not specified, cells were adapted from SDM79 to the medium indicated for one to three weeks. **a** Social motility on semi-solid plates was assessed in SDM80 with varying glucose availability. Left: SoMo plates on day 6 after inoculation, prior to performing community lifts. Cells were adapted from SDM80+G6 to the corresponding liquid medium for 5 days before inoculation. Right: merge of the corresponding community lifts probed for GPEET (red) and EP (green). **b** Flow cytometric analysis. The percentage of GPEET-positive cells (shaded area) is indicated. Cells grown in SDM79 were used as a control (2° antibody only, grey). **c** Cumulative growth in different media. Growth was monitored after transferring cells from SDM79 to the corresponding medium 3 days earlier. Numbers indicate population doubling time averaged over 7 days (mean ± SD). **d** Steady-state ATP levels in different media. Cell lysates were harvested in two independent experiments (replicate 1 or 2, respectively). Cell densities at harvest were between 6.2 - 8.0 or 7.3 - 8.9 x 10^6 cells ml^{-1}, respectively. Appropriate dilutions of cell lysates were assayed; the graph depicts means and error bars (SD) of technical replicates shown as circles (n=12 or 9, respectively). Data are shown relative to SDM80+G6. **e** Directional motility in different media. Cell densities at harvest were between 5.9 - 6.5 x 10^6 cells ml^{-1}. Centred motility tracks of individual cells are shown at identical scale. > 6000 tracks in each medium were pooled from nine movies acquired from three technical replicates.

f pH measurement of liquid cultures. Culture medium (medium only) and supernatants from log-phase (log) or dense culture (dense). Cell titres per ml in SDM79, SDM80+G6 or SDM80+N6, respectively: 3.2 x 10^6, 3.8 x 10^6 or 3.3 x 10^6 (log) and 5.1x 10^7, 4 x 10^7or 3.6 x 10^7 (dense). Three technical replicates are shown for each condition. The corresponding H$^+$ concentration is indicated on right y axis. **g** pH measurements on SoMo plates inoculated with cells pre-adapted to the corresponding liquid medium. ΔpH (centre – periphery) refers to pH in centre of the community minus the pH at the plate periphery. Each data point represents the mean ΔpH from replicate plates, error bars depict the range (n=3). The fold change in H$^+$ concentration is indicated on the right y axis.

Fig. 6 CARP3 is required for acid sensing in vitro and for establishment of infection in flies.

a Merged EP and GPEET signals of Lister 427 wt and PDEB1 knockout (KO). **b** Merged EP and GPEET signals of 427 Cas9 parental line, CARP3 knockout and CARP3-HA addback. Upper panel: untreated. Lower panel: exposed to acid and alkali. **c** Teneral flies were infected with wild-type Lister 427 procyclic forms (WT), the CARP3 double knockout (CARP3-KO) and the addback (AB). Dissections were performed 13-14 days post infection. Total numbers are derived from two replicates per cell line (Supplementary Table 3). The intensity of midgut infections was scored as heavy, intermediate or weak as described[36]. The p-value is shown for Fisher's exact test, two-sided. **d** Western blot analysis of immunoprecipitations with anti-HA and anti-myc antibodies (representative blot from three independent experiments). Myc-tagged ACP3 and ACP5 were expressed independently in the CARP3-HA background. The parental line 427 Cas9 was used as a control. Input samples have 40 times fewer cell equivalents than the immunoprecipitated samples. IP: immunoprecipitation with anti-HA or anti-myc antibodies. **e** Merged EP and GPEET signals of the 427 Cas9 parental line and 3 independent ACP5 single allele knockouts; panels as in **b**. **f** Myc-tagged ACP5 is localised to the flagellar tip in a cell line co-expressing CARP3-HA. Immunofluorescence was performed with an anti-myc antibody (red). DNA was stained with Hoechst dye. Scale bar = 10 μm. **g** Summary of mutants, SoMo phenotypes and responses to pH. -/-: double knockouts; +/-: single knockouts.

Supplementary Fig. 1 Different chemicals were tested for their impact on migrating cells on semi-solid surfaces. Procyclic forms of EATRO1125 were incubated on DTM plates. **a** All chemicals were dissolved or diluted in water to give 1 M solutions. Chemicals highlighted in red could not be dissolved completely, but were used anyway. **b** Effect on migrating cells and pH of 1M solutions. **c** SoMo assays. Chemicals were spotted on day 4 after inoculation and photographs were taken 16-20 h later. Photographs were processed to improve contrast and converted to black and white.

Supplementary Fig. 2 Independent CRAM knockout clones exposed to acid and alkali. **a** Upper panel: community lifts of CRAM knockout clones and their 427 Cas9 parental cell line. Lower

panel: community lifts of knockouts and the parental line exposed to acid (H^+) and alkali (OH^-).

The images show merged EP and GPEET signals. **b** Growth comparison of three individual CRAM

knockouts and parental 427 Cas9.

Supplementary Fig. 3 Characterisation of CARP3 and CARP4 knockouts. **a** Genotyping of CARP3

and CARP4 knockouts by PCR. Genomic DNA was isolated from individual clones and amplified

with the primers listed in Supplementary Table 5. **b** Community lifts of two individual CARP4

knockout clones. **c** Growth comparison of CARP3 and CARP4 knockouts and parental 427 Cas9. **d**

Determination of cell numbers on plates. Starting with an inoculum of 2 x 10^5 cells, cell numbers

were determined for 427 Cas9, the CARP3 knockout and the HA-tagged addback on days shown

in the figure. Cells were harvested by flushing the surface of the plate with PBS and cell numbers

were determined using a haemocytometer. Cells from all three lines were observed to giggle

under the microscope **e** Percentage of GPEET-positive cells (shaded area) is indicated. Cells

without prior incubation with anti-GPEET antiserum were used as a negative control (2° antibody

only, grey).

Supplementary Fig. 4 Motility and localization studies of CARP3 knockout. **a** Motility assay of

CARP3 knockouts. Centred motility tracks of individual cells are shown at identical scale. > 8000

tracks for each cell line (427 Cas9, CARP3 KO Cl.1, CARP3 KO Cl.2 and Trypanin KO, respectively)

were pooled from nine movies acquired from three technical replicates. The trypanin knockout

was used as a motility-deficient control[29]. **b** Velocity and speed of each cell line. The colour code

corresponds to the cell lines in d. Data points are displayed as box plot with whiskers representing

2.5 - 97.5 percentile, filled circles represent points outside that range. InterQuartileRange IQR

(25 - 75 percentile), with Median (50 percentile). **c** Localisation of CARP3-HA (green) by

immunofluorescence. DNA was stained with Hoechst dye. Scale bar = 10 μm.

Supplementary Fig. 5 Genotyping of the CARP3 knockout (**a**) and addback (**b**) used for the fly

experiments.

Supplementary Fig. 6 Characterisation of ACP knockouts. **a** Genotyping of ACP knockouts by PCR.

Genomic DNA was isolated and PCRs were performed with the primers listed in Supplementary

Table 5. **b** SoMo and pH taxis assays. Plates were inoculated with the cell lines indicated and 4

days later HCl and NaOH were applied to the plates (indicated by dots). Pictures were taken 16-

20 h after spotting of chemicals. **c** Growth of ACP5 single knockout clones compared to the

parental 427 Cas9 cell line. **d** Motility assays of ACP5 single knockouts. Centred motility tracks of

individual cells are shown at identical scale. > 7000 tracks for each clone were pooled from nine

movies acquired from three technical replicates. The parental line 427 Cas9 and Trypanin KO

were used as controls. **e** Velocity and speed of individual cell lines. The colour code corresponds

to the cell lines in d. Data points are displayed as box plot with whiskers representing 2.5 - 97.5

percentile, filled circles represent points outside that range. InterQuartileRange IQR (25 - 75

percentile), with Median (50 percentile). **f** Flow cytometric analysis of percentages of GPEET-

positive cells.

Supplementary Table 1 RNA-seq data. **a** Comparison of the roots and tips of projections.

Transcripts from the tips either upregulated >1 Log2FC or downregulated <1 Log2FC in

comparison to the root. FC: fold change. **b** Pairwise comparisons of untreated communities and

communities exposed to acid or alkali. Transcripts from tips that were either upregulated >1

Log2FC or downregulated <1 Log2FC. FC: fold change. Raw read files are deposited at the

European Nucleotide Archives (ENA) http://www.ebi.ac.uk/ena as study PRJEB41935.

Supplementary Table 2 Measurements of distances migrated by 427 WT and PDEB1 knockout

parasites after exposure to acid and alkali. Parasites were inoculated as described for SoMo

assays and incubated for four days. 1M HCl and NaOH (30 µl each) were added to the plates at

a distance of 1.5 cm from the edge of the community for the PDEB1 knockout and from the tips

of the protrusions for 427 WT (Fig. 3a, D = distance). The plates were sealed with parafilm and

incubated at 27 °C for 16-20 hours. Positive numbers are migration towards and negative

numbers are migration away from the spotted acid or alkali. Standard deviations were

calculated from 5 samples each.

Supplementary Table 3 Fly infections. Midgut and proventriculus infection rates from two

independent experiments with 427 Cas9, the CARP3 knockout and the untagged CARP3

addback.

Supplementary Table 4 Proteins interacting with CARP3-HA, ACP3-Myc and ACP5-Myc

identified by liquid chromatography – mass spectrometry (LC-MS).

Supplementary Table 5 List of primers used for knockouts and genotyping.

Supplementary Video 1 Early procyclic form parasites exhibit social motility.

Supplementary Video 2 Response of early procyclic forms to hydrochloric acid (HCl). Black arrows

indicate where solutions were spotted. The video is shown as a rewind loop.

Supplementary Video 3 Response of early procyclic forms to sodium hydroxide (NaOH). Black

arrows indicate where solutions were spotted. The video is shown as a rewind loop.

Supplementary Video 4 Response of late procyclic forms to sodium hydroxide (NaOH). Black

arrows indicate where solutions were spotted. The video is shown as a rewind loop.

Figure 1

Figure 2

Figure 3

a

b

untreated	OH- treatment	H+ treatment

GPEET

EP

OH⁻   OH⁻

H⁺   H⁺

Merge

c

Plate 1	Plate 2	Plate 3

GPEET

EP

H⁺   OH⁻   H⁺   OH⁻   H⁺   OH⁻

Merge

Figure 5

Figure 6

a

427 WT   PDEB1 KO

H+   OH-   H+   OH-

b

CARP3 KO   CARP3-AB-HA   427 Cas9

H+   OH-   H+   OH-   H+   OH-

c

p < .00001

Infection rates (%)

n=152   n=144   n=136

☐ Weak
☐ Intermediate
☐ Heavy

WT   CARP3 KO   AB

d

Input   HA-IP   Myc-IP

1   2   3   1   2   3   1   2   3

CARP3 — Anti-HA

ACP3/5 — Anti-Myc

1: ACP3 lysate
2: ACP5 lysate
3: Cas9 lysate

e

ACP5 sKO Cl. 2   ACP5 sKO Cl. 3   ACP5 sKO Cl. 5   427 Cas9

H+   OH-   H+   OH-   H+   OH-   H+   OH-

f

427 WT   CARP3-AB-HA+ACP5-Myc

PH

Hoechst

α-myc

merged

g

SoMo defect   ↓ Glucose   ACP5 +/-   CARP3 -/-   PDEB1 -/-   ACP1 +/-   ACP6 -/-   Hyper SoMo

Exposure to pH gradients

H+   OH-