An integrative model of plant gravitropism linking statoliths position and auxin transport

Nicolas Levernier [1,*]**, Olivier Pouliquen** [1] **and Yoël Forterre** [1]

[1]*Aix Marseille Univ, CNRS, IUSTI, Marseille, France*

Correspondence*:
IUSTI, 5 rue Enrico Fermi, 13453 Marseille cedex 13, France
nicolas.levernier@univ-amu.fr

2 ABSTRACT

Gravity is a major cue for the proper growth and development of plants. The response of plants to gravity implies starch-filled plastids, the statoliths, which sediments at the bottom of the gravisensing cells, the statocytes. Statoliths are assumed to modify the transport of the growth hormone, auxin, by acting on specific auxin transporters, PIN proteins. However, the complete gravitropic signaling pathway from the intracellular signal associated to statoliths to the plant bending is still not well understood. In this article, we build on recent experimental results showing that statoliths do not act as gravitational force sensor, but as position sensor, to develop a bottom-up theory of plant gravitropism. The main hypothesis of the model is that the presence of statoliths modifies PIN trafficking close to the cell membrane. This basic assumption, coupled with auxin transport and growth in an idealized tissue made of a one-dimensional array of cells, recovers several major features of the gravitropic response of plants. First, the model provides a new interpretation for the response of a plant to a steady stimulus, the so-called sine-law of plant gravitropism. Second, it predicts the existence of a gravity-independent memory process as observed recently in experiments studying the response to transient stimulus. The model suggests that the timescale of this process is associated to PIN turnover, calling for new experimental studies.

Keywords: Plant tropism, Gravity sensing, Auxin signaling, PIN trafficking, Modeling

1 INTRODUCTION

The detection of gravity by plants and the resulting growth response (gravitropism) offer a fascinating illustration of a multi-scale perception mechanism in living organisms (Fig. 1) (Moulia and Fournier, 2009; Morita, 2010; Toyota and Gilroy, 2013). It originates in specific cells, called statocytes, where tiny starch-accumulating amyloplasts acting as statoliths sediment under gravity at the bottom of the cells (Fig. 1C). When the plant is inclined, the repositioning of statoliths under gravity induces a relocalisation of auxin transporters (PIN proteins) at the membrane of statocytes, which generates a lateral transport of auxin toward the lower side of the shoot or the root (Cholodni–Went hypothesis) (Fig. 1B). In turn, this asymmetry in auxin concentration induces a differential growth across the plant organ, and thus its bending toward the gravity vector (Fig. 1A). Since the pioneering works of the Darwins and Sachs on plant tropisms (Darwin and Darwin, 1880; Sachs, 1887), progress has been made on every step of this gravitropic signaling pathway. Yet, basic questions remain unanswered. In particular, it is still not clear how the first physical signal generated by the sedimentation of statoliths is converted into biochemical signals downstream, to eventually produce the growth response at the plant scale (Nakamura et al., 2019).

Figure 1. Multiscale description of gravitropism. At the macroscopic scale **(A)**, the response to gravity of a shoot or a stem is achieved by differential growth across the organ, which induces a curvature of the organ. At the tissue scale **(B)**, differential growth results from a net flux of the auxin across the width (large green arrows), owing to the asymmetric distribution of auxin transporters (PINs, red circles). The local auxin fluxes are shown by the small green arrows. At the cell scale **(C)**, PIN asymmetry results from the asymmetric distribution of the statoliths position after sedimentation under gravity, which modifies PIN trafficking close to the cell membrane.

Recently, insights into the sensing mechanism and the transduction pathway have been obtained from
experiments both at the macroscopic and microscopic levels. First, the gravitropic response to permanent
stimuli (inclination of the plant), the so-called sine-law of gravitropism (Sachs, 1887; Larsen, 1969; Iino
et al., 1996; Galland, 2002; Dumais, 2013), was found to depend on the inclination but, surprisingly,
not on the intensity of gravity (Chauvet et al., 2016). Hence, statocytes behave like inclination sensors
not force sensors as previously believed. An important consequence is that the initial gravity stimulus
for gravitropism should be the position of the statoliths within statocytes (Pouliquen et al., 2017). This
position-sensor hypothesis gained a mechanistic support from the direct observation of statoliths motion
under gravity stimulation (Bérut et al., 2018). Unlike a pile of macroscopic grains like sand, statoliths were
found to move and flow at any inclination. This liquid-like behavior comes from the random agitation of
the statoliths, whose origin is not thermal but arises from the interaction of statoliths with the acto-myosin
cytoskeleton inside the cell (Sack et al., 1986; Saito et al., 2005; Nakamura et al., 2011). A second insight
came from dose-response like experiments on wheat coleoptiles, in which the gravity stimulus was applied
during a transient period only (Chauvet et al., 2019). When the shoots were inclined for short period of
time, the gravitropic response was found to deviate from the steady response and decay. The transition
occurred for a time $\tau_{\text{memory}} \sim 15$ min, which was independent of gravity and much larger than the
statoliths sedimentation time. This observation suggested the existence of a memory-integration process
in the gravitropic signaling pathway, independent of the statoliths dynamics, which integrates the initial
signal induced by statoliths displacement.

To account for these observations (position-sensor hypothesis, memory time independent of g), Chauvet
et al. (2019) built a mathematical model of gravitropism in which the gravitropic signal controlling the

differential growth was linked to the statoliths position by an integrative process of timescale τ_{memory} (a
similar approach was used in Meroz et al. (2019)). Once coupled to the statoliths dynamics and the tropic
growth motion, the model was able to reproduce the transient gravitropic response observed experimentally.
However, Chauvet et al. (2019)'s model was built on two *ad-hoc* postulates. First, it assumed that the
relation between the gravitropic signal and the statoliths position is known and given by the sine-law.
Second, it postulated the existence of the integrative process and time scale τ_{memory}, without explaining
its origin. The spatio-temporal dynamics of the molecular processes acting between the statoliths and the
growth response, such as the dynamics of PIN proteins and auxin transport, was not described.

The objective of this paper is to fill this gap by building an integrative model of plant gravitropism that
bridges the different scales of the process: (i) the initial intracellular gravitropic signal encoded in the
statoliths position, (ii) PINs dynamics at the cellular level, (ii) auxin transport at the tissue level and, finally,
(iv) differential growth and curvature at the plant organ scale. Previous models of plant gravitropism mainly
focused either on the macroscopic scale, describing how the complex spatio-temporal evolution of the
organ shape results from the interplay between differential growth and the slender geometry of the organ
(Bastien et al., 2013, 2014; Chelakkot and Mahadevan, 2017), or on the tissue level, modeling growth
mechanics (Dyson et al., 2014) or auxin transport (Band et al., 2012; Fendrych et al., 2018; Retzer et al.,
2019) in realistic 1D or 3D tissue geometries. In these latter models, the distribution of PINs in response
to plant inclination was prescribed and not linked to the intracellular dynamics of the statoliths. This is
precisely the goal of our study. Building on the recent position-sensor hypothesis, we propose a simple but
generic model of interaction between statoliths and PINs trafficking at the cell membrane, that we couple
with the classical equations of auxin transport and tissue growth. We then study the gravitropic response
predicted by the model for steady and unsteady gravity stimuli, comparing the results with the experiments
of Chauvet et al. (2016) and Chauvet et al. (2019).

2 MATERIAL AND METHODS

2.1 Link between gravitropic curvature, differential growth and auxin concentration gradient

At the plant scale, the gravitropic response is characterized by the curvature of the organ resulting from
differential growth, which itself results from auxin gradients (Cholodny–Went hypothesis). The first step
of the model is thus to relate those three quantities. For a slender organ like a shoot or a stem, the rate of
change of the local curvature $\mathcal{C}$ is related to differential growth through the following kinematic relationship:
$R\frac{\mathrm{d}\mathcal{C}}{\mathrm{d}t} = \frac{1}{\tau_g} \times \frac{\dot{\epsilon}_{\mathrm{bottom}} - \dot{\epsilon}_{\mathrm{top}}}{2\dot{\epsilon}_{\mathrm{mean}}}$, where R is the radius of the organ, $\dot{\epsilon}_{\mathrm{bottom}} - \dot{\epsilon}_{\mathrm{top}}$ is the difference of growth
rate between both sides , $\dot{\epsilon}_{\mathrm{mean}}$ is the mean growth rate and $\tau_g = \dot{\epsilon}_{\mathrm{mean}}^{-1}$ is the growth timescale (Silk,
1984; Moulia and Fournier, 2009; Bastien et al., 2013) (Fig. 1). The growth rate of plant cells is known
to be controlled by auxin, the so-called growth hormone. Auxin stimulates cell elongation by loosening
cell walls. To the best of our knowledge, the link between the local auxin concentration in walls and the
local growth rate of cells has not been robustly determined and only the response of the whole tissue to
an external addition of auxin has been investigated. It is however often assumed that growth is mainly
controlled by the auxin concentration in the vicinity of the 'skin' of the organ, as epidermal tissues are
stiffer than inner tissues (Kutschera and Niklas, 2007; Dyson et al., 2014). For the sake of simplicity, we
will here assume that the local growth rate is simply proportional to the local auxin concentration c, $\dot{\epsilon} = kc$
(Galston and Hand, 1949; Hopkins and Hüner, 2009), such that:

$$R\frac{\mathrm{d}\mathcal{C}}{\mathrm{d}t} = \frac{1}{\tau_g} \times \frac{c_{\mathrm{bottom}} - c_{\mathrm{top}}}{2c_{\mathrm{mean}}}, \tag{1}$$

94 where c_{bottom} and c_{top} are the auxin concentrations on both sides of the organ and c_{mean} the mean auxin
95 concentration. Under this assumption, the dimensionless gravitropic response deduced from the curvature
96 dynamics, $\tilde{\Delta}(t) \equiv R\tau_g \frac{dC}{dt}$, is equal to the relative auxin gradient across the organ, $\tilde{\Delta}(t) = \frac{c_{\text{bottom}} - c_{\text{top}}}{2c_{\text{mean}}}$. The
97 goal of the model is to predict how this auxin gradient establishes when the plant is tilted.

2.2 Auxin transport

Auxin transport plays a key role in shaping plants development and, as such, has been the topic of
extensive research over the past decades. Auxin transport is based on two distinct mechanisms (Goldsmith,
1977; Hopkins and Hüner, 2009; Runions et al., 2014). On the one hand, auxin in cell walls (mostly in a
protonated form) enters the neighboring cell passively, or thanks to Aux/Lax influx carriers that are evenly
distributed throughout the membrane. On the other hand, auxin inside cells (mostly in an anionic form)
can only exit thanks to active auxin efflux carriers, such as PIN proteins (Krecek et al., 2009) or ABCB
transporters (Zažímalová et al., 2010). While ABCB are evenly distributed throughout the membrane, PIN
proteins are usually polarized and can be redistributed in response to external stimuli such as gravity (in
particular PIN3, which is known to be implied in gravitropic response, see Friml et al. (2002)). Hence, an
asymmetric distribution of PIN carriers on each side of the cell can generate an active transport of auxin
from one cell to the other, resulting in a stable auxin gradient.

   To model this situation, we provide a simplified description of auxin transport in which the different
forms of auxin (proton-associated or not) are not taken into account. The tissue across the shoot or stem
(width $2R$) is modeled as a one-dimensional array of N cells of width W, separated by a cell wall of
width w (Fig. 1 b and Fig. 2). We denote c_n the auxin concentration inside the n-th cell and C_n the auxin
concentration inside the n-th wall, which are both assumed uniform (the equilibrium time of auxin in each
compartment is very fast, about 0.1 s in the cell wall and few s inside the cell taking typical values of auxin

Figure 2. One-dimensional, discrete model of auxin transport across the tissue (in reality $w \ll W$). Efflux
of auxin (solid green arrow) occurs through efflux carriers (PIN: red circle, ABCB: blue circle), whose
distribution (and thus permeabilities $P_n^{l,r}$) can be different on the right and left membrane of the cell. By
contrast, influx of auxin (green dotted arrow) occurs with a symmetrical permeability P^{in} on both side of
the cell. An asymmetry of efflux permeabilities $P^l \neq P^r$ can generate a net flux of auxin across the tissue,
yielding an auxin concentration gradient (background color gradient).

diffusion coefficients, see Kramer et al. (2007)). We also neglect auxin dilution due to cell growth and assume that auxin is neither degraded nor created, as the degradation time of auxin (of the order of the day, see Grieneisen et al. (2012)) is much longer than the minutes to hours timescales we are interested in. The efflux current of auxin (number of auxin molecules per unit time and unit surface) from the n-th cell to the left wall (resp. right wall) is given by $P_n^l c_n$ (resp. $P_n^r c_n$), where P_n^l (resp. P_n^r) is the permeability of the left (resp. right) membrane (unit m/s). Conversely, the influx current of auxin from the n-th wall to both adjacent cells is $P^{\text{in}} C_n$, where the influx permeability P^{in} is assumed uniform for all cells (see Fig. 2). The time-evolution of the concentration is then:

$$w\frac{\mathrm{d}C_n}{\mathrm{d}t} = -2P^{\text{in}}C_n + P_n^r c_n + P_{n+1}^l c_{n+1}\,, \tag{2}$$

$$W\frac{\mathrm{d}c_n}{\mathrm{d}t} = -(P_n^l + P_n^r)c_n + P^{\text{in}}(C_{n-1} + C_n)\,. \tag{3}$$

110 The cell wall size being much smaller than the cell width ($w \ll W$), the auxin concentration in the cell
111 wall can be assumed quasi-steady, $2P^{\text{in}}C_n \simeq P_n^r c_n + P_{n+1}^l c_{n+1}$, yielding:

$$2W\frac{\mathrm{d}c_n}{\mathrm{d}t} = P_{n+1}^l c_{n+1} - P_n^l c_n + P_{n-1}^r c_{n-1} - P_n^r c_n\,. \tag{4}$$

In the following, we assume that the distribution of auxin efflux carriers is the same in each cell, so that P^l
and P^r are independent of n. This is the case of shoot coleoptiles where all cells in the growing region
are similar and contain statoliths, but not the case of stems like the inflorescence of *Arabidopsis*, where
statoliths are only present on an external ring in the endodermal cells (the modification of the equation in
this case of inhomogeneous tissue is given in Appendix A). We also assume that auxin gradients occur
over a length scale much larger than the cell size. In the continuum limit [$c_n(t) \to c(x,t)$], equation (4) for
auxin transport then reduces to an advection-diffusion equation given by:

$$\frac{\partial c}{\partial t} = \mathcal{D}\frac{\partial^2 c}{\partial x^2} - v\frac{\partial c}{\partial x}\,, \tag{5}$$

where $\mathcal{D} = WP/2$ is the coefficient of diffusion and $v = \delta P/2$ the advection speed, with $P = (P^l + P^r)/2$
and $\delta P = P^r - P^l$.

The advective part of equation (5), which is responsible for auxin transport from one side to the other and
thus to the differentiated growth and the curvature of the organ, is entirely controlled by the asymmetry of
efflux permeabilities δP. Since ABCB carriers are evenly distributed, this asymmetry comes solely from
the asymmetry of PINs distribution between the right and left side of the cells:

$$\delta P = P^r - P^l = \frac{1}{S}\left(\alpha\int_S [PIN]\Big|_{\text{right}} - \alpha\int_S [PIN]\Big|_{\text{left}}\right)\,, \tag{6}$$

where α is the conductance of a single PIN carrier (unit m^3/s), S the lateral surface of the cells and
$[PIN]$ the surface concentration of PINs attached to the membrane. In the following, we assume that the
efflux permeability due to ABCB carriers is much larger than the efflux permeability due to PINs, such
that $P = (P^l + P^r)/2 \simeq (1/S)\beta\int[ABCB]$, where β is the conductance of a single ABCB carrier, is
independent of PINs concentration. This enables us to take a constant coefficient of diffusion $\mathcal{D}$ in the

model, which simplify the results without affecting much the conclusions.

131
2.3   Coupling PIN dynamics to statoliths position: biased efflux at cell scale

The previous section relates auxin transport to the asymmetry of PINs distribution at the cellular level.
We now model how this asymmetry emerges when the plant is tilted under gravity. Recently, it has been
demonstrated that the relevant gravitropic stimulus for graviperception is the statoliths position within
the statocytes (position-sensor hypothesis), and not their weight as previously believed (Chauvet et al.,
2016; Pouliquen et al., 2017). Statoliths have also been identified as key actors in the relocalisation of
PIN-proteins in response to change of gravity direction (Nakamura et al., 2019). Yet, how statoliths
position is detected and read to modify PIN polarity remains largely unknown. PINs trafficking involves
synthesis in the endoplasmic reticulum, degradation in the vacuole and recycling (Kleine-Vehn and Friml,
2008). Recycling is achieved by endocytosis, i.e., the deallocation of PIN proteins formerly attached to
the cell membrane toward the cytoplasm inside a vesicle, or by exocytosis, i.e., the reallocation of the
vesicle-carried PINs from the cytoplasm back to the cell membrane.

Following the position-sensor hypothesis, we assume that the presence of statoliths, either through direct
steric constraints or through indirect molecular signaling, modify the trafficking of PIN proteins, so that
PINs tend to enter the cell membrane preferentially on places where statoliths are in contact with it. This
mechanism is formalized as follows. The endocytosis rate of PINs, $\mathrm{d}[PIN]_i/\mathrm{d}t|_{\mathrm{endo}} = -k_{\mathrm{off},i}[PIN]_i$
where $[PIN]_i$ is the surface concentration of PIN attached to the membrane, is assumed to depend on the
presence of statoliths, with $i = 0$ if no statoliths are present and $i = 1$ if they are (see Fig. 3). Similarly,
the rate of exocytosis is written as $\mathrm{d}[PIN]_i/\mathrm{d}t|_{\mathrm{exo}} = +k_{\mathrm{on},i}[PIN]_{\mathrm{vol}}$, where $[PIN]_{\mathrm{vol}}$ is the volumic
concentration of PINs molecules inside the cell of volume WS. Two cases will be distinguished in the
model, depending on whether PINs can attach to any side of the cell ("apical/basal/lateral binding") or
only on lateral sides ("lateral binding") (see Fig. 4A). Assuming that the total number N_{tot} of PINs is
conserved during gravistimulation (Kleine-Vehn et al., 2010) leads to the following set of equations for the
PIN concentration attached to the membrane, $[PIN]_{i=0,1}$:

Figure 3. Interaction between PIN trafficking and statoliths position. The rate of reallocation k_{on} and deallocation k_{off} of PINs (bold red circles: PINs attached to the cell membrane, light red circles: PINs in bulk) depends on the presence of statoliths (grey).When the cell is tilted, the asymmetric distribution of the position of the statoliths induces a bias in the distribution of the PINs attached to the membrane.

Figure 4. Sketch of different scenario of PIN-binding. PINs are represented in red and the region with statoliths in grey. **(A)** Apical/basal/lateral-binding vs lateral-binding. **(B)** Infinite pool versus limiting pool. In the first case, the surface density of PINs is conserved whereas in the second one, the total number of PINs is conserved. **(C)** Low sensitivity of PIN to statoliths ($\mathcal{R} \simeq 1$) or high sensitivity ($\mathcal{R} \gg 1$).

$$\frac{\mathrm{d}[PIN]_0}{\mathrm{d}t} = -k_{\mathrm{off},0}\,[PIN]_0 + k_{\mathrm{on},0}\left(\frac{N_{\mathrm{tot}} - (S_0[PIN]_0 + S_1[PIN]_1)}{WS}\right)\,,$$

$$\frac{\mathrm{d}[PIN]_1}{\mathrm{d}t} = -k_{\mathrm{off},1}\,[PIN]_1 + k_{\mathrm{on},1}\left(\frac{N_{\mathrm{tot}} - (S_0[PIN]_0 + S_1[PIN]_1)}{WS}\right)\,, \tag{7}$$

where S_1 (resp. S_0) denotes the total surface in contact (resp. not in contact) with statoliths in case of
apical/basal/lateral binding, or only the lateral surfaces in contact (resp. not in contact) with statoliths in
the lateral binding case.

The form of equation (7) shows that two regimes can be distinguished, depending on whether W is large
or small compared to $k_{\mathrm{on}}/k_{\mathrm{off}}$. The first regime, called 'infinite-pool' regime in the following, corresponds
to the case where $Wk_{\mathrm{off}}/k_{\mathrm{on}}$ is so large that binding is only limited by k_{off}. The second regime, called
'limiting-pool' regime, corresponds to the opposite situation where the total number of attached carriers
is almost equal to N_{tot}: $S_0[PIN]_0 + S_1[PIN]_1 \simeq N_{\mathrm{tot}}$ (see Fig. 4B). Finally, we note that in writing
equation (7), we have neglected the diffusion of PINs inside the membrane. This is justified since, over the
time scales we are interesting in (running from minutes to one hour), PINs diffuse only over a distance
of about few micrometers, which is much smaller than the cell size (taking $0.1\mu m^2$ min for the diffusion
coefficient of a PIN, see Kleine-Vehn et al. (2011)).

Equations (1 , 5 , 7) represent a complete model describing the time-evolution of the gravitropic response
once the statoliths is known, from the PIN localization to auxin transport and bending of the plant. In

the following, we study the predictions of the model for a steady and transient stimulus. Table 1 and 2
summarizes all the physical quantities and dimensionless parameters defined from the model.

3 RESULTS

3.1 Steady gravitropic response: revisiting the sine-law

We first study the gravitropic response predicted by our model in the case of a steady inclination of
the plant θ, for long timescales when the system reaches a steady state. This situation corresponds to the
usual protocol for measuring the sensitivity of plant to gravity under steady condition, when the plant is
suddenly inclined to a fixed angle θ and its curvature (or tip angle) measured over time. After a transient,
the rate of change of curvature is found to be constant (Chauvet et al., 2016), which enables to measure the
steady gravitropic response $\tilde{\Delta}^{\text{steady}}(\theta) = R\tau_g \frac{d\mathcal{C}}{dt}$ (see Material and Methods) for each imposed angle θ.
For many plants, this relationship between the gravitropic response and the inclination angle has sine-like
shape (the response is null for $\theta = 0°$ or $\theta = 180°$ and maximal for $\theta = 90°$) and is called the 'sine-law'
in the literature (Sachs, 1887; Larsen, 1969; Iino et al., 1996; Galland, 2002; Dumais, 2013). Below, we
determine the steady gravitropic response $\tilde{\Delta}^{\text{steady}}(\theta)$ predicted by the model (eqs. 1 , 5 , 7) and compare
with measurements of the sine-law obtained previously for wheat coleoptiles over a wide range of angles
(Chauvet et al., 2016).

In the steady regime, the auxin transport equation (eq. 5) reduces to: $dJ/dx = 0$, where $J = \mathcal{D}(dc/dx) -$
vc is the auxin flux. For impermeable boundaries at $x = 0$ and $x = 2R$, the flux is null and the auxin
concentration profile is given by:

$$c(x) = c_{\text{mean}} \frac{\text{Pe} \times \exp\left(\text{Pe}\frac{x}{2R}\right)}{\exp\left(\text{Pe}\right) - 1}, \tag{8}$$

188 where $c_{\text{mean}} = (1/2R)\int_0^{2R} c(x)dx$ is the mean concentration of auxin and Pe is the Peclet number
189 comparing advection to diffusion and given by:

$$\text{Pe} \equiv \frac{2Rv}{\mathcal{D}} = N\frac{\delta P}{P} = N\frac{\alpha\left(\int_S[PIN]_{|\text{right}} - \int_S[PIN]_{|\text{left}}\right)}{\beta\int[ABCB]}. \tag{9}$$

190 For Pe $\ll 1$, the profile is linear and the auxin level in the middle of the stem is unchanged, whereas for
191 Pe $\gg 1$ the profile is strongly asymmetric with most auxin concentrated on the right, corresponding to the
192 lower side of the shoot (Fig. 5). From this steady profile of auxin, the gravitropic response can be computed
193 as $\tilde{\Delta}^{\text{steady}} = \frac{c(x=2R)-c(x=0)}{2c_{\text{mean}}}$ (eq. 1), which gives:

$$\tilde{\Delta}^{\text{steady}} = \frac{\text{Pe}}{2}. \tag{10}$$

In the steady state, the gravitropic response of the plant is thus given by the value of the Peclet number.
Previous measurements in various plant species representative of land angiosperm showed that $\tilde{\Delta}^{\text{steady}}$ is
typically of the order 1 (Chauvet et al., 2016) (for e.g. in wheat coleoptile, the maximal value of $\tilde{\Delta}^{\text{steady}}$
obtained for a 90 degrees inclination is about 0.7), meaning that the Peclet number is typically of the
order 1. Therefore, the auxin profile across the shoot is expected to be close to linear (Fig. 5A) (and not
a pronounced exponential, Fig. 5B). A consequence is that growth, which was assumed proportional to
the auxin concentration, varies also linearly from one side of the shoot to the other during the gravitropic
response.

Figure 5. Stationary auxin profile for **(A)** small and **(B)** large Peclet number.

202 The next step to determine $\tilde{\Delta}^{\text{steady}}$ is to compute the Peclet number given by equation (9), i.e., the relative
203 distribution of PIN between the left and right side of the cell. In the steady regime, the concentration of PIN
204 attached to a membrane not covered by statoliths ($[PIN]_0^{\text{steady}}$), or covered by statoliths ($[PIN]_1^{\text{steady}}$), is
205 given by (see eq. 7):

$$[PIN]_0^{\text{steady}} = \frac{N_{\text{tot}}}{S}\left(\frac{k_{\text{off},0}W}{k_{\text{on},0}} + \frac{1}{S}\left[S_0 + \frac{k_{\text{on},1}\,k_{\text{off},0}}{k_{\text{on},0}\,k_{\text{off},1}}S_1\right]\right)^{-1},$$

$$[PIN]_1^{\text{steady}} = \frac{N_{\text{tot}}}{S}\left(\frac{k_{\text{off},1}W}{k_{\text{on},1}} + \frac{1}{S}\left[S_1 + \frac{k_{\text{on},0}\,k_{\text{off},1}}{k_{\text{on},1}\,k_{\text{off},0}}S_0\right]\right)^{-1}. \tag{11}$$

206 Noting S_i^l (resp. S_i^r) the surface of the left (resp. right) side of the cell not covered ($i = 0$) or covered ($i = 1$)
207 by statoliths, we have $\int_S[PIN]_{|\text{right}} - \int_S[PIN]_{|\text{left}} = [PIN]_0^{\text{steady}}(S_0^r - S_0^l) + [PIN]_1^{\text{steady}}(S_1^r - S_1^l)$.
208 Finally, since $S_0^l + S_1^r = S_0^r + S_0^l = S$, where S is the lateral surface of the cells and using equation (9),
209 we have:

$$\tilde{\Delta}^{\text{steady}} = \frac{\alpha N(\mathcal{R}-1)(S_1^r - S_1^l)[PIN]_0^{\text{steady}}}{2\beta\int[ABCB]}, \tag{12}$$

210 with

$$[PIN]_0^{\text{steady}} = \frac{N_{\text{tot}}}{S}\frac{1}{\mathcal{R}}\left(\mathcal{P} + \frac{S_1}{S}\left(1 + \frac{S_0}{S_1\mathcal{R}}\right)\right)^{-1}, \tag{13}$$

211 and:

$$\mathcal{R} = \frac{[PIN]_1^{\text{steady}}}{[PIN]_0^{\text{steady}}} = \frac{k_{\text{on},1}\,k_{\text{off},0}}{k_{\text{on},0}\,k_{\text{off},1}}, \qquad \mathcal{P} = \frac{Wk_{\text{off},1}}{k_{\text{on},1}}. \tag{14}$$

The expressions (12–14) show that the steady gravitropic response is proportional to $(\mathcal{R}-1)(S_1^r - S_1^l)$.
For $\theta > 0$ as in figure 1, the difference $(S_1^r - S_1^l)$ is positive since statoliths sediment toward the right
side of the cell. Therefore, to obtain a 'normal' gravitropic response ($\tilde{\Delta}^{\text{steady}} > 0$, i.e., a larger auxin
concentration at the bottom side of the shoot), the ratio $\mathcal{R}$ must be larger than 1. The parameter $\mathcal{R}$ is the
ratio between the concentration of PINs in a zone with statolith and in a zone without statolith, hence the
larger $\mathcal{R}$, the more PINs in the region with statoliths compared to region without statolith (see Fig 4C). The
other main parameter controlling the gravitropic response is the pool number $\mathcal{P}$ characterizing the ratio of

Figure 6. Steady gravitropic response $\tilde{\Delta}^{\text{steady}}$ (arbitrary amplitude) as a function of the inclination angle θ, for $\mathcal{R}$ either large or close to 1, in the case of **(A)** infinite pool ($\mathcal{P} \gg 1$), **(B)** limiting pool ($\mathcal{P} \ll 1$) with apical/basal/lateral binding, **(C)** limiting pool ($\mathcal{P} \ll 1$) with lateral binding only. Note that in the case of an infinite pool, results are the same in the a/b/l-binding or l-binding case. Geometrical parameters used are $H_0 = 4d$, $W = 10d$, $H = 25d$ where d stands for the diameter of a statolith.

endocytosis to exocytosis, where $\mathcal{P} \gg 1$ corresponds to the infinite-pool regime and $\mathcal{P} \ll 1$ corresponds
to the limiting-pool regime (see Fig 4B).

Once $\mathcal{R}$ and $\mathcal{P}$ are set, the final step is to compute how the different surfaces covered (and not covered)
by the statoliths vary as function of the inclination angle θ. To this end, one has to know the final position of
statoliths when a cell is tilted. Recently, we addressed this question and showed that statoliths at the bottom
of statocytes behave like an effective liquid on long timescale, due to the agitation of statoliths by the cell
activity (Bérut et al., 2018). Therefore, the final free surface of the statoliths pile is horizontal, as sketched
in figure 3. This key feature of the flowing behavior of statoliths allows us to reduce the computation of the
surfaces touched or not by the statoliths ($S_{0,1}^{r,l}$, S_0 and S_1 in eqs. 12, 13) to a purely geometrical problem,
which depends on three parameters: the angle of inclination θ, the aspect ratio of the cell H/W and the
initial aspect ratio of the statolith pile H_0/W (see Fig. 3). The corresponding relationships are given in
Appendix B.

Figure 6 presents the typical steady gravitropic response $\tilde{\Delta}^{\text{steady}}(\theta)$ predicted by the model (eqs. 12, 13,
14) as a function of θ, in the case of an infinite pool ($\mathcal{P} \gg 1$) or a limiting pool ($\mathcal{P} \ll 1$), and for two
extreme values of $\mathcal{R}$: $\mathcal{R} \simeq 1$ (red curve, low influence of statoliths on PIN binding) and $\mathcal{R} \gg 1$ (blue curve,
strong influence of statoliths on PIN binding). The geometry used for the cell aspect ratio and the statoliths
pile ratio is taken from experimental observations of wheat coleoptile statocytes, with a typical aspect ratio
$H/W = 2.5$ and $H_0/W = 1/2.5$. In the infinite pool regime (Fig. 6A), the gravitropic response presents
a convex shape with a strong peak close to $90°$ whatever the value of $\mathcal{R}$, in disagreement with the usual
sine-law shape. The amplitude of this peak in this regime also strongly depends on the cell geometry: the
more elongated the cell, the higher the peak because in this case more lateral surface is available for PIN
attachment without any PIN dilution (see Fig 4B). By contrast, in the limiting-pool regime (Fig. 6B,C),
the response strongly depends on $\mathcal{R}$, the peak at $90°$ being much smaller for $\mathcal{R} \gg 1$ than for $\mathcal{R} \simeq 1$.
Interestingly, the response in this case also depends on whether PIN can attach on every sides of the cell
(apical/basal/lateral binding) or only on the lateral sides (lateral binding case), as attachment to the 'useless'
apical and basal sides contribute to deplete PIN from the available pool. Overall, we see that only one
case is compatible with the concave 'sine-law' shape observed experimentally: a limiting pool of PIN with
lateral binding and $\mathcal{R} \gg 1$ (blue curve in Fig. 6C).

Figure 7. Modified sine-law $\tilde{\Delta}^{\text{steady}}$ as a function of the inclination angle θ. Comparison between the model prediction for $\mathcal{P} \ll 1$ (limiting pool) and l-binding (eq. 15 with $\mathcal{R} = 25$ and $\mathcal{A} = 1.3$, red line) and experiments on wheat coleoptiles (symbols, Chauvet et al. (2016)). Geometrical parameters used in the model for the statocyte are $H_0 = 4d$, $W = 10d$, $H = 25d$ where d stands for the diameter of a statolith. Error bars are the mean value of the data by binning the [0,180] interval into 20 boxes.

247   In the following, we thus assume that PIN recycling occurs in the limited-pool regime ($\mathcal{P} \gg 1$), with
248 lateral binding only. In this limit, and using expression of the surfaces given in Appendix B, the steady
249 gravitropic response given by equations (12, 13, 14) can be written as:

$$\tilde{\Delta}^{\text{steady}}(\theta) = \frac{1}{2}\mathcal{A}(\mathcal{R}-1) \begin{cases} \dfrac{W\tan\theta}{2H + 2(\mathcal{R}-1)H_0} & \text{if } W\tan\theta < 2H_0 \\ \dfrac{\sqrt{2H_0 W\tan\theta}}{2H + (\mathcal{R}-1)\sqrt{2H_0 W\tan\theta}} & \text{if } 2H_0 < W\tan\theta < H^2/(2H_0) \\ \dfrac{1}{\mathcal{R}+1} & \text{else} \end{cases} \quad (15)$$

with $\mathcal{A} = \frac{\alpha N_{\text{tot}} N}{\beta \int [ABCB]}$. The different conditions stand for cases where statoliths are totally absent of the
left side, or totally covering the right side (see Fig. 7). Once the geometry of the cell and of the statoliths
pile are fixed, the predicted gravitropic response depends on two dimensionless parameters: $\mathcal{R}$ and $\mathcal{A}$.
Figure 7 presents the experimental measurements of the 'sine-law' obtained by Chauvet et al. (2016) on
wheat coleoptiles, together with the best fit of the data using a least-square method. Reasonable agreement
between theory and experiments is obtained with $\mathcal{R} \simeq 25$ (or larger values as the shape converges in this
case) and $\mathcal{A} \simeq 1.3$. It is interesting to note that within our position-sensor framework, the predicted steady
gravitropic response is not a simple 'sine-law', but rather a piece-wise curve with an overall concave shape.
This law is also not universal and can be affected by several anatomical and physiological properties, such
as the geometry of the cell H/W, the amount of statoliths H_0/W, or the molecular signaling machinery
(embedded in the parameters $\mathcal{R}$ and $\mathcal{A}$).

3.2 Transient gravitropic response: dose-response law

The previous results deal with the steady gravitropic response obtained when the gravity stimulus (the
angle of inclination θ of the plant) is permanent. We now turn to the study of the transient gravitropic
response, i.e., when the system has not yet reached the steady state. This situation typically corresponds
to 'dose-response' like experiments, in which the gravity stimulus is applied during a transient time ΔT
only. Using such protocol on wheat coleoptiles, Chauvet et al. (2019) revealed the existence of an intrinsic
'memory' time τ_{memory} in the gravitropic response. For $\Delta T \gg \tau_{\mathrm{memory}}$, the response was constant and
equal to the steady response $\tilde{\Delta}^{\mathrm{steady}}$. However, for $\Delta T \lesssim \tau_{\mathrm{memory}}$, the response was smaller and became
proportional to ΔT (Fig. 8). The memory time $\tau_{\mathrm{memory}} \sim 15$ min identified in these experiments was
longer than the sediment time of statoliths (~ 2 min) but shorter than the growth timescale (hours). It thus
reflects a temporal process in the gravitropic signaling pathway that remains to be identified. We address
below this question in the framework of the model.

The set of equations (5)-(7) give a complete description of the transient gravitropic response (we assume
that sedimentation of statoliths is fast enough that the surfaces in equation 7 can be computed from their
equilibrium values – see Appendix B). Two different typical times control the dynamics: τ_{aux}, describing
the transport of auxin across the tissue of length $2R$, and τ_{PIN}, describing the dynamics of PIN at molecular
scale. The time scale associated to auxin transport τ_{aux} can be estimated using equation (5) for a constant
coefficient of diffusion $\mathcal{D}$ and transport velocity v (i.e., PIN distribution). In this case, relaxation towards
the stationary profile (eq. 8) is exponential and occurs on a time scale set by the inverse of the shortest
non-vanishing eigen-mode of equation (5) (Mohsen and Baluch, 1983):

$$\tau_{\mathrm{aux}} = \frac{1}{\pi^2 + (\mathrm{Pe}^2/4)} \times \frac{(2R)^2}{\mathcal{D}} = \frac{1}{\pi^2 + (\mathrm{Pe}^2/4)} \times \frac{(2R)\mathrm{Pe}}{v} \qquad (16)$$

Estimating the Peclet number from the steady gravitropic response ($\mathrm{Pe} = 2\tilde{\Delta}^{\mathrm{steady}} \approx 1$, see Fig. 7), and
the auxin transport speed v from measurements of the speed of auxin pulses in plant tissues ($v \simeq 3~\mu\mathrm{m/s}$,
Goldsmith (1977); Rashotte et al. (2003)), gives $\tau_{\mathrm{aux}} \simeq 30$ s (taking $2R \sim 1$ mm). This is much shorter
than the memory time $\tau_{\mathrm{memory}} \sim 15$ min evidenced in dose-response experiments, and even shorter than
the statoliths sedimentation time. This suggests that the dynamics of the gravitropic response is controlled
by τ_{PIN}, rather than by the auxin diffusion time τ_{aux}.

The time scale τ_{PIN} is set by the slowest characteristic time of the system (eq. 7) describing PIN dynamics.
From the eigenvalues of this linear system, two timescales are obtained, which are solutions of:

$$\tau^{-2} - (k_{\mathrm{off},0} + k_{\mathrm{off},1} + [k_{\mathrm{on},0}S_0 + k_{\mathrm{on},1}S_1]/WS)\tau^{-1} \qquad (17)$$
$$+ (k_{\mathrm{off},0} + k_{\mathrm{on},0}S_0/WS)(k_{\mathrm{off},1} + k_{\mathrm{on},1}S_1/WS) - k_{\mathrm{on},0}k_{\mathrm{on},1}S_0S_1/W^2S^2 = 0\,.$$

289 In the limiting pool case ($WSk_{\mathrm{off,i}}/k_{\mathrm{on,i}}S_i \ll 1$), the two solutions are:

$$\tau_1 = \frac{WS}{k_{\mathrm{on},0}S_0 + k_{\mathrm{on},1}S_1} \qquad \text{and} \qquad \tau_2 = \frac{k_{\mathrm{on},0}S_0 + k_{\mathrm{on},1}S_1}{k_{\mathrm{off},0}k_{\mathrm{on},1}S_1 + k_{\mathrm{off},1}k_{\mathrm{on},0}S_0}, \qquad (18)$$

290 with $\tau_1 \ll \tau_2$. Therefore, the slowest timescale of the gravitropic signaling pathway, which sets τ_{memory},
291 should be given by $\tau_{\mathrm{PIN}} = \tau_2$. Note that when preferential attachment in region with statoliths is achieved via
292 a strongly increased attachment rate ($k_{\mathrm{on},1} \gg k_{\mathrm{on},0}$), and not by change of detachment rate ($k_{\mathrm{off},0} \simeq k_{\mathrm{off},1}$),
293 then $\tau_{\mathrm{PIN}} = k_{\mathrm{off}}^{-1}$. Conversely, if it is achieved by decreased detachment rate ($k_{\mathrm{off},0} \gg k_{\mathrm{off},1}$) and not by

Figure 8. Gravitropic response to a transient inclination (dose-response like protocol). **(A)** Maximal gravitropic response reached during the dynamics as function of the inclination time ΔT for $\theta = 45°$. The blue solid line is the model prediction using $\tau_{\text{PIN}} = 13$ min ($\tau_{\text{aux}}/\tau_{\text{PIN}}$ and $\tau_1/\tau_{\text{PIN}} \ll 1$, other parameters are fixed as in Fig. 7). Symbols correspond to the aggregation results of (Chauvet et al., 2019) under $1g$ and $3g$. **(B)** Evolution of the tip angle after an inclination $\theta = 50°$ predicted by the model with the same parameters (blue solid line) and in the experiments of Chauvet et al. (2016) (orange thick line). The predicted model must be shifted by a constant time $\tau_{\text{reac}} = 13$ min to match the experimental curve (dashed blue line). Inset: early time behavior of the gravitropic response predicted by the model in log-log scale.

change of attachment rate ($k_{\text{on},0} \simeq k_{\text{on},1}$), then $\tau_{\text{PIN}} = k_{\text{off},0}^{-1}(1 + S_1/S_0) \simeq k_{\text{off},0}^{-1}$. Remarkably, in this last
case, the equilibration time of PIN is not controlled by the slowest rate of detachment ($k_{\text{off},1}$), but by the
fastest one ($k_{\text{off},0}$), due to the limiting pool.

To check these predictions, we solve the model for a transient gravitropic stimulus that reproduces the
protocol of Chauvet et al. (2019). The inclination θ is set to $45°$ for a transient time ΔT and then put back
to zero, the gravitropic response being defined as the maximal auxin gradient reached during the dynamics:
$\tilde{\Delta}^{\text{trans}} = \text{Max}[\frac{c_{\text{bottom}}(t) - c_{\text{top}}(t)}{2c_{\text{mean}}}]$. Figure 8A compares the experimental data of Chauvet et al. (2019) with
the prediction of the model for $\tau_{\text{aux}}/\tau_{\text{PIN}} \ll 1$, using the parameters $\mathcal{R}$ and $\mathcal{A}$ already fixed by the steady
response (see Fig. 7). Good agreement is obtained using $\tau_{\text{PIN}} = 13$ min as the only fitting parameter. This
result shows that τ_{PIN} is playing the role of the memory time evidenced by the experiments of Chauvet et al.
(2019). If $\Delta T \ll \tau_{\text{PIN}}$, PIN transporters have not enough time to rearrange before the end of the stimulus,
and the response is weak. Conversely if $\Delta T \gg \tau_{\text{PIN}}$, PINs have time to rearrange and reach their steady
repartition before the end of the stimulus, and the response is maximal, similar to the one of a permanent
stimulus.

We finish our analysis by investigating the full temporal dynamics of the gravitropic response after a
sudden inclination $\theta = 50°$. Figure 8B presents the time evolution of the shoot curvature $\mathcal{C}(t)$ (or similarly
$\theta_{tip}(t)$ for small curvatures) predicted by the model for $\tau_{\text{PIN}} = 13$ min (blue curve), together with the
experimental data of Chauvet et al. (2016) (yellow curve). The time scale of the curvature change τ_{PIN} is
well captured by the model. However, to match the experiments, the model has to be shifted in time by a
constant time $\tau_{\text{reac}} \approx 13$ min. Such delay or reaction time of the gravitropic response after the stimulus was
already noticed by Chauvet et al. (2019), but does not seem to be captured by the model. Actually, a careful
analysis of the temporal behavior of the model (assuming $\tau_1 \ll \tau_{\text{aux}} \ll \tau_{\text{PIN}} = \tau_2$) shows that the auxin
gradient at early times increases as $\Delta c \sim t^{3/2}$ (so that $\mathcal{C} \propto \int_0^t \Delta c\, dt \propto t^{5/2}$) as long as $t < \tau_{\text{aux}}$, before

varying as $\Delta c \sim t$ for $\tau_{\mathrm{aux}} < t < \tau_{\mathrm{PIN}}$ (Inset of Fig. 8B). These scaling laws are thus not compatible with
a very flat initial response.

4   DISCUSSION

In this paper, we have derived a multi-scale model of plant gravitropism which links the different steps of
the gravitropic signaling pathway: (i) the initial intracellular perception of gravity by statoliths, (ii) the
transduction of this physical signal into a biochemical signal through the reorganization of PINs at the
membrane of statocytes, (iii) the intercellular signal transmission via auxin transport and (iv) asymmetric
organ growth. The main originality of the model lies in the mechanistic link we propose between the
statoliths position and the dynamics of PINs, based on the recent position-sensor hypothesis (Pouliquen
et al., 2017). This basic assumption coupled to auxin transport and growth in an idealized tissue made of a
one-dimensional array of cells recovers several major features of the gravitropic response of plants.

4.1   A new interpretation of the sine-law of plant gravitropism

The first main result concerns the steady gravitropic response to a permanent gravity stimulus, $\tilde{\Delta}^{\mathrm{steady}}(\theta)$.
For many plants, this response takes the form of an inclination-dependent law with a sine-like shape, called
for this reason the sine-law (Sachs, 1887; Larsen, 1969; Iino et al., 1996; Galland, 2002; Dumais, 2013).
This sine-law has long been interpreted in terms of a force sensor mechanism, for the projected weight of the
statoliths on the lateral surface of the cell varies with the sine of the inclination angle (Audus, 1969; Barlow,
1995). However, recent experiments showing that the response is independent of the gravity intensity
have dismissed this force-sensing hypothesis, calling for a new interpretation of the sine law (Chauvet
et al., 2016; Pouliquen et al., 2017). A key result of the model is that it predicts an inclination-dependent
steady gravitropic response $\tilde{\Delta}^{\mathrm{steady}}(\theta)$ without invoking a force-based mechanism. In the model, the initial
gravitropic stimulus is the statoliths position at the cell membrane, not their weight. Since statoliths behave
on long time like a liquid (Bérut et al., 2018), their position in steady state is a purely geometrical cue,
which depends only on the cell inclination. As a result, the steady response predicted by the model depends
on the inclination but not on the gravity intensity, in agreement with the observations.

In the model, the actual shape of the gravitropic response $\tilde{\Delta}^{\mathrm{steady}}(\theta)$ is never a pure sine law. It depends
on several parameters, related either to geometric factors, such as the aspect ratios of the statocytes and of
the sedimented statoliths pile, or to molecular processes: intensity of coupling between statoliths position
and PINs through parameter $\mathcal{R}$; ratio of endocytosis to exocytosis through parameter $\mathcal{P}$; ratio between
the conductance of PIN carriers to the total conductance of auxin transporters through parameter $\mathcal{A}$. For
elongated cells and shallow statoliths piles such as those of wheat coleoptile statocytes, the shape of the
response tends to be sine-like only in the case of a strong coupling between the statoliths and PINs ($\mathcal{R} \gg 1$),
and for a number of PINs conserved along the cell membrane (limiting pool regime, $\mathcal{P} \ll 1$). This latter
assumption is common in models of auxin transport (Runions et al., 2014; Retzer et al., 2019), while the
strong coupling assumption is compatible with the large asymmetry in PINs localization observed upon
gravity stimulation (Friml et al., 2002; Harrison and Masson, 2008; Kleine-Vehn et al., 2010). Interestingly,
although gravity is absent from the model, the gravitropic response depends on the amount of statoliths
through the geometrical aspect ratio of the statoliths pile H_0/W. The model could thus account for previous
experiments using starch-less and starch-excess mutants, which showed a modified gravitropic response
when the number of statoliths is changed (Kiss et al., 1997; Vitha et al., 2007; Pouliquen et al., 2017).
Finally, it is worth noting that the model assumes that the statoliths form a static pile at the bottom of
the cell, while statoliths actually exhibit a dynamic and random agitation due to the interaction with the
cytoskeleton (Sack et al., 1986; Saito et al., 2005; Nakamura et al., 2011; Bérut et al., 2018). We might
expect this agitation to reduce the averaged contact time between the statoliths and the cell membrane,

thereby decreasing the coupling between statoliths and PINs. It would be interesting to extend the model in
order to incorporate such effect of agitation on the gravitropic response. The model could then be compared
with the behavior of agravitropic mutants in *Arabidopsis thaliana* like *sgr9*, whose weaker response is
likely associated to an abnormally strong agitation of the statoliths (Nakamura et al., 2011).

Overall, our model suggests that the classical sine-law of plant gravitropism might not be universal, as its
shape and amplitude could depend on several anatomical and physiological parameters. Full measurements
of the gravitropic response of plants over a wide range of inclination are scarce and mostly performed on
shoot coleoptiles. It would be interesting to perform systematic measurements of the sine-law on other
plant organs (root, stem), to see if the shape of the sine-law is affected by different statocyte geometries
and tissue organization.

4.2 The gravity-independent memory process in dose-response laws is likely
associated to PIN dynamics

The second main result of our study concerns the gravitropic response to a transient gravity stimulus.
For a sudden inclination applied at time $t = 0$, the model predicts that the response reaches the steady
response $\tilde{\Delta}^{\text{steady}}(\theta)$ only after a time large compared to a 'memory' timescale τ_{memory}, corresponding to
the slowest timescale introduced in the model. Therefore, when a stimulus is applied only during a transient
time $\Delta T < \tau_{\text{memory}}$, a weaker gravitropic response is predicted, following a dose-response like law. In the
model, the memory time is not associated with the sediment time of the statoliths, which is assumed to be
much shorter than the other timescales of the gravitropic signaling pathway (a valid assumption for a gravity
intensity like Earth gravity). Our model is therefore compatible with the recent experiments of Chauvet
et al. (2019) performed on wheat coleoptiles, which show a dose-like behavior of the gravitropic response
with a memory time τ_{memory} independent of gravity. The model also provides the explicit origin of this
memory process, which was postulated in Chauvet et al. (2019). In the model, two different processes can
lead to the temporal filtration of the initial signal associated with statoliths position: auxin transport across
the tissue and the dynamics of PIN turnover at the molecular scale. Our study suggests that the limiting
process is actually controlled by PIN dynamics ($\tau_{\text{memory}} = \tau_{\text{PIN}}$), auxin transport being too fast to account
for the memory time measured experimentally ($\sim$ 10-20 min). Interestingly, visualization of the PIN3
auxin efflux carrier in root columella cells after a sudden change in the gravity vector indicates a time scale
of about 10 min for complete relocation (Friml et al., 2002), a duration very close to the memory time
measured by (Chauvet et al., 2019). Although our model successfully captures the existence and origin
of a gravity-independent memory process in the signaling pathway, it is not able to describe the delay
time $\tau_{\text{reaction}} \sim$ 10 min observed between the application of the stimulus and the first gravitropic response
(Chauvet et al., 2019). This delay may have different origins: the time needed to reorganize the cytoskeleton
implied in the transport of PIN carriers, the time needed by a PIN to go from the pool towards the plasma
membrane, or the time of incorporation of a PIN into the membrane. It is worth noting that a similar
timescale of about 10 min was identified in the gravity-sensing columella cells for the internalization of
PIN3 from the plasma membrane into vesicles (Kleine-Vehn et al., 2010). Further experiments combining
a transient stimulus with pharmacological and genetic approaches would be needed to confirm the key role
of the PIN turnover timescale in the gravitropic response. Besides, it would be interesting to see what is the
influence of this hierarchy of time-scales on the response of a plant to oscillatory stimulus (e. g. wind) or
on the dynamic competition between gravitropism and proprioception (Bastien et al., 2013).

4.3 Back to the statoliths/PIN coupling assumption

We conclude by discussing the possible origin of the coupling between statoliths and PINs, which is
at the core of our model. Although the respective roles of statoliths and PIN auxin transporters in plant

gravitropism are well established, the link between the two is still not clear (see Nakamura et al. (2019)
for a recent review of the possible molecular actors involved). In our model, we have used a very general
hypothesis for this coupling based on the recent finding that the relevant gravitropic stimulus is the statoliths
position inside the gravisensing cells (Chauvet et al., 2016; Pouliquen et al., 2017; Bérut et al., 2018).
We have postulated that statoliths in contact with the cell membrane bias the exocytosis and endocytosis
rate of PIN recycling, therefore inducing an asymmetry of PIN distribution when statoliths reposition
in response to plant inclination. Our results suggest that this interaction between PINs and statoliths is
strong, as large values of the parameter $\mathcal{R}$ are needed to match the experimental gravitropic response.
This interaction between statoliths and PINs could involve a complex molecular pathway that remains
to be unveiled (Strohm et al., 2014). However, more direct mechanisms of interaction could occur. For
example, statoliths could modify PIN vesicle-mediated transportation to the membrane by modifying the
architecture of the actin cytoskeleton. Another possibility would be that internalization of PINs from the
membrane is reduced by the presence of statoliths, for example by simple steric effects. Indeed, direct
visualizations reveal a length scale of $\sim 1\,\mu$m for the endosome formation, which is not far from statolith
size (Kleine-Vehn et al., 2010). Interestingly, such direct interaction of statoliths and the cytoskeleton
machinery was put forward as an explanation of gravitropism in rhizoids and protonemata, such as the
single-cell alga *Chara* (Sievers et al., 1996). A last possibility could be that PINs cluster in regions where
there is no statolith, while they do not cluster when statoliths are present. Indeed, for a conserved number
of PIN proteins, clustering reduces the efflux efficiency, as this diffusion process scales not linearly but
as the square-root of the number of carriers (Bénichou and Voituriez, 2014; Valet et al., 2019). Such a
clustering has been highlighted by Kleine-Vehn et al. (2011), but, to our knowledge, no comparison has
been done between regions of the membrane in contact or not in contact with statoliths. Further studies
would be needed to discriminate between these molecular mechanisms.

CONFLICT OF INTEREST STATEMENT

427 The authors declare that the research was conducted in the absence of any commercial or financial
428 relationships that could be construed as a potential conflict of interest.

AUTHOR CONTRIBUTIONS

429 N.L., O.P., and Y.F. designed research. N.L. built and analyzed the model with inputs from O.P. and Y.F.
430 N.L., O.P., and Y.F. wrote the paper..

FUNDING

431 This work was supported by the European Research Council (ERC) under the European Union's Horizon
432 2020 research and innovation programme (grant agreement N°647384) and by the French National Agency
433 (ANR) under the program Blanc Grap2 (ANR-13-BSV5-0005-01).

ACKNOWLEDGMENTS

434 The authors would like to thank Valérie Legué and Bruno Moulia for the many stimulating discussions we
435 have continually had on this subject.

APPENDIX A: CASE OF STATOLITHS ONLY PRESENT ON A RING

Let be a central region between l and $2R - l$ with no statolith (and constant efflux rate P_0). We can redo the
calculation leading to eq. (4) in this geometry. Clearly, equations will be similar in each region, with $v = 0$ in
the central one. We have to take care to the continuity of the flux at the jonction of each regions. We can show
that $J(l) = (P + \delta P)c(l^-) - P_0 c(l^+)$ and $J(2R - l) = P_0 c((2R - l)^-) - Pc((2R - l)^+)$. At steady state, we

thus get a jump of auxin concentration: $c(l^+) = (P+\delta P)/P_0 \; c(l^-)$ and $c((2R-l)^+) = P_0/P \; c((2R-l)^-)$.
In the limit of small Peclet number, we thus get $\Delta c/c = (P_0/P)(vl/\mathcal{D})$.

APPENDIX B: EXPRESSION OF THE GEOMETRIC PARAMETERS AS A FUNCTION OF θ

We give here the value of the surfaces $S_i^{r,l}(\theta)$ that are necessary to compute the Peclet number as a function of the angle (see eq. 15). As statoliths are solid, the shaded area in Fig. 3 is conserved and is thus equal to $H_0 W$. We note $\theta_1 = \arctan 2H_0/W$ the angle for which the horizontal level meets the lower left corner, and $\theta_2 = \arctan H^2/(2H_0 W)$ the one for which it meets the upper right one (we assume $H_0 < H/2$). Elementary geometry calculations give:

$$S_0^l = \begin{cases} H - H_0 + \dfrac{W \tan \theta}{2} & \text{if } \theta < \theta_1 \\ H & \text{else} \end{cases} \tag{19}$$

$$S_1^l = \begin{cases} H_0 - \dfrac{W \tan \theta}{2} & \text{if } \theta < \theta_1 \\ 0 & \text{else} \end{cases} \tag{20}$$

$$S_0^r = \begin{cases} H - H_0 - \dfrac{W \tan \theta}{2} & \text{if } \theta < \theta_1 \\ H - \sqrt{2H_0 W \tan \theta} & \text{if } \theta_1 < \theta < \theta_2 \\ 0 & \text{else} \end{cases} \tag{21}$$

$$S_1^r = \begin{cases} H_0 + \dfrac{W \tan \theta}{2} & \text{if } \theta < \theta_1 \\ \sqrt{2H_0 W \tan \theta} & \text{if } \theta_1 < \theta < \theta_2 \\ H & \text{else} \end{cases} \tag{22}$$

In the case of apical binding, this leads to

$$S_0 = \begin{cases} 2(H - H_0) & \text{if } \theta < \theta_1 \\ 2H - \sqrt{2H_0 W \tan \theta} & \text{if } \theta_1 < \theta < \theta_2 \\ H & \text{else} \end{cases} \tag{23}$$

$$S_1 = \begin{cases} 2H_0 & \text{if } \theta < \theta_1 \\ \sqrt{2H_0 W \tan \theta} & \text{if } \theta_1 < \theta < \theta_2 \\ H & \text{else} \end{cases} \tag{24}$$

442 In the case of apical/basal/lateral binding, the expressions for the total surfaces are slightly modified due to
443 attachment at apical and basal side.

REFERENCES

Audus, L. (1969). Geotropism. In *The physiology of plant growth and development* (McGraw-Hill, London). 203–241

Band, L. R., Wells, D. M., Larrieu, A., Sun, J., Middleton, A. M., French, A. P., et al. (2012). Root gravitropism is regulated by a transient lateral auxin gradient controlled by a tipping-point mechanism. *Proceedings of the National Academy of Sciences* 109, 4668

Barlow, P. W. (1995). Gravity perception in plants: a multiplicity of systems derived by evolution? *Plant, Cell & Environment* 18, 951–962

Bastien, R., Bohr, T., Moulia, B., and Douady, S. (2013). Unifying model of shoot gravitropism reveals proprioception as a central feature of posture control in plants. *Proceedings of the National Academy of Sciences* 110, 755–760

Bastien, R., Douady, S., and Moulia, B. (2014). A unifying modeling of plant shoot gravitropism with an explicit account of the effects of growth. *Frontiers in plant science* 5, 136

Bénichou, O. and Voituriez, R. (2014). From first-passage times of random walks in confinement to geometry-controlled kinetics. *Physics Reports* 539, 225–284

Bérut, A., Chauvet, H., Legué, V., Moulia, B., Pouliquen, O., and Forterre, Y. (2018). Gravisensors in plant cells behave like an active granular liquid. *Proceedings of the National Academy of Sciences* 115, 5123

Chauvet, H., Moulia, B., Legué, V., Forterre, Y., and Pouliquen, O. (2019). Revealing the hierarchy of processes and time-scales that control the tropic response of shoots to gravi-stimulations. *Journal of Experimental Botany* 70, 1955–1967

Chauvet, H., Pouliquen, O., Forterre, Y., Legué, V., and Moulia, B. (2016). Inclination not force is sensed by plants during shoot gravitropism. *Scientific reports* 6, 35431–35431

Chelakkot, R. and Mahadevan, L. (2017). On the growth and form of shoots. *Journal of The Royal Society Interface* 14, 20170001

Darwin, C. and Darwin, S. F. (1880). *The power of movements in plants* (Murray)

Dumais, J. (2013). Beyond the sine law of plant gravitropism. *Proceedings of the National Academy of Sciences* 110, 391–392

Dyson, R. J., Vizcay-Barrena, G., Band, L. R., Fernandes, A. N., French, A. P., Fozard, J. A., et al. (2014). Mechanical modelling quantifies the functional importance of outer tissue layers during root elongation and bending. *New Phytologist* 202, 1212–1222

Fendrych, M., Akhmanova, M., Merrin, J., Glanc, M., Hagihara, S., Takahashi, K., et al. (2018). Rapid and reversible root growth inhibition by tir1 auxin signalling. *Nature plants* 4, 453–459

Friml, J., Wiśniewska, J., Benková, E., Mendgen, K., and Palme, K. (2002). Lateral relocation of auxin efflux regulator pin3 mediates tropism in arabidopsis. *Nature* 415, 806–809

Galland, P. (2002). Tropisms of avena coleoptiles: sine law for gravitropism, exponential law for photogravitropic equilibrium. *Planta* 215, 779–784

Galston, A. W. and Hand, M. E. (1949). Studies on the physiology of light action; auxin and the light inhibition of growth. *Am J Bot* 36, 85–94

Goldsmith, M. H. M. (1977). The polar transport of auxin. *Annual Review of Plant Physiology* 28, 439–478

Grieneisen, V. A., Scheres, B., Hogeweg, P., and M Marée, A. F. (2012). Morphogengineering roots: comparing mechanisms of morphogen gradient formation. *BMC Systems Biology* 6, 37

Harrison, B. R. and Masson, P. H. (2008). Arl2, arg1 and pin3 define a gravity signal transduction pathway in root statocytes. *The Plant Journal* 53, 380–392

Hopkins, W. G. and Hüner, N. P. (2009). *Introduction to Plant Physiology 4th ed*

Iino, M., Tarui, Y., and Uematsu, C. (1996). Gravitropism of maize and rice coleoptiles: dependence on the stimulation angle. *Plant, cell & environment* 19, 1160–1168

Kiss, J. Z., Guisinger, M. M., Miller, A. J., and Stackhouse, K. S. (1997). Reduced gravitropism in hypocotyls of starch-deficient mutants of arabidopsis. *Plant and Cell Physiology* 38, 518–525

Kleine-Vehn, J., Ding, Z., Jones, A. R., Tasaka, M., Morita, M. T., and Friml, J. (2010). Gravity-induced pin transcytosis for polarization of auxin fluxes in gravity-sensing root cells. *Proceedings of the National Academy of Sciences* 107, 22344–22349

Kleine-Vehn, J. and Friml, J. (2008). Polar targeting and endocytic recycling in auxin-dependent plant development. *Annual Review of Cell and Developmental Biology* 24, 447–473

Kleine-Vehn, J., Wabnik, K., Martinière, A., Łangowski, Ł., Willig, K., Naramoto, S., et al. (2011). Recycling, clustering, and endocytosis jointly maintain pin auxin carrier polarity at the plasma membrane. *Molecular systems biology* 7, 540–540

Kramer, E. M., Frazer, N. L., and Baskin, T. I. (2007). Measurement of diffusion within the cell wall in living roots of arabidopsis thaliana. *Journal of experimental botany* 58, 3005–3015

Krecek, P., Skupa, P., Libus, J., Naramoto, S., Tejos, R., Friml, J., et al. (2009). The pin-formed (pin) protein family of auxin transporters. *Genome biology* 10, 249–249

Kutschera, U. and Niklas, K. (2007). The epidermal-growth-control theory of stem elongation: an old and a new perspective. *Journal of plant physiology* 164, 1395–1409

Larsen, P. (1969). The optimum angle of geotropic stimulation and its relation to the starch statolith hypothesis. *Physiologia Plantarum* 22, 469–488

Meroz, Y., Bastien, R., and Mahadevan, L. (2019). Spatio-temporal integration in plant tropisms. *Journal of the Royal Society Interface* 16, 20190038

Mohsen, M. F. N. and Baluch, M. H. (1983). An analytical solution of the diffusion-convection equation over a finite domain. *Applied mathematical modelling* 7, 285–287

Morita, M. T. (2010). Directional gravity sensing in gravitropism. *Annual review of plant biology* 61, 705–720

Moulia, B. and Fournier, M. (2009). The power and control of gravitropic movements in plants: a biomechanical and systems biology view. *Journal of experimental botany* 60, 461–486

Nakamura, M., Nishimura, T., and Morita, M. T. (2019). Gravity sensing and signal conversion in plant gravitropism. *Journal of experimental botany* 70, 3495–3506

Nakamura, M., Toyota, M., Tasaka, M., and Morita, M. T. (2011). An arabidopsis e3 ligase, shoot gravitropism9, modulates the interaction between statoliths and f-actin in gravity sensing. *The Plant Cell* 23, 1830–1848

Pouliquen, O., Forterre, Y., Bérut, A., Chauvet, H., Bizet, F., Legue, V., et al. (2017). A new scenario for gravity detection in plants: the position sensor hypothesis. *Physical Biology* 14, 035005

Rashotte, A. M., Poupart, J., Waddell, C. S., and Muday, G. K. (2003). Transport of the two natural auxins, indole-3-butyric acid and indole-3-acetic acid, in arabidopsis. *Plant Physiology* 133, 761

Retzer, K., Akhmanova, M., Konstantinova, N., Malínská, K., Leitner, J., Petrášek, J., et al. (2019). Brassinosteroid signaling delimits root gravitropism via sorting of the arabidopsis pin2 auxin transporter. *Nature communications* 10, 1–15

Runions, A., Smith, R. S., and Prusinkiewicz, P. (2014). Computational models of auxin-driven development. In *Auxin and its role in plant development* (Springer). 315–357

Sachs, J. (1887). *Lectures on the Physiology of Plants* (Clarendon Press)

Sack, F. D., Suyemoto, M. M., and Leopold, A. C. (1986). Amyloplast sedimentation and organelle saltation in living corn columella cells. *American journal of botany* 73, 1692–1698

Saito, C., Morita, M. T., Kato, T., and Tasaka, M. (2005). Amyloplasts and vacuolar membrane dynamics in the living graviperceptive cell of the arabidopsis inflorescence stem. *The Plant Cell* 17, 548–558

Sievers, A., Buchen, B., and Hodick, D. (1996). Gravity sensing in tip-growing cells. *Trends in plant science* 1, 249–250

Silk, W. K. (1984). Quantitative descriptions of development. *Ann. Rev. Plant Physiol.* 35, 479–518

Strohm, A. K., Barrett-Wilt, G. A., and Masson, P. H. (2014). A functional toc complex contributes to gravity signal transduction in arabidopsis. *Frontiers in plant science* 5, 148

Toyota, M. and Gilroy, S. (2013). Gravitropism and mechanical signaling in plants. *American journal of botany* 100, 111–125

Valet, M., Pontani, L.-L., Voituriez, R., Wandersman, E., and Prevost, A. M. (2019). Diffusion through nanopores in connected lipid bilayer networks. *Phys. Rev. Lett.* 123, 088101

Vitha, S., Yang, M., Sack, F. D., and Kiss, J. Z. (2007). Gravitropism in the starch excess mutant of arabidopsis thaliana. *American journal of botany* 94, 590–598

Zažímalová, E., Murphy, A. S., Yang, H., Hoyerová, K., and Hošek, P. (2010). Auxin transporters–why so many? *Cold Spring Harbor perspectives in biology* 2, a001552

Physical quantities	Definition	Unit
R	radius of the organ	(m)
$\mathcal{C}$	bending curvature of the organ	(m^{-1})
W	cell width	(m)
w	cell wall width	(m)
H	cell height	(m)
H_0	statolith pile height before inclination	(m)
S	cell lateral surface	(m^2)
$S_{0,1}$	total surface of the cell not (0) or (1) in contact with statoliths	(m^2)
$S_0^{l,r}$	surface of the left (l) or right (r) side of the cell not in contact with statoliths	(m^2)
$S_1^{l,r}$	surface of the left (l) or right (r) side of the cell in contact with statoliths	(m^2)
$\dot{\epsilon}$	growth rate	(s^{-1})
τ_g	$\tau_g = \dot{\epsilon}_{\text{mean}}^{-1}$ growth timescale	(s)
c	auxin concentration inside the cell	(mol m^{-3})
C	auxin concentration in the cell wall	(mol m^{-3})
$P^{l,r}$	auxin efflux permeability of the left (l) or right (r) side of the cell	(m^{-1})
P^{in}	auxin influx permeability	(m^{-1})
α	conductance of a single PIN carrier	$(\text{m}^3\ \text{s}^{-1})$
β	conductance of a single ABCB carrier	$(\text{m}^3\ \text{s}^{-1})$
P	$P = \frac{P^l+P^r}{2} = \frac{1}{S}\beta\int[ABCB]$	(m^{-1})
δP	$\delta P = P^r - P^l = \frac{1}{S}\left(\alpha\int_S[PIN]_{|\text{right}} - \alpha\int_S[PIN]_{|\text{left}}\right)$	(m^{-1})
$\mathcal{D}$	auxin coefficient of diffusion $\mathcal{D} = \frac{WP}{2}$	$(\text{m}^2\ \text{s}^{-1})$
v	auxin advection speed $v = \frac{\delta P}{2}$	(m^{-1})
$k_{\text{off},0,1}$	endocytosis rate when statoliths are (1) or are not (0) in contact with the membrane	(s^{-1})
$k_{\text{on},0,1}$	exocytosis speed when statoliths are (1) or are not (0) in contact with the membrane	(m s^{-1})
τ_{aux}	timescale for auxin transport across the tissue $\tau_{\text{aux}} = \frac{1}{\pi^2+(\text{Pe}^2/4)} \times \frac{(2R)^2}{\mathcal{D}}$	(s)
τ_{PIN}	timescale for PIN turnover $\tau_{\text{PIN}} = \frac{k_{\text{on},0}S_0+k_{\text{on},1}S_1}{k_{\text{off},0}k_{\text{on},1}S_1+k_{\text{off},1}k_{\text{on},0}S_0}$	(s)

Table 1. List of dimensional quantities in the model with their definition and unit.

Dimensionless parameters	Definition	Meaning
θ		Inclination of the plant
N		Number of cells across the tissue
N_{tot}		Total number of PIN carriers per cell
$\tilde{\Delta}$	$\tilde{\Delta} = \dfrac{c_{\text{bottom}} - c_{\text{top}}}{2c_{\text{mean}}}$	Gravitropic response
Pe	$\text{Pe} = \dfrac{2Rv}{\mathcal{D}}$	Peclet number comparing auxin advection to diffusion
$\mathcal{A}$	$\mathcal{A} = \dfrac{\alpha N_{\text{tot}} N}{\beta \int [ABCB]}$	Effect of PIN conductance vs ABCB conductance on auxin transport
$\mathcal{R}$	$\mathcal{R} = \dfrac{[PIN]_1^{\text{steady}}}{[PIN]_0^{\text{steady}}} = \dfrac{k_{\text{on},1}\, k_{\text{off},0}}{k_{\text{on},0}\, k_{\text{off},1}}$	Intensity of statoliths/PINs coupling
$\mathcal{P}$	$\mathcal{P} = \dfrac{W k_{\text{off},1}}{k_{\text{on},1}}$	Ratio of endocytosis to exocytosis (pool number)

Table 2. List of the dimensionless parameters used in the model with their definition and meaning.